Sulfur reduction coupling with anaerobic ammonium oxidation power the origin of

life

Peng Bao[1, 2, *], Guo-Xiang Li[1, 2, 3], Hu Li[1]

[1]Key Lab of Urban Environment and Health, Institute of Urban Environment, Chinese

Academy of Sciences, Xiamen 361021, P. R. China

[2]Ningbo Urban Environment Observation and Station, Chinese Academy of Sciences,

Ningbo 315800, P. R. China

[3]University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Address correspondence to:

Dr. Peng Bao

Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, P. R.

China

E-mail: pbao@iue.ac.cn; neilbao17@163.com

Abstract

The geochemical energy that drives the transition from non-life to life is as yet unknown. We found that sulfurous species reduction coupling with anaerobic ammonium oxidation (Sammox), could provide the primordial redox equivalents and energy for the reductive acetyl-CoA pathway combined with incomplete reductive tricarboxylic acid (rTCA) cycle and reductive amination. Fe-S mineral/metal alloy catalysis and thiols/thioesters as energy couplers could enhance the efficiency of Sammox-powered proto-anabolic networks. Genomic analysis of Sammox bacterium *Ralstonia thioammoniphilus* GX3-BWBA implied that this ancient metabolism in modern microbes should contain two stages according to ammonium transformation, —oxidation of ammonium to nitrite and denitrification. The incomplete rTCA cycle and reductive acetyl-CoA pathway were all identified in *R. thioammoniphilus* metabolic networks, which were responsible for chemolithotrophic metabolism. Sammox drove the coupling of the biochemical transformation of C, H, O, N, S, and/or Fe simultaneously in Hadean alkaline hydrothermal systems, thereby permitting the emergence of life. The results bridged the gap in the transition from geochemistry to biochemistry.

Introduction

The transition from non-life to life occurred in the context of geochemical energy derived from element-coupled transformation[1-3]. The chemistry of life is based on redox reactions, that is, successive transfers of electrons and protons from the six major elements, i.e., C, H, O, N, S, and P[1,4]. The H_2/CO_2 redox couple, which can occur in submarine hydrothermal vents, has been proposed to drive the reductive acetyl-CoA pathway, an ancient metabolic route[3,5-7]. Nevertheless, the primordial energy source of the H_2/CO_2 redox couple suffers from the difficulty that the exergonic reaction competes with the endergonic reaction for available H_2[8]. The subsequent surface metabolism and thioester world theories still could not answer the question, how the required reduced carbon compounds have been synthesized[9,10]. Despite this limitation, these theories have emphasized the important roles of thioesters and Fe-S mineral catalysis for driving the primordial rTCA cycle, a central anabolic biochemical pathway whose origins are proposed to trace back to geochemistry[9,11,12].

Native iron/metals were recently shown to promote the reductive acetyl-CoA pathway and rTCA cycle and strongly support the feasibility of these two primordial synthetic pathways[13,14], although they were generally considered to be rare near the Earth's surface[13] and cannot support the long elemental transition from geochemistry to biochemistry. Proto-anabolic networks consisting of the reductive acetyl-CoA pathway together with the complete/incomplete rTCA cycle as primordial synthetic pathways are therefore more logical[3,15-18]. However, the initial driving force for the rise of proto-anabolic networks is still unclear. Moreover, the roles of N and S

geochemical transformation in the origin of life have been largely ignored because

computational analysis has implied that N and S are essential for thermodynamically

feasible phosphate-independent metabolism before the rise of last universal common

ancestor (LUCA)[19]. The proto-anabolic networks would necessarily be driven by C, H,

O, N, S, and/or Fe coupling transformation at their earliest stage of the transition from

geochemistry to biochemistry. The co-evolution of the metabolism of those elements

may provide strong explanatory power for the origin of life and explain why the

structure of metabolic networks is as it is. We therefore speculate that

thermodynamically feasible sulfurous species reduction coupling with anaerobic

ammonium oxidation reaction[20,21] (Eqs. 1, 2), with/without the catalysis of Fe-S

minerals and thioesters, may have been the primordial power force for the rise of

proto-anabolic networks.

$$8NH_4^+ + 2SO_4^{2-} + 2HCO_3^- \rightarrow 4N_2 + 2HS^- + CH_3COO^- + 12H_2O + 5H^+ \quad \Delta G_r = -20.0 \text{ kJ mol}^{-1} \quad (1)$$

$$4NH_4^+ + S_2O_3^{2-} + 2HCO_3^- \rightarrow 2N_2 + 2HS^- + 2HCOO^- + 5H_2O + 4H^+ \quad \Delta G_r = -13.4 \text{ kJ mol}^{-1} \quad (2)$$

On early Earth, elemental sulfur, sulfite, and thiosulfate were produced

abundantly from volcanic and hydrothermal SO_2 or from H_2S oxidation by iron oxides

in sulfide-rich hydrothermal fluid[2,15,22]. In simulated hydrothermal systems with

conditions of 300 °C and high pressure, nitrite was readily converted to NH_3[23], and

Ni-Fe metals and alloys were also effective catalysts of N_2 reduction to NH_3 in

Hadean hydrothermal systems[24]. There would have been much higher CO_2

concentrations in the oceans on early Earth because there was perhaps up to 1000

times more CO_2 in the atmosphere than that today[25].

Hence, when sulfurous species and NH_3 in Hadean hydrothermal systems made

contact with CO_2, there were spontaneous electron and proton transfers for energy

generation and organic molecule synthesis via sulfurous species reduction coupling

with ammonium oxidation. We termed this process Sammox, which falls outside of

previous studies of the coupling elemental biogeochemical cycles[1,26]. A prebiotic

reaction should occur with the ability to branch out into S and N biochemistry, which

could contribute to the autocatalysis and evolution of primordial metabolic networks.

Thus, Sammox may facilitate the synthesis of thioesters and amino acids, which are

essential for the self-amplification of phosphate-independent metabolic networks

before the rise of LUCA. We expect that prebiotic chemical evidence of

Sammox-powered CO_2 fixation, thioesters, and amino acid synthesis, combined with

genetic analysis of a representative Sammox microbe, will provide profound insights

into the earliest origins of life and fill in the missing link of the emergence of

biochemistry from geochemistry.

Sammox and the construction of proto-anabolic networks

We set out to verify the feasibility of Sammox-driven prebiotic reductive

acetyl-CoA pathway, rTCA cycle, and reductive amination and to determine how

methanethiol and Fe-S mineral catalysis enhance the reactions in both thiosulfate- and

sulfate-fueled Sammox systems. Formate and acetate were the products of

Sammox-driven reductive acetyl-CoA pathway (Fig. 1). We haven't detected pyruvate

as the end-product of Sammox-driven reductive acetyl-CoA pathway because

pyruvate quickly entered the incomplete rTCA cycle as a reaction substrate (Figs. 2,

3). We qualitatively identified methyl acetate as a product in Sammox reaction

systems (Extended Data Fig. 1), implying that methanol should be an intermediate of

Sammox-driven reductive acetyl-CoA pathway (Fig. 3a). More importantly, this result

confirmed the capacity for Sammox-powered ester bond formation, which was critical

for the synthesis of complex organic molecules, such as lipids and RNA[27].

We quantitatively identified succinate and α-ketoglutarate, and qualitatively

identified polypyrrole (data not shown) and glutamate as products of Sammox-driven

incomplete rTCA cycle (Figs. 1, 2, Extended Data Fig. 2). Glutamate came from the

reductive amination of α-ketoglutarate with ammonium, while the existence of

polypyrrole implied that there should be pyrrole derived from succinate. This

provided the possibility of biocatalysis. We haven't detected other intermediates of

the incomplete rTCA cycle, potentially because of their low productivity. There were

no side reactions of pyruvate to lactate or α-ketoglutarate to α-hydroxyglutarate[16],

which allowed Sammox-driven proto-anabolic networks to persist and self-amplify.

Sulfate-fueled Sammox was more effective than thiosulfate-fueled Sammox

(Figs. 1, 2), due to the higher energy yield of the former reaction (Eqs. 1, 2). Both

Fe-S minerals and methanethiol could enhance the efficiency of the thiosulfate-fueled

Sammox reaction but not the sulfate-fueled Sammox reaction (Fig 1), logically

implying that the thiosulfate-fueled Sammox reaction should be the primordial

reaction driving the construction of proto-anabolic networks. Note that methanethiol

was more effective than Fe-S minerals for promoting the thiosulfate-fueled Sammox

reaction (Fig. 1a). It incorporates into C, H, O and N biochemical transformation via

thiol-thioester exchange[9], transforming itself and other organic reaction products into

new organic products[10], therefore resulted in the coupling biochemical transformation

of C, H, O, N, and S and the expansion of primordial metabolic networks. We

deduced that when reaction conditions were met in Hadean hydrothermal systems,

Sammox-generated proton gradient could modulate reduction potential similar to

applying a voltage, leading to electrocatalytic CO_2 reduction with/without catalysis of

Fe-S minerals and thioesters[28,29]. Thus, almost all of the essential metabolic

precursors for biosynthesis could be provided in one geological setting. The organics

formed within the vent pores in Hadean hydrothermal systems, where they should

concentrate via processes, such as thermophoresis, and potentially form structures,

such as lipid membranes lining hydrophobic walls[7,15,30]. Thereby, Sammox gave rise

to the origin of not free-living organic precursor on the surface of Fe-S minerals[31] (Fig.

3c).

As one of the products of Sammox, sulfide began to increase in the Hadean

oceans, facilitating thiols synthesis on Fe-S mineral surface[28] and leading to de

Duve's thioester world and the boom of S biochemistry. As a result, the S isotope

geochemical evidence can be traced back to 3.8 Ga and has been well

preserved[4,22,32,33]. Biogenic sulfide reacted with soluble Fe^{2+} and/or Ni^{2+}/Co^{2+}/Se to

maintain a continuous self-supply of freshly precipitated FeS and/or metals alloys and

even subsequent enzyme and redox protein active centers, such as ferredoxin[34]. This

may represent a route for S and Fe assimilation, which would facilitate escape of the

not free-living organic precursor from the surface of Fe-S minerals. As catalytic

properties improved, the yield of proto-anabolic networks increased, ultimately

extending the pathway through to C_6 tricarboxylic acids. Sulfur biogeochemistry

stimulated iron redox geochemistry; therefore, ferrous iron was released due to the

abiotic reduction of iron oxyhydroxide by biogenic sulfide, thus resulting in phosphite

liberation from ferruginous sediments[35]. In this study, we will not discuss how P

incorporated into primordial metabolic networks, but infer that Sammox-driven

proto-anabolic networks could facilitate self-evolution through optimizing the

ambient environment to make it sufficiently stable and habitable for life.

Discovery of Sammox microbe

We successfully isolated a pure Sammox culture (designated GX3-BWBA) that

could conserve energy from the reduction of sulfurous species (thiosulfate, sulfate,

sulfite, and elemental sulfur) coupled to anaerobic ammonium oxidation. Strain

GX3-BWBA appeared as bean-shaped cells with a diameter of 0.6 µm and a length of

1.0 µm and harbored a single flagella with a bacterial capsule (Fig. 4a, b, Extended

Data Fig. 3). Comparative sequence analysis of 16S rRNA genes revealed a 99%

query cover and 99% sequence identity between strain GX3-BWBA and a series of

uncultured *Ralstonia* bacterial clones (Fig. 4c). Thus, strain GX3-BWBA was

phylogenetically clustered within the *Ralstonia* genus (Fig. 4c). GX3-BWBA grew to

a maximal density of 1.6×10^5 CFU ml^{-1} in anaerobic freshwater mineral medium

containing bicarbonate, ammonium, and thiosulfate/sulfate as the sole carbon and

energy sources, suggesting chemoautotrophy (Fig. 4b). Approximately 8 µM

ammonium was sufficient to support the chemoautotrophic growth of strain

GX3-BWBA, demonstrating high affinity for ammonium (data not shown).

Thiosulfate was the preferred electron acceptor over sulfate by GX3-BWBA (Fig. 4b).

There was no significant growth of GX3-BWBA under conditions without ammonium

or sulfurous species (Fig. 4b); thus, ammonium and sulfurous species were essential

factors for GX3-BWBA chemoautotrophic growth. We provisionally classified this

Sammox bacterium as "*Ralstonia thioammoniphilus*" (thi.o.am.mo.ni' phi.lus. Gr. n.

thion sulfur; L. neut. n. *sal ammoniacum* salt of Ammon (NH_4Cl); Gr. adj. *phylos*

loving; N.L. neut. adj. *thioammoniphilus* sulfur-ammonium-loving).

As a probable ancient chemoautotroph, Sammox microbe preferred other

sulfurous species, except for sulfate, potentially because elemental sulfur, thiosulfate,

and sulfite were produced abundantly from ancient volcanic and hydrothermal SO_2 or

from sulfide oxidation by iron oxides in sulfide-rich hydrothermal fluid[2,15,22].

Thiosulfate constituted 68–78% of the immediate HS^- oxidation products and was

involved in a dynamic HS^--$S_2O_3^{2-}$ cycle in anoxic marine and freshwater sediments[36].

In contrast, sulfate would have had only very limited, localized significance[2].

We further tested thiosulfate- and sulfate-fueled Sammox reactions using *R.*

thioammoniphilus, and the end product N_2 generation rate was measured (Fig. 4c, d).

The generation rate of N_2 in the thiosulfate treatment group was slightly higher than

that in the sulfate treatment group (Fig. 4c, d). The production rate of $^{29}N_2$ was about

2.5 nmol ml^{-1} day^{-1}, which was significantly higher than $^{30}N_2$ (approximately 2.5 pmol

ml^{-1} day^{-1}; Fig. 4c, d). Figure 5 shows the coupled dynamic transformation of

sulfurous species and ammonium during Sammox process. In thiosulfate-fueled

Sammox, sulfide was the product of the thiosulfate reduction, whereas sulfite was an

intermediate (Fig. 5a). Ammonium oxidation accompanied by thiosulfate reduction

was significant compared with that in the control group. As intermediate of

ammonium oxidation, nitrite significantly increased on day 27 (Fig. 5b). Based on the

low concentration, we did not determine the dynamic generation of dinitrogen to

avoid leakage. There was weak but significant sulfate reduction in the sulfate-fueled

Sammox group, accompanied by sulfide generation (Fig. 5c). The concentration of

nitrite also showed a slight increase during the ammonium oxidation process (Fig. 5d).

The concentrations of sulfurous species and ammonium showed no significant

changes in the control group (Fig. 5). The assumed net thiosulfate-/sulfate-fueled

Sammox reactions are shown in Eqs. 1 and 2. The generation of precursors of CO_2

fixation was thermodynamically favorable by Sammox.

Notably, quite a few iron oxides could be abiotically reduced by sulfide

generated from Sammox in an iron-rich environment, then an illusion of ferric iron

reduction coupled to anaerobic ammonium oxidation (Feammox) arises[37-39]. Indeed, it

is still unclear to what extent Sammox overlaps with Feammox; however, we can

infer that Sammox may be widely distributed in different anoxic environments based

on the ubiquity of *Ralstonia* spp. and its high affinity for ammonium. Moreover, as an

ancient type of metabolism, there may be a variety of microbes that still preserved this

living strategy.

Genomic and N, S, and C metabolic gene analyses

To elucidate the mechanisms of Sammox reaction in modern microorganisms, we sequenced the genome of *R. thioammoniphilus* (Extended Data Table 1, Extended Data Fig. 4). There were denitrification genes in the *R. thioammoniphilus* genome, including copper-containing nitrite reductase (*nirK*), nitric oxide reductase (*norB*), and nitrous oxide reductase (*nosZ*) (Fig. 6a). This result confirmed that nitrite was the expected intermediate product of Sammox and that the thiosulfate-/sulfate-fueled Sammox reactions in *R. thioammoniphilus* contained two stages: oxidation of ammonium to nitrite (Eqs. 3 and 4)[21], followed by the final step of the denitrification pathway which is reduction of nitrite to dinitrogen (Eq. 5).

$$4NH_4^+ + 3S_2O_3^{2-} \rightarrow 4NO_2^- + 6HS^- + H_2O + 8H^+ \tag{3}$$

$$4NH_4^+ + 3SO_4^{2-} \rightarrow 4NO_2^- + 3HS^- + 4H_2O + 5H^+ \tag{4}$$

$$+2NO_2^- + 8H^+ + 6e^- \rightarrow N_2 + 4H_2O \tag{5}$$

We did not find any other anaerobic ammonium oxidation-related functional genes in the genome of *R. thioammoniphilus* (Fig. 6a). Thus, the ammonium oxidation in Sammox was different from that in complete ammonium oxidation (Comammox) and anaerobic ammonium oxidation (Anammox)[26].

Our main concern is how ammonium was oxidized by thiosulfate/sulfate into nitrite. Sammox in *R. thioammoniphilus* may still need an iron-sulfur catalyst that inherited from proto-anabolic networks. As lacking of genetic evidence about Sammox, we predicted that ferredoxin oxidoreductase (e.g., nitrite/sulfite reductase

ferredoxin domain protein and ferredoxin-nitrite reductase) or other iron-sulfur

enzymes encoded by genes with unknown sequences may mediate sulfur reduction

coupling with ammonium oxidation to nitrite (Eqs. 3–6, Extended Data Schematic 1).

Here, thiosulfate should be first transformed into sulfite, which could be reduced by

ferredoxin. Hence, Sammox reversed the nitrite reduction process. This process was

quite similar to a nitrite generation pathway in Feammox process (Eq. 7)[26,37,38],

implying its theoretical feasibility. Electrons and protons generated in the first stage of

Sammox reaction were used to drive nitrite reduction to dinitrogen. As a highly

exergonic reaction, energy released from nitrite reduction drove CO_2 fixation.

$$NH_4^+ + 6 \text{ oxidized ferredoxin} + 2H_2O \rightarrow NO_2^- + 6 \text{ reduced ferredoxin} + 8H^+ \qquad (6)$$

$$NH_4^+ + 6Fe^{3+} + 2H_2O \rightarrow NO_2^- + 6Fe^{2+} + 8H^+ \qquad (7)$$

However, so far we do not clearly know the molecular mechanisms of sulfate

reduction coupling with ammonium oxidation to nitrite. *R. thioammoniphilus* contains

no ATP sulfurylase (Sat)-coding gene (Fig. 6a), suggesting that sulfate reduction in

sulfate-fueled Sammox is different from dissimilatory sulfate reduction. Dissimilatory

sulfate reduction may be later than sulfate reduction in sulfate-fueled Sammox,

according to isotope geochemical evidence[22,40,41]. This is why sulfite reductase

(Extended Data Table 2, Extended Data Fig. 5) showed a much more extensive

distribution than Sat (Extended Data Table 3, Extended Data Fig. 6) and, also the

evidence for ancient metabolic networks led to the emergence of living systems prior

to the incorporation of P[19].

As we have inferred that the transition of geochemistry to biochemistry of S and

N should start at the same starting point. Sulfite and nitrite reductases belong to the

only class of enzymes that share a common architecture as well as a requirement for a

siroheme cofactor[42]. The phylogenetic tree indicated that the putative multifunctional

sulfite/nitrite reductase of archaea and eukaryota gathered into one and three clusters,

respectively. Putative interdomain lateral gene transfer may result in the distribution

of some archaea and eukaryota multifunctional sulfite/nitrite reductases into bacterial

clusters (Fig. 6b). Similar to sulfite reductase, nitrite reductase is widely distributed in

all types of modern organisms (Extended Data Table 4, Extended Data Fig. 7), and

indeed, sulfite reductases from some sources can catalyze the reduction of both sulfite

and nitrite[42]. This phenomenon suggested that S and N biochemistry may have a

common evolutionary origin derived from Sammox.

Similar to the sulfite/nitrite reductase, some biomolecules as remnants of

Sammox and ancient metabolism may be hidden in the architecture of the metabolic

networks of *R. thioammoniphilus*. In theory, such "metabolic fossils" for the origin of

the coupled transformation of elements should be widely distributed in all types of

modern organisms as well. Thiosulfate may be the most preferential electron acceptor

for Sammox in Hadean hydrothermal systems, leading to the wide distribution of

multifunctional rhodanese (thiosulfate sulfurtransferase) in prokaryotes, eukaryotes,

and archaea (Extended Data Table 5, Extended Data Fig. 8). Rhodanese has been

proposed to have an assimilatory role using dithiol dihydrolipoate as the sulfur

acceptor and acting as a sulfur insertase involved in the formation of prosthetic groups

in iron-sulfur proteins, such as ferredoxin[43,44]. This implies that rhodanese may be the

primary mechanism for the formation of the iron-sulfur center of primordial enzymes

to catalyze proto-anabolic networks in LUCA, thus indicating its ancient nature.

Rhodanese can also catalyze a sulfur dissimilatory metabolic reaction which is

thiosulfate cleavage to sulfite[44] (Scheme 1), thereby facilitate thiosulfate involve in

the first stage of Sammox.

$$S_2O_3^{2-} + 2R\text{-}S^- \rightarrow SO_3^{2-} + R\text{-}S\text{-}S\text{-}R + S^{2-}$$

287   Scheme 1

The structural similarity between rhodanese and Cdc25 phosphatases indicated

the common evolutionary origin of the two enzyme families[45]; alternatively,

phosphatases may originate from the rhodanese family because P incorporation

occurred later than S incorporation in metabolic networks. This evidence implied the

relationship between S and P metabolic evolution.

Six candidate genes encoding components of the reductive acetyl-CoA pathway

were identified as formate dehydrogenase (fdoI), 5,10-methenyl-H4 folate

cyclohydrolase (folD), 5,10-methylene-H4 folate dehydrogenase (folD),

5,10-methylene-H4 folate reductase (metF), CO dehydrogenase/acetyl-CoA synthase

(acs), and pyruvate:ferredoxin oxidoreductase (PFOR) (Fig. 6a). Three gene

homologs 10-formyl-H4 folate synthetase, methyl-H4 folate:corrinoid iron-sulfur

protein methyltransferase, and corrinoid iron-sulfur protein appeared to be missing. It

is possible that the present genomic analysis was insufficient to distinguish whether R.

$thioammoniphilus$ could facilitate CO_2 fixation via the incomplete reductive

acetyl-CoA pathway or there were functional related enzymes encoded by genes with

unknown sequences[46].

In annotations of the rTCA cycle, two essential gene homologs appeared to be

missing (Fig. 6a). One of the missing genes, α-oxoglutarate synthase, catalyzes

reductive carboxylation from succinate to α-ketoglutarate, and the other is citryl-CoA

synthetase. Two potential reversible ATP-dependent citrate lyase subunits (ACL)

(citE, mdcC) were identified (Fig. 6a), which was a key indication for autotrophic

CO_2 fixation via the rTCA cycle instead of citrate synthase[47]. These results suggested

the potential of *R. thioammoniphilus* for carbon fixation via the incomplete reductive

acetyl-CoA pathway and rTCA cycle.

Final remarks: from metabolic innovation to evolution

This finding, Sammox powered the origin of life, provided strong evidence of

prebiotic Sammox-driven proto-anabolic networks and a Sammox microbe, *R.*

thioammoniphilus. Our study was partially supported by the theory of a

chemoautotrophic origin of life supported by surface metabolism and a primordial

iron-sulfur world[9,11]. Here, Fe-S minerals and/or metal alloy catalysis enhanced

product generation of Sammox rather than providing hydrogen for CO_2 reduction. The

Fe-S mineral surfaces, where transition metal ions, such as Ni^{2+}, Co^{2+}, or Se, are

catalytically active[48], and also the very place facilitated Sammox-driven amnio acids

and thioesters synthesis. Therefore, a mixture of Fe-S minerals and transition metal

ions may have been the evolutionary precursor of the enzymatic active center, when

chelated by amnio acids, converted to biocatalysts[29,34]. In this regard, our study could

explain the occurrence of enzymes with Fe-S reaction centers in the electron transport

chains of most known extant bioenergy flows in all three kingdoms of life. Thioesters

derived during Sammox reaction provided the energetic and catalytic framework of

Sammox, and feedback and feedforward to Sammox-driven proto-anabolic networks,

ultimately leading to metabolic reproduction and innovation[48]. Thus, Sammox drove

the emergence of the primordial structure and function of the not free-living organic

precursor, a ligand sphere, held together by bonding to the surfaces of Fe-S minerals

(Fig. 3c). The feedback coupling between the primordial metabolic networks and their

environment could shape the evolution of both[49]. As a self-regulating system, C, H, O,

N, S, and/or Fe coupling metabolic networks could facilitate self-evolution through

optimization of the ambient environment, making it stable and habitable for life. The

first and foremost goal of primordial metabolic networks was to obtain sufficient

phosphorus. Sammox-driven sulfur biochemistry launched iron redox geochemistry;

therefore, phosphite was liberated from ferruginous sediments[35], permitting

biochemistry feedback to geochemistry. If and only if P reached a certain

concentration in the Hadean oceans, sulfur biochemistry could lead to the emergence

of phosphorus biochemistry, and the RNA world could have replaced the thioester

world.

Our findings regarding Sammox-driven proto-anabolic networks suggested that

primordial metabolic networks would rise from coupling transformation of C, H, O, N,

S, and/or Fe at their earliest stage of the transition from geochemistry to biochemistry.

The elements of the chemical components that supporting Sammox-driven

proto-anabolic networks ultimately became essential or trace elements of life,

providing a strong explanation for the unique features of life. Exploration of Sammox

provides a new perspective for understanding the emergence of biochemistry from

geochemistry and highlights the fundamental significance of Sammox for the origins

of life on planetary systems.

References

1. Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive earth's biogeochemical cycles. *Science* **320**, 1034–1039 (2008).

2. Sousa, F. L. et al. Early bioenergetic evolution. *Philos. Trans. R. Soc. B-Biol. Sci.* **368**, (2013).

3. Weiss, M. C. et al. The physiology and habitat of the last universal common ancestor. *Nat. Microbiol.* **1**, (2016).

4. Moore, E. K. et al. Metal availability and the expanding network of microbial metabolisms in the Archaean eon. *Nat. Geosci.* **10**, 629–636 (2017).

5. Ljungdah, L., Irion, E. & Wood, H. G. Total synthesis of acetate from CO2. I. Co-methylcobyric acid and CO-(methyl)-5-Methoxybenzimidazolylcobamide as intermediates with *Clostridium thermoaceticum*. *Biochemistry* **4**, 2771–2780 (1965).

6. Kelley, D. S. et al. A serpentinite-hosted ecosystem: the lost city hydrothermal field. *Science* **307**, 1428–1434 (2005).

7. Lane, N. & Martin, W. F. The origin of membrane bioenergetics. *Cell* **151**,

1406–1416 (2012).

8.   Wächtershäuser, G. Pyrite formation, the first energy source for life: a

hypothesis. *Syst. Appl. Microbiol.* **10**, 207–210, (1988).

9.   Wächtershäuser, G. Evolution of the first metabolic cycles. *PHAS J.* **1**,

200–204 (1990).

10. deDuve, C. Blueprint for a cell: the nature and origin of life. (Neil Patterson ,

1991).

11. Evans, M. C. W., Buchanan, B. B. & Arnon, D. I. A new ferredoxin-dependent

carbon reduction cycle in a photosynthetic bacterium. *Proc. Natl. Acad. Sci. U.*

S. A. **55**, 928–934 (1966).

12. Smith, E. & Morowitz, H. J. Universality in intermediary metabolism. *Proc.*

Natl. Acad. Sci. U. S. A. **101**, 13168–13173 (2004).

13. Muchowska, K. B. et al. Metals promote sequences of the reverse krebs cycle.

Nat. Ecol. Evol. **1**, 1716–1721 (2017).

14. Varma, S. J. et al. Native iron reduces CO_2 to intermediates and end-products

of the acetyl-CoA pathway. *Nat. Ecol. Evol.* **2**, 1019–1024 (2018).

15. Martin, W. & Russell, M. J. On the origin of biochemistry at an alkaline

hydrothermal vent. *Philos. Trans. R. Soc. B-Biol. Sci.* **362**, 1887–1925 (2007).

16. Orgel, L. E. The implausibility of metabolic cycles on the prebiotic earth.

PLoS. Biol. **6**, 5–13 (2008).

17. Braakman, R. & Smith, E. The emergence and early evolution of biological

carbon-fixation. *PLoS Comput. Biol.* **8**, (2012).

18. Camprubi, E. et al. Iron catalysis at the origin of life. *IUBMB Life* **69**, 373–381

(2017).

19. Goldford, J. E. et al. Remnants of an ancient metabolism without phosphate.

Cell **168**, 1126–1134 (2017).

20. Fdz-Polanco, F. et al. Combining the biological nitrogen and sulfur cycles in

anaerobic conditions. *Water Sci. Technol.* **44**, 77–84 (2001).

21. Amend, J. P. et al. Energetics of chemolithoautotrophy in the hydrothermal

system of Vulcano Island, southern Italy. *Geobiology* **1**, 37–58 (2003).

22. Philippot, P. et al. Early archaean microorganisms preferred elemental sulfur,

not sulfate. *Science* **317**, 1534–1537 (2007).

23. Brandes, J. A. et al. Abiotic nitrogen reduction on the early earth. *Nature* **395**,

365–367 (1998).

24. Smirnov, A. et al. Abiotic ammonium formation in the presence of Ni-Fe

metals and alloys and its implications for the hadean nitrogen cycle. *Geochem.*

Trans. **9**, (2008).

25. Sleep, N. H. The hadean-archaean environment. *Cold Spring Harbor Perspect.*

Biol. **2**, (2010).

26. Kuypers, M. et al. The microbial nitrogen-cycling network. *Nat. Rev.*

Microbiol. **16**, 263–276 (2018).

27. Eigen, M. Self-organization of matter and the evolution of biological

macromolecules. *Naturwissenschaften* **58**, 465–562 (1971).

28. Heinen, W. & Lauwers, A. M. Organic sulfur compounds resulting from the

interaction of iron sulfide, hydrogen sulfide and carbon dioxide in an

anaerobic aqueous environment. *Orig. Life Evol. Biosph.* **26**, 131–150 (1996).

29. Yamaguchi, A. et al. Electrochemical CO_2 reduction by Ni-containing iron

sulfides: how is CO_2 electrochemically reduced at bisulfide-bearing deep-sea

hydrothermal precipitates? *Electrochim. Acta* **141**, 311–318 (2014).

30. Martin, W. F., Sousa, F. L. & Lane, N. Evolution energy at life's origin.

Science **344**, 1092–1093 (2014).

31. Wächtershäuser, G. Before enzymes and templates-theory of surface

metabolism. *Microbiol. Rev.* **52**, 452–484 (1988).

32. Shen, Y. A., Buick, R. & Canfield, D. E. Isotopic evidence for microbial

sulphate reduction in the early Archaean era. *Nature* **410**, 77–81 (2001).

33. Mojzsis, S. J. et al.. Mass-independent isotope effects in Archean (2.5 to 3.8

Ga) sedimentary sulfides determined by ion microprobe analysis. *Geochim.*

Cosmochim. Acta **67**, 1635–1658 (2003).

34. Eck, R. V. & Dayhoff, M. O. Evolution of the structure of ferredoxin based on

living relics of primitive amino acid sequences. *Science* **152**, 363–366 (1966).

35. Herschy, B. et al. Archean phosphorus liberation induced by iron redox

geochemistry. *Nat. Commun.* **9**, (2018).

36. Jorgensen, B. B. A thiosulfate shunt in the sulfur cycle of marine-sediments.

Science **249**, (1990).

37. Clement, J. C. et al. Ammonium oxidation coupled to dissimilatory reduction

of iron under anaerobic conditions in wetland soils. *Soil Biol. Biochem.* **37**,

2323–2328 (2005).

38.   Yang, W. H., Weber, K. A. & Silver, W. L. Nitrogen loss from soil through

437   anaerobic ammonium oxidation coupled to iron reduction. *Nat. Geosci.* **5**,

438   538–541 (2012).

439   39.   Bao, P. & Li, G. X. Sulfur-driven iron reduction coupled to anaerobic

440   ammonium oxidation. *Environ. Sci. Technol.* **51**, 6691–6698 (2017).

441   40.   Postgate, J. R. The sulphate-reducing bacteria (Cambridge University, 1984).

442   41.   Wacey, D. et al. Microfossils of sulphur-metabolizing cells in

443   3.4-billion-year-old rocks of Western Australia. *Nat. Geosci.* **4**, 698–702

444   (2011).

445   42.   Crane, B. R. & Getzoff, E. D. The relationship between structure and function

446   for the sulfite reductases. *Curr. Opin. Struct. Biol.* **6**, 744–756 (1996).

447   43.   Bonomi, F., Werth, M. T. & Kurtz, D. M. Assembly of [FenSn(SR)]2-(n=2, 4)

448   in aqueous media from iron salts, thiols and sulfur, sulfide, thiosulfide plus

449   rhodonase. *Inorg. Chem.* **24**, 4331–4335 (1985).

450   44.   Lefaou, A. et al. Thiosulfate, polythionates and elemental sulfur assimilation

451   and reduction in the bacterial world. *FEMS Microbiol. Lett.* **75**, 351–382

452   (1990).

453   45.   Bordo, D. & Bork, P. The rhodanese/Cdc25 phosphatase superfamily-

454   sequence-structure-function relations. *EMBO Rep.* **3**, 741–746 (2002).

455   46.   Zhuang, W. Q. et al. Incomplete wood-ljungdahl pathway facilitates

456   one-carbon metabolism in organohalide-respiring *Dehalococcoides mccartyi*.

457   *Proc. Natl. Acad. Sci. U. S. A.* **111**, 6419–6424 (2014).

458 47. Mall, A. et al. Reversibility of citrate synthase allows autotrophic growth of a

459   thermophilic bacterium. *Science* **359**, 563–567 (2018).

460 48. Huber, C. & Wächtershäuser, G. Activated acetic acid by carbon fixation on

461   (Fe,Ni)S under primordial conditions. *Science* **276**, 245–247 (1997).

462 49. Lovelock, J. E. The Ages of Gaia (Oxford University, 1995).

464 **Acknowledgements** This research was financially supported by the National Natural

465 Science Foundation of China (General Program Nos. 41571240 and 41571130063).

467 **Competing interests** The authors declare no competing interests.

469 **Methods**

470 **Enrichment and isolation of Sammox bacterium.**

471   Surface paddy soil mud (0–20 cm depth) in the upper section of a paddy field

472 was collected from Guangxi Dahuanjiang region (24°53'52" N, 108°17'43" E), in the

473 southwest of China. Soil materials were air-dried, ground with a mortar and crushed

474 to pass through a 2.0 mm sieve for character determination. Soil organic carbon

475 (19.81 g kg^{-1}) was determined by potassium dichromate oxidation titration, and soil

476 total Fe (31.9 g kg^{-1}) was determined by ICP-OES. Plant available sulfur (29.1 mg

477 kg^{-1}) was determined by turbidimetry. Ammonium was 120.5 mg kg^{-1}.

478   For enrichment, approximately 0.5 g of paddy soil was transferred to 100 ml of

double distilled water. After shaking, five milliliters of the suspension was inoculated

into 100 ml of anaerobic freshwater medium[39,50] with additional sulfur (thiosulfate or

sufate) (3.0 mM) and ammonium ($^{15}NH_4Cl$, 1.0 mM) and incubated statically for one

week. The basal medium contained (g l^{-1}): KH_2PO_4 0.2, $MgCl·6H_2O$ 0.4, $CaCl_2·2H_2O$

0.1, KCl 0.5. The trace elemental mixture contained (l^{-1}): double distilled water 987

ml, HCl (25%) 12.5 ml (100 mM), $FeCl_2·4H_2O$ 957 mg (7.5 mM), H_3BO_3 30 mg (0.5

mM), $MnCl_2·4H_2O$ 100 mg (0.5 mM), $CoCl_2·6H_2O$ 190 mg (0.8 mM), $NiCl·6H_2O$ 24

mg (0.1 mM), $CuCl_2·2H_2O$ 2 mg (0.01 mM), $ZnSO_4·7H_2O$ 144 mg (0.5 mM),

$Na_2MoO_4·2H_2O$ 36 mg (0.15 mM). Selenite-tungstate solution (l^{-1}): NaOH 0.4 g,

$Na_2SeO_3·5H_2O$ 6 mg, $Na_2WO_4·2H_2O$ 8 mg. Bicarbonate solution $NaHCO_3$ 84 g l^{-1} 30

ml. The above mentioned stock solutions or aliquots were aseptically added to the

basal medium (l^{-1}): trace element solution (1.0 ml), selenite-tungstate solution (0.1 ml),

bicarbonate solution (10.0 ml), biotin, 4-aminobenzoic acid, 10 mg l^{-1} pantothenate,

pyridoxamine, nicotinic acid and 20 mg l^{-1} thiamine, 5 ml l^{-1}; vitamin B_{12} solution

with 50 mg l^{-1}, 1.0 ml l^{-1}. Sodium sulfate plus thiosulfate solution (1.0 ml, 3.0 mM

final concentration) and ammonium (0.5 ml, 1.0 mM final concentration) were added

into medium if necessary. The pH was adjusted to 8.0.

A volume of 0.5 ml of a positive Sammox culture, in which sulfur reduction

coupled to anaerobic ammonium oxidation, was sub-inoculated onto Acidovorax

complete agar medium under 5% CO_2 atmosphere. Single colony was selected and

inoculated back into anaerobic freshwater medium. Positive Sammox samples were

continuously sub-inoculated using the Acidovorax complete agar medium. A single

Sammox bacterium (designated GX3-BWBA) was isolated after sub-cultivation for

five times, and was continuously cultivated for one year in anaerobic freshwater

medium. The purity of GX3-BWBA was confirmed by promoting the growth of

heterotrophic bacteria by addition of peptone and yeast extract to the defined

freshwater mineral medium (data not shown)[51]. Finally, the purity of GX3-BWBA

was confirmed by deep Illumina sequencing.

This study was performed using a series of experiments, for Sammox by

GX3-BWBA: (I) 3.0 mM thiosulfate + 1.0 mM $^{15}NH_4Cl$, (II) 3.0 mM thiosulfate + 1.0

mM $^{15}NH_4Cl$ + GX3-BWBA, (III) 3.0 mM sulfate + 1.0 mM $^{15}NH_4Cl$, (IV) 3.0 mM

sulfate + 1.0 mM $^{15}NH_4Cl$ + GX3-BWBA. Experiments started and were sampled at

the sampled at the 0, 3, 6, 9, 12, 15, 18, 21, 24, 27, 34 day. Experiments were

performed anaerobically in 100-ml serum bottles in the dark at 30 ^{o}C. Each batch of

experiments was established in triplicate, inoculated with 1.0 ml GX3-BWBA

exponential phase culture if necessary.

Genome sequencing and assembly

The genome of *R. thioammoniphilus* (strain GX3-BWBA) was sequenced using

an Illumina HiSeq 4000 system (Illumina, SanDiego, CA, USA) at the Beijing

Genomics Institute (Shenzhen, China). Genomic DNA was sheared randomly to

construct three read libraries with lengths of (150:150) by a Bioruptor ultrasonicator

(Diagenode, Denville, NJ, USA) and physico-chemical methods. The paired-end

fragments libraries were sequenced according to the Illumina HiSeq 4000 system's

protocol. Raw reads of low quality from paired-end sequencing (those with

consecutive bases covered by fewer than five reads) were discarded. The sequenced

reads were assembled using SOAPdenovo v1.05 software. This Whole Genome

Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession

QMKS00000000. The version described in this paper is version QMKS01000000.

Genome Component prediction

Gene prediction was performed on the *R. thioammoniphilus* genome assembly by

glimmer3 (http://www.cbcb.umd.edu/software/glimmer/) with Hidden Markov models.

tRNA, rRNA and sRNAs recognition made use of tRNAscan-SE[52], RNAmmer, and

the Rfam database. The tandem repeats annotation was obtained using the Tandem

Repeat Finder (http://tandem.bu.edu/trf/trf.html), and the minisatellite DNA and

microsatellite DNA selected based on the number and length of repeat units. The

Genomic Island Suite of Tools (GIST) used for genomic islands analysis

(http://www5.esu.edu/cpsc/bioinfo/software/GIST/) with Island Path-DIOMB,

SIGI-HMM, Island Picker method. Prophage regions were predicted using the PHAge

Search Tool (PHAST) web server (http://phast.wishartlab.com/) and CRISPR

identification using CRISPRFinder.

Protein sequences will be downloaded from NCBI, and get sequences in the

taxonomy of species classification information; The sequences using mafft software

for multiple sequence alignment. After comparing the sequences, using Fasttree

software construct phylogenetic tree (neighbor joining algorithm). Using R language

ggtree package for visualization mapping.

Gene annotation and protein classification

The best hit abstracted using Blast alignment tool for function annotation. Seven

databases which are KEGG (Kyoto Encyclopedia of Genes and Genomes), COG

(Clusters of Orthologous Groups), NR (Non-Redundant Protein Database databases),

Swiss-Prot[53], and GO (Gene Ontology), TrEMBL, EggNOG are used for general

function annotation. Representative metabolic gene clusters were displayed with

Easyfig[54].

General procedure for prebiotic Sammox-driven CO_2 fixation

Each 50 ml basal solution was transferred into 60 ml serum bottles and sealed

with butyl rubber stoppers and aluminium crimp caps. The solution was autoclaved

and cooled as room temperature after washed by He gas (purity=99.999%). Additional

trace elemental mixture solutions were filter sterilized or autoclaved individually.

Additional sulfur (thiosulfate or sufate) (3.0 mM) and/or ammonium (NH_4Cl, 1.0 mM)

was added into serum bottle. The mixed solution was termed as Sammox reaction

system.

The above mentioned stock solutions or aliquots were aseptically added to the

basal medium (l^{-1}): trace element solution (2.0 ml), selenite-tungstate solution (0.2 ml),

bicarbonate solution (10.0 ml). Sodium sulfate, thiosulfate (1.0 ml), ammonium

solution (0.5 ml), and fresh precipitated Fe-S mineral were added into medium if

necessary. Serum bottles were kept 100 ^{o}C in a water bath in the dark for 24 h, then

maintain at 70 ^{o}C in the dark for another 24 h, and sampling at 48 h to determinate

organic products.

This study was performed using a series of experiments: (I) 3.0 mM thiosulfate +

1.0 mM NH_4Cl + 20 mM HCO_3^-, (II) 3.0 mM thiosulfate + 1.0 mM NH_4Cl + 20 mM

HCO_3^- + 1.0 mM Fe-S, (III) 3.0 mM thiosulfate + 20 mM HCO_3^-, (IV) 3.0 mM sulfate

+ 1.0 mM NH_4Cl + 20 mM HCO_3^-, (IV) 3.0 mM sulfate + 1.0 mM NH_4Cl + 20 mM

HCO_3^- + 1.0 mM Fe-S, (V) 3.0 mM sulfate + 20 mM HCO_3^-, (VI) 1.0 mM NH_4Cl +

20 mM HCO_3^-.

General procedure for thiol/thioester promoted Sammox-driven CO_2 fixation.

A solution of methanethiol was added to Sammox reaction system. The reaction

systems were heated at 100 ^oC in a water bath in the dark for 24 h, then maintain at 70

^oC in the dark for another 24 h, and removed from the water bath and allowed to cool

to room temperature before derivatization and gas chromatography–mass

spectrometry analysis.

This study was performed using a series of experiments: (I) 3.0 mM thiosulfate +

1.0 mM NH_4Cl + 20 mM HCO_3^- + 0.8 mM methanethiol, (II) 3.0 mM thiosulfate +

1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM Fe-S + 0.8 mM methanethiol, (III) 3.0

mM sulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 0.8 mM methanethiol, (IV) 3.0 mM

sulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM Fe-S + 0.8 mM methanethiol.

General procedure for verification of Sammox-driven combination of reductive

acetyl-CoA pathway and incomplete rTCA cycle.

We design this experimental set to verify if Sammox-driven reductive

acetyl-CoA pathway could go into Sammox-driven incomplete rTCA cycle. The

general procedure is the same as above. Pyruvate (1.0 mM, final concentration) was

added into Sammox reaction system as substrate. Serum bottles were heated at 100 ^oC

in a water bath in the dark for 24 h, then maintain at 70 $^{\circ}$C in the dark for another 24 h,

and allowed to cool to room temperature before derivatization and gas

chromatography–mass spectrometry analysis.

This study was performed using a series of experiments: (I) 3.0 mM thiosulfate +

1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM pyruvate + 1.0 mM Fe-S + methanethiol,

(II) 3.0 mM thiosulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM Fe-S + 1.0 mM

pyruvate, (III) 3.0 mM thiosulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM

pyruvate, (IV) 3.0 mM thiosulfate + 20 mM HCO_3^- + 1.0 mM pyruvate, (V) 3.0 mM

sulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM Fe-S + 1.0 mM pyruvate +

methanethiol, (VI) 3.0 mM sulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM Fe-S

+ 1.0 mM pyruvate, (VII)3.0 mM sulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0

mM pyruvate, (VIII) 3.0 mM sulfate + 20 mM HCO_3^- + 1.0 mM pyruvate, (IX) 1.0

mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM pyruvate.

Chemicals

All reagents and solvents were purchased from commercial suppliers and used

without further purification unless otherwise noted.

Sampling analytical methods

(i) **Enumeration of microbe viable counts.** The spread-agar-plate method was used

for enumeration of bacterial counts in GX3-BWBA culture[55]. The Acidovorax

complete agar medium, pH 7.0, contained 5.0 g l^{-1} peptone (Difco), 3.0 g l^{-1} beef

extract and 15 g l^{-1} agar (Difco). A 50 µl sample was taken from GX3-BWBA culture

at the 0, 3, 6, 9, 12, 15, 18, 21, 24, 27 and 34 days and inoculated onto plates for cell

counting. Visible colonies produced on the agar plate were counted at 48 h. Bacterial

numbers were expressed as colony-forming units (CFU) per milliliter of culture[55].

(ii) **Electron microscopy.** Morphology of GX3-BWBA was investigated by

transmission electron microscope (TEM) (TEM, HITACHI H-7500). Cells from the

culture were collected by centrifugation, washed and diluted with phosphate buffer

(pH 7.5), and dropped on to specimens. The material was examined after dry by

airing.

(iii) **Sulfide.** The analysis of sulfide in filtered samples was performed

photometrically by the methylene blue method[56]. Absorption at 660 nm was measured

(Infinite 200PRO; TECAN).

(iV) **Sulfate, sulfite, nitrite, acetate, formate and ammonium.** To determine sulfate,

sulfite, nitrite, acetate and formate, 0.5 ml of the sample was filtered (0.22 μm) to

remove particulates that could interfere with ion chromatography. The ion

chromatography system consisted of an ICS-5000$^+$ SP pump (Thermo Fisher

Scientific Inc. Sunnyvale, CA, USA), a column oven ICS-5000$^+$ DC, an

electrochemical detector DC-5. The ion chromatography column system used a

Dionex Ionpac AS11-HC column. The operating condition was with an eluent of 30

mM KOH at a flow rate of 1.0 ml min^{-1}. For determination of ammonium, the ion

chromatography column system used a Dionex Ionpac CS12A column.

(V) **Thiosulfate.** Thiosulfate was determined by an Agilent 1260 infinity HPLC

system, equipped with a quaternary pump (Agilent, USA). Thiosulfate was separated

by a Zorbax SB-C18 column (150×4.6 mm, 5 μm), and detected by using of

DAD detector at 215 nm. All analyses were performed at 40 °C with a flow rate of 1

ml min^{-1}. Na_2HPO_4 was used as solvent. The pH of the solvent was adjusted with 1.0

M HCl. Samples were filtered through 0.45 μm Cosmonice Filters (Millipore, Tokyo,

Japan) and immediately injected into the HPLC system[57].

(Vi) Gas Analysis. At each sampling time point, each serum bottle was shaken

vigorously to equilibrate the N_2 between dissolved and gas phases. Dinitrogen

sampling and determination was performed according to a literature procedure[39].

(Vii) Derivatization procedure and product identification

For optimal GC resolution, the carboxylic acids were converted to ethyl esters

using a mixture of ethanol/ethyl chloroformate (EtOH/ECF). Derivatization of

carboxylic acids to esters was performed according to a literature procedure[13].

Reaction products derivatized to ethyl esters of carboxylic acids were identified by

comparing the mass spectra and retention times against analogously derivatized

authentic samples. ECF derivatization was preferred for small molecule substrates

(pyruvate, lactate, malate, fumarate, succinate, α-ketoglutarate, amino acid). The

carboxylic acids might also convert to methyl esters using a mixture of

methanol/methyl chloroformate (MeOH/MCF), following the same procedure to ECF

derivatization to detect *cis*-aconitate, tricarballylate, isocitrate and citrate.

(Viii) Gas chromatography–mass spectrometry (GC–MS) analysis for rTCA

metabolites determination.

GC–MS analysis was performed on a GC System 7890B connected to a MSD

block 5977A, using Agilent High Resolution Gas Chromatography Column: PN

19091S–433, HP–5MS, 28 m×0.25 mm, 0.25 Micron, SN USN 462366H. Samples

were prepared in ethyl acetate (200 μl sample volume). The analysis was carried out

on a splitless 1 μl injection volume with an injection port temperature 250 °C.

Column oven temperature program was as follows: 60 °C for 1.0 min, ramped at

30 °C min^{-1} to 310 °C with 3.0 min hold, with a total running time of 12.33 min. The

mass spectrometer was turned on after 3 min and was operated at the electron

ionization mode with quadrupole temperature of 150 °C. Data acquisition was

performed in the full-scan mode (50-500). Hydrogen (99.999 % purity) was used as

carrier gas at a constant flow rate of 1.5 ml min^{-1}.

50.   Strous, M. et al. Deciphering the evolution and metabolism of an anammox

bacterium from a community genome. *Nature* **440**, 790–794 (2006).

51.   Konneke, M. et al. Isolation of an autotrophic ammonia oxidizing marine

archaeon. *Nature* **437**, 543–546 (2005).

52.   Lowe, T. M. & Eddy, S. R. tRNAscan-SE: a program for improved detection

of transfer RNA genes in genomic sequence. *Nucl. Acids Res.* **25** (5),

0955–0964 (1997).

53.   Torto-Alalibo, T., Collmer, C. W. & Gwinn-Giglio, M. The plant-associated

microbe gene ontology (PAMGO) consortium: community development of

new gene ontology terms describing biological processes involved in

microbe-host interactions. *BMC microbiol.* **9** (suppl 1), (2009).

54.   Sullivan, M. J., Nicola K. P. & Scott A. B. Easyfig: a genome comparison

visualizer. *Bioinformatics* **27** (7), 1009–1010 (2011).

55.   Kataoka, N. et al. Enrichment culture and isolation of slow growing bacteria.

Appl. Microbiol. *Biotechnol.* **45**, 771–777 (1996).

56.   Cline, J. D. Spectrophotometric determination of hydrogen sulfide in natural

waters. imnol. *Oceanogr*. **14** (3), 454–458 (1969).

57.   Kondo, R. et al. Determination of thiosulfate in a meromictic lake. *Fisheries*

Sci. **66**, 1076–1081 (2000).

Legends of figures

Figure. 1 Sammox-driven proto-anabolic networks under hydrothermal conditions.

Treatments were as follows (**a**, thiosulfate; **b**, sulfate): (i) sulfurous species,

ammonium, Fe-S minerals, and thiols; (ii) sulfurous species, ammonium, and Fe-S

minerals; (iii) sulfurous species, ammonium, and thiols; (iv) sulfurous species and

ammonium; (v) ammonium; (vi) sulfurous species (see experimental details). The bar

chart shows the yield of formate, acetate, succinate and α-ketoglutarate in each

treatment group after 48 h. Error bars represent standard deviations of three replicates.

Figure. 2 Sammox-driven incomplete rTCA with pyruvate amendment under

hydrothermal conditions. Treatments were as follows (**a**, thiosulfate; **b**, sulfate): (i)

sulfurous species, ammonium, Fe-S minerals, and thiols; (ii) sulfurous species,

ammonium, and Fe-S minerals; (iii) sulfurous species, ammonium, and thiols; (iv)

sulfurous species and ammonium; (v) ammonium; (vi) sulfurous species (see

experimental details). The bar chart shows the concentration of pyruvate, succinate

and α-ketoglutarate in each treatment group after 48 h. Error bars represent standard

deviations of three replicates.

Figure. 3 Hypothetical prebiotic reductive acetyl-CoA pathway powered by Sammox,

a. Hypothetical proto-anabolic networks, a combination of acetyl-CoA pathway

together with incomplete rTCA cycle powered by Sammox, and the role of its

intermediates as universal biosynthetic precursors. Bar represents the final step of

incomplete rTCA cycle, **b**. Conceptual model of Sammox-driven the origin of not

free-living organic precursor on the surface of iron-sulfur minerals in Hadean alkaline

hydrothermal systems. Cubes represent iron-sulfur minerals/metals alloy. Globes

represent the not free-living organic precursor, ligand sphere, **c**.

Figure. 4 Characters of *R. thioammoniphilus* strain GX3-BWBA. **a**, TEM image of

cells. Scale bar represents 0.2 µm. **b**, Growth of GX3-BWBA under combined

condition of ammonium and sulfurous species, with bicarbonate as sole carbon

sources. Phylogenetic tree constructed by the maximum likelihood method, using

1478 nucleotides from 16S rDNA sequences, showing the position of strain

GX3-BWBA in relation to members of the genus Ralstonia. The bar represents 0.1

changes per nucleotide position, **c**. Production rate of isotopically labeled dinitrogen

for thiosulfate and sulfate-fueled Sammox by *R. thioammoniphilus* strain

GX3-BWBA (**d**, **e**).

Figure. 5 Thiosulfate- and sulfate-fueled Sammox by *R. thioammoniphilus* strain

GX3-BWBA. **a**, thiosulfate and reduction products. **b**, ammonium oxidation and

products in thiosulfate-fueled Sammox. **c,** sulfate and reduction products (sulfide and

sulfite as products of sulfate in control group are below detection limit). **d**,

ammonium oxidation and products in sulfate-fueled Sammox. Error bars represent

standard deviations of three biological replicates.

Figure. 6 Representative metabolic gene clusters from *R. thioammoniphilus* strain

GX3-BWBA (**a**). The standard configuration of metabolic gene clusters (MGCs) in

primary metabolism located in two scaffolds. MGCs contain enzymes of ▮ CO_2

fixation genes, including 31 rTCA cycle, and 30 reductive acetyl-CoA pathway genes,

■ 6 dissimilatory sulfur metabolic genes, and ■ 5 nitrification genes. Gene size

and spacing are not to scale. Each gene was not condensed lie in genome. Protein

phylogenetic tree derived from 26443 amino acid sequences of sulfite/nitrite reductase

(**b**). Scale bars represent estimated sequence divergence or amino acid changes. ●

Archaea, ▲ Bacteria, ■ Eukaryota.

a

b

acetyl-CoA pathway

rTCA cycle

Aconitate

Isocitrate

Oxalosuccinate

Citrate

Lipids

CO_2

AcCoA

Me

Sammox

$2e^- + 2H^+$

CO_2

Pyruvate

Me

Sugars
Alanine

Sammox

Oxaloacetate

CO_2

Pyrimidines
Aspartate
Other amino acids

Sammox

$2e^- + 2H^+$

Malate

Fumarate

Succinate

α-ketoglutarate

$2e^- + 2H^+$ Sammox

CO_2

Glutamate
Other amino acids

Pyrroles

$2e^- + 2H^+$

Sammox

Sammox

c

$S_xO_y^{n-}$

NH_4^+

a

b