Sulfur reduction coupling with anaerobic ammonium oxidation drives proto-anabolic

networks

Peng Bao[1, 2, *], Guo-Xiang Li[1, 2, 3], Hu Li[1]

[1]Key Lab of Urban Environment and Health, Institute of Urban Environment, Chinese

Academy of Sciences, Xiamen 361021, P. R. China

[2]Ningbo Urban Environment Observation and Station, Chinese Academy of Sciences,

Ningbo 315800, P. R. China

[3]University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Address correspondence to:

Dr. Peng Bao

Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, P. R.

China

E-mail: pbao@iue.ac.cn; neilbao17@163.com

Abstract

The geochemical energy that drives the transition from non-life to life is as yet unknown. Here we show that thiosulfate/sulfate reduction coupling with anaerobic ammonium oxidation (Sammox), could provide the primordial redox equivalents and energy for prebiotic proto-anabolic networks, the reductive acetyl-CoA pathway combined with incomplete reductive tricarboxylic acid (rTCA) cycle, reductive amination and pyrrole synthesis. Fe-S mineral catalysis and thiols/thioesters as energy couplers enhance the efficiency of prebiotic proto-anabolic networks in thiosulfate-fueled Sammox reaction systems under hydrothermal conditions. Results implied that thiosulfate-fueled Sammox should be the primordial reaction driving the construction of proto-anabolic networks rather than sulfate-fueled Sammox, as it could be catalyzed, and also sulfate would have been severely limiting in ancient oceans. To confirm our findings, we isolated and identified a mixtrophic Sammox bacterium, *Ralstonia thioammoniphilus* GX3-BWBA, which prefer thiosulfate to sulfate. Genomic analysis of *R. thioammoniphilus* GX3-BWBA implied that this ancient metabolism in modern microbes should contain two stages according to ammonium transformation, —oxidation of ammonium to nitrite and denitrification. The incomplete rTCA cycle and reductive acetyl-CoA pathway were all identified in *R. thioammoniphilus* GX3-BWBA metabolic networks, which were responsible for chemolithotrophic metabolism. We inferred that Sammox drove the coupling of the biochemical transformation of C, H, O, N, S, and/or Fe simultaneously in Hadean alkaline hydrothermal systems, thereby permitting the emergence of life. The results

bridged the gap in the transition from geochemistry to biochemistry.

The transition from non-life to life occurred in the context of geochemical energy

derived from element-coupled transformation[1-3]. The chemistry of life is based on

redox reactions, that is, successive transfers of electrons and protons from the six

major elements, i.e., C, H, O, N, S, and P[1,4]. The H_2/CO_2 redox couple, which can

occur in submarine hydrothermal vents, has been proposed to drive the reductive

acetyl-CoA pathway, an ancient metabolic route[3,5-7]. Nevertheless, the primordial

energy source of the H_2/CO_2 redox couple suffers from the difficulty that the

exergonic reaction competes with the endergonic reaction for available H_2[8]. The

subsequent surface metabolism and thioester world theories still could not answer the

question, how the required reduced carbon compounds have been synthesized[9,10].

Despite this limitation, these theories have emphasized the important roles of

thioesters and Fe-S mineral catalysis for driving the primordial rTCA cycle, a central

anabolic biochemical pathway whose origins are proposed to trace back to

geochemistry[9,11,12].

Native iron/metals were recently shown to promote the reductive acetyl-CoA

pathway and rTCA cycle and strongly support the feasibility of these two primordial

synthetic pathways[13,14], although they were generally considered to be rare near the

Earth's surface[13] and cannot support the long elemental transition from geochemistry

to biochemistry. Proto-anabolic networks consisting of the reductive acetyl-CoA

pathway together with the complete/incomplete rTCA cycle as primordial synthetic

pathways are therefore more logical[3,15-18]. However, the initial driving force for the

rise of proto-anabolic networks is still unclear. Moreover, the roles of N and S

geochemical transformation in the origin of life have been largely ignored because

computational analysis has implied that N and S are essential for thermodynamically

feasible phosphate-independent metabolism before the rise of last universal common

ancestor (LUCA)[19]. The proto-anabolic networks might be driven by C, H, O, N, S,

and/or Fe coupling transformation at their earliest stage of the transition from

geochemistry to biochemistry. The co-evolution of the metabolism of those elements

may provide strong explanatory power for the origin of life and explain why the

structure of metabolic networks is as it is. We therefore speculate that

thermodynamically feasible sulfurous species reduction coupling with anaerobic

ammonium oxidation reaction[20,21,22,23,24] (Eqs. 1, 2; pH=8.0), with/without the

catalysis of Fe-S minerals and thioesters, may have been the primordial power force

for the rise of proto-anabolic networks.

$$8NH_4^+ + 2SO_4^{2-} + 2HCO_3^- \rightarrow 4N_2 + 2HS^- + CH_3COO^- + 12H_2O + 5H^+ \quad \Delta G_r = -20.0 \text{ kJ mol}^{-1} \quad (1)$$

$$4NH_4^+ + S_2O_3^{2-} + 2HCO_3^- \rightarrow 2N_2 + 2HS^- + 2HCOO^- + 5H_2O + 4H^+ \quad \Delta G_r = -13.4 \text{ kJ mol}^{-1} \quad (2)$$

On early Earth, elemental sulfur, sulfite, and thiosulfate were produced

abundantly from volcanic and hydrothermal SO_2 or from H_2S oxidation by iron oxides

in sulfide-rich hydrothermal fluid[2,15,25]. In simulated hydrothermal systems with

conditions of 300 °C and high pressure, nitrite was readily converted to NH_3[26], and

Ni-Fe metals and alloys were also effective catalysts of N_2 reduction to NH_3 in

Hadean hydrothermal systems[27]. There would have been much higher CO_2

concentrations in the oceans on early Earth because there was perhaps up to 1000

times more CO_2 in the atmosphere than that today[28].

Hence, when sulfurous species and NH_3 in Hadean hydrothermal systems made

contact with CO_2, there were spontaneous electron and proton transfers for energy

generation and organic molecule synthesis via sulfurous species reduction coupling

with ammonium oxidation. As sulfate would have been severely limiting in ancient

oceans[2,29], we termed this process Sammox, and a Sammox microbe is more likely

use thiosulfate, elemental sulfur and sulfite as electron acceptor besides sulfate,

distinguishing it from previous studies of the coupling elemental biogeochemical

cycles[1,30]. A prebiotic reaction should occur with the ability to branch out into S and

N biochemistry, which could contribute to the autocatalysis and evolution of

primordial metabolic networks. Thus, Sammox may facilitate the synthesis of

thioesters and amino acids, which are essential for the self-amplification of

phosphate-independent metabolic networks before the rise of LUCA. We expect that

prebiotic chemical evidence of Sammox-powered CO_2 fixation, thioesters, amino acid

and co-factors synthesis, combined with genetic analysis of a representative Sammox

microbe, will provide profound insights into the earliest origins of life and fill in the

missing link of the emergence of biochemistry from geochemistry.

Results and discussion

Sammox drives the combination of abiotic reductive acetyl-CoA pathway and

incomplete rTCA cycle

First, we aimed to verify the feasibility of Sammox-driven abiotic reductive

acetyl-CoA pathway, rTCA cycle, and off-cycle reactions (reductive amination and

co-factors synthesis) and to determine how methanethiol and Fe-S mineral catalysis

enhanced the reactions in both thiosulfate- and sulfate-fueled Sammox systems.

Formate and acetate were the products of Sammox-driven abiotic reductive

acetyl-CoA pathway in both thiosulfate- and sulfate-fueled Sammox systems with

bicarbonate as the carbon source (Fig. 1). We have not determined methanol but

qualitatively identified methyl acetate as a product in both thiosulfate- and

sulfate-fueled Sammox systems (Extended Data Fig. 1), implying that methanol

should be an intermediate of Sammox-driven abiotic reductive acetyl-CoA pathway.

More importantly, this result confirmed the capacity for Sammox-powered ester bond

formation, which was critical for the synthesis of complex organic molecules, such as

lipids and RNA[31].

Figure 1. Products of simulated prebiotic Sammox-driven proto-anabolic

networks with bicarbonate as the sole carbon source under hydrothermal

conditions. Treatments were as follows (a, thiosulfate; b, sulfate), from left to

right: (i) sulfurous species, ammonium, Fe-S minerals, and methanethiol; (ii)

sulfurous species, ammonium, and Fe-S minerals; (iii) sulfurous species,

ammonium, and methanethiol; (iv) sulfurous species and ammonium; (v)

ammonium; and (vi) sulfurous species. The bar chart shows the yields of formate,

acetate, succinate, and α-ketoglutarate in each treatment group. Error bars

represent standard deviations of three replicates.

We haven't detected pyruvate as the end-product of Sammox-driven abiotic

reductive acetyl-CoA pathway. When pyruvate was added into Sammox systems, it

quickly entered the incomplete rTCA cycle and was consumed as a reaction substrate

in both thiosulfate- and sulfate-fueled Sammox systems (Fig. 2, Fig. 3a, b). We

quantitatively identified succinate and α-ketoglutarate (Figs. 1, 2, Extended Data Fig.

2), and qualitatively identified polypyrrole (Extended Data Fig. 3) and glutamate

(Extended Data Fig. 4) as products of Sammox-driven abiotic incomplete rTCA cycle.

To further prove the feasibility of Sammox-driven abiotic incomplete rTCA cycle, we

adopted oxaloacetate, malate, fumarate and succinate, respectively, as substrates in

both thiosulfate- and sulfate-fueled Sammox systems, and all eventually produced

α-ketoglutarate and glutamate (Extended Data Fig. 5).

It is worth noting that the concentrations of α-ketoglutarate in N/S/FeS and N/S

groups were lower than the detection limit in thiosulfate-fueled Sammox systems with

bicarbonate as carbon source shown in figure 1 and 2. This was due to the reductive

amination of α-ketoglutarate to glutamate (Fig. 3b, Extended Data Fig 4). The

generation of polypyrrole implied that there should be pyrrole derived from succinate.

This might provide the possibility of biocatalysis emergence, as pyrrole was the

co-factors of enzyme. The above results implied that Sammox-driven abiotic

reductive acetyl-CoA pathway and abiotic incomplete rTCA cycle were combined and

tended to accumulate C_4 and C_5 products. We haven't detected other intermediates of

the incomplete rTCA cycle, potentially because of their low productivity. There were

no reductions of pyruvate to lactate and α-ketoglutarate to α-hydroxyglutarate in both

thiosulfate- and sulfate-fueled Sammox systems, which would disrupt the incomplete

rTCA cycle[16], therefore allowed Sammox-driven proto-anabolic networks to produce

major organic macromolecular products effectively.

Figure 2. Products of simulated prebiotic Sammox-driven incomplete rTCA cycle

with bicarbonate and pyruvate as carbon sources under hydrothermal

conditions. Treatments were as follows (a, thiosulfate; b, sulfate), from left to

right: (i) sulfurous species, ammonium, Fe-S minerals, and methanethiol; (ii)

sulfurous species, ammonium, and Fe-S minerals; (iii) sulfurous species,

ammonium, and methanethiol; (iv) sulfurous species and ammonium; (v)

ammonium; and (vi) sulfurous species. The bar chart shows the concentrations

of pyruvate, succinate, and α-ketoglutarate in each treatment group. Error bars

represent standard deviations of three replicates.

Figure. 3 Hypothetical prebiotic Sammox-driven reductive acetyl-CoA pathway

(a). Hypothetical proto-anabolic networks, a combination of acetyl-CoA pathway

together with incomplete rTCA cycle driven by Sammox, and the role of its

intermediates as universal biosynthetic precursors. Bar represents the final step

of incomplete rTCA cycle (b). Conceptual model of Sammox-driven the origin of

not free-living organic precursor on the surface of Fe-S minerals in Hadean

alkaline hydrothermal systems. Cubes represent Fe-S minerals/metals alloy.

Globes represent the not free-living organic precursor, ligand sphere (c).

Sulfate-fueled Sammox was more effective than thiosulfate-fueled Sammox

(Figs. 1, 2), due to the higher energy yield of the former reaction (Eqs. 1, 2). Both

Fe-S minerals and methanethiol could enhance the efficiency of the thiosulfate-fueled

Sammox reaction but not sulfate-fueled Sammox reaction (Fig 1), logically implying

that the thiosulfate-fueled Sammox reaction should be the primordial reaction driving

the construction of proto-anabolic networks, as it could be catalyzed. Note that

methanethiol was more effective than Fe-S minerals for promoting the

thiosulfate-fueled Sammox reaction (Fig. 1a). It incorporates into C, H, O and N

biochemical transformation via thiol-thioester exchange[9], transforming itself and

other organic reaction products into new organic products[10], therefore resulted in the

coupling biochemical transformation of C, H, O, N, and S and the expansion of

primordial metabolic networks. We deduced that when reaction conditions were met

in Hadean hydrothermal systems, Sammox-generated proton gradient could

modulate reduction potential similar to applying a voltage, leading to electrocatalytic

CO_2 reduction with/without catalysis of Fe-S minerals and thioesters[32,33]. Thus,

almost all kinds of the essential metabolic precursors for biosynthesis could be

provided in one geological setting. The organics formed within the vent pores in

Hadean hydrothermal systems, where they should concentrate via processes, such as

thermophoresis, and potentially form structures, such as lipid membranes lining

hydrophobic walls[7,15,34]. Thereby, Sammox gave rise to the origin of not free-living

organic precursor on the surface of Fe-S minerals[35] (Fig. 3c).

As one of the products of Sammox, sulfide began to increase in the Hadean

oceans, facilitating thiols synthesis on Fe-S mineral surface[32] and leading to de

Duve's thioester world and the boom of S biochemistry. As a result, the S isotope

geochemical evidence can be traced back to 3.8 Ga and has been well

preserved[4,25,36,37]. Biogenic sulfide reacted with soluble Fe^{2+} and/or Ni^{2+}/Co^{2+}/Se to

maintain a continuous self-supply of freshly precipitated FeS and/or metals alloys and

even subsequent enzyme and redox protein active centers, such as ferredoxin[38]. This

may represent a route for S and Fe assimilation, which would facilitate escape of the

not free-living organic precursor from the surface of Fe-S minerals. As catalytic

properties improved, the yield of proto-anabolic networks increased, ultimately

extending the pathway through to C_6 tricarboxylic acids. Sulfur biogeochemistry

prompted iron redox geochemistry, therefore, ferrous iron was released due to the

abiotic reduction of iron oxyhydroxide by biogenic sulfide, thus resulting in phosphite

liberation from ferruginous sediments[39]. In this study, we will not discuss how P

incorporated into primordial metabolic networks, but infer that Sammox-driven

proto-anabolic networks could facilitate self-evolution through optimizing the

ambient environment to make it sufficiently stable and habitable for life.

Discovery of Sammox microbe

To prove the existence of Sammox process in modern microbes, we successfully isolated a pure Sammox culture (designated GX3-BWBA) that could conserve energy from the reduction of sulfurous species (thiosulfate, sulfate, sulfite, and elemental sulfur) coupled to anaerobic ammonium oxidation. Strain GX3-BWBA appeared as bean-shaped cells with a diameter of 0.6 μm and a length of 1.0 μm and harbored a single flagella with a bacterial capsule (Fig. 4a, b, Extended Data Fig. 6). Comparative sequence analysis of 16S rRNA genes revealed a 99% query cover and 99% sequence identity between strain GX3-BWBA and a series of uncultured *Ralstonia* bacterial clones (Fig. 4c). Thus, strain GX3-BWBA was phylogenetically clustered within the *Ralstonia* genus (Fig. 4c). GX3-BWBA grew to a maximal density of 1.6×10^5 CFU ml^{-1} in anaerobic freshwater mineral medium containing bicarbonate, ammonium, and thiosulfate/sulfate as the sole carbon and energy sources, suggesting chemoautotrophy (Fig. 4b). Approximately 8 μM ammonium was sufficient to support the chemoautotrophic growth of strain GX3-BWBA, demonstrating high affinity for ammonium (data not shown). Thiosulfate was the preferred electron acceptor over sulfate by GX3-BWBA (Fig. 4b). There was no significant growth of GX3-BWBA under conditions without ammonium or sulfurous species (Fig. 4b); thus, ammonium and sulfurous species were essential factors for GX3-BWBA chemoautotrophic growth. We provisionally classified this Sammox bacterium as "*Ralstonia thioammoniphilus*" (thi.o.am.mo.ni′ phi.lus. Gr. n. *thion* sulfur; L. neut. n. *sal ammoniacum* salt of Ammon (NH_4Cl); Gr. adj. *phylos* loving;

   N.L. neut. adj. *thioammoniphilus* sulfur-ammonium-loving).

Figure. 4 Characters of *R. thioammoniphilus* GX3-BWBA. TEM image of cells (a). Scale bar represents 0.2 μm. Growth of GX3-BWBA under combined condition of ammonium and sulfurous species, with bicarbonate as sole carbon sources (b). Production rate of isotopically labeled dinitrogen for thiosulfate and sulfate-fueled Sammox by GX3-BWBA (c, d). Phylogenetic tree constructed by the maximum likelihood method, using 1478 nucleotides from 16S rDNA sequences, showing the position of strain GX3-BWBA in relation to members of the genus Ralstonia. The bar represents 0.1 changes per nucleotide position (e).

As a probable ancient metabolism, Sammox microbe preferred other sulfurous species, except for sulfate, potentially because elemental sulfur, thiosulfate, and sulfite were produced abundantly from ancient volcanic and hydrothermal SO_2 or from sulfide oxidation by iron oxides in sulfide-rich hydrothermal fluid[2,15,25]. Thiosulfate constituted 68–78% of the immediate HS^- oxidation products and was involved in a dynamic HS^--$S_2O_3^{2-}$ cycle in anoxic marine and freshwater sediments[40]. In contrast, sulfate would have had only very limited, localized significance[2].

We further tested thiosulfate- and sulfate-fueled Sammox metabolism using *R. thioammoniphilus*, and the end product N_2 generation rate was measured (Fig. 4c, d). The generation rate of N_2 in the thiosulfate treatment group was slightly higher than that in the sulfate treatment group (Fig. 4c, d). The production rate of $^{29}N_2$ was about 2.5 nmol ml^{-1} day^{-1}, which was significantly higher than $^{30}N_2$ (approximately 2.5 pmol ml^{-1} day^{-1}; Fig. 4c, d). Figure 5 shows the coupled dynamic transformation of

sulfurous species and ammonium during Sammox process. In thiosulfate-fueled

Sammox, sulfide was the product of the thiosulfate reduction, whereas sulfite was an

intermediate (Fig. 5a). Ammonium oxidation accompanied by thiosulfate reduction

was significant compared with that in the control group. As intermediate of

ammonium oxidation, nitrite significantly increased on day 27 (Fig. 5b). Based on the

low concentration, we did not determine the dynamic generation of dinitrogen to

avoid leakage. There was weak but significant sulfate reduction in the sulfate-fueled

Sammox group, accompanied by sulfide generation (Fig. 5c). The concentration of

nitrite also showed a slight increase during the ammonium oxidation process (Fig. 5d).

The concentrations of sulfurous species and ammonium showed no significant

changes in the control group (Fig. 5). The assumed net thiosulfate-/sulfate-fueled

Sammox reactions are shown in Eqs. 1 and 2. The generation of precursors of CO_2

fixation was thermodynamically favorable by Sammox.

Figure. 5 Thiosulfate- and sulfate-fueled Sammox by *R. thioammoniphilus*

GX3-BWBA. (a) thiosulfate and reduction products. (b) ammonium and its

oxidation products in thiosulfate-fueled Sammox. (c) sulfate and reduction

products (sulfide and sulfite as products of sulfate in control group are below

detection limit). (d) ammonium and its oxidation products in sulfate-fueled

Sammox. Error bars represent standard deviations of three biological replicates.

Notably, quite a few iron oxides could be abiotically reduced by sulfide

generated from Sammox in an iron-rich environment, then an illusion of ferric iron

reduction coupled to anaerobic ammonium oxidation (Feammox) arises[41-43]. Indeed, it

is still unclear to what extent Sammox overlaps with Feammox; however, we can

infer that Sammox may be widely distributed in different anoxic environments based

on the ubiquity of *Ralstonia* spp. and its high affinity for ammonium. Moreover, as an

ancient type of metabolism, there may be a variety of microbes that still preserved this

living strategy.

Genomic and N, S, and C metabolic gene analyses

To elucidate the mechanisms of Sammox reaction in modern microorganisms,

we sequenced the genome of *R. thioammoniphilus* (Extended Data Table 1, Extended

Data Fig. 7). There were denitrification genes in the *R. thioammoniphilus* genome,

including copper-containing nitrite reductase (*nirK*), nitric oxide reductase (*norB*),

and nitrous oxide reductase (*nosZ*) (Fig. 6a). This result confirmed that nitrite was the

expected intermediate product of Sammox and that the thiosulfate-/sulfate-fueled

Sammox reactions in *R. thioammoniphilus* contained two stages: oxidation of

ammonium to nitrite (Eqs. 3 and 4)[21], followed by the final step of the denitrification

pathway which is reduction of nitrite to dinitrogen (Eq. 5).

$$4NH_4^+ + 3S_2O_3^{2-} \rightarrow 4NO_2^- + 6HS^- + H_2O + 8H^+ \qquad (3)$$

$$4NH_4^+ + 3SO_4^{2-} \rightarrow 4NO_2^- + 3HS^- + 4H_2O + 5H^+ \qquad (4)$$

$$+2NO_2^- + 8H^+ + 6e^- \rightarrow N_2 + 4H_2O \qquad (5)$$

We did not find any other anaerobic ammonium oxidation-related functional

genes in the genome of *R. thioammoniphilus* (Fig. 6a). Thus, the ammonium oxidation

in Sammox was different from that in complete ammonium oxidation (Comammox)

and anaerobic ammonium oxidation (Anammox)[30].

Figure. 6 Representative metabolic gene clusters from _R. thioammoniphilus_

strain GX3-BWBA (a). The standard configuration of metabolic gene clusters

(MGCs) in primary metabolism located in two scaffolds. MGCs contain enzymes

of ■ CO$_2$ fixation genes, including 31 rTCA cycle, and 30 reductive

acetyl-CoA pathway genes, ■ 6 dissimilatory sulfur metabolic genes, and ■

5 denitrification genes. Gene size and spacing are not to scale. Each gene was not

condensed lie in genome. Protein phylogenetic tree derived from 26443 amino

acid sequences of sulfite/nitrite reductase (b). Scale bars represent estimated

sequence divergence or amino acid changes. ● Archaea, ▲ Bacteria, ■

Eukaryota.

Our main concern is how ammonium was oxidized by thiosulfate/sulfate into

nitrite. Sammox in *R. thioammoniphilus* may still need an iron-sulfur catalyst that

inherited from proto-anabolic networks. As lacking of genetic evidence about

Sammox, we predicted that ferredoxin oxidoreductase (e.g., nitrite/sulfite reductase

ferredoxin domain protein and ferredoxin-nitrite reductase) or other iron-sulfur

enzymes encoded by genes with unknown sequences may mediate sulfur reduction

coupling with ammonium oxidation to nitrite (Eqs. 3–6, Extended Data Schematic 1).

Here, thiosulfate should be first transformed into sulfite, which could be reduced by

ferredoxin. Hence, Sammox reversed the nitrite reduction process. This process was

quite similar to a nitrite generation pathway in Feammox process (Eq. 7)[30,41,42],

implying its theoretical feasibility. Electrons and protons generated in the first stage of

Sammox reaction were used to drive nitrite reduction to dinitrogen. As a highly

exergonic reaction, energy released from nitrite reduction drove CO_2 fixation.

$$NH_4^+ + 6 \text{ oxidized ferredoxin} + 2H_2O \rightarrow NO_2^- + 6 \text{ reduced ferredoxin} + 8H^+ \quad (6)$$

$$NH_4^+ + 6Fe^{3+} + 2H_2O \rightarrow NO_2^- + 6Fe^{2+} + 8H^+ \quad (7)$$

However, so far we do not clearly know the molecular mechanisms of sulfate

reduction coupling with ammonium oxidation to nitrite. *R. thioammoniphilus* contains

no ATP sulfurylase (Sat)-coding gene (Fig. 6a), suggesting that sulfate reduction in

sulfate-fueled Sammox is different from dissimilatory sulfate reduction. Dissimilatory

sulfate reduction may be later than sulfate reduction in sulfate-fueled Sammox,

according to isotope geochemical evidence[25,44,45]. That may be the reason sulfite

reductase (Extended Data Table 2, Extended Data Fig. 8) showed a much more

extensive distribution than Sat (Extended Data Table 3, Extended Data Fig. 9) and,

also the evidence for ancient metabolic networks led to the emergence of living

systems prior to the incorporation of P[19].

Our results thus far suggest that the transition from geochemistry to biochemistry

of S and N may have started from the same point. Sulfite and nitrite reductases belong

to the only class of enzymes that share a common architecture as well as a

requirement for a siroheme cofactor[46]. The phylogenetic tree indicated that the

putative multifunctional sulfite/nitrite reductase of archaea and eukaryota gathered

into one and three clusters, respectively. Putative interdomain lateral gene transfer

may result in the distribution of some archaea and eukaryota multifunctional

sulfite/nitrite reductases into bacterial clusters (Fig. 6b). Similar to sulfite reductase,

nitrite reductase is widely distributed in all types of modern organisms (Extended

Data Table 4, Extended Data Fig. 10), and indeed, sulfite reductases from some

sources can catalyze the reduction of both sulfite and nitrite[46]. This phenomenon

suggested that S and N biochemistry may have a common evolutionary origin derived

from Sammox.

Similar to the sulfite/nitrite reductase, some biomolecules as remnants of

Sammox and ancient metabolism may be hidden in the architecture of the metabolic

networks of *R. thioammoniphilus*. In theory, such "metabolic fossils" for the origin of

the coupled transformation of elements should be widely distributed in all types of

modern organisms as well. Thiosulfate may be the most preferential electron acceptor

for Sammox in Hadean hydrothermal systems, leading to the wide distribution of

multifunctional rhodanese (thiosulfate sulfurtransferase) in prokaryotes, eukaryotes,

and archaea (Extended Data Table 5, Extended Data Fig. 11). Rhodanese has been

proposed to have an assimilatory role using dithiol dihydrolipoate as the sulfur

acceptor and acting as a sulfur insertase involved in the formation of prosthetic groups

in iron-sulfur proteins, such as ferredoxin[47,48]. This implies that rhodanese may be the

primary mechanism for the formation of the iron-sulfur center of primordial enzymes

to catalyze proto-anabolic networks in LUCA, thus indicating its ancient nature.

Rhodanese can also catalyze a sulfur dissimilatory metabolic reaction which is

thiosulfate cleavage to sulfite[48] (Scheme 1), thereby facilitate thiosulfate involve in

the first stage of Sammox.

$$S_2O_3^{2-}+2R\text{-}S^- \rightarrow SO_3^{2-}+R\text{-}S\text{-}S\text{-}R+S^{2-}$$

375   Scheme 1

The structural similarity between rhodanese and Cdc25 phosphatases indicated

the common evolutionary origin of the two enzyme families[49]; alternatively,

phosphatases may originate from the rhodanese family because P incorporation

occurred later than S incorporation in metabolic networks. This evidence implied the

relationship between S and P metabolic evolution.

Six candidate genes encoding components of the reductive acetyl-CoA pathway

were identified as formate dehydrogenase (fdoI), 5,10-methenyl-H4 folate

cyclohydrolase (folD), 5,10-methylene-H4 folate dehydrogenase (folD),

5,10-methylene-H4 folate reductase (metF), CO dehydrogenase/acetyl-CoA synthase

(acs), and pyruvate:ferredoxin oxidoreductase (PFOR) (Fig. 6a). Three gene

homologs 10-formyl-H4 folate synthetase, methyl-H4 folate: corrinoid iron-sulfur

protein methyltransferase, and corrinoid iron-sulfur protein appeared to be missing. It

is possible that the present genomic analysis was insufficient to distinguish whether *R.*

thioammoniphilus could facilitate CO_2 fixation via the incomplete reductive

acetyl-CoA pathway or there were functional related enzymes encoded by genes with

unknown sequences[50].

In annotations of the rTCA cycle, two essential gene homologs appeared to be

missing (Fig. 6a). One of the missing genes, α-oxoglutarate synthase, catalyzes

reductive carboxylation from succinate to α-ketoglutarate, and the other is citryl-CoA

synthetase. Two potential reversible ATP-dependent citrate lyase subunits (ACL)

(citE, mdcC) were identified (Fig. 6a), which was a key indication for autotrophic

CO_2 fixation via the rTCA cycle instead of citrate synthase[51]. These results suggested

the potential of *R. thioammoniphilus* for carbon fixation via the incomplete reductive

acetyl-CoA pathway and rTCA cycle.

Final remarks: from metabolic innovation to evolution

In this study, we provided strong evidence of abiotic Sammox-driven

proto-anabolic networks and a Sammox microbe, *R. thioammoniphilus*. Fe-S mineral

catalysis and thiols/thioesters as energy couplers enhance the efficiency of prebiotic

proto-anabolic networks in thiosulfate-fueled Sammox reaction systems but not in

sulfate-fueled reaction systems under simulative hydrothermal conditions. Results

implied that the prebiotic thiosulfate-fueled Sammox reaction should be the

primordial reaction driving proto-anabolic networks, as it could be catalyzed. As a

probable ancient metabolism, Sammox microbe prefer thiosulfate to sulfate,

potentially because thiosulfate and/or elemental sulfur, and sulfite were produced

abundantly from ancient volcanic and hydrothermal SO_2. Our study was partially

supported by the theory of a chemoautotrophic origin of life supported by surface

metabolism and a primordial iron-sulfur world[9,11]. Here, Fe-S minerals and/or metal

alloy catalysis enhanced product generation of prebiotic thiosulfate-fueled Sammox

rather than providing hydrogen for CO_2 reduction. The Fe-S mineral surfaces, where

transition metal ions, such as Ni^{2+}, Co^{2+}, or Se, are catalytically active[52], and also the

very place facilitated prebiotic Sammox-driven amnio acids, thioesters and co-factors

synthesis. Therefore, a mixture of Fe-S minerals and transition metal ions may have

been the evolutionary precursor of the enzymatic active center, when chelated by

amnio acids and/or co-factors, converted to biocatalysts[33,38]. In this regard, our study

could explain the occurrence of enzymes with Fe-S reaction centers in the electron

transport chains of most known extant bioenergy flows in all three kingdoms of life.

Thioesters derived during prebiotic Sammox reaction provided the energetic and

catalytic framework of prebiotic Sammox, and feedback and feedforward to prebiotic

Sammox-driven proto-anabolic networks, ultimately leading to metabolic

reproduction and innovation[52]. Thus, prebiotic Sammox drove the emergence of the

primordial structure and function of the not free-living organic precursor, a ligand

sphere, held together by bonding to the surfaces of Fe-S minerals (Fig. 3c). The

feedback coupling between the primordial metabolic networks and their environment

could shape the evolution of both[53]. As a self-regulating system, C, H, O, N, S, and/or

Fe coupling metabolic networks could facilitate self-evolution through optimization

of the ambient environment, making it stable and habitable for life. The first and

foremost goal of Sammox-driven primordial metabolic networks was to obtain

sufficient phosphorus. Prebiotic Sammox-driven sulfur biochemistry prompted iron

redox geochemistry; therefore, phosphite was liberated from ferruginous sediments[39],

permitting biochemistry feedback to geochemistry. When P reached a certain

concentration in the Hadean oceans, sulfur biochemistry could lead to the emergence

of phosphorus biochemistry, and the RNA world could have replaced the thioester

world.

Our findings regarding Sammox-driven proto-anabolic networks suggested that

primordial metabolic networks might rise from coupling transformation of C, H, O, N,

S, and/or Fe at their earliest stage of the transition from geochemistry to biochemistry.

The elements of the chemical components that supporting prebiotic Sammox-driven

proto-anabolic networks ultimately became essential or trace elements of life,

providing a strong explanation for the unique features of life. Exploration of Sammox

provides a new perspective for understanding the emergence of biochemistry from

geochemistry and highlights the fundamental significance of Sammox for the origins

of life in hydrothermal environment on planetary systems.

References

1. Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive earth's biogeochemical cycles. *Science* **320**, 1034–1039 (2008).

2. Sousa, F. L. et al. Early bioenergetic evolution. *Philos. Trans. R. Soc. B-Biol. Sci.* **368**, (2013).

3. Weiss, M. C. et al. The physiology and habitat of the last universal common ancestor. *Nat. Microbiol.* **1**, (2016).

4. Moore, E. K. et al. Metal availability and the expanding network of microbial metabolisms in the Archaean eon. *Nat. Geosci.* **10**, 629–636 (2017).

5. Ljungdah, L., Irion, E. & Wood, H. G. Total synthesis of acetate from CO_2. I. Co-methylcobyric acid and CO-(methyl)-5-Methoxybenzimidazolylcobamide as intermediates with *Clostridium thermoaceticum. Biochemistry* **4**, 2771–2780 (1965).

6. Kelley, D. S. et al. A serpentinite-hosted ecosystem: the lost city hydrothermal field. *Science* **307**, 1428–1434 (2005).

7. Lane, N. & Martin, W. F. The origin of membrane bioenergetics. *Cell* **151**, 1406–1416 (2012).

8. Wächtershäuser, G. Pyrite formation, the first energy source for life: a hypothesis. *Syst. Appl. Microbiol.* **10**, 207–210, (1988).

9. Wächtershäuser, G. Evolution of the first metabolic cycles. *PHAS J.* **1**, 200–204 (1990).

10. de Duve, C. Blueprint for a cell: the nature and origin of life. (Neil Patterson ,

1991).

11. Evans, M. C. W., Buchanan, B. B. & Arnon, D. I. A new ferredoxin-dependent

carbon reduction cycle in a photosynthetic bacterium. *Proc. Natl. Acad. Sci. U. S.*

A. **55**, 928–934 (1966).

12. Smith, E. & Morowitz, H. J. Universality in intermediary metabolism. *Proc.*

Natl. Acad. Sci. U. S. A. **101**, 13168–13173 (2004).

13. Muchowska, K. B. et al. Metals promote sequences of the reverse krebs cycle.

Nat. Ecol. Evol. **1**, 1716–1721 (2017).

14. Varma, S. J. et al. Native iron reduces CO_2 to intermediates and end-products of

the acetyl-CoA pathway. *Nat. Ecol. Evol.* **2**, 1019–1024 (2018).

15. Martin, W. & Russell, M. J. On the origin of biochemistry at an alkaline

hydrothermal vent. *Philos. Trans. R. Soc. B-Biol. Sci.* **362**, 1887–1925 (2007).

16. Orgel, L. E. The implausibility of metabolic cycles on the prebiotic earth. *PLoS.*

Biol. **6**, 5–13 (2008).

17. Braakman, R. & Smith, E. The emergence and early evolution of biological

carbon-fixation. *PLoS Comput. Biol.* **8**, (2012).

18. Camprubi, E. et al. Iron catalysis at the origin of life. *IUBMB Life* **69**, 373–381

(2017).

19. Goldford, J. E. et al. Remnants of an ancient metabolism without phosphate. *Cell*

168, 1126–1134 (2017).

20. Fdz-Polanco, F. et al. Combining the biological nitrogen and sulfur cycles in

anaerobic conditions. *Water Sci. Technol.* **44**, 77–84 (2001).

21. Amend, J. P. et al. Energetics of chemolithoautotrophy in the hydrothermal

498   system of Vulcano Island, southern Italy. *Geobiology* **1**, 37–58 (2003).

499   22. Schrum, H. N. et al. Sulfate-reducing ammonium oxidation: A thermodynamically

500   feasible metabolic pathway in subseafloor sediment. *Geology* **37**(10), 939–942

501   (2009).

502   23. Rios-Del, T. E. E. et al. Anaerobic ammonium oxidation linked to sulfate and

503   ferric iron reduction fuels nitrogen loss in marine sediments. *Biodegradation* **29**,

504   429–442 (2018).

505   24. Cai, J. et al. Isolation and identification of bacteria responsible for simultaneous

506   anaerobic ammonium and sulfate removal. *Sci. China Chem.* **53**, 645–650

507   (2010).

508   25. Philippot, P. et al. Early archaean microorganisms preferred elemental sulfur, not

509   sulfate. *Science* **317**, 1534–1537 (2007).

510   26. Brandes, J. A. et al. Abiotic nitrogen reduction on the early earth. *Nature* **395**,

511   365–367 (1998).

512   27. Smirnov, A. et al. Abiotic ammonium formation in the presence of Ni-Fe metals

513   and alloys and its implications for the hadean nitrogen cycle. *Geochem. Trans.* **9**,

514   (2008).

515   28. Sleep, N. H. The hadean-archaean environment. *Cold Spring Harbor Perspect.*

516   *Biol.* **2**, (2010).

517   29. Crowe SA, et al. Sulfate was a trace constituent of Archean seawater. Science

518   346(6210):735–739 (2014).

30. Kuypers, M. et al. The microbial nitrogen-cycling network. *Nat. Rev. Microbiol.*

16, 263–276 (2018).

31. Eigen, M. Self-organization of matter and the evolution of biological

macromolecules. *Naturwissenschaften* **58**, 465–562 (1971).

32. Heinen, W. & Lauwers, A. M. Organic sulfur compounds resulting from the

interaction of iron sulfide, hydrogen sulfide and carbon dioxide in an anaerobic

aqueous environment. *Orig. Life Evol. Biosph.* **26**, 131–150 (1996).

33. Yamaguchi, A. et al. Electrochemical CO_2 reduction by Ni-containing iron

sulfides: how is CO_2 electrochemically reduced at bisulfide-bearing deep-sea

hydrothermal precipitates? *Electrochim. Acta* **141**, 311–318 (2014).

34. Martin, W. F., Sousa, F. L. & Lane, N. Evolution energy at life's origin. *Science*

344, 1092–1093 (2014).

35. Wächtershäuser, G. Before enzymes and templates-theory of surface metabolism.

Microbiol. Rev. **52**, 452–484 (1988).

36. Shen, Y. A., Buick, R. & Canfield, D. E. Isotopic evidence for microbial sulphate

reduction in the early Archaean era. *Nature* **410**, 77–81 (2001).

37. Mojzsis, S. J. et al.. Mass-independent isotope effects in Archean (2.5 to 3.8 Ga)

sedimentary sulfides determined by ion microprobe analysis. *Geochim.*

Cosmochim. Acta **67**, 1635–1658 (2003).

38. Eck, R. V. & Dayhoff, M. O. Evolution of the structure of ferredoxin based on

living relics of primitive amino acid sequences. *Science* **152**, 363–366 (1966).

39. Herschy, B. et al. Archean phosphorus liberation induced by iron redox

geochemistry. *Nat. Commun.* **9**, (2018).

40.   Jorgensen, B. B. A thiosulfate shunt in the sulfur cycle of marine-sediments.

Science **249**, (1990).

41.   Clement, J. C. et al. Ammonium oxidation coupled to dissimilatory reduction of

iron under anaerobic conditions in wetland soils. *Soil Biol. Biochem.* **37**,

2323–2328 (2005).

42.   Yang, W. H., Weber, K. A. & Silver, W. L. Nitrogen loss from soil through

anaerobic ammonium oxidation coupled to iron reduction. *Nat. Geosci.* **5**,

538–541 (2012).

43.   Bao, P. & Li, G. X. Sulfur-driven iron reduction coupled to anaerobic ammonium

oxidation. *Environ. Sci. Technol.* **51**, 6691–6698 (2017).

44.   Postgate, J. R. The sulphate-reducing bacteria (Cambridge University, 1984).

45.   Wacey, D. et al. Microfossils of sulphur-metabolizing cells in

3.4-billion-year-old rocks of Western Australia. *Nat. Geosci.* **4**, 698–702 (2011).

46.   Crane, B. R. & Getzoff, E. D. The relationship between structure and function

for the sulfite reductases. *Curr. Opin. Struct. Biol.* **6**, 744–756 (1996).

47.   Bonomi, F., Werth, M. T. & Kurtz, D. M. Assembly of [FenSn(SR)]2-(n=2, 4) in

aqueous media from iron salts, thiols and sulfur, sulfide, thiosulfide plus

rhodonase. *Inorg. Chem.* **24**, 4331–4335 (1985).

48.   Lefaou, A. et al. Thiosulfate, polythionates and elemental sulfur assimilation and

reduction in the bacterial world. *FEMS Microbiol. Lett.* **75**, 351–382 (1990).

49.   Bordo, D. & Bork, P. The rhodanese/Cdc25 phosphatase superfamily-

sequence-structure-function relations. *EMBO Rep.* **3**, 741–746 (2002).

50. Zhuang, W. Q. et al. Incomplete wood-ljungdahl pathway facilitates one-carbon

metabolism in organohalide-respiring *Dehalococcoides mccartyi. Proc. Natl.*

Acad. Sci. U. S. A. **111**, 6419–6424 (2014).

51. Mall, A. et al. Reversibility of citrate synthase allows autotrophic growth of a

thermophilic bacterium. *Science* **359**, 563–567 (2018).

52. Huber, C. & Wächtershäuser, G. Activated acetic acid by carbon fixation on

(Fe,Ni)S under primordial conditions. *Science* **276**, 245–247 (1997).

53. Lovelock, J. E. The Ages of Gaia (Oxford University, 1995).

Acknowledgements This research was financially supported by the National Natural

Science Foundation of China (General Program Nos. 41571240 and 41571130063).

Author contributions Peng Bao conceived the study, designed the experiment and

wrote the manuscript. Hu Li performed gas analysis. Guo-Xiang Li carried out all

other experiments and analysis. Peng Bao and Guo-Xiang Li contributed to

interpreting the data.

Competing interests The authors declare no competing interests.

Additional information

Extended data is available for this paper at

Supplementary information is available for this paper at

METHODS

Enrichment and isolation of Sammox bacterium.

Surface paddy soil mud (0–20 cm depth) in the upper section of a paddy field was collected from Guangxi Dahuanjiang region (24°53'52" N, 108°17'43" E), in the southwest of China. Soil materials were air-dried, ground with a mortar and crushed to pass through a 2.0 mm sieve for character determination. Soil organic carbon (19.81 g kg^{-1}) was determined by potassium dichromate oxidation titration, and soil total Fe (31.9 g kg^{-1}) was determined by ICP-OES. Plant available sulfur (29.1 mg kg^{-1}) was determined by turbidimetry. Ammonium was 120.5 mg kg^{-1}.

For enrichment, approximately 0.5 g of paddy soil was transferred to 100 ml of double distilled water. After shaking, five milliliters of the suspension was inoculated into 100 ml of anaerobic freshwater medium[43,54] with additional sulfur (thiosulfate or sufate) (3.0 mM) and ammonium ($^{15}NH_4Cl$, 1.0 mM) and incubated statically for one week. The basal medium contained (g l^{-1}): KH_2PO_4 0.2, $MgCl\cdot6H_2O$ 0.4, $CaCl_2\cdot2H_2O$ 0.1, KCl 0.5. The trace elemental mixture contained (l^{-1}): double distilled water 987 ml, HCl (25%) 12.5 ml (100 mM), $FeCl_2\cdot4H_2O$ 957 mg (7.5 mM), H_3BO_3 30 mg (0.5 mM), $MnCl_2\cdot4H_2O$ 100 mg (0.5 mM), $CoCl_2\cdot6H_2O$ 190 mg (0.8 mM), $NiCl\cdot6H_2O$ 24 mg (0.1 mM), $CuCl_2\cdot2H_2O$ 2 mg (0.01 mM), $ZnSO_4\cdot7H_2O$ 144 mg (0.5 mM), $Na_2MoO_4\cdot2H_2O$ 36 mg (0.15 mM). Selenite-tungstate solution (l^{-1}): NaOH 0.4 g, $Na_2SeO_3\cdot5H_2O$ 6 mg, $Na_2WO_4\cdot2H_2O$ 8 mg. Bicarbonate solution $NaHCO_3$ 84 g l^{-1} 30 ml. The above mentioned stock solutions or aliquots were aseptically added to the basal medium (l^{-1}): trace element solution (1.0 ml), selenite-tungstate solution (0.1 ml),

bicarbonate solution (10.0 ml), biotin, 4-aminobenzoic acid, 10 mg l^{-1} pantothenate,

pyridoxamine, nicotinic acid and 20 mg l^{-1} thiamine, 5 ml l^{-1}; vitamin B_{12} solution

with 50 mg l^{-1}, 1.0 ml l^{-1}. Sodium sulfate plus thiosulfate solution (1.0 ml, 3.0 mM

final concentration) and ammonium (0.5 ml, 1.0 mM final concentration) were added

into medium if necessary. The pH was adjusted to 8.0.

A volume of 0.5 ml of a positive Sammox culture, in which sulfur reduction

coupled to anaerobic ammonium oxidation, was sub-inoculated onto Acidovorax

complete agar medium under 5% CO_2 atmosphere. Single colony was selected and

inoculated back into anaerobic freshwater medium. Positive Sammox samples were

continuously sub-inoculated using the Acidovorax complete agar medium. A single

Sammox bacterium (designated GX3-BWBA) was isolated after sub-cultivation for

five times, and was continuously cultivated for one year in anaerobic freshwater

medium. The purity of GX3-BWBA was confirmed by promoting the growth of

heterotrophic bacteria by addition of peptone and yeast extract to the defined

freshwater mineral medium (data not shown)[55]. Finally, the purity of GX3-BWBA

was confirmed by deep Illumina sequencing.

This study was performed using a series of experiments, for Sammox by

GX3-BWBA: (I) 3.0 mM thiosulfate + 1.0 mM $^{15}NH_4Cl$, (II) 3.0 mM thiosulfate + 1.0

mM $^{15}NH_4Cl$ + GX3-BWBA, (III) 3.0 mM sulfate + 1.0 mM $^{15}NH_4Cl$, (IV) 3.0 mM

sulfate + 1.0 mM $^{15}NH_4Cl$ + GX3-BWBA. Experiments started and were sampled at

the sampled at the 0, 3, 6, 9, 12, 15, 18, 21, 24, 27, 34 day. Experiments were

performed anaerobically in 100-ml serum bottles in the dark at 30 ^{o}C. Each batch of

experiments was established in triplicate, inoculated with 1.0 ml GX3-BWBA

exponential phase culture if necessary.

Genome sequencing and assembly

The genome of *R. thioammoniphilus* (strain GX3-BWBA) was sequenced using

an Illumina HiSeq 4000 system (Illumina, SanDiego, CA, USA) at the Beijing

Genomics Institute (Shenzhen, China). Genomic DNA was sheared randomly to

construct three read libraries with lengths of (150:150) by a Bioruptor ultrasonicator

(Diagenode, Denville, NJ, USA) and physico-chemical methods. The paired-end

fragments libraries were sequenced according to the Illumina HiSeq 4000 system's

protocol. Raw reads of low quality from paired-end sequencing (those with

consecutive bases covered by fewer than five reads) were discarded. The sequenced

reads were assembled using SOAPdenovo v1.05 software. This Whole Genome

Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession

QMKS00000000. The version described in this paper is version QMKS01000000.

Genome Component prediction

Gene prediction was performed on the *R. thioammoniphilus* genome assembly by

glimmer3 (http://www.cbcb.umd.edu/software/glimmer/) with Hidden Markov models.

tRNA, rRNA and sRNAs recognition made use of tRNAscan-SE[56], RNAmmer, and

the Rfam database. The tandem repeats annotation was obtained using the Tandem

Repeat Finder (http://tandem.bu.edu/trf/trf.html), and the minisatellite DNA and

microsatellite DNA selected based on the number and length of repeat units. The

Genomic Island Suite of Tools (GIST) used for genomic islands analysis

(http://www5.esu.edu/cpsc/bioinfo/software/GIST/) with Island Path-DIOMB,

SIGI-HMM, Island Picker method. Prophage regions were predicted using the PHAge

Search Tool (PHAST) web server (http://phast.wishartlab.com/) and CRISPR

identification using CRISPRFinder.

Protein sequences will be downloaded from NCBI, and get sequences in the

taxonomy of species classification information; The sequences using mafft software

for multiple sequence alignment. After comparing the sequences, using Fasttree

software construct phylogenetic tree (neighbor joining algorithm). Using R language

ggtree package for visualization mapping.

Gene annotation and protein classification

The best hit abstracted using Blast alignment tool for function annotation. Seven

databases which are KEGG (Kyoto Encyclopedia of Genes and Genomes), COG

(Clusters of Orthologous Groups), NR (Non-Redundant Protein Database databases),

Swiss-Prot[57], and GO (Gene Ontology), TrEMBL, EggNOG are used for general

function annotation. Representative metabolic gene clusters were displayed with

Easyfig[58].

General procedure for prebiotic Sammox-driven CO_2 fixation

Each 50 ml basal solution was transferred into 60 ml serum bottles and sealed

with butyl rubber stoppers and aluminium crimp caps. The solution was autoclaved

and cooled as room temperature after washed by He gas (purity=99.999%). Additional

trace elemental mixture solutions were filter sterilized or autoclaved individually.

Additional sulfur (thiosulfate or sufate) (3.0 mM) and/or ammonium (NH_4Cl, 1.0 mM)

was added into serum bottle. The mixed solution was termed as Sammox reaction

system.

The basal solution contained (l^{-1}): KCl 0.1 g. The trace elemental mixture

contained (l^{-1}): double distilled water 987 ml, HCl (25%) 12.5 ml (100 mM), H_3BO_3

30 mg (0.5 mM), $MnCl_2·4H_2O$ 100 mg (0.5 mM), $CoCl_2·6H_2O$ 190 mg (0.8 mM),

$NiCl·6H_2O$ 24 mg (0.1 mM), $CuCl_2·2H_2O$ 2 mg (0.01 mM), $FeCl_2·4H_2O$ 957 mg (7.5

mM), $ZnCl_2$ 68 mg (0.5 mM), $Na_2MoO_4·2H_2O$ 36 mg (0.15 mM). Selenite-tungstate

solution (l^{-1}): NaOH 0.4 g, $Na_2SeO_3·5H_2O$ 6 mg, $Na_2WO_4·2H_2O$ 8 mg. Bicarbonate

solution $NaHCO_3$ 84 g l^{-1} 30 ml. The above mentioned stock solutions or aliquots

were aseptically added to the basal medium (l^{-1}): trace element solution (2.0 ml),

selenite-tungstate solution (0.2 ml), bicarbonate solution (10.0 ml). Sodium sulfate,

thiosulfate (1.0 ml, 3.0 mM final concentration), ammonium solution (0.5 ml, 1.0 mM

final concentration), and fresh precipitated Fe-S mineral (1.0 mM) were added into

medium if necessary. Serum bottles were kept 100 ^{o}C in a water bath in the dark for

24 h, then maintain at 70 ^{o}C in the dark for another 24 h, and sampling at 48 h to

determinate organic products.

This study was performed using a series of experiments: (I) 3.0 mM thiosulfate +

1.0 mM NH_4Cl + 20 mM HCO_3^-, (II) 3.0 mM thiosulfate + 1.0 mM NH_4Cl + 20 mM

HCO_3^- + 1.0 mM Fe-S, (III) 3.0 mM thiosulfate + 20 mM HCO_3^-, (IV) 3.0 mM sulfate

+ 1.0 mM NH_4Cl + 20 mM HCO_3^-, (IV) 3.0 mM sulfate + 1.0 mM NH_4Cl + 20 mM

HCO_3^- + 1.0 mM Fe-S, (V) 3.0 mM sulfate + 20 mM HCO_3^-, (VI) 1.0 mM NH_4Cl +

20 mM HCO_3^-.

General procedure for thiol/thioester promoted Sammox-driven CO₂ fixation.

A solution of methanethiol (0.8 mM, final concentration) was added to Sammox reaction system. The reaction systems were heated at 100 °C in a water bath in the dark for 24 h, then maintain at 70 °C in the dark for another 24 h, and removed from the water bath and allowed to cool to room temperature before derivatization and gas chromatography–mass spectrometry analysis.

This study was performed using a series of experiments: (I) 3.0 mM thiosulfate + 1.0 mM NH₄Cl + 20 mM HCO₃⁻ + 0.8 mM methanethiol, (II) 3.0 mM thiosulfate + 1.0 mM NH₄Cl + 20 mM HCO₃⁻ + 1.0 mM Fe-S + 0.8 mM methanethiol, (III) 3.0 mM sulfate + 1.0 mM NH₄Cl + 20 mM HCO₃⁻ + 0.8 mM methanethiol, (IV) 3.0 mM sulfate + 1.0 mM NH₄Cl + 20 mM HCO₃⁻ + 1.0 mM Fe-S + 0.8 mM methanethiol.

General procedure for verification of Sammox-driven combination of reductive acetyl-CoA pathway and incomplete rTCA cycle.

We design this experimental set to verify if Sammox-driven reductive acetyl-CoA pathway could go into Sammox-driven incomplete rTCA cycle. The general procedure is the same as above. Pyruvate (1.0 mM, final concentration) was added into Sammox reaction system as substrate. Serum bottles were heated at 100 °C in a water bath in the dark for 24 h, then maintain at 70 °C in the dark for another 24 h, and allowed to cool to room temperature before derivatization and gas chromatography–mass spectrometry analysis.

This study was performed using a series of experiments: (I) 3.0 mM thiosulfate + 1.0 mM NH₄Cl + 20 mM HCO₃⁻ + 1.0 mM pyruvate + 1.0 mM Fe-S + methanethiol,

(II) 3.0 mM thiosulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM Fe-S + 1.0 mM

pyruvate, (III) 3.0 mM thiosulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM

pyruvate, (IV) 3.0 mM thiosulfate + 20 mM HCO_3^- + 1.0 mM pyruvate, (V) 3.0 mM

sulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM Fe-S + 1.0 mM pyruvate +

methanethiol, (VI) 3.0 mM sulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM Fe-S

+ 1.0 mM pyruvate, (VII)3.0 mM sulfate + 1.0 mM NH_4Cl + 20 mM HCO_3^- + 1.0

mM pyruvate, (VIII) 3.0 mM sulfate + 20 mM HCO_3^- + 1.0 mM pyruvate, (IX) 1.0

mM NH_4Cl + 20 mM HCO_3^- + 1.0 mM pyruvate.

Chemicals

All reagents and solvents were purchased from commercial suppliers and used

without further purification unless otherwise noted.

Sampling analytical methods

(i) **Enumeration of microbe viable counts.** The spread-agar-plate method was used

for enumeration of bacterial counts in GX3-BWBA culture[59]. The Acidovorax

complete agar medium, pH 7.0, contained 5.0 g l^{-1} peptone (Difco), 3.0 g l^{-1} beef

extract and 15 g l^{-1} agar (Difco). A 50 µl sample was taken from GX3-BWBA culture

at the 0, 3, 6, 9, 12, 15, 18, 21, 24, 27 and 34 days and inoculated onto plates for cell

counting. Visible colonies produced on the agar plate were counted at 48 h. Bacterial

numbers were expressed as colony-forming units (CFU) per milliliter of culture[59].

(ii) **Electron microscopy.** Morphology of GX3-BWBA was investigated by

transmission electron microscope (TEM) (TEM, HITACHI H-7500). Cells from the

culture were collected by centrifugation, washed and diluted with phosphate buffer

(pH 7.5), and dropped on to specimens. The material was examined after dry by

airing.

(iii) **Sulfide.** The analysis of sulfide in filtered samples was performed

photometrically by the methylene blue method[60]. Absorption at 660 nm was measured

(Infinite 200PRO; TECAN).

(iV) **Sulfate, sulfite, nitrite, acetate, formate and ammonium.** To determine sulfate,

sulfite, nitrite, acetate and formate, 0.5 ml of the sample was filtered (0.22 μm) to

remove particulates that could interfere with ion chromatography. The ion

chromatography system consisted of an ICS-5000$^+$ SP pump (Thermo Fisher

Scientific Inc. Sunnyvale, CA, USA), a column oven ICS-5000$^+$ DC, an

electrochemical detector DC-5. The ion chromatography column system used a

Dionex Ionpac AS11-HC column. The operating condition was with an eluent of 30

mM KOH at a flow rate of 1.0 ml min^{-1}. For determination of ammonium, the ion

chromatography column system used a Dionex Ionpac CS12A column.

(V) **Thiosulfate.** Thiosulfate was determined by an Agilent 1260 infinity HPLC

system, equipped with a quaternary pump (Agilent, USA). Thiosulfate was separated

by a Zorbax SB-C18 column (150×4.6 mm, 5 μm), and detected by using of

DAD detector at 215 nm. All analyses were performed at 40 °C with a flow rate of 1

ml min^{-1}. Na_2HPO_4 was used as solvent. The pH of the solvent was adjusted with 1.0

M HCl. Samples were filtered through 0.45 μm Cosmonice Filters (Millipore, Tokyo,

Japan) and immediately injected into the HPLC system[61].

(Vi) Gas Analysis. At each sampling time point, each serum bottle was shaken

vigorously to equilibrate the N_2 between dissolved and gas phases. Dinitrogen

sampling and determination was performed according to a literature procedure[41].

(Vii) Derivatization procedure and product identification

For optimal GC resolution, the carboxylic acids were converted to ethyl esters

using a mixture of ethanol/ethyl chloroformate (EtOH/ECF). Derivatization of

carboxylic acids to esters was performed according to a literature procedure[13].

Reaction products derivatized to ethyl esters of carboxylic acids were identified by

comparing the mass spectra and retention times against analogously derivatized

authentic samples. ECF derivatization was preferred for small molecule substrates

(pyruvate, lactate, malate, fumarate, succinate, α-ketoglutarate, amino acid). The

carboxylic acids might also convert to methyl esters using a mixture of

methanol/methyl chloroformate (MeOH/MCF), following the same procedure to ECF

derivatization to detect *cis*-aconitate, tricarballylate, isocitrate and citrate.

(Viii) Gas chromatography–mass spectrometry (GC–MS) analysis for rTCA

metabolites determination.

GC–MS analysis was performed on a GC System 7890B connected to a MSD

block 5977A, using Agilent High Resolution Gas Chromatography Column: PN

19091S–433, HP–5MS, 28 m×0.25 mm, 0.25 Micron, SN USN 462366H. Samples

were prepared in ethyl acetate (200 μl sample volume). The analysis was carried out

on a splitless 1 μl injection volume with an injection port temperature 250 °C.

Column oven temperature program was as follows: 60 °C for 1.0 min, ramped at

30 °C min^{-1} to 310 °C with 3.0 min hold, with a total running time of 12.33 min. The

mass spectrometer was turned on after 3 min and was operated at the electron

ionization mode with quadrupole temperature of 150 °C. Data acquisition was

performed in the full-scan mode (50-500). Hydrogen (99.999 % purity) was used as

carrier gas at a constant flow rate of 1.5 ml min^{-1}.

(iX) liquid chromatography-mass spectrometry (LC-MS) method for polypyrrole

determination.

For qualitative detection of polypyrrole, Waters ACQUITY UPLC system with

an online coupled SYNAPT G2 mass spectrometer Q-TOF was used. Sample

separation was achieved on a ACQUITY UPLC HSS T3 column (1.8 μ m, 2.1 mm ×

100 mm; column temperature, 30 °C). Solvent A contained 2.5% methanol, 0.2%

formic acid in UPLC-grade water, and Solvent B was 97.5% UPLC-grade water with

0.2% formic acid. Injection of 2.0 μl of sample onto the column at 0.2 ml min^{-1} was

followed by gradient elution. Mass spectrometric qualitative detection of polypyrrole

was performed in resolution mode. Polypyrrole was identified by comparing the mass

spectra and retention times against pure commercially available pyrrole (100 μl ml^{-1})

heated at 100 °C in a water bath in the dark for 24 h, and maintain at 70 °C in the dark

for another 24 h. Tandem mass spectrometry data were analyzed using MassLynx

v4.1.

54.   Strous, M. et al. Deciphering the evolution and metabolism of an anammox

bacterium from a community genome. *Nature* **440**, 790–794 (2006).

55.   Konneke, M. et al. Isolation of an autotrophic ammonia oxidizing marine

archaeon. *Nature* **437**, 543–546 (2005).

56.   Lowe, T. M. & Eddy, S. R. tRNAscan-SE: a program for improved detection

of transfer RNA genes in genomic sequence. *Nucl. Acids Res*. **25** (5),

0955–0964 (1997).

57.   Torto-Alalibo, T., Collmer, C. W. & Gwinn-Giglio, M. The plant-associated

microbe gene ontology (PAMGO) consortium: community development of

new gene ontology terms describing biological processes involved in

microbe-host interactions. *BMC microbiol.* **9** (suppl 1), (2009).

58.   Sullivan, M. J., Nicola K. P. & Scott A. B. Easyfig: a genome comparison

visualizer. *Bioinformatics* **27** (7), 1009–1010 (2011).

59.   Kataoka, N. et al. Enrichment culture and isolation of slow growing bacteria.

Appl. Microbiol. *Biotechnol.* **45**, 771–777 (1996).

60.   Cline, J. D. Spectrophotometric determination of hydrogen sulfide in natural

waters. imnol. *Oceanogr*. **14** (3), 454–458 (1969).

61.   Kondo, R. et al. Determination of thiosulfate in a meromictic lake. *Fisheries*

Sci. **66**, 1076–1081 (2000).

Legends of figures

Figure 1. Products of simulated prebiotic Sammox-driven proto-anabolic

networks with bicarbonate as the sole carbon source under hydrothermal

conditions. Treatments were as follows (a, thiosulfate; b, sulfate), from left to

right: (i) sulfurous species, ammonium, Fe-S minerals, and methanethiol; (ii)

sulfurous species, ammonium, and Fe-S minerals; (iii) sulfurous species,

ammonium, and methanethiol; (iv) sulfurous species and ammonium; (v)

ammonium; and (vi) sulfurous species. The bar chart shows the yields of formate,

acetate, succinate, and α-ketoglutarate in each treatment group. Error bars

represent standard deviations of three replicates.

Figure 2. Products of simulated prebiotic Sammox-driven incomplete rTCA cycle

with bicarbonate and pyruvate as carbon sources under hydrothermal

conditions. Treatments were as follows (a, thiosulfate; b, sulfate), from left to

right: (i) sulfurous species, ammonium, Fe-S minerals, and methanethiol; (ii)

sulfurous species, ammonium, and Fe-S minerals; (iii) sulfurous species,

ammonium, and methanethiol; (iv) sulfurous species and ammonium; (v)

ammonium; and (vi) sulfurous species. The bar chart shows the concentrations

of pyruvate, succinate, and α-ketoglutarate in each treatment group. Error bars

represent standard deviations of three replicates.

Figure. 3 Hypothetical prebiotic Sammox-driven reductive acetyl-CoA pathway

(a). Hypothetical proto-anabolic networks, a combination of acetyl-CoA pathway

together with incomplete rTCA cycle driven by Sammox, and the role of its

intermediates as universal biosynthetic precursors. Bar represents the final step

of incomplete rTCA cycle (b). Conceptual model of Sammox-driven the origin of

not free-living organic precursor on the surface of Fe-S minerals in Hadean

alkaline hydrothermal systems. Cubes represent Fe-S minerals/metals alloy.

Globes represent the not free-living organic precursor, ligand sphere (c).

Figure. 4 Characters of *R. thioammoniphilus* GX3-BWBA. TEM image of cells

(a). Scale bar represents 0.2 μm. Growth of GX3-BWBA under combined

condition of ammonium and sulfurous species, with bicarbonate as sole carbon

sources (b). Production rate of isotopically labeled dinitrogen for thiosulfate and

sulfate-fueled Sammox by GX3-BWBA (c, d). Phylogenetic tree constructed by

the maximum likelihood method, using 1478 nucleotides from 16S rDNA

sequences, showing the position of strain GX3-BWBA in relation to members of

the genus Ralstonia. The bar represents 0.1 changes per nucleotide position (e).

Figure. 5 Thiosulfate- and sulfate-fueled Sammox by *R. thioammoniphilus*

GX3-BWBA. (a) thiosulfate and reduction products. (b) ammonium and its

oxidation products in thiosulfate-fueled Sammox. (c) sulfate and reduction

products (sulfide and sulfite as products of sulfate in control group are below

detection limit). (d) ammonium and its oxidation products in sulfate-fueled

Sammox. Error bars represent standard deviations of three biological replicates.

Figure. 6 Representative metabolic gene clusters from _R. thioammoniphilus_

strain GX3-BWBA (a). The standard configuration of metabolic gene clusters

(MGCs) in primary metabolism located in two scaffolds. MGCs contain enzymes

of ▮ CO_2 fixation genes, including 31 rTCA cycle, and 30 reductive

acetyl-CoA pathway genes, ▮ 6 dissimilatory sulfur metabolic genes, and ▮

5 denitrification genes. Gene size and spacing are not to scale. Each gene was not

condensed lie in genome. Protein phylogenetic tree derived from 26443 amino

acid sequences of sulfite/nitrite reductase (b). Scale bars represent estimated

sequence divergence or amino acid changes. ● Archaea, ▲ Bacteria, ▮

Eukaryota.

890