1 **Differential gene expression supports a resource-intensive, defensive role for colony**
2 **production in the bloom-forming haptophyte, *Phaeocystis globosa***
3

4 Margaret Mars Brisbin[a] and Satoshi Mitarai[a]
5
6 a Marine Biophysics Unit, Okinawa Institute of Science and Technology Graduate University,
7 Okinawa, Japan
8
9 **Correspondence**
10 M. Mars Brisbin, Marine Biophysics Unit, Okinawa Institute of Science and Technology
11 Graduate University, 1919-1 Tancha, Onna-son, Okinawa, Japan
12 e-mail: margaret.marsbrisbin@oist.jp
13

14 **Abstract**
15

Phaeocystis globosa forms dense, monospecific blooms in temperate, northern waters. Blooms
are usually dominated by the colonial morphotype—non-flagellated cells embedded in a secreted
mucilaginous mass. Colonial *Phaeocystis* blooms significantly affect food-web structure and
function and negatively impact fisheries and aquaculture, but factors initiating colony production
remain enigmatic. Destructive *P. globosa* blooms have been reported in tropical and subtropical
regions more recently and warm-water blooms could become more common with continued
climate change and coastal eutrophication. We therefore assessed genetic pathways associated
with colony production by investigating differential gene expression between colonial and
solitary cells in a warm-water *Phaeocystis globosa* strain. Our results illustrate a transcriptional
shift in colonial cells with most of the differentially expressed genes downregulated, supporting a
reallocation of resources associated with colony production. Dimethyl sulfide and acrylate
production and pathogen interaction pathways were upregulated in colonial cells, suggesting a
defensive role for colony production. We identify several protein kinase signaling pathways that
may influence the transition between morphotypes, providing targets for future research into
factors triggering colony production. This study provides novel insights into genetic mechanisms
involved in *Phaeocystis* colony formation and provides new evidence supporting a defensive role
for *Phaeocystis* colonies.
33

34 **Keywords**
35 Plankton; phytoplankton; DMS; DMSP; algal bloom; algae bloom; RNA-seq; transcriptome;
36 transcriptomics; colonial morphotype
37
38 **Introduction**
39
Phaeocystis is a cosmopolitan bloom-forming haptophyte genus encompassing 6 species
(Schoemann et al 2005, Andersen et al. 2015). Most *Phaeocystis* species (*P. globosa, P.*
antarctica, P. pouchetii, and *P. jahnii*) exhibit a polymorphic life-cycle, alternating between
colonial and free-living morphotypes. *Phaeocystis* blooms are usually dominated by the colonial
morphotype and are typically very dense, produce large biomasses, and impact food-web
structure and function (Schoemann et al. 2005). *Phaeocystis* is a major contributor to
Dimethylsulfoniopropionate (DMSP) and dimethyl sulfide (DMS) production globally (Liss et

al. 1994) and regional peaks in DMS production are closely correlated with colonial *Phaeocystis*
blooms (Van Duyl et al. 1998). DMS produced in the surface ocean is aerosolized and its
oxidation products promote cloud formation, increase albedo, and affect global climate
(Charlson et al. 1987). In algal cells, DMSP and its cleavage products, DMS and acrylate,
contribute to osmotic balance, neutralize reactive oxygen species, and deter grazing (Sunda et al.
2002, Noordkamp et al. 2000). Despite the ecological importance of colony formation in
Phaeocystis, triggers for transition to the colonial morphotype remain enigmatic, and the
functional role of colony formation in the *Phaeocystis* life-cycle is not clearly delineated
(Peperzak & Gabler-Schwarz 2012).
56
Myriad factors have been studied in regard to their roles in instigating colony formation in
Phaeocystis species, including nutrient and light availability (Cariou et al. 1994, Wang et al.
2011, Bender et al. 2018), temperature (Verity and Medlin 2003), mechanical stress (Cariou et
al. 1994), grazing cues (Tang 2003, Long et al. 2007, Wang et al. 2015), and viral infection
(Brussaard et al. 2005, Brussaard et al. 2007). However, these studies used different *Phaeocystis*
species, strains, and morphotypes with a range of experimental conditions, which yielded
variable and sometimes contradictory results. Nonetheless, several lines of evidence suggest that
colony formation serves a defensive role. First, while viruses can cause 30-100% cell lysis in
solitary *Phaeocystis,* viruses rarely infect colonial cells, which lyse primarily due to nutrient
limitation (Brussaard et al. 2005, 2007). Second, ciliates and other microzooplankton that graze
solitary *Phaeocystis* are unable to graze on colonies (Tang et al. 2001) and chemical cues from
these grazers induce colony formation and promote increased colony size (Tang 2003, Long et
al. 2007). Third, acrylate, which is produced with DMS when DMSP is cleaved, accumulates
within colonies and may further deter macro- and micro-grazers and heterotrophic bacteria
(Noordkamp et al. 2000, Hamm et al. 2000). However, while cellular growth rate increases in
colonial cells relative to solitary cells if colonies are induced in nutrient rich conditions, it
decreases when colonies are induced under nutrient limiting conditions (Wang et al. 2015). Thus,
colony formation can defend against pathogens and grazers, but it is costly (Wang et al. 2015),
suggesting that colony formation is likely a complex response to interacting biotic and abiotic
factors (Long et al. 2007).
77
Colony formation may also play a fundamental role in *Phaeocystis* reproduction. *Phaeocystis* has
one of the most complex and polymorphic life cycles among phytoplankton genera, and despite
extensive study, it remains largely unresolved in most species. Studies have implicated at least 6
different life stages and up to 15 functional components to the life-cycle (Gaebler-Schwarz et al.
2010). In *P. globosa,* four morphotypes are believed to exist: diploid colonial cells devoid of
scales and flagella, diploid scale-free flagellates arising from mechanically disrupted colonies,
and two types of small, scaled, haploid flagellates—those that produce vesicles containing star-
shaped alpha-chitin filaments and those that do not (Rousseau et al. 2007). Haploid flagellates
may fuse (syngamy) to produce diploid colony-forming cells, which in turn undergo meiosis and
produce haploid flagellates (Rousseau et al. 2013). Haploid flagellates are often observed
swarming inside colonies, suggesting that colonial bloom formation may contribute to successful
sexual reproduction in *Phaeocystis* (Peperzak et al. 2000, Rousseau et al. 2013). However,
neither syngamy nor meiosis have been directly observed in *Phaeocystis* spp. (Peperzak and
Gabler-Schwarz 2012), even though both events have been documented in several other

haptophyte genera (Houdan et al. 2003). If colonial *Phaeocystis* blooms are necessary for sexual
reproduction, it would further justify the resource costs associated with colony formation.
94
Historically, colonial *Phaeocystis* blooms have been restricted to cold, high-latitude waters—*P.*
globosa blooms in the English Channel and North Sea, *P. pouchetii* in the North Atlantic and
Arctic, and *P. antarctica* in the Southern Ocean (reviewed in Schoemann et al. 2005). In the last
two decades, however, blooms have increasingly been reported in tropical and subtropical
regions, including the subtropical N. Atlantic (Long et al. 2007) and the subtropical and tropical
South China Sea (Chen et al. 2002, Doan-Nhu et al. 2010, Liu et al. 2015). Decaying colonial
biomass sinks and produces anoxic conditions, making *Phaeocystis* blooms detrimental to
benthic fisheries and aquaculture (Desroy & Denis 2008, Peperzak & Poelman 2008, Spilmont et
al. 2009). In warmer waters, *Phaeocystis globosa* blooms have been especially catastrophic to
local aquaculture (Chen et al. 2002, Doan-Nhu et al. 2010), possibly because the hemolytic
activity of *P. globosa* liposaccharides increases with temperature (Peng et al. 2005). Global
climate change and increasing nutrient pollution in coastal regions may mean harmful
Phaeocystis blooms will continue to increase in range and frequency. Given the ecological
impact of colonial *Phaeocystis* blooms and their complex and enigmatic initiating triggers,
particularly in warm waters, it is imperative to better understand the regulation of colony
formation.
111
Transcriptional approaches have become an exceptionally useful tool to illuminate physiological
responses to environmental cues and genes associated with specific life-stages in algae and other
protists (Caron et al. 2017). In this study, we investigated genetic regulation of colony formation
by analyzing gene expression in colonial and flagellated morphotypes of a warm-water
Phaeocystis globosa strain. Since *Phaeocystis* is an important marine producer of DMSP and
DMS—and because these molecules may be associated with colonial defense, we queried our
dataset for algal genes involved in DMSP production (*DSYB*, Curson et al. 2018) and its
cleavage to DMS and acrylate (*Alma1*, Alcolombri et al. 2015). *DSYB* and *Alma1* are the only
algal genes that encode proteins experimentally proven to catalyze DMSP, DMS, and acrylate
production, but neither has been identified in *P. globosa* previously. Overall, our results
demonstrate a dramatic transcriptional shift in colonial *P. globosa*, with the vast majority of
differentially expressed genes downregulated in colonial cells. Such a strong transcriptional shift
supports an allocation of resources toward colony formation and away from other cellular
processes such as translation, cell growth, and cell division. Genes associated with DMSP
production (*DSYB*-like) were not differentially expressed, but an *Alma* family-like gene was
upregulated in colonies, suggesting colonies may produce more DMS and acrylate than do
solitary cells. Results further indicate that pathogen interaction affects colony formation, further
supporting a defensive role for colonies, and implicating several cell-signaling pathways as
important to colony formation. This study provides new insights into the functional role of
Phaeocystis colonies and physiological processes associated with colony formation. These
insights will guide future investigations into triggers that initiate colony formation in harmful
Phaeocystis blooms.
134
135 **Materials and Methods:**
136
137 ***Culture strain and maintenance***

138
Starter cultures of *Phaeocystis globosa* CCMP1528, a warm-water, colony forming *P. globosa*
strain (Wang et al. 2011), were purchased from the National Center for Marine Algae and
Microbiota (NCMA, Maine, U.S.A.) in January 2016. CCMP1528 did not initially form colonies
in our culture conditions, but it is known that *P. globosa* sometimes stops forming colonies in
culture (Janse et al. 1996). We maintained non-colonial cultures in replicate 150 mL Erlenmeyer
flasks with 100 mL L1-Si media in ambient light and temperature on a gently rotating twist
mixer (TM-300, AS ONE, Osaka, Japan, speed setting 1). Culture media was prepared by
enriching 34 ppt autoclaved artificial seawater prepared from milliQ water and sea salts (Marine
Art SF-1) with the NCMA L1 media kit (-Si) and filtering through sterile 0.22 μm pore-size
filters. Cultures were diluted biweekly with freshly prepared media. When one culture replicate
began producing colonies in January 2017, experimental culture conditions were promptly
initiated.
151
152   ***Experimental culture conditions***
153
We prepared four biological replicates each of colony-forming and non-colonial *P. globosa*
CCMP1528 by inoculating 45 mL of sterile L1 media with 1 mL stock culture in 50 mL
Erlenmeyer flasks. Replicates were placed on a gently rotating twist mixer in a plant growth
chamber with cool white fluorescent lamps (CLE-305, TOMY, Tokyo, Japan) set to 22°C with
light level 4 and a 12:12 day:night ratio. A HOBO temperature and light logger (Onset, MA,
U.S.A.) was kept in the growth chamber during the experiment. The daytime temperature was
21°C with about 1900-2000 Lux (~30 μmol m^{-2} s^{-1}) light intensity, and the nighttime temperature
was 22°C. Positions of replicates were rotated daily to prevent position in the chamber from
systematically affecting replicates. On days 1, 3, and 4, chlorophyll fluorescence was measured
at the middle of the light period by transferring 200 μL aliquots to a black 96-well plate
(ThermoFisher, MA, U.S.A) and recording fluorescence (excitation: 440 nm, emission: 685 nm)
with a Tecan Ultra Evolution microplate reader (Tecan, Switzerland). Exponential growth phase
was determined by comparing measured fluorescence to a growth curve for *Phaeocystis globosa*
CCMP 1528 grown under identical conditions. On day 4 of the first round, 1 mL of each
replicate was transferred to 45 mL of sterile L1 media and the experimental setup was repeated,
which allowed for adaptation to experimental culture conditions. Algal cells were harvested for
RNA extraction on day 4 of the second experimental culture round, when replicates were in
middle to late exponential growth phase (Supplemental Figure S1).
172
Prior to RNA extraction, each culture replicate was imaged with light microscopy (Olympus
CKX53) to ensure that colony-forming replicates were indeed producing colonies and that non-
colonial replicates were not (Supplemental Figure S2). Flagellates present in the colonial and
non-colonial replicates were actively swimming, suggesting that flagellates in this study were
scaled haploid flagellates, rather than scale-free diploid flagellates originating from disrupted
colonies, but neither flow-cytometry nor electron microscopy were performed (Rousseau et al.
2007). Colony-forming culture replicates were filtered through polytetrafluoroethylene (PTFE)
filters (pore size, 10 μm) (Millipore) under gentle vacuum. Colonies were visible by eye on the
filter surface and swimming flagellates were observed in the flow-through when viewed with
light microscopy. The non-colonial culture replicates were first filtered through 50 μm nylon
mesh to remove culture debris, and then filtered through 1.0 μm pore-size PTFE filters. No

flagellates were visible when the filtrates were viewed with light microscopy. Filters were
immediately flash frozen in liquid nitrogen and stored at -80°C until RNA extraction.
186
187 ***RNA Extraction, Library Preparation and Sequencing***
188
Total RNA was extracted from filters by following the manufacturer's protocols for the MoBio
PowerWater RNA extraction kit (Qiagen), including the optional initial heating step. Following
extraction, we assessed RNA quality and concentration. RNA extracts were diluted so that 10 ng
of RNA were used for each sample with the SMART-seq v4 Ultra Low Input RNA Kit
(Clonetech) along with 2 µL of a 1:10,000 dilution of External RNA Controls Consortium
(ERCC) spike-in mix 1 (Ambion), an internal quality control. The SMART-seq kit employs
poly-A priming to target eukaryotic mRNA and to reduce the amount of ribosomal and bacterial
RNA present in sequencing libraries. The quality and concentration of the resulting cDNA was
assessed before continuing with the manufacturer's protocols for the Nextera XT DNA Library
Prep Kit (Illumina). Finally, we checked cDNA fragment size before submitting libraries to the
Okinawa Institute of Science and Technology (OIST) DNA Sequencing Section for paired-end
150x150 bp sequencing across 8 lanes of an Illumina Hiseq4000 flow-cell.
201
202 ***Bioinformatic Processing and Quality Control***
203
Sequencing reads were processed with Trimmomatic software to remove adapter sequences and
to filter low-quality sequences (Bolger et al. 2014). Read quality was checked with FastQC
before and after trimming to ensure that adapters were removed (Andrews 2010). Remaining
reads were mapped to the ERCC reference sequences (Cronin et al. 2004) and mapped reads
were counted with RSEM software (Li & Collin, 2011). Counts were further analyzed in the R
statistical environment (R Core Team, 2013). Reads mapping to the ERCC reference sequences
were then removed from each sample with SAMtools (Li et al. 2009) and BEDTools (Quinlan &
Hall, 2010).
212
213 ***Transcriptome Assembly, Assessment, and Functional Annotation***
214
The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP, Keeling et al.
2014) assembled a transcriptome for *Phaeocystis sp.* CCMP2710, which groups with the
Phaeocystis globosa species complex in phylogenetic analyses (Supplemental Figure S3). Only
25% of our reads, however, mapped to this reference transcriptome. We therefore assembled a *de*
novo transcriptome for *Phaeocystis globosa* CCMP1528 to serve as a reference for read mapping
in this study. We used Trinity software for transcriptome assembly (Grabherr et al. 2013) and
dereplicated the transcriptome by removing reads with 95% similarity using CD-HIT-EST (Fu et
al. 2012). Bacterial contamination was removed by performing a blastn query against the NCBI
nucleotide database (downloaded March 2018, ncbi-blast v2.6.0+, Camacho et al. 2009) and
parsing results to identify and remove bacterial contigs. The final assembly was assessed for
completeness with Benchmarking Universal Single-Copy Orthologs (BUSCO v3, Simao et al.
2015) and results were compared with those for the MMETSP *Phaeocystis sp.* CCMP2710
transcriptome.
228

We annotated the CCMP1528 transcriptome using two different databases, Pfam (Finn et al.
2010) and KEGG (Kanehisa et al. 2016). Pfam annotation was performed with the dammit
software (Scott 2018), which wraps Transdecoder to translate transcriptome contigs to the
longest possible amino acid sequence (Haas et al. 2013), and HMMER to assign protein
homologs to sequences (Eddy 2011). After discarding annotations with e-values greater than 1E-
5, the annotation with the lowest e-value was selected for each contig. Gene Ontology (GO)
terms were assigned to Pfam annotations using the Gene Ontology Consortium's Pfam2GO
mapping (geneontology.org/external2go/pfam2go, version 07/14/2018, Mitchell et al. 2015).
KEGG annotation was performed with the GhostKOALA tool and translated amino acid
sequences (kegg.jp/ghostkoala, 05/21/2018, Kanehisa et al. 2016). Annotated K numbers were
then used to assign KEGG pathways by accessing the KEGG API
(kegg.jp/kegg/rest/keggapi.html, July 2018).
241
242 ***Differential Gene Expression Analysis***
243
Quality filtered sequences from each sample were mapped to the assembled *P. globosa*
CCMP1528 transcriptome and counted with RSEM software. Counts for each sample were
imported into the R statistical environment, where differential gene expression between colonial
and solitary culture replicates was tested with the DESeq function in the Bioconductor package
DESeq2 (Love et al. 2014). Genes that were differentially expressed were considered statistically
significant if the False Discovery Rate (FDR) adjusted p-value (padj) was less than 0.05.
250
251 ***Gene Set Enrichment Testing***
252
We identified GO terms enriched among significantly upregulated and downregulated genes by
applying a hypergeometric test in the R package GOstats (Falcon and Gentleman 2007). GOstats
accommodates user-defined GO annotations, which are necessary when studying non-model
organisms like *Phaeocystis*. Likewise, a hypergeometric test for significant enrichment of KEGG
pathways was applied using the enricher function from the R package ClusterProfiler (Yu et al.
2012). Because of the lower annotation rate, KEGG pathway enrichment was further investigated
by additionally applying linear model analysis with the kegga function in the R package edgeR
(Robinson et al. 2010). GO terms and KEGG pathways were considered significantly enriched
when the statistical test returned a p-value less than 0.05.
262
263 ***Genes associated with DMSP and DMS production***
264
Because *Phaeocystis* is a profusive producer of DMSP and DMS, we specifically queried our
dataset for recently discovered algal genes involved in DMSP production (DSYB) and its
cleavage to DMS and acrylate (Alma family genes). We performed blastp queries with curated
DSYB protein sequences (provided by Curson et al. 2018) and *E. huxleyi* and *Symbiodinium*
Alma family protein sequences downloaded from UniProt (July 2018) against *Phaeocystis*
globosa CCMP1528 amino acid sequences. Expression levels of putative *Phaeocystis globosa*
DSYB and *Alma* family genes were then checked in colonial and solitary culture replicates.
272
273 **Results**
274

275   *Bioinformatic Processing and Quality Control*
276
Sequencing for this project produced over 1.9 billion read pairs with 159-383 million read pairs
per sample. Following quality filtering with Trimmomatic, 1.7 billion read pairs remained, with
140-341 million read pairs per sample (Supplemental Table S1). After mapping reads from each
sample to ERCC reference sequences, we plotted the Log_2 FPKM for each sequence against the
Log_2 of its concentration in the standard mix. A simple linear regression was fitted for each
sample and R^2 values ranged from 0.93-0.937 for each sample (Supplementary Figure S4). The
strong correlation between observed FPKM and initial concentration for ERCC sequences
indicates that minimal bias was introduced during PCR amplification, library preparation, and
sequencing.
286
287   *Transcriptome Assembly, Assessment, and Functional Annotation*
288
The final assembly of the *Phaeocystis globosa* CCMP1528 transcriptome included 69,528
contigs and a total of 43.9 Mbp. The CCMP1528 transcriptome was about 3 times larger than the
MMETSP CCMP2710 transcriptome, but the minimum, maximum, and mean contig lengths
were about the same for both (Supplemental Table S2). When Transrate was used to align the
two transcriptomes, only 18% of CCMP1528 contigs aligned to the CCMP2710 transcriptome,
but 55% of the CCMP2710 contigs aligned to the CCMP1528 transcriptome. BUSCO software
was utilized to assess completeness of the *P. globosa* CCMP1528 transcriptome. It included
more complete eukaryote and protist BUSCOs than the MMETSP *Phaeocystis sp.* CCMP2710
transcriptome (Figure 1). Together, these results demonstrate that the transcriptome generated in
this study is more complete than the MMETSP transcriptome and is a better reference for this
study. The results also indicate that CCMP2710 and CCMP1528 are more genetically distant
than was expected based upon ribosomal RNA gene sequences (Supplemental Figure S3).
301

302
Figure 1. Percent of eukaryote and protist Benchmarking Universal Single Copy Orthologs (BUSCOs)
complete, fragmented, or missing in *Phaeocystis globosa* CCMP1528 and *Phaeocystis sp.* CCMP2710
transcriptomes. BUSCO software was used to determine the percent of eukaryotic and protistan BUSCOs
represented by complete single copies, complete but duplicated copies, copies that were fragmented or missing in
the *Phaeocystis globosa* CCMP1528 transcriptome assembled for this study and the *Phaeocystis sp.* CCMP2710
transcriptome assembled for the Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP). More

eukaryotic and protistan BUSCOs were represented in the *Phaeocystis globosa* CCMP1528 transcriptome than the
Phaeocystis sp. CCMP2710 transcriptome. The plot was rendered with the R package ggplot2.
311
312 Annotation was possible for relatively few of the contigs in the *P. globosa* CCMP1528
313 transcriptome assembly, but more genes were annotated with the Pfam and GO annotation
314 pipeline (26%) than with the KEGG pipeline (14%). Additionally, both annotation methods
315 annotated the significantly differentially expressed (DE) genes at a higher rate than the whole
316 transcriptome (Table 1).
317

	Pfam / GO	KEGG
Total genes annotated	17,826 (26%)	9,967 (14%)
Total w. pathway or GO	8,962 (13%)	5,585 (8%)
DE genes annotated	5,180 (66%)	3,305 (43%)
DE genes w. pathway or GO	2,764 (35%)	1,883 (24%)

318
319 **Table 1. Pfam, Gene Ontology (GO), and Kyoto Encyclopedia of Genes and Genomes (KEGG) annotation**
320 **statistics for the *Phaeocystis globosa* CCMP1528 transcriptome assembly and differentially expressed (DE)**
321 **genes.** Pfam annotations were assigned with dammit software and corresponding GO terms were determined with
322 Pfam2GO mapping provided by the GO Consortium. KEGG K numbers were assigned with GhostKOALA tool and
323 KEGG pathways were determined by accessing the KEGG API.
324
325 *Differential Gene Expression Analysis*
326
Gene expression patterns in colonial and solitary replicates were explored with a principal
component analysis (PCA) and an expression heatmap. Initial data exploration revealed the
colonial replicate 'C2' as an outlier to other colonial replicates and solitary replicates
(Supplemental Figures S5A, B) and this sample was excluded from further analyses. The
remaining 3 colonial replicates clustered separately from the 4 solitary replicates in a PCA plot
(Figure 2A). Differential expression analysis identified 535 genes as significantly upregulated
and 7,234 genes as significantly downregulated in colonial replicates. An expression heatmap of
the most differentially expressed genes sorted by FDR adjusted p-value (padj) clearly illustrates
the overall expression pattern—the majority of significantly differentially expressed genes are
downregulated in colonial replicates (Figure 2B).
337

338
Figure 2. Principal component analysis (PCA) and heatmap demonstrating gene expression patterns in
colonial and solitary *Phaeocystis globosa*. A. PCA performed on distances between samples derived from
regularized log transformed counts. Colonial and solitary replicates cluster separately, and the majority of variance
is between sample type rather than within replicates. Results plotted with R package ggplot2. **B.** Heatmap includes
the 1000 significantly differentially expressed genes with the lowest FDR adjusted p-values. Heatmap color
represents difference from the mean regularized log transformed count for each contig in each sample. The majority
of differentially expressed genes are downregulated in colonial replicates, and replicates cluster by sample type.
Results plotted with R package pheatmap.

347
348 *Gene Set Enrichment Analysis*
349

In order to identify Biological Process (BP) GO terms over-represented in significantly up- and
downregulated gene sets, we applied a hypergeometric test with a significance cut-off of p <
0.05. Twenty BP GO term were over-represented among significantly upregulated genes and
were primarily involved in cell signal transduction in response to external stimuli (Figure 3,
Supplemental Table S3). Notably, GO terms involving arabinose, a component of the colonial
matrix, were also enriched among upregulated genes. In the downregulated gene set, 48 BP GO
terms were enriched, including several involved in cation transport, response to oxygen-
containing compounds, translation and protein transport, and vacuolar transport and exocytosis
(Figure 3, Supplemental Table S4). REVIGO software was used to remove redundant GO terms
from lists of enriched terms and to visualize results in a Multidimensional Scaling (MDS) plot
(Figure 3) based on GO term semantic similarities (Supek et al. 2011) as determined by shared
ancestry (Pesquita et al. 2009).
362

363
364
**Figure 3. Multidimensional scaling plot of semantic similarities between non-redundant GO terms over-
represented in significantly up- and downregulated gene sets.** Analysis performed using the REVIGO tool
(http://revigo.irb.hr/) with the allowed similarity set to 0.7 (to remove redundant GO terms) and the SimRel metric
selected to calculate similarities. REVIGO results were exported to the R statistical environment and plotted with
ggplot2. Representative GO terms were manually selected and labeled on the plot. To view all GO term labels, an
interactive version of the plot made with R package ggplotly is available at:
https://brisbin.shinyapps.io/shinycolsol/.

When we applied hypergeometric testing to KEGG pathways, only the cGMP-PKG signaling
pathway (cyclic guanosine monophosphate-protein kinase G pathway) was enriched in the
upregulated gene set (p<0.05) (Supplemental Table S5). Five pathways were enriched among
downregulated genes: Lysosome, Autophagy, MAPK signaling pathway (mitogen activated
protein kinase signaling pathway), AMPK signaling pathway (adenosine monophosphate-
activated protein kinase signaling pathway), and Epidermal growth factor receptor (EGFR)
tyrosine kinase inhibitor resistance (Supplemental Table S6). The lower annotation rate for
KEGG pathways compared with GO terms contributed to the difference in enrichment testing
results. We therefore also applied a linear model test for KEGG pathway enrichment, which
identified several additional pathways as being significantly enriched in the up- (6) and
downregulated (7) gene sets. With this additional test, the PI3K-Akt signaling (phosphoinositide
3-kinase-protein kinase B signaling pathway), Glycosphingolipid biosynthesis, Ferroptosis,
Plant-pathogen interaction, Circadian rhythm, Viral carcinogenesis pathways were also enriched
among upregulated genes (Supplemental Table S7). The Protein processing in endoplasmic
reticulum, Oxidative phosphorylation, Ras signaling pathway, Sphingolipid metabolism, Steroid
biosynthesis, Fatty acid degradation, and Taste transduction pathways were additionally
identified as enriched in downregulated genes (Supplemental Table S8).

390
391 ***Genes associated with DMSP and DMS production***
392
A blastp query against curated DSYB protein sequences from Curson et al. (2018) and Alma
family protein sequences from Alcolombri et al. (2015) identified 4 *Phaeocystis globosa* contigs
as putative *DSYB* or *Alma* family genes (Table 2). The *P. globosa* amino acid (AA) sequence
translated from Transcript_30752 aligned with the sequence for *Prymnesium parvum*
CCAP946/1B DSYB protein, which is experimentally proven to be highly active. It is therefore
likely that this gene is actively involved in DMSP biosynthesis in *Phaeocystis globosa*. A second
P. globosa AA sequence, from Transcript_31221, aligned with the *Pseudonitzchia fraudulenta*
DYSB protein sequence, making it also a possible *DSYB* gene. The *Pseudonitzchia* DSYB
protein has not been experimentally proven to be active, but its sequence is phylogenetically
close to the *Fragillariopsis* DSYB, which has been proven to be active. Neither putative *P.*
globosa DSYB genes were differentially expressed between solitary and colonial culture
replicates in this study, but both were expressed at relatively high levels in both sample types
(Figure 4).
406
Two *Alma* family-like genes were identified in the *Phaeocystis globosa* transcriptome. One *P.*
globosa AA sequence, from Transcript_36000, aligned with *Emiliania huxleyi* Alma7. All 4
Alma homologs identified from the MMETSP *Phaeocystis antarctica* transcriptome are
phylogenetically closest to the *E. huxleyi Alma7*, but *E. huxleyi* Alma7 has not been proven to
have DMSP-lyase activity. Another *P. globosa* AA sequence, from Transcript_68879, aligned
with *E. huxleyi* Alma4, which also has not been experimentally proven active. Both putative *P.*
globosa Alma family genes were expressed at lower rates than *DSYB*-like genes. The *P. globosa*
Alma4-like gene was not differentially expressed in solitary and colonial culture replicates in this
study (Figure 4). The *P. globosa Alma7*-like gene, however, was significantly upregulated in
colonial replicates (padj = 1.44E-12, logFC = 3.55), suggesting that DMSP biosynthesis is
occurring in both colonial and solitary cells, but colonial cells may be cleaving DMSP to DMS
and acrylate more actively than solitary cells.
419

	Database Sequence	%ID	Alignment Length (AAs)	Gaps	E-value	Bit Score
Transcript_30752	*Prymnesium parvum* CCAP946/1B DSYB	74	314	2	1.15E-170	472
Transcript_31221	*Pseudonitzschia fraudulenta* WWA7 DSYB	45	263	8	1.92E-65	210
Transcript_36000	*Emiliania huxleyi* Alma7	61	272	4	3.70E-119	336
Transcript_68879	*Emiliania huxleyi* Alma4	40	253	4	2.64E-48	166

420 **Table 2. Top *Phaeocystis globosa* CCMP1528 blastp results against *DSYB* and *Alma* family reference**
421 **sequences.** The *Phaeocystis globosa* CCMP1528 transcriptome assembly was queried against curated DSYB
422 reference sequences (Curson et al. 2018) and Alma family reference sequences from *Emiliania huxleyi* and
423 *Symbiodinium* (Alcolombri et al. 2015). Blastp hits with e-values < 1E-30 are included in the table.

424
425 **Figure 4. Normalized expression levels of *DSYB*-like and *Alma*-like genes in *Phaeocystis globosa* colonial and
426 solitary cell cultures.** Box plots show the range, quartiles and median of the log normalized counts for each gene in
427 colonial and solitary culture replicates. Only the *E. huxleyi Alma7*-like gene was significantly differentially
428 expressed and was upregulated in colonial samples (padj = 1.44E-12, log fold-change = 3.55) and is represented
429 with an asterisk (*) on the plot. Plots made with R package ggplot2.
430
431 **Discussion**
432
Colonial *Phaeocystis* blooms widely impact ecosystem function and can be extremely
detrimental in some systems, particularly in subtropical and tropical coastal regions. Although
many hypotheses exist, factors initiating colonial blooms and the ecological function of colonial
formation remain enigmatic. We investigated gene expression associated with colony formation
in a warm-water colony-forming strain of *Phaeocystis globosa* to identify cellular processes
associated with colony formation and potentially provide clues for what initiates colony
formation and the functional role of colonies in the *Phaeocystis* life-cycle. Overall, we observed
a transcriptional shift in colonial cultures compared to solitary cell cultures, with vastly more
genes significantly downregulated in colonial cells than upregulated (Figure 2). This shift
suggests that there are trade-offs associated with colony production and resources must be
diverted to construct and maintain the colonial matrix. A relatively small number of genes are
upregulated to produce colonies, but the low annotation rate of these genes, and the
transcriptome overall, make it challenging to fully interpret the results (Table 1). Gene set
enrichment analyses inherently relies on how many and which genes are annotated and
systematic biases likely in gene annotation will influence results (Haynes et al. 2018). However,
these analyses still assist in identifying pathways and functions that may be important to the
question at hand and indicate genes and pathways that should be followed up in future studies.
The results presented here highlight genes involved in constructing the colonial matrix, changes

in cellular morphology, responding to external stimuli, cellular proliferation, and producing
DMSP, DMS, and acrylate. Results from this study support a defensive role for colony formation
in *Phaeocystis globosa*.
454
455 ***Colony matrix carbohydrates and colonial cell morphology***
456
Differential expression of genes associated with clearly observable changes between treatment
groups can serve to "ground-truth" results from RNA-seq experiments and therefore increase
confidence in expression changes detected for genes for less observable traits. In this study,
changes in cellular morphology and colony formation itself are clearly observable differences for
which several associated genes are differentially expressed. The observed expression patterns for
these genes can additionally provide new insight into the construction of the colonial matrix and
pathways associated with morphological changes in colonial *Phaeocystis* cells. Many different
polysaccharides are recognized as contributors to the matrix structure of *Phaeocystis globosa*
colonies, including arabinose, rhamnose, xylose, mannose, galactose, glucose, gluconuronate,
and O-methylated pentose sugars (Janse et al. 1996). *Phaeocystis* isolated from different
locations tends to have distinct matrix carbohydrate fingerprints, which may be due to genetic
attributes of different strains or which could arise from different environmental conditions, such
as light or nutrient availability. For example, arabinose is the most abundant matric carbohydrate
in *P. globosa* sampled from the North Sea (Janse et al. 1996). In this study, the GO term for
Arabinose metabolic process was enriched among upregulated genes in colonial cells (Figure 3).
These results indicate that arabinose is likely the dominant matrix polysaccharide in *P. globosa*
CCMP1528 and that arabinose production is specifically associated with colony formation in this
strain. The colonial matrix also contains nitrogen (Hamm 2000), which is likely included in
amino sugars (Solomon et al. 2003). Our results, however, did not indicate that amino sugar
biosynthesis or metabolism was upregulated in colonial cells.
477
Divalent cations, particularly Mg^{2+} or Ca^{2+}, are required for colonial polymers to gel and
contribute to the stability of the colonial matrix (van Boekel 1992). GO terms for Divalent
Inorganic cation transport, Magnesium ion transport, and Divalent metal ion transport, however
were enriched in downregulated genes in colonial replicates (Figure 3). Similarly, Bender et al.
(2018) found that *Phaeocystis antarctica* produces more calcium-binding proteins when iron
limitation decreases colony formation. These results may be due to the importance of divalent
cations for flagellate motility. Actively swimming *Phaeocystis globosa* flagellates, as observed
in this study, may require continuous transport of divalent cations to the point of masking their
shared importance in colony formation. Calcium signaling also induces secretion of vesicles
containing gel forming polymers (Chin et al. 2004), which further confounds the observed
downregulation of divalent cation transport genes in colonies. However, flagellates also secrete
vesicles, but instead of gel polymers they contain star-shaped structures composed of chitinous
filaments (Chretiennot-Dinet et al. 1997). The exact function of these structures is unknown, but
they may be involved in mating or defense (Dutz and Koski 2006). In addition to divalent cation
transport, a number of other GO-terms enriched in downregulated genes may be involved in
secreting these structures, such as Exocytosis, Secretion, Vesicle mediated transport, and
Vacuolar transport (Figure 3). Alternatively, these GO terms may be involved in scale formation
and secretion (Taylor et al. 2007), as scales are only observed on *Phaeocystis globosa* flagellates
and not colonial cells (Rousseau et al. 2007).

497
498 ***A defensive role for colony formation: resource allocation, pathogen interaction, and***
499 ***DMS/acrylate production***
500
Out of 7,769 genes that were significantly differentially expressed between colonial and solitary
replicates, 7,234 genes were downregulated in colonial cells. This dramatic transcriptional shift
in colonial cells supports a high resource cost associated with producing colonies (Wang et al.
2015). Specifically, our results indicate that resources are being diverted from protein translation
and transport and cell division in order to produce the colonial matrix. Several GO terms
involved in the synthesis of larger nitrogenous compounds and their transport, including
Translation initiation, Protein metabolic process, Protein N-linked glycosylation, and Protein
transport were significantly enriched in downregulated genes in colonial cells (Figure 3,
Supplemental Table S4). Similarly, the KEGG pathway, Protein processing in endoplasmic
reticulum, was also enriched in downregulated genes in colonies (Supplemental Table S7).
Likewise, several mitosis-associated GO terms (Chromatin assembly and disassembly,
Cytoskeleton organization, Cellular component biogenesis) were also enriched among
downregulated genes in colonial cells (Figure 3, Supplemental Table S4). However, the
downregulation of mitosis-associated genes in colonial cells conflicts with observations in
previous studies. Veldhius et al. (2005) observed that colonial cells divide at a higher rate than
solitary cells and proposed that in addition to experiencing less grazing and viral lysis, colonial
cells may dominate blooms because they outgrow solitary cells. We believe the difference in our
results may be due to the type of solitary cells observed—the solitary cells in previous studies
could have been diploid flagellates, especially if they were derived from disrupted colonies,
whereas solitary cells in our study are likely haploid flagellates, which have been reported to
divide extremely rapidly (Rousseau et al. 2007).
522
There were also several signaling pathways represented in the results suggesting that colonial
cells are exposed to fewer general stressors, but may be responding to more strongly to specific
pathogens. In plants, the MAPK (mitogen activated protein kinase) pathway primarily transduces
signals from extracellular stressors to the nucleus or cytoplasm and initiates an appropriate
response (Taj et al. 2010). In our results, the MAPK signaling pathway was significantly
enriched in genes downregulated in colonies. The downregulated genes in this pathway encode
MAP3Ks, MAP2Ks, and MAPKs, which are activated in response to pathogen attack and
infection, phytohormones, cold and salt stress, and reactive oxygen species (Taj et al. 2010).
Downregulation of genes associated with stress response is also evidenced by related GO terms
enriched among downregulated genes, specifically Response to wounding and Response to
oxygen-containing compounds. These results support the hypothesis that colony formation
serves a defensive purpose. Defense responses regulated by the MAPK pathway are unneeded
because the colony skin is protecting cells from these stressors. However, the Plant-pathogen
interaction pathway was enriched in upregulated genes in colonial cells, indicating that specific
pathogens may penetrate the colonial fortress or that pathogen interaction may play a role in
stimulating colony formation. Genes upregulated in this pathway were for calcium-dependent
protein kinases and calcium-binding protein CML (calmodulin-like protein), immune response
genes that are activated following recognition of specific pathogen-associated molecular patterns
(Cheval et al. 2013). Specific bacterial interactions are known to influence transitions between
life-cycle stages in several other protists: specific bacteria stimulate growth in marine diatoms

(Amin et al. 2015) and specific bacterial signaling molecules are responsible for inducing both
colony formation (Woznica et al. 2016) and sexual reproduction (Woznica et al. 2017) in
choanoflagellates, another single-celled colony-forming marine plankton. In *Phaeocystis*, axenic
cultures exhibit decreased growth rates (Solomon et al. 2003), but the effects of specific bacteria
on colony formation have not yet been investigated.
548
Phaeocystis is a copious producer of DMSP and its cleavage products, DMS and acrylate. DMS
and acrylate have been indicated as grazer-deterrents and antimicrobials (Wolfe and Steinke
1996, Hamm 2000, Noordkamp et al. 2000). While algal genes associated with DMSP
biosynthesis (*DSYB*, Curson et al. 2018) and cleavage (*Alma* family genes, Alcolombri et al.
2015) have been identified in many algal transcriptomes, including *Phaeocystis antarctica*, this
study is the first to identify these genes for *Phaeocystis globosa*. We found that colonial and
solitary *P. globosa* expressed *DSYB*-like genes at the similar levels, suggesting that the two cell
types produce similar amounts of DMSP. *DSYB* expression in *Prymnesium parvum*, the
haptophyte in which DSYB was discovered, is affected only by salinity, potentially indicating
that DMSP production functions primarily in osmoregulation rather than as a defensive or stress
response (Curson et al. 2018). Similar *DSYB*-like gene expression levels in colonial and solitary
P. globosa cells support a basic, shared function for DMSP in the two cell types. Contrastingly,
an *Alma* family gene was upregulated in colonial cells. Acrylate accumulates in *Phaeocystis*
colonies and may serve to deter grazers and pathogens from disrupting the colonial matrix
(Noordkamp et al. 2000). While acrylate may accumulate in colonies simply because it cannot
escape through the colonial skin, the upregulation of an *Alma*-like gene in colonies suggests that
colonial *Phaeocystis* cells may actively produce excess acrylate and DMS than solitary cells,
further supporting a defensive role for colony production in *P. globosa*.
567
568 ***Role of colonies in* Phaeocystis *reproduction***
569
Colony formation is believed to be involved in sexual reproduction in *Phaeocystis* since
swarming flagellates have been observed within senescent colonies (Peperzak et al. 2000,
Rousseau et al. 2013). However, we did not find meiosis or sexual reproduction GO terms or
KEGG pathways enriched in up- (or down-) regulated genes in this study. These results may
arise from RNA being extracted during mid- to late exponential growth phase. Previous
observations suggest that colonies produce flagellates during bloom decay, so we might have
found meiosis genes upregulated in colonial cells if we had sampled toward late stationary phase
instead of exponential phase.
578
579 ***Signaling pathways associated with colony formation***
580
Processes and pathways involved in cell-signaling, cell communication, and response to stimuli
that are enriched in upregulated-genes are particularly interesting because they shed some light
on factors stimulating colony formation in *Phaeocystis globosa*. The cGMP-PKG signaling
pathway was the only KEGG pathway significantly enriched among upregulated genes when a
hypergeometric test was used. Three genes in this pathway were upregulated: 1) cGMP-
dependent protein kinase, which phosphorylates biologically important targets, has been
implicated in cell division and nucleic acid synthesis, and reduces cytoplasmic Ca^{2+}
concentrations (Lincoln et al. 2001); 2) cAMP-dependent protein kinase regulator; and 3) a

589 cAMP-responsive element-binding protein (CREB), which binds to DNA to increase or decrease
590 transcription and is associated with increased cell survival (Chrivia et al. 1993). The PI3K-Akt
591 signaling pathway was also significantly enriched in upregulated genes when the additional
592 linear model test was used. Within this pathway, two Extracellular Matrix (ECM) focal adhesion
593 genes, for Tenascin (a glycoprotein) and Type IV collagen, were significantly upregulated. Focal
594 adhesion proteins connect cells to extracellular matrices both literally and figuratively, by
595 holding cells in place and by initiating cellular responses to external conditions (Wozniak et al.
596 2004). Bender et al. (2018) also found focal adhesion proteins, specifically glycoproteins,
597 upregulated in colonial *Phaeocystis antarctica*. It is therefore likely that these proteins have an
598 important function in structurally maintaining cell positions in the colonial matrix and signaling
599 between colonial cells. Focal adhesion proteins may be mediating interactions with protein
600 kinases in colonial cells, which go on to promote cell proliferation and differentiation into the
601 colonial morphotype. These signaling pathways represent important candidates for continued
602 study of molecular mechanisms regulating colony formation.
603
604 ***Conclusions and Future Directions***
605
606 This study investigated gene expression associated with colony formation in *Phaeocystis globosa*
607 for the first time and discovered a large transcriptional shift associated with colony production.
608 Differentially expressed genes were mostly downregulated in colonies, providing evidence for
609 extensive resource allocation toward colony formation. Together, activation of pathogen
610 interaction pathways, reduced expression of stress-response pathways, and increased expression
611 of a DMSP-lyase, which produces DMS and acrylate, supporting a defensive role for colony
612 formation. Future studies may extend this work by investigating *P. globosa* gene expression in
613 colonial and solitary cells in a time course study through the waxing and waning of a bloom and
614 under different nutrient and grazing regimes, potentially by using mesocosms or
615 metatranscriptomic methods in natural communities. While our ability to fully interpret the
616 results was inhibited by an overall lack of annotated genomes and transcriptomes for diverse
617 protist lineages, this study represents a step in the right direction by contributing a new and
618 deeply sequenced transcriptome for *Phaeocystis globosa*. Identification of *DSYB* and *Alma*
619 family-like genes in this transcriptome will additionally allow for further investigation into *P.*
620 *globosa* DMSP and DMS production in the oceans. We were also able to identify several protein
621 kinase signaling pathways that are potentially important for regulating colony formation and
622 should be experimentally investigated in follow-up studies. The results presented here will guide
623 and facilitate continued efforts to unravel the complex factors responsible for triggering harmful
624 colonial *Phaeocystis* blooms, which will likely increase with continued climate change and
625 nutrient pollution in the future.
626
627 **Data Availability**
628
629 Sequence data is available in the NCBI Sequence Read Archive (SRA) with accession numbers
630 SRR7811979–SRR7811986. The *Phaeocystis globosa* CCMP1528 *de novo* transcriptome
631 produced and used in this study, data files, and data analysis scripts can be accessed at
632 https://doi.org/10.5281/zenodo.147649.
633
634 **Acknowledgements**

This work was supported by funding from the Marine Biophysics Unit of the Okinawa Institute of Science and Technology Graduate University. MMB is supported by a JSPS DC1 graduate student fellowship. We thank Angela Ares Pita for valuable discussion and suggestions and for helpful feedback on manuscript drafts. Hiroki Goto and the OIST sequencing section provided important guidance for RNA library preparation and other sequencing considerations. Steven D. Aird edited the manuscript and provided helpful comments.

Works Cited

Alcolombri, U., Ben-dor, S., Feldmesser, E., Levin, Y., Tawfik, D. S., and Vardi, A. (2015). Identification of the algal dimethyl sulfide-releasing enzyme: a missing link in the marine sulfur cycle. *Science (80)*. 348, 1–4.

Andrews S. (2010) *FastQC: a quality control tool for high throughput sequence data*. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

Amin, S. A., Hmelo, L. R., van Tol, H. M., Durham, B. P., Carlson, L. T., Heal, K. R., et al. (2015). Interaction and signalling between a cosmopolitan phytoplankton and associated bacteria. *Nature* 522, 98–101. doi:10.1038/nature14488.

Andersen, R. A., Bailey, J. C., Decelle, J., and Probert, I. (2015). *Phaeocystis rex* sp. nov. (Phaeocystales, Prymnesiophyceae): a new solitary species that produces a multilayered scale cell covering. *Eur. J. Phycol.* 50, 207–222. doi:10.1080/09670262.2015.1024287.

Bender, S. J., Moran, D. M., McIlvin, M. R., Zheng, H., McCrow, J. P., Badger, J., et al. (2018). Iron triggers colony formation in *Phaeocystis antarctica*: connecting molecular mechanisms with iron biogeochemistry. *Biogeosciences*, 1–52. doi:10.5194/bg-2017-558.

Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30, 2114–2120. doi:10.1093/bioinformatics/btu170.

Brussaard, C. P. D., Bratbak, G., Baudoux, A. C., and Ruardij, P. (2007). *Phaeocystis* and its interaction with viruses. *Phaeocystis, Major Link Biogeochem. Cycl. Clim. Elem.*, 201–215. doi:10.1007/978-1-4020-6214-8_15.

Brussaard, C. P. D., Kuipers, B., and Veldhuis, M. J. W. (2005). A mesocosm study of *Phaeocystis globosa* population dynamics: I. regulatory role of viruses in bloom control. *Harmful Algae* 4, 859–874. doi:10.1016/j.hal.2004.12.015.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., et al. (2009). BLAST+: architecture and applications. *BMC Bioinformatics* 10, 1–9. doi:10.1186/1471-2105-10-421.

Cariou, V., Casotti, R., Birrien, J. L., and Vaulot, D. (1994). The initiation of *Phaeocystis* colonies. *J. Plankton Res.* 16, 457–470. doi:10.1093/plankt/16.5.457.

Caron, D. A., Alexander, H., Allen, A. E., Archibald, J. M., Armbrust, E. V., Bachy, C., et al. (2017). Probing the evolution, ecology and physiology of marine protists using transcriptomics. *Nat. Rev. Microbiol.* 15, 6–20. doi:10.1038/nrmicro.2016.160.

Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G. (1987). Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate. *Nature* 326, 655–661. doi:10.1038/326655a0.

Chen, Y. Q., Wang, N., Zhang, P., Zhou, H., and Qu, L. H. (2002). Molecular evidence identifies bloom-forming *Phaeocystis* (Prymnesiophyta) from coastal waters of southeast China as Phaeocystis globosa. *Biochem. Syst. Ecol.* 30, 15–22. doi:10.1016/S0305-1978(01)00054-0.

Cheval, C., Aldon, D., Galaud, J.-P., and Ranty, B. (2013). Calcium/calmodulin-mediated regulation of plant immunity. *Biochim. Biophys. Acta - Mol. Cell Res.* 1833, 1766–1771. doi: 10.1016/j.bbamcr.2013.01.031.

Chin, W. C., Orellana, M. V., Quesada, I., and Verdugo, P. (2004). Secretion in unicellular marine phytoplankton: demonstration of regulated exocytosis in Phaeocystis globosa. *Plant Cell Physiol.* 45, 535–542. doi:10.1093/pcp/pch062.

Chrétiennot-Dinet, M.-J., Giraud-Guille, M.-M., Vaulot, D., Putaux, J.-L., Saito, Y., and Chanzy, H. (1997). The chitinous nature of filaments ejected by *Phaeocystis* (Prymnesiophyceae). *J. Phycol.* 33, 666–672.

Chrivia, J. C., Kwok, R. P. S., Lamb, N., Hagiwara, M., Montminy, M. R., and Goodman, R. H. (1993). Phosphorylated CREB binds specifically to the nuclear protein CBP. *Nature* 365, 855.

Cronin, M., Ghosh, K., Sistare, F., Quackenbush, J., Vilker, V., and O'Connell, C. (2004). Universal RNA reference materials for gene expression. *Clin. Chem.* 50, 1464–1471. doi:10.1373/clinchem.2004.035675.

Curson, A. R. J., Williams, B. T., Pinchbeck, B. J., Sims, L. P., Martínez, A. B., Rivera, P. P. L., et al. (2018).
DSYB catalyses the key step of dimethylsulfoniopropionate biosynthesis in many phytoplankton. *Nat.*
Microbiol. 3, 430–439. doi:10.1038/s41564-018-0119-5.
Desroy, N., and Denis, L. (2004). Influence of spring phytodetritus sedimentation on intertidal macrozoobenthos in
the eastern English Channel. *Mar. Ecol. Prog. Ser.* 270, 41–53. doi:10.3354/meps270041.
Doan-Nhu, H., Nguyen-Ngoc, L., and Dippner, J. W. (2010). Development of *Phaeocystis globosa* blooms in the
upwelling waters of the South Central coast of Vietnam. *J. Mar. Syst.* 83, 253–261.
doi:10.1016/j.jmarsys.2010.04.015.
Dutz, J., and Koski, M. (2006). Trophic significance of solitary cells of the prymnesiophyte *Phaeocystis globosa*
depends on cell type. *Limnol. Oceanogr.* 51, 1230–1238. doi:10.4319/lo.2006.51.3.1230.
Eddy, S. R. (2011). Accelerated Profile HMM Searches. *PLOS Comput. Biol.* 7, e1002195.
doi:10.1371/journal.pcbi.1002195.
Falcon, S., and Gentleman, R. (2007). Using GOstats to test gene lists for GO term association. *Bioinformatics* 23,
257–258. doi:10.1093/bioinformatics/btl567.
Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. (2012). CD-HIT: Accelerated for clustering the next-generation
sequencing data. *Bioinformatics* 28, 3150–3152. doi:10.1093/bioinformatics/bts565.
Gaebler-Schwarz, S., Davidson, A., Assmy, P., Chen, J., Henjes, J., Nöthig, E. M., et al. (2010). A new cell stage in
the haploid-diploid life cycle of the colony-forming haptophyte *Phaeocystis antarctica* and its ecological
implications. *J. Phycol.* 46, 1006–1016. doi:10.1111/j.1529-8817.2010.00875.x.
Grabherr, M. G ., Brian J. Haas, Moran Yassour Joshua Z. Levin, Dawn A. Thompson, Ido Amit, Xian Adiconis,
Lin Fan, Raktima Raychowdhury, Qiandong Zeng, Zehua Chen, Evan Mauceli, Nir Hacohen, Andreas Gnirke,
Nicholas Rhind, Federica di Palma, Bruce W., N., and Friedman, and A. R. (2013). Trinity: reconstructing a
full-length transcriptome without a genome from RNA-Seq data. *Nat. Biotechnol.* 29, 644–652.
doi:10.1038/nbt.1883.Trinity.
Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Philip, D., Bowden, J., et al. (2013). De novo transcript
sequence reconstruction from RNA-seq: reference generation and analysis with Trinity. *Nat Protoc.* 8, 1–43.
doi:10.1038/nprot.2013.084.De.
Hamm, C. E. (2000). Architecture, ecology and biogeochemistry of *Phaeocystis* colonies. *J. Sea Res.* 43, 307–315.
doi:10.1016/S1385-1101(00)00014-9.
Haynes, W. A., Tomczak, A., and Khatri, P. (2018). Gene annotation bias impedes biomedical research. *Sci. Rep.* 8,
1–7. doi:10.1038/s41598-018-19333-x.
Houdan, A., Billard, C., Marie, D., Not, F., Sâez, A. G., Young, J. R., et al. (2004). Holococcolithophore-
heterococcolithophore (Haptophyta) life cycles: Flow cytometric analysis of relative ploidy levels. *Syst.*
Biodivers. 1, 453–465. doi:10.1017/S1477200003001270.
Janse, I., Van, R., M, G., Jc, L., C, G., and Wwc (1996). Carbohydrates in the North Sea during spring bloom of
Phaeocystis: A specific fingerprint. *Aquat Microb Ecol* 10, 97–103.
Kanehisa, M., Sato, Y., and Morishima, K. (2016). BlastKOALA and GhostKOALA: KEGG tools for functional
characterization of genome and metagenome sequences. *J. Mol. Biol.* 428, 726–731.
doi:10.1016/j.jmb.2015.11.006.
Keeling, P. J., Burki, F., Wilcox, H. M., Allam, B., Allen, E. E., Amaral-Zettler, L. a., et al. (2014). The Marine
Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): illuminating the functional diversity of
eukaryotic life in the oceans through transcriptome sequencing. *PLoS Biol.* 12, e1001889.
doi:10.1371/journal.pbio.1001889.
Li, B., and Dewey, C. N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a
reference genome. *BMC Bioinformatics* 12, 323. doi:10.1186/1471-2105-12-323.
Lincoln, T. M., Dey, N., and Sellak, H. (2001). Signal transduction in smooth muscle invited review: cGMP-
dependent protein kinase signaling mechanisms in smooth muscle: from the regulation of tone to gene
expression. *J. Appl. Physiol.* 91, 1421–1430. doi:10.1152/japplphysiol.01002.2001.
Liss, P. S., Malin, G., Turner, S. M., and Holligan, P. M. (1994). Dimethyl sulphide and *Phaeocystis*: A review. *J.*
Mar. Syst. 5, 41–53. doi:10.1016/0924-7963(94)90015-9.
Liu, H.-X., Huang, H.-H., Xu, S.-N., Dai, M., and Shen, P.-P. (2015). Planktonic community structure during a
harmful bloom of Phaeocystis globosa in a subtropical bay, with special reference to the ciliate assemblages.
Ecotoxicology. doi:10.1007/s10646-015-1464-2.
Long, J. D., Smalley, G. W., Barsby, T., Anderson, J. T., and Hay, M. E. (2007). Chemical cues induce consumer-
specific defenses in a bloom-forming marine phytoplankton. *Proc. Natl. Acad. Sci.* 104, 10512–10517.
doi:10.1073/pnas.0611600104.

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol.* 15, 1–21. doi:10.1186/s13059-014-0550-8.
Mitchell, A., Chang, H.-Y., Daugherty, L., Fraser, M., Hunter, S., Lopez, R., et al. (2015). The InterPro protein
families database: the classification resource after 15 years. *Nucleic Acids Res.* 43, D213–D221. Available at:
http://dx.doi.org/10.1093/nar/gku1243.
Noordkamp, D. J. B., Gieskes, W. W. C., Gottschal, J. C., Forney, L. J., and Van Rijssel, M. (2000). Acrylate in
Phaeocystis colonies does not affect the surrounding bacteria. *J. Sea Res.* 43, 287–296. doi:10.1016/S1385-
1101(00)00021-6.
Peng, X. C., Yang, W. D., Liu, J. S., Peng, Z. Y., Lü, S. H., and Ding, W. Z. (2005). Characterization of the
hemolytic properties of an extract from *Phaeocystis globosa* Scherffel. *J. Integr. Plant Biol.* 47, 165–171.
doi:10.1111/j.1744-7909.2005.00039.x.
Peperzak, L., Colijn, F., Vrieling, E. G., Gieskes, W. W. C., and Peeters, J. C. H. (2000). Observations of flagellates
in colonies of *Phaeocystis globosa* (Prymnesiophyceae); a hypothesis for their position in the life cycle. *J.*
Plankton Res. 22, 2181–2203. doi:10.1093/plankt/22.12.2181.
Peperzak, L., and Gabler-Schwarz, S. (2012). Current knowledge of the life cycles of *Phaeocystis globosa* and
Phaeocystis antarctica (prymnesiophyceae). *J. Phycol.* 48, 514–517. doi:10.1111/j.1529-8817.2012.01136.x.
Peperzak, L., and Poelman, M. (2008). Mass mussel mortality in The Netherlands after a bloom of *Phaeocystis*
globosa (prymnesiophyceae). *J. Sea Res.* 60, 220–222. doi:10.1016/j.seares.2008.06.001.
Pesquita, C., Faria, D., Falcão, A. O., Lord, P., and Couto, F. M. (2009). Semantic similarity in biomedical
ontologies. *PLoS Comput. Biol.* 5. doi:10.1371/journal.pcbi.1000443.
Quinlan, A. R., and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics 26, 841–842. Available at: http://dx.doi.org/10.1093/bioinformatics/btq033.
Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2009). edgeR: A Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* 26, 139–140.
doi:10.1093/bioinformatics/btp616.
Rousseau, V., Chrétiennot-Dinet, M.-J., Jacobsen, A., Verity, P., and Whipple, S. (2007). The life cycle of
Phaeocystis: state of knowledge and presumptive role in ecology. *Biogeochemistry* 83, 29–47.
Rousseau, V., Lantoine, F., Rodriguez, F., LeGall, F., Chretienot-Dinet, M. J., and Lancelot, C. (2013).
Characterization of *Phaeocystis globosa* (Prymnesiophyceae), the blooming species in the Southern North
Sea. *J. Sea Res.* 76, 105–113. doi:10.1016/j.seares.2012.07.011.
Schoemann, V., Becquevort, S., Stefels, J., Rousseau, V., and Lancelot, C. (2005). *Phaeocystis* blooms in the global
ocean and their controlling mechanisms: A review. *J. Sea Res.* 53, 43–66. doi:10.1016/j.seares.2004.01.008.
Scott, C. (2018). Dammit: an open and accessible de novo transcriptome annotator. *in prep.* Available at:
www.camillescott.org/dammit.
Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., and Zdobnov, E. M. (2015). BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31, 3210–3212.
doi:10.1093/bioinformatics/btv351.
Solomon, C. M., Lessard, E. J., Keil, R. G., and Foy, M. S. (2003). Characterization of extracellular polymers of
Phaeocystis globosa and *P. antarctica*. *Mar. Ecol. Prog. Ser.* 250, 81–89. doi:10.3354/meps250081.
Spilmont, N., Denis, L., Artigas, L. F., Caloin, F., Courcot, L., Créach, A., et al. (2009). Impact of the *Phaeocystis*
globosa spring bloom on the intertidal benthic compartment in the eastern English Channel: A synthesis. *Mar.*
Pollut. Bull. 58, 55–63. doi:10.1016/j.marpolbul.2008.09.007.
Supek, F., Bošnjak, M., Škunca, N., and Šmuc, T. (2011). Revigo summarizes and visualizes long lists of gene
ontology terms. *PLoS One* 6. doi:10.1371/journal.pone.0021800.
Taj, G., Agarwal, P., Grant, M., and Kumar, A. (2010). MAPK machinery in plants. *Plant Signal. Behav.* 5, 1370–
1378. doi:10.4161/psb.5.11.13020.
Tang, K. W. (2003). Grazing and colony size development in *Phaeocysis globosa* (Prymnesiophyceae): the role of
chemical signal. *J. Plankton Res.* 35, 831–842.
Tang, K. W., Jakobsen, H. H., and Visser, A. W. (2001). *Phaeocystis globosa* (Prymnesiophyceae) and the
planktonic food web: feeding, growth, and trophic interactions among grazers. *Limnol. Oceanogr.* 46, 1860–
1870. doi:10.4319/lo.2001.46.8.1860.
Taylor, A. R., Russell, M. A., Harper, G. M., Collins, T. F. T., and Brownlee, C. (2007). Dynamics of formation and
secretion of heterococcoliths by *Coccolithus pelagicus* ssp. braarudii. *Eur. J. Phycol.* 42, 125–136.
doi:10.1080/09670260601159346.
Team, R Core. 2013. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for
Statistical Computing. http://www.r-project.org/.

Van Boekel, W. H. M. (1992). *Phaeocystis* colony mucus components and the importance of calcium ions for
colony stability. *Mar. Ecol. Prog. Ser.* 87, 301–305. doi:10.3354/meps087301.
Van Duyl, F. C., Gieskes, W. W. C., Kop, A. J., and Lewis, W. E. (1998). Biological control of short-term variations
in the concentration of DMSP and DMS during a *Phaeocystis* spring bloom. *J. Sea Res.* 40, 221–231.
doi:10.1016/S1385-1101(98)00024-0.
Veldhuis, M. J. W., Brussaard, C. P. D., and Noordeloos, A. A. M. (2005). Living in a *Phaeocystis* colony: a way to
be a successful algal species. *Harmful Algae* 4, 841–858. doi:10.1016/j.hal.2004.12.013.
Verity, P. G., and Medlin, L. K. (2003). Observations on colony formation by the cosmopolitan phytoplankton
genus *Phaeocystis*. *J. Mar. Syst.* 43, 153–164. doi:10.1016/j.jmarsys.2003.09.001.
Wang, X., Wang, Y., Ou, L., He, X., and Chen, D. (2015). Allocation costs associated with induced defense in
Phaeocystis globosa (Prymnesiophyceae): the effects of nutrient availability. *Sci. Rep.* 5, 10850.
doi:10.1038/srep10850.
Wang, X., Wang, Y., and Smith, W. O. (2011). The role of nitrogen on the growth and colony development of
Phaeocystis globosa (Prymnesiophyceae). *Eur. J. Phycol.* 46, 305–314. doi:10.1080/09670262.2011.602430.
Wolfe, G. V, and Steinke, M. (1996). Grazing-activated production of dimethyl sulfide (DMS) by two clones of
Emiliania huxleyi. *Limnol. Oceanogr.* 41, 1151–1160.
Wozniak, M. A., Modzelewska, K., Kwong, L., and Keely, P. J. (2004). Focal adhesion regulation of cell behavior.
Biochim. Biophys. Acta - Mol. Cell Res. 1692, 103–119. doi:10.1016/j.bbamcr.2004.04.007
Woznica, A., Cantley, A. M., Beemelmanns, C., Freinkman, E., Clardy, J., and King, N. (2016). Bacterial lipids
activate, synergize, and inhibit a developmental switch in choanoflagellates. *Proc. Natl. Acad. Sci.* 113, 7894–
7899. doi:10.1073/pnas.1605015113.
Woznica, A., Gerdt, J. P., Hulett, R. E., Clardy, J., and King, N. (2017). Mating in the closest living relatives of
animals is induced by a bacterial chondroitinase. *Cell* 170, 1175–1183.e11. doi:10.1016/j.cell.2017.08.005.
Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for comparing biological themes
among gene clusters. *Omi. a J. Integr. Biol.* 16, 284–287.
826