Modeling the relative role of human mobility, land-use and climate factors on dengue outbreak

emergence in Sri Lanka

Ying Zhang[1,2], Jefferson Riera[3], Kayla Ostrow[1], Sauleh Siddiqui[1,4], Harendra de Silva[5], Sahotra Sarkar[6],

Lakkumar Fernando[7], Lauren Gardner[1,8*]

[1]Department of Civil Engineering, Johns Hopkins University, Baltimore, MD 21218, USA

[2]Department of Earth and Planetary Sciences, Johns Hopkins University, Baltimore, MD 21218, USA

[3]Department of Environmental Health and Engineering, Johns Hopkins University, Baltimore, MD

21205, USA

[4]Department of Applied Mathematics and Statistics, Johns Hopkins University, Baltimore, MD 21218,

USA

[5]Department of Pediatrics, University of Colombo, Colombo, 00900, Sri Lanka

[6]Department of Philosophy, Department of Integrative Biology, University of Texas at Austin

Austin, TX, 78712

[7]Centre for Clinical Management of Dengue and Dengue Haemorrhagic Fever, Negombo, 11500, Sri

Lanka

[8]School of Civil and Environmental Engineering, University of New South Wales (UNSW) Sydney,

Sydney, NSW, 2052 Australia

*Corresponding Author: l.gardner@jhu.edu

Email of other authors:

Ying Zhang: ying.zhang@jhu.edu

Jefferson Riera: jriera1@jhu.edu

Kayla Ostrow: kostrow5@jhu.edu

Sauleh Siddiqui: siddiqui@jhu.edu

Harendra de Silva: harendra51@gmail.com

Sahotra Sarkar: sarkar@austin.utexas.edu

Lakkumar Fernando: lakkumar@gmail.com

Abstract

Background: More than 80,000 dengue cases including 215 deaths were reported nationally in less than

seven months between 2016-2017, a fourfold increase in the number of reported cases compared to the

average number over 2010-2016. The region of Negombo, located in the Western province, experienced

the greatest number of dengue cases in the country and is the focus area of our study, where we aim to

capture the spatial-temporal dynamics of dengue transmission.

Methods: We present a statistical modeling framework to evaluate the spatial-temporal dynamics of the

2016-2017 dengue outbreak in the Negombo region of Sri Lanka as a function of human mobility, land-

use, and climate patterns. The analysis was conducted at a $1 \text{ km} \times 1 \text{ km}$ spatial resolution and a weekly

temporal resolution.

Results: Our results indicate human mobility to be a stronger indicator for local outbreak clusters than

land-use or climate variables. The minimum daily temperature was identified as the most influential

climate variable on dengue cases in the region; while among the set of land-use patterns considered, urban

areas were found to be most prone to dengue outbreak, followed by areas with stagnant water and then

coastal areas. The results are shown to be robust across spatial resolutions.

Conclusions: Our study highlights the potential value of using travel data to target vector control within a

region. In addition to illustrating the relative relationship between various potential risk factors for dengue

outbreaks, the results of our study can be used to inform where and when new cases of dengue are likely

to occur within a region, and thus help more effectively and innovatively, plan for disease surveillance

and vector control.

Keywords: dengue; outbreaks; risk factors; human mobility; climate; land-use; spatial-temporal

dynamics; statistical modeling, Sri Lanka

1. Background

Dengue is a mosquito-borne viral disease that infects approximately 390 million people globally every year, particularly in tropical and subtropical countries (1, 2). The high number of infections combined with the lack, as yet, of a routinely used effective vaccine has made dengue a notorious public health problem (2, 3).

Dengue spreads through the bite of infected *Ades* mosquitoes, especially *Aedes aegypti*– the primary vector, with an estimated 15 to 17-day delay between the primary and secondary human infections (4). Dengue outbreak control is a challenge for policy makers because *Aedes aegypti* mosquitoes are well adapted to high density urban environments and actively feed during the day (5-7), thus presenting an elevated risk to humans. Urban settings provide an ideal habitat for *Aedes aegypti* breeding due to an abundance of discarded trash bags, plastic bottles, tires, and other containers that enable the formation of stagnant shallow water surfaces after precipitation (8). Urban regions in developing countries are particularly vulnerable due to a lack of indoor plumbing infrastructure that, in conjunction with a lack of air-conditioning, results in higher human-mosquito exposure rates during the day. Additionally, because of the daytime feeding behaviors of *Aedes aegypti*, common vector control measures that work for night-biting mosquitoes, such as bed nets, fail to effectively control dengue transmission. Given these challenges, there is a need to better understand and predict dengue outbreaks and transmission risk within urban regions in developing countries so that vector control and surveillance resources can be optimally allocated.

Previous studies highlighted human mobility as a critical factor for dengue transmission (9-15), which contrasts the more minor role travel plays in the spread of vector-borne diseases transmitted by night-biting mosquitoes (15). While *Aedes aegypti* mosquitoes have a hard time dispersing geographically

across large areas because they rarely travel more than 400m from where they emerge as adults (16-19),

humans regularly travel much longer distances on a daily basis. As new dengue cases and clusters are

regularly reported kilometers apart, it is likely that human mobility play a critical role in the spread of

dengue outbreaks, *i.e.,* infected humans introduce dengue into new mosquito populations at their trip

ends. As an example, Vazquez-Prokopec, Montgomery (12) studied the pattern of dengue transmission

using location-based contact tracing on infected dengue patients during a dengue outbreak centered at

Cairns, Australia. They collected locations that the patients frequently traveled to during the daytime and

2-4 weeks prior to the onset of symptoms through phone interviews. The contact locations with a

proximity of 100 meters and a separation of 20 days were spatial-temporally linked into pairs and then

chains to identify the plausible sites of dengue virus transmission. They showed that the complex pattern

of dengue transmission was primarily driven by human mobility, and that targeted residual spaying could

potentially reduce the probability of dengue transmission up to 96%. Their study highlights the

importance of understanding dengue transmission patterns to optimize the allocation of dengue prevention

and vector-control measures.

In addition to human mobility, recent studies have pointed to a strong association between climate

conditions and dengue outbreaks at various locations and across different temporal resolutions (8, 20-24).

Precipitation, mean temperature and temperature fluctuation were revealed to affect the population

dynamics of *Aedes aegypti* mosquitoes and the dengue virus extrinsic incubation period (25-29).

Specifically, a suitable average temperature and moderate temperature fluctuations are often favorable for

dengue transmission (25), while an increase in precipitation is strongly associated with the onset of a

dengue outbreak (22). Humidity, a combined effect of precipitation and temperature, is also a common

climate index to evaluate the environmental capacity for dengue emergence (20, 21, 30, 31). Wesolowski,

Qureshi (13) accounted for both climate and mobility in a study of dengue virus transmission over a large

dengue outbreak period in Pakistan. They developed an epidemiological model that included temperature

and relative humidity as input parameters for mosquito dynamics, as well as biting rate to capture the

interactions between human and mosquito hosts. Human mobility was captured using mobile phone data

of ~40 million subscribers to estimate the spatially explicit travel volume, albeit not differentiating

infected and non-infected people. They showed that the emergence of dengue epidemics in a new region

could be predicted using aggregated travel patterns from endemic areas in combination with the

developed epidemiological model. While climatic factors were found to be significant for prediction, this

was in part due to the large study region, *i.e.*, country level, which has variable climatic suitability for the

mosquito vector. The study region considered in our work is much smaller and has minimal climactic

variability, thus alternative methods are required to distinguish site-specific risk.

Land-use patterns — indicators of human activities and potential breeding habitats — have also been

linked to dengue outbreaks (32-37). Previous studies investigated the effect of land-use patterns on the

spread of dengue and found that human settlements, water bodies, and mixed horticulture are the top three

associated land-use patterns for dengue emergence in Malaysia (35). In another study (36), areas

surrounded by rice paddies and marshes/swamps were associated with a significantly higher population of

dengue vectors during the rainy season in Thailand. Orchards (which often contain artificial water

containers) and irrigation fields have also been shown to play an important role in dengue infections;

however, their role varies given different local conditions. Sometimes, land-use type can be a proxy for

other features, such as socio-economic factors, which may have a contradictory effect on dengue

infections (37).

In this study, we present a statistical modeling framework to evaluate the relative role of human travel

patterns, climate conditions, and land-use patterns on dengue outbreak dynamics in Negombo, Sri Lanka

(Figure 1), where dengue outbreaks are increasing in scale and prevalence. Dengue has been endemic in

Sri Lanka since 1960s, and all four serotypes circulate in the region (38). Dengue hemorrhagic fever

(DHF) cases were rare until 1989, when the first major dengue outbreak occurred with approximately 200

cases and 20 deaths; DHF has since become endemic in the country (38-40). The 2017 outbreak involved

many severe DHF cases, with evidence showing a dengue virus type 2 (DENV-2) being the causative

serotype (41, 42). DENV-2 had only be detected infrequently over the recent decades of dengue

epidemics and is therefore associated with low immunity in the region (38). As a result, more than 80,000

dengue cases including 215 deaths were reported nationally in less than seven months between 2016-

2017, a fourfold increase in the number of reported cases compared to the average number over 2010-

2016 (43). The region of Negombo, located in the Western province, experienced the greatest number of

dengue cases in the country; approximately 45% of the cases nationwide by July 2017 (Figure 1), and is

the focus area of our study.

We applied a mixed-effects model, where the mobility data bridges the time-varying, spatially-invariant

climate variables and the rasterized spatially explicit, time-invariant population and land-use variables, to

capture the spatial-temporal dynamics of dengue transmission. Our model framework differs from

previous studies that simulated the transmission process (12, 13, 44), and instead focuses on estimating

the timing and location of new case introductions though a non-process based statistical model.

Specifically, we focus on modeling the home locations of (newly infected) dengue patients, and assume

dengue is introduced in new areas by infected individuals who travel to the area. This assumption is

consistent with previous studies that have shown visits to a household by infected people determines the

infection risk in that household (11). In addition, the study was conducted at a fine-grained spatial and

temporal resolution — 1 km × 1 km spatially and one week temporally — providing an improved

understanding of the role of mobility in the spread of dengue. While previous work studied the impact of

mobility (12, 13), climate (8, 20, 22, 25), and land-use (35-37) separately on dengue, the authors are

unaware of any existing study that considers these factors within a single integrated framework. Thus,

previous studies have been unable to quantify the relative contribution of each factor on the spatial-

temporal patterns of dengue transmission as we do. The results from our study indicate that mobility is a

more significant indicator of new dengue case clusters compared with land-use and climate factors.

Furthermore, the case study in Sri Lanka provides critical insights into effective application of dengue

prevention and vector control measures in developing regions.

Fig. 1. Land-use groups and the climate station in the study region of Negombo, Sri Lanka (left), and the
total number of dengue cases at the district level in Sri Lanka (45) during the months from October 2016
to June 2017 that cover our study period (right).

2. Methods

A statistical model is applied to investigate the spatial-temporal dynamics of dengue outbreak with respect to a range of potential explanatory variables. The complete set of potential explanatory variables is listed in Table 1. Detailed descriptions of the data followed by a description of the methodology are provided below.

(a) Case and Mobility Data

A patient travel survey was conducted among dengue patients in the Negombo region of Sri Lanka over an approximately 8-month period during a major outbreak spanning from end of October 2016 to early July 2017. Geolocation data were collected from all patients admitted to the special High Dependency Unit (HDU) for critically ill dengue patients within the Clinical Centre for Managing Dengue and Dengue Haemorrhagic Fever (CCMDDHF) at the Negombo Hospital in Negombo, Sri Lanka. Specifically, the date of admission, home address, the complete set of locations visited, and corresponding trips made between all locations during the 10-days prior to hospital admission were collected from all HDU CCMDDHF admitted patients for the entire study period. The case data provide spatial-temporal information on the outbreak patterns, while the mobility data collected captures daily travel activity of the admitted dengue patients. The data were collected by trained students and supervised by a Senior House Officer on site. For weekday admissions the patients were surveyed upon admittance, for night admissions data were collected the following day, and for weekend admissions on the following Monday. The majority of admissions were 48 to 72 hours following the onset of fever. Dengue infection was confirmed for the patient set using either NS1 antigen or IgM antibody diagnostic test.

(b) Climate Data

We used the Global Surface Summary of the Day (GSOD) daily weather data (46) from a station in

Negombo (Figure 1) to explore the impact of climate factors on the dengue outbreak. The location of the

weather station (7.18ºN, 79.87ºE) is approximately in the center of the study region and it is the only

station that falls into our study region with a comprehensive set of climate data available during the study

period. There are several global reanalysis products that provide spatial-explicit climate data during the

study period; however, upon evaluation against the station observations, these globally gridded data sets

did not provide accurate representations of the local climate variables, particularly at a daily time-step

(Figure S1). Hence, the weather data are assumed to be representative for the region which has relatively

homogeneous weather patterns (47). We selected a range of potential climate variables based on previous

studies (8, 20-22, 25-31), including daily mean temperature (*T*avg), daily maximum temperature (*T*max),

daily minimum temperature (*T*min), diurnal temperature range (*DTR*), precipitation (*Pre*), the number of

raining days (*RD*), and relative humidity (*RH*) to analyze climatic influence for the weeks before and

during the same period of analysis that the mobility data was collected.

(c) Population and Land-use Data

We used a global population data layer based on Landscan 2016 (48), that is available at an

approximately 1 km × 1 km resolution to represent the population distribution spatially. We aggregated

the data to 5 km × 5 km grid for additional analysis with a coarser spatial resolution. Land-use data (49)

were obtained from the Sri Lanka Survey Department which performed an initial survey in 2000 and has

since continuously updated the maps. The map was extracted for our region of interest and reclassified

into several groups (Figure 1): *Sea, Standing Water (StWtr), Flowing Water (FlwWtr), Coconut, Marsh,*

Paddy, Built-up (BuiltUp), Scrubland, Homesteads, Forest, Rubber, Rock/Sand (RockS), Other

Agriculture (OthAg), and Other. Water bodies were categorized depending on the potential effect on

dengue transmission dynamics. Additional details on land-use classification groupings and processing is

available in the supplementary material.

(d) Data Processing and Statistical Model

We divided the study region (Figure 1) into a grid at a 1 km × 1 km resolution and aggregated daily data

into a 1-weekly resolution. The number of patients who were admitted to the hospital during each week of

the recorded time period was used to generate the weekly number of newly admitted dengue patients in

each cell based on their home locations. This becomes our 'case' variable.

To incorporate the role of mobility into the model we used the travel itineraries provided by the patients

to generate a time-dependent connectivity matrix, which represented the total number of trips made by

dengue infected patients between each pair of cells for each week of the study period. The travel data

included all destinations visited each day during the 10 days preceding hospital admittance (the time

interval that the patient is assumed to be able to spread the disease) for each patient. The number of daily

trips between each pair of cells was summed over all patients, to provide daily trip volumes between cells,

and then aggregated to the weekly level. For each cell the total incoming weekly trips was summed to

define our 'trip' variable. Critically, we exclude all trips with a destination of 'home' when computing our

trip variable, in order to remove the inherent dependence between the 'case' variable, *i.e.,* the home

location of infected individuals, and the 'trip' variable (explanatory variable). Thus, the total number of

trips (excluding trips home) made by infected dengue patients entering a given cell *i* in a given week *t*,

V_t^i, was used as a spatial-temporal explanatory variable in the model. The same method was used for the 5

km × 5 km analysis.

Climate variables were averaged or aggregated temporally to a weekly resolution, including weekly

average Tavg, Tmin, Tmax, DTR, RH, weekly total Pre, and RD. Land-use data were aggregated spatially

to match the targeted spatial grid resolution. The population data were in an original resolution that

matched the 1 km × 1 km grid. For land-use, the percentage of occupied land of each type was determined

for each 1 km × 1km grid cell. Both were subsequently aggregated to a 5 km × 5 km grid.

A linear mixed-effects model combined with backward elimination of insignificant fixed effects (p-value

> 0.05, two-tail test) was applied to investigate the spatial-temporal dynamics of dengue outbreak with the

potential explanatory variables at a weekly time step and 1 km × 1 km spatial resolution. In building the

model we first conducted sensitivity analysis to identify the optimal set of climatic variables to include in

the model, and corresponding time lag for each of them.

Along with the chosen climate variable, the remaining set of potential explanatory variables (Table 1) was

normalized and then taken into the mixed-effects model initially, with population included in the spatial

random effects. Population density was incorporated using random effects in the model because

population is likely to have spatially heterogeneous effects on dengue outbreaks (44, 50). For example,

high population areas may imply access to tap water and better living conditions which could restrict

dengue transmission (51), while the higher density of population facilitates disease spread. Furthermore,

there could be spatial variance in the distribution of people living in a particular area. In addition to

mobility, climate, and land-use variables; the number of new cases in a given cell in the weeks prior were

added as explanatory variables to account for autocorrelations in the case data. Subsequently, the variable

with the most insignificant fixed-effects coefficient was eliminated each iteration, until only variables

with significant coefficients (at 95% significance level) remained in the model. A range of lead time for

V_t^i prior to the admitted week was also tested. A separate analogous process was conducted using a 5 km

× 5 km resolution, to test the sensitivity of model results across spatial resolutions, and the robustness of

the modeling framework and findings.

Thus, the mathematical representation of the model is given by:

5
$$N_t^i = \sum_{l \in \bar{L}} \alpha_l f_l^i + \sum_{c \in \bar{C}} \beta_c c_{t,d_c} + \sum_u \gamma_u V_{t-u}^i + \sum_w \delta_w N_{t-w}^i + a^i + b^i P^i + \varepsilon_t^i$$

Where

i is the cell index; $i = 1, 2, \dots.$

l is the land-use variable, which belongs to the land-use group set $\bar{L}$, where $\bar{L}$ includes *Sea, StWtr, FlwWtr,*

Coconut, Marsh, Paddy, BuiltUp, Scrubland, Homesteads, Forest, Rubber, RockS, OthAg, and Other.

f_l^i is the occupation fraction of land-use group l in cell i, time-invariant.

P^i is the population in cell i, time-invariant.

t is the time index at weekly resolution; $t = 1, 2, \dots.$

N_t^i is the number of patients who are admitted to the hospital during week t, whose home locations are in

cell i.

N_{t-w}^i is the number of patients who are admitted to the hospital during week t-w, whose home locations

are in cell i, where w is measured in weeks; $w = 1, 2, \dots$

V_{t-u}^i is the number of total number of trips made into cell i during the week t-u, where u is measured in

weeks; $u = 1, 2, \dots$

c is the climate variable which belongs to the climate variable set $\bar{C}$. $\bar{C}$ includes *T*avg, *T*max, *T*min, *DTR*,

Pre, *RD*, and *RH*.

c_{t,d_c} is the climate variable during the week that begins d_c days prior to the start of week t. d_c ranges

from 7 to 17 days and can be different for different climate variables. Multiple climate variables can be

included in the model.

ε_t^i is the model residual associated with cell i and week t.

α_l is the estimated fixed-effects coefficient for l.

β_c is the estimated fixed-effects coefficient for c.

γ_u is the estimated fixed-effects coefficient for V_{t-u}^i.

δ_w is the estimated fixed-effects coefficient for N_{t-w}^i.

a^i is the intercept associated with cell i.

b^i is the estimated spatial random-effects coefficient for P^i.

The data processing and modeling were performed using MATLAB R2017a.

(e) Sensitivity Analysis

We performed sensitivity analysis to evaluate the robustness of our model results. Firstly, a Jackknife

analysis was conducted, specifically the statistical model was fit to all but one week of data, iteratively

excluding one week at a time, over the entire time period modeled. The variability in estimated

parameters and their corresponding significance are presented in Supplementary Table S3. Second, we

evaluated the sensitivity of the model to fluctuations of the 'trip' variable. The motivation behind this

sensitivity analysis was the uncertainty resulting from potential human error in the 'trip' variable, which

is based on the patients' recollection of their travel in the 10-day prior to hospital admittance. To assess

the robustness of the model to error in the trip variable, we implemented Monte Carlo sampling, assuming

an error of 10% uniformly distributed around the original observations V_t^i. We performed 1000 random

simulations, and the results are presented in Supplementary Table S4.

5   3. Results

6   (a) Data Analysis

The number of admitted dengue patients aggregated over the study region peaks during December and

June (Figure 2), aligned with the monsoon months (38). Figure 2a illustrates the relationship between the

total number of dengue patients, N_t, admitted during each week t and the total number of recorded patient

trips (excluding the trips to home) during the same week (V_t). Figure 2b illustrates N_t and the weekly

averaged minimum daily temperature in week t ($T\min_t$). It shows a lagged relationship of N_t with $T\min_t$,

mostly in the same direction. For the purposes of these graphics, the variables are aggregated over the

entire study region.

(a)   (b)

Fig. 2. The number of admitted dengue patients in week t (N_t) and (a) the number of recorded trips in
week t (V_t) summed over the entire study region, and (b) the weekly averaged minimum daily temperature
($T\min_t$).
19

The travel destinations recorded in our study include medical facilities, homes, workplaces, schools, and

others (Figure S3). Additional analysis performed reveals that a vast majority of trips were longer than the

distance a mosquito can travel. Specifically, 96.6% of the trips were longer than 0.4 km (Table S1; Figure

S4), outside the range of a mosquito's maximum travel distance (16-19), further supporting the role

human mobility is likely to play in the outbreak.

Figure 3 illustrates both the spatial-temporal distribution of dengue patients' home locations over the

course of the outbreak, and the corresponding travel patterns of the patients during 5-week periods,

excluding the trips with a destination of 'home'. The patient home locations were well distributed over

the area of the study region for the first few months of the outbreak, with correspondingly scattered travel

patterns. However, as the outbreak progressed, the recorded case locations and the trip ends of newly

infected dengue patients became more concentrated near the town center and just above the lake. There

were also a large number of trips (>50) within the cell near the town center.

1 (a) Weeks 2-6   (b) Weeks 7-11   (c) Weeks 12-16

3 (d) Weeks 17-21   (e) Weeks 22-26   (f) Weeks 27-31

Fig. 3. Weekly number of patients and the number of trips summed over 5-week intervals for a 5 km × 5
km resolution (excluding trips with a destination of 'home'). Patient home locations are plotted as the
case location. The size of the circle indicates the number of patients admitted during the time period. The
color of the circle indicates the number of trips that end in the grid cell during the time period. The
thickness of the line is proportional to the number of trips made between two locations. Week 2 begins on
October 27, 2016 and week 31 begins on May 18, 2017. For visual clarity, the 5 km × 5 km resolution
was used for the figure, instead of 1 km × 1 km.

(b) Selection of Climate Variables

Among the climate variables, significant correlations were observed for weekly averaged Tavg, Tmin, and

DTR with a lead time ranging from 7 days to 17 days prior to the weekly admitted number of patients

(N_t), where the lead time (d_c) is the lag in days between the climate variable and N_t (Figure S2).

Regression models based on different combinations of the climate variables and lead time were developed

and compared; the best performance model was select based on F-test and adjusted-R^2. As a result, Tmin

with an optimal lead time of 10 days was included in the final set of mixed-effects models to account for

the partial influence of climate on the dengue outbreak ($R^2 = 0.248$; *adj.* $R^2 = 0.226$). This is consistent

with previous findings (29) that daily minimum temperature were associated with increase in the larval

abundance. We assumed a relatively homogenous climate over the study region, thus *Tmin* does not vary

spatially over the study region.

(c) Model Results

A mixed-effects model was developed to estimate the number of new dengue cases in a given cell in a

given week as a function of the mobility patterns of individuals infected with dengue in the preceding

week(s), as well as land-use and climate data from days prior.

Table 1: Summary of potential explanatory variables

Variables	Description	Properties
f_l^i	occupation fraction of land-use group l in cell i (%)	
BuiltUp	urban area	
Coconut	coconut cultivation land	
Homesteads	homesteads and garden	
Paddy	rice cultivation land	spatially-explicit, time-invariant
Sea	ocean	
StWtr	standing water	
FlwWtr	flowing water	
c_{t,d_c}	weekly value of climate variable in week t with a lag of d_c	
*T*avg	weekly averaged daily mean temperature	
*T*max	weekly averaged maximum daily temperature	
*T*min	weekly averaged minimum daily temperature	
DTR	weekly averaged diurnal temperature range	time-varying, spatially-invariant
Pre	weekly total precipitation	
RD	weekly number of raining days	
RH	weekly averaged daily relative humidity	
P^i	population in cell i	spatially -explicit, time-invariant
V_{t-u}^i	number of trips made to cell i in week t-u	spatially -explicit, time-varying
N_{t-w}^i	number of patients admitted to hospital who live in cell i in week t-w	spatially -explicit, time-varying

Note: t, u, and *w* in weeks, d_c in days. For notation, variable superscripts in Table 1 denote spatial indices and subscripts denote time indices.

Multiple models with explanatory variables representing land-use, climate, and mobility were created,

and the three representative models are presented here. The three models vary based on the type of

mobility variable included, specifically how far back in time travel is accounted for. The first model

includes the mobility patterns one-week prior ($u = 1$), the second model includes the mobility patterns

two-weeks prior ($u = 2$), and the third model excludes mobility altogether ("Exclude V"). The final set of

climate and land-use variables found to be significant varies between models. All explanatory variables

were normalized to a mean of zero and a standard deviation of one in the mixed-effects model. The fixed-

effects coefficients (Table 2) therefore reflect the relative influence of each explanatory variable on the

dengue outbreak dynamics.

The results (Table 2) for each of the three models are presented for both a 1 km × 1 km and 5 km × 5 km

resolution, and reveal that the mobility patterns of dengue patients, specifically the number of trips made

into a cell in a given week, to be the most dominant explanatory factor of new dengue cases in that cell

the following week. Under the spatial resolution of 1 km × 1 km, the fixed-effects coefficient for the trips

one week prior ($u = 1$) is 0.070, which is greater than the fixed-effects coefficients for other explanatory

variables, suggesting human mobility plays a critical role in dengue outbreak dynamics. Results also

illustrate a decrease in explanatory power of mobility patterns further than a week in advance, and are

insignificant for the trips variable with a lead time of two weeks ($u = 2$). When mobility data is excluded

from the model altogether, the adjusted R^2 decreases slightly. In general, the power of the number of trips

in predicting dengue cases deteriorates with longer lead time, with the number of trips two-weeks prior

showing little or no advantage over other explanatory variables. The same conclusion is applicable for the

results under the 5 km × 5 km resolution, as shown in Table 2. These results highlight the importance of

collecting and utilizing mobility data within an appropriate timeframe for the purposes of modeling

dengue outbreak risk. These conclusions are further supported by the sensitivity analysis, which show the

rank and magnitude of the estimated model parameters (and significance) are robust to changes in both

the outbreak period and also variability in the trips variable (Figure S6).

Among the seven land-use groups (see variable descriptions in Table 1) under the 1 km × 1 km resolution,

only *BuiltUp* shows significant positive fixed effects on dengue cases (Table 2). Under the coarser spatial

rasterization of 5 km × 5 km, *StWtr* and *Sea* also show significant positive fixed effects, in addition to

BuiltUp. Whereas *BuiltUp* and *StWtr* show significant fixed effects in all three models with the

coefficients ranging from 0.034 to 0.050 for *BuiltUp* and 0.026 to 0.032 for *StWtr*, , *Sea* shows the

significant coefficient of 0.029 only in the model with the trip variable excluded. It indicates that urban

areas, areas with standing water, and areas near the coastline are associated with a higher risk of dengue

infections; the effect of *StWtr* and *Sea* is stronger under the 5 km × 5 km spatial resolution. In contrast,

human mobility and *BuiltUp* are shown to be significant and robust indicators of dengue dynamics for

both spatial resolutions. The same conclusions hold under the sensitivity analysis performed (Table S3

and S4).

Table 2: Fixed-effects coefficients and standard error of the mixed-effects model outputs based on the 1 km × 1 km and 5 km × 5 km resolution, respectively. The presented results are post-completion of the backward elimination of nonsignificant fixed effects. Variables without coefficients listed in the table were eliminated during the backwards elimination procedure for each model (each column). Variable descriptions are listed in Table 1.

	1 km × 1 km			5 km × 5 km		
	$u = 1$	$u = 2$	Exclude V	$u = 1$	$u = 2$	Exclude V
BuiltUp	0.047***	0.055***	0.053***	0.037**	0.034*	0.050**
	(0.014)	(0.014)	(0.014)	(0.014)	(0.014)	(0.016)
Sea					0.029*	
					(0.013)	
StWtr				0.026*	0.032**	0.032**
				(0.011)	(0.011)	(0.011)
$Tmin_{t, d_{Tmin}}$	0.027***	0.026***	0.026***	0.024*	0.023*	0.021*
	(0.0073)	(0.0075)	(0.0074)	(0.0097)	(0.010)	(0.0098)
V_{t-1}^i	0.070***			0.113***		
	(0.0086)			(0.013)		
V_{t-2}^i					0.040**	
					(0.013)	
N_{t-1}^i	0.191***	0.196***	0.195***	0.331***	0.350***	0.0359***
	(0.0083)	(0.0084)	(0.0083)	(0.017)	(0.017)	(0.016)
N_{t-2}^i	0.214***	0.217***	0.217***	0.368***	0.398***	0.403***
	(0.0082)	(0.0084)	(0.0082)	(0.017)	(0.017)	(0.016)
R^2	0.265	0.265	0.262	0.733	0.728	0.727
Adj. R^2	0.265	0.264	0.262	0.733	0.728	0.727
No. obs	13532	13035	13532	2856	2772	2856

Standard errors are reported in parentheses.

t is in weeks, $d_{Tmin} = 10$ days, and all variables are normalized.

*, **, *** indicates significance at the 95%, 99%, and 99.9% level, respectively.

Results from the sensitivity analysis are presented in Supplementary Table S3 for jackknife, and Table S4

and Figure S6 for trip variable error. Both sets of analysis reveal minimal fluctuation in the estimated

model coefficients and set of significant variables, which illustrates the robustness of the model results.

Specifically, for both spatial resolutions, travel volumes into a given cell one week prior is always a

significant indicator of newly reported cases in the cell, while travel two-weeks prior is a less reliable

factor, and negligent at smaller spatial scales, *i.e.,* 1 km × 1 km. Further, travel patterns one-week prior is

consistently identified as a more significant indicator of new dengue cases than climate and land use.

4. Discussion

The results from this study illustrate the critical contribution of human mobility on the location and

timing of new dengue cases, relative to land-use and climate variables. The results are sensitive to the

temporal patterns of travel during the week immediately preceding the appearance of new case reports.

This was the variable with the greatest predictive power. Although, travel patterns two weeks prior were

found to have an insignificant effect on dengue outbreak dynamics. Our results are consistent with

Stoddard, Forshey (11), who concluded that visits to households by dengue infected individuals

determines the infection risk, further validating our use of patient home locations in the model.

Furthermore, the significance of mobility in outbreak prediction was found to be robust under both spatial

resolutions of 1 km × 1 km and 5 km × 5 km.

In contrast to the role of mobility, which we found to be a consistently significant indicator of new

dengue cases, the effect of land-use patterns on the number of new cases is sensitive to the spatial

resolution of the models. Land-use variables played a larger explanatory role at the coarser spatial

resolution of 5 km × 5 km (compared to the finer 1 km × 1 km resolution), particularly for smaller

spatially-dominant land-use patterns such as *Sea* and *StWtr*. *BuiltUp* showed the strongest positive effect

overall, indicating urbanization is associated with an increased risk of dengue outbreak (which is

consistent with multiple previous findings (52, 53)). *StWtr* also showed significant positive effect, which

is to be expected because standing water provides suitable mosquito breeding habitat (19). The positive

effect of *Sea* only appeared significant when human mobility was excluded from the model. Given the

significant positive correlation between *Sea* and the number of trips (Table S2), it is likely that the large

travel volume towards the area near the coastline makes the study region prone to dengue outbreaks.

Previous studies have also found evidence that dengue mosquitoes can breed in brackish water (54),

further supporting our finding of the distance to the sea as significant in some of the models. If this

pattern holds in other regions, as seems likely, that fact can be used for the spatial prioritization of

resource allocation for disease case and vector surveillance and control.

Among the climate factors, temperature-related variables including *T*avg, *T*min, and *DTR*, were more

strongly associated with the outbreak emergence than precipitation-related variables including *Pre* and

RD, or *RH*, which is related to both. This finding is in accordance with (22), which concluded that

"rainfall strongly modulates the timing of dengue (*e.g.*, epidemics occurred earlier during rainy years)

while temperature modulates the annual number of dengue fever cases." Based on regression analysis, we

found *T*min with a 10-day lead time to be the best climate-based predictor of new weekly dengue cases.

Given the likely robustness of this result in other regions, this fact can be used for the temporal

prioritization of resources.

In addition to human mobility, climate, and land-use variables, which were included as fixed effects,

population density was incorporated using random effects in the model because population is likely to

have spatially heterogeneous effects on dengue outbreaks, as noted in the *Methods*. Based on the model

results, the random-effects coefficients for population are mostly positive, as expected, indicating that higher population density is associated with a higher number of dengue cases (Figure S5). The most significant positive effect is seen north of the lagoon along the coastline, highlighting potentially high-risk areas, where higher populations are likely to facilitate the emergence of dengue outbreaks. A few cells resulted in negative random-effects coefficients, which may be due to confounding interactions between different variables included in the model, or alternative factors not captured in the model; these cells were few and only occurred in the model when the dominant mobility variable was included. It is possible the dominant role of mobility could over compensate for the impact of population, *e.g.,* because people are likely to travel to crowded downtown areas, along the lagoon, or near the ocean where the large number of trips made to those regions could offset the impact of population. That the random-effects coefficients for population density are positive and negative lends support to the modeling decision to treat it as having enough stochasticity to qualify as a random-effects variable.

The results of this analysis have implications that are relevant to the design of measures to control dengue cases, such as allocation of resources for mosquito vector control. Previous global modeling of ecological suitability for dengue vector mosquito species (both *Aedes aegypti* and *Aedes albopictus*) have shown that the entire study area is a prime habitat for these species (55, 56). This conclusion drawn from the global models finds validation in our analysis, which shows that climate and land-use variables are not the most strongly associated with dengue case outbreaks. Consequently, epidemiological risk based on vector ecology may be insufficient for the purposes of optimizing vector control resource allocation, as it is unable to distinguish between potential sites to target within the study area. Because travel into the sites is the most important predictor of new case clusters, it may well be time to optimize vector control resources based on mobility data, with the aim to prevent exposure to the day-biting mosquitoes, *i.e., Aedes aegypti,* at the highest risk locations. To the best of our knowledge, such a design for dengue control measures has not yet been tried in the field.

Finally, various limitations of this study should be noted. First, only dengue patients admitted to the

CCMDDHF at the Negombo Hospital were included and surveyed in this study. Thus mild or

asymptomatic cases, which account for the majority of dengue cases (1), were not accounted for in the

study. Second, some patients infected in the study region may have gone to hospitals in other districts and

would therefore not be included. Third, the mobility data was based on patients' recollections over a 10-

day period prior to hospital admission, and therefore may have inaccuracies due to human error in

recalling the information. However, detailed analysis of the travel data revealed most trips recorded

represent daily commuting routines. Thus, while some trips may be excluded due to human error, we

believe the relative connectivity between cells is accurately captured by the survey responses. In addition,

the sensitivity analysis revealed our results are robust to variation in the time of outbreak and trips

variable. Fourth, the distance traveled and the time spent in a certain location were not considered due to

the unavailability of relevant data; however these factors have been shown to have little influence on

dengue transmission (11), and thus their exclusion does not invalidate the methodology used in this

analysis. Fifth, by utilizing all the mobility data collected, we made an implicit assumption that the

patients were infectious during the entire 10-day period prior to hospital admittance. This period does fall

within the combined intrinsic incubation period (4-10 days) (57) and the early symptomatic period before

admitted to the hospital. A sixth assumption was that the patients were infected at or around their home

locations. This assumption is consistent with a wide variety of previous studies that revealed homes as the

primary point of contact for dengue transmission (11, 58, 59). Vazquez-Prokopec, Montgomery (12) tried

to identify the most plausible transmission locations based on reported contact locations from a dengue

outbreak in Cairns, Australia and found that only 10.2% of the identified transmission sites were at out-

of-home locations, and a notable portion of them were within 1 km of the home locations. Given that our

objective was not to model the transmission chains of dengue as in (12), assuming home locations as the

site of infection provides reasonable support for predicting where infected individuals reside, and

therefore the risk posed around homes of infected individuals. Lastly, the climate data were obtained from a single station, thus a homogenous climatic region was assumed for our study region. Therefore, the role of climate factors on the dengue outbreak may be underestimated.

While the modeling framework used here is readily applicable to other contexts, future work should investigate how widely transferable the model results are. More specifically using general mobility data (tracking movements for all residents); such as using mobile phone data as in (13), or transport planning data, which may be more readily available and cost effective; should be compared to the use of patient mobility surveys as in this study.

5. Conclusions

Our study highlights the potential value of using travel data to target vector control within a region. In addition to illustrating the relative relationship between various potential risk factors for dengue outbreaks, the results of our study can be used to inform where and when new cases of dengue are likely to occur within a region, and thus help more effectively and innovatively, plan for disease surveillance and vector control.

List of abbreviations

DENV-2   dengue virus type 2

HDU   High Dependency Unit

CCMDDHF   Clinical Centre for Managing Dengue and Dengue Haemorrhagic Fever

GSOD   Global Surface Summary of the Day

1	Tavg	daily mean temperature
2	Tmax	daily maximum temperature
3	Tmin	daily minimum temperature
4	DTR	diurnal temperature range
5	Pre	precipitation
6	RD	the number of raining days
7	RH	relative humidity
8	StWtr	Standing Water
9	FlwWtr	Flowing Water
10	BuiltUp	Built-up
11	RockS	Rock/Sand
12	OthAg	Other Agriculture

Declarations

Ethics approval and consent to participate. The protocol was reviewed and approved by the Johns

Hopkins Bloomberg School of Public Health Institutional Review Board (JHSPH IRB), #IRB00009485,

as well as approved by the Negombo Hospital. Patient consent was obtained in writing from all

participants for the purposes of this study.

Concent for publichation. Not applicable

Availability of data and material. Patient case reports and travel diaries can not be shared due to privacy

restrictions. The remaining data used in this study is provided as a supplementary file, along with the code

used to generate the results.

Competing interests. We have no competing interests.

Funding. We receieved no funding for this work**.**

Authors' contributions. L.G. conceived the study. Y.Z., S.Siddique and L.G. designed the experiments.

Y.Z., H.S., L.F. and J.R. collected the data. Y.Z. developed the model and performed the computational

analysis. All authors analyzed the data and model results. Y.Z., S.Siddiqui., S.Sarkar, J.R., K.O., and

L.G., contributed to the writing of the manuscript.

Acknowledgements. We thank Dhammika Silva from the Centre for Clinical Management of Dengue

and Dengue Haemorrhagic Fever who supervised the data collection and Benjamin Zaitchik at Johns

Hopkins University for advice on the climate data extraction.

References

1.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and burden of dengue. Nature. 2013;496:504.

2.	Simmons CP, Farrar JJ, van Vinh Chau N, Wills B. Dengue. New England Journal of Medicine. 2012;366(15):1423-32.

3.	Vannice KS, Roehrig JT, Hombach J. Next generation dengue vaccines: A review of the preclinical development pipeline. Vaccine. 2015;33(50):7091-9.

4.	Aldstadt J, Yoon I-K, Tannitisupawong D, Jarman RG, Thomas SJ, Gibbons RV, et al. Space-time analysis of hospitalised dengue patients in rural Thailand reveals important temporal intervals in the pattern of dengue virus transmission. Tropical Medicine & International Health. 2012;17(9):1076-85.

5.	Chadee DD. Landing periodicity of the mosquito Aedes aegypti in Trinidad in relation to the timing of insecticidal space-spraying. Medical and Veterinary Entomology. 1988;2(2):189-92.

6.	Yasuno M, Tonn RJ. A study of biting habits of Aedes aegypti in Bangkok, Thailand. Bulletin of the World Health Organization. 1970;43(2):319-25.

7.	Wong J, Astete H, Morrison AC, Scott TW. Sampling Considerations for Designing Aedes aegypti (Diptera: Culicidae) Oviposition Studies in Iquitos, Peru: Substrate Preference, Diurnal Periodicity, and Gonotrophic Cycle Length. Journal of Medical Entomology. 2011;48(1):45-52.

8.	Pathirana S, Kawabata M, Goonatilake R. Study of potential risk of dengue disease outbreak in Sri Lanka using GIS and statistical modelling. Journal of Rural and Tropical Public Health. 2009;8:8-17.

9.	Lana RM, Gomes MFdC, Lima TFMd, Honório NA, Codeço CT. The introduction of dengue follows transportation infrastructure changes in the state of Acre, Brazil: A network-based analysis. PLOS Neglected Tropical Diseases. 2017;11(11):e0006070.

10.	Reiner RC, Stoddard ST, Scott TW. Socially structured human movement shapes dengue transmission despite the diffusive effect of mosquito dispersal. Epidemics. 2014;6:30-6.

11.	Stoddard ST, Forshey BM, Morrison AC, Paz-Soldan VA, Vazquez-Prokopec GM, Astete H, et al. House-to-house human movement drives dengue virus transmission. Proceedings of the National Academy of Sciences. 2013;110(3):994-9.

12.	Vazquez-Prokopec GM, Montgomery BL, Horne P, Clennon JA, Ritchie SA. Combining contact tracing with targeted indoor residual spraying significantly reduces dengue transmission. Science Advances. 2017;3(2).

13.	Wesolowski A, Qureshi T, Boni MF, Sundsøy PR, Johansson MA, Rasheed SB, et al. Impact of human mobility on the emergence of dengue epidemics in Pakistan. Proceedings of the National Academy of Sciences. 2015;112(38):11887-92.

14.	Adams B, Kapan DD. Man Bites Mosquito: Understanding the Contribution of Human Movement to Vector-Borne Disease Dynamics. PLOS ONE. 2009;4(8):e6763.

15.	Stoddard ST, Morrison AC, Vazquez-Prokopec GM, Paz Soldan V, Kochel TJ, Kitron U, et al. The Role of Human Movement in the Transmission of Vector-Borne Pathogens. PLOS Neglected Tropical Diseases. 2009;3(7):e481.

16.   Harrington LC, Scott TW, Lerdthusnee K, Coleman RC, Costero A, Clark GG, et al. Dispersal of the
dengue vector Aedes aegypti within and between rural communities. American Journal of Tropical
Medicine and Hygiene. 2005;72(2):209-20.

17.   Hemme RR, Thomas CL, Chadee DD, Severson DW. Influence of Urban Landscapes on Population
Dynamics in a Short-Distance Migrant Mosquito: Evidence for the Dengue Vector Aedes aegypti. PLOS
Neglected Tropical Diseases. 2010;4(3):e634.

18.   Getis A, Morrison AC, Gray K, Scott TW. Characteristics of the spatial pattern of the dengue
vector, Aedes aegypti, in Iquitos, Peru. American Journal of Tropical Medicine and Hygiene.
2003;69(5):494-505.

19.   World Health Organization, Special Programme for Research, Training in Tropical Diseases,
World Health Organization. Department of Control of Neglected Tropical Diseases, World Health
Organization. Epidemic & Pandemic Alert. Dengue: guidelines for diagnosis, treatment, prevention and
control: World Health Organization; 2009.

20.   Xu L, Stige LC, Chan K-S, Zhou J, Yang J, Sang S, et al. Climate variation drives dengue dynamics.
Proceedings of the National Academy of Sciences. 2017;114(1):113-8.

21.   Sirisena P, Noordeen F, Kurukulasuriya H, Romesh TA, Fernando L. Effect of Climatic Factors and
Population Density on the Distribution of Dengue in Sri Lanka: A GIS Based Evaluation for Prediction of
Outbreaks. PLOS ONE. 2017;12(1):e0166806.

22.   Morin CW, Monaghan AJ, Hayden MH, Barrera R, Ernst K. Meteorologically Driven Simulations of
Dengue Epidemics in San Juan, PR. PLOS Neglected Tropical Diseases. 2015;9(8):e0004002.

23.   Metcalf CJE, Walter KS, Wesolowski A, Buckee CO, Shevliakova E, Tatem AJ, et al. Identifying
climate drivers of infectious disease dynamics: recent advances and challenges ahead. Proceedings of
the Royal Society B: Biological Sciences. 2017;284(1860).

24.   Gloria-Soria A, Armstrong PM, Powell JR, Turner PE. Infection rate of *Aedes aegypti*
mosquitoes with dengue virus depends on the interaction between temperature and mosquito
genotype. Proceedings of the Royal Society B: Biological Sciences. 2017;284(1864).

25.   Lambrechts L, Paaijmans KP, Fansiri T, Carrington LB, Kramer LD, Thomas MB, et al. Impact of
daily temperature fluctuations on dengue virus transmission by *Aedes aegypti*. Proceedings
of the National Academy of Sciences. 2011;108(18):7460-5.

26.   Bar-Zeev M. The Effect of Temperature on the Growth Rate and Survival of the Immature Stages
of Aëdes aegypti (L.). Bulletin of Entomological Research. 2009;49(1):157-63.

27.   Rueda LM, Patel KJ, Axtell RC, Stinner RE. Temperature-Dependent Development and Survival
Rates of Culex quinquefasciatus and Aedes aegypti (Diptera: Culicidae). Journal of Medical Entomology.
1990;27(5):892-8.

28.   Tun-Lin W, Burkot TR, Kay BH. Effects of temperature and larval diet on development rates and
survival of the dengue vector Aedes aegypti in north Queensland, Australia. Medical and Veterinary
Entomology. 2000;14(1):31-7.

29.   Nagao Y, Thavara U, Chitnumsup P, Tawatsin A, Chansang C, Campbell-Lendrum D. Climatic and social risk factors for Aedes infestation in rural Thailand. Tropical Medicine & International Health. 2003;8(7):650-9.

30.   Naish S, Dale P, Mackenzie JS, McBride J, Mengersen K, Tong S. Climate change and dengue: a critical and systematic review of quantitative modelling approaches. BMC Infectious Diseases. 2014;14(1):167.

31.   Chakravarti A, Kumaria R. Eco-epidemiological analysis of dengue infection during an outbreak of dengue fever, India. Virology Journal. 2005;2(1):32.

32.   Vanwambeke SO, Lambin EF, Eichhorn MP, Flasse SP, Harbach RE, Oskam L, et al. Impact of Land-use Change on Dengue and Malaria in Northern Thailand. EcoHealth. 2007;4(1):37-51.

33.   Patz JA, Daszak P, Tabor GM, Aguirre AA, Pearl M, Epstein J, et al. Unhealthy Landscapes: Policy Recommendations on Land Use Change and Infectious Disease Emergence. Environmental Health Perspectives. 2004;112(10):1092-8.

34.   Norris DE. Mosquito-borne Diseases as a Consequence of Land Use Change. EcoHealth. 2004;1(1):19-24.

35.   Cheong YL, Leitão PJ, Lakes T. Assessment of land use factors associated with dengue cases in Malaysia using Boosted Regression Trees. Spatial and Spatio-temporal Epidemiology. 2014;10:75-84.

36.   Sarfraz MS, Tripathi NK, Tipdecho T, Thongbu T, Kerdthong P, Souris M. Analyzing the spatio-temporal relationship between dengue vector larval density and land-use using factor analysis and spatial ring mapping. BMC Public Health. 2012;12(1):853.

37.   Vanwambeke SO, van Benthem BHB, Khantikul N, Burghoorn-Maas C, Panart K, Oskam L, et al. Multi-level analyses of spatial and temporal determinants for dengue infection. International Journal of Health Geographics. 2006;5(1):5.

38.   Sirisena PDNN, Noordeen F. Evolution of dengue in Sri Lanka—changes in the virus, vector, and climate. International Journal of Infectious Diseases. 2014;19:6-12.

39.   Vitarana T, Jayakuru W, Withane N. Historical account of Dengue Haemorrhagic Fever in Sri Lanka: WHO Regional Office for South-East Asia. ; 1997 [Available from: http://www.who.int/iris/handle/10665/148517.

40.   Messer WB, Vitarana UT, Sivananthan K, Elvtigala J, Preethimala LD, Ramesh R, et al. Epidemiology of dengue in Sri Lanka before and after the emergence of epidemic dengue hemorrhagic fever. The American Journal of Tropical Medicine and Hygiene. 2002;66(6):765-73.

41.   Ali S, Khan AW, Taylor-Robinson AW, Adnan M, Malik S, Gul S. The unprecedented magnitude of the 2017 dengue outbreak in Sri Lanka provides lessons for future mosquito-borne infection control and prevention. Infection, Disease & Health. 2018;23(2):114-20.

42.   World Health Organization. Situation Report #01, Sri Lanka dengue outbreak 2017 [Available from: http://www.searo.who.int/srilanka/areas/dengue/dengue_outbreak_2017_sit_rep_1.pdf.

43.   World Health Organization. Dengue fever - Sri Lanka: World Health Organization; 2017
[Available from: http://www.who.int/csr/don/19-july-2017-dengue-sri-lanka/en/.

44.   Favier C, Schmit D, Müller-Graf CDM, Cazelles B, Degallier N, Mondet B, et al. Influence of spatial
heterogeneity on an emerging infectious disease: the case of dengue epidemics. Proceedings of the
Royal Society B: Biological Sciences. 2005;272(1568):1171-7.

45.   Epidemiology Unit in Ministry of Health of Sri Lanka. Distribution of Notification(H399) Dengue
Cases by Month 2017 [Available from:
http://www.epid.gov.lk/web/index.php?option=com_casesanddeaths&Itemid=448&lang=en.

46.   National Centers for Environmental Information (NCEI). Global Surface Summary of the Day
(GSOD) U.S.: NOAA/NESDIS/NCEI; 2015 [updated February 8, 2018. Available from:
https://catalog.data.gov/dataset/global-surface-summary-of-the-day-gsod.

47.   Wickramagamage P. Seasonality and spatial pattern of rainfall of Sri Lanka: Exploratory factor
analysis. International Journal of Climatology. 2010;30(8):1235-45.

48.   Bright EA, Rose AN, Urban ML, McKee JJ. LandScan 2016. 2016 ed. Oak Ridge, TN: Oak Ridge
National Laboratory; 2017.

49.   Survey Department of Sri Lanka. Spatial Data Infrastructure 2018 [Available from:
https://www.survey.gov.lk/nsdi/lis/index_cad.php.

50.   Romeo-Aznar V, Paul R, Telle O, Pascual M. Mosquito-borne transmission in urban landscapes:
the missing link between vector abundance and human density. Proceedings of the Royal Society B:
Biological Sciences. 2018;285(1884).

51.   Schmidt W-P, Suzuki M, Dinh Thiem V, White RG, Tsuzuki A, Yoshida L-M, et al. Population
Density, Water Supply, and the Risk of Dengue Fever in Vietnam: Cohort Study and Spatial Analysis.
PLOS Medicine. 2011;8(8):e1001082.

52.   Gubler DJ. Dengue and Dengue Hemorrhagic Fever. Clinical Microbiology Reviews.
1998;11(3):480-96.

53.   Gubler DJ. Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic
problem in the 21st century. Trends in Microbiology. 2002;10(2):100-3.

54.   Ramasamy R, Surendran SN, Jude PJ, Dharshini S, Vinobaba M. Larval Development of Aedes
aegypti and Aedes albopictus in Peri-Urban Brackish Water and Its Implications for Transmission of
Arboviral Diseases. PLOS Neglected Tropical Diseases. 2011;5(11):e1369.

55.   Gardner L, Sarkar S. A Global Airport-Based Risk Model for the Spread of Dengue Infection via
the Air Transport Network. PLOS ONE. 2013;8(8):e72129.

56.   Gardner LM, Fajardo D, Waller ST, Wang O, Sarkar S. A Predictive Spatial Model to Quantify the
Risk of Air-Travel-Associated Dengue Importation into the United States and Europe. Journal of Tropical
Medicine. 2012;2012.

57.   World Health Organization. Dengue and severe dengue 2018 [Available from:
http://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue.

58.   Mammen MP, Jr., Pimgate C, Koenraadt CJM, Rothman AL, Aldstadt J, Nisalak A, et al. Spatial
and Temporal Clustering of Dengue Virus Transmission in Thai Villages. PLOS Medicine. 2008;5(11):e205.

59.   Yoon I-K, Getis A, Aldstadt J, Rothman AL, Tannitisupawong D, Koenraadt CJM, et al. Fine Scale
Spatiotemporal Clustering of Dengue Virus Transmission in Children and Aedes aegypti in Rural Thai
Villages. PLOS Neglected Tropical Diseases. 2012;6(7):e1730.