Activity-dependent production of ATP and ROS in mitochondria of myelinated axons in healthy and neuropathic conditions

Short title: ATP and ROS production in axonal mitochondria

Gerben van Hameren[1]*, Graham Campbell[1], Marie Deck[1], Jade Berthelot[1], Roman Chrast[2], Nicolas Tricaud[1]*

[1] Institut des Neurosciences de Montpellier, INSERM U1051, Université de Montpellier, Montpellier, France

[2] Departments of Neuroscience and Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

* Corresponding authors: gerben.van-hameren@inserm.fr

                             nicolas.tricaud@inserm.fr

Author contributions

GvH conducted the experiments. GvH and NT designed the experiments and wrote the paper with contributions from GC and RC. GC, MD and JB contributed to the experiments and to the design of the experiments. NT supervised the project.

Abstract

Mitochondria are critical for the function and the maintenance of myelinated axons notably through the ATP production. A by-product of this activity is reactive oxygen species (ROS), which are highly deleterious for neurons. While ROS and metabolism are involved in several neurodegenerative diseases, it is still unclear how axonal activity or myelin modulates ATP and ROS production in axonal mitochondria. We imaged and quantified mitochondrial ATP and hydrogen peroxide (H_2O_2) in resting or stimulated peripheral nerve myelinated axons *in vivo,* using genetically-encoded fluorescent probes, two-photon time-lapse and CARS imaging. ATP and H_2O_2 production is intrinsically higher in nodes of Ranvier even in resting conditions. Axonal firing increased both ATP and H_2O_2 productions but with different dynamics. In neuropathic $MFN2^{R94Q}$ mice, mimicking Charcot-Marie-Tooth 2A disease, defective mitochondria failed to upregulate ATP production following axonal activity such as control mice. However, H_2O_2 production was dramatically sustained. Moreover, mimicking demyelinating peripheral neuropathy resulted in a reduced production of ATP while H_2O_2 level soared. Taken together, our results suggest that ATP and ROS productions are decoupled under neuropathic conditions, which may compromise axonal function and integrity.

Keywords: Axons**,** axonal activity, mitochondria, ATP**,** ROS, PNS, demyelination, MFN2.

1. Introduction

While the nervous system, and the brain in particular, represents around 2% of the body mass, it consumes up to 20% of the glucose we mobilize every day [1]. This high energy

expenditure in the nervous system is firstly due to the synaptic activity that requires high

amounts of adenosine tri-phosphate (ATP) [1]. A second energy demanding process is the

propagation of action potentials (APs) along axons. Indeed, conduction of APs involves ion

exchanges through the plasma membrane, first through voltage-gated ion channels to

depolarize, then through the sodium-potassium ATPase to repolarize [2,3]. Therefore, the

production of ATP in axons is crucial for repeated regeneration of APs. Both in the central

nervous system (CNS) and in the peripheral nervous system (PNS), myelination significantly

reduces the energy cost of AP propagation through the sequestration of the AP firing

machinery at the node of Ranvier and axon initial segment [4].

Mitochondria appear to be the main source of cellular ATP and these organelles are abundant

in CNS [5,6] and PNS [7] axons. Some reports indicated that mitochondria were more

abundant in the node of Ranvier [8]. In addition, axonal AP activity and axo-glial junctions

regulates the recruitment of mitochondria in the nodal area [9]. Therefore, a well-accepted

idea is that axonal AP activity could locally stimulate ATP production in axonal mitochondria

[10]. However, this nodal enrichment in mitochondria remains controversial [11] and the

dynamic of local mitochondrial ATP production following AP is actually unknown.

As an intrinsic by-product of ATP production, mitochondria also produce reactive oxygen

species (ROS) [12,13]. The mitochondrial electron transport chain (ETC) and NAPDH

oxidases are the major sources of superoxide ($O_2^{\bullet-}$), a highly reactive type of ROS, within

cells [14]. $O_2^{\bullet-}$ is rapidly converted to H_2O_2 by superoxide dismutase (SOD) and H_2O_2 is then

reduced to H_2O and O_2 by mitochondrial glutathione peroxidase (GPx) and cytosolic enzymes

[15]. H_2O_2 can also react with iron or $O_2^{\bullet-}$ to form hydroxyl radicals (OH•), which is then

reduced by molecular hydrogen to H_2O [16]. If not reduced by antioxidants, all types of ROS

are highly toxic for the cell and in particular for mitochondria, where they are produced.

Indeed ROS can directly oxidize and damage DNA, carbohydrates, proteins or lipids [17].

In several neurodegenerative diseases and axonopathies, such as Parkinson's disease [18],

multiple sclerosis (MS) [19] and Charcot-Marie-Tooth diseases (CMT) [20], mitochondrial

dysfunction and increased levels of ROS have been shown to be involved. Whether axonal

mitochondria are involved is still unclear and investigating the production of ATP and ROS

by these mitochondria in active or diseased axons in real time *in vivo* would constitute a first

answer.

The importance of mitochondrial homeostasis for the normal function of myelinated axons

[9,21] is highlighted by mutations in the *mitofusin 2* (*MFN2*) gene. Indeed, MFN2 has several

functions in mitochondria physiology, among which mitochondrial fusion [22] and transport

along axons [23], and mutations in its gene result in axonal form of peripheral neuropathy

(CMT2A) in humans and in mice [24,25]. In particular, MFN2 loss-of-function in axonal

mitochondria decreases the expression of oxidative phosphorylation subunits [26], suggesting

that *MFN2* mutations also affect ATP and ROS production. However, this remains to be

shown *in vivo*.

MS is a chronic inflammatory disease affecting myelinated tracts in the brain [27]. MS is

characterized by demyelinated lesions resulting from the loss of the myelin sheath and the

progressive form of MS shows successive events of myelin degeneration and regeneration

that damage axons [27]. A common hypothesis is that demyelinated axons are more

demanding in ATP in order to maintain their functions and their integrity [28]. Recent data

showed that axonal mitochondria complex IV activity is increased in MS lesions and the total

mitochondrial mass is increased [29], suggesting that axonal mitochondria are indeed

producing more ATP. In addition, while activated microglia and macrophages are thought to

be the initial source of ROS, axonal mitochondria dysfunction and an increase of

mitochondrial ROS contribute to progressive MS [30]. Taken together this suggests that the

myelin sheath plays a significant role in the maintenance of axonal mitochondria homeostasis

and in particular in the production of ATP and ROS in mitochondria.

To answer these questions, we set up an *in vivo* approach using a H_2O_2-sensitive GFP

(roGFP-Orp1) [31] and a fluorescent ATP sensor (AT1.03) [32] targeted to mitochondria to

image and measure the dynamics of ATP and H_2O_2 production in mitochondria of myelinated

axons in myelinated axons of the PNS. Using this approach, we show that both ATP and H_2O_2

levels are increased in mitochondria residing in nodes of Ranvier and that nerve stimulation

sharply increases ATP and H_2O_2 production by mitochondria in few minutes. While MFN2

mutations had no effect on basal levels of ATP and H_2O_2 in axons, it prevented the increase of

ATP production after nerve stimulation. At the opposite, an increased H_2O_2 production was

observed. Moreover, pathological demyelination reduced ATP production while enhancing

mitochondrial ROS, showing that both neuropathic conditions decoupled the production of

ATP from the production of ROS.

2. Results

2.1. Validation of the mito-roGFP-Orp1 and mito-AT1.03 probes

We used AT1.03, which is a genetically-encoded fluorescence probe based on fluorescence

resonance energy transfer (FRET), to detects ATP [32]. This probe consists of the cyan

fluorescent protein (CFP) mseCFP, the yellow fluorescent protein (YFP) variant monomeric

Venus, both linked at the ε-subunit of the Bacillus subtilis FoF1-ATP synthase. In the

presence of ATP, the ε-subunit retracts to bring the two fluorescent proteins close to each

other, thereby increasing FRET efficiency. The conformational change is reversible. Relative

ATP levels are measured using the YFP/CFP fluorescence ratio.

Redox sensitive GFPs (roGFPs) harbor an engineered dithiol/disulfide switch on their

surface, which determines their wavelength of excitation [33]. RoGFP2 has been converted

into an H_2O_2 specific probes by fusion with the microbial H_2O_2 sensor Oxidant Receptor

Peroxidase 1 (Orp1) [31]. The reaction is reversible via reduction by cellular thioredoxin

(Trx) or GRx/GSH [34]. Relative H_2O_2 levels are measured using the ratio of emitted

fluorescence lights when excited at 800nm (oxidized form) or 940nm (reduced form). All

experiments were done in the same imaging conditions and analysis. As both probes are

ratiometric, the measured values are independent of the number of mitochondria or of the

absolute fluorescence intensity of the probe. Both probes were targeted to the mitochondrial

matrix.

We first validated mito-roGFP-Orp1 and mito-AT1.03 probes *in vivo*. Injection of AAV9-

mito-roGFP-Orp1 and AAV9-mito-AT1.03 in the spinal cord of mouse pups one day after

birth (P1) resulted in expression of the fluorescent probes in multiple axons that are part of

sciatic and saphenous nerves (Fig 1A). Coherent Anti-stokes Raman Scattering (CARS)

microscopy uses the non-linear interactions between light and molecules to generate light

from lipids without labeling [35]. As myelin is enriched with lipids, CARS microscopy allows

visualizing the myelin sheath around axons *in vivo* [36,37,38]. Using this technique, we

observed that the fluorescent probe signal was encompassed by the myelin sheath, which

shows the probes are expressed in myelinated axons (Fig 1A). We then performed an

immunostaining of teased fibers for the mitochondrial marker TOM20 and observed a partial

145 colocalisation between fluorescent probes and mitochondria (Fig 1B), showing probes are

146 expressed in axonal mitochondria, but not all of them. Consistently, mitochondria located in

147 Schwann cells surrounding axons were labeled with TOM20 antibody but not with fluorescent

148 probes. Both probes could be detected in axons of the mouse saphenous and sciatic nerves *in*

149 *vivo*.

150   Then the ratio of the fluorescent intensity of oxidized/reduced roGFP-Orp1 (R) was

151 followed over time (Fig 1C), showing a small and slow increase of the ratio (0.206 ± 0.002

152 per hour) probably due to air oxidation. Addition of reducer DTT on the nerve resulted in a

153 significant decrease of the fluorescence ratio, while injection of oxidizer diamide resulted in a

154 significant increase of fluorescence ratio (Fig 1D). These results show that the

155 dithiol/disulfide switch of mito-roGFP-Orp1 is an efficient and reliable indicator of relative

156 redox changes in axonal mitochondria of mouse peripheral nerves *in vivo*. The fluorescence

157 ratio YFP/CFP of mito-AT1.03 (R) was very stable over time (Fig 1E). Injection of 0.4M and

158 1M ATP into the sciatic nerve in an increase of AT1.03 fluorescence ratio (Fig 1F), showing

159 this probe is also functional *in vivo*.

Figure 1

Figure 1: *In vivo* validation of the fluorescent probes.

(A) Several axons expressing mito-roGFP-Orp1 or mito-AT1.03 probe can be observed in teased fibers of mouse sciatic nerve 1 month after the virus injection. CARS imaging (inserts) shows that axons expressing mito-roGFP-Orp1 or mito-AT1.03 (green) are surrounded by a myelin sheath (red). Scale bars = 3 µm. (B) mito-roGFP-Orp1 expression (green) partially colocalizes with mitochondrial marker TOM20 (red) in the axon. Similarly, mito-AT1.03 partially colocalizes with TOM20 in axonal mitochondria. Scale bars = 3 µm. (C) A fluorescent signal of mito-roGFP-Orp1 is detected when the probe is reduced and when it is oxidized by H_2O_2. The ratio of mito-roGFP-Orp1 fluorescence measured *in vivo* (n=6 axons; 6 mice) shows a slight increase over time. Scale bar = 10 µm (D) Mito-roGFP-Orp1 fluorescence ratio decreases with DTT (p=0.006; n=3 axons, 3 mice) and increases after injection of diamide into the nerve (p=0.003; n=4 axons, 4 mice). (E) A fluorescent signal of the CFP subunit and YFP subunit of mito-AT1.03 is detected. Mito-AT1.03 fluorescence ratio measured *in vivo* shows no significant change over time (n=18 axons; 3 mice). Scale bar = 10 µm. (F) Mito-AT1.03 fluorescence ratio increases after injection of 0.4M (p=0.06; n=3 axons, 3 mice) and 1M (p<0.001; n=6 axons, 4 mice) ATP into the sciatic nerve. All error bars show SEM. Statistical analysis shows Student two-tailed T-tests.

2.2 The effect of nerve stimulation on ATP and H_2O_2 production

While ATP is supposed to be critical for the firing of axons, ATP production in mitochondria

of functionally active axons has never been observed *in vivo*. We used the real-time imaging

approach previously described in combination with a setup for electrical nerve stimulation

based on previous work [39] (Fig 2A-D). We used the saphenous nerve, which is mostly

composed of sensory axons, hence its stimulation leads to only limited unwanted contraction

of the paw's muscles [40]. After recording the probe fluorescence for at least 20 minutes

without stimulation, APs were induced by an electrical burst stimulation protocol including 3

bursts of 50Hz for 30 seconds spaced by a 60 second recovery period (Fig 2E).

Figure 2: Setup of the saphenous nerve imaging experiment.

(A) After deep anesthesia, the mouse is placed on its back, the skin of its left inner tight is removed and the saphenous nerve (N) is placed on a plastic strip (S). Two stimulation microelectrodes (1 and 2) are inserted on both sides of the nerve on the plastic strip and one hook-shaped recording electrode (3) is holding the nerve on the opposite side. (B) The ground electrode (4) is inserted in the mouse tail and the negative electrode (5) in the groin area. (C) A glass coverslip (Co) is then placed on the nerve soaked in PBS and the mouse is then placed under the two-photon microscope immersion lens 20X (L) in water. (D) Schematic view of the imaging setup. (E) Schematic representation of the nerve stimulation pattern used to induce APs. The generation of APs was verified using the recording electrode. Negative electrode was used to correct for background signal.

173
174

The fluorescent signal of mito-AT1.03 was recorded every 5 minutes for 20 minutes after

stimulation (Fig 3A). At each time point the fluorescence ratio (R) was corrected for the mean

fluorescence ratio of that same axon before nerve stimulation (R0). After a first stimulation,

the fluorescence ratio of mito-AT1.03 increased in several axons, but not in all of them

(44%), resulting in a slight non-significant average increase (Fig 3a). After the second

stimulation, all axons responded significantly increasing mitochondrial ATP production (Fig

3a). In the same conditions, nerve stimulations significantly increased mitochondrial H_2O_2

production just 1 minute after the stimulation period (Fig 3B). H_2O_2 level then quickly went

back and stabilized at pre-stimulation values (Fig 3B). These data show that axonal

mitochondria very quickly adapt to the axonal activity up-regulating their production of ATP.

Spikes in mitochondrial H_2O_2 production were also observed just before the surge of ATP in

mitochondria (Fig 3C), suggesting that this H_2O_2 production reflects the oxidative

phosphorylation process.

Figure 3: Effect of burst nerve stimulation on mitochondrial ATP and H₂O₂ levels.

(A) Upper panels: Nerve stimulation induced changes in the fluorescence signal of both the CFP and YFP subunit of mito-AT1.03 as illustrated by the YFP/CFP overlay pictures. Lower panel: graph showing mito-AT1.03 fluorescence ratio normalized to pre-stimulation values (R/R0) following two successive nerve stimulation period (black bars at the top). The first stimulation results in a slight, non-significant increase; a significant increase is measured after a second stimulation ($p=0.0171$ at $t=45$, $p=0.0001$ at $t=50$, $p=0.049$ at $t=55$; F-value = 2.872; Df=15; n=9 axons in 3 mice). (B) Upper panels: Nerve stimulation induced changes in the fluorescence signal of both the oxidized and reduced forms of GFP in mito-roGFP-Orp1 as illustrated by the overlay pictures. Lower panel: graph showing mito-roGFP-Orp1 fluorescence ratio normalized on pre-stimulation values (R/R0) following two successive nerve stimulation (black bars at the top). Both stimulations result in a significant increase 5 minutes after the stimulation ($p=0.007$; $p=0.04$; F-value = 2.804; Df=14; n=14 axons in 7 mice). (C) Graphs described in a and b were overlaid to show the relative dynamics of ATP and ROS levels after nerve stimulations. Scale bars= 5 µm. Error bars show SEM. Statistical tests are one-way ANOVA.

2.3. ATP and H2O2 production are altered in CMT2A neuropathic mice

We recently showed that MFN2[R94Q] mice, a model for CMT2A disease [25] where MFN2 is

defective, displayed altered mitochondria motility and clustering in peripheral nerve axons

[41]. We used this model to measure the level of ATP in axonal mitochondria of myelinated

axons in neuropathic conditions. In non-stimulated conditions, control and mutant mice

axonal mitochondrial ATP production were similar (Fig 4A). After the first stimulation,

similar to control mice, mitochondrial ATP production increased in some axons whereas

others did not respond, resulting in statistically non-significant variation (Fig 4B). However,

after a second period of nerve stimulation, as opposed to controls, MFN2[R94Q] axonal

mitochondria failed to upregulate ATP production (Fig 4B-C). This indicates that

dysfunctional MFN2 does not hinder the basal production of ATP by axonal mitochondria but

impairs their ability to up-regulate their production in response to axonal activity.

Similar to ATP, we found no difference between controls and MFN2[R94Q] mice in the

basal H_2O_2 levels (Fig 4D). When nerves were stimulated, mitochondrial H_2O_2 significantly

increased after each stimulation period (Fig 4E). However, this increase was delayed

compared to controls and H_2O_2 levels remained high for longer time (Fig 4F), showing that

axonal mitochondria produce more deleterious H_2O_2 than control mice in response to axonal

activity. Since ROS production in the mitochondrial matrix reflects the oxidative

phosphorylation process [42,43], these data indicate that oxidative phosphorylation is

decoupled from ATP production in mitochondria of these neuropathic mice.

Figure 4

Figure 4: ATP and H$_2$O$_2$ levels of MFN2^{R94Q} mice.

(A) Graph showing mito-AT1.03 fluorescence ratio in resting axonal mitochondria of wild-type or mutant MFN2^{R94Q} mice (Student two tailed T-test p=0.809; n=18 axons, 7 mice per group). (B) Graph showing mito-AT1.03 fluorescence ratio normalized on pre-stimulation values (R/R0) following two successive nerve stimulation periods (black bars at the top). The first period of stimulation resulted in a slight, non-significant increase; the second stimulation period did not generate any change (n=14 axons in 5 mice). (C) Graphs showing mito-AT1.03 R/R0 in wild-type (Fig. 3A) and mutant MFN2^{R94Q} mice (B panel) were overlaid to show the relative dynamics of ATP levels in both genotypes after nerve stimulations (Two way ANOVA, p=0.011 at t=45, p<0.001 at t=50, p=0.02 at t=55, F-value=19,27; Df=1). (D) Graph showing mito-roGFP-Orp1 fluorescence ratio in resting axonal mitochondria of wild-type or mutant MFN2^{R94Q} mice (Student two tailed T-test p=0.337; n=16 axons, 8 mice per group). (E) Graph showing mito-roGFP-Orp1 fluorescence ratio normalized on pre-stimulation values (R/R0) following two successive nerve stimulation periods (black bars at the top). Both stimulations result in a significant increase (p=0.013 at t=15, p=0.020 at t=20, p=0.044 at t=25 and p=0.038 at t=45; F-value=3.095; Df=14; n=11 axons in 4 mice). (F) Graphs showing mito-roGFP-Orp1 R/R0 in wild-type (Fig. 3B) and mutant MFN2^{R94Q} mice (E panel) were overlaid to show the relative dynamics of H$_2$O$_2$ levels in both genotypes after nerve stimulations. A significant higher H$_2$O$_2$ level is detected in wild-type first (Two way ANOVA, p=0.007 at t=5, F-value= 5,484; Df=1), then a higher H$_2$O$_2$ level is detected in mutant MFN2^{R94Q} (p=0.014 at t=15, p=0.047 at t=40). Error bars show SEM. # = p-value<0.05. ## = p-value<0.01. ### = p-value<0.001.

2.4. ATP and H2O2 production in nodes of Ranvier and internodes

The enrichment of mitochondria at the node of Ranvier and whether these nodal mitochondria

are more metabolically active remains controversial. In order to detect potential spatial

differences between nodal and internodal mitochondria, we used CARS microscopy. Gaps in

CARS signal between two myelinated internodes are nodes of Ranvier (Fig 5A) [38,43,44].

Combining two-photon imaging of fluorescent probes with CARS imaging allowed us to

analyze mitochondrial physiology in nodes of Ranvier versus internodes defined as areas

distant of more than 5 μm from the node (Fig 5A). In all observed nodes of Ranvier, probes-

labeled mitochondria were present, but they were not more abundant or larger than in

internodes (Fig 5A). However, when measuring the mito-AT1.03 and mito-roGFP-Orp1

fluorescence ratio, we found significantly higher ATP and H_2O_2 levels in node of Ranvier

mitochondria (Fig 5B-C). These data, obtained in resting animal without nerve stimulations,

indicate that nodal mitochondria are intrinsically more metabolically active than internodal

ones. In neuropathic MFN2^{R94Q} mice, nodal mitochondria did not show a higher level of ATP

versus internodal mitochondria (Fig 5B). However, they still had higher level of H_2O_2 (Fig

5C), suggesting again a decoupling between oxidative phosphorylation and the production of

ATP in mitochondria of these mutant mice.

Figure 5: ATP and H$_2$O$_2$ levels are different along axons.

(A) Two-photons imaging of CFP (blue) and YFP (yellow) fluorescence of mito-AT1.03 was combined with CARS imaging of the myelin sheath (red). CARS imaging gap shows a node of Ranvier (arrowhead). Using the combined images, mitochondria located in the node of Ranvier and mitochondria located in internodes can be identified (right panel). Scale bar = 10 µm (B) Graph showing mito-AT1.03 fluorescence ratio in axonal mitochondria located in nodes of Ranvier (N) or internodes (IN) of wild-type (WT) or MFN2^{R94Q} mice (MFN2 mutant). In wild-type mice, ATP levels are higher in mitochondria in nodes of Ranvier than in internodes (p=0.035; n=14 axons in 7 mice). In MFN2^{R94Q} mice, ATP levels in node of Ranvier mitochondria are equal to internode mitochondria (n=7 nodes in 3 mice). (C) Graph showing mito-roGFP-Orp1 fluorescence ratio in axonal mitochondria located in nodes of Ranvier or internodes of wild-type or mutant MFN2^{R94Q} mice. In both wild-type mice and mutant MFN2^{R94Q} mice, H$_2$O$_2$ levels are higher in mitochondria in nodes of Ranvier than in internodes (WT: p=0.009; n=8 axons in 3 mice. MFN2^{R94Q}: p=0.019; n=9 axons in 3 mice). Error bars show SEM. Statistical tests are Paired T-tests.

2.5. ATP and H$_2$O$_2$ production in mitochondria of demyelinated axons

The myelin sheath is critical to maintain the node of Ranvier on myelinated axons and is

therefore likely to play a significant role in the homeostasis of axonal mitochondria. As

demonstrated in Fig 5a, the myelination state of the axons in the mouse sciatic nerve can be

assessed *in vivo* by CARS imaging. We therefore investigated how this demyelination affects

axonal mitochondria physiology *in vivo*. To induce demyelination we used lyso-

phosphatidylcholine (LPC) and we followed the production of mitochondrial ATP and H$_2$O$_2$

in axons during demyelination and during the restoration of the myelin sheath

(remyelination). LPC, which is a signaling molecule resulting from the hydrolysis of

phophatidylcholine by phospholipase A2 [45] acts in mSC and oligodendrocytes as a

demyelinating signal [46,47,48]. LPC was injected in the sciatic nerve of adult mice and

demyelination occurred immediately to culminate one week after the injection with the

formation of myelin ovoids and debris as seen using CARS (Fig 6A, week 1) [37,49]. As soon

as two weeks after LPC injection, remyelination occurred and some thin myelin sheaths

reappeared around axons (Fig 6A, week 2). Axons were completely remyelinated three weeks

after injection of LPC (Fig 6a, week 3). No demyelination occurred in control mice injected

with PBS (Fig 6A last panel). Axonal integrity is not directly affected by LPC (S1 Figure)

[50] and axonal mitochondria remained visible using either mito-AT1.03 or mito-roGFP-Orp1

probes during the entire process of demyelination and remyelination (Fig 6A inserts).

Figure 6: Impact of demyelination on axonal mitochondria ATP and H_2O_2.

(A) CARS imaging is used to visualize myelin. At week 0, before injection of LPC, most axons are myelinated. 1 week after LPC injection, axons are demyelinated and myelin debris and ovoids are observed. At week 2, thinly myelinated axons are observed in between the myelin debris (arrowheads) and at week 3, almost all axons are myelinated again, thus resembling a healthy nerve. PBS injection induces small conformational changes, but no formation of ovoids or debris. Inserts show that neuronal mitochondria remain visible throughout the whole process. (B) Graph showing mito-AT1.03 fluorescence ratio (R) in axonal mitochondria normalized on pre-demyelination values (R0) following demyelination. Following LPC injection, ATP levels are unchanged after 1 week (p=0.085; n=5 mice; 27 axons), decreased after 2 weeks (p=0.019; n=3 mice, 24 axons) and restored after 3 weeks (p=0.491; n=5 mice; 31 axons). PBS injection results in no significant change (p=0.799; n=3 mice; 23 axons). (C) Graph showing mito-roGFP-Orp1 fluorescence ratio (R) in axonal mitochondria normalized on pre-demyelination values (R0) following demyelination. Following LPC injection, H_2O_2 levels are increased after 1 week (p=0.031; n=10 mice, 17 axons), and unchanged after 2 weeks (p=0.663; n=5 mice, 9 axons) and after 3 weeks (p=0.450; n=8 mice, 13 axons). PBS injection results in no significant change (p=0.121; n=3 mice, 5 axons).

We therefore measured the fluorescence ratio for these probes in non-stimulated axonal

mitochondria during the process. One week after LPC injection, when axons are completely

demyelinated, ATP production was slightly lower than before demyelination, but not

significantly (Fig 6B), while H_2O_2 levels peaked (Fig 6C). ATP levels were significantly

lower in axonal mitochondria at two weeks after LPC injection, while H_2O_2 levels were back

to control and pre-demyelination levels (Fig 6B-C). Finally, three weeks after injection, when

remyelination is completed, ATP levels were back to control and pre-demyelination levels as

are the H_2O_2 levels (Fig 6B-C). No change either in ATP or H_2O_2 levels were detected at one

week after injection with PBS in control mice (Fig 6B-C). Moreover, increase in ROS did not

result from a general oxidative stress in the demyelinated nerve, since CellRox dye did not

show any significant increase, except at the LPC injection site where we did not image our

probe (S1 Fig). Taken together, these data indicate that the myelin sheath has a profound

impact on axonal mitochondria physiology. Mitochondria manage to maintain their ATP

production during myelin break-down, but they generate more H_2O_2. The process of

remyelination leads to decreased production of ROS, while ATP production is temporally

diminished before full remyelination. These data therefore indicate that demyelination induces

the decoupling of ATP production from oxidative phosphorylation.

3. Discussion

Although it has been shown that PNS axonal mitochondria, like most mitochondria, produce

both ATP and ROS, the regulations of their production are still not well understood *in vivo*

and *in situ*. We used a combination of viral delivery of mitochondria-targeted fluorescent

probes to active peripheral neurons and two-photon and CARS live imaging in mice to fill

that gap. The mito-AT1.03 and mito-roGFP-Orp1 fluorescent probes had been previously

validated *in vitro* [51] and *in vivo* [33] in other systems but we also validated them *in vivo* in

our system. We did not attempt to measure absolute values of ATP or H_2O_2 with each probe

fluorescence *in vivo* because firstly, this was difficult as probes are targeted to mitochondrial

matrix and secondly, it was not necessary as we aimed to compare different genotypes and

conditions in particular with time. So only relative values obtained in the same imaging

conditions are shown here. We found that the range of detection was large enough to cover

the changes occurring in axonal mitochondria *in vivo*. Moreover the probe sensitivity was

sufficient to observe variations in real-time using a 5 minutes delay between measures. This

delay was partly imposed by relative low speed of our imaging system (1 minute to create 10

scans of $200 \mu m^2$ over 40 μm depth for each wavelength), but also by the intrinsic limits of our

set-up: electrical stimulation induced slight contractions of saphenous nerve surrounding

muscles which made the pictures blurred during the stimulation. The probes alterations were

also reversible fast enough to observe changes in both directions. Mito-AT1.03 was stable,

but mito-roGFP-Orp1 showed a slight increase of its fluorescence ratio over time under our

experimental settings. These results are not explained by photobleaching or degradation of the

probe, since the total fluorescent signal does not decrease. This change in ratio of

fluorescence may be caused by air oxidation [52]. However, buffered artificial cerebrospinal

fluid as an incubation medium for the saphenous nerve did not prevent the probe oxidation.

Nonetheless, this slow oxidation of mito-roGFP-Orp1 had negligible effects on our

experiments, since it was too slow to cause significant changes in our relatively short term

imaging.

Axonal activity induced stereotypic changes in ATP and H_2O_2 production by axonal

mitochondria. Shortly after the stimulation, H_2O_2 production increased and dropped

immediately and then ATP increased for several minutes. This was stereotypic, because it

occurred almost identically after both stimulations we did. After the first stimulation, not all

axonal mitochondria produced more ATP, suggesting that either our stimulation protocol was

not mobilizing enough axons or axonal mitochondria are deactivated, similar to some ion

channels. This limitation was lifted by the second stimulation. On the attempts we did for

further stimulations, sometimes the ATP level was maintained at high range and sometimes

mitochondria did not respond anymore. The reason for these variations in the long term is

unknown and we did not pursue this further. However, taken together, this suggests that ATP

production in axonal mitochondria is positively correlated with the level of axonal activity

and can eventually reach a plateau in our conditions. Due to technical reasons we could not

measure H_2O_2 levels in mitochondria after several stimulations. However, looking at the first

two stimulations, we conclude that H_2O_2 production in mitochondria is also positively

correlated with axonal activity but not cumulative. It would be interesting to look further for

the molecular mechanisms that allow mitochondria to detect axolemma depolarization or ion

channels activity on a very short period of time (around one minute in our detection

capacities). This fast kinetic suggests the role of ions, such as calcium ions, which are

mobilized during AP firing at the node of Ranvier [53], or kinases. A mechanism involving

calcium has been shown to anchor mitochondria at the node of Ranvier [54], suggesting that

mitochondria are associated with the axolemma there. Consistently, we found nodal

mitochondria to be more metabolically active than internodal ones. However, our data

following stimulations were recorded on mitochondria independently of their location on

axons. As the node is much smaller than the internodes (1 μm versus 600 μm in average in the

mouse nerve) the probability we recorded nodal mitochondria is extremely small. So the

molecular mechanism that mobilizes mitochondria after AP firing has to exist also in

internodal mitochondria.

Looking at the known origin of ROS in mitochondria, our data suggest that the H_2O_2

production we detected in axonal mitochondria reflects the loading of the mitochondrial

matrix with protons before the production of ATP. However, since the H_2O_2 production drops

very quickly and, contrary to the ATP production, it is not cumulative, a more complex

scheme has to be taken in consideration. Indeed, H_2O_2 is formed by SOD enzyme from $O_2^{\bullet -}$,

which directly results from the ETC. Moreover, H_2O_2 is then changed in H_2O and O_2 by

several reducing enzymes, such as mitochondrial GPx, so actually the amount of H_2O_2 we

detected is the result of equilibrium between $O_2^{\bullet -}$ production and SOD and reducing enzymes'

activities. As these enzymes are highly expressed in mitochondria [55], they probably do not

constitute a limiting factor and the H_2O_2 levels we detected still reflect axonal mitochondria

metabolic activation. However, H_2O_2 levels drop quickly, probably because reducing enzymes

are removing it quickly. In addition, these enzymes' activities change following fast post-

translational modifications such as phosphorylation [56] or environmental changes [57].

While this may explain why H_2O_2 levels are not incremental following several stimulations,

further investigation will be needed to clarify underlying mechanisms.

Several CNS and PNS neuropathies are linked to mitochondrial dysfunctions in neurons

[18,19,20]. Our data show that the electrical activity of neurons directly regulates the

physiology and the respiration of axonal mitochondria. As axonal mitochondria are the most

abundant among neuronal mitochondria, this suggests that axonal mitochondria dysfunctions

may severely impact neuronal physiology and survival. We evaluated this hypothesis using a

couple of peripheral neuropathy mouse model, such as a model of CMT2A disease expressing

mutant MFN2[R94Q] [25]. While *in vitro* data had shown increased H_2O_2 levels [58], but no

change in ATP production [59] in neuronal mitochondria of these mice, *in vivo* we did not

measure significant differences in these parameters between resting MFN2[R94Q] mice and

control mice. A first explanation for this discrepancy is that cultured cells are living in an

environment that is significantly different from the *in vivo* conditions. This is especially true

in terms of metabolism [60]. Another explanation is that mitochondria physiology is altered,

but this can only be seen when mitochondria are challenged. Environmental changes in

culture may constitute this challenge, while more physiologically relevant challenges have to

be found *in vivo*. Our electrical stimulations were physiologically meaningful challenges for

axons and, similar to control mice, we observed ATP levels increasing in several axons of

mutant mice, but not all, after a first nerve stimulation. However, while control mice

mitochondria increased their ATP production further following a second stimulation, no

increase was observed in mutant mice mitochondria. We have shown that Mitochondria

Associated Membranes (MAM), which mediate interactions between the endoplasmic

reticulum and mitochondria [41], are impaired in these mutant mice. So we suggest that an

impaired Ca^{2+} uptake by mitochondria through MAM [61] could alter the ATP production as

it has been shown that calcium is required for the TCA cycle [62]. However, mutant mice

mitochondria were still able to produce a large amount of H_2O_2 following both stimulations,

suggesting the ETC and therefore the TCA cycle are working properly. Yet this H_2O_2

production was completely abnormal: it was largely delayed and was sustained for over at

least 10 minutes. As discussed before the levels of H_2O_2 result from an equilibrium between

several parameters, so it is difficult to explain this abnormal production. Nevertheless, a

relationship between mitochondrial shape and ROS has been shown *in vitro* [63]. It has been

suggested that smaller mitochondria result in intra-mitochondrial redistribution of cytochrome

c and the pro-aging molecule p66shc that acts trough ROS upregulation [64,65]. Smaller

fragmented mitochondria, such as MFN2 mutants [63], correlate with a malfunctioning

antioxidant system and an increased production of ROS [58,64]. In any case, our data show

that an altered MFN2 function lead to the dysfunction of axonal mitochondria only when

axons are firing. This dysfunction is strongly deleterious as mitochondria don't produce more

ATP but show a sustained production of deleterious ROS. This functional decoupling between

ATP and H_2O_2 production, occurring in the most abundant mitochondria of the neurons and in

conditions of normal firing activity, is likely to be a significant cause of the neuronal

dysfunction in this peripheral neuropathy.

Looking at the node of Ranvier of MFN2 mutant mice, we found that, similar to control mice,

nodal mitochondria were more metabolically active than internodal ones. As MFN2 mutant

mitochondria do not increase ATP production in active axons, this suggests that nodal

mitochondria are intrinsically more active even in absence of firing. This is consistent with

the fact that motile mitochondria slow down in the nodes of Ranvier [6] and that they have a

different morphology [66]. Moreover, nodal mitochondria may participate to axonal

degeneration since this process starts in nodes of Ranvier due to high ROS levels [67]. It

would have been interesting to investigate changes occurring in ATP and H_2O_2 production of

nodal mitochondria after electrical stimulation but this was impossible due to the small size of

the node and the change of focus that occurs following stimulations.

The myelin sheath that covers axons appears to be an important regulator of axonal

metabolism and of mitochondria physiology. Indeed in MS, demyelination alters axonal

mitochondria [19,29,68] and we show here that nodes of Ranvier devoid of myelin house

more metabolically active mitochondria. To go further, we therefore investigated the impact

of the loss of myelin on axonal mitochondria. LPC was the right tool because it induced

demyelination without destroying axons [46,47,48]. LPC-induced demyelination resulted in a

decrease of ATP levels in axonal mitochondria. This decrease was strongest after two weeks

and was reversed by 3 weeks, showing that ATP production within neurons is impaired until

complete remyelination. This decrease was unexpected, since AP propagation in

demyelinated axons is more energy demanding: the nodal machinery and the Na^+/K^+ ATPase

are diffused all along the axolemma and all the energy-effective benefits of myelination are

lost [69]. In demyelinated area of brains affected by MS, an upregulation of ETC complexes

protein expression and more complexes I and IV activity [68] was observed. Moreover,

demyelinated axons showed more mitochondria [29], which is consistent with a higher energy

demand.

Nevertheless, H_2O_2 levels strongly increased in axonal mitochondria of fully demyelinated

axons in absence of general oxidative stress in the tissue. This decoupling of mitochondrial

oxidative phosphorylation and ATP production is in accordance with previous studies that

reported a decreased ATP production, together with increased ROS levels in demyelinating

diseases [70]. This suggests that actually, mitochondrial ETC complexes are more active and

produce more ROS, but this does not lead to more ATP production. Such a mechanism has

been shown in particular conditions where the goal of mitochondrial activity is to produce

heat. Uncoupling proteins such as UCP2 permeabilize the inner mitochondrial membrane to

dissipate the proton gradient [71]. This may not be the sole utility of uncoupling as this

process has been shown to occur in several neurodegenerative diseases [72]. However, so far

we failed to detect unusual UCP2 expression in demyelinated nerves. The high production of

H_2O_2 during demyelination is probably not without consequences as ROS can alter the

function of the ATP synthase [73]. This may explain why ATP production remains weak in

mitochondria of axons that are being remyelinated two weeks after LPC injection, despite that

ROS production is back to normal. This suggests that during demyelination, axons may

recruit more mitochondria [29], or use another metabolism, such as aerobic glycolysis, in

order to cover their need in ATP.

While very little amount of data exists on the homeostasis of axonal mitochondria in

demyelinated axons of the PNS, in the CNS, demyelination in progressive MS results in

axonal mitochondria dysfunction and ROS increase [74]. The data we collected in PNS

myelinated axons on ROS production by axonal mitochondria are consistent with this. This

suggests that, despite the large difference that exists between PNS and CNS myelinating glia,

their similar role in segregating axonal firing machinery at nodes of Ranvier is essential for

mitochondria homeostasis in axons. Nevertheless, the electrical isolation of the axon and the

formation of the node of Ranvier is not the only function of the myelin sheath as it also

largely participates to the metabolic homeostasis of the axon through the lactate shuttle

process [75,76]. This is particularly true for the CNS myelin [76]. In addition, MS disease is

significantly different from peripheral nerve diseases, in particular because of the important

role played by CNS glial cells astrocytes and microglia. So, it would be essential to confirm

the data we obtained in PNS myelinated axons in CNS myelinated axons *in vivo*, in order to

definitely conclude on the alteration of ATP and ROS production by axonal mitochondria in

demyelinated lesion of progressive MS.

To conclude, combining several novel techniques such as two-photon and CARS non-linear

live imaging coupled to electrical nerve stimulation and genetically-encoded fluorescent

probes delivered by AAV vectors, we were able to observe the production of ATP and H_2O_2

by axonal mitochondria in real time and in living and active axons of mouse peripheral

nerves. We demonstrated that mitochondria are more metabolically active at the node of

Ranvier and we characterized the role of AP firing in the dynamics of mitochondrial ATP and

H_2O_2 production. Moreover we show the deleterious alterations that occur in axonal

mitochondria physiology of two mouse models of neuropathy, peripheral axonal CMT2A and

demyelinating peripheral neuropathies. Together, our data thus provide new insight into the

role of axonal mitochondria under physiological and pathological conditions.

4. Materials and methods

4.1. Cloning

pLPCX mito-roGFP-Orp1 (Addgene; #64992) was digested with XhoI/ClaI, blunted and

cloned into pAAV-MCS (Cell Biolabs, Inc.) under the control of a CMV promoter. Clones

were validated by sequencing. Likewise, pcDNA-mito-AT1.03 (from H. Imamura, Tokyo,

Japan) was digested with XhoI/HindIII, blunted and cloned into the CMV promoter controlled

pAAV-MCS. Mitochondria-targeting tags were CoxVIII for roGFP-Orp1 and two tandem

copies of CoxVIII for mito-AT1.03. AAV viral particles were produced at the viral vector

production centre at the Centre de Biotecnologia Animal I Teràpia Gènica in Barcelona, Spain

or the University of Nantes, France.

4.2. Mouse strains

Either Swiss mice (Janvier, France) or MFN2^{R94Q} mice [25] were used in the performed

experiments. Mice were kept in the animal facility of the Institute for Neurosciences of

Montpellier in clear plastic boxes and subjected to standard light cycles. All animal

experiments were conducted in accordance with the French Institutional and National

Regulation CEEA-LR-11032.

4.3. In vivo virus injection in spinal cord

A thin borosilicate glass capillary (Harvard Apparatus, Ref. 30-0016) was pulled with a

Vertical Micropipette Puller (Sutter Instruments, P30-682) to form a glass needle. The glass

needle was filled with viral solution (1 μl) and a 1 day old mouse is restrained in a position

that exposes the lower back. The needle is injected through the skin using a micromanipulator

and introduced into the spinal cord. The viral solution was injected over 2 minutes with short

pressure pulses using a Picopump (World Precision Instrument) coupled to a pulse generator.

After injection, the injection site is cleaned with betadine solution (Vetoquinol, cat. no.

3042413) for disinfection and the pup is placed back with its mother and littermates.

4.4. Immunohistochemistry

One month after the viral injection, the sciatic nerve was dissected and fixed in Zamboni's

fixative [77] for 10 min at room temperature. After fixation, the dissected sciatic nerves are

washed in PBS and incubated in successive glycerol baths (15, 45, 60, 66% in PBS) for 18 to

24 h each before freezing at –20 °C. The nerves were cut in small pieces in 66% glycerol and

the epineurium sheaths removed. Small bundles of fibers were teased in double-distilled water

on Superfrost slides and dried for 3 hours at room temperature. Some nerves were frozen in

O.C.T. Compound (Tissue-Tek, Ref. 4583) and longitudinal sections were cut using a cryostat

(Leica Biosystems, CM3050). For immunostaining, the teased fibers or longitudinal sections

were incubated for 1 h at room temperature in blocking solution (10% goat serum and 0.3%

TritonX100 in PBS). Then, the samples were incubated with TOM20 (FL-145) primary rabbit

antibody (1/500, Santa Cruz, Ref. sc-11415), mouse anti-Myelin Basic Protein (1/500, Merck,

Ref. NE1019) or rabbit NF-200 (1/500, Sigma, Ref. N4142) in blocking solution overnight at

4 °C. The next day, the samples were washed in PBS and incubated for 1h at room

temperature with secondary donkey antibodies coupled to Alexa568 (1/1000, Invitrogen, Ref.

A10042) or Alexa488 (1/1000, Invitrogen, Ref. A21202). Finally the samples were washed in

PBS and mounted in Immu-mount (Thermo Scientific). Images were acquired at room

temperature using a 20× or 40× objective, a Zeiss confocal microscope LSM710, and its

associated software.

4.5. Imaging of sciatic nerve in living mice

One month after the viral injection, the mice were anesthetized with a constant flow (1.5

l/min) of oxygen + 5% of isoflurane in an anesthesia box (World Precision Instruments, Ref.

EZ-B800) for 5 min. Thereafter the anesthesia was maintained with a mask delivering 2%

isoflurane at 0.8 l/min. The eyes were protected by eye protection gel (Ocry-gel, TVM, cat.

no. 48026T613/3). Intraperitoneal injection of 0.1 mg/kg buprenorphine was used for pre-

surgery analgesia. The mouse was placed in a silicone mold, lying on its belly, shaved on his

hind paw and the paws immobilized using small pins. The incision area was disinfected with

betadine solution (Vetoquinol, cat. no. 3042413). The skin was cut using scissors, fat tissue

removed, the gluteus superficialis and biceps femoris muscles were separated to reveal a

cavity crossed by the sciatic nerve, and the sciatic nerve was lifted up using a small spatula. A

long plastic strip was placed underneath the sciatic nerve and this strip fixed using magnets.

The sciatic nerve was kept in an aqueous environment of either sterile PBS buffer or artificial

cerebrospinal fluid (148 mM NaCl, 3 mM KCl, 1.4 mM $CaCl_2 \cdot 2H_2O$, 0.8 mM $MgCl_2 \cdot 6H_2O$,

0.2 mM $NaPO_4 \cdot H_2O$ in sterile H_2O) to prevent drying. At that point the mouse was placed

under the two-photon microscope objective lens, a glass, 12mm diameter, 0.25mm thick

microscope coverslip put on top of the nerve and a drop of deionized water placed on it to

immerse the objective lens (20X, Carl Zeiss Microscopy, LD CApochromat, Ref. 421887–

9970).

4.6. Demyelination procedure

After imaging of the sciatic nerve in vivo for at least 30 minutes, 5 µl of 1 mg/ml

lysophotphatidyl choline (LPC) was injected directly into the sciatic nerve using a Hamilton

syringe (Ref. 80930) to induce demyelination. Injection of 5 µl PBS is used as a negative

control. After injection, the nerve was placed back to its original location in the body. The

skin of the incision was realigned together using a blunt scalpel and stapled along the wound

with two clips (Fine Science Tools; 12020-00). The area around the wound was disinfected

again with betadine solution. The anesthesia mask was removed and the mouse was

monitored until it had woken up. Motor function of the injected and non-injected hindlimb

was followed by lifting the animal by the tail. At 1 week after LPC injection, the non-injected

hindlimb showed a normal postural reflex characterized by spreading of leg whereas the

injected hindlimb showed abnormal reflexes such as tremors, clasping or retracting of the

paw. This behavior has been described and linked to deficient myelination in the PNS in

previous studies [78]. The severity of abnormal hindlimb function decreased at 2 weeks after

LPC injection and no difference in leg function between the injected and non-injected

hindlimb could be observed anymore at 3 weeks after LPC injection.

4.7. Saphenous nerve stimulation in living mice

After skin incision and removal of connecting tissue, the saphenous nerve is lifted up and

isolated using a plastic strip (Fig 2a). To stimulate the saphenous nerve, two platinum kapton

microelectrodes (World Precision Instruments, PTM23B05KT) were inserted at the posterior

side of the plastic strip using a micromanipulator (Fig 2A). One hook-shaped recording

electrode (AD Instruments, MLA 1203) was placed at the anterior side (Fig 2A). The ground

electrode was inserted in the tail of the mouse and the negative electrode in the groin area (Fig

2B). The mouse was placed under the two-photon microscope (Fig 2C), a microscope glass

coverslip was placed on top of the nerve and the electrodes were connected to a Powerlab 26T

(AD Instruments; ML4856). A drop of deionized water was then placed on top of the

microscope glass to immerse the 20× objective lens.

4.8. Two-photon image acquisition

All in vivo images were obtained with a two-photon microscope LSM 7 MP OPO (Zeiss,

France) coupled to a dark microscope incubator (L S1 Dark, Zeiss) in which the temperature

was maintained at 37 °C (Heating Unit XL S, Zeiss, France). Mitochondria images were

acquired by time-lapse recording varying from one image every minute to one image every 5

minutes during 1 hour. Each image is a stack at Maximum Intensity (ZEN software, Zeiss) of

10 scans over 40 μm depth. For AT1.03 imaging, a single track at 850 nm excitation

wavelength is used to obtain both the CFP (em. 475 nm) and YFP (em. 527 nm) image at the

same time point. For roGFP imaging, the two images were acquired for each time point using

alternating tracks at 940nm and 800nm. Change of track was set after each stack. For

Coherent Anti-Stokes Raman Scattering (CARS) imaging, 2 synchronized laser lines at

excitation wavelengths 836 nm and 1097 nm are used simultaneously thanks to the OPO

system. Each scan was acquired with constant laser intensity (20% for 940 nm, 10% for 850

nm, 10% for 800nm, 15% for 836nm, 4% 1097nm) at a 512×512 pixel resolution and

microscope imaging parameters were maintained over all different regions we imaged.

Images, acquired with ZEN software (Zeiss), were saved in .czi format.

4.9. Data and statistical analysis

We used ImageJ software to analyze the relative ATP or H_2O_2 levels in mitochondria of

peripheral axons. The acquired images for each wavelength were aligned using the Template

Matching plugin. We defined a Region of Interest (ROI) encompassing all labeled

mitochondria of the same axon and the mean fluorescent intensity on the ROI was measured

on both images. These light intensities were then corrected for background light intensity

determined as an area within the nerve where no fluorescent signal from the viral probe can

be observed. Either the AT1.03 YFP/CFP ratio or the oxidized roGFP/reduced roGFP ratio

was then calculated from the 2 values for mean light intensity.

Statistical significance for the effect of time on probe stability was determined using linear

regression. The effects of the positive controls and negative controls on probe validation were

determined using Student two-tailed T-tests. The effect of nerve stimulation was determined

using one-way ANOVA and Dunnett's post-hoc tests. In addition, the threshold for a

responding axon was set at 20% change in probe fluorescence ratio. The differences between

MFN2 mutants and wild-type mice were determined using either Student two-tailed T-tests or

two- way ANOVA with Sidak post-hoc tests. Differences between internodes and node of

Ranvier mitochondria were determined using paired two-tailed T-tests. The effect of

demyelination was determined using paired two-tailed T-tests.

4.10. Data and statistical analysis

The data that support the findings of this study are available from the corresponding author

upon reasonable request.

5. Financial disclosure

This work has benefited from support by an ERC consolidator grant to NT and by the Labex

EpiGenMed ANR-10-LABX-12-01.

6. Acknowledgements

In addition we thank the MRI imaging platform, supported by the French National Research

Agency (ANR-10-INBS-04), in particular Hassan Boukhaddaoui, and the Animal Facility of

the INM. We also would like to thank Patrice Quintana for his help with the nerve stimulation

protocol and Volker Baeker for the image analysis protocol.

7. Competing interests

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

8. References
627

1.   Harris JJ, Jolivet R, Attwell D. Synaptic Energy Use and Supply. Neuron. 2012 Sep 6;75(5):762–
77.

2.   Attwell D, Laughlin SB. An Energy Budget for Signaling in the Grey Matter of the Brain. J Cereb
Blood Flow Metab. 2001 Oct 1;21(10):1133–45.

3.   Ames A. CNS energy metabolism as related to function. Brain Res Rev. 2000 Nov 1;34(1):42–68.

4.   Salzer JL, Zalc B. Myelination. Curr Biol. 2016 Oct 24;26(20):R971–5.

5.   Fischer TD, Dash PK, Liu J, Waxham MN. Morphology of mitochondria in spatially restricted
axons revealed by cryo-electron tomography. PLOS Biol. 2018 Sep 17;16(9):e2006169.

6.   Misgeld T, Kerschensteiner M, Bareyre FM, Burgess RW, Lichtman JW. Imaging axonal transport
of mitochondria in vivo. Nat Methods. 2007 Jun 10;4:559.

7.   Zhou B, Yu P, Lin M-Y, Sun T, Chen Y, Sheng Z-H. Facilitation of axon regeneration by enhancing
mitochondrial transport and rescuing energy deficits. J Cell Biol. 2016;214(1):103–119.

8.   Ohno N, Kidd GJ, Mahad D, Kiryu-Seo S, Avishai A, Komuro H, et al. Myelination and axonal
electrical activity modulate the distribution and motility of mitochondria at CNS nodes of
Ranvier. J Neurosci Off J Soc Neurosci. 2011 May 18;31(20):7249–58.

9.   Perkins GA, Ellisman MH. Mitochondrial Configurations in Peripheral Nerve Suggest Differential
ATP Production. J Struct Biol. 2011 Jan;173(1):117–27.

10.   Tarasov AI, Griffiths EJ, Rutter GA. Regulation of ATP production by mitochondrial Ca(2+). Cell
Calcium. 2012 Jul;52(1):28–35.

11.   Edgar JM, McCulloch MC, Thomson CE, Griffiths IR. Distribution of mitochondria along small-
diameter myelinated central nervous system axons. J Neurosci Res. 2008;86(10):2250–7.

12.   Giorgio M, Trinei M, Migliaccio E, Pelicci PG. Hydrogen peroxide: a metabolic by-product or a
common mediator of ageing signals? Nat Rev Mol Cell Biol. 2007 Sep 1;8:722.

13.   Murphy MP. How mitochondria produce reactive oxygen species. Biochem J. 2009 Jan 1;417(Pt
1):1–13.

14.   Hamanaka RB, Chandel NS. Mitochondrial reactive oxygen species regulate cellular signaling
and dictate biological outcomes. Trends Biochem Sci. 2010 Sep;35(9):505–13.

15.   Tormos KV, Chandel NS. Seeing the Light: Probing ROS In Vivo Using Redox GFP. Cell Metab.
2011 Dec 7;14(6):720–1.

16.   Dixon BJ, Tang J, Zhang JH. The evolution of molecular hydrogen: a noteworthy potential
therapy with clinical significance. Med Gas Res. 2013;3:10–10.

17.   Tomanek L. Proteomic responses to environmentally induced oxidative stress. J Exp Biol.
2015;218(12):1867–1879.

18.   Dauer W, Przedborski S. Parkinson's disease: mechanisms and models. Neuron. 39th ed.
2003;889–909.

19.   Su KG, Banker G, Bourdette D, Forte M. Axonal degeneration in multiple sclerosis: The
mitochondrial hypothesis. Curr Neurol Neurosci Rep. 2009 Sep;9(5):411–7.

20.   Palau F, Estela A, Pla-Martín D, Sánchez-Piris M. The Role of Mitochondrial Network Dynamics
in the Pathogenesis of Charcot-Marie-Tooth Disease. In: Espinós C, Felipo V, Palau F, editors.
Inherited Neuromuscular Diseases: Translation from Pathomechanisms to Therapies [Internet].
Dordrecht: Springer Netherlands; 2009. p. 129–37.

21.   Smith GM, Gallo G. The role of mitochondria in axon development and regeneration. Dev
Neurobiol. 2017 Oct 14;78(3):221–37.

22.   Chen H, Chan DC. Emerging functions of mammalian mitochondrial fusion and fission. Hum Mol
Genet. 2005 Oct 15;14(suppl_2):R283–9.

23.   Misko A, Jiang S, Wegorzewska I, Milbrandt J, Baloh RH. Mitofusin 2 is necessary for transport
of axonal mitochondria and interacts with the Miro/Milton complex. J Neurosci Off J Soc
Neurosci. 2010 Mar 24;30(12):4232–40.

24.   Züchner S, Mersiyanova IV, Muglia M, Bissar-Tadmouri N, Rochelle J, Dadali EL, et al. Mutations
in the mitochondrial GTPase mitofusin 2 cause Charcot-Marie-Tooth neuropathy type 2A. Nat
Genet. 2004 Apr 4;36:449.

25.   Cartoni R, Arnaud E, Médard J-J, Poirot O, Courvoisier DS, Chrast R, et al. Expression of
mitofusin 2R94Q in a transgenic mouse leads to Charcot–Marie–Tooth neuropathy type 2A.
Brain. 2010 May 1;133(5):1460–9.

26.   Pich S, Bach D, Briones P, Liesa M, Camps M, Testar X, et al. The Charcot–Marie–Tooth type 2A
gene product, Mfn2, up-regulates fuel oxidation through expression of OXPHOS system. Hum
Mol Genet. 2005 Jun 1;14(11):1405–15.

27.   Frohman EM, Racke MK, Raine CS. Multiple Sclerosis — The Plaque and Its Pathogenesis. N Engl
J Med. 2006 Mar 2;354(9):942–55.

28.   Sedel F, Bernard D, Mock DM, Tourbah A. Targeting demyelination and virtual hypoxia with
688   high-dose biotin as a treatment for progressive multiple sclerosis. Oligodendrocytes Health Dis.
689   2016 Nov 1;110:644–53.

690   29.   Mahad DJ, Ziabreva I, Campbell G, Lax N, White K, Hanson PS, et al. Mitochondrial changes
691   within axons in multiple sclerosis. Brain J Neurol. 2009 May;132(Pt 5):1161–74.

692   30.   Campbell G, Mahad DJ. Mitochondrial dysfunction and axon degeneration in progressive
693   multiple sclerosis. FEBS Lett. 2018 Feb 17;592(7):1113–21.

694   31.   Gutscher M, Sobotta MC, Wabnitz GH, Ballikaya S, Meyer AJ, Samstag Y, et al. Proximity-based
695   Protein Thiol Oxidation by H(2)O(2)-scavenging Peroxidases. J Biol Chem. 2009 Nov
696   13;284(46):31532–40.

697   32.   Imamura H, Huynh Nhat KP, Togawa H, Saito K, Iino R, Kato-Yamada Y, et al. Visualization of
698   ATP levels inside single living cells with fluorescence resonance energy transfer-based
699   genetically encoded indicators. Proc Natl Acad Sci U S A. 2009 Sep 15;106(37):15651–6.

700   33.   Albrecht SC, Barata AG, Großhans J, Teleman AA, Dick TP. In Vivo Mapping of Hydrogen
701   Peroxide and Oxidized Glutathione Reveals Chemical and Regional Specificity of Redox
702   Homeostasis. Cell Metab. 2011 Dec 7;14(6):819–29.

703   34.   Ren W, Ai H-W. Genetically Encoded Fluorescent Redox Probes. Sensors. 13th ed. 2013;15422–
704   33.

705   35.   Tu H, Boppart SA. Coherent anti-Stokes Raman scattering microscopy: overcoming technical
706   barriers for clinical translation. J Biophotonics. 2014 Jan;7(0):9–22.

707   36.   Mytskaniuk V, Bardin F, Boukhaddaoui H, Rigneault H, Tricaud N. Implementation of a Coherent
708   Anti-Stokes Raman Scattering (CARS) System on a Ti:Sapphire and OPO Laser Based Standard
709   Laser Scanning Microscope. J Vis Exp. 113th ed. 2016;doi: 10.3791/54262.

710   37.   Hajjar H, Boukhaddaoui H, Rizgui A, Sar C, Berthelot J, Perrin-Tricaud C, et al. Label-free non-
711   linear microscopy to measure myelin outcome in a rodent model of Charcot-Marie-Tooth
712   diseases. J Biophotonics. 2018 Aug 9;0(ja):e201800186.

713   38.   Huff TB, Cheng J-X. In vivo coherent anti-Stokes Raman scattering imaging of sciatic nerve
714   tissue. J Microsc. 2007 Feb;225(Pt 2):175–82.

715   39.   Gonzalez S, Fernando R, Berthelot J, Perrin-Tricaud C, Sarzi E, Chrast R, et al. In vivo time-lapse
716   imaging of mitochondria in healthy and diseased peripheral myelin sheath. Mitochondrion.
717   2015 Jul 1;23:32–41.

718   40.   Ino D, Sagara H, Suzuki J, Kanemaru K, Okubo Y, Iino M. Neuronal Regulation of Schwann Cell
719   Mitochondrial Ca2+ Signaling during Myelination. Cell Rep. 2015 Sep 29;12(12):1951–9.

720   41.   Bernard-Marissal N, Van Hameren G, Juneja M, Pellegrino C, Rochat C, El Mansour O, et al.
721   Endoplasmic reticulum and mitochondria dysfunction underlies reversible Charcot–Marie–
722   Tooth type 2A neuropathy. (Under Review). under review;

723   42.   Bazil JN, Beard DA, Vinnakota KC. Catalytic Coupling of Oxidative Phosphorylation, ATP
724   Demand, and Reactive Oxygen Species Generation. Biophys J. 2016 Feb 23;110(4):962–71.

725   43.   Cadenas E, Davies KJA. Mitochondrial free radical generation, oxidative stress, and aging. Free
726   Radic Biol Med. 2000 Aug 1;29(3):222–30.

727   44.   Bélanger E, Henry FP, Vallée R, Randolph MA, Kochevar IE, Winograd JM, et al. In vivo
728   evaluation of demyelination and remyelination in a nerve crush injury model. Biomed Opt
729   Express. 2011 Sep;2(9):2698–2708.

730   45.   Cooper MF, Webster GR. The differentiation of phospholipase A1 and A2 in rat and human
731   nervous tissues. J Neurochem. 1970 Nov;17(11):1543–54.

732   46.   Allt G, Ghabriel MN, Sikri K. Lysophosphatidyl choline-induced demyelination. Acta Neuropathol
733   (Berl). 1988 Sep 1;75(5):456–64.

734   47.   Plemel JR, Michaels NJ, Weishaupt N, Caprariello AV, Keough MB, Rogers JA, et al. Mechanisms
735   of lysophosphatidylcholine-induced demyelination: A primary lipid disrupting myelinopathy.
736   Glia. 2017 Oct 25;66(2):327–47.

737   48.   Kiryu-Seo S, Ohno N, Kidd GJ, Komuro H, Trapp BD. Demyelination increases axonal stationary
738   mitochondrial size and the speed of axonal mitochondrial transport. J Neurosci Off J Soc
739   Neurosci. 2010 May 12;30(19):6658–66.

740   49.   Tricaud N, Park H. Wallerian demyelination: chronicle of a cellular cataclysm. Cell Mol Life Sci.
741   74th ed. 2017;4049–57.

742   50.   Meiri H, Steinberg R, Medalion B. Detection of Sodium Channel Distribution in Rat Sciatic Nerve
743   Following Lysophosphatidylcholine-Induced Demyelination. J Membrane Biol. 92nd ed.
744   1986;47–56.

745   51.   Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY, et al. Investigating
746   Mitochondrial Redox Potential with Redox-sensitive Green Fluorescent Protein Indicators. J Biol
747   Chem. 2004 Mar 26;279(13):13044–53.

748   52.   Dooley CT, Dore TM, Hanson GT, Jackson WC, Remington SJ, Tsien RY. Imaging Dynamic Redox
749   Changes in Mammalian Cells with Green Fluorescent Protein Indicators. J Biol Chem. 2004 May
750   21;279(21):22284–93.

751   53.   Samara C, Poirot O, Domènech-Estévez E, Chrast R. Neuronal activity in the hub of
752   extrasynaptic Schwann cell-axon interactions. Front Cell Neurosci. 2013;7:228.

753   54.   Yi M, Weaver D, Hajnóczky G. Control of mitochondrial motility and distribution by the calcium
754   signal: a homeostatic circuit. J Cell Biol. 2004 Nov 22;167(4):661–72.

755   55.   Fukai T, Ushio-Fukai M. Superoxide Dismutases: Role in Redox Signaling, Vascular Function, and
756   Diseases. Antioxid Redox Signal. 2011 Sep 15;15(6):1583–606.

757   56.   Yamakura F, Kawasaki H. Post-translational modifications of superoxide dismutase. Biochim
758   Biophys Acta. 1804th ed. 2010;318–25.

759   57.   Little C, Olinescu R, Reid KG, O'Brien PJ. Properties and Regulation of Glutathione Peroxidase. J
760   Biol Chem. 1970 Jul 25;245(14):3632–6.

58.   Nie Q, Wang C, Song G, Ma H, Kong D, Zhang X, et al. Mitofusin 2 deficiency leads to oxidative
stress that contributes to insulin resistance in rat skeletal muscle cells. Mol Biol Rep. 2014 Oct
1;41(10):6975–83.

59.   Baloh RH, Schmidt RE, Pestronk A, Milbrandt J. Altered Axonal Mitochondrial Transport in the
Pathogenesis of Charcot-Marie-Tooth Disease from Mitofusin 2 Mutations. J Neurosci.
2007;27(2):422–430.

60.   Zilberter Y, Zilberter T, Bregestovski P. Neuronal activity in vitro and the in vivo reality: the role
of energy homeostasis. Trends Pharmacol Sci. 2010 Sep 1;31(9):394–401.

61.   Ainbinder A, Boncompagni S, Protasi F, Dirksen RT. Role of Mitofusin-2 in Mitochondrial
Localization and Calcium Uptake in Skeletal Muscle. Cell Calcium. 2015 Jan;57(1):14–24.

62.   Chen Y, Csordás G, Jowdy C, Schneider TG, Csordás N, Wang W, et al. Mitofusin 2-containing
Mitochondrial-Reticular Microdomains Direct Rapid Cardiomyocyte Bioenergetic Responses via
Inter-Organelle Ca(2+) Crosstalk. Circ Res. 2012 Sep 14;111(7):863–75.

63.   Yu T, Robotham JL, Yoon Y. Increased production of reactive oxygen species in hyperglycemic
conditions requires dynamic change of mitochondrial morphology. Proc Natl Acad Sci U S A.
2006 Feb 21;103(8):2653–8.

64.   de Brito OM, Scorrano L. Mitofusin 2: A Mitochondria-Shaping Protein with Signaling Roles
Beyond Fusion. Antioxid Redox Signal. 2007 Dec 20;10(3):621–34.

65.   Giorgio M, Migliaccio E, Orsini F, Paolucci D, Moroni M, Contursi C, et al. Electron Transfer
between Cytochrome c and p66Shc Generates Reactive Oxygen Species that Trigger
Mitochondrial Apoptosis. Cell. 2005 Jul 29;122(2):221–33.

66.   Chen H, Chan DC. Mitochondrial dynamics–fusion, fission, movement, and mitophagy–in
neurodegenerative diseases. Hum Mol Genet. 2009 Oct 15;18(R2):R169–76.

67.   Bros H, Millward JM, Paul F, Niesner R, Infante-Duarte C. Oxidative damage to mitochondria at
the nodes of Ranvier precedes axon degeneration in ex vivo transected axons. Exp Neurol. 2014
Nov 1;261:127–35.

68.   Witte ME, Bø L, Rodenburg RJ, Belien JA, Musters R, Hazes T, et al. Enhanced number and
activity of mitochondria in multiple sclerosis lesions. J Pathol. 2009 May 26;219(2):193–204.

69.   Young EA, Fowler CD, Kidd GJ, Chang A, Rudick R, Fisher E, et al. Imaging correlates of
decreased axonal Na+/K+ ATPase in chronic multiple sclerosis lesions. Ann Neurol.
2008;63(4):428–35.

70.   Gilgun-Sherki Y, Melamed E, Offen D. The role of oxidative stress in thepathogenesis of multiple
sclerosis: The need for effectiveantioxidant therapy. J Neurol. 2004 Mar 1;251(3):261–8.

71.   Rousset S, Alves-Guerra M-C, Mozo J, Miroux B, Cassard-Doulcier A-M, Bouillaud F, et al. The
Biology of Mitochondrial Uncoupling Proteins. Diabetes. 2004;53(suppl 1):S130–S135.

72.   Dalgaard LT, Pedersen O. Uncoupling proteins: functional characteristics and role in the
pathogenesis of obesity and Type II diabetes. Diabetologia. 2001 Aug 1;44(8):946–65.

73.   Rexroth S, Poetsch A, Rögner M, Hamann A, Werner A, Osiewacz HD, et al. Reactive oxygen
799   species target specific tryptophan site in the mitochondrial ATP synthase. Biochim Biophys Acta
800   BBA - Bioenerg. 2012 Feb 1;1817(2):381–7.

801   74.   Yin X, Kidd GJ, Ohno N, Perkins GA, Ellisman MH, Bastian C, et al. Proteolipid protein–deficient
802   myelin promotes axonal mitochondrial dysfunction via altered metabolic coupling. J Cell Biol.
803   2016 Nov 21;215(4):531–42.

804   75.   Yellen G. Fueling thought: Management of glycolysis and oxidative phosphorylation in neuronal
805   metabolism. J Cell Biol. 2018;217(7):2235–2246.

806   76.   Funfschilling U, Supplie LM, Mahad D, Boretius S, Saab AS, Edgar J, et al. Glycolytic
807   oligodendrocytes maintain myelin and long-term axonal integrity. Nature. 2012 Apr
808   29;485(7399):517–21.

809   77.   Stefanini M, Martino CD, Zamboni L. Fixation of Ejaculated Spermatozoa for Electron
810   Microscopy. Nature. 1967 Oct 14;216:173.

811   78.   Hung H, Kohnken R, Svaren J. The NURD chromatin remodeling complex is required for
812   peripheral nerve myelination. J Neurosci. 2012 Feb 1;32(5):1517–27.