1   Effects of genome size on pollen performance

5   How does genome size affect the evolution of pollen tube growth rate, a haploid

6   performance trait?

11   John B. Reese[1,2] and Joseph H. Williams[2]

12   Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN

13   37996, U.S.A.

17   [1]Author for correspondence:

18   John B. Reese

19   Tel: 865 974 9371

20   Email: jreese11@vols.utk.edu

ABSTRACT

Premise of the Study – Male gametophytes of most seed plants deliver sperm to eggs via a

pollen tube. Pollen tube growth rates (*PTGRs*) of angiosperms are exceptionally rapid, a pattern

attributed to more effective haploid selection under stronger pollen competition. Paradoxically,

whole genome duplication (WGD) has been common in angiosperms but rare in gymnosperms.

Pollen tube polyploidy should initially accelerate *PTGR* because increased heterozygosity and

gene dosage should increase metabolic rates, however polyploidy should also independently

increase tube cell size, causing more work which should decelerate growth. We asked how

genome size changes have affected the evolution of seed plant *PTGRs*.

Methods - We assembled a phylogenetic tree of 451 species with known *PTGRs*. We then used

comparative phylogenetic methods to detect effects of neo-polyploidy (within-genus origins),

DNA content, and WGD history on *PTGR*, and correlated evolution of *PTGR* and DNA content.

Key Results - Gymnosperms had significantly higher DNA content and slower *PTGR* optima

than angiosperms, and their *PTGR* and DNA content were negatively correlated. For

angiosperms, 89% of model weight favored Ornstein-Uhlenbeck models with a faster *PTGR*

optimum for neo-polyploids, but *PTGR* and DNA content were not correlated. In comparisons of

within-genus and intraspecific-cytotype pairs, *PTGRs* of neo-polyploids ≤ paleo-polyploids.

Conclusions – Genome size increases should negatively affect *PTGR* when genetic

consequences of WGDs are minimized, as found in intra-specific autopolyploids (low heterosis)

and gymnosperms (few WGDs). But in angiosperms, the higher *PTGR* optimum of neo-

polyploids and non-negative *PTGR*-DNA content correlation suggest that recurrent WGDs have

caused substantial *PTGR* evolution in a non-haploid state.

Keywords: DNA content, evolution of development, gametophyte, macroevolution, pollen

competition, pollen tube growth rate, polyploidy, whole genome duplication.

INTRODUCTION

In seed plants, the male gametophyte is a highly-reduced, haploid organism that develops within the pollen grain and completes its life cycle after pollination by growing a pollen tube that invades female reproductive tissues. The pollen tube functions to attach the male gametophyte and to absorb nutrients from female tissues, and in most seed plants (conifers, Gnetales, and angiosperms), it has the novel function of transporting the sperm cells to the egg-bearing female gametophyte (siphonogamy) (Friedman, 1993). Pollen tube growth rate (*PTGR*) is a central aspect of male gametophyte performance that can evolve due to changes in the time between pollination and fertilization, and due to changes in the intensity of pollen tube competition. Strikingly, angiosperms are known to have much shorter reproductive cycles (Williams and Reese, 2019), much higher potential for pollen competition (Mulcahy, 1979), and orders of magnitude faster *PTGRs* (Williams, 2012) relative to gymnosperms. The pattern of exceptionally fast angiosperm *PTGRs* is thought to have evolved rapidly via haploid selection on pollen-expressed genes (Mulcahy, 1979; Arunkumar et al., 2013; Otto et al., 2015), which constitute a large portion of the genome (Tanksley et al., 1981; Rutley and Twell, 2015; Hafidh et al., 2016).

If the dramatic and rapid acceleration of *PTGR*s in angiosperms has been driven by haploid selection on pollen performance genes, then one might expect polyploidy to be rare in angiosperms. Evolution above the haploid level is expected to reduce the efficiency of selection on pollen (Otto et al., 2015). Yet, the opposite is true – ancient whole genome duplications (WGDs), recent polyploids, and speciation by polyploidy have been especially common in angiosperms, whereas in gymnosperms genome size has evolved largely by other processes (Wood et al., 2009; Mayrose et al., 2011; Leitch & Leitch, 2012, 2013; Landis et al., 2018). In fact, large changes genome size can have a number of immediate effects on *PTGR*. First, *PTGR*

might be faster in a neo-diploid pollen tube since increases in gene number cause: 1) heterosis,

due to sheltering of deleterious pollen-expressed alleles and/or new allelic interactions upon loss

of haploidy (Lande and Schemske, 1985; Husband and Schemske, 1997; Comai, 2005; Birchler

et al. 2010; Husband, 2016), and 2) gene dosage effects, due to increased capacity for protein

synthesis and hence the possibility for higher metabolic rates (Stebbins, 1974; Comai, 2005;

Conant and Wolfe, 2008). On the other hand, substantial increases in DNA content (whether by

WGD or other processes) are known to increase nuclear size, cell size, and the duration of the

cell cycle, independent of the effects of genes (Bennett, 1971, 1972; Cavalier-Smith, 1978; Price

1988; Cavalier-Smith, 2005). The phenotypic effects of increased bulk DNA, hereafter referred

to as "nucleotypic" effects (Bennett, 1971; Snodgrass et al, 2016; Doyle & Coate, 2019), cause

more work for the growing pollen tube cell and should therefore negatively affect *PTGR*,

counteracting the positive "genotypic" effects of heterozygosity and gene dosage.

As shown in Figure 1, if genome size expansion occurs without increasing the number of

genes, then nucleotypic effects will predominate and slower *PTGRs* should evolve. But if

genome size increase occurs by WGD, then altered gene expression patterns (due to dosage and

heterozygosity effects) should counteract nucleotypic effects in the stabilized neo-polyploid (Fig.

1). In the latter case, the balance of nucleotypic and genotypic effects varies depending on the

magnitude of potential heterosis, which depends directly on the amount of standing genetic

variation (Birchler et al. 2010). In general, at inception tetraploid sporophytes are expected to

have higher heterozygosity than their diploid progenitors, irrespective of mode of

polyploidization (auto- to allo-polyploidy) or mating system (Lande and Schemske 1985; Soltis

and Soltis 2000). Thus, at inception, autopolyploids that arise from outcrossing progenitors and

allopolyploids will have a higher potential for heterosis, relative to autopolyploids that arose

from selfing ancestors (Fig. 1).

After the initial effects of WGD, genotypic effects continue to evolve under both

stabilizing and directional selection on *PTGR*, mediated by shifts in mating system and

phenomena such as genome downsizing, biased gene retentions, recombination, and ultimately

the return to disomic inheritance (Conant and Wolfe, 2008; De Smet et al. 2013; Conant et al.,

2014; Freeling et al., 2015; Dodsworth et al., 2016; Panchy et al. 2016; Wendel et al., 2018).

Nucleotypic effects by definition can only evolve by changes in genome size, which after WGD

tend to be biased to small losses relative to the size of the WGD (Dodsworth et al. 2016). Hence,

with time, genotypic effects are predicted to overwhelm nucleotypic effects, irrespective of

initial effects and the direction of *PTGR* evolution.

In this study, we used model-based comparative phylogenetic analyses to determine if

polyploidy, DNA C-value, and WGD history have affected the evolution of *PTGRs* in seed

plants, and whether genome size effects have occurred predominantly during polyploid periods

of history or during subsequent periods of more or less diploid evolution. Because all seed plants

have at least one WGD in their history, we defined neo-polyploids as having a higher

chromosome multiple than the base chromosome number of their genus, and paleo-polyploids

(hereafter, "diploids") as having similar chromosome number as the genus base number (as in

Wood et al. 2009; Mayrose et al. 2011). This allowed us to determine if, 1) neo-polyploids have

faster *PTGRs* than diploids, as predicted if WGDs generally produce strong initial genotypic

effects that persist in the polyploid condition, or 2) neo-polyploids have slower *PTGRs* than

diploids, as predicted if nucleotypic effects initially outweigh genotypic effects and if fast

PTGRs generally evolve after diploidization (eg. in paleopolyploids). We also predict an

underlying negative correlation between *PTGR* and genome size due to nucleotypic effects,

which should be most apparent in intraspecific neo-polyploids and in lineages with little history

of WGD.

121   **MATERIALS AND METHODS**

Tree Construction and Dating – GenBank accessions for 16 gene regions (*rbcL, matK, trnL-F,*

18s_rDNA, atpB, ndhF, adh, trnL, rpl32, trnT-L, psbA-trnH, rpl32-trnL, ITS, 5.8s_rRNA, rps16,

and *26s_rDNA*) for 451 seed plant species with pollen tube growth rate data were retrieved,

cleaned, and assembled into a multiple gene alignment (length – 9263 base pairs, 16 partitions,

69.6% missing data) using PHLAWD and phyutility (Smith and Donoghue, 2008; Smith and

Dunn, 2008). Tree inference was performed using maximum likelihood in RAxML version 8

(Stamatakis, 2014) on CIPRES. A pruned version of the seed plant tree from Magallón et al.

(2015) was used as a guide tree to enforce topology of major clades. The resulting maximum

likelihood estimate of the tree was rooted and ultrametricized using the *ape* (Paradis et al., 2004)

and *geiger* packages in R (Harmon et al., 2008). Time-calibration was performed with the

Congruification method (Eastman et al., 2013), using the Magallón et al. (2015) phylogeny as the

reference tree.

Data collection and character scoring – Data on *PTGRs* were taken from Williams

(2012) and more recent literature (cited in Appendix S1; see the Supplementary Data with this

article). The *PTGR* value used for each species represents an estimate of maximum sustained

growth rate, which is consistent with other comparative analyses of physiological traits, and with

the fact that researchers almost always measure *PTGRs* from the longest pollen tube(s). Thus,

PTGR values for each species represent an average of maximum in vivo growth rates, or if there

was more than one report for a species the average of those values (as in Williams, 2012).

PTGRs were taken exclusively from within-ploidy level crosses (i.e., never from inter-ploid

crosses), in keeping with our overall goal of finding mechanisms underlying the pattern of *PTGR*

evolution within stabilized polyploids.

DNA content was analyzed using 1C-value: the amount of nuclear DNA in the

unreplicated gametic nucleus, irrespective of ploidy level (Swift, 1950; Bennett and Leitch,

2012). As we were primarily interested in the nucleotypic effects of bulk DNA amount, we use

the terms C-value, DNA content, and genome size interchangeably throughout. C-value data was

collected from the Kew Royal Botanic Gardens Plant C-value Database (Bennett and Leitch,

2012). Chromosome counts were obtained from the Index to Plant Chromosome Numbers

(IPCN). To examine the effect of recent polyploidy (defined as occurring at or within the genus

level; Wood et al., 2009; Mayrose et al., 2011) on *PTGR*, we scored taxa as "neo-polyploid" if

their chromosome counts were $\geq$ 1.5 times that of their generic 1x base count (from Wood et al.,

2009) and "diploid" (paleo-polyploid) if < 1.5 times that value (N = 206 angiosperms, 23

gymnosperms). To examine the effect of ancient (deeper than genus-level) duplication events on

PTGR, the number of WGDs in each genus-to-root lineage was counted for each angiosperm

(found in Appendix S1 of Landis et al. 2018) and gymnosperm (Li et al. 2015).

Phylogenetic Comparative Analyses - To visualize changes in DNA content and *PTGR*

along tree branches and to generate estimates of node states, ancestral state reconstructions were

performed and plotted using the contMap function in *phytools* (Felsenstein, 1985; Revell,

2012)(Appendix S2). Given many known biological differences between gymnosperms and

angiosperms in pollen tube growth (Friedman, 1993; Williams, 2008) and in mechanisms of

genome size change (see Discussion) (Ohri and Khoshoo, 1986; Leitch et al., 1998), all

comparative analyses were performed on gymnosperms only, angiosperms only, and the full

dataset (all spermatophytes). C-value and *PTGR* were $\log_{10}$-transformed for all analyses.

Model-based analyses were used to examine patterns of *PTGR* and C-value evolution

separately. The *OUwie* function was implemented in R (Beaulieu and O'Meara, 2014), and the

following models were tested: single- and multi-rate Brownian motion (BM1, BMS,

respectively), single-regime Ornstein-Uhlenbeck (OU1), and multi-regime OU models with

either one global α and σ^2 estimate (OUM), one α and multiple σ^2 (OUMV), or multiple α and

one σ^2 (OUMA). In all models, σ^2 represents the rate of random evolution and α, the strength of

attraction to an optimum, θ. The single- and multiple-regime models were compared to test

whether or not, 1) angiosperms and gymnosperms evolve around different *PTGR* or C-value

optima, and 2) diploids and neo-polyploids (within all three groups) evolve around different

PTGR or C-value optima. For all analyses, AICc values were used to calculate model weights

and the weighted average of parameter values was then calculated using all models that

contributed > 1% of the model weight (Burnham & Anderson, 2002). Unless otherwise noted, all

measures of uncertainty around parameter estimates are standard errors.

Since each *PTGR* value represents a species mean obtained from multiple measurements,

we attempted to incorporate error into phylogenetic comparative analyses. Since species means

were $\log_{10}$-transformed for analysis, $\log_{10}$-transformed *SE*s are also required. As there is no

reliable way to calculate the $\log_{10}$-transformed *SE* from the literature without the original data for

each species, we took the following approach. First, we assumed all species had similar *SE*s in

PTGR, and we applied an empirically-determined *SE* from an exemplar species to all. *Magnolia*

grandiflora has an average *PTGR* of 828 ± 141 μm h^{-1} ($N = 25$ outcrosses), close to the

angiosperm median of 587 μm h^{-1} (Williams, 2012 and this study) (Appendix S3). The standard

deviation (*SD*) of $\log_{10}$-transformed data was calculated and divided by the mean of the $\log_{10}$-

transformed data to acquire a coefficient of variation (*CV*) of 0.0237. We then multiplied the

$\log_{10}$-transformed mean *PTGR* of each species by 0.0237 to provide an estimate of the $\log_{10}$

taxon-specific standard deviation. The standard deviation (*SD*) was used as a conservative

estimate of error because sample sizes were generally not available for calculating *SE*. Secondly,

we performed a sensitivity analysis by evaluating each evolutionary model in *OUwie* with *SDs*

calculated from hypothetical global *CV*s of 0.00, 0.05, 0.10, 0.25, and 0.50 (Appendix S4).

The association between recent polyploidy and *PTGR* was also assessed among 10

diploid-polyploid near-relative pairs (appearing as sister taxa on the tree at the within-genus

level). Only polyploid taxa with a single diploid sister on the tree were used. The *PTGRs* of 11

intraspecific diploid-autopolyploid pairs from the literature were also compared. A two-tailed

binomial (sign) test was used to test significance in both.

The cumulative effect of ancient polyploid events was explored with phylogenetic

generalized least squares (*PGLS*) regression using the *phylolm* package in R (Ho & Ane, 2014).

The number of ancient duplication events in the history of each tip taxon (inferred from Landis et

al., 2018) was used as the predictor variable with *PTGR* as the response variable.

The relationship between pollen tube growth rate and gametophytic DNA content was

also assessed with *PGLS* regression. Gametophytic DNA content was used as the predictor

variable and *PTGR* the response variable. BM (Grafen, 1989) and OU (Martins and Hansen,

1997) models were both used, in addition to Pagel's lambda, kappa, and delta models (Pagel,

1997, 1999). To examine the effect of ploidy and the interaction between ploidy and C-value on

PTGR, a phylogenetic ANCOVA was implemented with C-value as the covariate in *phylolm*.

Shifts among convergent *PTGR* and C-value optima were determined with a maximum

likelihood approach to detect multiple optima within seed plants, using *SURFACE* in R (Ingram

and Mahler, 2013). Using an OU model with a global α and σ^2, a single-optimum model was

subdivided into multiple-optima models in a stepwise fashion until adding another optimum

decreased the model likelihood by ΔAIC > -2. Separate optima were then collapsed (i.e. two

regimes were assigned the same optimum) in a pairwise fashion until further collapses decreased

model likelihood. Shifts in *PTGR* and C-value optima that occurred at the same node, or within

two nodes of each other, were identified manually. Nodes with *PTGR* or C-value regime shifts

were also manually compared to the Landis et al. (2018) WGD map to see if a WGD had

occurred at that node or up to two nodes *prior* to the regime shift.

219   **RESULTS**

PTGR evolution and C-value evolution in angiosperms versus gymnosperms – The *PTGR* tree

comprised 451 seed plants, with 28 species from 7 of 8 gymnosperm orders (Christenhusz et al.

2011) including Cycads, *Ginkgo*, conifers and Gnetales; and 423 species from 38 of 64 (59%) of

angiosperm orders (APG IV 2016), including representatives from all three ANA grade lineages,

Chloranthaceae, eumagnoliids, and a broad distribution of both monocots and eudicots (full tree

in Appendix S2). Gymnosperm *PTGR*s ranged from < 1 to 19 μm h^{-1} (mean ± *SD* = 3.29 ± 4.34,

median = 1.49 μm h^{-1}), whereas angiosperm *PTGR*s ranged from < 5 to > 30,000 μm h^{-1} (mean ±

SD = 1744 ± 3576 μm h^{-1}, median = 587 μm h^{-1}). The maximum likelihood (ML) reconstruction

indicated that ancestral $\log_{10}$ *PTGR* of extant angiosperms was 2.44 μm h^{-1} (95% CI: 1.09-3.69)

versus 0.215 μm h^{-1} (95% CI: -1.48-1.92) for extant gymnosperms (Appendix S2).

Model-based analyses of seed plant *PTGRs* and C-values favored OU models with

separate optima for angiosperms and gymnosperms, accounting for > 99.9 % of the model

weight in both analyses (Appendix S5, S6). Log_{10} *PTGR* optima were more than an order of

magnitude faster in angiosperms (2.69 ± 0.048 μm h^{-1}) than in gymnosperms (0.187 ± 0.123 μm

h^{-1}).

The C-value tree included 183 species from the *PTGR* tree for which DNA content data

could be obtained. The resulting log_{10} C-value optimum for angiosperms (0.184 ± 0.051 pg) was

more than a magnitude of order smaller than that of gymnosperms (1.231 ± 0.041 pg). Ancestral

Log_{10} DNA content was also smaller for angiosperms than for gymnosperms, 0.29 pg (95% CI: -

0.45-1.04) versus 1.10 pg (95% CI: -0.28-2.47), consistent with larger comparative analyses of

DNA content (see Leitch and Leitch, 2013).

Joint evolution of PTGR and ploidy – In model-based analyses of angiosperms using the

empirical error rate, 89% of the model weight favored a separate and higher optimum for neo-

polyploids (*N* = 68) than for diploids (*N* = 138) (model averaged log_{10} *PTGR* = 3.2 ± 0.23 vs. 2.8

± 0.08 μm h^{-1}; Table 2). In the sensitivity analysis, OU models with separate and faster *PTGR*

optima for neo-polyploids than diploids received > 50% of model weight when the error

calculated from *CVs* ranged from 0 to 25 %, but above 25% single-regime and BM models had

the majority of the weight (Appendix S4). These are conservative results, since *SDs*, not *SEs*,

were used to model error on the tree. The gymnosperm-only analysis was not performed due to

low sample size (2 of 23 species were polyploid).

A survey of intraspecific cytotypes found autopolyploids had slower *PTGR* than diploids

in 9 of 11 pairs and no difference in the remaining two (Binomial test, *P* = 0.002; Appendix

S7b). In the nearest-relative comparisons, within-genus polyploids had slower *PTGR* than

diploids in four pairs, faster *PTGR* in five, and no difference in one (Two-tailed binomial test, *P*

= 0.623)(Appendix S7a).

The historical effect of number of ancient genome duplications on *PTGR* was non-

significant, whether or not recent (within-genus) WGDs were included (kappa model weight >

99.9%, $N = 451$; $P > 0.3$ in both analyses).

Joint evolution of PTGR, DNA content and ploidy - For seed plants, ordinary least

squares (*OLS*) regression showed a significant negative correlation between DNA content and

PTGR ($N = 183$, $P<0.0001$), but that result was clearly driven by the large DNA contents and

slow *PTGRs* of gymnosperms relative to angiosperms (Fig. 2), because the *PGLS* regression was

non-significant (Table 1). Taking these two clades separately, DNA content was negatively

correlated with *PTGR* in gymnosperms in the *PGLS* regression ($N = 23$; model-averaged slope: -

$1.09 \pm 0.49 \log_{10}$ *PTGR*; Table 1). In angiosperms, a positive correlation using *OLS* ($N = 161$;

$P=0.0005$), was non-significant using *PGLS* (Table 1). The patterns of *PTGR* and C-value

evolution in seed plants can be visualized in Figure 3. In a smaller phylogenetic ANCOVA

analysis, after controlling for C-value, the effect of ploidy on *PTGR* was non-significant in

angiosperms ($N = 100$) and seed plants ($N = 118$) (non-significant ploidy x C-value interaction

removed; Appendix S8).

Coincident regime shifts in PTGR and DNA content – Maximum likelihood analysis of

convergent evolution of *PTGRs* detected 13 distinct optima ($N = 451$ taxon tree), with 51 shifts

(22 to faster and 29 to slower optima). For C-value, there were 9 distinct optima ($N = 184$ taxon

tree), with 4 shifts to larger and 7 shifts to smaller optima. Regime shifts in both traits were

coincident at only two nodes: a *PTGR* acceleration (from $\theta = 0.147$ to $\theta = 2.47 \log_{10} \mu m \ h^{-1}$) and

genome downsizing ($\theta = 2.71$ to $\theta = 0.702 \log_{10}$ pg) in the CA of extant angiosperms; and a

PTGR slowdown ($\theta = 2.78$ to $\theta = 2.47$ $\log_{10}$ μm h^{-1}) and genome size decrease ($\theta = 0.209$ to $\theta =$

-0.386 $\log_{10}$ pg) in the CA of rosids and Saxifragales (i.e. superrosids; Fig. 4). When the search

was relaxed to include adjacent nodes, an additional coincidence occurred, with shift to higher

PTGR followed by a shift to higher C-value near the base of monocots. Ancient WGDs

coincided with the shifts in *PTGR* and C-value at the CA of angiosperms (above) and with a

decrease in C-value in the CA of eudicots.

283   **DISCUSSION**

The impact of genome size on *PTGR* is determined by the magnitudes of conflicting nucleotypic

and genotypic effects. Such effects depend on the mechanism of genome size change.

Nucleotypic effects decelerate *PTGR* and are always present irrespective of mode of genome size

change, whereas large-scale genetic effects are only possible after WGD. We predicted that

angiosperms and gymnosperms should have different patterns of *PTGR* evolution based on their

contrasting patterns of genome size change. Gymnosperm *PTGRs* should be most susceptible to

nucleotypic effects because they have evolved large genomes sizes and WGDs have been rare. In

contrast, angiosperms have evolved smaller genome sizes despite recurrent WGDs and

widespread present-day polyploidy. Thus, gene duplication and sorting have played a much

greater role in the evolution of angiosperm *PTGRs*, allowing genotypic effects to counterbalance

or overwhelm nucleotypic effects. Below we discuss our findings in light of these expected

patterns.

The evolution of* PTGR *in angiosperms versus gymnosperms - We found that seed plant

PTGRs best fit an OU model, indicating less *PTGR* variation among lineages than expected

under a Brownian motion evolutionary model, with a faster optimum for angiosperms than for

gymnosperms. Phylogenetic half-lives were similar (5.6 and 5.7 MY, respectively) and very

short (only 3.9% and 2.3 % of their respective crown ages), indicating a strong attraction to their

optimum values. Such a pattern is consistent with stabilizing selection on *PTGR* imposed by

slower evolution of linked sporophytic traits, such as the timing of stigma receptivity relative to

egg receptivity, pollen tube pathway length, or maternal provisioning. Gymnosperm *PTGRs* may

have been constrained by a hard boundary such as by biophysical or physiological limitations, or

a soft boundary, such as by lack of selection for fast rates. Angiosperms have clearly not been

bound by those same limitations, given their much higher *PTGR* optimum, the convergent

evolution of extremely fast *PTGRs* in many unrelated derived lineages of monocots and eudicots,

and occasionally large within-genus differences in *PTGR*.

Our results suggest that most of the accelerations of angiosperm *PTGR*, and their higher

PTGR variance relative to gymnosperms, have largely evolved *after* the origin of angiosperms

and their novel pollen tube cell biology. First, estimates of angiosperm ancestral *PTGR* and

ancestral optimum under OU (275 and 295 μm h^{-1}, respectively) are slower than the angiosperm-

wide OU optimum of 490 μm h^{-1} and the angiosperm median of 587 μm h^{-1}. Secondly, the higher

among-lineage variance is due to many transitions to both faster and slower *PTGR* optima within

extant angiosperms. Transitions to slower rates within angiosperms are concentrated on lineages

that have evolved delayed fertilization, such Fagales, orchids and others, or high selfing rates,

which suggests relaxation of directional selection on *PTGR* (Williams and Reese, 2019). In

contrast, gymnosperm *PTGRs* were likely ancestrally slow (Figure 4).

There are several non-mutually exclusive hypotheses for what triggered the evolution of

fast *PTGRs* in angiosperms. First, Mulcahy (1979) invoked a shift to much higher intensity of

pollen competition in angiosperms as a driver of the origin and continued evolution of faster

growth rates. Notably, no other type of tip-growing cell in land plants (whether gametophytic or

sporophytic) has evolved comparably fast tip-growth rates and none of those cell types, including

gymnosperm pollen tubes, experience intense competition for resources (Williams et al., 2016).

Secondly, gymnosperm *PTGRs* may be slow because they lack novel biophysical or

physiological attributes of pollen tubes and/or those attributes enabled faster *PTGRs* to evolve in

angiosperms (Hoekstra, 1983; Derksen et al., 1999; Fernando et al., 2005; Williams, 2008,

2009). Thirdly, with or without pollen competition, rapid *PTGRs* may have been necessary as

angiosperm sporophytes transitioned to a much faster reproductive cycle (Stebbins, 1974;

Williams, 2012; Williams and Reese, 2019). Finally, our results suggest a new possibility, that

strong differences in genome-level processes have impacted the evolution of angiosperm *PTGRs*

relative to their living and extinct seed plant relatives.

Mechanisms of genome size change and PTGR evolution within seed plants – A major finding

of this study is that angiosperm neo-polyploids evolved around a much faster *PTGR* optimum

(1648 μm h^{-1}) than diploids (595 μm h^{-1}), despite several sources of variation in the data. First,

neo-polyploids were by definition derived within genera, and their smaller sample size and

shorter branch lengths reduced the power to estimate parameters relative to diploids, as reflected

by the larger standard error around the neo-polyploid optimum. Nevertheless, the proportion of

neo-polyploids in our data set (33% of angiosperms) is almost exactly that found in the full

Wood et al. (2009) data set and similar to that in other studies (Mayrose et al., 2011; Barker et

al., 2016; Landis et al., 2018).

There was also biological variability in our dataset. In our taxon sampling, we were

agnostic to variation in mating systems and modes of polyploid origins, since our interest was in

how *PTGR* has evolved in natural stabilized polyploids. In retrospect, our sample does seem

representative. Of 14 angiosperm polyploids whose mode of origin has been studied, seven were

autopolyploid and seven allopolyploid, similar to the nearly-equal proportions found by Barker

et al. (2016). Furthermore, among 16 polyploids for which mating system has been studied, eight

were fully outcrossing, seven were self-compatible (two autogamous, two mixed mating, and

three unknown), and one was apomictic – a not unusual distribution (Goodwillie et al., 2005;

Gibbs, 2014; Ashman et al., 2014). Thus, our taxon sampling seems not be have been greatly

biased. Even with such information, predicting the magnitude of genetic variation in polyploids

is not so simple. For example, autotetraploids originate with a subset of the genetic variation in

the diploid progenitor population but they often outcross and hybridize, whereas allopolyploids

can be highly heterozygous when they originate, but often are highly selfing (Stebbins, 1974;

Soltis and Soltis, 1999; Barringer 2007; Whitney et al., 2010). Hence, despite several sources of

heterogeneity, the faster *PTGR* optimum of neo-polyploids indicates that *PTGR* acceleration

evolves either at the time of WGDs or during the time period in which the descendant species

retain a polyploid chromosome number.

The closest approximation of the initial effect of polyploidy on *PTGR* is the comparison

of diploids with their intraspecific, autopolyploid cytotypes. In all 11 pairs, *PTGRs* of

autopolyploid cytotypes were slower than or equal to those of their intraspecific diploid

progenitors. We should re-emphasize that all studies involved in vivo crosses among diploid

sporophytes (1x pollen on 2x pistils) compared to crosses among tetraploid sporophytes (2x

pollen on 4x pistils), in keeping with our goal of generalizing effects on *PTGR* in stabilized

polyploids. Nucleotypic effects acting to slow *PTGR* should be most apparent in autopolyploids

at inception, because there is lower potential for heterosis. Thus, the lack of any examples of

faster *PTGR* in neo-autotetraploid cytotypes than in their diploid progenitors suggests that

increased gene dosage by itself generally does not initially fully offset nucleotypic effects.

Nucleotypic effects on *PTGR* could be substantial. Tube size affects *PTGR* in a linear

fashion, because larger tubes must make more tube wall per unit time, and since tube diameter is

constant during growth, the rate of wall production is directly proportional to tip extension rate

(Williams et al., 2016). Kostoff & Prokofieva (1935) reported in vivo pollen tubes to be 39%

larger in diameter in an allotetraploid *Nicotiana* relative to the mean of its presumed diploid

progenitors, and Iyengar (1938) found 8-53% larger tube diameters in tetraploid versus diploid

species of *Gossypium*.

Taken together our results suggest that nucleotypic effects are strong and act as a brake

on *PTGR* at inception (intraspecific polyploid analysis), but as neo-polyploids become stabilized

and persist over time, nucleotypic effects are more than offset by genotypic effects (within-genus

pairs and model-based analyses) which often produce faster *PTGRs* in angiosperms.

We found that DNA content has evolved around a significantly lower optimum in

angiosperms than in gymnosperms, even though angiosperms have a broad range of DNA C-

values that encompass the entire range of seed plant genome sizes (Fig. 3; see Leitch and Leitch,

2013 for a larger survey). Angiosperms also have great variation in ploidy level, a history of

speciation by polyploidy, and much evidence of past genome duplication (Ahuja, 2005; Wood et

al., 2009; Husband et al., 2013; Van de Peer et al., 2017; Landis et al., 2018). There were at least

1-7 WGDs in the lineages leading from the seed plant root to each of the tips in our *PTGR* tree,

and 33% of taxa (68/206 angiosperms versus 2/23 gymnosperms) were identified as neo-

polyploids. The often low DNA content and high ploidy levels of angiosperms are not surprising

given that genome duplication is commonly followed by rapid loss of DNA sequences, gene

fractionation by large-scale deletions, biased retention of genes with beneficial dosage effects,

and ultimately a return to an apparent diploid state in sporophytes (Conant and Wolfe, 2008;

Conant et al., 2014; Freeling et al., 2015; Dodsworth et al., 2016; Wendel et al., 2018). Thus, one

explanation for the much faster *PTGRs* of angiosperms relative to gymnosperms is that

widespread gene duplication by WGDs have often enabled transgressive evolution of faster

PTGRs leading to the observed pattern of convergent evolution of extremely fast *PTGRs* in many

unrelated lineages of monocots and eudicots.

WGDs have been rare in gymnosperms (Ahuja, 2005; Leitch et al., 2005; Wood et al.,

2009; Soltis et al., 2009; Husband et al., 2013; Leitch and Leitch, 2013; Lee and Kim, 2014) and

their high DNA contents are thought to be due mainly to high transposon activity without

repeated rounds of genome duplication (Leitch & Leitch, 2013; Lee and Kim, 2014). Hence,

gymnosperms may have experienced the nucleotypic effects of higher DNA content on pollen

tube dimensions, which is predicted to reduce *PTGR*, without the potential for counter-balancing

effects, such as initially higher gene dosage and heterozygosity followed by gene sorting during

the diploidization process. Our finding of a negative correlation between *PTGR* and DNA

content in gymnosperms, but not in angiosperms supports that hypothesis.

Though gymnosperm *PTGRs* are likely affected by tube sizes, nucleotypic effects do not

account for the magnitude of the difference in their slow *PTGRs* relative to those of angiosperms.

Gymnosperm pollen tubes can range up to 300 μm in diameter (Coulter and Chamberlain, 1928;

Gifford and Foster, 1989), but many species of siphonogamous conifers and Gnetales have

angiosperm-like pollen tube diameters in the 10 to 20 μm range. Yet no gymnosperm has

evolved a *PTGR* faster than 20 μm h^{-1}. It has been argued that their pecto-cellulosic wall

structure is a limitation relative to angiosperm pollen tube walls, which use the plasma

membrane-bound enzymes callose synthase and pectin-methylesterase in a novel way to more

rapidly synthesize a strong and durable tube cell wall and callose plugs (Derksen, 1999,

Abercrombie et al., 2012; Wallace and Williams, 2017). However, other types of pecto-cellosic

tip-growing cells, such as root hairs, grow faster than gymnosperm pollen tubes (Williams et al.,

2016). Thus, it seems likely that the extremely slow growth rates of gymnosperm pollen tubes

reflect an ancestrally antagonistic relationship between maternal tissues and pollen tubes that

functioned as invasively growing rhizoids, coupled with a lack of selection for faster growth rate

due to the absence of pollen competition and a long period between pollination and fertilization.

Our results also suggest a lack of opportunity for genotypic effects to evolve due to the rarity of

WGDs.

Conclusions - Studies across the tree of life have consistently shown that ploidy level and DNA

content are correlated with cell size and metabolic rate (Cavalier-Smith, 1978; Gregory, 2001;

Cavalier-Smith, 2005). Pollen tube dimensions and energetics affect the amount of cell wall

material produced per unit of growth and the rate at which cell wall is produced, which together

determine *PTGR*. In gymnosperms, *PTGR* was negatively correlated with genome size, but in

angiosperms, where the effects of WGDs are much more prevalent, there was no such

correlation, and neo-polyploids evolved around a higher *PTGR* optimum than diploids. These

results support the expectation that genome size increases incur nucleotypic effects that act as a

brake on growth rate. The degree to which genotypic effects counterbalance nucleotypic effects

depends on the historical nature and time since genome size increase in any particular lineage.

Understanding causal relationships between genome size, ploidy and *PTGR* will involve

mechanistic studies of tube cell dimensions and wall synthesis rates in haploid and polyploid

gametophytes. On the other hand, there appears to be great variation in the tug of war between

genotypic and nucleotypic effects, and there are likely to be deeper evolutionary patterns

underlying that variation.

ACKNOWLEDGEMENTS

We thank B. O'Meara and J. Beaulieu for advice on phylogenetic analyses, I. Leitch for data on

DNA content, and J. Edwards and M. Rankin for assistance in the lab. We are tremendously

grateful to several anonymous reviewers for their perceptive and useful comments. Partial

support to J.B.R. was provided by National Science Foundation award IOS 1052291 to J.H.W.

Authors Contributions: J.B.R. and J.H.W. jointly conceived of the study and wrote the paper;

J.B.R. collected data on genome sizes and ploidy levels, constructed the phylogenetic tree and

performed all comparative analyses; J.H.W. collected data on *PTGRs* and diploid-autopolyploid

PTGRs.

Data Accessibility Statement: Scripts written during the creation of this manuscript are

available on GitHub: https://github.com/jbr1848/PTGR.genome.evolution. The phylogenetic tree

created during this study can be found on TreeBase:

http://purl.org/phylo/treebase/phylows/study/TB2:S24291.

LITERATURE CITED

Abercrombie, J. M., B. C. O'Meara, A. R. Moffatt, and J. H. Williams. 2011. Developmental

evolution of flowering plant pollen tube cell walls: callose synthase (CalS) gene expression

patterns. *EvoDevo* 2: 14.

Ahuja, M. R. 2005. Polyploidy in gymnosperms: revisited. *Silvae Genetica* 54:2:59-69.

Arunkumar, R., E. B. Josephs, R. J. Williamson, and S. I. Wright. 2013. Pollen-specific, but not

sperm-specific, genes show stronger purifying selection and higher rates of positive selection

than sporophytic genes in *Capsella grandiflora*. *Molecular Biology and Evolution* 30: 2475–

2486.

Ashman, T-L., D. Bachtrog, H. Blackmon, E. E. Goldberg, M. W. Hahn, M. Kirkpatrick, J.

Kitano, J. E. Mank, et al. 2014. Tree of Sex: A database of sexual systems. *Scientific Data* 1:

140015.

Barker, M. S., N. Arrigo, A. E. Baniaga, Z. Li, & D. A. Levin. (2016). On the relative abundance

of autopolyploids and allopolyploids. *New Phytologist*, *210*(2), 391-398.

Beaulieu, J. M., and B. O'Meara. 2014. *OUwie*: analysis of evolutionary rates in an OU

framework. *R package version* 1.

Bennett, M. D. 1971. The duration of meiosis. *Proceedings of the Royal Society of London B:*

Biological Sciences 178: 277–299.

Bennett, M. D. 1972. Nuclear DNA content and minimum generation time in herbaceous plants.

Proceedings of the Royal Society of London B: Biological Sciences 181: 109–135.

Bennett, M. D., and I. J. Leitch. 2012. Plant DNA C-values Database (Release 6.0).

Birchler, J. A., H. Yao, S. Chudalayandi, D. Vaiman, and R. A. Veitia. 2010. Heterosis. *Plant*

Cell: 110.076133.

Blomberg, S. P., T. Garland Jr., and A. R. Ives. 2003. Testing for phylogenetic signal in

comparative data: behavioral traits are more labile. *Evolution* 57: 717–745.

Burnham, K. P., and D. R. Anderson. 2002. Model selection and multimodel inference: a

practical information-theoretic approach. *New York*. *Springer-Verlag*.

Cavalier-Smith, T. 1978. Nuclear volume control by nucleoskeletal DNA, selection for cell

volume and cell growth rate, and the solution of the DNA C-value paradox. *Journal of Cell*

Science 34: 247–278.

Cavalier-Smith, T. 2005. Economy, speed and size matter: evolutionary forces driving nuclear

genome miniaturization and expansion. *Annals of Botany* 95: 147–175.

Christenhusz, M. J., J. L. Reveal, A. Farjon, M. F. Gardner, R. R. Mill, and M. W. Chase. 2011.

A new classification and linear sequence of extant gymnosperms. *Phytotaxa* 19: 55–70.

Comai, L. 2005. The advantages and disadvantages of being polyploid. *Nature Reviews Genetics*

6: 836.

Conant, G. C., J. A. Birchler, and J. C. Pires. 2014. Dosage, duplication, and diploidization:

clarifying the interplay of multiple models for duplicate gene evolution over time. *Current*

Opinion in Plant Biology 19: 91–98.

Conant, G. C., and K. H. Wolfe. 2008. Turning a hobby into a job: how duplicated genes find

new functions. *Nature Reviews Genetics* 9: 938.

Coulter, J. M., and C. J. Chamberlain. 1928. *Morphology of Gymnosperms* (4th ed.). University

of Chicago Press, Chicago.

Cosgrove, D. J. 2005. Growth of the plant cell wall. *Nature Reviews Molecular Cell Biology* 6:

850.

De Smet, R., K. L. Adams, K. Vandepoele, M. C. Van Montagu, S. Maere, and Y. Van de Peer.

2013. Convergent gene loss following gene and genome duplications creates single-copy

families in flowering plants. *Proceedings of the National Academy of Sciences* 110: 2898–2903.

Derksen, J., Y. Li, B. Knuiman, and H. Geurts. 1999. The wall of *Pinus sylvestris* L. pollen

tubes. *Protoplasma* 208: 26–36.

Dodsworth, S., M. W. Chase, and A. R. Leitch. 2016. Is post-polyploidization diploidization the

key to the evolutionary success of angiosperms? *Botanical Journal of the Linnean Society* 180:

1–5.

Doyle, J. J., and J. E. Coate. 2019. Polyploidy, the Nucleotype, and Novelty: The Impact of

Genome Doubling on the Biology of the Cell. *International Journal of Plant Sciences* 180: 1–52.

Eastman, J. M., L. J. Harmon, and D. C. Tank. 2013. Congruification: support for time scaling

large phylogenetic trees. *Methods in Ecology and Evolution* 4: 688–691.

Felsenstein, J. 1985. Phylogenies and the comparative method. *American Naturalist* 125: 1–15.

Fernando, D. D., M. D. Lazzaro, and J. N. Owens. 2005. Growth and development of conifer

pollen tubes. *Sexual Plant Reproduction* 18: 149–162.

Freeling, M., M. J. Scanlon, and J. E. Fowler. 2015. Fractionation and subfunctionalization

following genome duplications: mechanisms that drive gene content and their consequences.

Current Opinion in Genetics & Development 35: 110–118.

Friedman, W. E. 1993. The evolutionary history of the seed plant male gametophyte. *Trends in*

Ecology & Evolution 8: 15–21.

Gibbs, P. E. 2014. Late-acting self-incompatibility–the pariah breeding system in flowering

plants. *New Phytologist* 203: 717–734.

Gifford, E. M., and A. S. Foster. 1989. Morphology and Evolution of Vascular Plants. W. H.

Freeman, New York.

Goodwillie, C., S. Kalisz, and C. G. Eckert. 2005. The evolutionary enigma of mixed mating

systems in plants: occurrence, theoretical explanations, and empirical evidence. *Annual Review*

of Ecology Evolution and Systematics. 36: 47–79.

Grafen, A. 1989. The phylogenetic regression. *Philosophical Transactions of the Royal Society*

of London. *Series B, Biological Sciences* 326: 119–157.

Gregory, T. R. 2001. The bigger the C-value, the larger the cell: genome size and red blood cell

size in vertebrates. *Blood Cells, Molecules, and Diseases* 27: 830–843.

Hafidh, S., J. Fíla, and D. Honys. 2016. Male gametophyte development and function in

angiosperms: a general concept. *Plant Reproduction* 29: 31-51.

Harmon, L. J., J. T. Weir, C. D. Brock, R. E. Glor, and W. Challenger. 2008. GEIGER:

investigating evolutionary radiations. *Bioinformatics* 24: 129–131.

Ho, L. S. T., and C. Anné. 2014. *Phylolm: phylogenetic linear regression*. *R package version*

2.1.

Hoekstra, F. A. 1983. Physiological evolution in angiosperm pollen: possible role of pollen

vigour. In: Mulcahy DL, Ottaviano E, eds. Pollen: Biology and Implications for Plant Breeding,

35–41. Elsevier Science, Amsterdam.

Husband, B. C. 2016. Effect of inbreeding on pollen tube growth in diploid and tetraploid

Chamerion angustifolium: Do polyploids mask mutational load in pollen? *American Journal of*

Botany 103: 532–540.

Husband, B. C., S. J. Baldwin, and J. Suda. 2013. The incidence of polyploidy in natural plant

populations: major patterns and evolutionary processes. *In* J. Greilhuber, J. Doležel, and J. F.

Wendel [eds.], Plant Genome Diversity Volume 2, 255–276. Springer.

Husband, B. C., and D. W. Schemske. 1997. The effect of inbreeding in diploid and tetraploid

populations of *Epilobium angustifolium* (Onagraceae): implications for the genetic basis of

inbreeding depression. *Evolution* 51: 737–746.

Ingram, T., and D. L. Mahler. 2013. SURFACE: detecting convergent evolution from

comparative data by fitting Ornstein-Uhlenbeck models with stepwise Akaike Information

Criterion. *Methods in Ecology and Evolution* 4: 416–425.

Iyengar N. K. 1938. Pollen-tube studies in *Gossypium*. *Journal of Genetics* 37: 69–106.

Jörgensen A, and C. Rydin. 2015. Reproductive morphology in the *Gnetum cuspidatum* group

(Gnetales) and its implications for pollination biology in the Gnetales. *Plant Ecology and*

Evolution 148: 387–396.

Kostoff, D., and A. Prokofieva. 1935. Studies on the pollen-tubes. I. The growth potency of the

pollen-tubes in *Nicotiana* in connection with the length of the styles and some other factors. *Bul*.

Inst. *Genetics*, *Acad*. *Sci*. *Leningrad* 10: 65–82.

Lande, R., and D. W. Schemske. 1985. The evolution of self-fertilization and inbreeding

depression in plants. I. Genetic models. *Evolution* 39: 24–40.

Landis, J. B., D. E. Soltis, Z. Li, H. E. Marx, M. S. Barker, D. C. Tank, and P. S. Soltis. 2018.

Impact of whole-genome duplication events on diversification rates in angiosperms. *American*

Journal of Botany.

Lee, S-I., and N-S. Kim. 2014. Transposable elements and genome size variations in plants.

Genomics & Informatics 12: 87–97.

Leitch, I. J., M. W. Chase, and M. D. Bennett. 1998. Phylogenetic analysis of DNA C-values

provides evidence for a small ancestral genome size in flowering plants. *Annals of Botany* 82:

85–94.

Leitch, A. R., and I. J. Leitch. 2012. Ecological and genetic factors linked to contrasting genome

dynamics in seed plants. *New Phytologist* 194: 629–646.

Leitch, I. J., and A. R. Leitch. 2013. Genome size diversity and evolution in land plants. *In* J.

Greilhuber, J. Doležel, and J. F. Wendel [eds.], Plant Genome Diversity Volume 2, 307–322.

Springer.

Leitch, I. J., D. E. Soltis, P. S. Soltis, and M. D. Bennett. 2005. Evolution of DNA amounts

across land plants (Embryophyta). *Annals of Botany* 95: 207–217.

Li, Z., A. E. Baniaga, E. B. Sessa, M. Scascitelli, S. W. Graham, L. H. Rieseberg, and M. S.

Barker 2015. Early genome duplications in conifers and other seed plants. *Science Advances* 1:

e1501084.

Magallón, S., S. Gómez-Acevedo, L. L. Sánchez-Reyes, and T. Hernández-Hernández. 2015. A

metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New*

Phytologist 207: 437–453.

Martins, E. P., T. F. Hansen. 1997. Phylogenies and the comparative method: a general approach

to incorporating phylogenetic information into the analysis of interspecific data. *The American*

Naturalist 149: 646–667.

Mayrose, I., Zhan, S. H., Rothfels, C. J., Magnuson-Ford, K., Barker, M. S., Rieseberg, L. H.,

and S. P. Otto. 2011. Recently formed polyploid plants diversify at lower rates. *Science* 333:

1257-1257.

Mulcahy, D.L. 1979. The rise of the angiosperms: a genecological factor. *Science* 206: 20–23.

Ohri, D., and T. N. Khoshoo. 1986. Genome size in gymnosperms. *Plant Systematics and*

Evolution 153: 119–132.

Otto, S. P., M. F. Scott, and S. Immler. 2015. Evolution of haploid selection in predominantly

diploid organisms. *Proceedings of the National Academy of Sciences* 112: 15952–15957.

Owens, J. N., T. Takaso, and C. J. Runions. 1998. Pollination in conifers. *Trends in Plant*

Science 3: 479–485.

Pagel, M. 1997. Inferring evolutionary processes from phylogenies. *Zoologica Scripta* 26: 331–

348.

Pagel, M. 1999. Inferring the historical patterns of biological evolution. *Nature* 401: 877.

Panchy, N., M. Lehti-Shiu, and S-H. Shiu. 2016. Evolution of gene duplication in plants. *Plant*
Physiology 171: 2294–2316.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: analyses of phylogenetics and evolution in
R language. *Bioinformatics* 20: 289–290.

Price, H. 1988. DNA Content Variation Among Higher-Plants. *Annals of the Missouri Botanical*
Garden 75: 1248–1257.

Revell, L. J. 2012. Phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* 3: 217–223.

Rutley, N., and D. Twell. 2015. A decade of pollen transcriptomics. *Plant Reproduction* 28: 73-
89.

Smith, S. A., and M. J. Donoghue. 2008. Rates of molecular evolution are linked to life history
in flowering plants. *Science* 322: 86–89.

Smith, S. A., and C. W. Dunn. 2008. Phyutility: a phyloinformatics tool for trees, alignments and
molecular data. *Bioinformatics* 24: 715–716.

Snodgrass, S. J., J. Jaraczek and J. F. Wendel. 2016. An examination of nucleotypic effects in
diploid and polyploid cotton. *AoB PLANTS*. 8: plw082.

Soltis, D. E., V. A. Albert, J. Leebens-Mack, C. D. Bell, A. H. Paterson, C. Zheng, D. Sankoff et
al. 2009. Polyploidy and angiosperm diversification. *American Journal of Botany* 96: 336–348.

Soltis, D. E., and P. S. Soltis. 1999. Polyploidy: recurrent formation and genome evolution.
Trends in Ecology & Evolution 14: 348–352.

Soltis, P. S., & Soltis, D. E. 2000. The role of genetic and genomic attributes in the success of
polyploids. *Proceedings of the National Academy of Sciences* 97: 7051-7057.

Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* 30: 1312–1313.

Stebbins, G. L. 1974. Flowering plants: evolution above the species level. *London: Arnold xviii,*

399p. Illustrations. General (KR, 197500089).

Swift, H. 1950. The constancy of desoxyribose nucleic acid in plant nuclei. *Proceedings of the*

National Academy of Sciences 36: 643–654.

Tanksley, S. D., D. Zamir, and C. M. Rick. 1981. Evidence for extensive overlap of sporophytic

and gametophytic gene expression in *Lycopersicon esculentum. Science* 213: 453–455.

Van de Peer, Y., E. Mizrachi, and K. Marchal. 2017. The evolutionary significance of

polyploidy. *Nature Reviews Genetics* 18: 411.

Wallace, S., and J. H. Williams. 2017. Evolutionary origins of pectin methylesterase genes

associated with novel aspects of angiosperm pollen tube walls. *Biochemical and biophysical*

research communications 487: 509–516.

Wendel, J. F., D. Lisch, G. Hu, and A. S. Mason. 2018. The long and short of doubling down:

polyploidy, epigenetics, and the temporal dynamics of genome fractionation. *Current Opinion in*

Genetics & Development 49: 1–7.

Whitney, K. D., E. J. Baack, J. L. Hamrick, M. J. W. Godt, B. C. Barringer, M. D. Bennett, C. G.

Eckert et al. 2010. A role for nonadaptive processes in plant genome size evolution? *Evolution:*

64: 2097–2109.

Williams, J. H. 2008. Novelties of the flowering plant pollen tube underlie diversification of a

key life history stage. *Proceedings of the National Academy of Sciences* 105: 11259–11263.

Williams, J. H. 2009. *Amborella trichopoda* (Amborellaceae) and the evolutionary

developmental origins of the angiosperm progamic phase. *American Journal of Botany* 96: 144–

165.

Williams, J. H. 2012. Pollen tube growth rates and the diversification of flowering plant

reproductive cycles. *International Journal of Plant Sciences* 173: 649–661.

Williams, J. H., J. A. Edwards, and A. J. Ramsey. 2016. Economy, efficiency, and the evolution

of pollen tube growth rates. *American Journal of Botany* 103: 471–483.

Williams, J. H., and J. B. Reese. 2019. Evolution of development of pollen performance. *In* U.

Grossinklaus [ed.], Plant Development and Evolution. Current Topics in Developmental

Biology, Volume 131. Chapter 12. 299-336. Elsevier.

Williams, J. H., M. L. Taylor, and B. C. O'Meara. 2014. Repeated evolution of tricellular (and

bicellular) pollen. *American Journal of Botany* 101: 559–571.

Wood, T. E., N. Takebayashi, M. S. Barker, I. Mayrose, P. B. Greenspoon, and L. H. Rieseberg.

2009. The frequency of polyploid speciation in vascular plants. *Proceedings of the National*

Academy of Sciences 106: 13875–13879.

Table 1: Phylogenetic generalized least squares regression of $\log_{10}$ *PTGR* as a function of $\log_{10}$
C-value. Only models contributing more than 1% of total weight are included. *P* values are for
the slope of the regression. Gymnosperm averaged model: $\log_{10}$ *PTGR* = 1.46 (± 0.62) - 1.09 (±
0.49)*($\log_{10}$ C-value).

Seed plants (*N* = 183)			Angiosperms (*N* = 161)			Gymnosperms (*N* = 23)		
Model	**Weight**	*P*	**Model**	**Weight**	*P*	**Model**	**Weight**	*P*
kappa	0.999	0.463	kappa	0.975	0.284	OU	0.265	0.020
			lambda	0.024	0.221	delta	0.257	0.006
						kappa	0.193	0.445
						BM	0.121	0.001
						lambda	0.119	0.327
						EB	0.044	0.001

Table 2*: Parameter estimates for angiosperm *PTGR* analyses under different evolutionary models. Note that OU1 is a single optimum model, and the rest specify separate "diploid" (paleo-polyploid) and neo-polyploid optima. BM1 and BMS models contributed <1% model weight and were excluded.

Model	ΔAICc	Model weight	Diploid σ^2	Polyploid σ^2	Diploid α	Polyploid α	Diploid optimum	Polyploid optimum
OUMA	[332.76]	0.373	0.099		0.077	0.074	2.776	3.285
OUMV	0.23	0.333	0.099	0.057	0.077		2.768	3.262
OUM	1.53	0.174	0.090		0.080		2.760	3.263
OU1	2.26	0.120	0.089		0.077		2.812	
AVERAGED MODEL		~1.0	0.096 ± 0.177	0.082 ± 0.220	0.078 ± 0.213	0.077 ± 0.217	2.775 ± 0.079	3.217 ± 0.231

 FIGURE LEGENDS

Figure 1. Predicted initial effects of large increases in genome size on pollen tube growth

rate (*PTGR*). The dashed line indicates an ancestral haploid (1x) *PTGR*. Upon transition to a

larger (> 1x) genome size, nucleotypic effects should act to decrease *PTGR* regardless of

mechanism of change. Genotypic effects are only present after WGD or large-scale gene

duplications and are predicted to increase *PTGR* via increased gene dosage and heterozygosity.

The magnitude of heterosis due to initial increase in heterozygosity is expected to scale with

genetic variation in the descendent taxon. The ancestral haploid *PTGR* can only be conserved

when genotypic and nucleotypic effects perfectly offset each other.

Figure 2: Relationship between pollen tube growth rate (*PTGR*) and DNA content (1C-

value) in seed plants. The model- averaged slope of the PGLS regression is shown for

gymnosperms (green points, $N = 161$), whereas slopes for seed plants (all points, $N = 183$) and

angiosperms (purple points, $N = 23$) were non-significant. Optima (with standard error bars) for

each group (from model-based analyses in Tables S3, S4) are included for illustrative purposes.

Figure 3: Inferred pattern of pollen tube growth rate (*PTGR*) and genome size changes in

seed plants. Contour plot comparing *PTGR* evolution (left, $\mu m\ h^{-1}$) and C-value evolution (right,

picograms) ($N = 183$). Scale bar at the bottom of each phylogeny indicates 100 million years.

GYM = gymnosperms; ANA = Amborellales, Nymphaeales, Austrobaileyales, Chloranthales,

eumagnoliids; MONO = monocots.

Figure 4: Coincident evolution of pollen tube growth rate (*PTGR*) and DNA content (C-value). Paired *SURFACE* plot showing regime shifts in *PTGR* (left) versus DNA content (right) (*N* = 183). Nodes which have experienced a regime shift along the stem leading to it are marked with magenta diamonds (not all *PTGR* shifts are shown, since *PTGR* tree has been pruned to match C-value tree). Branch colors: *gray* = seed plant ancestral optimum (*PTGR* θ = 0.147; C-value θ = 2.71); *green* = ancestral optimum for angiosperms (*PTGR* θ = 2.47; C-value θ = 0.702); *red* = derived lineages following a shift to a higher optimum than previously; *blue* = derived lineages following a shift to a lower optimum than previously. Black arrows indicate instances where shifts in *PTGR* and C-value coincide. Scale bar at the bottom of each phylogeny indicates 100 million years. GYM = gymnosperms; A = Amborellales, Nymphaeales, Autrobaileyales, Chloranthales, eumagnoliids; MONO = monocots.

Additional Supporting Information may be found online in the supporting information section at the end of the article:

Appendix S1: **119 additional *PTGR* values and references not reported in Williams 2012.**

Appendix S2: Pollen tube growth rate (*PTGR*) evolution across Spermatophytes.

Appendix S3: Summary statistics for pollen tube growth rate (*PTGR*) of *Magnolia grandiflora*.

Appendix S4: Sensitivity analysis for the magnitude of log$_{10}$ *PTGR* error estimates.

Appendix S5: *PTGR* evolution in gymnosperms vs. angiosperms.

Appendix S6: C-value evolution in gymnosperms vs. angiosperms.

747   **Appendix S7: Closely-related taxon analyses**.

749   **Appendix S8: Phylogenetic ANCOVA results.**

ROSIDAE

ASTERIDAE

MONO

ANA

GYM

slower   faster
PTGR

smaller   larger
C-value

ROSIDAE

ASTERIDAE

MONO

A

GYM

Appendix S1: 119 additional *PTGR* values and references not reported in Williams 2012.

Taxon	*PTGR* (*µm h⁻¹*)	Reference
Abelmoschus esculentus	5217	Patil et al 2013
Abutilon x hybridum	1292	Cited in Sears 1937
Acacia mangium	46.5	Ngheim et al 2013
Acacia_auriculiformis	116	Ngheim et al 2013
Acer rubrum	182	van Ryn et al 1988, Radford et al 1968
Aegle marmelos	181	Bhardwaj and Tandon 2013
Albuca canadensis	1000	Johnson et al 2012
Albuca setosa	2477	Johnson et al 2012
Alstroemeria aurea	3091	Aizen and Raffaele 1998, De Jeu et al. 1996
Alstroemeria pelegrina	574	De Jeu et al. 1996
Anathallis	4320	Gontijo et al 2010
Antirrhinum controversum	131	Cario and Guemes 2014
Antirrhinum valentinum	160	Cario and Guemes 2014
Aureolaria pedicularia	15.2	Ramstetter and Mulcahy 1986
Bertholletia excels	800	Moritz and Ludders 1993
Betula papyrifera	11.1	Williams unpbl.
Boswellia serrata	182	Sunnichan et al 2005
Brassica rapa	371	Hiroi et al 2013
Calluna vulgaris	387	Behrend et al 2012, Mahy and Jacquemart 1999
Cambessedesia	587	dos Santos et al 2012
Camellia oleifera	338	Gao et al 2015
Carica papaya	214	Traub and O'Rork 1939
Ceiba pentandra	2500	Gribel et al 1999
Ceratonia siliqua	152	Von Haselberg et al. 2004
Chamaecrista fasciculata	637	Tucker 1996, Fenster and Sork 1988
Citrullus lanatus	1151	Sedgley and Buttrose 1978
Citrus maxima	58.3	Distephano et al 2012
Citrus medica	122	Distefano et al 2012
Citrus reticulata	51.3	Distefano et al 2012
Clarkia xantiana	630	Hove and Mazer 2013
Commiphora wrightii	42.4	Geetha et al 2013
Cornus florida	70.8	Reed 2004
Corylus heterophylla	62.5	Liu et al 2014
Cucumis anguria	774	Matsumoto et al. 2012
Cucumis melo	870	Matsumoto et al. 2012
Cucumis metulifer	1009	Matsumoto et al. 2012
Cybistax antisyphilitica	583	Bittencourt et al 2010

Cyrtandra kauaiensis	261	Johnson et al 2015
Cyrtandra longifolia	221	Johnson et al 2015
Cyrtandra platyphylla	305	Johnson et al 2015
Cytisus multiflorus	4.48	Valtueña et al 2010
Cytisus striatus	19.8	Rodriguez-Riaño et al 1999
Dalzellia zeylanica	1159	Sehgal et al 2011
Dianthus caryophyllus	3002	Larsen et al 1995
Downingia bacigalupii	853	Kaplan 1969
Echium vulgare	590	Melser et al 1997
Eruca vesicaria	275	Cited in Sears 1937
Eucalyptus globulus	58.3	Gore et al 1990
Faramea occidentalis	2183	Travers 1999
Ficus carica	47.6	Beck and Lord 1988
Fumana	273	Carrio and Guemes 2013
Guihaiothamnus acaulis	188	Xie et al 2013
Haberlea rhodopensis	152	Bogacheva-Milkoteva 2013
Handroanthus ochraceus	1617	Oliveira, pers. comm.
Handroanthus serratifolius	1617	Oliveira, pers. comm.
Hedyosmum brasiliense	97.2	Williams and Edwards, unpbl.
Hedyotis acutangula	991	Wu et al 2010
Helleborus foetidus	514	Vesprini and Pacini 2000
Heuchera micrantha	181	Rabe and Soltis 1999
Hippophae rhamnoides	20.9	Mangla et al 2013
Hymenaea	1667	Gibbs et al 1999
Ipomoea purpurea	7450	Shu-Mei Chang, pers. comm. 2014
Ipomopsis aggregata	2409	Sage et al 2006, Wolf et al 2001
Jathropa curcas	915	Abdelgadir et al 2012
Lactoris fernandeziana	40	Bernardello et al 1999
Lactuca sativa	3085	Einset 1944
Lagerstroemia indica	1175	Pounders et al 2006
Lathyrus chloranthus	271	Herrick et al 1993
Lathyrus odoratus	321	Herrick et al 1993
Limnocharis	467	Hall 1902
Linaria	392	Cited in Sears 1937
Lupinus arizonicus	442	Wainwright 1978
Magnolia grandiflora	828	Edwards, Rankin, and Williams, unpbl. 2014
Medicago rigidula	82.2	Sangduen et al 1983
Medicago sativa	192	Barnes and Cleveland 1963
Morinda parvifolia	957	Liu et al 2012
Mussaenda kwangtungensis	963	Luo et al 2015
Mussaenda shikokiana	816	Chen et al 2014
Nemesia strumosa	333	Sears 1937

Nivenia corymbosa	873	Goldblatt and Bernhardt 1990
Nivenia stokeii	1217	Goldblatt and Bernhardt 1990
Nuphar advena	835	Taylor and Williams, unpbl.
Nyctanthes arbor tristis	526	Bhatnagar and Uma 1969
Orchis anthropophora	357	Luca et al 2015
Orchis italica	357	Luca et al 2015
Oreocharis acaulis	1318	Guo et al 2013
Oroxylum indicum	3000	Gautam et al 2009
Paeonia brownii	20.8	Bernhardt et al 2013
Parthenium	1333	Gerstel and Riner 1950
Passiflora edulis	2174	Rego et al 2000
Phalaenopsis	208	Zhang and O-Neill 1993
Phoenix dactylifera	315	Reuveni et al 1986
Platanthera	462	Stickler et al 2013 (poster)
Plumbago zeylanica	12741	Russell 1985
Polypleurum stylosum	153	Khosla et al 2000
Potamogeton intortusifolius	267	Zhang et al 2010
Potamogeton perfoliatus	1585	Zhang et al 2010
Potamogeton wrightii	1483	Zhang et al 2010
Pseudopiptadenia	39	Pires and Freitas 2008
Restrepia	99	Millner et al 2015
Schisandra sphenanthera	88.9	Du et al 2012
Silene vulgaris	2323	Glaeti 2006
Solanum chacoense	396	Liu et al 2012
Solanum laxum	333	Lewis and Crowe 1958
Sorghum bicolor	3638	Heslop-Harrison et al 1984, Hodnett et al 2005
Spathodea campanulata	4028	Bittencourt et al 2003
Sporobolus anglicus	8943	Li et al 2008
Thryptomene calycina	320	Beardsell et al 1993
Ticodendron incognitum	382	Sogo and Tobe 2008
Torenia baillonii	3200	Kikuchi et al 2007
Torenia concolor	1900	Kikuchi et al 2007
Trimezia	2000	Bystedt and Vennigerholz 1991
Vaccinium corybosum	191	Knight and Scott 1964
Vaccinium myrtillus	67.1	Jacquemart and Thompson 1996
Vaccinium uliginosum	54.3	Jacquemart and Thompson 1996
Vaccinium vitis idaea	98.6	Jacquemart and Thompson 1996
Zeyheria montana	2554	Bittencourt and Semir 2004
Zeylanidium lichenoides	354	Chaudhary et al 2014, Sehgal et al 2014

References

Abdelgadir, H., S. Johnson, and J. Van Staden. 2012. Pollen viability, pollen germination and pollen tube growth in the biofuel seed crop *Jatropha curcas* (Euphorbiaceae). *South African Journal of Botany* 79:132-139.

Aizen, M. A., and E. Raffaele. 1998. Flowering-shoot defoliation affects pollen grain size and postpollination pollen performance in *Alstroemeria aurea*. *Ecology* 79:2133-2142.

Barnes, D., and R. Cleveland. 1963. Genetic evidence for nonrandom fertilization in alfalfa as influenced by differential pollen tube growth 1. *Journal of Crop Science* 3:295-297.

Beardsell, D., R. Knox, and E. Williams. 1993. Breeding system and reproductive success of *Thryptomene calycina* (Myrtaceae). *Australian Journal of Botany* 41:333-353.

Beck, N., and E. Lord. 1988. Breeding system in *Ficus carica*, the common fig. II. Pollination events. *American Journal of Botany* 75:1913-1922.

Behrend, A., T. Borchert, A. Müller, J. Tänzer, and A. Hohe. 2013. Malformation of gynoecia impedes fertilisation in bud-flowering *Calluna vulgaris*. *Plant Biology* 15:226-232.

Bernardello, G., G. J. Anderson, P. Lopez, M. A. Cleland, T. F. Stuessy, and D. J. Crawford. 1999. Reproductive biology of *Lactoris fernandeziana* (Lactoridaceae). *American Journal Of Botany* 86:829-840.

Bernhardt, P., R. Meier, and N. Vance. 2013. Pollination ecology and floral function of Brown's peony (*Paeonia brownii*) in the Blue Mountains of northeastern Oregon. *Journal of Pollination* Ecology 2:9-20.

Bhardwaj, V., and R. Tandon. 2013. Self-incompatibility and post-fertilization maternal regulation cause low fecundity in *Aegle marmelos* (Rutaceae). *Botanical Journal of the Linnean Society* 172:572-585.

Bhatnagar, S. P., and M. C. Uma. 1969. The structure of style and stigma in some Tubiflorae. *Phytomorphology* 19:99-109.

Bittencourt Jr, N. S., E. J. Pereira Jr, P. de Souza São-Thiago, and J. Semir. 2011. The reproductive biology of *Cybistax antisyphilitica* (Bignoniaceae), a characteristic tree of the South American savannah-like "Cerrado" vegetation. *Flora-Morphology, Distribution, Functional Ecology of Plants* 206:872-886.

Bittencourt, N. S., and J. Semir. 2004. Pollination biology and breeding system of *Zeyheria montana* (Bignoniaceae). *Plant Systematics and Evolution* 247:241-254.

Bittencourt, N. S. J., P. E. Gibbs, and J. Semir. 2003. Histological study of post-pollination events in *Spathodea campanulata* Beauv. (Bignoniaceae), a species with late-acting self-incompatibility. *Annals of Botany* 91:827-834.

Bogacheva-Milkoteva, K., E. Kozuharova, R. Claßen-Bockhoff, and A. Gogala. 2013. Pollination ecology of *Haberlea rhodopensis* Friv. (Gesneriaceae), a Tertiary relict endemic to the Balkan Peninsula. *Journal Comptes rendus de l'Académie bulgare des Sciences* 66.

Bystedt, P. A., and F. Vennigerholz. 1991. The transmitting tract in *Trimezia fosteriana* (Iridaceae). III. Pollen tube growth in the stigma, style and ovary. *Nordic Journal of Botany* 11:459-464.

Carrió, E., and J. Güemes. 2013. The role of a mixed mating system in the reproduction of a Mediterranean subshrub (*Fumana hispidula*, Cistaceae). *Journal of Plant Research* 126:33-40.

Carrió, E., and J. Güemes. 2014. The effectiveness of pre-and post-zygotic barriers in avoiding hybridization between two snapdragons (*Antirrhinum* L.: Plantaginaceae). 176:159-172.

Chaudhary, A., P. Khanduri, R. Tandon, P. Uniyal, and H. M. Ram. 2014. Central cell degeneration leads to three-celled female gametophyte in *Zeylanidium lichenoides* Engl. (Podostemaceae). *South African Journal of Botany* 91:99-106.

Chen, S., Z. Luo, and D. Zhang. 2014. Pre-and post-zygotic reproductive isolation between co-occurring *Mussaenda pubescens var. alba* and *M. shikokiana* (Rubiaceae*). Journal of Integrative Plant Biology* 56:411-419.

de Assis Pires, J. P., and L. Freitas. 2008. Reproductive biology of two tree species of Leguminosae in a Montane Rain Forest in southeastern Brazil. *Flora-Morphology, Distribution, Functional Ecology of Plants* 203:491-498.

de Jeu, M. J., F. G. Caldere, and J. L. van Went. 1996. Sporogenesis, gametogenesis, and progamic phase in *Alstroemeria. Canadian Journal of Botany* 74:1354-1361.

Distefano, G., A. Hedhly, G. Las Casas, S. La Malfa, M. Herrero, and A. Gentile. 2012. Male–female interaction and temperature variation affect pollen performance in *Citrus. Scientia Horticulturae* 140:1-7.

dos Santos, A. P. M., C. M. Fracasso, M. Luciene dos Santos, R. Romero, M. Sazima, and P. E. Oliveira. 2012. Reproductive biology and species geographical distribution in the Melastomataceae: a survey based on New World taxa. *Annals of Botany* 110:667-679.

Du, W., L. J. Huang, and X. F. Wang. 2012. Deceit pollination and the effect of deforestation on reproduction in dioecious *Schisandra sphenanthera* (Schisandraceae) in central China. *Journal of Systematics and Evolution* 50:36-44.

Einset, J. 1944. Cytological basis for sterility in induced autotetraploid lettuce (*Lactuca sativa* L.). *American Journal of Botany* :336-342.

Fenster, C. B., and V. L. Sork. 1988. Effect of crossing distance and male parent on *in vivo* pollen tube growth in *Chamaecrista fasciculata. American Journal of Botany* 75:1898-1903.

Gao, C., D. Yuan, Y. Yang, B. Wang, D. Liu, and F. Zou. 2015. Pollen tube growth and double fertilization in *Camellia oleifera. Journal of the American Society for Horticultural Science* 140:12-18.

Gautam, M., R. Tandon, and H. M. Ram. 2009. Pollination ecology and breeding system of *Oroxylum indicum* (Bignoniaceae) in the foothills of the Western Himalaya. *Journal of Tropical Ecology* 25:93-96.

Geetha, K., A. Kawane, A. K. Bishoyi, A. Phurailatpam, C. Ankita, S. Malik, R. Srinivasan, and S. Bhat. 2013. Characterization of mode of reproduction in *Commiphora wightii* [(Arnot) Bhandari] reveals novel pollen–pistil interaction and occurrence of obligate sexual female plants. *Trees* 27:567-581.

Gerstel, D. U., and M. E. Riner. 1950. Self-Incompatibility studies in *Guayule* .1. Pollen-tube behavior. *Journal of Heredity* 41:49-55.

Gibbs, P. E., P. E. Oliveira, and M. B. Bianchi. 1999. Postzygotic control of selfing in *Hymenaea stigonocarpa* (Leguminosae-Caesalpinioideae), a bat-pollinated tree of the Brazilian Cerrados. *International Journal of Plant Sciences* 160:72-78.

Glaettli, M., and J. Goudet. 2006. Variation in the intensity of inbreeding depression among successive life-cycle stages and generations in gynodioecious *Silene vulgaris* (Caryophyllaceae). *Journal of Evolutionary Biology* 19:1995-2005.

Goldblatt, P., and P. Bernhardt. 1990. Pollination biology of *Nivenia* (Iridaceae) and the presence of heterostylous self-compatibility. *Israeli Journal of Plant Sciences* 39:93-111.

Gontijo, S. L., A. R. Barbosa, M. C. de Melo, and E. L. Borba. 2010. Occurrence of different sites of self-incompatibility reaction in four *Anathallis* (Orchidaceae, Pleurothallidinae) species. *Plant Species Biology* 25:129-135.

Gore, P. L., B. M. Potts, P. W. Volker, and J. Megalos. 1990. Unilateral cross-incompatibility in *Eucalyptus*: the case of hybridization between *E. globulus* and *E. nitens*. *Australian Journal of Botany* 38:383-394.

Gribel, R., P. E. Gibbs, and A. L. Queiróz. 1999. Flowering phenology and pollination biology of *Ceiba pentandra* (Bombacaceae) in Central Amazonia. *Journal of Tropical Ecology* 15:247-263.

Guo, Y.-F., Y.-Q. Wang, and A. Weber. 2013. Floral ecology of *Oreocharis acaulis* (Gesneriaceae): An exceptional case of "preanthetic" protogyny combined with approach herkogamy. *Flora-Morphology, Distribution, Functional Ecology of Plants* 208:58-67.

Hall, J. G. 1902. An embryological study of *Limnocharis emarginata*. *Botanical Gazette* 33:214-219.

Herrick, J., B. Murray, and K. Hammett. 1993. Barriers preventing hybridisation of Lathyrus odoratus with *L. chloranthus* and *L. chrysanthus*. *New Zealand Journal of Crop Horticultural Science* 21:115-121.

Heslop-Harrison, Y., B. Reger, and J. Heslop-Harrison. 1984. The pollen-stigma interaction in the grasses. 6. The stigma ('silk') of *Zea mays* L. as host to the pollens of *Sorghum bicolor* (L.) Moench and *Pennisetum americanum* (L.) Leeke. *Acta Botanica Nederlandica* 33:205-227.

Hiroi, K., M. Sone, S. Sakazono, M. Osaka, H. Masuko-Suzuki, T. Matsuda, G. Suzuki, K. Suwabe, and M. Watanabe. 2013. Time-lapse imaging of self-and cross-pollinations in *Brassica rapa*. *Annals of Botany* 112:115-122.

Hodnett, G. L., B. L. Burson, W. L. Rooney, S. L. Dillon, and H. J. Price. 2005. Pollen-pistil interactions result in reproductive isolation between *Sorghum bicolor* and divergent *Sorghum* species. *Crop science* 45:1403-1409.

Hove, A. A., and S. J. Mazer. 2013. Pollen performance in *Clarkia* taxa with contrasting mating systems: implications for male gametophytic evolution in selfers and outcrossers. *Plants* 2:248-278.

Jacquemart, A.-L., and J. Thompson. 1996. Floral and pollination biology of three sympatric *Vaccinium* (Ericaceae) species in the Upper Ardennes, Belgium. *Canadian Journal of Botany* 74:210-221.

Johnson, M. A., D. K. Price, J. P. Price, and E. A. Stacy. 2015. Postzygotic barriers isolate sympatric species of *Cyrtandra* (Gesneriaceae) in Hawaiian montane forest understories. *American Journal Of Botany* 102:1870-1882.

Johnson, S. D., A. Jürgens, and M. Kuhlmann. 2012. Pollination function transferred: modified tepals of *Albuca* (Hyacinthaceae) serve as secondary stigmas. *Annals of Botany* 110:565-572.

Kaplan, D. R. 1969. Sporogenesis and gametogenesis in *Dowingia* (Campanulaceae; Loelioideae). *Bulletin of the Torrey Botanical Club* 96:418-434.

Khosla, C., K. Shivanna, and H. M. Ram. 2000. Reproductive biology of *Polypleurum stylosum* (Podostemaceae). *Journal of Aquatic Botany* 67:143-154.

Kikuchi, S., H. Kino, H. Tanaka, and H. Tsujimoto. 2007. Pollen tube growth in cross combinations between *Torenia fournieri* and fourteen related species. *Breeding Science* 57:117-122.

Knight Jr, R., and D. Scott. 1964. Effects of temperatures on self-and cross-pollination and fruiting of four highbush blueberry varieties. *Proceedings of the American Society for Horticultural Science*.

Larsen, P. B., E. N. Ashworth, M. L. Jones, and W. R. Woodson. 1995. Pollination-induced ethylene in carnation (role of pollen tube growth and sexual compatibility). *Plant Physiology* 108:1405-1412.

Lewis, D., and L. K. Crowe. 1958. Unilateral interspecific incompatibility in flowering plants. *Heredity* 12:233-256.

Li, H., S. An, Y. Zhi, C. Yan, L. Zhao, C. Zhou, Z. Deng, W. Su, and Y. Liu. 2008. Protogynous, pollen limitation and low seed production reasoned for the dieback of *Spartina anglica* in coastal China. *Plant Science* 174:299-309.

Liu, B., N. Boivin, D. Morse, and M. Cappadocia. 2012. A time course of GFP expression and mRNA stability in pollen tubes following compatible and incompatible pollinations in *Solanum chacoense. Sexual Plant Reproduction* 25:205-213.

Liu, J., H. Zhang, Y. Cheng, S. Kafkas, and M. Güney. 2014. Pistillate flower development and pollen tube growth mode during the delayed fertilization stage in *Corylus heterophylla* Fisch. *Plant Reproduction* 27:145-152.

Liu, Y., Z. L. Luo, X. Q. Wu, X. F. Bai, and D. X. Zhang. Functional dioecy in *Morinda parvifolia* (Rubiaceae), a species with stigma-height dimorphism. *Plant Systematics and Evolution* 298:775-785.

Luca, A., A. Palermo, F. Bellusci, and G. Pellegrino. 2015. Pollen competition between two sympatric *Orchis* species (Orchidaceae): the overtaking of conspecific of heterospecific pollen as a reproductive barrier. *Plant Biolo*gy 17:219-225.

Luo, Y., L. Lu, A. H. Wortley, D.-Z. Li, H. Wang, and S. Blackmore. 2015. Evolution of angiosperm pollen. 3. Monocots. *Annals of the Missouri Botanical Garden* 101:406-455.

Mahy, G., and A. L. Jacquemart. 1999. Early inbreeding depression and pollen competition in *Calluna vulgaris* (L.) Hull. *Annals of Botany* 83:697-704.

Mangla, Y., R. Tandon, S. Goel, and S. Raina. 2013. Structural organization of the gynoecium and pollen tube path in Himalayan sea buckthorn, *Hippophae rhamnoides* (Elaeagnaceae). *AoB Plants* 5.

Matsumoto, Y., M. Miyagi, N. Watanabe, and T. Kuboyama. 2012. Temperature-dependent enhancement of pollen tube growth observed in interspecific crosses between wild *Cucumis spp.* and melon (*C. melo* L.). *Scientia Horticulturae* 138:144-150.

Melser, C., M. C. Rademaker, and P. G. Klinkhamer. 1997. Selection on pollen donors by *Echium vulgare* (Boraginaceae). *Sexual Plant Reproduction* 10:305-312.

Millner, H. J., A. R. McCrea, and T. C. Baldwin. 2015. An investigation of self-incompatibility within the genus *Restrepia. American Journal Of Botany* 102:487-494.

Moritz, A., and P. Ludders. 1993. Pollen germination, pollen-tube growth and fertilization behavior of different Brazil nut clones (*Bertholletia excelsa* Humb And Bonpl). *Angewandte Botanik* 67:107-112.

Nghiem, Q., J. Harbard, C. Harwood, A. Griffin, T. Ha, and A. Koutoulis. 2013. Pollen-pistil interactions between autotetraploid and diploid *Acacia mangium* and diploid *A. auriculiformis. Journal of Tropical Forest Science*:96-110.

Patil, P., S. K. Malik, K. S. Negi, J. John, S. Yadav, G. Chaudhari, and K. V. Bhat. 2013. Pollen germination characteristics, pollen-pistil interaction and reproductive behaviour in interspecific crosses among *Abelmoschus esculentus* Moench and its wild relatives. *Grana* 52:1-14.

Pounders, C., S. Reed, and M. Pooler. 2006. Pollination biology of *Lagerstroemia indica* and several interspecific hybrids. *HortScience* 413:575-578.

Rabe, A. J., and D. E. Soltis. 1999. Pollen tube growth and self-incompatibility in *Heuchera micrantha var. diversifolia* (Saxifragaceae). *International Journal of Plant Sciences* 160:1157-1162.

Radford, A. E., H. E. Ahles, and C. R. Bell. 1968. Manual of the vascular flora of the Carolinas. University of North Carolina Press, Chapel Hill, NC:.

Ramstetter, J., and D. Mulcahy. 1986. Pollen competition in *Aureolaria pedicularia*. Pages 411-416 *Biotechnology and Ecology of Pollen*. Springer.

Reed, S. M. 2004. Self-incompatibility in *Cornus florida*. HortScience 39:335-338.

Rêgo, M., E. Rêgo, C. Bruckner, E. Da Silva, F. Finger, and K. Pereira. 2000. Pollen tube behavior in yellow passion fruit following compatible and incompatible crosses. *Theoretical and Applied Genetics* 101:685-689.

Reuveni, O., S. Abu, and S. Golobovitz. 1985. Date palm pollen germination and tube elongation on pistillate flowers cultured at different temperatures. Pages 91-96 *in* Symposium on Physiology of Productivity of Subtropical and Tropical Tree Fruits 175.

Rodríguez-Riaño, T., A. Ortega-Olivencia, and J. A. Devesa. 1999. Reproductive biology in two Genisteae (Papilionoideae) endemic of the western Mediterranean region: *Cytisus striatus* and *Retama sphaerocarpa*. *Canadian Journal of Botany* 77:809-820.

Russell, S. D. 1985. Preferential fertilization in Plumbago - Ultrastrucural evidencefor gamete-level recognition in an angiosperm. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 82:6129-6132.

Sage, T. L., M. V. Price, and N. M. Waser. 2006. Self-sterility in *Ipomopsis aggregata* (Polemoniaceae) is due to prezygotic ovule degeneration. *American Journal Of Botany* 93:254-262.

Sangduen, N., E. L. Sorensen, and G. H. Liang. 1983. Pollen germination and pollen tube growth following self-pollination and intra- and interspecific pollination of *Medicago* species. *Euphytica* 32:527-534.

Sears, E. R. 1937. Cytological phenomena connected with self-sterility in the flowering plants. *Genetics* 22:130.

Sedgley, M., and M. Buttrose. 1978. Some effects of light intensity, daylength and temperature on flowering and pollen tube growth in the watermelon (*Citrullus lanatus*). *Annals of botany* 42:609-616.

Sehgal, A., J. P. Khurana, M. Sethi, and H. Ara. 2011. Occurrence of unique three-celled megagametophyte and single fertilization in an aquatic angiosperm-*Dalzellia zeylanica* (Podostemaceae-Tristichoideae). *Sexual Plant Reproduction* 24:199-210.

Sehgal, A., N. Mann, and H. M. Ram. 2014. Structural and developmental variability in the female gametophyte of *Griffithella hookeriana*, *Polypleurum stylosum*, and *Zeylanidium lichenoides* and its bearing on the occurrence of single fertilization in Podostemaceae. *Plant reproduction* 27:205-223.

Sogo, A., and H. Tobe. 2008. Mode of pollen tube growth in pistils of *Ticodendron incognitum* (Ticodendraceae, Fagales) and the evolution of chalazogamy. *Botanical Journal of the Linnean Society* 157:621-631.

Sunnichan, V. G., H. Y. M. Ram, and K. R. Shivanna. 2005. Reproductive biology of Boswellia serrata, the source of salai guggul, an important gum-resin. *Botanical Journal of the Linnean Society* 147:73-82.

Traub, H. P., and C. T. O'Rork. 1939. Course of pollen tube growth in *Carica papaya* and *Cucurbita spp*. *Nature* 143:562-562.

Travers, S. E. 1999. Environmental effects on components of pollen performance in *Faramea occidentalis* (L.) A. Rich.(Rubiaceae) 1. *Biotropica* 31:159-166.

Tucker, S. C. 1996. Trends in evolution of floral ontogeny in *Cassia sensu stricto*, *Senna*, and *Chamaecrista* (Leguminosae: Caesalpinioideae: Cassieae: Cassiinae); a study in convergence. *American Journal Of Botany* 83:687-711.

Valtueña, F. J., T. Rodríguez-Riaño, F. Espinosa, and A. Ortega-Olivencia. 2010. Self-sterility in two *Cytisus* species (Leguminosae, Papilionoideae) due to early-acting inbreeding depression. *American Journal Of Botany* 97:123-135.

van Ryn, D., J. Lassoie, and J. Jacobson. 1988. Effects of acid mist on in vivo pollen tube growth in red maple. *Canadian Journal of Forest Research* 18:1049-1052.

Vesprini, J. L., and E. Pacini. 2000. Breeding systems in two species of the genus *Helleborus* (Ranunculaceae). *Plant Biosystems* 134:193-197.

von Haselberg, C., P. Ludders, and R. Stosser. 2004. Pollen tube growth, fertilization and ovule longevity in the carob tree (*Ceratonia siliqua* L.). Angew. Bot 78:32-40.

Wainwright, C. M. 1978. Floral biology and pollination ecology of two desert Lupines. *Bulletin of the Torrey Botanical Club* 105:24-38.

Williams, J. H. 2012. Pollen tube growth rates and the diversification of flowering plant reproductive cycles. *International Journal of Plant Sciences* 173:649-661.

Wolf, P. G., D. R. Campbell, N. M. Waser, S. D. Sipes, T. R. Toler, and J. K. Archibald. 2001. Tests of pre- and postpollination barriers to hybridization between sympatric species of Ipomopsis (Polemoniaceae). *American Journal Of Botany* 88:213-219.

Wu, X., A. Li, and D. Zhang. 2010. Cryptic self-incompatibility and distyly in *Hedyotis acutangula* Champ.(Rubiaceae). *Journal of Plant Biology* 12:484-494.

Xie, P. W., Z. L. Luo, and D. X. Zhang. 2013. Syrphid fly pollination of *Guihaiothamnus acaulis* (Rubiaceae), a species with "butterfly" flowers. *Journal of Systematics and Evolution* 51:86-93.

Zhang, X. L., R. W. Gituru, C. F. Yang, and Y. H. Guo. 2010. Exposure to water increased pollen longevity of pondweed (*Potamogeton* spp.) indicates different mechanisms ensuring pollination success of angiosperms in aquatic habitat. *Evolutionary Ecology* 24:939-953.

Zhang, X. S., and S. D. O'Neill. 1993. Ovary and gametophyte development are coordinately regulated by auxin and ethylene following pollination. *Plant Cell 5:403-418*.

Appendix S2: Pollen tube growth rate (*PTGR*) evolution across Spermatophytes. Contour plot showing reconstructed history of *PTGR*. Cool colors indicate *PTGR*s closer to the minimum value in seed plants while warm colors indicate *PTGR*s closer to the maximum value in seed plants. Scale bar indicates millions of years before present.

Figure 3.

Appendix S3: Summary statistics for pollen tube growth rate (*PTGR*) of *Magnolia grandiflora*.

Statistic	*PTGR*
raw mean (*N = 25* crosses)	827.6 μm h^{-1}
raw SD (*N = 25*)	141.3 μm h^{-1}
raw CV	0.1708
Log(10) mean	2.912 μm h^{-1}
transformed SD	0.0689 μm h^{-1}
transformed CV	0.0237

Appendix S4: Sensitivity analysis for the magnitude of log$_{10}$ *PTGR* error estimates. Values in each column represent model weights from separate analyses of angiosperm diploids ($N =$ 138) vs. polyploids ($N = 68$). Column headings indicate the coefficient of variation (CV), ranging from zero to 0.50, used to calculate estimated species-specific standard deviations around *PTGRs* in each analysis. The best-fitting model at each CV is indicated in bold. [a], Empirically-determined CV of *Magnolia grandiflora*.

	Coefficient of Variation					
Model	**0.00**	**0.0237[a]**	**0.05**	**0.10**	**0.25**	**0.50**
OUMV	**0.376**	**0.373**	**0.459**	**0.461**	0.119	0.045
OUMA	0.363	0.333	0.307	N/A	0.172	0.070
OUM	0.157	0.174	0.137	0.309	0.212	0.080
OU1	0.104	0.120	0.097	0.230	**0.390**	0.188
BM1	1.74E-31	1.77E-25	3.20E-20	4.58E-14	0.089	**0.528**
BMS	5.45E-32	5.52E-26	4.81E-24	1.30E-12	0.017	0.089

Appendix S5: *PTGR* evolution in gymnosperms vs. angiosperms. Selective regime 1 represents gymnosperms (*N* = 28) and selective regime 2 represents angiosperms (*N* = 423). Models representing <1% of the model weight are excluded.

model	ΔAICc	model weight	sigma sq1	alpha1	sigma sq2	alpha2	optimum1	se1	optimum2	se2
OUMV	(822.1)	0.338	0.091	0.122	0.139	0.122	0.187	0.123	2.690	0.048
OUM	0.040	0.332	0.137	0.124	0.137	0.124	0.188	0.150	2.690	0.047
OUMA	0.051	0.330	0.091	0.120	0.091	0.123	0.187	0.123	2.690	0.048
AVERAGED MODEL		~1	0.106	0.122	0.122	0.123	0.187	0.132	2.690	0.048

Appendix S6: C-value evolution in gymnosperms vs. angiosperms. Selective regime 1 represents gymnosperms ($N = 23$) and selective regime 2 represents angiosperms ($N = 161$). Models representing <1% of the model weight are excluded.

model	ΔAICc	model weight	sigma sq1	alpha1	sigma sq2	alpha2	optimum1	se1	optimum2	se2
OUMV	(172.4)	0.502	0.007	0.095	0.005	0.095	1.231	0.041	0.184	0.051
OUMA	0.02	0.498	0.006	0.085	0.006	0.096	1.231	0.042	0.184	0.051
AVERAGED MODEL	~1		0.007	0.09	0.005	0.095	1.231	0.041	0.184	0.051

Appendix S7: Closely-related taxon analyses.

Appendix S7a. Closely-related species pairs extracted from ploidy dataset. *PTGRs* in μm h^{-1}.
Binomial test ($P = 0.623$; $N = 10$).

DIPLOID TAXON	POLYPLOID TAXON	DIPLOID *PTGR*	POLYPLOID *PTGR*	FASTER TAXON
Anagallis arvensis	*Anagallis monelli*	233.33	105.56	diploid
Hemerocallis thunbergii	*Hemerocallis fulva*	4166.67	6266.67	polyploid
Ipomoea purpurea	*Ipomoea batatas*	7450	4625	diploid
Iris mandshurica	*Iris pseudacorus*	278.65	4255.50	polyploid
Lythrum junceum	*Lythrum salicaria*	722.22	493.60	diploid
Medicago rigidula	*Medicago sativa*	82.23	192.17	polyploid
Prunus avium	*Prunus domestica*	260.88	177.5	diploid
Tabebuia rosea	*Tabebuia chrysotricha*	1111.11	1342.45	polyploid
Trifolium pratense	*Trifolium polymorphum*	103.89	444.44	polyploid
Ulmus pumila	*Ulmus americana*	56.25	56.25	equivocal

Appendix S7b. Intraspecific diploid-polyploid cytotypes taken from the literature. All are autopolyploids. Binomial test, $N = 11$, $P = 0.0020$. Percent difference is calculated relative to the diploid.

REF.	TAXON	DIPLOID PTGR	POLYPLOID PTGR	POLYPLOID +/- (% DIFF)
1	*Beta vulgaris* 2x,4x	241.2 µm/h	142.7 µm/h	slower (-69%)
2	*Cucumis melo* 2x,4x	"no difference"		equivocal
3	*Datura stramonium* 2x,4x	2953.7 µm/h	2812.5 µm/h	slower (-4.8%)
4	*Lactuca sativa* 2x,4x	"faster"	"slower"	slower
5	*Malus domestica* 2x,4x	3.8 units/96 h	3.1 units/96 h	slower (-18.4%)
6	*Malus domestica* 2x,3x	682 µm/h	465 µm/h	slower (-31.8%)
7	*Secale cereale* 2x,4x	12.24 units/h	12.08 units/h	slower (-1.3%)
8	*Solanum sp.* 2x,4x	"faster"	"slower"	slower
9	*Trifolium pratense* 2x,4x	2322 µm/h	1950 µm/h	slower (-16%)
10,11	*Zea mays* 2x,4x	Slower pollen germination and pollen tube growth rate in 4x		slower

References

1. **Matsumura S, Mochizuki A. 1953.** Improvement of sugar beet by means of induced triploidy. *The Japanese Journal of Genetics* **28**(2): 47-56.
2. **Susin I, Álvarez JM. 1997.** Fertility and pollen tube growth in polyploid melons (*Cucumis melo* L.). *Euphytica* **93**(3): 369-373.
3. **Buchholz JT, Blakeslee AF. 1929.** Pollen-tube growth in crosses between balanced chromosomal types of *Datura stramonium*. *Genetics* **14**: 538-568.

4. **Einset J. 1944.** Cytological basis for sterility in induced autotetraploid lettuce (*Lactuca sativa* L.). *American Journal of Botany*: 336-342.

5. **Adachi Y, Komori S, Hoshikawa Y, Tanaka N, Abe K, Bessho H, Watanabe M, Suzuki A. 2009.** Characteristics of fruiting and pollen tube growth of apple autotetraploid cultivars showing self-compatibility. *Journal of the Japanese Society for Horticultural Science* **78**(4): 402-409.

6. **Modlibowska I. 1945.** *Pollen tube growth and embryo-sac development in apples and pears*. Ph.D. dissertation, University of London, London.

7. **Chin T. 1943.** Cytology of the autotetraploid rye. *Botanical Gazette* **104**(4): 627-632.

8. **Modlibowska I. 1945.** *Pollen tube growth and embryo-sac development in apples and pears*. Ph.D. dissertation, University of London, London.

9. **Evans AM. 1962.** Species hybridization in *Trifolium*. 2. Investigating pre-fertilization barriers to compatibility. *Euphytica* **11**(3): 256-262.

10. **Randolph L. 1935.** Cytogenetics of tetraploid maize. *J. agric. Res* **50**: 591-605.

11. **Green JM. 1946.** Comparative rates of pollen tube establishment in diploid and tetraploid maize. *Journal of Heredity* **37**(4): 117-121.

Appendix S8: Phylogenetic ANCOVA results. Models comprising <1% of the model weight are excluded.
Appendix S8a: Angiosperms only ($N = 100$).

		Full model		
Model	**Weight**	$P_{\text{C-value}}$	P_{ploidy}	$P_{\text{interaction}}$
kappa	0.688	0.028	0.565	0.334
OU	0.287	0.005	0.679	0.414
lambda	0.025	0.076	0.867	0.626

Model averaged slope for C-value: 0.399 ± 0.167

		No interaction	
Model	**Weight**	$P_{\text{C-value}}$	P_{ploidy}
kappa	0.652	0.046	0.748
OU	0.313	0.006	0.895
lambda	0.035	0.083	0.979

Model averaged slope for C-value: 0.344 ± 0.153

		C-value only
Model	**Weight**	$P_{\text{C-value}}$
kappa	0.642	0.045
OU	0.322	0.005
lambda	0.036	0.079

Model averaged slope for C-value: 0.344 ± 0.153

Appendix S8b: All seed plants ($N = 118$).

		Full model		
Model	**Weight**	$P_{\text{C-value}}$	P_{ploidy}	$P_{\text{interaction}}$
kappa	0.998	0.080	0.486	0.624

Slope for C-value: 0.288 ± 0.163

		No interaction	
Model	**Weight**	$P_{\text{C-value}}$	P_{ploidy}
kappa	0.997	0.092	0.562

Slope for C-value: 0.263 ± 0.155

		C-value only
Model	**Weight**	$P_{\text{C-value}}$
kappa	0.997	0.090

Slope for C-value: 0.264 ± 0.154