1   Effects of genome size on pollen performance

6   How does genome size affect the evolution of pollen tube growth rate, a haploid

7   performance trait?

John B. Reese[1,2] and Joseph H. Williams[1]

Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN

37996, U.S.A.

[1]Author for correspondence:

John B. Reese

Tel: 865 974 9371

Email: jreese11@vols.utk.edu

25   **ABSTRACT**

Premise of the Study - Male gametophytes of seed plants deliver sperm to eggs via a pollen

tube. Pollen tube growth rate (*PTGR*) may evolve rapidly due to pollen competition and haploid

selection, but many angiosperms are currently polyploid and all have polyploid histories.

Polyploidy should initially accelerate *PTGR* via "genotypic effects" of increased gene dosage

and heterozygosity on metabolic rates, but "nucleotypic effects" of genome size on cell size

should reduce *PTGR*. How are such cell-level consequences of genome size change related to the

evolution of orders-of-magnitude faster *PTGRs* in angiosperms versus gymnosperms?

Methods - We assembled a phylogenetic tree of 451 species with known *PTGRs* and added

ploidy and DNA content values from the literature. We then used comparative phylogenetic

methods to detect phylogenetic signal, differences in selective optima, and correlated evolution

of *PTGR* and genome size.

Key Results - Gymnosperms had significantly higher C-value and slower *PTGR* optima than

angiosperms. DNA content was negatively correlated with *PTGR* in gymnosperms, but non-

significant in angiosperms. Among angiosperms, model-based analyses indicated a single *PTGR*

selective optimum for diploids and polyploids. Sister-taxon and intraspecific-cytotype

comparisons indicated either no difference or slower *PTGRs* in polyploids than diploids.

Conclusions – Results from closely-related taxa indicate nucleotypic effects are initially equal to

or outweigh genotypic effects, and none of the analyses found that genome duplication

accelerated *PTGR*. Thus, our results suggest that gradual and sustained expansion of *PTGRs* in

angiosperms has mostly occurred in diploid or diploidized species, where pollen tubes are most

subject to haploid selection.

Keywords: DNA content, evolution of development, gametophyte, growth rate,

macroevolution, pollen competition, pollen tube, polyploidy, whole genome duplication.

INTRODUCTION

In seed plants, the male gametophyte is a highly-reduced haploid organism, developing within the pollen grain until pollination, when it germinates and a pollen tube emerges to invade maternal tissues. The pollen tube has been recruited for the novel function of siphonogamy, the transport of non-motile sperm cells to the egg-bearing female gametophyte, in conifers and Gnetales (two related gymnosperm groups) and in angiosperms (Friedman, 1993). In siphonogamous groups, pollen germination speed and pollen tube growth rate (*PTGR*) together determine fertilization timing. Much of the haploid genome of the male gametophyte is expressed during pollen tube development and also after fertilization, in the diploid sporophyte generation (Tanksley et al., 1981; Twell et al.*,* 2006; Otto et al. 2015). Pollen competition is thought to be much more intense in angiosperms, causing greater exposure to haploid selection and faster evolution of genes for pollen germination speed and tube growth rate (Mulcahy, 1979; Snow, 1990; Arunkumar et al. 2013; Otto et al. 2015). Indeed, angiosperms have evolved orders-of-magnitude faster pollen tube growth rates than gymnosperms (Williams 2009, 2012).

One complication with the idea that pollen performance genes have evolved rapidly under haploid selection is that whole genome duplications (WGDs) have been especially common in angiosperms relative to gymnosperms (Leitch & Leitch, 2012, 2013). WGDs elevate pollen tube gene expression above the haploid level, which has consequences for growth rates. First, WGDs mask haploid selection on pollen-expressed genes via the effects of heterosis (sheltering of deleterious alleles and/or new allelic interactions) (Lande and Schemske, 1985; Husband and Schemske, 1997; Comai, 2005; Birchler et al. 2010; Husband, 2016). Secondly, WGDs enable gene dosage effects that could increase the capacity for protein synthesis and hence increase metabolic rates (Stebbins, 1974; Comai, 2005; Conant and Wolfe, 2008). Thirdly,

large changes in DNA content (whether by WGD or other processes) are known to have

"nucleotypic" effects on nuclear size, cell size and the duration of the cell cycle, independent of

"genotypic" effects at individual loci (Bennett, 1971, 1972; Cavalier-Smith, 1978; Price 1988;

Cavalier-Smith, 2005).

We know surprisingly little about the mechanistic effects of WGDs on gametophyte

performance in general, and how they might balance each other out in any one species, or if there

are any universal effects of WGDs on pollen performance specifically. Almost all studies of the

effects of increased DNA content on cell size and growth rate have been on relatively isotropic

cells with isotropic growth (Commoner, 1964; Holm-Hansen, 1969; Price et al., 1973; Grime,

1983; Shuter et al., 1983; Beaulieu et al., 2008). In these, increases in DNA content, whether by

WGDs or other mechanisms are known to have the following effects: 1) a longer S-phase is

required to replicate more DNA (Van't Hof, 1965; Bennett, 1972), and 2) a longer period of cell

growth is needed to grow a larger cell, because more DNA requires a larger nucleus and a

proportionally larger cytoplasm (Gregory, 2001; Cavalier-Smith, 2005). As a result, in cell

populations, including multicellular tissues, the *population (tissue) growth rate* is slower because

the cell cycles of cell lineages are longer. Importantly, the *volumetric growth rate* of individual

cells is rarely of interest, and may in fact stay the same after WGD, if the longer duration of the

cell cycle is proportional to the increase in cell size.

If increases in cell cycle duration and cell size are disproportionate after an increase in

DNA content, then volumetric growth rate of an individual isotropic cell must also evolve. Most

roughly isotropic plant cells grow by diffuse growth, in which cell wall construction occurs

simultaneously across the entire expanding plasma membrane surface (Albersheim et al., 2011).

In such cells, surface area increases as the 2/3 power of volume, and consequently ever more cell

wall material must be made per unit time during growth. This means that wall and plasma

membrane production rates must either keep increasing just to maintain a constant rate of

volumetric growth or that volumetric growth rate slows down as surface area increases. Diffuse-

growing plant cells can partially reduce this problem by delaying some of their cell wall

production until after they reach mature size (Cosgrove, 2005; Albersheim et al., 2011). But a

large increase in cell size, such as after WGD, is expected to cause a disproportionate amount of

extra construction costs, making it likely that a slower cell growth (expansion) rate will be the

consequence.

The problem is quite different for anisometric, tip-growing cells, such root hairs and

pollen tubes. First, pollen tubes are terminally-differentiated cells, so their cell growth rate is

entirely restricted to G1 of the cell cycle. Secondly, as shown in Fig. 1, new cell wall production

in a pollen tube occurs only in a very small region near the elongating tube apex (Chebli et al.,

2012), and growth rate is directly determined by the rate of wall production (Rounds et al.,

2011). Thus, in a pollen tube, cell surface area and cell volume are directly proportional to each

other throughout growth, because the area of active wall synthesis remains constant during

growth (Fig. 1). Given a constant nutrient supply and temperature, *PTGR* is also constant

(Brewbaker and Majumder, 1961). Tube diameter is determined as the tube tip is formed and

after germination a tube of constant diameter is maintained by self-similar growth (Fayant et al.,

2010; Geitmann, 2011; Nezhad et al., 2013). This means the same amount of work is done at all

stages of elongation, and consequently wall production rate does not become limiting at later

stages and larger sizes as it does in isotropic, diffuse-growing cells. Furthermore, since a pollen

tube only functions during growth, rather than as a mature cell, wall production is not delayed

until after mature size is attained. In sum, the volumetric growth rate of a pollen tube directly

reflects the amount of wall material being produced per unit of time (Williams *et al.*, 2016). For

a neo-diploid pollen tube relative to its haploid ancestor, the nucleotypic effect on *PTGR* will not

depend on cell cycle duration or mature cell size per se, but only on the degree to which the

amount of wall materials has been modified to accommodate its new DNA content (Fig. 1).

How should the nucleotypic effect play out in a cell that only functions during growth? In

isotropic, diffuse-growing cells, genome size scales with cell size because the nucleo-

cytoplasmic ratio is thought to be conserved for optimal function of the mature cell (Bennett,

1972; Cavalier-Smith, 1978; Price, 1988; Cavalier-Smith, 2005; Barow, 2006). In a pollen tube

(Fig. 1), most of the functional cytoplasm, including the sperm cells, tube nucleus, and

organelles, are confined to a space of constant size between the leading edge of the enlarging

tube vacuole and the elongating tip (Hepler and Winship, 2015). Hence, a reasonable expectation

is that the volume *per unit of tube length* of a growing diploid pollen tube would be up to double

that of its progenitor haploid tube. Doubling volume per unit length results in a 41% increase in

tube diameter and circumference (Fig. 1). That value is consistent with in vivo observations of a

39% larger diameter tube in an allotetraploid *Nicotiana* relative to the mean of its presumed

diploid progenitors (Kostoff & Prokofieva, 1935), or of 8-53% larger tube diameters in tetraploid

versus diploid species of *Gossypium* (Iyengar, 1938).

Tube shank wall thickness is generally seen as being near its minimal value to function in

resisting turgor pressure and compression stress (Parre & Geitmann, 2005). Wall thickness did

not scale with tube diameter in several angiosperms (Williams et al. 2016), and turgor did not

scale with either tube diameter in fungal hyphae (Harold *et al.*, 1996) or with *PTGR* lily (Benkert

et al., 1997). Therefore, all else being equal, nucleotypic effects of genome doubling are

expected to cause up to a 41% increase in the amount of wall material needed to extend a pollen

tube tip a unit of length, which in turn is predicted to slow *PTGR* by up to 41%,.

Genome doubling can also affect *PTGR* by its genotypic effects on pollen tube

energetics, which ultimately affect the rate of wall synthesis. Pollen tube energetics are still not

well understood, and there are redundant aerobic and anaerobic respiratory systems (Rounds et

al. 2010, 2011; Colaço et al., 2012; Obermeyer et al., 2013; Selinski and Scheibe, 2014). Yet it is

well accepted that most of the metabolic energy expended by a growing pollen tube is allocated

in one way or another to the production of new wall material (Bove et al., 2008). Genome

duplication might be expected to increase wall production rates because of the effect of increased

gene dosage on metabolic processes. Although gene expression levels can become rapidly

modified during and after polyploidization due to gene loss, gene silencing, and dosage

compensation (Freeling et al., 2015; Schoenfelder and Fox, 2015; Dong et al., 2016; Panchy et

al., 2016), there is evidence that dosage effects have played an important role in the evolution of

metabolic rates. For example, in yeast cells, alcohol dehydrogenase activity per cell increased

linearly with ploidy across four ploidy levels (Dilorio *et al.*, 1987). In plant tissues,

endopolyploidy is common in cells that have secretory or transfer functions and endopolyploid

cells have higher metabolic rates (d'Amato, 1984; Galbraith *et al.*, 1991; Scholes & Paige,

2015). Finally, WGD-generated paralogs involved in metabolic pathways, such as glycolysis,

can be preferentially retained as genomes undergo gene losses after WGD (Conant and Wolfe

2007).

Another genetic consequence of WGD on *PTGR* occurs via the fitness effects of diploidy

over haploidy at the gene level. Diploidy is expected to initiate or maintain faster *PTGRs* for two

reasons. First, via the sheltering of deleterious performance alleles that were present in one or

both haploid ancestors or that arose after WGD (Husband and Schemske, 1997; Husband, 2016).

Secondly, via the effect of heterozygosity (overdominance), which is known to cause heterosis

for growth rates of sporophytes in intra- and inter-specific hybrids (Lippman and Zamir, 2007;

Feys et al., 2018). These potential effects of diploidy on *PTGR* are expected in all neo-polyploids

(Lande and Schemske, 1985), but the heterotic effects of allelic interactions and sheltering

become more evident as genetic variation in the polyploid increases. Still, in all but the special

case where there is no genetic variation, diploidization of an ancestrally haploid pollen tube

predicts genetic effects on performance speed to result in greater than or equal to ancestral

haploid speed.

The predicted effects of WGDs on pollen performance speed are summarized in Fig. 2.

Increases in DNA content, by WGD or any other process, are expected to *reduce PTGR* by

nucleotypic effects, primarily that of increased tube cell size on wall volume. In contrast,

increases in DNA content by WGD involves genome-wide gene duplication, and are expected to

increase PTGR through the genotypic effects of gene dosage and heterosis. Given that haploid

selection on pollen tubes has been seen as an important arbiter of the evolution of flowering plant

sporophytes, how robust has pollen tube growth and development been to WGDs? Is there

evidence that WGDs been involved in generating the pattern of accelerated *PTGRs* within

angiosperms?

In this study, we test the hypothesis that changes in DNA content have affected *PTGR* by

asking if there are consistent macroevolutionary patterns that might distinguish among the

alternative predictions described above. Specifically, we tested for differences in *PTGR* between

diploids and polyploids at the intraspecific, sister-taxon, and macroevolutionary scale; and for

correlations between *PTGR* and DNA content at the macroevolutionary scale. We examined the

macroevolutionary pattern at the level of all plants that have pollen tubes (seed plants), and

within the two major lineages that have different pollen tube cell biology and vastly different

levels of species diversity, gymnosperms and angiosperms.

<div align="center">**MATERIALS AND METHODS**</div>

Data collection – Data on *PTGRs* were taken from Williams (2012) and more recent literature

(cited in Supplemental Information). Consistent with other comparative analyses of

physiological traits, and with the way researchers measure *PTGR* from the longest pollen tubes,

PTGR values for each species represent an average of maximum in vivo growth rates, or if there

was more than one report for a species the average of those values (see Williams 2012 for

details). *PTGRs* were taken from within-diploid or within-polyploid (i.e., never interploidy)

crosses, in keeping with our overall goal of finding mechanisms underlying the pattern of *PTGR*

evolution in stabilized polyploids. DNA content was analyzed using C-value: the amount of

nuclear DNA in the unreplicated gametic nucleus, irrespective of ploidy level (Swift, 1950;

Bennett and Leitch, 2012). While DNA content is equivalent to genome size in diploid

organisms, it is a multiple of genome size in polyploids, thus we use the term DNA content

throughout. C-value data was collected from the Kew Royal Botanic Gardens Plant C-Value

Database (Bennett and Leitch, 2012). Ploidy level data was extracted from the same source as

PTGR data where possible, and otherwise from the Kew Plant C-Value Database, TRY

databases, or the Index to Plant Chromosome Numbers. As ploidy level is determined either by

chromosome count or segregation pattern in these data, paleopolyploids and ancient duplication

events are not included.

Tree Construction and Dating – GenBank accessions for 16 gene regions (rbcL, matK,

trnL-F, 18s_rDNA, atpB, ndhF, adh, trnL, rpl32, trnT-L, psbA-trnH, rpl32-trnL, ITS,

5.8s_rRNA, rps16, and 26s_rDNA) for the 451 seed plant species with pollen tube growth rate

data were retrieved, cleaned, and assembled into multiple gene alignments using PHLAWD and

phyutility (Smith and Donoghue, 2008; Smith and Dunn, 2008). Tree inference was performed

using maximum likelihood in RAxML version 8 (Stamatakis, 2014) on CIPRES. A pruned

version of the seed plant tree from Magallón et al. (2015) was used as a guide tree to enforce

topology of major clades. The resulting maximum likelihood estimate of the tree was rooted and

ultrametricized using the *ape* (Paradis et al., 2004) and *geiger* packages in R (Harmon et al.,

2008). Time-calibration was performed using the Congruification method (Eastman et al., 2013).

219   ***Character scoring***- The *PTGR* value used for each species represents an estimate of

maximum sustained growth rate. Since each *PTGR* value represents a species mean obtained

from multiple measurements, we attempted to incorporate error into phylogenetic comparative

analyses. *OUwie* allows for incorporating estimates of standard error (*SE*), and since species

means were log-transformed for analysis, log-transformed *SE*s are required. As there is no

reliable way to calculate the log-transformed *SE* from the literature without the original data for

each species, we used the following conservative approximation. First, we assumed all species

had similar SEs in *PTGR*, and we applied an empirically-determined *SE* from an exemplar

species to all. *Magnolia grandiflora* has an average *PTGR* of 828 (±141) μm h^{-1} (*N* = 25

outcrosses), close to the angiosperm median of 587 μm h^{-1} (Williams, 2012 and this study)

(Table S1). The standard deviation (*SD*) of log-transformed data was calculated and divided by

the mean of the log-transformed data to acquire a coefficient of variation (*CV*) of 0.0237. We

then multiplied the log-transformed mean *PTGR* of each species by 0.0237 to provide an

estimate of taxon-specific standard deviation. The standard deviation was used as a conservative

estimate of error because sample sizes were generally not available for calculating *SE*. We also

performed a sensitivity analysis by running the *OUwie* analyses with hypothetical error values

obtained set as 0, 0.05, 0.1, 0.25, and 0.5 (Table S2). Unless otherwise noted, all measures of

uncertainty around parameter estimates are in standard error.

We analyzed two different sets of ploidy data. In the first, 'restricted dataset' ($N = 273$

taxa), we recognized only ploidy levels as stated in the literature. In the second 'expanded

dataset' ($N = 335$ taxa), we inferred ploidy levels of 62 additional species by comparing their

chromosome count to their generic base count as reported in Wood et al. (2009). Taxa with a

chromosome number equal to or greater than twice the generic base count were considered

polyploid.

Phylogenetic Comparative Analyses - Phylogenetic signal was assessed for both *PTGR*

and gametophytic DNA content using Blomberg's K in the *picante* package in R (Blomberg et

al., 2003; Kembel et al., 2010). C-values and *PTGRs* were $\log_{10}$ transformed in this and all other

phylogentic comparative analyses below. To visualize changes in DNA content and *PTGR* along

tree branches and to generate estimates of node states, ancestral state reconstructions were

performed and plotted using the contMap function in *phytools* (Felsenstein, 1985; Revell, 2012).

Given many known biological differences between gymnosperms and angiosperms for both

pollen tube growth (Friedman, 1993; Williams, 2008) and DNA content /polyploidy (see

Discussion) (Ohri and Khoshoo, 1986; Leitch et al., 1998), all analyses were performed on

gymnosperms only, angiosperms only, and the full dataset (all spermatophytes). Whether *PTGR*

evolves under different selective regimes between the two seed plant groups was also evaluated

using the *OUwie* function (*OUwie* package) in R (Beaulieu and O'Meara, 2014) to compare

different models of evolution (detailed below). In multiple-regime models, the regimes were

either gymnosperms (1) or angiosperms (2). The same analysis was used to test for differences in

DNA content evolution between gymnosperms and angiosperms as well.

To detect whether pollen tube growth rates of polyploid species evolve under a different

selective regime than diploid species, *OUwie* was used to compare single-regime and two-regime

models. The following models were tested: single-rate Brownian motion (BM1), multi-rate

Brownian Motion (BMS), single-regime Ornstein-Uhlenbeck (OU1), multi-regime OU with

single *alpha* and *sigma^2* estimates (OUM), multi-regime OU with single *alpha* and multiple

sigma^2 (OUMV), and multi-regime OU with multiple *alpha* and single *sigma^2* (OUMA).

The association between polyploidy and *PTGR* was also assessed among 20 diploid-

polyploid sister taxa on the tree (within-genus or within-family level). Only polyploid taxa with a

single diploid sister were used. The *PTGRs* of 11 intraspecific diploid-polyploid pairs from the

literature were also compared. A two-tailed binomial (sign) test was used to test significance.

The relationship between the continuous traits, pollen tube growth rate and gametophytic

DNA content, was assessed with phylogenetic generalized least squares (*PGLS*) regression using

the *phylolm* package in R. Gametophytic DNA content was used as the predictor variable and

PTGR the response variable. BM (Grafen, 1989) and OU (Martins and Hansen, 1997) models

were both used, in addition to Pagel's lambda, kappa, and delta models (Pagel, 1997, 1999).

To explore the level of convergent evolution within each trait and determine coincidence

of selective regime shifts in *PTGR* and DNA content, the *SURFACE* package in R was used

(Ingram and Mahler, 2013). *SURFACE* adds selective regime shifts to a single regime model in a

stepwise fashion until adding another regime shift decreases the model likelihood. Regimes are

then collapsed in a pairwise fashion until further collapses decrease the likelihood. Nodes that

included a regime shift in both traits were identified.

RESULTS

PTGR evolution and C-value evolution in angiosperms versus gymnosperms - Blomberg's K

indicated significant phylogenetic signal across seed plants and within angiosperms, for both the

PTGR and C-value datasets (Table 1). *OUwie* analyses of seed plants rejected both the Brownian

Motion (BM) and Ornstein-Uhlenbeck (OU) single-optimum models for the evolution of *PTGR*

and C-value. OU models that allowed separate selective regimes for angiosperms and

gymnosperms accounted for > 99.9 % of the model weight in both *PTGR* and C-value (Tables

S3, S4). Log_{10} *PTGR* selective optima were more than a magnitude of order higher in

angiosperms (2.69 ± 0.048 μm h^{-1}) than in gymnosperms (0.187 ± 0.123 μm h^{-1}). Log_{10} DNA

content selective optima were more than a magnitude of order smaller in angiosperms (0.184 ±

0.051 pg) than in gymnosperms (1.231 ± 0.041 pg). The maximum likelihood (ML)

reconstructions of angiosperm and gymnosperm common ancestors (CAs) showed a similar

pattern. Ancestral Log_{10} *PTGR* for angiosperms was higher than that of gymnosperms, 2.44 μm

h^{-1} (95% CI: 1.09-3.69) versus 0.215 μm h^{-1} (95% CI: -1.48-1.92), respectively (Fig. 3), and

ancestral Log_{10} DNA content was 0.29 pg (95% CI: -0.45-1.04) for angiosperms and 1.10 pg

(95% CI: -0.28-2.47) for gymnosperms.

  Joint evolution of PTGR and ploidy - In the restricted ploidy dataset, the best fitting

OUwie model for angiosperms using the *Magnolia* error estimate of 0.024 was the single-rate

OU model which had 50.0% of the model weight (Table 2a); whereas the three two-regime OU

models accounted for the remaining 50.0% of the model weight (both BM models were rejected).

This pattern held up for a wide range of error values in the sensitivity analysis (Table S2), except

that when error reached 0.5, a value > 20 times that seen in *Magnolia*. In the expanded ploidy

dataset (Table 2b), the weight of the single-regime OU model was 57.2% compared to a

combined 42.8% for the three two-regime models. However, the estimates for the polyploid

selective optima are different from those of the restricted dataset; diploid optima in both datasets

were 2.7 ± 0.07 $\log_{10}$ μm h^{-1}, whereas polyploid optima were 1.4 ± 1.5 $\log_{10}$ μm h^{-1} in the

restricted dataset versus 2.8 ± 1.00 l og$_{10}$ μm h^{-1} in the expanded dataset.

A survey of intraspecific cytotypes found autopolyploids had slower *PTGR* than diploids

in 9 of 11 pairs and no difference in the remaining two (Binomial test, $P = 0.002$; Table S4b). In

the within-genus sister-taxon comparison, polyploids had slower *PTGR* than diploids in 9 pairs,

faster *PTGR* in 2, and no difference in two (Two-tailed binomial test, $P = 0.065$). If expanded to

within-family sisters, there were twelve, six, and two pairs, respectively ($P = 0.238$; Table S4a).

Joint evolution of PTGR and DNA content - For seed plants, simple linear regression

(*SLR*) showed a significant negative correlation between DNA content and *PTGR* ($P<0.0001$),

but that result was clearly driven by the large DNA contents and slow *PTGRs* of gymnosperms

relative to angiosperms (Fig. 5), because the *PGLS* regression was non-significant ($P=0.463$;

Table 3). Taking these two clades separately, DNA content was negatively correlated with *PTGR*

in gymnosperms in the *PGLS* regression (*SLR*: $P = 0.200$; *PGLS*: $P \leq 0.02$, OU model; Table 3),

but not *SLR* (phylogenetic signal was not significant for either trait in gymnosperms; Table 1).

For angiosperms, it was positively correlated using *SLR* ($P=0.0005$), but non-significant using

PGLS ($P = 0.284$; Table 3). In angiosperms, the *PGLS* regression was non-significant when

polyploids were excluded ($P = 0.457$, kappa model).

Detecting coincident regime shifts in PTGR and DNA content within seed plants - The

SURFACE analysis of *PTGRs* found 13 distinct selective regimes ($N = 451$ taxon tree), with 51

selective regime shifts (22 to faster and 29 to slower regimes). For C-value, there were 9 distinct

selective regimes ($N = 184$ taxon tree), with 4 shifts to larger and 7 shifts to smaller regimes.

Regime shifts in *PTGR* and DNA content were coincident at only two nodes, a *PTGR*

acceleration and genome downsizing in the CA of extant angiosperms and a *PTGR* slowdown

and genome size decrease in the CA of rosids and (Fig. 6). Excluding terminal, branches, all

other regime shifts were at least 3 nodes away from each other. *PTGR* shifts on the smaller tree

(Fig. 6) were in the same locations as on the larger *PTGR* tree (not shown).

332   **DISCUSSION**

DNA content variation and the evolution of PTGR in angiosperms versus gymnosperms - We

found that *PTGR* has evolved around a significantly faster selective optimum in angiosperms

than in gymnosperms. Though it has long been known that most angiosperm *PTGRs* far exceed

those of gymnosperms, this result provides quantitative support to the near-universal assumption

that shifts to faster angiosperm-like *PTGRs* began *prior* to the origin of extant angiosperms.

However, the slow angiosperm-wide selective optimum and reconstructed ancestral *PTGRs* (both

< 490 μm/h), are consistent with gradual evolution of faster *PTGRs* in angiosperms.

340   There are several hypotheses for how and why angiosperms evolved such fast *PTGRs*.

First, Mulcahy (1979) invoked a shift to much higher intensity of pollen competition in

angiosperms as a driver of the origin and continued evolution of faster growth rates. Notably, no

other type of tip-growing cell (haploid or diploid) in land plants has evolved comparably fast tip-

growth rates and none of those cell types, including gymnosperm pollen tubes, experience

intense competition for resources (Williams et al., 2016). Secondly, rapid *PTGR* may have been

advantageous as angiosperms transitioned to a much faster reproductive cycle (Stebbins, 1974;

Williams, 2012; Williams and Reese, *in review*). Thirdly, biophysical or physiological attributes

of pollen tubes may have either constrained *PTGRs* of gymnosperms to be slow or enabled faster

PTGRs in angiosperms, or both (Hoekstra, 1983; Derksen et al., 1999; Fernando et al., 2005;

Williams, 2008, 2009). A fourth possibility, tested here, is that strong differences in genome-

level processes between angiosperms and gymnosperms have impacted the evolution of

angiosperm *PTGRs* relative to their living and extinct seed plant relatives.

We found that DNA content has evolved around a significantly lower selective optimum

in angiosperms than in gymnosperms, even though angiosperms have a broad range of DNA C-

values that encompass the entire range of seed plant genome sizes (Fig. 5; see Leitch and Leitch,

2013 for a larger survey). Paradoxically, angiosperms have great variation in ploidy level, a

history of speciation by polyploidy, and much evidence of past genome duplication in both

diploids and polyploids, including a WGD in a common ancestor of extant angiosperms (Wood

et al., 2009; Husband et al., 2013; Van de Peer et al*.,* 2017; Landis et al*.,* 2018). Evidence for

more recent WGDs is found in most modern diploids, which is consistent with studies of

experimental WGDs that show that genome duplication is commonly accompanied by rapid loss

of DNA sequences, gene fractionation by large-scale deletions, biased retention of genes with

beneficial dosage effects, and ultimately a return to the diploid state (Conant and Wolfe, 2008;

Conant et al., 2014; Freeling et al., 2015; Dodsworth et al*.,* 2016; Wendel et al., 2018).

In contrast, WGDs have been very rare in gymnosperms (Leitch et al., 2005; Wood et al.,

2009; Soltis et al., 2009; Husband et al., 2013; Leitch and Leitch, 2013), and their high DNA

contents are thought to be due mainly to high transposon activity without repeated rounds of

genome duplication (Leitch & Leitch, 2013; Lee and Kim, 2014). Hence, gymnosperms have

experienced the effects of higher DNA content on pollen tube dimensions, which is predicted to

reduce *PTGR,* without the potential counter-balancing effects of gene dosage and heterosis

incurred by repeated rounds of genome duplications (Fig. 2), gene sorting, and diploidization.

372   Angiosperm pollen (Hoekstra, 1983) and sporophytes (Bond, 1989; Feild and Arens,

2005; Beaulieu et al., 2007) are known to have evolved much higher respiration rates than the

pollen and sporophytes of gymnosperms. Respiration rates of angiosperm pollen are ~10 times

higher than those of sporophytic tissues (Tadege and Kuhlemeier, 1997), and other energetic

pathways also function (Rounds et al., 2011; Obermeyer et al., 2013). A major consequence of

WGDs over single-gene duplication events is that WGDs can increase gene dosage of whole

metabolic pathways at once (Conant and Wolfe, 2007). When particular metabolic pathways are

limiting to growth rate, duplication can upregulate the pathway via gene dosage effects, and

positive selection on dosage can shelter gene duplicates, allowing time for sub- or neo-

functionalization, and hence much faster evolution of those genes (Kondrashov and Kondrashov,

2006; Conant and Wolfe, 2007). Pollen tube growth rate genes are widespread in the genome, for

example, occurring on every chromosome arm of corn (Sari-Gorla et al., 1992). Thus, *PTGR* is a

trait governed by many interacting genes that are widespread in the genome, which is exactly the

kind of trait in which WGDs are favored over individual gene duplications or segmental

duplications of chromosomes, due to the need to conserve gene dosage balance (Birchler et al.,

2012; Conant et al., 2014; Freeling et al., 2015).

Growth rate evolution in sporophytic systems is consequent on growth of tissues,

populations of cells, and rates and durations of many individual cell cycles. In contrast, *PTGR*

genes with gametophytic expression govern a very restricted aspect of individual cell growth –

mostly the intracellular transport system and cell wall synthesis machinery operating within a

single, terminal cell cycle stage (G_1, or G_0) (Russell et al., 2012; Geitmann and Nebenführ,

2015). This means that the evolution of faster *PTGR* in angiosperms is likely to have imposed

selection on a restricted set of gametophytically-expressed genes in metabolic and intracellular

transport pathways (Arunkumar et al., 2015; Gossmann et al., 2016). Genes within such

biochemical pathways might be preferentially retained after WGD due to the need to maintain

balanced gene dosage effects (Birchler and Veita, 2012; Conant et al., 2014). Starting with the

origin of flowering plants, WGDs have provided repeated opportunities for genome-wide, gene-

level effects (gene sorting via positive and purifying selection, sheltering of deleterious alleles,

and heterosis) to affect *PTGR* evolution. At the same time, given the small genome sizes of so

many plants with a history of WGDs, the nucleotypic effect of doubled DNA content on cell size

must often have been tempered by rapid genome downsizing following WGD (Freeling et al.,

2015; Dodsworth et al., 2016; Wendel et al., 2018).

In summary, increases in DNA content by any mechanism are expected to initially

decrease *PTGR* via the nucleotypic effects of larger genome size on cell size and growth,

whereas increases in DNA content by WGD are expected to initially increase *PTGR* via the

genotypic effects of increased gene/allele numbers (Fig. 2). *PTGR* evolution of gymnosperms

may have been relatively stagnant for many reasons, such as lack of pollen competition, relaxed

selection on *PTGR* due to slow reproductive cycles, and developmental constraints on their

pollen tube structure and physiology. However, it may also be that their high degree of

heterochromatin and rarity of WGDs have given them all the limitations of high DNA content

but none of the evolutionary opportunities provided by large-scale redundancy due to WGD and

fractionation cycles. Angiosperm *PTGRs* have evolved via repeated cycles WGDs, followed by

fractionation, gene retentions due to advantageous dosage effects, sub- or neo-functionalization

of paralogs, and eventually diploidization. Consequently, growth rate genes have alternately

experienced periods of relaxed selection due to WGD and haploid selection on performance. Our

findings that angiosperms have evolved around a higher optimum *PTGR* but a lower optimum C-

value than gymnosperms, no coincident regime shifts to higher *PTGR* and higher C-value, and

the lack of evidence for faster *PTGRs* in polyploids than diploids, suggests that the rapid *PTGRs*

of angiosperms have largely evolved at the haploid level, in diploid or diploidized species.

Polyploidy and PTGR evolution within angiosperms - Diploid and polyploid

angiosperms evolved under a single selective regime, irrespective of how ploidy level was

scored. However, in both analyses the three two-regime OU models also accounted for

substantial model weight (43% and 50%), and in these, the polyploid *PTGR* optimum was

similar or slightly slower than the diploid optimum. Thus, we find no evidence that WGDs result

in higher *PTGRs*. Two caveats are first, only 13-14% of the taxa in our two analyses were scored

as polyploids versus 25-35% estimated for angiosperms (Wood et al*.,* 2009; Landis et al., 2018);

and secondly, on our tree polyploid taxa were generally recently-derived with relatively short

branches, comprising a very small proportion of the total branch length of the tree. Hence, we

may have had lower power to estimate parameters for polyploids relative to diploids. Still, the

sister-taxon comparisons avoided these weaknesses and were consistent with the model-based

results – polyploids had similar or slightly slower *PTGR* than diploids in most sister taxa on our

tree.

Neo-polyploid *PTGR* is affected by the balance between genotypic and nucleotypic

effects (Fig. 2). Since polyploid genetic variation can vary but affects only the potential for

genotypic effects, our species might not have been a random sample of polyploid variation.

Mating systems, modes of polyploid origins, and patterns of chromosomal segregation have

effects on the degree of genetic variation and the nature of recombination in neo-polyploids. Yet,

among 16 species scored as polyploid in both our analyses, eight were fully outcrossing, seven

were self-compatible (two autogamous, two mixed mating, and four unknown), and 1 was

apomictic – a not unusual distribution (Goodwillie et al., 2005; Gibbs, 2014; Ashman et al.,

2014). Whitney et al. (2010) found only a weak correlation between DNA content and mating

system, and this may reflect the fact that genetic variation in polyploids cannot be predicted

easily. For example, autotetraploids originate with a subset of the genetic variation in the diploid

progenitor population but they often outcross and hybridize, whereas allopolyploids can be

highly heterozygous when they originate, but often are highly selfing (Stebbins, 1974; Soltis and

Soltis, 1999; Whitney et al., 2010). In the special case in which there is no genetic variation in a

diploid progenitor and its neo-autopolyploid descendant, *PTGR* would be solely determined by

the balance between nucleotypic effects and gene dosage effects (which also depend on the

degree of dosage compensation; Guo et al., 1996) (Fig. 2).

The closest approximation of the initial effect of genome duplication on *PTGR,*

independent of levels of genetic variation, is the comparison of diploids with their intraspecific,

autopolyploid cytotypes. In all 11 pairs, *PTGRs* of autopolyploid cytotypes were slower than or

equal to those of their intraspecific diploid progenitors. We should re-emphasize that all studies

involved in vivo diploid crosses (1x pollen on 2x pistils) compared to tetraploid crosses (2x

pollen on 4x pistils). The lack of any examples of faster *PTGR* in neo-autotetraploid cytotypes

than in their diploid progenitors (in which the effects of heterosis are minimized) seems to

suggest that increased gene dosage generally cannot fully offset nucleotypic effects, causing

slower or at best similar *PTGR* upon autotetraploid formation.

Disentangling nucleotypic from genotypic effects on* PTGR *evolution – Genome size is

positively correlated with cell size and often negatively correlated with organismal growth rate

across all kingdoms of life (Cavalier-Smith, 1978; Gregory, 2001). One major proposed

explanation for the correlation between DNA content (which strongly influences nuclear size)

and cell size in isotropic cells is that there is an optimal ratio of nuclear to cytoplasmic volume

for metabolic efficiency (nucleoskeletal theory) (Cavalier-Smith, 1978, 2005). "Growth rate" in

most studies refers to the doubling time of lineages of cells, and hence to the average duration of

cell cycles, not to individual cell growth rates per se. Cell cycles are longer after genome

duplication due to a longer DNA replication phase and also to the longer growth period needed

to reach a larger size. Thus, the speed of cell enlargement in such studies is conflated with the

duration of S-phase. This is not the case for a pollen tube, which is a terminally-differentiated

cell that only grows and functions in the G_1-phase of its cell cycle. Therefore, in pollen tubes,

unlike in diffuse-growing cells of multicellular tissues, volumetric cell growth rate is the primary

target of natural or sexual selection.

Large-scale genome size increases, whether by WGD or any other method, are expected

to increase cell size to maintain the nucleo-cytoplasmic ratio. *PTGR* is directly determined by the

amount and rate of cell wall production, which depends on tube size (Winship et al. 2010;

Williams et al. 2016). A neo-diploid pollen tube with doubled tube volume must produce 41%

more cell wall material per unit time to maintain its ancestral haploid *PTGR* (Fig. 1). Size

increases of that order have been observed in diploid over haploid pollen tubes (Kostoff &

Prokofieva, 1935; Iyengar, 1938). Hence, without increases in metabolic rates that could

compensate for that extra work, *PTGR* and genome size are expected to be negatively correlated.

Importantly, we found such a negative correlation in gymnosperms. Nucleotypic effects

should outweigh genotypic effects in gymnosperms, since the group has experienced large

genome expansions without widespread gene duplication by WGD after their origin (Leitch &

Leitch, 2013; Lee and Kim, 2014). Similar support for the existence of nucleotypic effects is that

PTGR was generally slower in the youngest angiosperm polyploids (eg. intraspecific cytotypes

or within-genus sisters). Over time, nucleotypic effects remain more or less constant in

polyploids, but positive genotypic effects can accumulate via all the usual population genetic

processes, counterbalancing or outweighing nucleotypic effects. Hence, the lack of correlation

between DNA content and *PTGR* in angiosperms as a whole should be interpreted as a

consequence of variation in the degree of compensating genotypic effects on *PTGR,* due to time

since origin and genetic variation.

A weakness of comparative analyses of the question of DNA content and pollen

performance is that all angiosperms, including "diploids" as determined by diploid segregation

of chromosomes or by diploid base chromosome number, have at least one WGD in their

ancestry (Landis, 2018, and reference therein). Thus, the most relevant measures of ploidy level

are probably cryptic. For example, is pollen tube gene expression in a particular taxon haploid or

polyploid? Perhaps a more relevant measure for comparisons is the number of WGDs among

different lineages, traced back from each tip. Such data are already becoming available as more

genomic data is accumulated.

Conclusions - Studies across the tree of life have consistently shown that ploidy level and

DNA content are correlated with cell size and metabolic rate. Pollen tube dimensions and

energetics affect the amount of cell wall material produced per unit of growth and the rate at

which cell wall is produced, which together determine *PTGR.* In gymnosperms, *PTGR* was

negatively correlated with genome size, but in angiosperms, where the effects of WGDs are more

prevalent, *PTGR* seems to be somewhat robust to genome duplication. Neo-autopolyploids

inherited similar or slower *PTGRs* than their diploid ancestors, and polyploids compared to their

diploid near-relatives seem to follow the same pattern. With changes in genome size, nucleotypic

effects act as a brake on growth rate and are always present, but the degree to which genotypic

effects counterbalance these depends on the historical nature and time since genome size

increase in any particular lineage. Understanding causal relationships between genome size,

ploidy and *PTGR* will involve mechanistic studies of tube cell dimensions and wall synthesis

rates in haploid and polyploid gametophytes. On the other hand, there appears to be great

variation in the tug of war between genotypic and nucleotypic effects, and there are likely to be

deeper evolutionary patterns underlying that variation.

ACKNOWLEDGEMENTS

We thank B. O'Meara and J. Beaulieu for advice on phylogenetic analyses, I. Leitch for data on

DNA content, and J. Edwards and M. Rankin for assistance in the lab. Several anonymous

reviewers also provided useful advice. Partial support to J.B.R. was provided by National

Science Foundation award IOS 1052291 to J.H.W.

Authors Contributions: J.B.R. and J.H.W. jointly conceived of the study and wrote the paper;

J.B.R. collected data on genome sizes and ploidy levels, constructed the phylogenetic tree and

performed all comparative analyses; J.H.W. collected data on *PTGRs* and diploid-autopolyploid

PTGRs.

LITERATURE CITED

Albersheim, P., A. Darvill, K. Roberts, R. Sederoff, and A. Staehelin. 2011. Plant Cell Walls:

from Chemistry to Biology. Garland Science Google Scholar, New York, New York, USA.

d'Amato, F. 1984. Role of polyploidy in reproductive organs and tissues. *In* B. M. Johri [ed.]

Embryology of Angiosperm*, 519–566. Springer-Verlag, Berlin.

Arunkumar, R., E. B. Josephs, R. J. Williamson, and S. I. Wright. 2013. Pollen-specific, but not

sperm-specific, genes show stronger purifying selection and higher rates of positive selection

than sporophytic genes in *Capsella grandiflora. Molecular Biology and Evolution* 30: 2475–

2486.

Ashman, T-L., D. Bachtrog, H. Blackmon, E. E. Goldberg, M. W. Hahn, M. Kirkpatrick, J.

Kitano, J. E. Mank, et al. 2014. Tree of Sex: A database of sexual systems. *Scientific Data* 1:

140015.

Barow, M. 2006. Endopolyploidy in seed plants*. Bioessays* 28:271-281.
543
Beaulieu, J. M., I. J. Leitch, and C. A. Knight. 2007. Genome size evolution in relation to leaf

strategy and metabolic rates revisited. *Annals of Botany*, 99: 495-505.

Beaulieu, J. M., I. J. Leitch, S. Patel, A. Pendharkar, and C. A. Knight. 2008. Genome size is a

strong predictor of cell size and stomatal density in angiosperms. *New Phytologist* 179: 975–986.

Beaulieu, J. M., and B. O'Meara. 2014. *OUwie*: analysis of evolutionary rates in an OU

framework. *R package version* 1.

Benkert, R., G. Obermeyer, and F-W. Bentrup. 1997. The turgor pressure of growing lily pollen

tubes. *Protoplasma* 198: 1–8.

Bennett, M. D. 1971. The duration of meiosis. *Proceedings of the Royal Society of London B:*

Biological Sciences 178: 277–299.

Bennett, M. D. 1972. Nuclear DNA content and minimum generation time in herbaceous plants.

Proceedings of the Royal Society of London B: Biological Sciences 181: 109–135.

Bennett, M. D. 1987. Variation in genomic form in plants and its ecological implications. *New*

Phytologist 106: 177–200.

Bennett, M. D. 2004. Perspectives on polyploidy in plants–ancient and neo. *Biological Journal*

of the Linnean Society 82: 411–423.

Bennett, M. D., and I. J. Leitch. 2012. Plant DNA C-values Database (Release 6.0).

Birchler, J. A., H. Yao, S. Chudalayandi, D. Vaiman, and R. A. Veitia. 2010. Heterosis. *Plant*

Cell: 110.076133.

Birchler, J. A., and R. A. Veitia. 2012. Gene balance hypothesis: connecting issues of dosage

sensitivity across biological disciplines. *Proceedings of the National Academy of Sciences*:

201207726.

Blomberg, S. P., T. Garland Jr., and A. R. Ives. 2003. Testing for phylogenetic signal in

comparative data: behavioral traits are more labile. *Evolution* 57: 717–745.

Bond, W. J. 1989. The tortoise and the hare: ecology of angiosperm dominance and gymnosperm

persistence. *Biological Journal of the Linnean Society* 36: 227–249.

Bove, J., B. Vaillancourt, J. Kroeger, P. K. Hepler, P. W. Wiseman, and A. Geitmann. 2008.

Magnitude and direction of vesicle dynamics in growing pollen tubes using spatiotemporal

image correlation spectroscopy and fluorescence recovery after photobleaching. *Plant*

Physiology 147: 1646–1658.

Bretagnolle, F, and J. D. Thompson. 1995. Gametes with the somatic chromosome number:

mechanisms of their formation and role in the evolution of autopolyploid plants. *New Phytologist*

129: 1–22.

Brewbaker, J. L. 1967. The distribution and phylogenetic significance of binucleate and

trinucleate pollen grains in the angiosperms. *American Journal of Botany* 54: 1069–1083.

Brewbaker, J. L, and S. K. Majumder. 1961. Cultural studies of the pollen population effect and

the self-incompatibility inhibition. *American Journal of Botany* 48: 457–464.

Cavalier-Smith, T. 1978. Nuclear volume control by nucleoskeletal DNA, selection for cell

volume and cell growth rate, and the solution of the DNA C-value paradox. *Journal of Cell*

Science 34: 247–278.

Cavalier-Smith, T. 2005. Economy, speed and size matter: evolutionary forces driving nuclear

genome miniaturization and expansion. *Annals of Botany* 95: 147–175.

Chebli, Y., M. Kaneda, R. Zerzour, and A. Geitmann. 2012. The cell wall of the *Arabidopsis*

thaliana pollen tube-spatial distribution, recycling and network formation of polysaccharides.

Plant Physiology: pp–112.

Colaço, R, N. Moreno, and J.A. Feijó. 2012. On the fast lane: mitochondria structure, dynamics

and function in growing pollen tubes. *Journal of Microscopy* 247: 106–118.

Comai, L. 2005. The advantages and disadvantages of being polyploid. *Nature Reviews Genetics*

6: 836.

Commoner, B. 1964. Roles of Deoxyribonucleic Acid in Inheritance. *Nature* 202: 960-.

Conant, G. C., J. A. Birchler, and J. C. Pires. 2014. Dosage, duplication, and diploidization:

clarifying the interplay of multiple models for duplicate gene evolution over time. *Current*

Opinion in Plant Biology 19: 91–98.

Conant, G. C., and K. H. Wolfe. 2007. Increased glycolytic flux as an outcome of whole-genome

duplication in yeast. *Molecular systems biology* 3: 129.

Conant, G. C., and K. H. Wolfe. 2008. Turning a hobby into a job: how duplicated genes find

new functions. *Nature Reviews Genetics* 9: 938.

Cosgrove, D. J. 2005. Growth of the plant cell wall. *Nature Reviews Molecular Cell Biology* 6:

850.

Derksen, J., Y. Li, B. Knuiman, and H. Geurts. 1999. The wall of *Pinus sylvestris* L. pollen

tubes. *Protoplasma* 208: 26–36.

Dilorio, A. A., P. J. Weathers, and D. A. Campbele. 1987. Comparative enzyme and ethanol

production in an isogenic yeast ploidy series. *Current Genetics* 12: 9–14.

Dodsworth, S., M. W. Chase, and A. R. Leitch. 2016. Is post-polyploidization diploidization the

key to the evolutionary success of angiosperms? *Botanical Journal of the Linnean Society* 180:

1–5.

Dong, Y., M. Deng, Z. Zhao, and G. Fan. 2016. Quantitative proteomic and transcriptomic study

on autotetraploid *Paulownia* and its diploid parent reveal key metabolic processes associated

with *Paulownia* autotetraploidization. *Frontiers in Plant Science* 7: 892.

Eastman, J. M., L. J. Harmon, and D. C. Tank. 2013. Congruification: support for time scaling

large phylogenetic trees. *Methods in Ecology and Evolution* 4: 688–691.

Fayant, P., O. Girlanda, Y. Chebli, C-É. Aubin, I. Villemure, and A. Geitmann. 2010. Finite

element model of polar growth in pollen tubes. *The Plant Cell*: tpc–110.

Field, T. S., and N. C. Arens. 2005. Form, function and environments of the early angiosperms:

merging extant phylogeny and ecophysiology with fossils. *New Phytologist* 166: 383–408.

Felsenstein, J. 1985. Phylogenies and the comparative method. *American Naturalist* 125: 1–15.

Fernando, D. D., M. D. Lazzaro, and J. N. Owens. 2005. Growth and development of conifer

pollen tubes. *Sexual Plant Reproduction* 18: 149–162.

Fey, K., K. Demuynck, J. De Block, A. Bisht, A. De Vliegher, D. Inzé, and H. Nelissen. 2018.

Growth rate rather than growth duration drives growth heterosis in maize B104 hybrids. *Plant,*

Cell & Environment 41: 374–382.

Freeling, M., M. J. Scanlon, and J. E. Fowler. 2015. Fractionation and subfunctionalization

following genome duplications: mechanisms that drive gene content and their consequences.

Current Opinion in Genetics & Development 35: 110–118.

Friedman, W. E. 1993. The evolutionary history of the seed plant male gametophyte. *Trends in*

Ecology & Evolution 8: 15–21.

Galbraith, D. W., K. R. Harkins, and S. Knapp. 1991. Systemic endopolyploidy in *Arabidopsis*
thaliana. *Plant Physiology* 96: 985–989.

Geitmann, A. 2011. Generating a cellular protuberance: mechanics of tip growth. *In* P.
Wojtaszek [ed.] Mechanical integration of plant cells and plants117–132. Springer, Heidelburg.

Geitmann, A, and A. Nebenführ. 2015. Navigating the plant cell: intracellular transport logistics
in the green kingdom. *Molecular Biology of the Cell* 26: 3373–3378.

Gibbs, P. E. 2014. Late-acting self-incompatibility–the pariah breeding system in flowering
plants. *New Phytologist* 203: 717–734.

Goodwillie, C., S. Kalisz, and C. G. Eckert. 2005. The evolutionary enigma of mixed mating
systems in plants: occurrence, theoretical explanations, and empirical evidence. *Annual Review*
of Ecology Evolution and Systematics. 36: 47–79.

Gossmann, T. I., D. Saleh, M. W. Schmid, M. A. Spence, and K. J. Schmid. 2016.
Transcriptomes of plant gametophytes have a higher proportion of rapidly evolving and young
genes than sporophytes. *Molecular Biology and Evolution* 33: 1669–1678.

Grafen, A. 1989. The phylogenetic regression. *Philosophical Transactions of the Royal Society*
of London. Series B, Biological Sciences 326: 119–157.

Gregory, T. R. 2001. The bigger the C-value, the larger the cell: genome size and red blood cell
size in vertebrates. *Blood Cells, Molecules, and Diseases* 27: 830–843.

Grime, J. P. 1983. Prediction of weed and crop response to climate based upon measurements of
nuclear DNA content. *In* Aspects of Applied Biology 4. Influence of environmental factors on
herbicide performance and crop and weed biology, 87–98.

Harmon, L. J., J. T. Weir, C. D. Brock, R. E. Glor, and W. Challenger. 2008. GEIGER:
investigating evolutionary radiations. *Bioinformatics* 24: 129–131.

Harold, R. L., N. P. Money, and F. M. Harold. 1996. Growth and morphogenesis in *Saprolegnia*
ferax: Is turgor required? *Protoplasma* 191: 105–114.

Hepler, P. K., and L. J. Winship. 2015. The pollen tube clear zone: clues to the mechanism of

polarized growth. *Journal of Integrative Plant Biology* 57: 79–92.

Hoekstra, F. A. 1983. Physiological evolution in angiosperm pollen: possible role of pollen

vigour. In: Mulcahy DL, Ottaviano E, eds. Pollen: Biology and Implications for Plant Breeding,

35–41. Elsevier Science, Amsterdam.

Holm-Hansen, O. 1969. Algae: amounts of DNA and organic carbon in single cells. *Science* 163:

87–88.

Husband, B. C. 2016. Effect of inbreeding on pollen tube growth in diploid and tetraploid

Chamerion angustifolium: Do polyploids mask mutational load in pollen? *American Journal of*

Botany 103: 532–540.

Husband, B. C., S. J. Baldwin, and J. Suda. 2013. The incidence of polyploidy in natural plant

populations: major patterns and evolutionary processes. *In* J. Greilhuber, J. Doležel, and J. F.

Wendel [eds.], Plant Genome Diversity Volume 2, 255–276. Springer.

Husband, B. C., and D. W. Schemske. 1997. The effect of inbreeding in diploid and tetraploid

populations of *Epilobium angustifolium* (Onagraceae): implications for the genetic basis of

inbreeding depression. *Evolution* 51: 737–746.

Husband, B. C., and D. W. Schemske. 2000. Ecological mechanisms of reproductive isolation

between diploid and tetraploid *Chamerion angustifolium*. *Journal of Ecology* 88: 689–701.

Ingram, T., and D. L. Mahler. 2013. SURFACE: detecting convergent evolution from

comparative data by fitting Ornstein-Uhlenbeck models with stepwise Akaike Information

Criterion. *Methods in Ecology and Evolution* 4: 416–425.

Iyengar N. K. 1938. Pollen-tube studies in *Gossypium*. *Journal of Genetics* 37: 69–106.

Kembel, S.W., P. D. Cowan, M. R. Helmus, W. K. Cornwell, H. Morlon, D. D. Ackerly, S. P.

Blomberg, C. O. Webb. 2010. Picante: R tools for integrating phylogenies and ecology.

Bioinformatics 26: 1463–1464.

Knight, C. A., R. B. Clancy, L. Götzenberger, L. Dann, J. M. Beaulieu. 2010. On the relationship

between pollen size and genome size. *Journal of Botany* 2010.

Knight, T. M., J. A. Steets, J. C. Vamosi, S. J. Mazer, M. Burd, D. R. Campbell, M. R. Dudash,

et al. 2005. Pollen limitation of plant reproduction: pattern and process. *Annual Review Ecology*

Evolution and Systematics. 36: 467–497.

Kondrashov, F. A., and A. S. Kondrashov. 2006. Role of selection in fixation of gene

duplications. *Journal of Theoretical Biology* 239: 141–151.

Kostoff, D., and A. Prokofieva. 1935. Studies on the pollen-tubes. I. The growth potency of the

pollen-tubes in *Nicotiana* in connection with the length of the styles and some other factors. *Bul.*

Inst. Genetics, Acad. Sci. Leningrad 10: 65–82.

Lande, R., and D. W. Schemske. 1985. The evolution of self-fertilization and inbreeding

depression in plants. I. Genetic models. *Evolution* 39: 24–40.

Landis, J. B., D. E. Soltis, Z. Li, H. E. Marx, M. S. Barker, D. C. Tank, and P. S. Soltis. 2018.

Impact of whole-genome duplication events on diversification rates in angiosperms. *American*

Journal of Botany.

Lee, S-I., and N-S. Kim. 2014. Transposable elements and genome size variations in plants.

Genomics & Informatics 12: 87–97.

Leitch, I. J., M. W. Chase, and M. D. Bennett. 1998. Phylogenetic analysis of DNA C-values

provides evidence for a small ancestral genome size in flowering plants. *Annals of Botany* 82:

85–94.

Leitch, A. R., and I. J. Leitch. 2012. Ecological and genetic factors linked to contrasting genome

dynamics in seed plants. *New Phytologist* 194: 629–646.

Leitch, I. J., and A. R. Leitch. 2013. Genome size diversity and evolution in land plants. *In* J.
Greilhuber, J. Doležel, and J. F. Wendel [eds.], Plant Genome Diversity Volume 2, 307–322.
Springer.

Leitch, I. J., D. E. Soltis, P. S. Soltis, and M. D. Bennett. 2005. Evolution of DNA amounts
across land plants (Embryophyta). *Annals of Botany* 95: 207–217.

Lippman, Z. B., and D. Zamir. 2007. Heterosis: revisiting the magic. *Trends in Genetics* 23: 60–
66.

Magallón, S., S. Gómez-Acevedo, L. L. Sánchez-Reyes, and T. Hernández-Hernández. 2015. A
metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New*
Phytologist 207: 437–453.

Mandakova, T., and M. A. Lysak. 2018. Post-polyploid diploidization and diversification
through dysploid changes. *Current Opinion in Plant Biology* 42: 55–65.

Martins, E. P., T. F. Hansen. 1997. Phylogenies and the comparative method: a general approach
to incorporating phylogenetic information into the analysis of interspecific data. *The American*
Naturalist 149: 646–667.

Mazer, S. J., A. A. Hove, B. S. Miller, and M. Barbet-Massin. 2010. The joint evolution of
mating system and pollen performance: predictions regarding male gametophytic evolution in
selfers vs. outcrossers. *Perspectives in Plant Ecology, Evolution and Systematics* 12: 31–41.

Möller, M. 2018. Nuclear DNA C-values are correlated with pollen size at tetraploid but not
diploid level and linked to phylogenetic descent in *Streptocarpus* (Gesneriaceae). *South African*
Journal of Botany 114: 323–344.

Mulcahy, D.L. 1979. The rise of the angiosperms: a genecological factor. *Science* 206: 20–23.

Nezhad, A. S., M. Packirisamy, R. Bhat, and A. Geitmann. 2013. In vitro study of oscillatory
growth dynamics of *Camellia* pollen tubes in microfluidic environment. *IEEE Transactions on*
Biomedical Engineering 60: 3185–3193.

Obermeyer, G., L. Fragner, V. Lang, and W. Weckwerth. 2013. Dynamic adaption of metabolic

pathways during germination and growth of lily pollen tubes after inhibition of the electron

transport chain. *Plant Physiology* 162: 1822–1833.

Ohri, D., and T. N. Khoshoo. 1986. Genome size in gymnosperms. *Plant Systematics and*

Evolution 153: 119–132.

Otto, S. P., M. F. Scott, and S. Immler. 2015. Evolution of haploid selection in predominantly

diploid organisms. *Proceedings of the National Academy of Sciences* 112: 15952–15957.

Pagel, M. 1997. Inferring evolutionary processes from phylogenies. *Zoologica Scripta* 26: 331–

348.

Pagel, M. 1999. Inferring the historical patterns of biological evolution. *Nature* 401: 877.

Panchy, N., M. Lehti-Shiu, and S-H. Shiu. 2016. Evolution of gene duplication in plants. *Plant*

Physiology 171: 2294–2316.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: analyses of phylogenetics and evolution in

R language. *Bioinformatics* 20: 289–290.

Parre, E., and A. Geitmann. 2005. More than a leak sealant. The mechanical properties of callose

in pollen tubes. *Plant Physiology* 137: 274–286.

Price, H. 1988. DNA Content Variation Among Higher-Plants. *Annals of the Missouri Botanical*

Garden 75: 1248–1257.

Price, H. J., A. H. Sparrow, and A. F. Nauman. 1973. Correlations between nuclear volume, cell

volume and DNA content in meristematic cells of herbaceous angiosperms. *Experientia* 29:

1028–1029.

Ramsey, J., and D. W. Schemske. 1998. Pathways, mechanisms, and rates of polyploid formation

in flowering plants. *Annual Review of Ecology and Systematics* 29: 467–501.

Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other

things). *Methods in Ecology and Evolution* 3: 217–223.

Rounds, C. M., P. K. Hepler, S. J. Fuller, and L. J. Winship. 2010. Oscillatory growth in lily

pollen tubes does not require aerobic energy metabolism. *Plant Physiology* 152: 736–746.

Rounds, C. M., L. J. Winship, and P. K. Hepler. 2011. Pollen tube energetics: respiration,

fermentation and the race to the ovule. *AoB Plants* 2011.

Russell, S. D., X. Gou, C. E. Wong, X. Wang, T. Yuan, X. Wei, P. Bhalla, and M. B. Singh.

2012. Genomic profiling of rice sperm cell transcripts reveals conserved and distinct elements in

the flowering plant male germ lineage. *New Phytologist* 195: 560–573.

Sari-Gorla, M., M. E. Pe, D. L. Mulcahy, and E. Ottaviano. 1992. Genetic dissection of pollen

competitive ability in maize. *Heredity* 69: 423.

Schoenfelder, K. P., and D. T. Fox. 2015. The expanding implications of polyploidy. *Journal of*

Cell Biology: 209: 485–491.

Scholes, D. R., and K. N. Paige. 2015. Plasticity in ploidy: a generalized response to stress.

Trends in Plant Science 20: 165–175.

Selinski, J., and R. Scheibe. 2014. Pollen tube growth: where does the energy come from? *Plant*

Signaling & Behavior 9: e977200.

Shuter, B. J., J. E. Thomas, W. D. Taylor, and A. M. Zimmerman. 1983. Phenotypic correlates of

genomic DNA content in unicellular eukaryotes and other cells. *The American Naturalist* 122:

26–44.

Smith, S. A., and M. J. Donoghue. 2008. Rates of molecular evolution are linked to life history

in flowering plants. *Science* 322: 86–89.

Smith, S. A., and C. W. Dunn. 2008. Phyutility: a phyloinformatics tool for trees, alignments and

molecular data. *Bioinformatics* 24: 715–716.

Snow, A. A. 1990. Effects of pollen-load size and number of donors on sporophyte fitness in

wild radish (*Raphanus raphanistrum*). *The American Naturalist* 136: 742–758.

Soltis, D. E., V. A. Albert, J. Leebens-Mack, C. D. Bell, A. H. Paterson, C. Zheng, D. Sankoff et

al. 2009. Polyploidy and angiosperm diversification. *American Journal of Botany* 96: 336–348.

Soltis, D. E., and P. S. Soltis. 1999. Polyploidy: recurrent formation and genome evolution.

Trends in Ecology & Evolution 14: 348–352.

Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics* 30: 1312–1313.

Stebbins, G. L. 1974. Flowering plants: evolution above the species level. *London: Arnold xviii,*

399p. Illustrations. General (KR, 197500089).

Swift, H. 1950. The constancy of desoxyribose nucleic acid in plant nuclei. *Proceedings of the*

National Academy of Sciences 36: 643–654.

Tadege, M, and C. Kuhlemeier. 1997. Aerobic fermentation during tobacco pollen development.

Plant Molecular Biology 35: 343–354.

Tanksley, S. D., D. Zamir, and C. M. Rick. 1981. Evidence for extensive overlap of sporophytic

and gametophytic gene expression in *Lycopersicon esculentum. Science* 213: 453–455.

Twell, D., S-A. Oh, and D. Honys. 2006. Pollen development, a genetic and transcriptomic view.

In The Pollen Tube, 15–45. Springer, Heidelburg.

Van de Peer, Y., E. Mizrachi, and K. Marchal. 2017. The evolutionary significance of

polyploidy. *Nature Reviews Genetics* 18: 411.

Van't Hof, J. 1965. Relationships between mitotic cycle duration, S-period duration and the

average rate of DNA synthesis in the root meristem cells of several plants. *Experimental Cell*

Research 39: 48–58.

Walsh, N. E., and D. Charlesworth. 1992. Evolutionary interpretations of differences in pollen

tube growth rates. *The Quarterly Review of Biology* 67: 19–37.

Wendel, J. F., D. Lisch, G. Hu, and A. S. Mason. 2018. The long and short of doubling down:

polyploidy, epigenetics, and the temporal dynamics of genome fractionation. *Current Opinion in*

Genetics & Development 49: 1–7.

Whitney, K. D., E. J. Baack, J. L. Hamrick, M. J. W. Godt, B. C. Barringer, M. D. Bennett, C. G.

Eckert et al. 2010. A role for nonadaptive processes in plant genome size evolution? *Evolution:*

64: 2097–2109.

Williams, J. H. 2008. Novelties of the flowering plant pollen tube underlie diversification of a

key life history stage. *Proceedings of the National Academy of Sciences* 105: 11259–11263.

Williams, J. H. 2012. Pollen tube growth rates and the diversification of flowering plant

reproductive cycles. *International Journal of Plant Sciences* 173: 649–661.

Williams, J. H., J. A. Edwards, and A. J. Ramsey. 2016. Economy, efficiency, and the evolution

of pollen tube growth rates. *American Journal of Botany* 103: 471–483.

Williams, J. H., W. E. Friedman, and M. L. Arnold. 1999. Developmental selection within the

angiosperm style: using gamete DNA to visualize interspecific pollen competition. *Proceedings*

of the National Academy of Sciences 96: 9201–9206.

Williams, J. H., M. L. Taylor, and B. C. O'Meara. 2014. Repeated evolution of tricellular (and

bicellular) pollen. *American Journal of Botany* 101: 559–571.

Williams, E. G., and J. L. Rouse. 1990. Relationships of pollen size, pistil length and pollen tube

growth rates in *Rhododendron* and their influence on hybridization. *Sexual Plant Reproduction*

3: 7–17.

Winship, L. J., G. Obermeyer, A. Geitmann, and P. K. Hepler. 2010. Under

pressure, cell walls set the pace. *Trends in Plant Science* 15: 363 – 369.

Wood, T. E., N. Takebayashi, M. S. Barker, I. Mayrose, P. B. Greenspoon, and L. H. Rieseberg.

2009. The frequency of polyploid speciation in vascular plants. *Proceedings of the National*

Academy of Sciences 106: 13875–13879.

 FIGURE LEGENDS

Figure 1: Pollen tube wall construction. A primary cell wall is generated at the tip (red/blue

areas), whereas a secondary wall (angiosperms only) is synthesized in a short region behind the

tip (yellow area). Genome duplication is predicted to double the volume of a mature cell, but

since the tube cell is constantly growing and its cytoplasm is largely confined to the tip region, a

pollen tube can only change its effective volume by changing its fixed diameter (circumference)

or its length per time (pollen tube growth rate, *PTGR*). If wall thickness (*W*) and *PTGR* (*L/t*) are

held constant, a diploid tube cell with doubled volume per unit time will have a 41% larger

circumference. As a result, wall production rate would increase 1.41-fold, not two-fold, over that

of its haploid ancestor: WPR_{1x} (μm^3 h^{-1}) $= C_0$ x *W* x *L/t*; WPR_{2x} (μm^3 h^{-1}) $= 1.41 C_0$ x *W* x *L/t*

Figure 2. Predicted direction of effects of WGD on pollen tube growth rate (*PTGR*).

Genotypic effects (gene dosage and heterozygosity) are predicted to increase *PTGR,* whereas

nucleotypic effects should decrease *PTGR,* after transition from a haploid (1x) to a diploid (2x)

pollen tube. The magnitude of genotypic effects is expected to scale with genetic variation in the

polyploid. The ancestral haploid *PTGR* will be conserved in the derived diploid male

gametophyte when genotypic and nucleotypic effects perfectly offset each other.

Figure 3: Pollen tube growth rate (*PTGR*) evolution across Spermatophytes. Contour plot

showing reconstructed history of *PTGR*. Cool colors indicate *PTGR*s closer to the minimum

value in seed plants while warm colors indicate *PTGR*s closer to the maximum value in seed

plants. Scale bar indicates millions of years before present.

Figure 4: Pollen tube growth rate (*PTGR*) and genome size evolution across seed plants.

Contour plot comparing *PTGR* evolution (left, μm h^{-1}) and c-value evolution (right, picograms).

Scale bars at the bottom of each phylogeny indicate 100 million years. Center bars indicate the

following major groups: gray = gymnosperms, blue = early-divergent angiosperms and

eumagnoliids, green = monocots, orange = asterids, pink = rosids.

Figure 5: Relationship between pollen tube growth rate (*PTGR*) and genome size (C-value)

in seed plants. Separate analyses of gymnosperms (green) and angiosperms (blue) were non-

significant. Selective optima (with standard errors) for each group (from *OUwie* analyses; Tables

S3-S4) are included for illustrative purposes.

Figure 6: Coincident evolution of pollen tube growth rate (*PTGR*) and DNA content (C-

value). Paired *SURFACE* plot showing selective regime shifts in *PTGR* (left) versus genome size

(right). Nodes which have experienced a regime shift along the stem leading to it are marked

with magenta x's. Branch colors – *gray* = seed plant ancestral selective regime; *green* = ancestral

selective regime for angiosperms; *red* = derived selective regimes faster/larger than angiosperm

ancestral regime; and *blue* = derived selective regimes slower/smaller than angiosperm ancestral

regime. Yellow arrows indicate instances where shifts in *PTGR* and genome size coincide. Scale

bars at the bottom of each phylogeny indicate 100 million years. Center bars indicate the

following major groups: gray = gymnosperms, blue = early-divergent angiosperms and

eumagnoliids, green = monocots, orange = asterids, pink = rosids.

Figure S1: Pollen tube growth rate and ploidy level evolution across Spermatophytes.

Table S1: Summary statistics for pollen tube growth rate (*PTGR*) of *Magnolia grandiflora.*

Table S2: Sensitivity analysis for pollen tube growth rate (*PTGR*) error estimates.

Table S3: Pollen tube growth rate (*PTGR*) evolution in gymnosperms vs. angiosperms.

Table S4: C-value evolution in gymnosperms vs. angiosperms.

Table S5: Sister taxon analyses.

Table 1: Phylogenetic signal (Blomberg's *K*) for *PTGR* and C-value.

	K	*P*-value
PTGR seed plants	0.213	<0.0001
PTGR angiosperms	0.075	<0.0001
PTGR gymnosperms	0.060	0.0553
C-value seed plants	0.146	<0.0001
C-value angiosperms	0.099	<0.0001
C-value gymnosperms	0.371	0.169

888 **Table 2**: *OUwie* **model comparison and parameter estimates**.

890 **a.** Results using "restricted" dataset.

model	ΔAICc	model weight	Diploid σ^2	Diploid α	Polyploid σ^2	Polyploid α	Optimum ($\pm$ *SE*)	Diploid optimum ($\pm$ *SE*)	Polyploid optimum ($\pm$ *SE*)
OU1	[418.4]	0.500	0.078	0.067	0.078	0.067	2.728 $\pm$ 0.068		
OUM	1.315	0.259	0.078	0.067	0.078	0.067		2.742 $\pm$ 0.070	1.421 $\pm$ 1.509
OUMA	2.765	0.125	0.082	0.070	0.082	0.068		2.744 $\pm$ 0.069	1.246 $\pm$ 1.480
OUMV	2.921	0.116	0.079	0.066	0.059	0.066		2.742 $\pm$ 0.070	1.375 $\pm$ 1.343
BM1	130.164	2.72E-29	0.033	NA	0.033	NA	2.489 $\pm$ 0.533		
BMS	131.651	1.29E-29	0.034	NA	0.000	NA		2.503 $\pm$ 0.556	1.297 $\pm$ 7.097

892 **b.** Results using "expanded" dataset.

model	ΔAICc	model weight	Diploid σ^2	Diploid α	Polyploid σ^2	Polyploid α	Optimum ($\pm$ *SE*)	Diploid optimum ($\pm$ *SE*)	Polyploid optimum ($\pm$ *SE*)
OU1	[530.8]	0.572	0.076	0.061	0.076	0.061	2.714 $\pm$ 0.064		
OUM	2.045	0.206	0.076	0.061	0.076	0.061		2.713 $\pm$ 0.066	2.805 $\pm$ 1.060
OUMA	3.155	0.118	0.078	0.061	0.078	0.060		2.713 $\pm$ 0.066	2.797 $\pm$ 1.085
OUMV	3.393	0.105	0.077	0.060	0.058	0.060		2.712 $\pm$ 0.067	2.760 $\pm$ 0.973
BM1	153.391	2.81E-34	0.036	NA	0.036	NA	2.469 $\pm$ 0.552		
BMS	156.317	6.51E-35	0.037	NA	0.021	NA		2.514 $\pm$ 0.572	-0.589 $\pm$ 8.214

895 **Table 3:** Phylogenetic generalized least squares regression model comparisons of $\log_{10}$ DNA
896 content versus $\log_{10}$ *PTGR*.
897

Seed plants			Angiosperms only			Gymnosperms only		
Model	**Weight**	***P***	**Model**	**Weight**	***P***	**Model**	**Weight**	***P***
kappa	0.999	0.463	kappa	0.975	0.284	OU	0.265	0.020
lambda	0.001	0.368	lambda	0.024	0.221	delta	0.257	0.006
OU	6.26E-28	0.644	OU	2.90E-04	0.005	kappa	0.193	0.445
delta	1.22E-35	0.471	delta	3.08E-21	0.276	BM	0.121	0.001
BM	1.48E-42	0.373	BM	2.18E-33	0.365	lambda	0.119	0.327
EB	5.45E-43	0.373	EB	8.03E-34	0.365	EB	0.044	0.001

900   **Figure 1**

901
902

903 **Figure 2**

904
905

906 **Figure 3**

907
908

Figure 4

913   **Figure 5**

914
915

916 **Figure 6**