Three decades of native bee biodiversity surveys at Pinnacles National Park

highlight the importance of monitoring natural areas over time

Meiners, Joan M.[1*], Griswold, Terry L.[2], Carril, Olivia Messinger [3]

[1] School of Natural Resources and Environment, University of Florida, Gainesville, FL 32611

[2] USDA-ARS Pollinating Insects Research Unit (PIRU), Utah State University, Logan, UT

84322

[3] 3 Juego Place, Santa Fe, NM, 87508

*Corresponding author

Email: joan.meiners@gmail.com

Keywords: native bees, biodiversity hotspot, conservation, inventory, survey, refugia

Abstract

Thousands of species of bees are in global decline, yet research addressing the ecology and status of these wild pollinators lags far behind work being done to address similar impacts on the managed honey bee. This knowledge gap is especially glaring in natural areas, despite knowledge that protected habitats harbor and export diverse bee communities into nearby croplands where their pollination services have been valued at over $3 billion per year. Surrounded by ranch and farmlands, Pinnacles National Park in the Inner South Coast Range of California contains intact Mediterranean chaparral shrubland. This habitat type is among the most valuable for bee biodiversity worldwide, as well as one of the most vulnerable to agricultural conversion, urbanization and climate change. Pinnacles National Park is also one of a very few locations where extensive native bee inventory efforts have been repeated over time. This park thus presents a valuable and rare opportunity to monitor long-term trends of native bees in natural habitats. Fifteen years after a species inventory marked Pinnacles as a biodiversity hotspot for native bees, we resurveyed these native bee communities over two flowering seasons using a systematic, plot-based design. Combining results, we report a total of 450 bee species within this 109km^2 natural area of California, including 48 new species records as of 2012 and 95 species not seen since 1999. As far as we are aware, this species richness marks Pinnacles National Park as one of the most densely diverse places known for native bees. We explore patterns of bee diversity across this protected landscape, compare results to other surveyed natural areas, and highlight the need for additional repeated inventories of bee biodiversity in protected areas over time amid widespread concerns of bee declines.

Introduction

The importance of bees as critical ecosystem service providers can scarcely be exaggerated. Twenty thousand species of bees worldwide provide the pollination services required for reproduction in 85% of wild and cultivated plants [1,2]. In the United States, the economic importance of bees to agriculture has been valued at up to $14.6 billion annually [3], with $3.08 billion and up to 30% of the U.S. diet specifically credited to the four thousand North American species of native, non-honey bees [4]. Diverse assemblages of native bees have been found capable of enhancing fruit set and yield in the presence of imported honey bees, and of providing adequate pollination for a majority of crops in their absence [5–7]. In natural areas, without the manpower of imported, managed honey bee hives, native bees play a key role in maintaining plant communities that provide soil structure, shelter other invertebrate ecosystem service providers, and establish the base of the food chain [8,9].

Although native bees are often observed pollinating agricultural fields, they seldom nest there. Instead, they rely on nearby remnant patches of semi-natural habitat, a resource that is rapidly disappearing with increasing agricultural intensification, habitat fragmentation, and urban development [10–12]. Despite recognition of natural areas as valuable reservoirs of pollinators [13,14], research on native bee ecology remains concentrated in urban or agricultural settings where baselines may already reflect impacts of degraded ecosystems. Compared to massive honey bee research efforts, progress towards a holistic understanding of how to protect wild bee communities or the habitats they require has not matched their value as pollinators or the known risks they face [15–17].

The relative paucity of research on native bees is due, in part, to the complexity of their biology and behaviors, particularly in wild landscapes. Efforts to monitor wild bees must

contend with the 'taxonomic impediment' of expertise required to evaluate their vast global

biodiversity, and the logistics of sampling a taxon with rapid spatiotemporal turnover, short

lifespans, and solitary, elusive habits [18–21]. Unlike many taxa that follow a latitudinal

biodiversity gradient [22], bee diversity is highest in xeric and Mediterranean environments,

owing to strong seasonal blooms and well-drained soils -- features which support a range of

foraging specializations and high temporal turnover of ground-nesting species [19,20,23]. When

environmental conditions signal a poor year for host plants, some ground-nesting, specialist bee

species can remain underground in diapause for additional years, necessitating multi-year

biodiversity monitoring efforts [24]. This fine and irregular partioning of space and time make

native bees challenging, time-consuming, and expensive to exhaustively sample in any habitat

[25]. Once found, many bee species are difficult to identify even with training and, given reports

of functional redundancy within highly-nested pollination networks, the benefit to ecology of

doing so may seem unclear [26,27]. However, links between non-random species loss and the

stability of ecosystems and mutualistic networks [10,28–34], highlight the merits of species-level

bee biodiversity monitoring. Long-term monitoring of native bee species in natural areas is

necessary to reliably assess trajectories of both thriving and struggling native bee communities

over time, and to forecast their resilience to future climates and perturbations.

Several large surveys of native bee faunas, particularly in the western United States, have

added to current knowledge of the diversity and variability of bee species across space [35–42].

A pair of studies comparing bee faunas from several Mediterranean climate zones concluded that

the chaparral habitats of California represent perhaps the greatest of global biodiversity hotspots

for native bees [43,44]. In the late 1990s, Messinger and Griswold [37] found Pinnacles National

Monument in California's Inner South Coast Range to be one of the most diverse areas known

for bees, with 393 bee species discovered in what was then a 68km^2 area. They attributed this

remarkable richness, in part, to Pinnacles' high floral diversity and habitat heterogeneity [37],

features which also make it an ideal place to investigate relationships between native bee

community dynamics and environmental variables. In 2002, Pinnacles staff conducted a native

bee survey of three changing habitats that added species and a time step to the record of bee

biodiversity in the monument.

Fifteen years after that initial species inventory effort and a decade after the smaller

survey, we returned to Pinnacles, which became a National Park in 2013, to reinventory its

native bee biodiversity and establish a more systematic bee monitoring program [45]. Though

several other bee biodiversity studies have spanned multiple years, as far as we are aware,

Pinnacles is the only region exhaustively surveyed in a previous decade that has since been

resurveyed and published, providing much-needed records of native bee biodiversity over longer

periods of time. As such, our study may aid efforts to understand and protect native bee

biodiversity in natural areas and help determine restoration goals for bee communities in

degraded habitats. Here we present patterns of species accumulation and resource use from three

decades of bee species inventories at Pinnacles National Park and examine how its biodiversity

density compares to other published large-scale bee inventories across the United States.

Materials and Methods

Site description and collecting history

Pinnacles National Park is small as national parks go, approximately 109km^2, yet the

topography is highly dynamic. The roughly oval-shaped park is bisected by a high rock-ridge

spine running north-south that creates a steep elevational gradient and divides the park into a

higher, coastal slope to the west and a drier, lower valley on the east. Initial sampling in 1996 by

TLG suggested a rich bee fauna and motivated the initiation of a more systematic effort to

inventory the bee species across the then-monument's 65km^2 undertaken the following year by

OMC. This first full inventory spanned 1996-1999 and was conducted along the trail network by

opportunistically collecting on a 10-14 day schedule using primarily active (handheld aerial nets)

but also passive (pan traps or "bee bowls") methods during the peak flowering season (locally

February through May). Efforts across these years varied in terms of collecting days (as few as 5

or as many as 56 per year), months covered, and locations sampled. In 2002, a passive pan

trapping study was conducted by a local park ranger in three central regions of the park, with

traps placed out every two weeks between March and mid-July, weather permitting.

In 2005, Pinnacles National Monument acquired an additional 15km^2 of privately-owned

land that expanded the park boundary primarily to the east, but also incorporated some relatively

inaccessible lands to the north and south. In 2010, TLG initiated a follow-up biodiversity survey

of the bees at Pinnacles, including the new lands to the east. In order to better track temporal

trajectories in native bee biodiversity and phenology, we adopted a more systematic park-wide

sampling protocol and established long-term monitoring plots where timed, regular collecting

events using both nets and pan traps were conducted by JMM across the 2011 and 2012

flowering seasons. The following methods and results are focused on this most recent systematic

survey, since a summary of the 1996-1999 inventory has previously been published [37].

Field methods

For the 2011-2012 re-inventory effort, we established ten 1-hectare long-term plots

across a diversity of habitat types and reasonably-accesssible areas of the park. We placed three

plots on the western side of the rocky spine divide: two in grasslands and one in a Blue Oak

woodland. On the larger, lower-elevation eastern side, we set up three plots in alluvial habitats,

two in Live Oak woodlands, and one in a Blue Oak woodland. We also established one plot in a

Blue Oak woodland along the high rock spine bisecting the park. One-hectare rectangular plots

were roughly 200m by 50m, which fit the constraints of the narrow canyon landscapes. In

addition to sampling within plots, we visited areas sampled during the original inventory as well

as newly-aquired lands to conduct opportunistic aerial net collecting and set out pan traps at the

same locations that were sampled using pan traps in 2002 (Fig 1).

Spatially, our collecting extended beyond previous efforts to capture bee biodiversity in

three main ways: by traveling off the trail network (along which most collecting was conducted

in the 1990s, except for one extensive burned area) for plot and opportunistic sampling, by

explicitly establishing repeatedly-sampled plots in a diversity of habitat types across the park,

and by venturing into the 15km^2 of new lands acquired by Pinnacles National Monument in 2005

for both opportunistic and systematic sampling (Fig 1). Temporally, whereas sampling in the

1990s was somewhat irregular, in 2011-12 we sought to capture the full bee community

phenology by sampling plots fortnightly throughout the entire flowering season, beginning in

February before bee activity began and continuing through late June after most bloom had faded

[46].

We sampled all ten plots, typically two per day, every fortnight on days that were mostly

sunny, without high winds, and over 15C°. We conducted additional opportunistic net collecting

along the trail network or in new off-trail areas in between plot efforts. Immediately before each

collecting event, we recorded the ambient temperature, wind speed, humidity, barometric

pressure, and a categorical cloud cover value. During plot sampling, two collectors used aerial

nets to perform thirty-minute timed collections of all bees visually or auditorily detected in plots

at consistent times in both the morning and afternoon. In order to sample the community as

evenly and systematically as possible, we walked a steady pace through plots rather than

focusing on activity at flowers. We placed all netted bees in vials according to their floral host

and collected a voucher plant when the floral host was unknown. At the end of sampling days,

we pinned and labeled all specimens and froze them for 48 hours to prevent beetle infestation.

In addition to net collecting, we also set out thirty colored pan traps, a common passive

collection method, between 9am and 4pm in each plot on the day we net collected there. Pan

traps were made prior to going into the field by spraying 2-oz Solo cups with one of three colors

of paint: fluorescent blue, fluorescent yellow, and white, as indicated by the protocol set up for

native bee monitoring by Lebuhn et al. [47]. Traps were placed directly on the ground across

rectangular plots approximately 10m apart in an "X" pattern in alternating colors and were filled

3/4 full of mildly soapy water to break the surface tension and cause bees visiting the bowls to

sink to the bottom. At 4pm, we strained insects from the water and immersed them in 75%

ethanol until they could be rinsed, pinned and labelled. Data for each pan-trapped specimen

includes the color of the bowl from which it was collected.

Data management and analysis

At the end of the field season, we brought all specimens to the USDA-ARS Pollinating

Insect Research Unit (PIRU) in Logan, Utah and incorporated them into its US National

Pollinating Insects Collection with the exception of small reference and display collections

returned to Pinnacles National Park. Bee identifications were completed by trained experts using

Leica dissecting microscopes, taxonomic literature, and the extensive reference collection housed

at PIRU (approximately 1.5 million curated bee specimen). After processing all 2011 and 2012

bee specimens, we reviewed all identifications for the Pinnacles bees from the 1996-1999 and

2002 collections (which are also housed at PIRU) to ensure nomenclature was current and

consistent with recent inventory identifications. We identified plant vouchers using appropriate

keys [48] and guidance from botanists at Pinnacles or the Utah State University Intermountain

Herbarium.

We entered field data into PIRU's existing relational database, assigned corresponding

individual ID numbers and barcodes to each specimen, and pinned labels with this information to

each bee. We conducted quality checks with multiple people at each step of the curation process.

We used SQL and Microsoft Access to query and manage data, and Microsoft Excel, R-Cran

statistical package version 0.99.879 or ARC-GIS to clean, analyze, and map data [49]. Data is

either included as supplementary tables or will be deposited with Dryad. Data and code for

analysis will be publicly available on Github.

Literature review

To place the bee biodiversity results at Pinnacles National Park in context with those of

other bee inventory efforts across the United States, we conducted a literature search for all

published studies that reported at least one hundred bee species from natural (non-agricultural,

non-urban) areas and methods indicative of an exhaustive, systematic diversity inventory. Using

Web of Science and Google Scholar, we identified nineteen published studies that met these

criteria, to which we added four unpublished studies that qualify. To allow for a quantitative

comparison of relative richness between exhaustive bee surveys, we used a novel metric to

calculate biodiversity density along the species-area curve based on the number of species and

genera reported in each publication as well as the total size of the area covered. For studies that

195 did not specify the area of land covered, we contacted authors for estimates and/or performed a

196 web search of the study place named to estimate total area surveyed.

Results

Pinnacles bee collections over time

200   Initial opportunistic trail collecting between 1996-1999 yielded 27,055 bee specimens

201 representing 382 species and 52 genera collected over 125 collector days at 32 different locations

202 within the old monument boundary (Table 1) (differences from results reported by Messinger

203 and Griswold in 2003 are a result of recent taxonomic changes) [37]. The smaller pan trapping

204 study by park empoyee Amy Fesnock over 10 days in 2002 yielded 7,255 bees representing 151

205 species and 38 genera from 3 different locations in the central lowlands of the eastern edge and

206 exterior of the monument boundary. In the recent inventory during the flowering seasons of 2011

207 and 2012, we captured 52,789 bees over 214 collector days (107 days with two collectors) at 90

208 different locations across all accessible areas of the park (Fig 1). This effort resulted in a

209 collection of 291 bee species across 45 genera in 2011 and 294 species across 49 genera in 2012

210 (Table 1a). There was a 79% overlap in species and an 94% overlap in genera between the two

211 years (Table 1b). The preservation and curation of older specimens enabled us to update species

212 determinations from previous inventories based on more recent taxonomic changes to compare

213 and combine biodiversity records across inventory efforts (Table 2).

214 **Table 1. Summary of bee sampling efforts at Pinnacles National Park.** (a) Specimen

215 collection statistics by year of sampling. (b) Proportion overlap between bee species and genera

216 collected during each year of sampling.

217 **(a)**

Bee collection statistics for Pinnacles Natl Park	Grand totals	Year						
		1996	1997	1998	1999	2002	2011	2012
Number of Specimens Collected	**87,099**	1362	8077	9382	8234	7255	20351	32438
Number of Species Collected	**450**	172	299	313	211	151	291	294
Number of Genera Collected	**54**	38	48	49	43	38	45	49
Number of New Species Records	--	all	140	60	10	20	22	26
Number of New Genus Records	--	all	11	1	0	0	0	3
Specimens per New Species Record	**177**	8	56	142	749	470	565	903
Specimens per New Genus Record	**1668**	36	734	9383	--	--	--	10839
Species Unique to that Year	--	4	22	21	2	5	15	26
Genera Unique to that Year	--	1	0	0	0	0	0	3
Days of Collecting	**246**	5	50	56	14	10	55	52
Methodology (equipment): *Since methodology and sampling effort vary widely between years and projects, comparisons should be interpreted with caution.*		Opportunistic trail collecting (aerial handheld net) + pan traps				Passive collecting (pan trap bowls)	Plot (N=10) sampling (aerial nets + pan traps); Trail collecting (nets); Resample of 2002 bowl sites (pan traps)	
Primary Collectors		Olivia Messinger Carril & Terry Griswold				Amy Fesnock	Joan Meiners & Therese Lamperty	

218 **(b)**

	1996	1997	1998	1999	2002	2011	2012	
1996	**1.0**	0.68	0.64	0.71	0.52	0.63	0.61	
1997	0.86	**1.0**	0.81	0.73	0.49	0.75	0.52	**Proportion of species in common (above daigonal)**
1998	0.85	0.99	**1.0**	0.72	0.52	0.78	0.75	
1999	0.89	0.95	0.93	**1.0**	0.52	0.69	0.67	
2002	0.87	0.88	0.87	0.91	**1.0**	0.57	0.59	
2011	0.89	0.95	0.96	0.93	0.92	**1.0**	0.79	
2012	0.85	0.93	0.94	0.91	0.87	0.94	**1.0**	
Proportion of genera in common (below diagonal)								

Fig 1. Map of Pinnacles National Park in Monterey and San Benito Counties, California.

As a national monument, established in 1908, it grew from 36 km^2 to 68km^2, shown by the

shaded region. The outlined area encompasses lands added in 2005 and represents the current

national park boundary (109 km^2). Locations sampled during the original native bee inventory of

1996-1999 are marked with filled black circles. The three locations where native bees were

sampled with pan traps in 2002 are marked by open circles around an 'x'. For the 2011-2012

survey, plus signs mark sites of opportunistic sampling and colored squares indicate the habitat

type and position (not sized to scale) of systematically-sampled hectare plots. Dense chaparral

shrubs, steep hillsides, homogenous landscapes, and few trail access points made the northern

and southern regions of the park relatively inaccessible and unpromising for bee diversity.

230 **Table 2**. **Overview of Pinncles National Park bee biodiversity and comparisons between**

231 **survey efforts**. Numbers of species unique to that survey timeframe are in parentheses. Due to

232 taxonomic changes, updated species determinations, and the addition of data from 2002, some

233 totals differ from those reported in Messinger and Griswold 2003. See S1 Table for additional

234 species details.

Family	*Genus*	Number of species in early surveys (1996-1999, 2002)	Number of species in recent survey (2011-2012)	Number of singleton species (represeted by only one specimen)	Number of species recorded only in new lands (acquired in 2005)	Cleptoparasitic (C); Oligolectic (O)
Andrenidae	*Ancylandrena*	1 (1)		1		O
	Andrena	60 (19)	49 (8)	7	3	
	Calliopsis	8 (2)	6			O
	Macrotera	1	1			O
	Panurginus	4 (1)	3			
	Perdita	13 (5)	10 (2)	3	1	O
Apidae	*Anthophora*	12 (4)	8	3		
	Anthophorula	2	2			
	Apis	1	1			
	Bombus	6 (1)	5			
	Brachynomada		1 (1)			C
	Centris	1 (1)				
	Ceratina	11	11			
	Diadasia	5	9 (3)	1		O
	Epeolus	3	4 (1)		1	C
	Eucera	9	9			
	Habropoda	3 (1)	2			
	Melecta	3	3			C
	Melissodes	9 (4)	8 (3)	3	3	
	Neopasites		1 (1)	1	1	C
	Nomada	26 (10)	21 (5)	6	1	C
	Oreopasites	2	2			C
	Peponapis		1 (1)			O
	Townsendiella	2 (1)	1	1		C
	Triepeolus	2 (1)	7 (5)	1	1	C
	Xeromelecta	2 (1)	1	1		C
	Xylocopa	1	1			
Colletidae	*Colletes*	5 (1)	5 (1)	2		O
	Hylaeus	15 (5)	10	2		

Family	Genus					
Halictidae	*Agapostemon*	2	2			
	Augochlorella	1	1			
	Conanthalictus	2	2			O
	Dufourea	6	7 (1)	1		O
	Halictus	4 (1)	3			
	Lasioglossum	28 (2)	29 (3)	2	2	
	Micralictoides	2	2			O
	Sphecodes	10 (5)	6 (1)	3		C
Megachilidae	*Anthidiellum*	1	1			
	Anthidium	6 (3)	5 (2)	1	1	
	Ashmeadiella	16 (5)	13 (2)	2	1	
	Atoposmia	3 (1)	3 (1)	1		O
	Chelostoma	7	7			O
	Coelioxys	4 (2)	3 (1)	1		C
	Dianthidium	5 (1)	4	1		
	Dioxys	4 (1)	3	1		C
	Heriades	1 (1)				
	Hoplitis	17 (2)	15			O
	Megachile	18 (5)	15 (2)	1		
	Osmia	38 (6)	35 (3)	5	2	
	Protosmia	1	1			
	Stelis	13 (2)	12 (1)		1	C
	Trachusa	2	2			
Melittidae	*Hesperapis*	2	2			O
Totals		**417 (95)**	**355 (48)**	**51**	**18**	

The combined results from all three inventories document a total of 450 species of bees across 53 genera and all six North American bee families within the modest 109km^2 of Pinnacles National Park (Table 2). The most recent survey documented 48 new species records for the Pinnacles National Park area and did not recapture 95 species that had been collected in earlier studies (S1 Table). Of the 48 species recorded for the first time in 2011 and 2012, 47 were rare (here defined as represented by fewer than ten specimens), and 20 were singletons (represented by a single specimen) (S1 Table). Thirty of the 48 new species were captured in areas previously sampled, while 18 were only captured in new lands added to the park since previous inventories (Table 2). Overall, 51 of the 450 species were singletons (Table 2), and 95 were present in only

one year of sampling, with the majority of these temporally rare species being from the families

Apidae and Andrenidae (Fig 2a). The family Megachilidae had the most species present in all

seven years of sampling (N=38 out of 68 total) (Fig 2a). Overlap in species lists between years

ranged from 49% to 81% and overlap in genera ranged from 85-99% between any two years

(Table 1b).

Despite extensive sampling of bee biodiversity within Pinnacles National Monument

between 1996-1999, subsequent sampling continued to add species richness to the overall

collection (Fig 2b). The 2002 effort added 20 new species to the park list. The 2011 collection

netted 22 bee species new to Pinnacles, and the 2012 collection, which sampled mostly the same

areas as 2011, resulted in 26 new species and 3 never-before recorded genera within Pinnacles

National Park (Table 1a). Between 2 and 26 species were unique to a particular year and not

recorded within the park during any of the other six years of surveys. The genus *Ancylandrea*

(family Andrenidae) was present only in the 1996 collection and 2012 was the only year that

three genera from the family Apidae (*Neopasites*, *Peponapis*, and *Brachynomada*) were

documented (S1 Table). For five out of six bee families, new species were added to the park list

nearly every year. Melittidae is represented by only two common species, both of which were

collected in the original year of sampling, and in every year thereafter (Fig 2b).

Fig 2. Comparison of bee species collections at Pinnacles National Park over seven years of surveys. (a) Numbers of species in each of six North American bee families represented in up to all seven years of collections. (b) Accumulation over time of number of species collected in each of six North American bee families from each additional year of collecting.

Recent Pinnacles bee survey details

During the 2011-2012 survey, we completed 150 plot samples across our ten one-hectare

plots, eighty in 2011 and seventy in 2012, sampling only on days that were sufficiently sunny,

calm, and warm to ensure adequate bee activity for comparisons between plots. In 2011, 80 plot

samples conducted over 55 days resulted in between 1 and 2088 bees from an individual plot

sample, with a mean of 368 bees per plot per day and a standard deviation of 398. In 2012, 70

plot samples conducted over 52 days resulted in between zero and 1317 bees collected in a day

and plot, with a mean of 370 and a standard deviation of 380 bees per plot per day.

A species accumulation curve for the observed rate of capture of the 334 species

collected in plots across 150 plot samples shows that our efforts captured a majority of the

estimated true bee biodiversity within these areas (Fig 3). The leveling off of the curve at the far

right indicates that additional plot sampling would be unlikely to yield many more species to the

collection, especially for organisms like insects for which observed richness rarely reaches an

asymptote [50]. Expanding collecting efforts into areas not extensively surveyed in the more

remote and homogenous chaparall habitats in the northern and southern ranges of the park may

yield additional bee biodiversity (Fig 1). However, the number of singleton or doubleton species

recorded across many genera illustrates the frequency of rare bee species at Pinnacles National

Park, which additional sampling efforts may or may not detect (S1 Table). The blue curve and

vertical confidence interval lines indicate the estimated rate of species accumulation for a

random community with the same number of species and samples (Fig 3). That the observed

curve has an initially steeper slope than expected is indicative of Pinnacles' rich biodiversity

resulting in rapid early accumulation of common species.

Fig 3. Species accumulation curve. Observed rate of accumulation of 334 species across 150

samples (black line, grey 95% confidence interval bands) compared to an expected rate of

species accumulation for a random community with the same number of species and samples

(blue line and 95% confidence interval bars).

Bee species richness in 150 plot samples was normally distributed (Shapiro-Wilk

normality test, p=0.8) and positively related to the floral richness of bee-visited plants by a

power-law linear regression model (Bee Richness = exp(2.79 + 0.38*log(FR)); R^2=0.37, p<0.01,

S1a Fig). To a lesser extent, bee abundance (square-root transformed to normalize distribution)

was also significantly positively correlated with the floral diversity of bee-visited plants in plot

samples (Bee Abundance = exp(2.26 + 0.23*log(FR)); R^2 = 0.16, p<0.01, S1b Fig).

Bee abundance, dominance, and floral activity varied between species and the two

consecutive years of sampling at Pinnacles National Park. Across all 150 plot samples over two

years, *Lasioglossum* (Halictidae) was the most abundant bee genus, followed by *Hesperapis*

(Melittidae), *Osmia* (Megachilidae)*, and *Halictus* (Halictidae). *Oreopasities*, *Peponapis*,

Xeromelecta, and *Townsendiella* (all Apidae) were among the rarest genera collected over the

two years of plot sampling; all but *Peponapis* are cleptoparasites.

Between years, rank abundance of the top twenty-five bee species reflects high

interannual species turnover, with *Hesperapis regularis* (Melittidae) occupying the top spot in

2011 and only ranking as the fourth most abundant species in 2012 (Table 3a). Similarly, *Osmia*

nemoris (Megachilidae) was the most abundant species collected in plot samples at Pinnacles in

2012, after having been ranked fifth most abundant in 2011. Halictidae was the bee family with

the highest number of most abundant species in both years, followed by Megachilidae in 2011

and Andrenidae in 2012 (Table 3a).

The most bee-popular plants also varied between years. In 2011, more bees visited

Clarkia unguiculata (Onagraceae), the host plant of 2011's most abundant bee, *Hesperapis*

regularis, than any other plant (N= 247, compared to 116 bees on this flower in 2012), and

Eriogonum fasciculatum (Polygoneaceae) was visited by the most bees in 2012 (N = 644,

compared to 109 bees on this flower in 2011) (Table 3b). *Adenostoma fasciculatum* (Rosaceae)

and *Eschscholzia californica* (Papaveraceae) maintained their positions as the second and third

most bee-popular plants, respectively, in both years of collecting. Floral species from the

Boraginaceae family dominated the list of top twenty-five most bee-popular plants in 2011 and

tied with Asteraceae and Fabaceae for most bee-popular family in 2012 (Table 3b).

Table 3. Most commonly-collected bees and most bee-popular plants in 2011 and 2012

surveys at Pinnacles National Park. (a) Twenty-five most commonly-collected bee species by

rank abundance per year. (b) Twenty-five most commonly recorded plants visited by bees,

ranked by popularity with bees per year. See S1 and S2 Tables for the complete taxa lists.

(a)

During the 2011 flowering season			Rank Abun.	During the 2012 flowering season		
Bee Family	*Genus*	*Species*		**Bee Family**	*Genus*	*Species*
Melittidae	*Hesperapis*	*regularis*	1	Megachilidae	*Osmia*	*nemoris*
Halictidae	*Halictus*	*tripartitus*	2	Halictidae	*Halictus*	*tripartitus*
Halictidae	*Lasioglossum*	*nigrescens*	3	Halictidae	*Lasioglossum*	*incompletum*
Halictidae	*Lasioglossum*	*brunneiventre*	4	Melittidae	*Hesperapis*	*regularis*
Megachilidae	*Osmia*	*nemoris*	5	Halictidae	*Halictus*	*farinosus*
Halictidae	*Lasioglossum*	*incompletum*	6	Halictidae	*Lasioglossum*	*nigrescens*
Apidae	*Apis*	*mellifera*	7	Apidae	*Melissodes*	*stearnsi*
Halictidae	*Lasioglossum*	*punctatoventre*	8	Apidae	*Apis*	*mellifera*
Halictidae	*Halictus*	*farinosus*	9	Halictidae	*Lasioglossum*	*brunneiventre*
Halictidae	*Lasioglossum*	*sp. 9*	10	Halictidae	*Lasioglossum*	*punctatoventre*
Halictidae	*Lasioglossum*	*imbrex*	11	Apidae	*Eucera*	*actuosa*
Apidae	*Ceratina*	*arizonensis*	12	Andrenidae	*Panurginus*	*gracilis*
Andrenidae	*Andrena*	*aff. cerasifolii*	13	Halictidae	*Agapostemon*	*angelicus/texanus*
Andrenidae	*Andrena*	*sp.*	14	Halictidae	*Lasioglossum*	*sp. 9*
Halictidae	*Agapostemon*	*angelicus/texanus*	15	Apidae	*Diadasia*	*bituberculata*
Andrenidae	*Andrena*	*crudeni*	16	Apidae	*Melissodes*	*sp.*
Halictidae	*Lasioglossum*	*nevadense*	17	Andrenidae	*Perdita*	*distropica*
Megachilidae	*Protosmia*	*rubifloris*	18	Halictidae	*Lasioglossum*	*sp.*
Apidae	*Eucera*	*actuosa*	19	Andrenidae	*Andrena*	*aff. cerasifolii*
Megachilidae	*Osmia*	*brevis*	20	Megachilidae	*Osmia*	*aglaia*
Andrenidae	*Panurginus*	*gracilis*	21	Megachilidae	*Osmia*	*regulina*
Halictidae	*Lasioglossum*	*sisymbrii*	22	Halictidae	*Lasioglossum*	*nevadense*
Apidae	*Diadasia*	*angusticeps*	23	Andrenidae	*Andrena*	*macrocephala*
Megachilidae	*Trachusa*	*perdita*	24	Andrenidae	*Andrena*	*w-scripta*
Megachilidae	*Osmia*	*regulina*	25	Apidae	*Ceratina*	*arizonensis*

328 **(b)**

During the 2011 flowering season		Popul. Rank	During the 2012 flowering season	
Plant Name	Plant Family		*Plant Name*	Plant Family
Clarkia unguiculata	Onagraceae	1	*Eriogonum fasciculatum*	Polygonaceae
Adenostoma fasciculatum	Rosaceae	2	*Adenostoma fasciculatum*	Rosaceae
Eschscholzia californica	Papaveraceae	3	*Eschscholzia californica*	Papaveraceae
Clarkia purpurea	Onagraceae	4	*Clarkia unguiculata*	Onagraceae
Chaenactis glabriuscula	Asteraceae	5	*Hirschfeldia incana*	Brassicaceae
Lotus scoparius var.scoparius	Fabaceae	6	*Marrubium vulgare*	Lamiaceae
Ranunculus californicus	Ranunculaceae	7	*Eriodictyon tomentosum*	Boraginaceae
Eriogonum fasciculatum	Polygonaceae	8	*Chaenactis glabriuscula*	Asteraceae
Hirschfeldia incana	Brassicaceae	9	*Amsinckia menziesii*	Boraginaceae
Salix exigua	Salicaceae	10	*Salix lasiolepis*	Salicaceae
Lupinus albifrons	Fabaceae	11	*Clarkia purpurea*	Onagraceae
Vicia villosa	Fabaceae	12	*Lasthenia californica*	Asteraceae
Eriodictyon tomentosum	Boraginaceae	13	*Lupinus albifrons*	Fabaceae
Viola pedunculata	Violaceae	14	*Calochortus venustus*	Liliaceae
Quercus agrifolia var.agrifolia	Fagaceae	15	*Ceanothus cuneatus*	Rhamnaceae
Lasthenia californica	Asteraceae	16	*Chorizanthe douglasii*	Polygonaceae
Marrubium vulgare	Lamiaceae	17	*Erodium cicutarium*	Geraniaceae
Pholistoma auritum	Boraginaceae	18	*Salix exigua*	Salicaceae
Arctostaphylos pungens	Ericaceae	19	*Penstemon heterophyllus*	Plantaginaceae
Amsinckia menziesii	Boraginaceae	20	*Lotus scoparius var.scoparius*	Fabaceae
Ceanothus cuneatus	Rhamnaceae	21	*Baccharis salicifolia*	Asteraceae
Bloomeria crocea	Liliaceae	22	*Vicia villosa*	Fabaceae
Heliotropium curassavicum	Boraginaceae	23	*Malacothamnus aboriginum*	Malvaceae
Erodium brachycarpum	Geraniaceae	24	*Ranunculus californicus*	Ranunculaceae
Salix lasiolepis	Salicaceae	25	*Heliotropium curassavicum*	Boraginaceae

Pinnacles bee biodiversity in context

To enable an analysis of how remarkable the bee biodiversity at Pinnacles really is, we

used literature searches and expert opinions to compile a list of 23 studies within the United

States that matched our criteria for comparison (N>100 species, extensive inventory-style

sampling in a natural area) (S3 Table). It is worth visualizing that, while efforts to survey native

bees have increased in recent years, these published inventories still only cover a small

proportion of natural areas and habitat types across the United States, and thus offer only a small

window into the status of native bees across the country (Fig 4).

Fig 4. **Map of the location, size, and number of bee species recorded for all exhaustive bee**

inventory efforts undertaken across the United States for which data is published or

reported. The black arrow points to Pinnacles National Park. See S3 Table for more details on

projects shown.

Without controlling for the area sampled, Pinnacles' 450 bee species place it fourth

among 23 completed studies reporting high numbers of bee species within a natural area. Studies

with more total bee species include Grand Staircase Escalante National Monument, where OMC

recorded 656 different species of bees between 2000-2003 [38], a study conducted by TLG in

Clark County, Nevada that documented 598 bee species over three years [35], and an

unpublished study in Yosemite National Park in the mid-2000s that found 554 species (*Griswold,*

unpublished data). A variety of additional systematic inventories conducted in natural lands also

report stunning bee biodiversity, including 393 bee species found over seven years in San

Bernardino Valley, Arizona [23], previously thought to have the highest biodiversity of native

bees by area.

A meaningful biodiversity comparison between this list of bee inventories is hindered by

the vastly different areas covered by each. After removing one study that covered less than

$10km^2$ from comparisons to avoid errors in overestimating beta richness across larger areas by

scaling-up alpha diversity sampled at smaller-than-local scales [51], we calcuated the number of

bee species recorded per km^2 for each listed survey (S3 Table). By this crude comparison, the

areas with the highest density of bee species in the United States include the Black Rock Forest

Preserve, where 144 species were discovered within less than $16km^2$ [42], MPG Ranch in

Montana, where 229 species were discovered over $39km^2$ [52], and Archibald Biological Station

in Florida, where 113 species have been recorded across $21km^2$ [53]. With 450 species of native

bees detected over the past three decades across just $109km^2$, Pinnacles National Park again

ranked fourth in the measure of most densely-diverse areas for native bees.

Because species density is known to not scale linearly across increasing area, we then

incorporated the knowledge of increasing diversity over space modeled by the species-area

curve. We plotted a power-law species-area relationship based on the reported species richness

and area covered by known bee inventories (S3 Table) to calculate which of the 23 listed studies

found lower-than-expected bee richness based on their size and which studies were likely true

hotspots of native bee biodiversity (Fig 5).

a

b

Fig 5. **Species-Area relationships and trend line for all major, exhaustive bee inventory**

studies conducted in the United States in natural or semi-natural habitats. (a) The black

trend line delineates expectations for how the number of species will increase with increasing

area size based on the (log-transformed) species-area relationship. Studies above the trend line

(grey points) recorded more bee species than expected for the area of the site; those below the

line (black points) recorded fewer bee species than might be expected on average for that size

area. Pinnacles National Park is circled in red. (b) Barplot of the difference in the number of bee

species observed in each study relative to the number of bee species predicted by the trend line

plotted in panel (a). Pinnacles National Park is outlined in red. Studies are listed in S3 Table.

Based on this difference between observed and expected species richness per area (the

positive or negative distance of the point to the trend line in Fig 5), we conclude that Pinnacles

National Park is home to the highest bee biodiversity per area surveyed of any published or

known exhaustive bee biodiversity survey (with over 100 species) in natural areas across the

United States. Grand Staircase Escalante National Monument (GSENM) also contains more bee

biodiversity than would be expected by even its vast size, as does Yosemite National Park;

Carlsbad Caverns National Park; Clark County, Nevada; San Bernardino, Arizona; Carlinville,

Illinois; MPG Ranch, Montana; Curlew Valley, Idaho; Indiana Dunes, Indiana; and San Rafael

Desert, Utah. Studies that reported bee biodiversity lower than what would be expected by our

species-area relationship included Black Belt Prairie, Missouri; Hattiesville, Missouri; Tonasket

Ranger District, Washington; and the Black Hills of South Dakota and Wyoming, among other

natural areas (Fig 5, S3 Table). Many more studies will be necessary to fill in the map of bee

biodiversity in natural areas (Fig 4) and interpret how the bee species-area relationship relates to

ecosystem, climate, or habitat stage (Fig 5).

Discussion

Wild, native bees are key ecosystem service providers in both natural and agricultural

landscapes [5–7,54]. Compared to the unstable European honey bee, on which United States

agriculture is heavily dependent, little is known about the four thousand North American species

of native bees, who may also be vulnerable to the parasites, pesticides, and habitat modification

plaguing the honey bee [3,16,17,34,55,56]. One of the reasons for this lack of attention to native

pollinators is the expense, time, and skill required to collect and identify native bees, which are

spatiotemporally variabile, short-lived, diverse in their taxonomy and nesting habits, and often

difficult to see. Even when extensive bee inventories are conducted at intensities and intervals

sufficient to capture local diversity in native bees, our literature review found that they are rarely

replicated later, resulting in few datasets that allow for robust assessment of trends in native bee

populations over ecologically relevant time scales.

With three separate inventories conducted over three decades, the native bee inventory

efforts at Pinnacles National Park in the Inner South Coast Range of California represent an

exception to this lack of temporal knowledge. Combined results from seven years of sampling

suggest that Pinnacles National Park may harbor the highest density of bee species currently

known anywhere in the United States, and potentially the world, since California is already

recognized as a global bee biodiversity hotspot [20]. In comparison to Pinnacles' 450 species

across an area of 109km^2, only 388 species of bees have been recorded in the state of Wisconsin

and only 40 species on the entire two large islands of New Zealand [57,58]. The closest

comparison by habitat type outside of the United States may be a survey conducted 1983-1987

over a Mediterranean area of unspecified size outside Athens, Greece that reported 661 species

of bees [59]. However, the fact that substantial species diversity was added to the bee inventory

list for Pinnacles even after five prior years of surveys (Figs 2b and 3) suggest that inventories in

other locations over shorter timespans may grossly undercount rare species. Furthermore, the

variability in bee species over time at Pinnacles (Fig 2a) underscores the the importance of long-

term sampling to meet the research challenge of detecting the signal amidst the noise of bee

community variability.

Until that time when more comparable data are available, our review of exhaustive bee

surveys conducted in the continental United States identified only three natural areas with more

recorded bee species than Pinnacles, all conducted across areas larger than Pinnacles by at least

an order of magnitude (S3 Table). When plotted along a species-area curve, Pinnacles National

Park also has the greatest deviation above the curve compared to any of these other studies,

suggesting it contains far more species per area sampled than would be expected in other

locations of similar size (Fig 6). Bee diversity in GSENM in Utah, with 1.5 time the number of

bees as Pinnacles over 70 times the area, also exceeds the diversity predicted by the species-area

curve, though not by the same margin. These results suggest that, while GSENM and several

other protected regions sampled are remarkably diverse for the area covered, Pinnacles National

Park is home to the most exceptional bee species density of any known place.

We attribute the extraordinarily rich bee fauna of Pinnacles National Park to its

Mediterranean climate, steep environmental gradients, and high habitat heterogeneity, the last of

which has been found in other research to be a stronger predictor of species richness than the

species-area relationship [60,61]. Habitat heterogeneity can occur over both space and time.

Mediterranean habitats, like those at Pinnacles, are known for rich 'flash-bloom' cycles during

spring months, followed by hot, dry summers and mild, wet winters, an environment that tends

to support a high biodiversity of many taxa by creating many temporal habitat niches [9,62].

Among bees, the rapid turnover of floral resources in these areas may favor solitary species,

whose shorter flight periods and more specialized foraging behaviors may allow many species to

coexist in a single area, as each occupies a narrower temporal and foraging niche space than

longer-lived social or generalist species, which are more common in temperate areas [19,23].

Across space, habitats at Pinnacles change rapidly from the western, coastally-influenced

slopes, up the 500m elevational gradient to the rock ridge, and down the different aspects and

microclimates of the drier east side. Pinnacles spans several fault lines, the geologic movements

of which may have contributed to its elevational variation and broader array of soil types than

would typically be found in such a small area [63]. Perhaps because of this soil heterogeneity,

Pinnacles is also considered to be a transitional zone between the floral ecotones of northern and

southern California [64] and boasts an extraordinary plant list of nearly 700 species, many of

them flowering ones [65]. We found bee richness to be highly correlated with the richness of

bee-visited angiosperms on any given day and site at Pinnacles (S1 Fig), which corroborates

results from previous studies [9,38]. Indeed, our conclusion is that the extraordinary diversity of

native bees at Pinnacles is a function of the dynamic climate, rich wildflower flora, and

landscape patchiness creating a wide array of spatiotemporal habitat niches. These factors may

allow more diverse bee communities to coexist across space than has been found anywhere else.

The unparallelled biodiversity of native bees at Pinnacles National Park is especially

intriguing given its juxtaposition with nearby agricultural intensity. Salinas Valley, at the

doorstep of Pinnacles National Park, produces most of the strawberries, tomatoes, spinach,

lettuce, celery, and garlic for the country, among other smaller crops. Many of the lands

surrounding the park that aren't irrigated for crops are grazed by cows, reducing floral diversity

and thus their suitability for diverse native bees [66]. Native bees are most diverse in natural,

undisturbed areas, proximity to which has been directly linked to crop pollination success

because of the constant influx of wild pollinating insect populations into arated lands inhospitible

to long-term residence [11,13]. Agricultural habitats fail to support diverse native bees due to

impacts of pesticides, nutritional deficits resulting from monocultures offering only one type of

bloom, and the practice of tilling and turning over the soil where many native bee species

overwinter [5,30,67]. The native bees known to pollinate crops persist not within the fields but in

nearby patches of natural, uncultivated land. California has increased efforts to restore habitat for

wild bees in agricultural lands. But less attention has been paid to bee source populations in

adjacent natural areas, even though source-sink dynamics have recently been determined to

influence bee population sensitivity to decline [68].

If Pinnacles National Park is indeed a biological refuge for native bee populations within

a highly-altered landscape, it will be even more important to track trends in its bee biodiversity

over time. Our establishment of ten 1-hectare plots and repeatable methodology will facilitate

ongoing monitoring activities and better comparisons of bee biodiversity and population stability

over time than are currently possible. During 2011 and 2012, we recorded 355 species of bees at

Pinnacles National Park, 48 of which were new records for the park. Initial inventories in the

1990s recorded 382 species, 95 of which we did not encounter during the recent inventory. After

six prior years of sampling and a clear leveling of the species accumulation curve, we still

recorded three new genera in 2012. These results illustrate the difficulty in deciphering

ecological trends from inventories conducted using different methods or in different locations.

Long-term, systematic monitoring studies in consistent locations will enable improved

understanding of species turnover, range extensions (invasions), local extinctions, baseline states,

and how to differentiate natural community variability from bee biodiversity decline, a question

we consider a research priority towards assessing trajectories of native bee decline.

The need for multi-year, temporally replicated bee surveys to better quantify trends and

declines in native bees over time is further highlighted by the recent increase in the use of

chronosequences, which substitute space as a proxy for time in restored habitats to model

changes in native bee dynamics [69,70]. This is a clever approach but increasing efforts to repeat

surveys using the same methodology in the same natural areas over actual timespans would be

better. Spatial coverage of published bee inventory studies is sparse (Fig 5), and temporal

coverage is worse. Expanding long-term bee biodiversity monitoring to additional habitats and

supporting the museum work and collection maintenance that enable these important temporal

comparisons will bolster our chances of protecting native bee biodiversity and agricultural

stability.

Conclusions

Here we reported details of the third extensive bee inventory effort at Pinnacles Naitonal

Park in California over three decades in order to share ongoing findings from a native bee

biodiversity hotspot and to highlight the need for more studies like this to evaluate trends among

our most important polliantors. Our study is the first to compile and compare information on

native bee biodiversity from published inventory studies in natural areas across the United States.

With 450 species of native bees, we found that Pinnacles National Park houses a higher density

of species than any other natural area studied, and compared to seven California urban areas

surveyed (species N = 60-80) [54]. This result indicates that America's newest national park may

be a substantial exporter of free, native pollinators into economically valuable agricultural lands.

However, our discovery that Pinnacles is the only area to have been extensively and repeatedly

surveyed for bee biodiversity over multiple decades underscores our call for increased repeated

monitoring efforts to facilitate research on bee population decline and variability at its source.

Acknowledgements

We are grateful to Therese Lamperty for dedication in the field, and to Harold Ikerd and

Skyler Burrows for assistance in the lab. This work would not have been possible without

generous guidance from USU co-P.I. Edward W. Evans and from Pinnacles wildlife biologist

Paul G. Johnson. We thank Michael Orr, Skyler Burrows, Harold Ikerd, Karen Wright, Zachary

Portman, Brian Rozick, and Ethan Frehner for help with bee identifications; Valerie Nuttman,

Brent Johnson, and Denise Louie for support at Pinnacles; Amy Fesnock for her 2002 work on

Pinnacles bees; Eugene Schupp, Hao Ye, Erica Christensen, and Kenny Anderson for comments

on drafts; and Jereme Gaeta, Cody Griffin, and Audrey Wilson for assistance with figures.

References

1. Ollerton J, Winfree R, Tarrant S. How many flowering plants are pollinated by animals?
    Oikos. 2011;120: 321–326. doi:10.1111/j.1600-0706.2010.18644.x

2. Michener CD. The Bees of the World. Baltimore: Johns Hopkins University Press; 2007.

3. Morse R, Calderone NW. The value of honey bees as pollinators of U.S. Crops in 2000.
    Bee Cult. 2001;128. Available: http://agris.fao.org/agris-
    search/search.do?recordID=XE20122002449

4. Losey JE, Vaughan M. The Economic Value of Ecological Services Provided by Insects.
    BioScience. 2006;56: 311–323. doi:10.1641/0006-3568(2006)56[311:TEVOES]2.0.CO;2

5. Winfree R, Williams NM, Dushoff J, Kremen C. Native bees provide insurance against
    ongoing honey bee losses. Ecol Lett. 2007;10: 1105–1113. doi:10.1111/j.1461-
    0248.2007.01110.x

6. Garibaldi LA, Steffan-Dewenter I, Winfree R, Aizen MA, Bommarco R, Cunningham SA,
    et al. Wild Pollinators Enhance Fruit Set of Crops Regardless of Honey Bee Abundance.
    Science. 2013;339: 1608–1611. doi:10.1126/science.1230200

7. Greenleaf SS, Kremen C. Wild bees enhance honey bees' pollination of hybrid sunflower.
    Proc Natl Acad Sci. 2006;103: 13890–13895. doi:10.1073/pnas.0600929103

8. Tepedino VJ. The importance of bees and other insect pollinators in maintaining floral
    species composition. Gt Basin Nat Mem. 1979; 139–150.

9. Potts SG, Vulliamy B, Dafni A, Ne'eman G, Willmer P. Linking Bees and Flowers: How
    Do Floral Communities Structure Pollinator Communities? Ecology. 2003;84: 2628–2642.
    doi:10.2307/3450108

10. Cane JH, Minckley RL, Kervin LJ, Roulston TH, Williams NM. Complex responses within
    a desert bee guild (Hymenoptera : Apiformes) to urban habitat fragmentation. Ecol Appl.
    2006;16: 632–644. doi:10.1890/1051-0761(2006)016[0632:CRWADB]2.0.CO;2

11.   Kremen C, Williams NM, Bugg RL, Fay JP, Thorp RW. The area requirements of an
ecosystem service: crop pollination by native bee communities in California. Ecol Lett.
2004;7: 1109–1119. doi:10.1111/j.1461-0248.2004.00662.x

12.   Tscharntke T, Steffan-Dewenter I, Kruess A, Thies C. Characteristics of insect populations
on habitat fragments: A mini review. Ecol Res. 2002;17: 229–239. doi:10.1046/j.1440-
1703.2002.00482.x

13.   Öckinger E, Smith HG. Semi-natural grasslands as population sources for pollinating
insects in agricultural landscapes. J Appl Ecol. 2007;44: 50–59. doi:10.1111/j.1365-
2664.2006.01250.x

14.   Morandin LA, Kremen C. Hedgerow restoration promotes pollinator populations and
exports native bees to adjacent fields. Ecol Appl. 2013;23: 829–839. doi:10.1890/12-1051.1

15.   Chaplin-Kramer R, Dombeck E, Gerber J, Knuth KA, Mueller ND, Mueller M, et al. Global
malnutrition overlaps with pollinator-dependent micronutrient production. Proc R Soc B.
2014;281: 20141799. doi:10.1098/rspb.2014.1799

16.   Goulson D, Nicholls E, Botías C, Rotheray EL. Bee declines driven by combined stress
from parasites, pesticides, and lack of flowers. Science. 2015;347: 1255957.
doi:10.1126/science.1255957

17.   Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. Global
pollinator declines: trends, impacts and drivers. Trends Ecol Evol. 2010;25: 345–353.
doi:10.1016/j.tree.2010.01.007

18.   Gonzalez VH, Griswold T, Engel MS. Obtaining a better taxonomic understanding of
native bees: where do we start? Syst Entomol. 2013;38: 645–653. doi:10.1111/syen.12029

19.   Linsley EG. The Ecology of Solitary Bees. Hilgardia. 1958;27: 543–599.

20.   Michener CD. Biogeography of the bees. Ann Mo Bot Gard. 1979;66: 277–347.

21.   Cardoso P, Erwin TL, Borges PAV, New TR. The seven impediments in invertebrate
conservation and how to overcome them. Biol Conserv. 2011;144: 2647–2655.
doi:10.1016/j.biocon.2011.07.024

22.   Fischer AG. Latitudinal Variations in Organic Diversity. Evolution. 1960;14: 64–81.
doi:10.2307/2405923

23.   Minckley R. Faunal composition and species richness differences of bees (Hymenoptera:
Apiformes) from two north American regions. Apidologie. 2008;39: 176–188.
doi:10.1051/apido:2007062

24.   Minckley RL, Roulston TH, Williams NM. Resource assurance predicts specialist and
generalist bee activity in drought. Proc R Soc B Biol Sci. 2013;280.
doi:10.1098/rspb.2012.2703

25.   Williams N, Minckley R, Silveira F. Variation in native bee faunas and its implications for
detecting community changes. Conserv Ecol. 2001;5: 1–24.

26.   Bascompte J, Jordano P, Olesen J. Asymmetric coevolutionary networks facilitate
biodiversity maintenance. SCIENCE. 2006;312: 431–433. doi:10.1126/science.1123412

27.   Bascompte J, Jordano P, Melián CJ, Olesen JM. The nested assembly of plant–animal
mutualistic networks. Proc Natl Acad Sci. 2003;100: 9383–9387.
doi:10.1073/pnas.1633576100

28.   Kremen C. Managing ecosystem services: what do we need to know about their ecology?
Ecol Lett. 2005;8: 468–479. doi:10.1111/j.1461-0248.2005.00751.x

29.   Larsen TH, Williams NM, Kremen C. Extinction order and altered community structure
rapidly disrupt ecosystem functioning. Ecol Lett. 2005;8: 538–547. doi:10.1111/j.1461-
0248.2005.00749.x

30.   Williams NM, Crone EE, Roulston TH, Minckley RL, Packer L, Potts SG. Ecological and
life-history traits predict bee species responses to environmental disturbances. Biol
Conserv. 2010;143: 2280–2291. doi:10.1016/j.biocon.2010.03.024

31.   Aizen MA, Sabatino M, Tylianakis JM. Specialization and Rarity Predict Nonrandom Loss
of Interactions from Mutualist Networks. Science. 2012;335: 1486–1489.
doi:10.1126/science.1215320

32.   Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, et al. Historical
changes in northeastern US bee pollinators related to shared ecological traits. Proc Natl
Acad Sci. 2013;110: 4656–4660. doi:10.1073/pnas.1218503110

33.   Bommarco R, Biesmeijer JC, Meyer B, Potts SG, Pöyry J, Roberts SPM, et al. Dispersal
capacity and diet breadth modify the response of wild bees to habitat loss. Proc R Soc Lond
B Biol Sci. 2010; rspb20092221. doi:10.1098/rspb.2009.2221

34.   Memmott J, Craze PG, Waser NM, Price MV. Global warming and the disruption of plant–
pollinator interactions. Ecol Lett. 2007;10: 710–717. doi:10.1111/j.1461-
0248.2007.01061.x

35.   Griswold TL, Andres M, Andrus R, Garvin G, Keen K, Kervin L, et al. A survey of the rare
bees of Clark County, Nevada. Final Rep Nat Conserv Las Vegas NV. 1999; Available:
http://works.bepress.com/terry_griswold/63

36.   Marlin JC, LaBerge WE. The Native Bee Fauna of Carlinville, Illinois, Revisited After 75
Years:a Case for Persistence. Ecol Soc. 2007; Available: http://agris.fao.org/agris-
search/search.do?recordID=XE20122002329

37.   Messinger O, Griswold TL. A Pinnacle of bees. Fremontia. 2003;30: 32–40.

38. Messinger O. A survey of the bees of Grand Staircase-Escalante National Monument,
Southern Utah: Incidence, Abundance, and Community dynamics. Masters of Science, Utah
State University. 2006.

39. Roubik DW. Ups and downs in pollinator populations: When is there a decline? 2001;
Available: http://dlc.dlib.indiana.edu/dlc/handle/10535/3364

40. Wilson JS, Messinger OJ, Griswold T. Variation between bee communities on a sand dune
complex in the Great Basin Desert, North America: Implications for sand dune
conservation. J Arid Environ. 2009;73: 666–671.

41. Drons DJ. An Inventory of Native Bees (Hymenoptera: Apiformes) in the Black Hills of
South Dakota and Wyoming [Internet]. South Dakota State University. 2012. Available:
http://gfp.sd.gov/images/WebMaps/Viewer/WAP/Website/SWGSummaries/Drons%20201
2_an%20inventory%20of%20native%20Black%20Hills%20bees%20acknowledge.pdf

42. Giles V, Ascher JS. A survey of the bees of the Black Rock Forest preserve, New York
(Hymenoptera: Apoidea). J Hymenopt Res. 2006;15: 208–231.

43. Moldenke AR. California pollination ecology and vegetation types. Phytologia. 1976;34:
305–361.

44. Moldenke AR. Evolutionary history and diversity of the bee faunas of Chile and Pacific
North America. Wasmann J Biol. 1976;34: 147–178.

45. Meiners JM, Griswold TL, Evans EW. Native Bees of Pinnacles National Park: Diversity
Inventory and Plot Sampling Final Report and Sampling Manual. National Park Service;
Utah State University; 2015 Sep p. 76.

46. Meiners JM. Biodiversity, Community Dynamics, and Novel Foraging Behaviors of a Rich
Native Bee Fauna across Habitats at Pinnacles National Park, California. Masters of
Science, Utah State University. 2016.

47. LeBuhn G, Griswold T, Minckley R, Droege S, Roulston T, Cane J, et al. A standardized
method for monitoring bee populations–the bee inventory (BI) plot. 2003; Available:
http://cybercemetery.unt.edu/archive/nbii/20120111121317/http://online.sfsu.edu/~beeplot/
pdfs/Bee%20Plot%202003.pdf

48. Baldwin BG, Goldman DH. The Jepson Manual: Vascular Plants of California. University
of California Press; 2012.

49. R Core Team. R: A language and environment for statistical computing. https://www.R-
project.org. Vienna, Austria; 2015.

50. Gotelli NJ, Colwell RK. Quantifying biodiversity: procedures and pitfalls in the
measurement and comparison of species richness. Ecol Lett. 2001;4: 379–391.
doi:10.1046/j.1461-0248.2001.00230.x

51.   Gering JC, Crist TO. The alpha–beta–regional relationship: providing new insights into
local–regional patterns of species richness and scale dependence of diversity components.
Ecol Lett. 2002;5: 433–444. doi:10.1046/j.1461-0248.2002.00335.x

52.   Kuhlman M, Burrows S. Checklist of bees (Apoidea) from a private conservation property
in west-central Montana. Biodivers Data J.

53.   Deyrup M, Edirisinghe J, Norden B. The diversity and floral hosts of bees at the Archbold
Biological Station, Florida (Hymenoptera: Apoidea). Available: http://publikationen.ub.uni-
frankfurt.de/frontdoor/index/index/year/2013/docId/23868

54.   Frankie GW, Thorp RW, Hernandez J, Rizzardi M, Ertter B, Pawelek JC, et al. Native bees
are a rich natural resource in urban California gardens. Calif Agric. 2009;63: 113–120.
doi:10.3733/ca.v063n03p113

55.   USDA - National Agricultural Statistics Service - Honey Bee Surveys and Reports
[Internet]. [cited 20 Nov 2016]. Available:
https://www.nass.usda.gov/Surveys/Guide_to_NASS_Surveys/Bee_and_Honey/

56.   Kremen C, Ricketts T. Global Perspectives on Pollination Disruptions. Conserv Biol.
2000;14: 1226–1228.

57.   Wolf AT, Ascher JS. Bees of Wisconsin (Hymenoptera: Apoidea: Anthophila). Gt Lakes
Entomol. 2009;41.1: 129–168.

58.   DONOVAN BJ. INTERACTIONS BETWEEN NATIVE AND INTRODUCED BEES IN
NEW ZEALAND. N Z J Ecol. 1980;3: 104–116.

59.   Petanidou T, Kallimanis AS, Tzanopoulos J, Sgardelis SP, Pantis JD. Long-term
observation of a pollination network: fluctuation in species and interactions, relative
invariance of network structure and implications for estimates of specialization. Ecol Lett.
2008;11: 564–575. doi:10.1111/j.1461-0248.2008.01170.x

60.   Báldi A. Habitat heterogeneity overrides the species–area relationship. J Biogeogr. 2008;35:
675–681. doi:10.1111/j.1365-2699.2007.01825.x

61.   Tews J, Brose U, Grimm V, Tielbörger K, Wichmann MC, Schwager M, et al. Animal
species diversity driven by habitat heterogeneity/diversity: the importance of keystone
structures: Animal species diversity driven by habitat heterogeneity. J Biogeogr. 2004;31:
79–92. doi:10.1046/j.0305-0270.2003.00994.x

62.   Klausmeyer KR, Shaw MR. Climate Change, Habitat Loss, Protected Areas and the
Climate Adaptation Potential of Species in Mediterranean Ecosystems Worldwide. PLoS
ONE. 2009;4: e6392. doi:10.1371/journal.pone.0006392

63.   Matthews VI. Correlation of Pinnacles and Neenach Volcanic Formations and Their
Bearing on San Andreas Fault Problem. AAPG Bull. 1976;60: 2128–2141.

64. Tucker S, Knudsen K, Robertson J. Additional lichen collections from Pinnacles National
Monument, San Benito County, California. Bull Calif Lichen Soc. 2006;13: 8–11.

65. NPS. Nature - Pinnacles National Park (U.S. National Park Service) [Internet]. 2015 [cited
15 Nov 2015]. Available: http://www.nps.gov/pinn/learn/nature/index.htm

66. Debano SJ. Effects of livestock grazing on aboveground insect communities in semi-arid
grasslands of southeastern Arizona. Biodivers Conserv. 2006;15: 2547.
doi:10.1007/s10531-005-2786-9

67. Winfree R, Aguilar R, Vázquez DP, LeBuhn G, Aizen MA. A meta-analysis of bees'
responses to anthropogenic disturbance. Ecology. 2009;90: 2068–2076. doi:10.1890/08-
1245.1

68. Iles DT, Williams NM, Crone EE. Source-sink dynamics of bumblebees in rapidly
changing landscapes. J Appl Ecol. 2018;55: 2802–2811. doi:10.1111/1365-2664.13175

69. Griffin SR, Bruninga-Socolar B, Kerr MA, Gibbs J, Winfree R. Wild bee community
change over a 26-year chronosequence of restored tallgrass prairie. Restor Ecol. 2017;
doi:10.1111/rec.12481

70. Albrecht M, Riesen M, Schmid B. Plant-pollinator network assembly along the
chronosequence of a glacier foreland. Oikos. 2010;119: 1610–1624. doi:10.1111/j.1600-
0706.2010.18376.x

71. Bohart G, Knowlton G. The Bees of Curlew Valley (Utah and Idaho). Proc Utah Acad Sci
Arts Lett. 1973; Available: http://digitalcommons.usu.edu/piru_pubs/790

72. Griswold T, Parker F, Tepedino V. The bees of the San Rafael Desert: Implications for the
bee fauna of the Grand Staircase-Escalante National Monument. In: Hill LM, editor.
Learning from the land: Grand Staircase Escalante National Monument Science
Symposium Proceedings. 1998. pp. 175–186.

73. Brown JW, Bahr SM. The Insect (Insecta) Fauna of Plummers Island, Maryland: Brief
Collecting History and Status of the Inventory. Bull Biol Soc Wash. 2008;15: 54–64.
doi:10.2988/0097-0298(2008)15[54:TIIFOP]2.0.CO;2

74. Grundel R, Jean RP, Frohnapple KJ, Gibbs J, Glowacki GA, Pavlovic NB. A Survey of
Bees (Hymenoptera: Apoidea) of the Indiana Dunes and Northwest Indiana, USA. J Kans
Entomol Soc. 2011;84: 105–138. doi:10.2317/JKES101027.1

75. Tepedino VJ, Stanton NL. Diversity and Competition in Bee-Plant Communities on Short-
Grass Prairie. Oikos. 1981;36: 35–44. doi:10.2307/3544376

76. Rhoades PR, Griswold T, Ikerd H, Waits L, Bosque-Pérez N, Eigenbrode S. The native bee
fauna of the Palouse Prairie (Hymenoptera: Apoidea). J Melittology. 2017;0: 1–20.
doi:10.17161/jom.v0i66.5703

77. Rust R, Menke A, Miller D. A biogeographic comparison of the bees, sphecid wasps, and
mealybugs of the California Channel Ilsnads (Hymenoptera, Homoptera). Entomology of
the California Channel Islands: proceedings of the first symposium. Santa Barbara, CA:
Santa Barbara Museum of Natural History; 1985. pp. 22–59.

78. Wilson JS, Wilson LE, Loftis LD, Griswold T. The Montane Bee Fauna of North Central
Washington, USA, with Floral Associations. West North Am Nat. 2010;70: 198–207.
doi:10.3398/064.070.0206

79. Smith BA, Brown RL, Laberge W, Griswold T. A Faunistic Survey of Bees (Hymenoptera:
Apoidea) in the Black Belt Prairie of Mississippi. J Kans Entomol Soc. 2012;85: 32–47.
doi:10.2317/JKES111025.1

80. Michener CD. Bees of a Limited Area in Southern Mississippi (Hymenoptera; Apoidea).
Am Midl Nat. 1947;38: 443–455. doi:10.2307/2421575

Supporting Information

S1 Fig. Relationship between floral richness (FR) and either (a) bee richness (BR) or (b)

bee abundance (BA, square-root transformed) at the plot-sample level (N=150) within

Pinnacles National Park (2011-12). Shown with power-law model (black line; (a) BR =

$\exp(2.79 + 0.38 \ast \log(FR))$; $R^2 = 0.37$, $p < 0.01$; (b) BA = $\exp(2.26 + 0.23 \ast \log(FR))$; $R^2 = 0.16$,

$p < 0.01$).

Bee Taxonomy				Early Inventory				Bowls	Recent Inventory			
				1996	1997	1998	1999	2002	2011	2012	Prop. years present	New/ Absent in 2011-12
Family	**Genus**	**Subgenus**	**Species**									
Andrenidae	Ancylandrena		atoposoma	S							0.14	A
	Andrena	(Anchandrena)	quercina			R			R	R	0.43	
		(Belandrena)	nemophilae			R			S	R	0.43	
			palpalis		R	R					0.29	A
		(Cremnandrena)	anisochlora			C			C	R	0.43	
		(Dasyandrena)	cristata			R					0.14	A
		(Derandrena)	arctostaphylae			S		S			0.29	A
			californiensis			R		S	S	R	0.57	
			n. sp.	S	S	R			R	C	0.71	
			vandykei			C		S	S		0.43	
			viridissima						R		0.14	N
		(Diandrena)	apasta					S			0.14	A
			chalybioides							S	0.14	N
			cuneilabris							S	0.14	N
			lewisorum	S	R	S	C		C	C	0.86	
			nothocalaidis			S					0.14	A
			puthua		R	C			C	R	0.57	
			subchalybea			R	R	R	C	C	0.71	
		(Erandrena)	principalis			R					0.14	A
			astragali		S	R			S	R	0.57	
			auricoma	R	R	C	R		C		0.71	
			caerulea			C			R	R	0.43	
			chlorura		R	C		R	C	C	0.71	
			dissimulans		R	C			C	C	0.57	
			misella			S					0.14	A
			nigrocaerulea	S	R	R			S	R	0.71	
			suavis			R			C	C	0.43	

Genus	Subgenus	Species									
		subdepressa			R			R	S	0.43	
	(Genyandrena)	mackieae		R	C			R		0.43	
	(Hesperandrena)	baeriae		S	S			R	R	0.57	
		compositarum						S	S	0.29	N
		escondida							R	0.14	N
	(Holandrena)	cressonii ssp. infasciata		S			S			0.29	A
	(Melandrena)	cerasifolii	C	C	C	S	S	C	C	1	
		aff. cerasifolii		C	C	C		C	C	0.71	
		perimelas					S			0.14	A
		sola	R	C	C	C		C	C	0.86	
	(Micrandrena)	annectens			R					0.14	A
		chlorogaster		R	C		R	C	C	0.71	
		aff. ishii			C					0.14	A
		microchlora		R	C	S		R	R	0.71	
		piperi		S	R		S	R	R	0.71	
	(Nemandrena)	crudeni			C		S	C	C	0.57	
		subnigripes			S					0.14	A
	(Oligandrena)	macrocephala	R	C	C		C	C	C	0.86	
	(Parandrena)	concinnula			S			R		0.29	
	(Pelicandrena)	atypica	S	R	C	R	R	R	R	1	
	(Plastandrena)	prunorum							S	0.14	N
	(Psammandrena)	congrua		R	R			R	R	0.57	
	(Ptilandrena)	pallidiscopa			R				R	0.29	
	(Scaphandrena)	lomatii			S					0.14	A
		plana			R					0.14	A
		santaclarae		S	R			R	R	0.57	
	(Scoliandrena)	cryptanthae	R					R		0.29	
		osmioides			R			R	S	0.43	
	(Scrapteropsis)	biareola			R					0.14	A
	(Simandrena)	angustitarsata	R	R	C	C		C	C	0.86	
		hypoleuca						S	R	0.29	N
		orthocarpi			R			R	R	0.43	
		pallidifovea	S	R	R	R		R	R	0.86	
	(Thysandrena)	candida	C	C	C	C	R	C	C	1	
		knuthiana					S	C	C	0.43	
		vierecki		S	S					0.29	A
		w-scripta	R	R	C	C			S	0.71	
	(Trachandrena)	fuscicauda			S	R				0.29	A
		semipunctata		R	S			S		0.43	
	(Tylandrena)	subaustralis						S		0.14	N
		subtilis			S					0.14	A
		waldmerei			R					0.14	A
Calliopsis	(Micronomadopsis)	fracta	R	S	R			S		0.57	
		helianthi					R	S		0.29	
		mellipes		S				R		0.29	
		trifolii	R			R				0.29	A

Family	Genus	Subgenus	Species									
		(Nomadopsis)	anthidia	S	R	R		S	C		0.71	
			obscurella	R	C	R	R	C	C	R	1	
			zonalis	S	S			S			0.43	A
			smithi	C	C	C	C			S	0.71	
	Macrotera	(Macroteropsis)	arcuata			S			S	R	0.43	
	Panurginus		gracilis			C	R	C	C	C	0.71	
			melanocephalus		R	R			C	C	0.57	
			nigrellus	C	C	C		R	C	C	0.86	
			aff. occidentalis					S			0.14	A
	Perdita	(Hesperoperdita)	trisignata ssp. ornata	R	C	C	C		S	S	0.86	
		(Perdita)	claypolei ssp. limatula		C	C					0.29	A
			hirticeps ssp. hirticeps					S		R	0.29	
			aff. holoxantha						S		0.14	N
			isocomae		S						0.14	A
			jucunda		R		S				0.29	A
			aff. linsleyi	R	R	R	S		R		0.71	
			rhois ssp. reducta	C	C	C	S			R	0.71	
			salicis ssp. personata						R		0.14	N
		(Pygoperdita)	aureovittata ssp. stenozona		S						0.14	A
			californica			R					0.14	A
			distropica	R	C	C	C	C	C	C	1	
			micheneri ssp. micheneri		C	R		S	R		0.57	
			montereyensis	C	C	C	C	S	C	C	1	
			nitens	C	C	R			C	R	0.71	
Apidae	Anthophora	(Anthophoroides)	californica	R	C	R	R	S	R	R	1	
		(Heliophila)	columbariae			R					0.14	A
			curta	R	R	S	R		R	R	0.86	
			estebana		S						0.14	A
			flavocincta					S			0.14	A
		(Lophanthophora)	pacifica		R	C			S		0.43	
		(Melea)	bomboides	S							0.14	A
		(Mystacanthophora)	urbana	R	C	C	C	S	C	C	1	
		(Paramegilla)	centriformis	R	C	R	C		R		0.71	
		(Pyganthophora)	crotchii	R	C	R	R	S	C	C	1	
			edwardsii	R	R	R			C	C	0.71	
			platti	C	C	C	R		C	C	0.86	
	Anthophorula	(Anthophorisca)	nitens		C	C			C	C	0.57	
		(Anthophorula)	albicans		C	C		R	S	C	0.71	
	Apis		mellifera	R	C	C	C	C	C	C	1	
	Bombus	(Crotchiibombus)	crotchii		C	C	S				0.43	A
		(Fervidobombus)	californicus	R	C	C	R	R	R	R	1	
		(Pyrobombus)	caliginosus		C				S	S	0.43	
			melanopygus	R	C	C	R	R	C	R	1	
			vandykei		R	R			R		0.43	
			vosnesenskii	R	C	C	S	S	C	R	1	
	Brachynomad	(Melanomada)	melanantha							R	0.14	N

a

Genus	Subgenus	Species									
Centris	(Paracentris)	aff. californica		C	C	R				0.43	A
Ceratina	(Ceratina)	arizonensis	C	C	C	C	C	C	C	1	
	(Euceratina)	dallatorreana			S		R		C	0.43	
	(Zadontomerus)	acantha	S	C	R		R	C	C	0.86	
		hurdi	S	R	R		R	C	C	0.86	
		nanula	R	C	C	C	C	C	C	1	
		aff. nanula	S	R	S		S	R	R	0.86	
		pacifica		R	S			S	R	0.57	
		punctigena	S	C	R	C		R	S	0.86	
		sequoiae	R	C	C	C	R	C	C	1	
		tejonensis		C	R			R		0.43	
		timberlakei		R	R	R		R	S	0.71	
Diadasia		aff. ochracea		S	S		S	R	C	0.71	
		angusticeps		R	R	C	R	C	C	0.86	
		australis					S	S		0.29	
		bituberculata	R	C	R	C	C	C	C	1	
		consociata							R	0.14	N
		laticauda		C	C	R	R	C	C	0.86	
		nigrifrons						S		0.14	N
		nitidifrons		R	C	R	R	C	C	0.86	
		rinconis							R	0.14	N
Epeolus		americanus	S	R			S	R	R	0.71	
		compactus			R	S			S	0.43	
		mesillae						S	R	0.29	N
		minimus				S	S		S	0.43	
Eucera	(Synhalonia)	actuosa		S	R	R	C	C	C	0.86	
		amsinckiae	R	R	R		R	R	R	0.86	
		cordleyi		R	C		S	C	R	0.71	
		delphinii		C	R	S	S	R	R	0.86	
		dorsata		C	R	R	S	C	C	0.86	
		edwardsii	S	R	R		S		R	0.71	
		lunata			R			R	R	0.43	
		venusta ssp. carinata	C	C	C	R		R	R	0.86	
		virgata			C	S	S	C	C	0.71	
Habropoda		dammersi			R					0.14	A
		depressa	S	C	C	R	S	C	C	1	
		tristissima	R	C	R	S	R	R	C	1	
Melecta	(Melecta)	pacifica		R	R			R		0.43	
		separata			S	R	R	R	C	0.71	
	(Melectomimus)	edwardsii		R				S		0.29	
Melissodes	(Callimelissodes)	clarkiae		R				R	R	0.43	
		composita							S	0.14	N
		lupina		R	C		R	C	C	0.71	
		lustra		S						0.14	A
		n. sp. 1		R		R				0.29	A

		n. sp. 2	R	R	R	S	R	C		0.86	
		nigracauda		S						0.14	A
		plumosa	R				R	C		0.43	
		stearnsi	R	S		C	C	C		0.71	
	(Eumelissodes)	paulula	R							0.14	A
		velutina						R		0.14	N
	(Melissodes)	tepida						C		0.14	N
Neopasites	(Micropasites)	sp. 4						S		0.14	N
Nomada	(Centrias)	crotchii spp. crotchii	R	S						0.29	A
		crotchii ssp. nigrior		C			R	R		0.43	
		sp. A	S	R						0.29	A
	(Holonomada)	edwardsii spp. edwardsii	R	R				S		0.43	
	(Nomada)	sp. A	R	R	S		R	S		0.71	
		sp. AA						S		0.14	N
		sp. B	S	S	S		R	R		0.71	
		sp. BB				S	R	R		0.43	
		sp. CC				S		S		0.29	
		sp. D	C	C	C	S				0.57	A
		sp. DD					S			0.14	N
		sp. E	C	R	S	R	R	R		0.86	
		sp. EE					R			0.14	N
		sp. F	S		R			S		0.43	
		sp. FF						S		0.14	N
		sp. G					R	R		0.29	N
		sp. GG			S			R		0.29	
		sp. HH				S	S			0.29	A
		sp. I				S				0.14	A
		sp. II					R		R	0.29	
		sp. J			S		S			0.29	
		sp. Q		S	S		S			0.43	
		sp. R		R	R					0.29	A
		sp. S		S						0.14	A
		sp. T		S						0.14	A
		sp. U	C	C			R	C		0.57	
		sp. V	S	R	R		R	R	R	0.86	
		sp. W	C	C			C	C		0.57	
		sp. X			S		R	R		0.43	
		sp. Y		R		S				0.29	A
		sp. Z			C					0.14	A
Oreopasites		aff. hurdi n.sp.	S	R	S		S	S		0.71	
		vanduzeei	R	R	S	S		S		0.71	
Peponapis	(Peponapis)	pruinosa						R		0.14	N
Townsendiella		ensifera	R	C	C		R			0.57	
		rufiventris		S						0.14	A
Triepeolus		heterurus						R		0.14	N
		melanarius						S		0.14	N

Family	Genus	Subgenus	Species									
			sp. P1						R	S	0.29	N
			sp. P2						R		0.14	N
			sp. P3							R	0.14	N
			sp. P4		R						0.14	A
			timberlakei		R	R	R		S	R	0.71	
			aff. timberlakei					S	R		0.29	
	Xeromelecta	(Melectomorpha)	californica	R	C	R	C		R	R	0.86	
		(Xeromelecta)	larreae	S							0.14	A
	Xylocopa	(Notoxylocopa)	tabaniformis ssp. orpifex	R	C	R	R	S	R	R	1	
Colletidae	Colletes consors		californicus							S	0.14	N
			consors ssp. pascoensis	C	C	R	R		C	R	0.86	
	daleae		aff. algarobiae	S	C	C	R		R	C	0.86	
	simulans		fulgidus ssp. fulgidus		R	R	R			R	0.57	
			simulans ssp. nevadensis		S						0.14	A
			slevini	R	C	C	C		R	R	0.86	
	Hylaeus	(Cephalylaeus)	nunenmacheri			S	R				0.29	A
		(Hylaeus)	bisinuatus		S						0.14	A
			conspicuus				R	R		R	0.43	
			granulatus		C	C	C	S	R	R	0.86	
			mesillae ssp. cressoni	R	C	C	C	S	C	C	1	
			rudbeckiae	R					R	R	0.43	
			verticalis	R	C	C	C		R	R	0.86	
		(Paraprosopis)	aff. cookii n.sp.		R	R	R				0.43	A
			calvus	R		C	C		R	C	0.71	
			coloradensis		C	C	R				0.43	A
			n. sp.2		C	R	R		R		0.57	
			nevadensis	C	C	C	C		R	C	0.86	
			polifolii		C	C	C		R		0.57	
		(Prosopis)	aff. episcopalis	C	C	C	C		R	R	0.86	
		(Spatulariella)	punctatus				S				0.14	A
Halictidae	Agapostemon	(Agapostemon)	femoratus					R	R		0.29	
			texanus	R	R	R	C	C	C	C	1	
	Augochlorella	(Augochlorella)	pomoniella		R	R				R	0.43	
	Conanthalictus	(Phaceliapis)	bakeri		S	R	S		R	S	0.71	
			seminiger				S			S	0.29	
	Dufourea		dentipes						S		0.14	N
			leachi	S	S	C	R	R	R	C	1	
			mulleri		R	R			R		0.43	
			rhamni	R	R	R				S	0.57	
			sandhouseae		C	C			C	C	0.57	
			sparsipunctata	C	C	R	R	C	C	C	1	
			virgata	C	R	S	S		R	R	0.86	
	Halictus	(Nealictus)	farinosus	C	C	C	C	C	C	C	1	
		(Odontalictus)	ligatus				R	C	C	C	0.57	
		(Protohalictus)	rubicundus					S			0.14	A
		(Seladonia)	tripartitus	C	C	C	C	C	C	C	1	

Family	Genus	Subgenus	Species									
	Lasioglossum	(Dialictus)	albohirtum				R		S	R	0.43	
			brunneiventre		R	R	R		C	C	0.71	
			diversopunctatum			S			R		0.29	
			hudsoniellum						S	R	0.29	N
			imbrex	R	C	C	C		C	C	0.86	
			cf. impavidum							S	0.14	N
			incompletum	R	R	C	C		C	C	0.86	
			megastictum			S			R		0.29	
			nevadense	R	C	C	C		C	C	0.86	
			perichlarum						R		0.14	N
			petrellum		S						0.14	A
			punctatoventre	R	C	C	C		C	C	0.86	
			aff. ruidosense	R	R		R				0.43	A
		(Evylaeus)	argemonis	R	C	C	C		C	C	0.86	
			giffardi					S	C	R	0.43	
			robustum		S	R	C		C	C	0.71	
		(Hemihalictus)	aspilurum		S		S		R	R	0.57	
			glabriventre	R	R	R	C		C	C	0.86	
			kincaidii	R	R	S	S		R	R	0.86	
			ovaliceps		S	R			S		0.43	
			ruficorne	R	R	C	C		C	R	0.86	
			sequoiae	S		R	C		C	C	0.71	
		(Lasioglossum)	egregium		S	R	R		C	C	0.71	
			mellipes					R	C	R	0.43	
			sisymbrii	R	C	C	C	C	C	C	1	
			titusi			S	R		C	C	0.57	
		(Sphecodogastra)	allonotum			S			R	R	0.43	
			aff. avalonense		S	R	C		C	C	0.71	
			miguelense		S		R		R	C	0.57	
			nigrescens	C	C	C	C		C	C	0.86	
			sp. 16				R		C	C	0.43	
	Micralictoides		altadenae		S				R		0.29	
			ruficaudus	R	R	S			C	R	0.71	
	Sphecodes		arvensiformis	S	R			S	S	R	0.71	
			sp. B	R	C	C	R		R	R	0.86	
			sp. C	R							0.14	A
			sp. D	R		S	R		R	R	0.71	
			sp. E	S	R	R	C	R	C	C	1	
			sp. F	S	R						0.29	A
			sp. I		S						0.14	A
			sp. J		R						0.14	A
			sp. K		S						0.14	A
			sp. L		S	R	C		R		0.57	
			sp. M							S	0.14	N
Megachilidae	Anthidiellum	(Loyalanthidium)	robertsoni		C	C			R		0.43	
	Anthidium	(Anthidium)	collectum	R	C	C	C	R	R	R	1	

Genus	Subgenus	Species									
		edwardsii							R	0.14	N
		jocosum							R	0.14	N
		maculosum		S						0.14	A
		mormonum	S	R	S	R				0.57	A
		pallidiclypeum	R	S	R	S	S			0.71	A
		utahense	C	C	C	C	C	C	C	1	
	(Callanthidium)	illustre		C	R	R		S	R	0.71	
Ashmeadiella	(Arogochila)	aff. salviae n. sp. 2						S		0.14	N
		australis		R	R	R			S	0.57	
		salviae	R	C	C	R	S	R	C	1	
		timberlakei	C	C	C	C		R	R	0.86	
	(Ashmeadiella)	altadenae						R	R	0.29	N
		aridula					R		C	0.29	
		bucconis		C	R				R	0.43	
		cactorum ssp. basalis		R					R	0.29	
		californica ssp. californica	R	C	C	C	R	R	C	1	
		difugita ssp. emarginatula		R	R			S	R	0.57	
		femorata		R		C				0.29	A
		foveata	S	C	C	C		S		0.71	
		gillettei ssp. cismontanica		S	S					0.29	A
		meliloti		C	C					0.29	A
		pronitens		S						0.14	A
		aff. rufitarsis	R	R	R	R				0.57	A
		sonora		R	S		S		R	0.57	
		titusi		C	R	S			S	0.57	
Atoposmia	(Atoposmia)	n. sp. 2	R	R	S			S		0.57	
		pycnognatha		C	R	S				0.43	A
	(Eremosmia)	hemizoniae						S		0.14	N
	(Hexosmia)	copelandica ssp. copelandica	R	C	R	R	S	R	R	1	
Chelostoma	(Chelostoma)	aff. minutum n.sp.		R	R	R			S	0.57	
		californicum	C	C	C	C	S	C	R	1	
		cockerelli	C	C	C	C	R	C	C	1	
		incisulum	C	C	C	C		C	R	0.86	
		marginatum ssp. incisuloides	C	C	C	R		C	R	0.86	
		phaceliae	C	C	C	C		C	R	0.86	
		tetramerum	R	C	C	R		S	R	0.86	
Coelioxys	(Boreocoelioxys)	octodentata		R	R	S		S		0.57	
	(Coelioxys)	hirsutissima						R		0.14	N
		serricaudata	S	R	R	R	S	R	R	1	
	(Cyrtocoelioxys)	gilensis		S						0.14	A
		gonaspis		R						0.14	A
Dianthidium	(Dianthidium)	dubium ssp. dilectum		C	C	R	R	R	C	0.86	
		parvum ssp. schwarzi					R	R	R	0.43	
		pudicum ssp. consimile	S	S	S	R	R	R	C	1	
		singulare		S						0.14	A

Genus	Subgenus	Species									
		ulkei ssp. ulkei		R	S		S		R	0.57	
Dioxys		aurifusca	S							0.14	A
		pacifica ssp. pacifica			S		S		R	0.43	
		pomonae ssp. pomonae	R	R	R	R		C	R	0.86	
		producta ssp. cismontanica	R	R	R	R		R	S	0.86	
Heriades	(Neotrypetes)	occidentalis		C	C					0.29	A
Hoplitis	(Acrosmia)	aff. emarginata		C	R	R		S		0.57	
	(Alcidamea)	colei	R	C	R	C	R	C	C	1	
		grinnelli	R	C	R	R			R	0.71	
		producta ssp. bernardina		C				C	S	0.43	
		producta ssp. gracilis	C	C	C	C	R	C	R	1	
		sambuci	S	C	R	S			S	0.71	
	(Cyrtosmia)	hypocrita	S	C	C	R		R	R	0.86	
	(Hoplitina)	bunocephala	S		S					0.29	A
		howardi	C	C	C	C		R	R	0.86	
	(Monumetha)	albifrons ssp. maura	C	C	C	C	C	C	C	1	
		fulgida ssp. platyura	C	C	C	C	R	C	R	1	
	(Penteriades)	remotula	R	C	R		S	R	S	0.86	
	(Proteriades)	cryptanthae	R	R			S			0.43	A
		jacintana	C	C	S	R		R		0.71	
		nanula	R	C	S	R		R		0.71	
		seminigra	S	C	R	S		S		0.71	
		semirubra	R	R	R	S	R	R	R	1	
Megachile	(Argyropile)	parallela		S						0.14	A
	(Chelostomoides)	angelarum		C	C			R	R	0.57	
		davidsoni		C				S	S	0.43	
		exilis							R	0.14	N
		spinotulata		C	R					0.29	A
	(Eutricharaea)	apicalis		C	R		R	S	C	0.71	
	(Litomegachile)	coquilletti	S	C	C	S	R	C	C	1	
		gentilis		R	S				R	0.43	
		lippiae							R	0.14	N
		onobrychidis			R	S	R	R	C	0.71	
		texana	S	R	R	R				0.57	A
	(Megachile)	montivaga		R			S	S	R	0.57	
	(Megachiloides)	gravita		R		C	R	C	C	0.71	
		pascoensis	R	C	R	C	R	C	C	1	
		pseudonigra	S	R				R		0.43	
		subnigra ssp. angelica			R		R	C	R	0.57	
	(Sayapis)	fidelis		S	R				S	0.43	
		frugalis ssp. pseudofrugalis		C	C			S	C	0.57	
		inimica ssp. jacumbensis		R						0.14	A
		newberryae		R	R					0.29	A
Osmia	(Acanthosmioides)	nigrifrons		S		R				0.29	A
		nigrobarbata	R	S	S	S	S	R	S	1	
		odontogaster	R	C	C	R	S	R	C	1	

Genus	Subgenus	species								ratio	
		sedula	S	C	R	R	C	R	S	1	
	(Cephalosmia)	californica	R	R	S	S		R	R	0.86	
		montana ssp. quadriceps	R	R	R	C	S	C	R	1	
	(Euthosmia)	glauca	R	C	C	C	C	C	C	1	
	(Helicosmia)	coloradensis		C	C	R		R	R	0.71	
		texana		C	S	R	R	R	R	0.86	
	(Melanosmia)	aglaia	R	C	C	C	C	C	C	1	
		atrocyanea	R	C	C	C	C	C	C	1	
		brevis	C	C	C	C	R	C	C	1	
		calla	R	C	C	C	R	C	C	1	
		cara	C	C	C	C	S	C	C	1	
		clarescens	R	C	C	R		R	R	0.86	
		cyanella	C	C	C	C	R	C	C	1	
		cyanopoda	S	C	R	R			S	0.71	
		densa	C	C	C	R	R	C	C	1	
		gabrielis	C	C	C	C	R	C	C	1	
		gaudiosa					R	C	R	0.43	
		granulosa	R	C	C	C	R	C	C	1	
		aff. hesperos						S		0.14	N
		inurbana			R	R	C	C	C	0.71	
		kincaidii	C	C	C	C	R	C	C	1	
		laeta	R	C	C	R	R	C	C	1	
		malina			S					0.14	A
		melanopleura	C	C	C	C	R	C	R	1	
		pusilla	R	C		R	R	R	R	0.86	
		aff. pusilla						S		0.14	N
		raritatis	R	R	R	R		R	R	0.86	
		regulina	R	C	C	C	C	C	C	1	
		rostrata				R				0.14	A
		sp. P1					R		R	0.29	
		tristella		R	S	R				0.43	A
		vandykei		R	R	R				0.43	A
		visenda	C	C	C	R	R	C	C	1	
	(Mystacosmia)	nemoris		R	C	C	C	C	C	0.86	
	(Osmia)	lignaria ssp. propinqua	R	C	C	S	R	C	C	1	
		ribifloris ssp. biedermannii		S	S					0.29	A
	(Pyrosmia)	nigricollis							S	0.14	N
	(Trichinosmia)	latisulcata		R	S	R		C	S	0.71	
Protosmia	(Chelostomopsis)	rubifloris	C	C	C	C	R	C	C	1	
Stelis	(Protostelis)	anthidioides	S		R		S	S		0.57	
		hurdi		R	R	R	R	R		0.71	
	(Stelis)	aff. foederalis n.sp.		S		R		R		0.43	
		ashmeadiellae	R	R	R			S	S	0.71	
		calliphorina	R							0.14	A
		chemsaki							R	0.14	N
		cockerelli				R		R	R	0.43	

Family	Genus	Subgenus	species									
			interrupta				S			S	0.29	
			lateralis		R	R	R		R		0.57	
			micheneri	S	R		S		R		0.57	
			montana	R	R	R	R	S	R	R	1	
			nigriventris		R	R					0.29	A
			occidentalis	S	R					S	0.43	
			subemarginata	R	R	R	R		S		0.71	
	Trachusa	(Heteranthidium)	timberlakei		R	C	C	R	R	R	0.86	
		(Trachusomimus)	perdita		C	C	C	C	C	R	0.86	
Melittidae	Hesperapis	(Amblyapis)	ilicifoliae	C	C	C	C	C	C	C	1	
		(Panurgomia)	regularis	C	C	C	C	C	C	C	1	

S2 Table. **Floral taxa visited by bees at Pinnacles National Park (unique groups, identified**

to lowest possible level), and their relative popularity by year. Plants are marked with "R" for

rare if bee visits were fewer than 10 in that year, with "U" for uncommon if bee visits ranged

between 10-100, and "C" for common when over 100 bees were collected on that plant. The last

row sums the plant taxa on which bees were collected per year. Dashed vertical line marks 2002

collection as separate from original 1996-9 study, and prior to the current study.

Plant Name	Early Inventory				Bowls	Recent Inventory	
	1996	1997	1998	1999	2002	2011	2012
Alliaceae Allium fimbriatum						R	
Alliaceae Allium lacunosum		U	U	R			
Alliaceae Allium lacunosum var.micranthum							R
Alliaceae Allium sp.	U	U	R				
Anacardiaceae Toxicodendron diversilobum						R	
Apiaceae Anthriscus caucalis						R	
Apiaceae Apiaceae sp.		R	U				
Apiaceae Apiaceae sp. (yellow)			R			R	
Apiaceae Lomatium dasycarpum							R
Apiaceae Lomatium sp.		U	R				R
Apiaceae Lomatium utriculatum			R			U	R
Apiaceae Perideridia californica			R				
Apiaceae Sanicula crassicaulis						R	
Apiaceae Sanicula sp.		R					
Apiaceae Sanicula tuberosa						R	R
Asclepiadaceae Asclepias sp.			U				
Asteraceae Achillea millefolium		R				R	R
Asteraceae Agoseris grandiflora						R	
Asteraceae Agoseris sp.		R					
Asteraceae Anaphalis margaritacea		R					
Asteraceae Asteraceae sp.		R	U				
Asteraceae Asteraceae sp. (yellow)			R				
Asteraceae Baccharis pilularis		U	U				
Asteraceae Baccharis salicifolia	U	R		U		U	U
Asteraceae Carduus tenuiflorus		U					
Asteraceae Centaurea melitensis						R	R
Asteraceae Centaurea solstitialis		U	U		R		R
Asteraceae Chaenactis glabriuscula		R				C	U
Asteraceae Cirsium occidentale		R	U	R		R	R
Asteraceae Cirsium sp.		U	R				

Species							
Asteraceae Cirsium vulgare		R	R				
Asteraceae Erigeron foliosus		U					
Asteraceae Erigeron foliosus var.foliosus							R
Asteraceae Erigeron petrophilus		U					
Asteraceae Eriophyllum confertiflorum		U	U	U		R	R
Asteraceae Eriophyllum lanatum			R				
Asteraceae Eriophyllum multicaule							R
Asteraceae Eriophyllum sp.		U					
Asteraceae Euthamia occidentalis		R					
Asteraceae Gnaphalium bicolor						R	
Asteraceae Gnaphalium californicum						R	R
Asteraceae Hemizonia lobbii		U					
Asteraceae Heterotheca sessiliflora		U					
Asteraceae Hypochaeris glabra						R	
Asteraceae Hypochaeris radicata						U	R
Asteraceae Lasthenia californica	R	U	C			U	U
Asteraceae Layia hieracioides		R					
Asteraceae Lessingia tenuis				U			
Asteraceae Madia sp.		R					
Asteraceae Malacothrix californica						R	
Asteraceae Microseris douglasii						R	
Asteraceae Packera breweri						R	
Asteraceae Pectis papposa		R					
Asteraceae Senecio flaccidus		R					
Asteraceae Senecio sp.		U					
Asteraceae Stephanomeria virgata ssp.pleurocarpa			R				
Asteraceae Wyethia helenioides	R					U	U
Asteraceae Wyethia sp.				R			
Boraginaceae Amsinckia menziesii	U			R	R	U	U
Boraginaceae Amsinckia menziesii var.menziesii		U	U	R		R	
Boraginaceae Amsinckia sp.					U		
Boraginaceae Cryptantha sp.	U	C	U	R			R
Boraginaceae Emmenanthe penduliflora			R	U			
Boraginaceae Eriodictyon sp.			R				
Boraginaceae Eriodictyon tomentosum	U	C	U	C		U	U
Boraginaceae Heliotropium curassavicum		U				U	U
Boraginaceae Nemophila menziesii var.integrifolia		U	U				
Boraginaceae Nemophila menziesii var.menziesii						R	R
Boraginaceae Phacelia brachyloba		U					
Boraginaceae Phacelia californica		U					
Boraginaceae Phacelia distans		U	U				R
Boraginaceae Phacelia imbricata		U	U	U		R	
Boraginaceae Phacelia malvifolia			R	U			
Boraginaceae Phacelia ramosissima		U	U	R			
Boraginaceae Phacelia ramosissima var.ramosissima						R	
Boraginaceae Phacelia sp.	U	C	U	U			R
Boraginaceae Phacelia sp. (white)	U						
Boraginaceae Pholistoma auritum	U	C	C	R			R
Boraginaceae Pholistoma auritum var.auritum						U	R
Boraginaceae Pholistoma membranaceum			U			R	R
Boraginaceae Plagiobothrys canescens						U	
Boraginaceae Plagiobothrys nothofulvus			C				
Boraginaceae Plagiobothrys sp.		R	U		U		R
Brassicaceae Brassica nigra		C	U	U			

Species							
Brassicaceae Brassicaceae sp.					R		
Brassicaceae Cardamine californica			U				
Brassicaceae Cardamine californica var.californica			U			R	
Brassicaceae Erysimum capitatum var.capitatum						R	
Brassicaceae Erysimum sp.		R					
Brassicaceae Hirschfeldia incana						C	C
Brassicaceae Rorippa nasturtium-aquaticum			U	R			
Brassicaceae Thysanocarpus curvipes			C				
Brassicaceae Thysanocarpus laciniatus							R
Caprifoliaceae Lonicera hispidula			U				
Caprifoliaceae Lonicera sp.			U				R
Caprifoliaceae Lonicera subspicata var.denudata						R	
Caprifoliaceae Sambucus mexicana						R	U
Chenopodiaceae Chenopodium californicum						R	
Convolvulaceae Calystegia collina						R	U
Convolvulaceae Calystegia collina ssp.venusta						R	
Convolvulaceae Calystegia purpurata							R
Convolvulaceae Calystegia sp.			R				
Convolvulaceae Calystegia subacaulis	R		U				
Convolvulaceae Convolvulus arvensis						R	
Crassulaceae Dudleya cymosa		R	R	R			
Crassulaceae Dudleya sp.		R					
Crassulaceae Sedum spathulifolium		R					
Cuscutaceae Cuscuta californica		R					
Ericaceae Arctostaphylos pungens			C			U	
Ericaceae Arctostaphylos sp.			R		U		
Euphorbiaceae Euphorbia sp.			R				
Fabaceae Glycyrrhiza lepidota		R					
Fabaceae Lotus humistratus/wragelianus						R	
Fabaceae Lotus micranthus						R	
Fabaceae Lotus purshianus			U			R	
Fabaceae Lotus scoparius	U	C	C				
Fabaceae Lotus scoparius var.scoparius						C	U
Fabaceae Lotus sp.			U	U	R		
Fabaceae Lotus wrangelianus			U				
Fabaceae Lupinus albifrons	R		U	U		R	U
Fabaceae Lupinus albifrons var.albifrons						U	U
Fabaceae Lupinus bicolor			R				
Fabaceae Lupinus concinnus						R	
Fabaceae Lupinus microcarpus var.densiflorus						R	
Fabaceae Lupinus sp.	R		R	R			
Fabaceae Melilotus indicus			R			U	R
Fabaceae Trifolium albopurpureum						R	
Fabaceae Trifolium depauperatum						R	
Fabaceae Trifolium gracilentum var.gracilentum						R	
Fabaceae Trifolium microcephalum			R				R
Fabaceae Trifolium sp.	U		U	R			
Fabaceae Trifolium willdenovii			U			R	R
Fabaceae Vicia sp.			R				
Fabaceae Vicia villosa	U			R		U	U
Fagaceae Quercus agrifolia						R	R
Fagaceae Quercus agrifolia var.agrifolia						U	U
Fagaceae Quercus douglasii						R	R
Fagaceae Quercus lobata						R	

Species							
Fagaceae Quercus sp.	U	R	R			R	R
Fumariaceae Dicentra chrysantha		U	R	U			R
Fumariaceae Dicentra sp.			U				
Geraniaceae Erodium botrys						R	R
Geraniaceae Erodium brachycarpum						U	U
Geraniaceae Erodium cicutarium			R			U	U
Geraniaceae Erodium sp.		R					
Hippocastanaceae Aesculus californica		R	R			R	
Lamiaceae Lamium amplexicaule							R
Lamiaceae Lepechinia calycina		U	U	R		R	U
Lamiaceae Marrubium vulgare						U	U
Lamiaceae Mentha spicata		U					
Lamiaceae Mentha suaveolens		U					
Lamiaceae Monardella lanceolata		R					
Lamiaceae Monardella sp.			R				
Lamiaceae Monardella villosa		R				R	R
Lamiaceae Salvia columbariae						R	
Lamiaceae Salvia mellifera	R	U	U	U		U	R
Lamiaceae Stachys bullata		U	R	U		R	U
Lamiaceae Trichostema lanatum	R	U	U	R		R	R
Lamiaceae Trichostema lanceolatum	R	U					R
Liliaceae Bloomeria crocea		U	U			U	
Liliaceae Brodiaea sp.		R					
Liliaceae Brodiaea terrestris		R	U			R	R
Liliaceae Calochortus venustus		U	U	U		U	U
Liliaceae Dichelostemma capitatum		U				R	U
Liliaceae Triteleia hyacinthina		R					
Liliaceae Triteleia lugens		R	U				U
Liliaceae Zigadenus fremontii		R	R				
Liliaceae Zigadenus venenosus							R
Malvaceae Eremalche parryi				U			
Malvaceae Malacothamnus aboriginum		U	U			U	U
Oleaceae Fraxinus dipetala		R				R	
Onagraceae Camissonia sp.	R	U	R	R			R
Onagraceae Clarkia affinis		R				R	
Onagraceae Clarkia cylindrica		R	R				R
Onagraceae Clarkia modesta		R	R	R		R	
Onagraceae Clarkia purpurea	R	U	U	U		C	U
Onagraceae Clarkia similis		R					
Onagraceae Clarkia sp.	U	U	U	U			
Onagraceae Clarkia speciosa						R	R
Onagraceae Clarkia unguiculata		C	C	U		C	C
Onagraceae Epilobium canum			R				
Orobanchaceae Castilleja affinis			R			R	
Orobanchaceae Castilleja exserta		R	R			R	
Orobanchaceae Castilleja sp.		R	R				
Orobanchaceae Pedicularis densiflora						R	
Orobanchaceae Pedicularis sp.			R				
Orobanchaceae Triphysaria pusilla						R	
Papaveraceae Dendromecon rigida		U	R	R		R	
Papaveraceae Eschscholzia californica	U	C	U	C	R	C	C
Papaveraceae Eschscholzia sp.					R		
Papaveraceae Meconella linearis		R					
Papaveraceae Platystemon sp.			R				

	Col1	Col2	Col3	Col4		Col6	Col7
Phyrmaceae Mimulus aurantiacus	R	C	R	C		U	R
Phyrmaceae Mimulus guttatus				R		U	R
Phyrmaceae Mimulus pilosus		U					
Phyrmaceae Mimulus sp.		R	R				
Pinaceae Pinus sabiniana						R	R
Plantaginaceae Antirrhinum multiflorum		U	R				
Plantaginaceae Antirrhinum sp.			R	R			
Plantaginaceae Collinsia heterophylla	U	C	C	U		R	U
Plantaginaceae Collinsia parviflora		U					
Plantaginaceae Keckiella breviflora		U	U				
Plantaginaceae Penstemon centranthifolius		U	R			U	U
Plantaginaceae Penstemon heterophyllus		R	U			U	U
Plantaginaceae Plantago erecta						R	
Plantaginaceae Veronica anagallis-aquatica		U				R	
Polemoniaceae Gilia achilleifolia			U				
Polemoniaceae Gilia angelensis		U					
Polemoniaceae Gilia capitata	R	R				R	
Polemoniaceae Gilia sp.		R	U				
Polemoniaceae Linanthus parviflorus						R	
Polemoniaceae Linanthus sp.		R	R				R
Polemoniaceae Navarretia hamata							U
Polemoniaceae Navarretia sp.		R					
Polygonaceae Chorizanthe douglasii	R	U		C		U	U
Polygonaceae Eriogonum elongatum		U					
Polygonaceae Eriogonum fasciculatum	U	C	C	U		R	
Polygonaceae Eriogonum fasciculatum var.foliolosum						C	C
Polygonaceae Eriogonum gracile		R					
Polygonaceae Eriogonum nortonii		R				R	
Polygonaceae Eriogonum sp.			U				
Polygonaceae Eriogonum vimineum		U					
Polygonaceae Polygonum punctatum		R					
Polygonaceae Polygonum sp.				U			
Portulacaceae Claytonia perfoliata		R	U			U	R
Portulacaceae Montia fontana		R					
Primulaceae Anagallis arvensis			R				R
Primulaceae Dodecatheon clevelandii							R
Primulaceae Dodecatheon clevelandii ssp.patulum						R	
Primulaceae Dodecatheon sp.		R					
Ranunculaceae Clematis lasiantha		R				R	R
Ranunculaceae Clematis sp.		R					
Ranunculaceae Delphinium hesperium							R
Ranunculaceae Delphinium hesperium ssp.pallescens						R	
Ranunculaceae Delphinium parryi						R	R
Ranunculaceae Delphinium parryi/patens						R	
Ranunculaceae Delphinium sp.		U	R	R			
Ranunculaceae Ranunculus californicus		R	U			C	U
Rhamnaceae Ceanothus cuneatus		R	C	U		U	
Rhamnaceae Ceanothus cuneatus var.cuneatus						U	U
Rhamnaceae Ceanothus sp.			R		R		
Rhamnaceae Rhamnus ilicifolia		R	U	C		R	R
Rhamnaceae Rhamnus sp.						R	
Rosaceae Adenostoma fasciculatum	U	U	C	R		C	C
Rosaceae Cercocarpus betuloides		R				R	R
Rosaceae Drymocallis glandulosa							R

Rosaceae Heteromeles arbutifolia		U				R	
Rosaceae Prunus ilicifolia		U	R	C		R	
Rosaceae Rosa californica		R				U	R
Rosaceae Rubus parviflorus				R			
Rosaceae Rubus sp.		R					
Rosaceae Rubus ursinus		U					
Rubiaceae Galium sp.		R					
Salicaceae Salix exigua						U	U
Salicaceae Salix laevigata						U	R
Salicaceae Salix lasiolepis						U	U
Salicaceae Salix sp.		U	C				
Saxifragaceae Lithophragma affine			R				
Saxifragaceae Saxifraga californica			U				
Scrophulariaceae Scrophularia californica		R		R			
Solanaceae Solanaceae sp.					R		
Solanaceae Solanum umbelliferum		U	R			U	U
Valerianaceae Plectritis ciliosa							R
Valerianaceae Plectritis macrocera			U			U	
Valerianaceae Plectritis sp.		R	U				
Verbenaceae Verbena lasiostachys var.scabrida						R	R
Verbenaceae Verbena sp.							R
Violaceae Viola pedunculata		R	U			U	U
Count of unique floral taxa on which bees were collected in each year of Pinnacles study (sampling effort not equal):	**30**	**142**	**115**	**49**	**11**	**128**	**102**

758 **S3 Table**. **Bee biodiversity density results for all known native bee inventory projects with**

759 **at least 100 species in natural or semi-natural areas across the United States (N= 23).**

Study Location	Date range	Species	Approx. Total Area (km^2)	Approx. Species Density (spp./ km^2)	References
Grand Staircase Escalante National Monument, UT	2000-2003	656	7,610	0.086	[38]
Clark County, NV	1998, 2005, 2006	598	20,487	0.029	[35]
Yosemite National Park, CA	2006-2009	554	3028	0.18	*pers. comm. T. Griswold*
Pinnacles National Park, CA	1996-1999, 2002, 2011-2012	450	109	4.1	*present results* & [37]
San Bernardino, AZ*	2000-2007	383	1,088*	0.35	[23]
Carlsbad Caverns National Park, NM	2010-2011	364	189	1.9	*pers. comm. T. Griswold*
Curlew Valley, ID	1969-1974	340	4,999	0.068	[71] & *updated totals by pers. comm. T. Griswold*
San Rafael Desert, UT	1979-1992	333	5,180	0.064	[72]
Mojave National Preserve, CA	1975-1995	305	6,475	0.047	*pers. comm. T. Griswold*
Black Hills of SD and WY	2010-2011	290	12,950	0.022	[41]
Carlinville, IL*	1884-1916	288	256*	1.1	[23]
Plummers Island, MD%	1920s-2006	232	0.15	1,547	[73]
MPG Ranch, MT	2013-2015	229	39	5.9	[52]
Indiana Dunes, IN	2003, 2004, 2010	204	60	3.4	[74]
Albany County, WY*	1995-1996	200	11,160*	0.018	[75]
Palouse Prairie, ID	2012-2013	174	2,122	0.082	[76]
Dugway Proving Ground, UT*	2003, 2005	163	3,243*	0.050	[40]
Channel Islands, CA	Not specified	154	904	0.17	[77]
Black Rock Forest Preserve, NY	2003	144	15.5	9.3	[42]
Tonasket Ranger District, WA*	2004	140	1,678*	0.083	[78]
Black Belt Prarie, MS*	1991-2001	118	803*	0.15	[79]
Archibold Biol. Station, FL		113	21	5.4	[53]
Hattiesburg, MS*	1943-1944	104	140*	0.74	[80]

760 *Area size was not specified by publication or through author communications, so was estimated

761 by calculating known size of map area named.

762 %The Plummer's Island study was eliminated as an outlier in the species-area relationship shown

763 in Fig 5 because of its extremely small size compared to other studies.