Title

Drift and directional selection are the evolutionary forces driving gene expression

divergence in eye and brain tissue of *Heliconius* butterflies.

Ana Catalán[1,3], Adriana Briscoe[2], Sebastian Höhna[3,4].

[1] Department of Evolutionary Biology, Evolutionary Biology Centre (EBC), Uppsala

University, Norbyvägen 14-18 75236, Uppsala, Sweden.

[2] Department of Ecology and Evolutionary Biology, University of California, Irvine, CA

92697, USA.

[3] Division of Evolutionary Biology, Ludwig-Maximilians-Universität München,

Grosshaderner Straße 2, Planegg-Martinsried 82152, Germany.

[4] GeoBio-Center, Ludwig-Maximilians-Universität München, Richard-Wagner Str. 10,

80333 Munich, Germany

Corresponding author: ana.catalan@gmail.com

Abstract

Investigating gene expression evolution over micro- and macroevolutionary timescales will expand our understanding of the role of gene expression in adaptation and speciation. In this study, we characterized which evolutionary forces are acting on gene expression levels in eye and brain tissue of five *Heliconius* butterflies with divergence times of ~5-12 MYA. We developed and applied Brownian motion and Ornstein-Uhlenbeck models to identify genes whose expression levels are evolving through drift, stabilizing selection, or a lineage-specific shift. We find that 81% of the genes evolve under genetic drift. When testing for branch-specific shifts in gene expression, we detected 368 (16%) shift events. Genes showing a shift towards up-regulation have significantly lower gene expression variance than those genes showing a shift leading towards down-regulation. We hypothesize that directional selection is acting in shifts causing up-regulation, since transcription is costly. We further uncover through simulations that parameter estimation of Ornstein-Uhlenbeck models is biased when using small phylogenies and only becomes reliable with phylogenies having at least 50 taxa. Therefore, we developed a new statistical test based on Brownian motion to identify highly conserved genes (i.e., evolving under strong stabilizing selection), which comprised 3% of the orthoclusters. In conclusion, we found that drift is the dominant evolutionary force driving gene expression evolution in eye and brain tissue in *Heliconius*. Nevertheless, the higher proportion of genes evolving under directional than under stabilizing selection might reflect species-specific selective pressures on vision and brain necessary to fulfill species-specific requirements.

Introduction

Species and populations diverge through the accumulation of genetic changes that affect coding or non-coding genomic regions. Genetic variation affecting gene expression has the potential of changing gene expression patterns in a spatiotemporal manner, by changing gene expression profiles in specific organs and cell types at particular developmental stages (Carroll 2005; Signor and Nuzhdin 2018). This spatiotemporal attribute of gene expression might enable evolutionary change in a compartmentalized way, allowing for change where it is required but also allowing for the needed processes to remain conserved. Phenotypic diversity caused by changes in gene expression encompasses a great variety of traits, including changes affecting an organism's coloration (Nadeau 2016), size and shape (Ahi et al. 2017), as well as sensory perception and behavior, amongst other phenotypes (Lee et al. 2000; Wanner et al. 2007). Even though major advances have been made in linking gene expression variation to a phenotype (Catalán et al. 2016; Glaser-Schmitt and Parsch 2018), discerning the evolutionary forces shaping gene expression level variation among closely related species is an area that needs further research.

To understand the evolutionary forces acting on gene expression it is necessary to model within and between species gene expression variance. Neutral gene expression divergence between species leads to gene expression difference through divergence alone. Thus, neutral changes in gene expression modeled by random drift provides a null hypothesis to detect deviations from the expected neutral gene expression divergence. A linear relationship between divergence time and gene expression variance difference has been proposed for closely related species, assuming a clock-like (i.e., constant through

time) rate of gene expression divergence (Khaitovich et al. 2004; Khaitovich 2005).

Another approach to studying the evolutionary forces shaping gene expression evolution,

which is motivated by statistical phylogenetics, is fitting Brownian motion (BM) models.

BM-models are often used to describe changes in continuous trait through time using

random drift with rate σ^2 and taking into account the known phylogeny of the taxa of

interest (Felsenstein 1985). Thus, in a BM context, σ^2 can also be described as the

volatility parameter that determines the rate at which a trait's value diffuses away from its

current state (Bedford and Hartl 2009). Fitting BM models to investigate gene expression

evolution has shown that stabilizing selection and evolution through genetic drift can be

readily characterized (Kalinka et al. 2010; Wong et al. 2015).

Ornstein-Uhlenbeck (OU) models have also been used to study continuous trait

evolution in a phylogenetic context (Hansen 1997; Butler and King 2004). Ornstein-

Uhlenbeck models, an extension to BM-models, include two extra parameters, α and θ,

for modeling the strength of stabilizing (α) selection towards a phenotypic optimum (θ).

As in a BM context, σ^2 is the rate at which a trait changes through time, and α is the

force pulling back the diffused trait to an optimum state. This is analogous to stabilizing

selection pulling back a trait to its optimum value after having experienced a departure

from it. Theta (θ) is described as the trait's optimum state at a particular time point

toward which the pull of α is aimed (Hansen 1997; Butler and King 2004). OU-models

offer a useful framework to generate hypothesis about the evolutionary forces acting on

transcriptome levels, whether it is drift, stabilizing selection or directional selection

(Bedford and Hartl 2009; Rohlfs and Nielsen 2015; Wong et al. 2015; Chen et al. 2017;

Stern and Crandall 2018).

In this study, we used five closely-related species of *Heliconius* butterflies to

explore the evolutionary forces shaping gene expression variation in eye and brain tissue.

Heliconius charithonia, *H. sara*, *H. erato*, *H. melpomene* and *H. doris* (Figure 1) belong

to four of the seven distinct *Heliconius* phylogenetic clades with divergence times

ranging from 5.5 to 11.8 MYA. Beside showing a great diversity in wing color patterns

(Kronforst and Papa 2015), *Heliconius* butterflies also show a diversity of life history

traits (Salcedo 2010; Merrill et al. 2015), mating systems (Beltrán et al. 2007; Walters et

al. 2012) and behavior (Mendoza-Cuenca and Macías-Ordóñez 2005). Since *Heliconius*

butterflies are diurnal species, visual stimuli provide key sources of information about the

environment. For example, flowers and oviposition sites, potential mates or predators are

all targets of interest to butterflies in which the first line of perception is visual

(Finkbeiner et al. 2014; Finkbeiner et al. 2017). After visual cues are detected by the

visual system, the detected information travels to the brain, where it is processed and its

output can result in a specific behavior or physiological response. Thus, the brain's

processing and output together with the visual system have the potential of being finely

tuned according to a species' life history. In the case of *Heliconius* butterflies, a high

diversity of adult compound eye retinal mosaics (between sexes and species) has been

discovered (McCulloch et al. 2017), as well as species-specific differences in brain

morphology (Montgomery et al. 2016). Which evolutionary forces are shaping adult eye

and brain expression in *Heliconius* is one question we seek to investigate, and in that

way, gain an understanding into the potential role of inter-species gene expression

differences in speciation and adaptation.

Therefore, in this study we investigated which evolutionary forces are driving

gene expression variation in eye and brain tissue. More specifically, we aimed to identify

if expression variation in individual genes is evolving, for example, through drift,

stabilizing selection or directional selection. To this end, we generated a set of

orthoclusters shared among our five butterfly species together with expression data for

each gene in each orthocluster. We characterized the selective forces acting on gene

expression levels thereby revealing the fraction of the transcriptome evolving under drift,

directional selection, as well stabilizing selection.

Methods

Data set

The data set used in this study was published in (Catalán et al. 2018) and retrieved

from ArrayExpress: E-MTAB-6810 and Dryad data identifier: DOI: doi: 10.5061/

dryad.ds21fv5. In Catalán et al (2018), whole transcriptomes were generated from eye

and brain tissue combined for *Heliconius charitonia*, *H. sara*, *H. erato*, *H. doris* and *H.*

melpomene. *De novo* transcriptome assemblies were generated for each species and the

corresponding reads were mapped back to their matching transcriptome using Bowtie

(version 1.0.0). Raw read counts and FPKM values were calculated for each species and

used for downstream analysis. TransDecoder (version 5.0.2) was used to identify

candidate coding regions from each *de novo* Trinity transcriptome. The predicted coding

sequences were utilized to annotate each transcriptome by identifying orthologous hits in

UniProt, Flybase and Pfam databases using blastp (2.2.30) and keeping only hits with an

e-value $< 10^{-3}$ (Altschul et al. 1990; Chintapalli et al. 2007; Punta et al. 2012).

Orthology assessment

The set of orthoclusters used to assess gene expression evolution across

Heliconius was retrieved from supplementary table S15 published in Catalán *et. al.* 2018.

Briefly, the Unrooted Phylogenetic Orthology (UPhO) pipeline and model was used to

assess orthologous relationships between the five *Heliconius* species (Ballesteros and

Hormiga 2016). UPhO uses an all species pairwise blastp search and a Markov clustering

algorithm (MCL) (version 1.0.0) (Enright et al. 2002) to cluster sequences according to

sequence similarity. Clustered sequences were aligned with MAFFT (version 7.3.05)

(Katoh and Toh 2008) and curated after alignment with trimAl (version 1.3) (Capella-

Gutiérrez et al. 2009). Phylogenetic inference for each sequence cluster was done using

RAxML (version 8.2.10) (Stamatakis 2006) and orthology was assessed for each

generated tree using the UPhO algorithm. A matrix with $\log_2$ FPKM values was

generated for each orthoclusters which was used to analyze gene expression variance.

Modelling Gene Expression Evolution

Table 1. Summary of the models implemented in this work

Model	Description	Parameters
Equal species means	All species have the same mean gene expression level.	μ – global mean gene expression level
Unequal species means	All species have their own independent mean gene expression level.	μ_i – mean gene expression level per species
Brownian motion (BM)	Random drift of the species mean gene expression level along the phylogeny.	σ^2 - rate of drift
Brownian motion (BM) with shift	Random drift with one branch having a different rate (directional selection).	σ^2_B - rate of drift background branch

		σ^2_F - rate of drift foreground branch
Ornstein-Uhlenbeck (OU)	Stabilizing selection of the species mean gene expression level evolving along the phylogeny.	σ^2 - rate of drift α - strength of selection θ - optimal gene expression level
Ornstein-Uhlenbeck (OU) with shift	Directional selection due to a shift in optimal gene expression level.	σ^2_B - rate of drift background branch σ^2_F - rate of drift foreground branch. θ_B - optimal gene expression level background branch θ_F - optimal gene expression level foreground branch α - strength of selection

156   To study the forces driving gene expression evolution, we implemented a set of

157 six different statistical models (Table 1). Each model models the mean species gene

158 expression level (between-species variance) and the gene expression levels of individual

159 samples per species (within-species variance). How these mean species gene expression

160 levels evolve, or not, along the phylogeny and over time, is specific and central to each

161 model. We estimated the parameters of each model and performed Bayesian model

162 selection using Bayes factors to establish which model describes the observed data best

163 and thus which process is most likely to drive gene expression evolution in the five

164 *Heliconius* species of our study (see below).

165   The simplest model of gene expression assumes that all species have the exact

166 same mean gene expression level. In this case, we only have one parameter μ which

167 defines the mean gene expression level of all species. The expression level X_{ij} of

168 individual *i* from species *j* is modeled using a normal distribution with $X_{ij} \sim \text{Norm}(\mu, \delta^2_i)$.

We chose a uniform prior distribution between -20 and 20 for the mean gene expression

parameter μ. Note that we assume that every species has its own gene expression variance

parameter δ^2_i (see below). This model assumes there is no evolution of gene expression

levels, i.e., gene expression levels are completely conserved among species.

The second model that we implemented was a model where each species has its

own gene expression mean μ_i. Thus, we model the gene expression level X_{ij} of gene *i* from

species *j* using a normal distribution with $X_{ij} \sim Norm(\mu_i, \delta^2_i)$. In this model, each species

has a different mean gene expression level, but these gene expression levels do not

evolve under an evolutionary model; they are intrinsically different without any

mechanistic reason (no phylogenetic signal). As with the first model, we assumed a

uniform prior distribution between -20 and 20 for each mean gene expression level μ_i.

The third model that we implemented was a phylogenetic Brownian motion

model (Felsenstein 1985). We assume that any gene expression value at the root of the

phylogeny is equally probable. Then, the mean gene expression levels μ evolve along the

lineages of the phylogeny. The Brownian motion model specifies that the focal variable,

μ in our case, is drawn from a normal distribution centered around the value of the

ancestor, μ_A. The amount of change, i.e., rate of random drift, is defined by the parameter

σ^2. We assumed a log-uniform prior distribution between 10E-5 and 10E5 for the drift

parameter σ^2. Thus, the mean gene expression levels μ_i for the species of the phylogeny

are distributed according to a multivariate normal distribution where the covariance

structure is defined by the phylogeny (Felsenstein 1985). This means that more closely-

related species are expected to have a more similar mean gene expression level because

they share more evolutionary history (i.e., they are more recently diverged), which is

modeled by the covariance structure. Such a phylogenetic model of gene expression

evolution has been applied previously by (Bedford and Hartl 2009). Importantly, the BM

model only defines how the mean gene expression levels evolve but does not allow for

any sample variance of the individuals of a species. Therefore, we extended the standard

phylogenetic BM model to allow for within-species sample variance where again the

expression level X_{ij} of individual i from species j is normally distributed with $X_{ij} \sim$

Norm(μ_i, δ^2_i) where δ^2_i is the within-species variance parameter (Ives et al. 2007; Rohlfs

and Nielsen 2015). This extension to allow for within-species variance was developed for

all phylogenetic models (BM, BM with shift, OU and OU with shift).

The fourth model that we implemented was a phylogenetic BM model with

branch-specific rates of evolution, thus detecting directional selection. The mean gene

expression level evolves under a BM model (i.e., random drift) where the rate of

evolution for branch k is given by σ^2_k (O'Meara et al. 2006; Eastman et al. 2011). Thus,

a branch with a higher rate of evolution σ^2_k signifies more change in gene expression

levels than under a constant rate random drift model (the BM model). Directional

selection can therefore be detected by inferring an elevated estimate of σ^2_k compared

with the background rate of drift σ^2. Specifically, we applied a background rate of drift

σ^2_B to all branches except the chosen foreground branch which received its own rate of

drift σ^2_F.

The fifth model we implemented was a phylogenetic Ornstein-Uhlenbeck process

(Hansen 1997). The Ornstein-Uhlenbeck (OU) process models, similar to BM, the

evolution of the mean gene expression level per species along a phylogenetic tree.

However, unlike BM, the mean expression level diffuses with rate σ^2 and is attracted

with strength α to an optimum level θ. Thus, the OU process has an expected variance of

$\sigma^2/2\alpha$ which is independent of time, i.e., does not increase with increasing time but

instead stabilizes (c.f. Figure 10). The variance becomes small if either the strength of

selection is large or the rate of drift is small. This is, in fact, a major problem for OU

models which cannot distinguish if attraction (or selection) is strong or diffusion is weak

(Ho and Ané 2014; Cooper et al. 2016).

The sixth model we implemented was an Ornstein-Uhlenbeck process with a

branch-specific shift in both the rate of drift σ^2 and the optimum gene expression level θ

(Rohlfs et al. 2014; Uyeda and Harmon 2014). Thus, this branch-specific OU model is

analogous to the branch-specific BM model, allowing for directional selection in an OU

framework. Specifically, we tested if there was a significant support for the chosen

foreground branch which received its own rate of drift σ^2_F and optimum θ_B to be different

from the background rate of drift σ^2_B and optimum θ_B. We used the same prior

distributions as before and assumed that both parameters for the background and

foreground branches are drawn from the identical prior distribution. This model has in

total five free additional parameters along with the five nuisance parameters (the within-

species variances). Thus, we expect that this model is more prone to be over-

parameterized for our dataset with five species. Nevertheless, our Bayesian approach for

parameter estimation and model selection integrates over parameter uncertainty and

penalizes extra parameters by integrating over the prior distribution.

Parameter Estimation and Model Selection

We estimated parameters for our different models in a Bayesian statistical

framework. Thus, we approximated the posterior distribution of the model parameters

using Markov chain Monte Carlo sampling (Metropolis et al. 1953; Hastings 1970). We

ran a separate MCMC analysis for each model and gene, 2393 analysis per model. Every

model parameter was updated twice per MCMC iteration where the order of parameter

updates was chosen randomly. We applied the same settings of the MCMC algorithm for

each model. First, a burn-in phase of the MCMC algorithm was run for 2,000 iterations

with auto-tuning every 100 iterations. Then, the actual MCMC simulation was run for

50,000 iterations with sampling 10 iterations, yielding 5,000 samples from the posterior

distribution (Höhna et al. 2017).

Model selection was performed using marginal likelihoods. Marginal likelihoods

are the probability of the data for a specific model integrated over all possible parameter

values. From the marginal likelihood we can then compute Bayes factors and model

probabilities (i.e., weights of a model being the true model generating the data given a set

of candidate models). We approximated the marginal likelihoods using stepping stone

sampling (Fan et al. 2011). The stepping stone algorithm implemented in RevBayes

consisted of 128 MCMC runs where each MCMC ran had the likelihood function raised

to the power of β computed by the quantiles of a beta probability distribution (Höhna et

al. 2017).

Data availability

The five different models that we used in our study are implemented in Bayesian

phylogenetic inference software RevBayes v1.0.8 (Höhna et al. 2016). For efficient

computations, we implemented the restricted maximum likelihood (REML) algorithm for

BM models (Felsenstein 1985) and OU models (Fitzjohn 2012; Freckleton 2012). The

source code and compiled executables of RevBayes are available from

https://github.com/revbayes/revbayes and tutorials about the analyses are available from

https://revbayes.github.io/tutorials/.

Results

To assess the evolutionary forces acting on gene expression levels in *H.*

charithonia, *H. sara*, *H. erato*, *H. doris* and *H. melpomene* (Figure 1), we used the gene

orthology dataset, composed of 2373 orthologous genes, published before (Catalán et al.

2018). From this previous work, we also obtained FPKM (Fragments per Kilobase per

Million mapped read) values for each gene and sample. The $\log_2$ transformed FPKM

values were used to build an expression matrix and to model gene expression evolution.

Figure 1 Phylogenetic relationship of the *Heliconius* species used in this study showing

divergence times at each node (Kozak et al. 2015).

Testing for equality of within-species variance in gene expression levels

Equality of variances among populations or samples drawn from a normal

distribution is usually assumed when testing for differences in mean values obtain from

continuous data or gene expression data as in our case (Warnefors and Eyre-Walker

2012; Rohlfs et al. 2014). Assuming equality of variances when it is not the case can lead

to high Type I error rates (Gastwirth et al. 2009). To avoid assuming equality of

variances, we computed the within-species variances for each orthocluster and checked

for the presence of a correlation across species. From a pairwise assessment of within-

species variance we found no significant correlation among all possible pairs, with

Pearsons's rho values ranging from 0.07 to 0.2 (Figure 2). Since gene expression variance

across species is heterogeneous, hence not correlated among species, we treated within-

species variance as a random variable when fitting BM and OU models.

Figure 2 Pairwise correlation between five *Heliconius* species and their gene expression

within-species variance. The correlation strength between within-species variances was

estimated by calculating Pearson's correlation coefficient, rho ranging from 0.07 to 0.2.

Testing for gene expression evolution through drift using Brownian motion models

We applied BM models to describe changes in gene expression levels through

random drift. As the alternative hypotheses, we used two non-phylogenetic models where

either all species had identical mean gene expression levels (Model 1) or all species had

their own independent mean gene expression levels (Model 2). For each gene we

computed the probability of the BM model having produced the observed data, i.e., a

high probability means that it is more probable that the gene expression levels evolved

under a BM model whereas a low probability means that it is more probable that the gene

expression levels evolved under a non-phylogenetic model (Model 1 and Model 2). A

model probability of >0.75 corresponds to a Bayes factor of >3 (positive support) and a

model probability of >0.95 corresponds to a Bayes factor of >20 (strong support).

Figure 3. Testing for random drift in gene expression levels of *Heliconius* using

Brownian motion. Significance is shown at model probability > 0.75 (solid red, Bayes

factor > 3, positive support) and model probability > 0.95 (solid red, Bayes factor > 20,

strong support). (A) Shows the comparison between the two non-phylogenetic models

(identical vs independent species mean). (B) Shows the model probability of the BM

model compared with the independent species mean model. (C) Shows the model

probability of the BM model compared with the identical species mean model.

Our results show that the majority of gene expression levels (2369 out of 2393)

are evolving as expected given the known *Heliconius* species phylogeny, i.e., that there is

a strong phylogenetic signal (Figure 3). These results indicate that drift is the dominating

evolutionary force driving transcriptome change. Only a small fraction of the

orthoclusters did not show a phylogenetic signal, opening the question of the putative

genetic forces shaping gene expression levels of these genes.

Testing for conserved gene expression level

The next question we explored was how prevalent conserved gene expression

levels are in eye and brain tissue of *Heliconius*. This question could be answered with our

previous results by computing how often Model 1, with identical species means, was

recovered (Figure 2C). However, our model selection procedure relied on computing

marginal likelihoods which are intrinsically sensitive to the choice of prior distribution

(Berger 1990; Kass and Raftery 1995; Sinharay and Stern 2002). Therefore, we

additionally performed a sensitivity analysis of $\sigma^2 = 0$ using Monte Carlo simulation as

follows (Goldman 1993). We estimated the posterior distribution of all parameters under

the identical species mean model (the only parameters were the within-species variances),

then we used 1,000 parameter samples from the posterior distribution to simulate gene

expression datasets (e.g., a dataset consisting of a single gene with five species and 6-12

individuals per species) under the identical species mean model, yielding 1,000 simulated

datasets per gene in total. Then, for every gene of the 2393 genes, we estimated σ^2 for

each simulated dataset as well as the original dataset, which amounted to a total of

2,395,393 MCMC analyses. Finally, we calculated if the mean posterior estimate of the

empirical dataset was larger than 95% of the mean posterior estimates of the simulated

datasets. In the cases when the mean posterior estimate of σ^2 was not larger than the

mean estimate of 95% of the simulated datasets we concluded that these genes are highly

conserved (Figure 4).

Figure 4. Posterior mean estimates of the drift rate σ^2 (blue) and the 95% threshold

computed (red) using Monte Carlo simulations. The genes were sorted by an ascending

estimate of σ^2. Inset: close-up of genes whose σ^2 is not significantly bigger than zero.

357   By using the described approach, we uncover a set of 83 orthoclusters whose gene

expression variance across species is highly conserved (Figure 4 and Figure S1). A sigma

squared not significantly different from zero, can be caused by stabilizing selection

hindering gene expression divergence, resulting in more similar gene expression patterns

across different *Heliconius* species.

Testing for stabilizing selection acting on gene expression levels

Subsequently, we moved forward into implementing an Ornstein-Uhlenbeck

model (OU) to investigate the strength of stabilizing selection (Bedford and Hartl 2009;

Rohlfs et al. 2014). OU-models are an extension of BM-models, in which they include

two extra parameters, α and θ. In a BM context, if σ^2 is the rate at which a trait changes

through time, α is then described as a force pulling back the diffused trait to an optimum

state (θ).

We estimated the marginal likelihood for each gene under a BM model and an

OU model. Then, we computed the probability (i.e., support) of an OU model over a BM

model using the marginal likelihoods. Our results show a very low support for stabilizing

selection (Figure 5). When the marginal likelihoods were examined, in 99.7% of the

cases a BM model explained our expression data better than an OU-model.

Figure 5. Model probability when testing the strength of alpha when fitting an OU model

for the assessment of stabilizing selection. Significance is shown at model probability >

0.75 (solid red, Bayes factor > 3, positive support) and model probability > 0.95 (solid red,

Bayes factor > 20, strong support). There are only 7 genes with a significant support for

stabilizing selection.

Testing the power to estimate stabilizing selection

Our results indicating that very few genes evolved under stabilizing selection

conflict with previous findings (Bedford and Hartl 2009). However, it has previously

been discussed that when working with small phylogenies (less than ten species) there is

a lack of power for parameter estimation when using an OU-model (Rohlfs et al. 2014).

By simulating data under an OU model using phylogenies with varying numbers of taxa,

we were able to show how parameter estimation is biased. The attraction parameter α

could only be estimated closely to the true values used for the simulations when the

phylogenies contained 50 or more taxa. Thus, we can observe that the bias observed for

parameter estimation drops considerably when the number of taxa composing the

phylogeny reaches 50 (Figure 6). This observation holds as well for the estimation of σ_2

under a range of sigma values (Figure S2). Our simulation study shows that attention

needs be paid when applying OU models to assess gene expression evolution for

phylogenies containing less than 50 taxa.

397

Figure 6. Simulation study for the assessment of parameter estimation bias under an OU-

model. The relative bias in estimates of the attraction/selection parameter (α) through

1000 simulations under sigma values ranging from 0.1 – 10 and alpha values ranging

from 0.01 – 10. Simulations were performed for phylogenies with sizes ranging from 5 to

1000 taxa.

Detection of branch specific shifts in gene expression

To reveal genes whose gene expression patterns have putatively been shaped by

directional selection, we tested for branch-specific shifts in evolutionary rates along the

Heliconius tree. To explore branch-specific shifts in gene expression, firstly we used a

BM model to test for the evolutionary rate (σ^2) of a focal branch being different from the

background rate (i.e., the rest of the branches in the phylogenetic tree) and assessed

significance by applying Bayes factors (Figure S3 and Figure 7). Secondly, we also tested

branch-specific shifts through an OU model and tested for a branch-specific shift in gene

expression level optimum (θ_F) vs the rest of the tree's θ_B (Figure S4).

With a BM approach, we were able to detect a total of 322 branch specific shifts,

when considering only tip branches (Figure 7). We found 112 branch specific shifts in the

HER linage, 70 in HAS, 67 in HCH, 44 in HDO and 29 in HME (Figure 7 and Figure

S4). HCH, HAS and HDO had more shifts towards a down-regulation, although only in

HCH and HAS was this difference significant (sign test, HCH: *P*-value 6.738e-05 and

HAS: *P*-value 1.653e-06). In HER and HME more up-regulated genes were causing a

branch specific shift, although no significant difference was found.

When implementing an OU model we recover a total of 75 genes showing a

branch specific shift in gene expression optimum (Figure 7 and Figure S4). From these

genes, 55 also show a branch specific shift when implementing a BM model and 20 genes

show uniquely a gene expression level shift in optimum when using an OU model (Figure

S4).

Figure 7. Barplot showing branch-specific shifts on gene expression levels in *Heliconius*.

Bars in light blue show branch shifts identified by BM and dark blue bars show branch-

shifts identified by OU models.

Next, we assessed within-species gene expression variance of all the genes

identified as having a branch-specific shift in gene expression through BM and OU

models. When we plotted the distribution of the within-species variance we found that

up-regulated genes have a significantly lower variance when compared to genes with a

gene expression shift towards a down-regulation (Figure 8). Different evolutionary forces

acting on shifts causing up- or down-regulation have the potential of maintaining

different levels of gene expression variation within species.

Figure 8. Boxplot showing the distribution of the within-species variance of gene

expression levels identified as having a shift towards up-regulation and shifts towards

down-regulation. Numbers above the boxplots show the total number of genes identified

with a BM and an OU model. Wilcoxon-test: * $P<0.05$, ** $P<0.01$, *** $P<0.001$.

Discussion

Gene expression evolution through genetic drift

Our study of the evolutionary forces acting on gene expression in eye and brain

tissue of *Heliconius* butterflies revealed that most transcriptome levels (81%) are

evolving under drift. According to neutral expectations, phenotypic changes are expected

to accumulate as a function of time, by drift and mutation alone (Lande 1976). As a

consequence, the change of transcriptomic levels through drift should reflect the

divergence history of the taxa of interest. From our BM analysis, we conclude that in

most of the gene expression levels on eye and brain a phylogenetic signal can be

recovered (Figure 3).

Consequently, we hypothesize that gene expression variation influencing

phenotypic variation across species mainly arises through random drift. Evolutionary

rates of gene expression evolution have been investigated at the population and at the

species level and it has been found that the proportion of the type of evolutionary force

acting on transcriptomic levels is not constant across taxa (Whitehead and Crawford

2006a; Nourmohammad et al. 2017; Stern and Crandall 2018). For example, when

examining the evolutionary forces acting on gene expression levels in several fish

populations, the authors reported that the dominant force driving expression changes was

genetic drift (Whitehead and Crawford 2006b). Comparably, in a study concerning

primates, genetic drift was the main force driving gene expression evolution (Khaitovich

et al. 2005). The proportion of gene expression levels evolving by drift depends on the

strength of natural selection acting on the interrogated transcriptome. For example, in a

comparison between different organ types in mammals, gonad gene expression showed

the lowest phylogenetic signal when compared to other organs like cerebellum or heart

(Brawand et al. 2011). In *Heliconius* butterflies, other organs would need to be tested in

order to get a more global understanding on how gene expression is evolving in the

whole organism.

We explored the gene expression data further by comparing the expected gene

expression divergence under a BM model to the observed data. Consequently, we

simulated expression levels for 10,000 genes along the known *Heliconius* phylogeny and

computed the mean of the pairwise species difference. Similarly, we computed the mean

pairwise difference of the observed gene expression data. Alternatively, we can also

derive the expected divergence in gene expression levels between two species over time

under Brownian motion. Both species evolve under random drift and, thus, their gene

expression values are normally distributed with variance σ^2 x T where T is the time since

the most recent common ancestor of the two species. Therefore, the difference in gene

expression levels between the two species is normally distributed with variance 2 x σ^2 x

T. Since we are only interested in the absolute value of the gene expression difference,

we use a truncated normal distribution instead. From this truncated normal distribution

with mean zero and variance 2 x σ^2 x T we compute the expected gene expression

difference through time (Figure 9). For the empirical data, we estimate σ^2 using a sum of

squares approach. We find that our observed gene expression data has a close fit to the

simulated data (Figure 9).

Figure 9. Between species gene expression variance plotted as a function of divergence

time according to the *Heliconius* phylogeny. Red: σ^2 from gene expression levels

observed in *Heliconius*. Blue: simulated gene expression difference under random drift

with different values of sigma.

Gene expression evolution through genetic stabilizing selection

Studies done in *Drosophila* and mammals have shown that stabilizing selection is

the evolutionary force dominating gene expression evolution (Rifkin et al. 2003; Rohlfs

and Nielsen 2015). In contrast to these studies, in *Heliconius* we uncovered that only 3%

of gene expression are either highly conserved (Figure 4) or evolving through stabilizing

selection (Figure 5). Factors like tissue type, gene functionality turnover or epistatic

levels, have the potential to influence the degree of stabilizing selection acting on the

transcriptome (Larracuente et al. 2008; Kalinka et al. 2010; Romero et al. 2012).

Additionally, in groups that have experienced an adaptive radiation, like in *Heliconius*

(Kozak et al. 2015), and thus have recently experienced an elevated rate of trait

evolution, directional selection might be more recurrent than stabilizing selection.

508   OU models are suitable models to study the force of stabilizing selection acting

on a phenotype since the α parameter simulates the strength of selection keeping a trait

close to an optimum (Beaulieu et al. 2012), as several studies exemplify (Kalinka et al.

2010; Brawand et al. 2011; Stern and Crandall 2018). When we applied an OU model to

identify stabilizing selection on gene expression, we detected parameter estimation biases

as shown by our simulation study (Figure 6). For small phylogenies, accurate parameter

estimation is challenging since statistical power is weak with small sample sizes (Rohlfs

et al. 2014). Specifically, it is very challenging with small phylogenies to distinguish

between conserved gene expression levels due to low values of drift (i.e., no change) and

high values of selection (i.e., drift is removed due to selection). Not only the number of

taxa, but also the depth of the phylogeny can influence the suitability of OU-models to

infer stabilizing selection (Fay and Wittkopp 2008; Bedford and Hartl 2009). Therefore,

we propose that for small phylogenies, testing for $\sigma^2 = 0$ under a BM framework and

assessing for significance by applying Monte Carlo simulations is a better model choice

to uncover stabilizing selection. When using this approach, we identified 83 genes with

conserved gene expression levels across species. These genes might be involved in

maintaining conserved processes that are essential for the function of eye and brain tissue

in *Heliconius*. For example, from the top ten genes with the most conserved gene

expression levels, we found the transcription factor *bobby sox* (*bbx*) (Group_674,

Appendix 1). BBX, belongs to the high mobility box domain superfamily, which are

involved in transcription, replication and chromatin remodeling (Chintapalli et al. 2007).

BBX has also been found to have orthologues in flies, human and mice (Nitta et al. 2015)

suggesting a high essentiality of *bbx* expression. Another highly conserved orthocluster

(Group 977, Appendix 1) was annotated as *glaikit* (Chintapalli et al. 2007), which is

known to be essential for the formation of epithelial polarity and nervous system

development (Dunlop et al. 2004).

We repeated our previous data exploration by simulating 10,000 genes under an

OU-model under a range of σ^2 and α values and computed the mean differences between

pairs of species. We observed a reasonably good fit to our data (Figure 10). Most

importantly, in Figure 10 we observed a steeper change in gene expression difference

between closely related species (low evolutionary distance). Consequently, adding more

species, including closely related species, to our model could not only improve OU

parameter estimation but could also help in disentangling the evolutionary forces acting

on gene expression divergence, specially between closely related species. Interestingly,

the observed differences (red dots) in gene expression levels (which are averaged over all

2393 genes) show a clear departure from the expected difference predicted by both a BM

and OU model (red lines in Figure 9 and Figure 10). This could be a strong indication of

gene expression divergence under different evolutionary forces (drift, stabilizing

selection or directional selection).

Figure 10. Between species gene expression variance plotted as a function of divergence

time according to the *Heliconius* phylogeny. Red: σ^2 from gene expression levels

observed in *Heliconius*. Blue: simulated gene expression variance under different values

of sigma. Each panel shows estimates for a different value of alpha (α).

Gene expression evolution through genetic directional selection

To reveal branch-specific shifts in gene expression levels we applied BM and OU

models, allowing for branch-specific shifts in the rest of the phylogeny. Using this

approach, we found that 16% of the genes show a branch-specific shift, towards either

up- or down-regulation, with increased expression levels showing lower variance than

expected (Figure 8). The direction of gene expression shifts might be influenced by its

mode of regulation. For example, in yeast it was found that regulatory mutations

affecting *trans*-regulatory factors were more likely to cause an increase in gene

expression. Conversely, mutations on *cis*-regulatory elements were found to be skewed

towards a decrease in expression (Metzger et al. 2016). On the other hand, in primates, a

higher proportion of species-specific gene expression shifts were found to be towards

down-regulation (Khaitovich et al. 2005). If directional selection is causing a branch-

specific shift in gene expression one would expect to see a low within-species variance,

whereas if the shift is caused by a relaxation of purifying selection or perhaps balancing

selection, a higher within-species variance would be expected.

When we looked at the degree of variability between genes showing a shift

towards a higher or a lower expression level, we observed that down-regulated genes

have a significantly higher variance than genes showing up-regulation (Figure 8). From

this observation, we hypothesize that relaxation of purifying selection might be driving

the shifts causing down-regulation on gene expression, a pattern which could eventually

lead to a loss of expression. However, balancing selection or experimental noise could

also lead to an elevated within-species variance. Because of the cost of gene expression,

it is expected that only those genes that are essential and have fitness effects will continue

to be expressed, whereas genes that are not will eventually stop being transcribed (Stern

and Crandall 2018). However, a shift towards down-regulation does not always have to

be a consequence of relaxed purifying selection. For example, in the orthocluster with id

Group_449_clean_0, a 7-fold lower expression shift was detected in the branch leading to

H. doris (Figure S5), and a significantly smaller variance than expected transcriptome-

wide (Fisher's exact test, P-value < 0.001). Directional selection favoring down-

regulation of gene expression can occur in a scenario where fine tuning of expression

levels are necessary for optimal cell or tissue function (Cayirlioglu et al. 2008; Catalán et

al. 2016).

On the other hand, genes showing a branch-specific shift towards up-regulation

have significantly lower variance when compared to expression level shifts towards

down-regulation (Figure 8). This observed pattern could be a result of directional

selection acting on gene expression levels leading to a reduction of the variation observed

in gene expression. It is possible that in order to achieve an increase in gene expression

levels, the selective forces leading to up-regulation would have to be sufficiently strong

to result in a greater investment in energy allocated to transcription costs (Wagner 2005;

Lang et al. 2009). Some of the genes having the most extreme branch shifts in expression,

either toward a higher or a lower expression, are involved in enzymatic activity

(Appendix I). Enzymes support biochemical and physiological processes helping in the

optimization of tissue function (Wagner and Altenberg 1996; Feller and Gerday 1997).

Thus, optimal enzymatic activity might be a key factor for species-specific brain and eye

function, which in turn might be optimized for the species-specific life history and

ecological environment.

A factor possibly influencing the proportion of transcriptome levels found to be

evolving through drift, stabilizing or directional selection is the methodology used for

orthology assessment. In our analysis of gene expression variation, we assessed variation

in orthoclusters where an orthologous hit was found for each of our five *Heliconius*

species. Genes with fast-evolving protein rates—to the point that orthology assessment

becomes challenging—might also show gene expression shifts, which would not be

detectable in our experimental design. For example, orthology assessment for genes

showing sex-biased gene expression might require an alternative method. In fact, from

the ortholclusters that we identified in this study, only two included genes with sex-

biased expression (Catalán et al. 2018). Additionally, gene expression shifts due to

duplication events need to be explored by applying an appropriate statistical approach, as

expression of genes where a duplication event has happened could contribute to the

fraction of the transcriptome evolving by directional selection.

With this work, we have generated a set of candidate genes that are putatively

evolving through directional selection and that have the potential of being involved in the

processes of adaptation and speciation. To test the role of these genes in such processes,

functional validation will be necessary to gain a deeper insight in the evolutionary

consequences of gene expression shifts. Techniques like *in situ* hybridization, RNAi and

CRISPR/Cas9 are adequate tools that can be used in shedding light into the functionality

of these genes. Particularly interesting could be those genes whose gene expression levels

have shifted to the degree of showing absence of expression (Figure S6). The evolution

of gain and loss of gene expression across a phylogeny requires a suitable theoretical

framework that should be explored particularly since such events have the potential to

accelerate evolution.

Acknowledgements

Thanks to Aline Rangel Olguin for technical assistance with the orthocluster analysis.

This work was partially supported by National Science Foundation grant IOS-1656260 to

A.D.B and by the Knut & Alice Wallenberg Foundation to Jochen Wolf.

References

Ahi EP, Richter F, Sefc KM. 2017. A gene expression study of ornamental fin shape in

Neolamprologus brichardi, an African cichlid species. Sci Rep 7:1–14.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment

search tool. J Mol Biol 215:403–410.

Ballesteros JA, Hormiga G. 2016. A new orthology assessment method for phylogenomic

data: Unrooted Phylogenetic Orthology. Mol Biol Evol 33:1–18.

Beaulieu JM, Jhwueng DC, Boettiger C, O'Meara BC. 2012. Modeling stabilizing

selection: Expanding the Ornstein-Uhlenbeck model of adaptive evolution.

Evolution (N Y) 66:2369–2383.

Bedford T, Hartl DL. 2009. Optimization of gene expression by natural selection. Proc

Natl Acad Sci U S A 106:1133–1138.

Beltrán M, Jiggins C, Brower A, Bermingham E, Mallet J. 2007. Do pollen feeding and

pupal-mating have a single origin in *Heliconius*? Inferences from multilocus

sequence data. Biol J Linn Soc 92:221–239.

Berger JO. 1990. Robust bayesian. J Stat Plan Inference 25:303–328.

Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M, Liechti

A, Aximu-Petri A, Kircher M, et al. 2011. The evolution of gene expression levels

in mammalian organs. Nature 478:343–348.

Butler MA, King AA. 2004. Phylogenetic Comparative Analysis: A Modeling Approach

for Adaptive Evolution. Am Nat 164:683–695.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: A tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–

1973.

Carroll SB. 2005. Evolution at two levels: on genes and form. PLoS Biol 3:e245.

Catalán A, Glaser-Schmitt A, Argyridou E, Duchen P, Parsch J. 2016. An Indel

Polymorphism in the *MtnA* 3' Untranslated Region Is Associated with Gene

Expression Variation and Local Adaptation in *Drosophila melanogaster*. PLOS

Genet 12:1–24.

Catalán A, Macias-Muñoz A, Briscoe AD. 2018. Evolution of Sex-Biased Gene

Expression and Dosage Compensation in the Eye and Brain of *Heliconius*

Butterflies. Mol Biol Evol 35:2120–2134.

Cayirlioglu P, Kadow IG, Zhan X, Okamura K, Suh GSB, Gunning D, Lai EC, Zipursky

SL. 2008. Hybrid neurons in a microRNA mutant are putative evolutionary

intermediates in insect CO_2 sensory systems. Science (80-) 510:3–8.

Chen J, Swofford R, Johnson J, Cummings BB, Rogel N, Lindblad-Toh K, Haerty W,

Palma F di, Regev A. 2017. A quantitative model for characterizing the evolutionary

history of mammalian gene expression. bioRxiv:229096.

Chintapalli VR, Wang J, Dow JAT. 2007. Using FlyAtlas to identify better *Drosophila*

melanogaster models of human disease. Nat Genet 39:715–720.

Cooper N, Thomas GH, Venditti C, Meade A, Freckleton RP. 2016. A cautionary note on

the use of Ornstein Uhlenbeck models in macroevolutionary studies. Biol J Linn Soc

118:64–77.

Dunlop J, Morin X, Corominas M, Serras F, Tear G. 2004. *glaikit* is essential for the

formation of epithelial polarity and neuronal development. Curr Biol 14:2039–2045.

Eastman JM, Alfaro ME, Joyce P, Hipp AL, Harmon LJ. 2011. A Novel comparative

method for identifying shifts in the rate of character evolution on trees. Evolution (N

Y) 65:3578–3589.

Enright AJ, Van Dongen S, Ouzounis CA. 2002. An efficient algorithm for large-scale

detection of protein families. Nucleic Acids Res 30:1575–1584.

Fan Y, Wu R, Chen MH, Kuo L, Lewis PO. 2011. Choosing among partition models in

Bayesian phylogenetics. Mol Biol Evol 28:523–532.

Fay JC, Wittkopp PJ. 2008. Evaluating the role of natural selection in the evolution of

gene regulation. Heredity (Edinb) 100:191–199.

Feller G, Gerday C. 1997. Psychrophilic enzymes: Molecular basis of cold adaptation.

Cell Mol Life Sci 53:830–841.

Felsenstein J. 1985. Phylogenies and the comparative method. Am Nat 125:1–15.

Finkbeiner SD, Briscoe AD, Reed RD. 2014. Warning signals are seductive: relative

contributions of color and pattern to predator avoidance and mate attraction in

Heliconius butterflies. Evolution 68:3410–3420.

Finkbeiner SD, Fishman DA, Osorio D, Briscoe AD. 2017. Ultraviolet and yellow

reflectance but not fluorescence is important for visual discrimination of

conspecifics by *Heliconius erato*. J Exp Biol:jeb.153593.

Fitzjohn RG. 2012. Diversitree: Comparative phylogenetic analyses of diversification in

R. Methods Ecol Evol 3:1084–1092.

Freckleton RP. 2012. Fast likelihood calculations for comparative analyses. Methods

Ecol Evol 3:940–947.

Gastwirth JL, Gel YR, Miao W. 2009. The Impact of Levene's Test of Equality of

Variances on Statistical Theory and Practice. Stat Sci 24:343–360.

Glaser-Schmitt A, Parsch J. 2018. Functional characterization of adaptive variation

within a *cis*-regulatory element influencing *Drosophila melanogaster* growth. PLoS

Biol 16:1–28.

Goldman N. 1993. Simple diagnostic statistcal tests of models for DNA substitution. J

Mol Evol 36:182–198.

Hansen TF. 1997. Stabilizing Selection and the Comparative Analysis of Adaptation.

Evolution (N Y) 51:1341.

Hastings WK. 1970. Monte Carlo simulation methods using Markov Chains and their

applications. Biometrika 57:97–109.

Ho LST, Ané C. 2014. Intrinsic inference difficulties for trait evolution with Ornstein-

Uhlenbeck models. Methods Ecol Evol 5:1133–1146.

Höhna S, Landis MJ, Heath TA. 2017. Phylogenetic inference using RevBayes. Curr

Protoc Bioinforma 2017:6.16.1-6.16.34.

Höhna S, Landis MJ, Heath TA, Boussau B, Lartillot N, Moore BR, Huelsenbeck JP,

Ronquist F. 2016. RevBayes: Bayesian phylogenetic inference using graphical

models and an interactive model-specification language. Syst Biol 65:726–736.

Ives AR, Midford PE, Garland T. 2007. Within-species variation and measurement error

in phylogenetic comparative methods. Syst Biol 56:252–270.

Kalinka AT, Varga KM, Gerrard DT, Preibisch S, Corcoran DL, Jarrells J, Ohler U,

Bergman CM, Tomancak P. 2010. Gene expression divergence recapitulates the

developmental hourglass model. Nature 468:811–814.

Kass KE, Raftery AE. 1995. Bayes Factors. J Am Stat Assoc 90:773–795.

Katoh K, Toh H. 2008. Recent developments in the MAFFT multiple sequence alignment

program. Brief Bioinform 9:286–298.

Khaitovich P. 2005. Parallel patterns of evolution in the genomes and transcriptomes of

humans and chimpanzees. Science (80-) 309:1850–1854.

Khaitovich P, Pääbo S, Weiss G. 2005. Toward a neutral evolutionary model of gene

expression. Genetics 170:929–939.

Khaitovich P, Weiss G, Lachmann M, Hellmann I, Enard W, Muetzel B, Wirkner U,

Ansorge W, Pääbo S. 2004. A neutral model of transcriptome evolution. PLoS Biol

2.

Kozak KM, Wahlberg N, Neild AFE, Dasmahapatra KK, Mallet J, Jiggins CD. 2015.

Multilocus species trees show the recent adaptive radiation of the mimetic

Heliconius. butterflies. Syst Biol 64:505–524.

Kronforst MR, Papa R. 2015. The functional basis of wing patterning in *Heliconius*

butterflies: The molecules behind mimicry. Genetics 200:1–19.

Lande R. 1976. Natural Selection and Random Genetic Drift in Phenotypic Evolution.

Evolution (N Y) 30:314–334.

Lang GI, Murray AW, Botstein D. 2009. The cost of gene expression underlies a fitness

trade-off in yeast. Proc Natl Acad Sci 106:5755–5760.

Larracuente AM, Sackton TB, Greenberg AJ, Wong A, Singh ND, Sturgill D, Zhang Y,

Oliver B, Clark AG. 2008. Evolution of protein-coding genes in Drosophila. Trends

Genet 24:114–123.

Lee G, Foss M, Goodwin SF, Carlo T, Taylor BJ, Hall JC. 2000. Spatial, temporal, and

sexually dimorphic expression patterns of the fruitless gene in the *Drosophila*

central nervous system. J Neurobiol 43:404–426.

McCulloch KJ, Yuan F, Zhen Y, Aardema ML, Smith G, Llorente-Bousquets J,

Andolfatto P, Briscoe AD. 2017. Sexual Dimorphism and Retinal Mosaic

Diversification Following the Evolution of a Violet Receptor in Butterflies. Mol

Biol Evol 34:1–14.

Mendoza-Cuenca L, Macías-Ordóñez R. 2005. Foraging polymorphism in Heliconius

charitonia (Lepidoptera: Nymphalidae): morphological constraints and behavioural

compensation. J Trop Ecol 21:407–415.

Merrill RM, Dasmahapatra KK, Davey JW, Dell'Aglio DD, Hanly JJ, Huber B, Jiggins

CD, Joron M, Kozak KM, Llaurens V, et al. 2015. The diversification of *Heliconius*

butterflies: What have we learned in 150 years? J Evol Biol 28:1417–1438.

Metropolis N, Rosenbluth AW, Rosenbluth MN, Teller AH, Teller E. 1953. Equation of

state calculations by fast computing machines. J Chem Phys 21:1087–1092.

Metzger BPH, Duveau F, Yuan DC, Tryban S, Yang B, Wittkopp PJ. 2016. Contrasting

Frequencies and Effects of *cis-* and *trans*-Regulatory Mutations Affecting Gene

Expression. Mol Biol Evol 33:1131–1146.

Montgomery SH, Merrill RM, Ott SR. 2016. Brain composition in *Heliconius* butterflies,

posteclosion growth and experience-dependent neuropil plasticity. J Comp Neurol

524:1747–1769.

Nadeau NJ. 2016. Genes controlling mimetic colour pattern variation in butterflies. Curr

Opin Insect Sci 17:24–31.

Nitta KR, Jolma A, Yin Y, Morgunova E, Kivioja T, Akhtar J, Hens K, Toivonen J,

Deplancke B, Furlong EEM, et al. 2015. Conservation of transcription factor

binding specificities across 600 million years of bilateria evolution. Elife 2015:1–20.

Nourmohammad A, Rambeau J, Held T, Kovacova V, Berg J, Lässig M. 2017. Adaptive

evolution of gene expression in *Drosophila*. Cell Rep 20:1385–1395.

O'Meara BC, Ané C, Sanderson MJ, Wainwright PC. 2006. Testing for different rates of

continuous trait evolution using likelihood. Evolution (N Y) 60:922–933.

Punta M, Coggill P, Eberhardt R, Mistry J, Tate J, Boursnell C, Pang N, Forslund K,

Ceric G, Clements J, et al. 2012. The Pfam protein families databases. Nucleic Acids

Res 30:1–12.

Rifkin SA, Kim J, White KP. 2003. Evolution of gene expression in the *Drosophila*

melanogaster subgroup. Nat Genet 33:138–144.

Rohlfs R V., Harrigan P, Nielsen R. 2014. Modeling gene expression evolution with an

extended Ornstein-Uhlenbeck process accounting for within-species variation. Mol

Biol Evol 31:201–211.

Rohlfs R V., Nielsen R. 2015. Phylogenetic ANOVA: The expression variance and

evolution model for quantitative trait evolution. Syst Biol 64:695–708.

Romero IG, Ruvinsky I, Gilad Y. 2012. Comparative studies of gene expression and the

evolution of gene regulation. Nat Rev Genet 13:505–516.

Salcedo C. 2010. Environmental Elements Involved in Communal Roosting in *Heliconius*

Butterflies (Lepidoptera: Nymphalidae). Environ Entomol 39:907–911.

Signor SA, Nuzhdin S V. 2018. The evolution of gene expression in *cis* and *trans*. Trends

Genet:1–13.

Sinharay S, Stern HS. 2002. On the sensitivity of Bayes factors to the prior distributions.

Am Stat 56:196–201.

Stamatakis A. 2006. RAxML-VI-HPC: Maximum likelihood-based phylogenetic

analyses with thousands of taxa and mixed models. Bioinformatics 22:2688–2690.

Stern DB, Crandall KA. 2018. The Evolution of Gene Expression Underlying Vision

Loss in Cave Animals. Mol Biol Evol 35:2005–2014.

Uyeda JC, Harmon LJ. 2014. A novel Bayesian method for inferring and interpreting the

dynamics of adaptive landscapes from phylogenetic comparative data. Syst Biol

63:902–918.

Wagner A. 2005. Energy constraints on the evolution of gene expression. Mol Biol Evol

22:1365–1374.

Wagner GP, Altenberg L. 1996. Complex Adaptations and the Evoluation of

Evolvability. Evolution (N Y) 50:967–976.

Walters JR, Stafford C, Hardcastle TJ, Jiggins CD. 2012. Evaluating female remating

rates in light of spermatophore degradation in *Heliconius* butterflies: Pupal-mating

monandry versus adult-mating polyandry. Ecol Entomol 37:257–268.

Wanner KW, Anderson AR, Trowell SC, Theilmann DA, Robertson HM, Newcomb RD.

2007. Female-biased expression of odourant receptor genes in the adult antennae of

the silkworm, *Bombyx mori*. Insect Mol Biol 16:107–119.

Warnefors M, Eyre-Walker A. 2012. A selection index for gene expression evolution and

its application to the divergence between humans and chimpanzees. PLoS One 7.

Whitehead A, Crawford DL. 2006a. Variation within and among species in gene

expression: Raw material for evolution. Mol Ecol 15:1197–1211.

Whitehead A, Crawford DL. 2006b. Neutral and adaptive variation in gene expression.

Proc Natl Acad Sci 103:5425–5430.

Wong ES, Thybert D, Schmitt BM, Stefflova K, Odom DT, Flicek P. 2015. Decoupling

of evolutionary changes in transcription factor binding and gene expression in

mammals. Genome Res 25:167–178.

1   **Supplementary figures**

Figure S1. Orthoclusters with conserved gene expression values across five *Heliconius*

species. The orthoclusters with σ^2 not significantly larger than zero from Figure 4 were used

here.

Figure S2. Simulation study for the assessment of parameter estimation bias under an OU-

model. Parameter (σ^2) bias estimation through 1000 simulations under sigma values ranging

from 0.1 – 10. Simulations were performed for phylogenies with sizes ranging from 5 to 1000

taxa.

Figure S3. Model probability when testing for directional selection. Significance is shown at

model probability > 0.75 (solid red, Bayes factor > 3, positive support) and model probability

> 0.95 (solid red, Bayes factor > 20, strong support).

21 **Figure S4.** Venn Diagram depicting the number of genes showing a branch specific shift in

22 σ^2 tested by an BM (blue) model and those genes showing a shift in gene expression optimum

23 assessed with an OU model (red). In parenthesis are shown the number of genes contributing

24 to the overlapping region identified by each of the method.

Figure S5. Orthoclusters where a shift in gene expression optima was detected along a particular terminal *Heliconius* species branch. Panels show shifts in the HER branch (A), in HSA (B), in HCH (C), in HDO (D) and in the HME branch (E).

Figure S6. Examples of branch specific shifts in evolutionary rates for each of the five

Heliconius species. Boxplots depict the distribution of log$_2$ transformed FPKM expression

values. Horizontal solid line indicates median expression level. Shift in: *H. erato*:

OrthoGroup_1567_clean, annotated as protein domain with unknown function (Appendix 1).

H. sara: OrthoGroup_530_clean, annotated as vinculin family involved in cell adhesion and

cytoskeleton. *H. charithonia*: OrthoGroup_1276_clean, annotated as photoreceptor

hydrogenase. *H. doris*: OrthoGroup_449_clean, annotated as Tropomyosin 1, involved in

dendrite morphogenesis.

Figure S6. *Heliconius* orthoclusters where an orthology was identified with zero expression

values. A total of 254 orthoclusters were found.