PerSort facilitates characterization and elimination of persister subpopulation in mycobacteria

Vivek Srinivas[1], Mario L. Arrieta-Ortiz[1], Eliza J.R. Peterson[1*], Nitin S. Baliga[1-4*]

[1]Institute for Systems Biology, Seattle, WA, USA
[2]Departments of Biology and Microbiology, University of Washington, Seattle, WA, USA
[3]Molecular Engineering Program, University of Washington, Seattle, WA, USA
[4]Lawrence Berkeley National Lab, Berkeley, CA, USA

*Corresponding authors
Eliza J.R. Peterson: eliza.peterson@isbscience.org;
Nitin S. Baliga: nitin.baliga@isbscience.org (lead contact)

Summary

We have developed a novel method, PerSort, to isolate translationally dormant cells that pre-exist in small numbers within *Mycobacterium spp.* cultures growing under naïve conditions (i.e., absence of antibiotic treatment), but dramatically increase in proportion under stress conditions. The translationally dormant cells have high tolerance to isoniazid and rifampicin, and can regenerate the parental population structure in standard media, albeit after a significantly longer lag phase, indicating they are persister cells. Single-cell expression profiling demonstrated that the translationally dormant persister subpopulation is a mixture of *vapC30*, *mazF*, and *relA/spoT* overexpressing cells, indicating there are multiple pathways to become a persister cell. Regardless of the mechanism by which they are generated, the persister cells have reduced oxidative metabolism, which is reversed upon addition of L-cysteine to effect complete clearance by INH and RIF under host-related stress.

Abstract

Mycobacterium tuberculosis (MTB) generates phenotypic diversity to persist and survive the harsh conditions encountered during infection. MTB avoids immune effectors and antibacterial killing by entering into distinct physiological states. The surviving cells, persisters, are a major barrier to the timely and relapse-free treatment of tuberculosis (TB). We present for the first time, PerSort, a method to isolate and characterize persisters in the absence of antibiotic, or other pressure. We demonstrate the value of PerSort to isolate translationally dormant cells that pre-exist in small numbers within *Mycobacterium spp.* cultures growing under optimal conditions, but which dramatically increased in proportion under stress conditions. The translationally dormant subpopulation exhibited multidrug tolerance and regrowth properties consistent with persister cells. Furthermore, PerSort enabled single-cell transcriptional profiling that provided evidence that the translationally dormant persisters were generated through a variety of mechanisms, including *vapC30*, *mazF*, and *relA/spoT* overexpression. Finally, we demonstrate that notwithstanding the varied mechanisms by which the persister cells were generated, they converge on a similar low oxygen metabolic state that was reversed through activation of respiration to rapidly eliminate persisters fostered under host-relevant stress conditions. We conclude that PerSort provides a new tool to study MTB persisters, enabling targeted strategies to improve and shorten the treatment of TB.

Keywords: Mycobacterium, phenotypic heterogeneity, persisters, antibiotic tolerance, nutrient starvation

Introduction

Studies have revealed cell-to-cell variability in organisms from all domains of life –unicellular to multicellular. Through cell-to-cell variability, cell populations are prepared for sudden environmental changes by harbouring subpopulations that are phenotypically pre-adapted. This evolutionary strategy, known as bet-hedging (1), confers fitness advantage to pathogens, such as *Mycobacterium tuberculosis* (MTB), which routinely experience varying environments within the host. In addition to spontaneous bet-hedging, MTB responds to host derived stresses with physiological changes (e.g., shifts in metabolism and respiration, induction of toxin-antitoxin systems, cell wall modifications) that allow it to survive and persist. These physiological changes (either stochastically- or environmentally-induced) result in antibiotic tolerance, in which MTB is genetically susceptible to antibiotics but exists in a physiological state rendering it refractory to drug killing. These persistent states are a major reason why long courses of antibiotic therapy are required to treat human tuberculosis (TB) (2); standard chemotherapy of TB requires 6 months of treatment and 5% are not cured even then (3, 4). There is an urgent need for new

strategies that shorten the duration of treatment and target drug tolerant MTB. Addressing this
gap requires a better understanding of how MTB generates phenotypic heterogeneity to
withstand pressures from the host environment and evade antibiotic therapy.

Recent work has studied phenotypic heterogeneity in MTB from single-cell analyses
using fluorescent reporter strains. Cell-to-cell variability of MTB was captured *in vitro* and
during murine infections using a reporter of 16s rRNA gene expression (5). The microscopy-
based platform was able to track heterogeneity in growth rate under standard growth
conditions and found heterogeneity was amplified by stress conditions and murine infection.
However, the non-growing subpopulation was not isolated and characterized further, most
likely due to low fluorescence levels of the reporter. In another study, Jain *et al* developed a
dual-reporter mycobacteriophage (φ^{2}DRM) system and used fluorescent activated cell sorting
(FACS) to isolate drug tolerant MTB cells from *in vitro* cultures and human sputa (6). However,
the necessity to re-infect daughter cells with φ^{2}DRM limited the ability to follow isolated cells
over generations and study their regrowth patterns. The study also used a reporter that
specifically enriched for persisters of isoniazid treatment (i.e., fluorescent protein fused to the
dnaK promoter), disregarding multidrug tolerant subpopulations which are known to exist
within the host environment (7).

Here, we sought to develop a fluorescent reporter system to isolate and characterize
multidrug tolerant subpopulations of mycobacteria from naïve growth conditions (i.e., absence
of antibiotic treatment). We wanted to avoid killing susceptible cells with antibiotics in order to
study both persister cells *and* actively growing cells from the same culture and prevent drug
pressure from confounding persister formation. Instead, based on a considerable body of work
linking bacterial stress pathways to the acquisition of drug tolerance via translation inhibition
(8-13), we developed "PerSort" to enrich for translationally dormant mycobacterium (i.e., cells
that are able to transcribe but not translate a fluorescent reporter) under naïve conditions. We
demonstrated the persister-like properties of the translationally dormant subpopulation and
their increased abundance during stress conditions. Moreover, we performed single-cell
transcriptional profiling of "PerSorted" cells and highlight various mechanisms that generate
translationally dormant mycobacteria. Finally, using these mechanistic insights we revealed a
physiological response common to persister cells and demonstrated that activating respiration
potentiated both isoniazid and rifampicin to rapidly clear mycobacteria. Importantly, this was
achieved from a host-relevant condition that fosters multidrug tolerant mycobacteria. This
study confirms that PerSort enables a better understanding of mycobacterial phenotypic
heterogeneity and can lead to new strategies for shortening TB treatment.

103
Results

Multidrug tolerant mycobacteria increase in nutrient starved conditions and show lag-
dormancy
To establish a system that enriches for multidrug tolerant mycobacteria, we grew *M.*
smegmatis (MSM) in nutrient rich (7H9 media, 0.2% glycerol, 0.05% Tween-80, and ADC
complementation) and nutrient starved (PBS and 0.05% Tween-80) conditions. Deprivation of

nutrients results in a marked slowing of mycobacterial growth and concurrent phenotypic tolerance to various antibiotics, including isoniazid (INH) and rifampicin (RIF) (14, 15). Moreover, studies have established the relevance of the bacterial physiological state adopted under nutrient starved conditions to TB infection (16). We incubated wild type MSM mc^{2}155 strain in nutrient rich or nutrient starved conditions for 12 hours (h) then treated with either 5x minimum inhibitory concentration (MIC) INH or RIF (40 and 15 µg/mL respectively). Antibiotic-treated MSM cultures were diluted and plated over a 20 h interval. Percent survival was calculated, in reference to time 0 h, and used to generate kill curves (**Figure 1a, 1b**). According to the bimodal model of bacterial persistence (17), multiple subpopulations were identified, either drug susceptible or drug tolerant, as distinguished by varying slopes of killing. We further estimated the abundance of drug tolerant subpopulations in either nutrient rich or nutrient starved conditions via the intersection of the y-axis with the extrapolated slope of the drug tolerant subpopulation. Thus, MSM cultures grown in standard conditions (i.e., nutrient rich) were ~45% tolerant to INH and ~30% tolerant to RIF. Whereas after nutrient starvation, the proportion of the drug tolerant subpopulations increased to ~75% for INH and ~50% for RIF. The increase in delay of RIF killing under nutrient starved conditions, might reflect cell wall alterations that are required for adaptation to nutrient deprivation, but also restrict entry of RIF (18, 19).

Another advantage of the nutrient deprivation model for enriching and characterizing multidrug tolerant mycobacteria is that nutrient starved bacteria can easily grow upon being returned to nutrient rich media, thus this model allows easy quantification of drug susceptibility and growth patterns. We characterized the population wide growth characteristics of the nutrient rich and nutrient starved cultures using ScanLag, a technique that combines cell plating with high-throughput imaging (20). Following 12 h incubation in nutrient rich or nutrient starved conditions, MSM mc^{2}155 cultures were plated onto nutrient rich plates (7H10 media, 0.2% glycerol, OADC) and observed for colony growth by imaging at 1 h intervals. The mean time of appearance (TOA) of individual colony forming units (CFUs) from the nutrient rich conditions was determined to be 37 h, whereas the TOA from the nutrient starved conditions was 41 h (**Figure 1c**). This significant delay in resuming growth is due to a greater abundance of cells with lag-dormancy, a phenotype well-established with drug tolerance (21, 22), which explains why nutrient deprived mycobacteria are multidrug tolerant.

PerSort isolates translationally dormant mycobacteria that are multidrug tolerant

To further characterize multidrug tolerant subpopulations, we developed a FACS-based protocol to isolate translationally dormant mycobacterial cells. In drug tolerance inducing conditions, translation is repressed via various mechanisms that downregulate rRNA (e.g. RelA and CarD) or degrade rRNA (e.g. VapC and other toxins) [10-15]. Therefore, we hypothesized

that a fluorescent reporter that relays translation activity could efficiently sort and isolate drug
tolerant mycobacterium that form in a multitude of ways. The reporter plasmid (Trans-mEos2,
Figure S1) was constructed by inserting the highly stable fluorescent gene, *mEos2*, under the
transcriptional control of an anhydrotetracycline (ATc)-inducible promoter and a strong
mycobacterial translation initiation sequence (Shine-Dalgarno sequence), then inserted into the
PstKi plasmid (23-25). The PstKi plasmid contains an integrase gene and thus eliminates copy
number variability among cells by integrating into the mycobacterium genome. Moreover, the
ATc-inducible promoter achieves detectable fluorescence from a genome-integrated reporter
while also allowing for modulation of reporter expression.
The Trans-mEos2 plasmid was transformed into wild type MSM (creating the MSM-
mEos2 strain) for differentiation of mEos2 fluorescence levels in mycobacterial cells using FACS.
To minimize clumps, MSM-mEos2 cultures (grown in 7H9, 0.2% glycerol, 0.05% Tween-80, ADC
complementation and high shaking conditions) were passed through a 10 µm filter before
running on the BD FACS Influx instrument. Parameters for single-cells were defined by 3 µm
rainbow fluorescent beads and the efficiency of single-cell sorting was assessed using a mixed
culture of ATc-induced MSM-mEos2 cells (resistant to kanamycin) and MSM cells expressing
mCherry (resistant to hygromycin) (26). Cells from the mixed fluorescent reporter cultures were
sorted and plated onto 7H10 plates with appropriate antibiotic selection. Sorting was found to
be 93% efficient (**Figure S2**).
Sort gates were defined for cultivable and non-cultivable cells (*i.e.*, dead cells and viable
but non-culturable cells). Viable but non-culturable cells (VBNCs) are antibiotic tolerant, but
they lack the ability to regain active growth in standard conditions and therefore do not
contribute to bacteria survival (27). To exclude non-cultivable cells, MSM-mEos2 cultures were
stained with propidium iodide (PI) and PI signal was used for gate selection. Heat-killed cultures
were used for evidence of non-cultivable cells in the PI(+) gate (**Figure S3)**. Gates for
translationally active and dormant cells were defined based on mEos2 fluorescence signal in the
PI(-) cultivable population (**Figure 2a**). Wild type MSM and MSM-mEos2 cultures were grown in
the presence of ATc (500 ng/mL) for 12 h and stained with 0.5% PI. Background fluorescence
from wild type MSM cells was used to define the gate for translationally dormant cells ("dim").
For clear distinction between subpopulations, gates were set at a higher mEoS2 fluorescence
signal to define MSM-mEos2 translationally active cells ("lit"), leaving ~25% of PI(-) cells
unselected (**Figure 2a**). Further, the PI signal gates were set such that thresholds for selection of
cultivable cells, PI(-), decreased with the mEos2 signal, as VBNCs were expected at high
proportions in low mEos2 signal range. Based on these FACS gates and following ATc-induction
of MSM-mEos2 cultures in standard 7H9 media (*i.e.*, same as nutrient rich conditions) for 12 h
(OD600 > 1) , a majority of cells were translationally active "lit" cells, while a consistent
subpopulation of translationally dormant "dim" cells was also present, reaching about 1% of
the population (**Figure 2a**). We sorted equal number of cells from the dim, lit, and PI(+) gates

and plated onto 7H10 media (**Figure 2b**). All of the cells from the dim and lit subpopulations
formed CFUs, whereas only 25% formed CFUs from the PI(+) subpopulation, confirming cells
from the PI(-) negative (i.e., dim and lit) gates were cultivable and devoid of VBNCs and dead
cells. Finally, we measured dim cell proportions following ATc-induction of MSM-mEos2
cultures in standard growth conditions until cultures were either in exponential (~6 h after ATc
addition) or stationary (12 h after ATc addition) phase of growth. Dim cells were found in low
numbers (~0.4% of population) during exponential phase and increased during stationary phase
to make up ~1% of the population (**Figure S4**). This indicates that dim cells pre-exist under low
stress conditions (i.e., exponential phase) and a density-dependent increase in dim cell
formation, likely caused by nutrient depletion of the culture during stationary phase (28).
Having developed the fluorescent reporter system and optimized the FACS procedure to
sort translationally active and dormant cells from naïve growth conditions (i.e., absence of
antibiotic treatment), we investigated the growth characteristics and drug susceptibility of the
dim and lit subpopulations. For growth characterization, cells were sorted from the PI(-) gate of
ATc-induced MSM-mEos2 cultures and ScanLag analysis was performed as described previously
(**Figure 2c**). The dim cells showed a longer TOA compared to lit cells, with a difference between
the subpopulations (~200 minutes) similar to that observed between nutrient rich and nutrient
starved cultures (**Figure 1c**). As expected, cells from the unselected PI(-) gate had a mean TOA
in between the dim and lit subpopulations, suggesting it is almost an equal mixture of
translationally active and inactive cells. To assess drug susceptibility, 100,000 dim and lit cells
from ATc-induced MSM-mEos2 cultures were sorted into 7H9 media containing either 5x MIC
INH or 5x MIC RIF. The survival of dim and lit subpopulations was determined by comparing
CFUs after 12 h INH or 8 h RIF treatment to CFUs before drug treatment (**Figure 2d, 2e**). The
drug treatment times were selected from the slowed phase of killing from the biphasic kill
curves (**Figure 1a, 1b**). A significantly larger percentage of dim cells survived both INH (p-value =
0.001) and RIF treatment (p-value = 0.006), indicating the dim cells are a multidrug tolerant
subpopulation. Further, the survival of dim cells was compared to the entire population of ATc-
induced MSM-mEos2 cultures (i.e., unsorted) grown in either nutrient rich or nutrient starved
conditions. The percentage of dim cells surviving INH and RIF treatment matched closely with
the nutrient starved cultures. These results reveal the striking similarities, both in terms of
growth characteristics (i.e., lag-dormancy) and antibiotic susceptibility (i.e., multidrug
tolerance), between dim cells and nutrient starved cultures. As such, we hypothesized the dim
cells are a phenotypically heterogeneous subpopulation, with translational dormancy and
multidrug tolerance, that increase in proportion under nutrient deprivation because their
physiological state is best suited to withstand such environmental stress.
222

Translationally dormant mycobacteria form at different probabilities in naïve versus nutrient starved conditions

We used the demonstrated capability of PerSort to isolate persister-like mycobacterial cells to investigate the population structure and regrowth characteristics of subpopulations from both naïve (i.e., absence of antibiotic treatment) and nutrient starved conditions. Following ATc induction of the MSM-mEos2 culture for 12 h, 100,000 dim and lit cells were PerSorted into standard 7H9 media. Sorted samples were grown to an OD600 of 0.6, induced with ATc again for 12 h and reanalyzed by FACS. Cultures generated from both dim and lit subpopulations had similar structure as the parent population vis-à-vis proportions of dim and lit cells (**Figure 3a**). These results demonstrate that translational dormancy of dim cells is not heritable and dim and lit cells interconvert under standard *in vitro* growth conditions. We further assessed regrowth by measuring OD600 over time after PerSorting 100,000 dim and lit cells into standard 7H9 media. As expected, dim cell cultures grew after a longer lag phase than lit cells (**Figure 3b**), but the subpopulations reached similar maximum growth rate and carrying capacity (**Figure 3c**). This demonstrates that dim cells can resume normal growth (*i.e.*, same as lit cells) in low stress conditions, which is a key characteristic of persister cells (29). Further, ATc-induced MSM-mEos2 cultures were grown in either naïve or nutrient starved conditions and then PerSorted to determine the population structure. Both dim and lit cells were present in the nutrient starved cultures, but there was a significant increase in the proportion of dim cells (**Figure 3d**). This confirmed our hypothesis, establishing that dim cells (i.e., translationally dormant mycobacteria) dramatically increase in proportion upon nutrient starvation, further suggesting that the likelihood of generating a dim cell in each cell division is distinct under low stress versus high stress conditions (**Figure 3e**). Whereas dim cells (and likely other phenotypically heterogeneous subpopulations) form at low probability under naïve conditions, nutrient deprivation favors the formation of dim cells (a translationally dormant and multidrug tolerant subpopulation) and enables mycobacteria to withstand stress, including drug treatment.

Translationally dormant mycobacteria are composed of three discernable subtypes of *vapC30, mazF* toxins and(or) *relA/spoT* over-expressing cells

To further characterize the formation of phenotypically heterogeneous mycobacteria, we profiled within individual cells of both dim and lit subpopulations following PerSort, the transcript levels of 45 genes that were previously implicated in persister formation and drug tolerance in *Mycobacterium* spp. and *E. coli* (**Table S1**). Single-cell gene expression profiling was performed with the Fluidigm Biomark 48x48 system per manufacturer instructions and assayed relative to single-cell genomic DNA signal. The DNA signal from single dim and lit cells was

found to have low variation in expression (**Figure S5a**), indicating that transcripts per genome
copy accurately reports differential gene expression. Transcript abundances were normalized to
a spike-in RNA control to account for experimental noise (**Figure S5b**).

Kernel PCA (kPCA) with radial basis function identified distinct clusters consisting of dim
and lit cells as well as some overlap between the subpopulations (**Figure S6**). The overlap could
indicate some phenotypic uncertainty, based on the expression of selected persister genes. We
used a tree-based feature selection (see *Methods*) to rank the persister genes for their ability to
differentiate the dim and lit subpopulations (30). Using the top features (**Table S2**), we
performed unsupervised hierarchical clustering and dynamic tree cutting to identify four
distinct clusters within the dim cells (**Figure 4a**) (31, 32). Specifically, clustering revealed
subtypes of translationally dormant mycobacteria with high *relA/spoT*, *vapC30*, or *mazF*
expression (and another subtype with no distinct signature of persister gene expression).
Importantly, similar clustering analysis did not detect any statistically significant subtypes
within the lit cells (**Figure 4a**). Increased expression of *relA/spoT* was observed in higher
proportions of dim cells, compared to lit cells. The bifunctional activity of the protein (*i.e.,*
hydrolase and synthase) (33) could suggest different functions of RelA/SpoT between the dim
and lit subpopulations.

We sought further evidence that one or some combination of the three gene features
identified from single-cell expression profiling (i.e., RelA/SpoT, VapC30, or MazF) were active in
the dim cell subpopulation. The VapC30 and MazF toxins belong to a family of type II toxin-
antitoxin (TA) systems, implicated in mycobacterial dormancy and persistence (28, 34, 35).
Activation of the type II TA system results in toxin-mediated cleavage of tRNAs (36), or rRNAs
(12, 36) causing translational dormancy (37). We analyzed 16S and 23S rRNA levels in dim and
lit cells by qRT-PCR. We discovered that indeed dim cells have lower rRNA content relative to lit
cells (**Figure 4b**, 16s rRNA log2 normalized FC = -4.17, 23s rRNA log2 normalized FC = -2.34).
There was no discernible decrease in transcript level of a highly expressed metabolic gene
(**Figure 4b**, phosphoglucoisomerase log2 FC = -0.87). These results support the notion that
some of the translationally dormant mycobacteria are formed via toxin-mediated cleavage of
rRNA.

Translationally dormant mycobacteria are in a low O$_2$ respiratory state

The single-cell expression data revealed that translationally dormant mycobacteria may
be formed by at least three distinct mechanisms. The multiplicity of mechanisms confers
robustness to the pathogen, but also thwarts therapeutic strategies to block formation of the
multidrug tolerant mycobacteria. Nonetheless, we found evidence from literature that the
mechanisms for persister formation converge on a common physiology and predicted the dim
cells are in a reduced state of respiration. Direct measurements of respiration rate (e.g., oxygen

electrode or Seahorse analysis) are problematic with low bacterial cell numbers. Therefore, to
test whether dim cells have lower O_2 metabolism we measured reactive oxygen species (ROS)
levels in single-cells from the PerSorted dim and lit subpopulations using CellRox-orange. It is
established that increases in ROS levels are linked with increased flux to the TCA cycle and
oxidative metabolism (38, 39). Cultures of ATc-induced MSM-mEos2 were grown in standard
7H9 media, stained with CellRox-orange (40), and PerSorted to compare ROS levels between
dim and lit cells (**Figure 4c**). As reference controls, MSM-mEos2 cultures were treated with Mn-
TBAP (ROS inducer) or N-acetyl-cysteine (ROS quencher) with Mn-TBAP. Cultures were
incubated in CellRox-orange stain for 1.5 h to ensure uniform dye permeation across the
subpopulations. The ROS levels were significantly lower in dim cells (**Figure 4c**), confirming the
translationally dormant mycobacteria have reduced oxidative metabolism.

Activation of oxidative metabolism eliminates translationally dormant mycobacteria and
achieves faster killing by INH and RIF in nutrient starved conditions

Since respiration may play an important role in generating and/or maintaining
translationally dormant mycobacteria, we explored the use of L-cysteine to activate oxidative
metabolism and reduce the proportion of dim cells. It is well-known that exogenous amino
acids fuel the TCA cycle and oxidative respiration in bacteria, especially molecules such as
cysteine and proline which are rapidly metabolized (41). Cultures of MSM-mEos2 were grown in
standard 7H9 media with 4mM L-cysteine (similar growth observed in 7H9 media with or
without 4 mM L-cysteine, data not shown), induced with ATc and PerSorted for dim and lit
subpopulations. The L-cysteine treated MSM-mEos2 cultures were found to be completely
devoid of dim cells (**Figure 5a**). Further, we supplemented nutrient starved MSM cultures with
4 mM L-cysteine and then treated with either 5x MIC INH or 5x MIC RIF. Kill-curves for the
antibiotic and L-cysteine treated cultures were generated as previously described (**Figure 5b, c**).
Activation of respiration by adding L-cysteine to nutrient starved cultures potentiated clearance
by INH and RIF at rates identical to or better than rates for nutrient rich cultures. This further
corroborates that, similar to dim cells, the expanded translationally dormant mycobacteria in
nutrient starved conditions share a low respiratory physiologic state. Ultimately, these results
demonstrate that drug adjuvants that activate respiration could shorten drug treatment,
particularly in environmental niches that foster drug tolerance in mycobacteria.

Discussion

The ability of microorganisms to survive sudden environmental changes stems from the
formation of phenotypically heterogeneous subpopulations. Phenotypic heterogeneity confers
fitness advantage to clonal microbial communities, such as infectious MTB, but impedes efforts

of the immune system to clear the pathogen as well as chemotherapeutic efforts to rapidly
treat TB. In this study, we developed a method to identify and characterize multidrug tolerant
subpopulations of mycobacterium cells. We demonstrated that our method enables a better
understanding of phenotypic heterogeneity in mycobacteria in naïve and stressed conditions,
and could lead towards novel strategies to shorten TB treatment.

We generated a fluorescent reporter system, PerSort, which sorts mycobacterial cells
based on translational activity. We found a small proportion of translationally dormant ("dim")
cells from naïve cultures, in the absence of stress such as antibiotic treatment. We confirmed
the translationally dormant subpopulation was tolerant to both INH and RIF, demonstrated a
longer lag phase upon regrowth, and could regenerate the original population structure upon
regrowth (i.e., a mixture of translationally active and dormant cells). These data indicate the
translationally dormant subpopulation identified by PerSort consist of cells with properties of
persisters. We report these cells pre-exist in low numbers in an isogenic MSM culture growing
without stress and exponential phase of growth. The translationally dormant subpopulation
(along with other phenotypically heterogeneous subpopulations) is generated stochastically, as
a bet-hedging strategy, for the mycobacterial population to withstand unpredictable
environmental stress.

Indeed, we discovered an increase in the proportion of the translationally dormant
subpopulation under density-dependent stress, and even more so under nutrient starvation.
This suggests an environmentally induced component that forms this subpopulation, in addition
to stochastic formation. It also begs the question of what is influencing the translational state of
this mycobacterial subpopulation. The PerSort method overcomes the challenge of
characterizing persisters at single cell resolution by sorting them and, importantly, not killing
the susceptible cells with antibiotics. This technological advancement has overcome the
confounding issue that antibiotic treatment itself induces persister cell formation (42, 43) and
allows comparative analysis of persister cells *and* actively growing drug susceptible cells from
the *same* culture. Because of these novel capabilities of PerSort, we were able to quantify
transcript abundance of 45 genes associated with persister formation and drug tolerance in
single-cells from both the translationally dormant (i.e., dim) and translationally active (i.e., lit)
subpopulations. Expression analysis revealed that translationally dormant persisters are a mix
of *vapC30*, *mazF*, and *relA/spoT* overexpressing cells. These results reinforce the hypothesis
that there are multiple pathways (both stochastically and deterministically activated) to
become a persister cell and reveals the complex and combinatorial schemes used by
mycobacteria to generate heterogeneous subpopulations.

Within the translationally dormant cells, we found that high expression of toxin *mazF*
was also associated with high *relA/spoT* expression. We suspect that *mazEF* could be regulated
by the alarmone response, elicited by RelA/SpoT synthesis of (p)ppGpp, in a manner that is
induced deterministically by stress conditions (44). In contrast, the single-cell expression data

suggest that *vapC30* overexpression can also act to induce persister formation, in a
spontaneous and alarmone-independent manner. This supports a recent study demonstrating
that a *relA/spoT* knockout mutant of MSM, with reduced alarmone response, still formed
persisters at levels similar to wildtype (45). This collection of evidence points toward multiple
mechanisms of generating the translationally dormant mycobacteria characterized here, some
of which are deterministically activated (controlled by *relA/spoT* in response to stress) and
some of which are stochastically activated (controlled by spontaneous *vapC30* activation) (46).
While MSM has a single VapBC-type TA system, MTB has 70 copies of VapBC (35), indicating the
human pathogen has evolved to increase phenotypic diversity and bet-hedging for survival in
the host environment. Much work is still needed, specifically using live cell monitoring
techniques, to understand how and when these TA systems are activated to form persisters,
and the contribution of other mechanisms, either stochastic or deterministic, to phenotypic
heterogeneity in MTB.

The activation of TA systems (47) and the alarmone response (48) has been shown to
decrease oxidative metabolism in bacterial persisters. Moreover, it is expected that
translational dormancy would be associated with low respiration (28). Given this knowledge,
we demonstrated lower ROS levels in the translationally dormant subpopulation compared to
translationally active cells, indicating reduced oxygen metabolism in the persister cells. In other
words, regardless of their mechanism of formation (i.e., *vapC30*, *mazF*, or *relA/spoT*
overexpression), the persister subpopulation shares a low oxygen respiratory state, which
presents a vulnerability that could be targeted to modulate this subpopulation (**Figure 6**). We
confirmed that addition of L-cysteine, which is known to activate oxidative metabolism (49),
dramatically reduced the proportion of translationally dormant cells in nutrient starved MSM
cultures, conditions where the population is abundant. This goes beyond previous studies by
directly demonstrating that promoting oxidative metabolism reduces the proportion of
multidrug tolerant persister cells that form independent of drug pressure. Furthermore, the
addition of L-cysteine was able to effect complete clearance by INH and RIF in nutrient starved
conditions. The addition of L-cysteine potentiates faster drug killing by converting (or limiting
the formation of) multidrug tolerant persister cells, cells with enhanced fitness advantage in the
host-relevant stress (nutrient starved) conditions, to a population with increased oxidative
metabolism and drug susceptibility. Our findings prove that until properties of heterogeneous
subpopulations are disrupted, we will not be able to successfully clear mycobacterial cells in
infected patients. This study highlights how novel methods to isolate and characterize
heterogeneous subpopulations can enable targeted strategies to eliminate detrimental
persister cells, and thereby shorten the course of treatment.
410
411 **Materials and Methods**
412

Bacterial growth and MSM-mEos2 strain development. *Mycobacterium smegmatis* mc^{2}155 strain (MSM) was obtained from ATCC and grown in 7H9 broth medium (Difco) with 0.2% glycerol, 0.05% Tween-80, and 10% ADC enrichment (BD biosciences). pSTKi-mEos2 plasmid was constructed from pSTKi plasmid and pRSETa mEos2 plasmid (23, 25). Synthetic oligo with translation initiation signal (mycoSD) was used to amplify mEos2 from pRSETa mEos2 plasmid, and amplified fragment was inserted into the pSTKi plasmid with restriction ligation at BamH1 and EcoR1 sites. The pSTKi-mEos2 transcript was electroporated into electrocompetent MSM cultures, transformed colonies were selected on 7H10 plates with 30 μg/ml kanamycin (KAN). MSM-mEos2 cells were cultured in 7H9 media supplemented with 30 μg/ml KAN.

Development of PerSort. The PerSort method was developed using BD FACS Influx. A 70 micron tip and sheath fluid from BD bioscience was used in sorting and FACS analysis. Green fluorescent beads (3.5 micron) and 5 micron Accudrop beads from BD biosciences were used to calibrate the instrument for laser alignment, compensation, and cell sorting. Propidium Iodide (PI) stain (SigmaAldridge, 0.5%) was used to stain dead cells. Heat killed cells (incubated at 70^o − 80^o C for 5 mins) were used as dead cell control for PI stain. MSM cells with pCHERRY3 plasmid (MSM-mCherry) (26) was used to optimize single-cell mycobacterium sorting. MSM-mEos2 (KAN resistant) cultures were grown to OD600 of 0.6 and induced with 500 ng/ml ATc. Induced cultures were incubated at 37 ^{o} C for 12 h. MSM-mCherry (Hygromycin resistant) were mixed with induced MSM-mEos2 cultures in equal proportions and single mEos2 and mCherry cells were sorted and plated on 7H10 plates with Kan and Hyg to determine the efficiency of single-cell sorting (**Figure S2**). Gates for sorting dim and lit cells of ATc induced MSM-mEos2 strains (induced for 12 h) were determined using uninduced MSM-mEos2 strains stained with PI. FACS data was analyzed in FlowJo software (ver. 10).

Antimicrobial tolerance assay. Minimum inhibitory concentrations (MIC) of the MSM-mEOS2 strains were determined with disk diffusion assay (50). To determine percent survival, cultures (bulk and PerSorted) were incubated in 7H9 media containing desired MIC concentrations of INH and RIF. Drug treated bacteria were washed with or diluted (1:100) in 7H9 media and plated (100 μl) or spotted (5 μl) on 7H10 media with 30 μg/ml KAN at time points 0, 8, 12, 16, 20, and 24 h after incubation. Percentage survival was calculated with respect to 0 h.

qRT-PCR of PerSorted samples. Dim and lit cells were sorted into 500 μl TriZol and RNA was extracted with DirectZol RNA purification kit (Zymo Research) with the manufacturer's instructions. qRTPCR was performed with primers mentioned in **Table S3** with Luna® Universal Probe One-Step RT-qPCR Kit (New England Biolabs). Log2 normalized RNA abundances were calculated using ValidPrime signal as reference (51).

Single-cell persister gene expression and analysis. The Fluidigm Biomark system with 48x48 plates was used for this study. Single dim and lit cells were PerSorted into 96 well plates with VILO^{TM} reaction mix (5x), SUPERase (Invitrogen^{TM}), and 10% NP40 in a pre-noted random order to avoid sampling bias. Sorted plates were spun down and freeze-thawed 3 times on dry ice to rupture cells. ValidPrime assay (51) for non-transcribed genomic DNA was used to determine the rupture efficiency and calculate the signal from a single nucleic-acid strand (used as

reference to calculate transcript abundance). Reverse transcription (RT) was performed on
freeze thawed cells with VILO cDNA preparation mixture, T4-Gene32 protein, and random
hexamer primers. RNA spike-in (ECC2_SpikeIn RNA, 10 pM) was included in the RT master mix.
cDNA of the genes of interest (**Table S1**) was pre-amplified with TaqMan® PreAmp master mix
(Invitrogen[TM]) and equimolar mixture of forward and reverse strand primers designed for the
genes of interest (**Table S3**). Primers were removed with Exonuclease I (Invitrogen[TM]). Primers
sets used for preamplification were primed into the 48x48 Biomark assay plates. Quantitative
PCR assay with Biomark prescribed protocol was run on diluted, pre-amplified Exo 1 treated
cDNA with Sso Fast EvaGreen Supermix (Bio-Rad laboratories). Quality control for determining,
sorting, cell lysis, and cDNA preparation was performed by comparing the CT values of genomic
DNA control and spike-in control in all the cells (**Figure S5a,b**). Expression levels of genes was
measured as ∂CT in individual cells with reference to genomic DNA control (expected to result
from 1 copy of genomic DNA), less than or equal expression (of genomic DNA control) was
considered as zero expression and assays with CT >40 were flagged as missing values. ∂CT
values for each cells were corrected by adding or subtracting the deviation from median ∂CT of
spike-in control for a particular cell. Hierarchical clustering was performed with Python Seaborn
package and the R package, PVclust.
474
Feature selection of single-cell clusters. Ratio of toxin expression to their respective antitoxin
expression was calculated and included along with the other persister gene expression values.
Key features that differentiate dim and lit subpopulations were selected with tree-based
feature selection tool (30). Weights of the features that signify its ability to differentiate
between dim and lit sub populations were estimated as an average over 100 iterations of
randomly selected subsets of the dataset (70% of gene expression values). Constraints of gene
regulation was also used in selection of features as most persisters specific responses are a part
of alarmone response, a broad, genome wide change in transcript levels (52). The three
features with the highest weights and evidence of additional criteria were selected for
identification of clusters within dim and lit cells.
485
In situ ROS level measurements with CellRox assay. MSM-mEos2 cells were induced with ATc
(500 ng/ml) and incubated for 12 h under standard growth conditions. 10 µM N-acetyl cysteine
(NAC), an oxidative agent that reduces level of in situ reactive oxygen species (ROS) was added
to the induced MSM-mEos2 cultures and incubated at 37° C for 90 mins to prepare the negative
control. Along with negative control, another set of induced cultures were treated with a ROS
inducing 100µM MnTBAP (Sigma Aldrich, 55266-18-7) and incubated for 15 mins to prepare
positive control. CellRox orange (Thermo Fisher, C10443), was added to the induced samples
and control and incubated for 30 mins at room temperature. FACS analysis was performed to
measure the CellROX orange intensity. ROS + and negative gates were determined with control
samples and ROS levels in dim and lit cell population were determined with reference to the
controls.
497
Dim cell proportion and antimicrobial tolerance following L-cysteine addition. MSM-mEos2
cells were grown in PBS and 0.05% Tween-80, with and without 4mM L-cysteine, to an OD600
of 0.6. Cultures were induced with ATc (500 ng/ml) and incubated for 12 h. Dim cell proportions

in both conditions were measured with PerSort assay and normalized to the dim cell
proportions observed in optimal growth conditions (growth in 7H9 media without L-cysteine).
Tolerance of cultures grown in various conditions was measured with time kill curve assays,
under 5x MIC RIF (17.5 µg/ml) and 5x MIC INH (20 µg/ml) treatment. The % survival was
measured at varying time point by counting number of CFUs on 7H10 plates in reference to the
time point 0 of the respective samples.
507
508
509

Acknowledgements: We thank members of the Baliga lab for critical discussions; Amardeep Kaur for her technical expertise; Tim Petersen and Monica Orellana for help with FACS. Funding was provided by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health [R01AI128215] and [U19AI10676, U19AI135976]; and National Science Foundation [1518261, 1565166, and 1616955].

Author contributions: V.S. designed research, performed experiments, analyzed data and wrote the paper. M.L.A-O. performed computational analyses. E.J.R.P and N.S.B. designed research, analyzed data, and wrote the paper.

Competing interests: The authors declare no competing financial interests.

References

1.   Philippi T, Seger J. Hedging one's evolutionary bets, revisited. Trends Ecol Evol. 1989;4(2):41-4.

2.   CARE T. International Standards for Tuberculosis Care. TB CARE I. The Hague, The Netherlands2014.

3.   Sarathy JP, Via LE, Weiner D, Blanc L, Boshoff H, Eugenin EA, et al. Extreme Drug Tolerance of Mycobacterium tuberculosis in Caseum. Antimicrob Agents Chemother. 2018;62(2).

4.   Chaulk CP, Kazandjian VA. Directly observed therapy for treatment completion of pulmonary tuberculosis: Consensus Statement of the Public Health Tuberculosis Guidelines Panel. JAMA. 1998;279(12):943-8.

5.   Manina G, Dhar N, McKinney JD. Stress and host immunity amplify Mycobacterium tuberculosis phenotypic heterogeneity and induce nongrowing metabolically active forms. Cell Host Microbe. 2015;17(1):32-46.

6.   Jain P, Weinrick BC, Kalivoda EJ, Yang H, Munsamy V, Vilcheze C, et al. Dual-Reporter Mycobacteriophages (Phi2DRMs) Reveal Preexisting Mycobacterium tuberculosis Persistent Cells in Human Sputum. MBio. 2016;7(5).

7.   Liu Y, Tan S, Huang L, Abramovitch RB, Rohde KH, Zimmerman MD, et al. Immune activation of the host cell induces drug tolerance in Mycobacterium tuberculosis both in vitro and in vivo. J Exp Med. 2016;213(5):809-25.

8.   Cho J, Rogers J, Kearns M, Leslie M, Hartson SD, Wilson KS. Escherichia coli persister cells suppress translation by selectively disassembling and degrading their ribosomes. Mol Microbiol. 2015;95(2):352-64.

9.   Shah D, Zhang Z, Khodursky A, Kaldalu N, Kurg K, Lewis K. Persisters: a distinct physiological state of E. coli. BMC Microbiol. 2006;6:53.

10.   Torrey HL, Keren I, Via LE, Lee JS, Lewis K. High Persister Mutants in Mycobacterium tuberculosis. PLoS One. 2016;11(5):e0155127.

11.   Ahidjo BA, Kuhnert D, McKenzie JL, Machowski EE, Gordhan BG, Arcus V, et al. VapC toxins from Mycobacterium tuberculosis are ribonucleases that differentially inhibit growth and are neutralized by cognate VapB antitoxins. PLoS One. 2011;6(6):e21738.

12.   Schifano JM, Edifor R, Sharp JD, Ouyang M, Konkimalla A, Husson RN, et al.
Mycobacterial toxin MazF-mt6 inhibits translation through cleavage of 23S rRNA at the
ribosomal A site. Proc Natl Acad Sci U S A. 2013;110(21):8501-6.
13.   Stallings CL, Stephanou NC, Chu L, Hochschild A, Nickels BE, Glickman MS. CarD is an
essential regulator of rRNA transcription required for Mycobacterium tuberculosis persistence.
Cell. 2009;138(1):146-59.
14.   Betts JC, Lukey PT, Robb LC, McAdam RA, Duncan K. Evaluation of a nutrient starvation
model of Mycobacterium tuberculosis persistence by gene and protein expression profiling.
Mol Microbiol. 2002;43(3):717-31.
15.   Xie Z, Siddiqi N, Rubin EJ. Differential antibiotic susceptibilities of starved
Mycobacterium tuberculosis isolates. Antimicrob Agents Chemother. 2005;49(11):4778-80.
16.   McKinney JD, Honer zu Bentrup K, Munoz-Elias EJ, Miczak A, Chen B, Chan WT, et al.
Persistence of Mycobacterium tuberculosis in macrophages and mice requires the glyoxylate
shunt enzyme isocitrate lyase. Nature. 2000;406(6797):735-8.
17.   Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. Bacterial persistence as a phenotypic
switch. Science. 2004;305(5690):1622-5.
18.   Sebastian JS, S., Ajitkumar, P. Reduced Permeability to Rifampicin by Capsular
Thickening as a Mechanism of Antibiotic Persistence in Mycobacterium tuberculosis. BioRxiv.
2019;624569.
19.   McNeil MBC, S.; Awasthi, D.; Parish, T. Cell wall inhibitors increase the accumulation of
rifampicin in Mycobacterium tuberculosis. Access Microbiology. 2019;1.
20.   Levin-Reisman I, Fridman O, Balaban NQ. ScanLag: high-throughput quantification of
colony growth and lag time. J Vis Exp. 2014(89).
21.   Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ. Antibiotic tolerance
facilitates the evolution of resistance. Science. 2017;355(6327):826-30.
22.   Fridman O, Goldberg A, Ronin I, Shoresh N, Balaban NQ. Optimization of lag time
underlies antibiotic tolerance in evolved bacterial populations. Nature. 2014;513(7518):418-21.
23.   McKinney SA, Murphy CS, Hazelwood KL, Davidson MW, Looger LL. A bright and
photostable photoconvertible fluorescent protein. Nat Methods. 2009;6(2):131-3.
24.   Sawyer EB, Grabowska AD, Cortes T. Translational regulation in mycobacteria and its
implications for pathogenicity. Nucleic Acids Res. 2018;46(14):6950-61.
25.   Parikh A, Kumar D, Chawla Y, Kurthkoti K, Khan S, Varshney U, et al. Development of a
new generation of vectors for gene expression, gene replacement, and protein-protein
interaction studies in mycobacteria. Appl Environ Microbiol. 2013;79(5):1718-29.
26.   Carroll P, Schreuder LJ, Muwanguzi-Karugaba J, Wiles S, Robertson BD, Ripoll J, et al.
Sensitive detection of gene expression in mycobacteria under replicating and non-replicating
conditions using optimized far-red reporters. PLoS One. 2010;5(3):e9823.
27.   Oliver JD. The viable but nonculturable state in bacteria. J Microbiol. 2005;43 Spec
No:93-100.
28.   Keren I, Minami S, Rubin E, Lewis K. Characterization and transcriptome analysis of
Mycobacterium tuberculosis persisters. MBio. 2011;2(3):e00100-11.
29.   Orman MA, Henry TC, DeCoste CJ, Brynildsen MP. Analyzing Persister Physiology with
Fluorescence-Activated Cell Sorting. Methods Mol Biol. 2016;1333:83-100.

30. Baranauskas JN, O.; Nozawa, S.; Macedo, A. A tree-based algorithm for attribute
selection. Applied Intelligence. 2018;48(4):821-33.

31. Suzuki R, Shimodaira H. Pvclust: an R package for assessing the uncertainty in
hierarchical clustering. Bioinformatics. 2006;22(12):1540-2.

32. Langfelder P, Zhang B, Horvath S. Defining clusters from a hierarchical cluster tree: the
Dynamic Tree Cut package for R. Bioinformatics. 2008;24(5):719-20.

33. Weiss LA, Stallings CL. Essential roles for Mycobacterium tuberculosis Rel beyond the
production of (p)ppGpp. J Bacteriol. 2013;195(24):5629-38.

34. Ramage HR, Connolly LE, Cox JS. Comprehensive functional analysis of Mycobacterium
tuberculosis toxin-antitoxin systems: implications for pathogenesis, stress responses, and
evolution. PLoS Genet. 2009;5(12):e1000767.

35. Sala A, Bordes P, Genevaux P. Multiple toxin-antitoxin systems in Mycobacterium
tuberculosis. Toxins (Basel). 2014;6(3):1002-20.

36. Winther KS, Gerdes K. Regulation of enteric vapBC transcription: induction by VapC
toxin dimer-breaking. Nucleic Acids Res. 2012;40(10):4347-57.

37. Germain E, Roghanian M, Gerdes K, Maisonneuve E. Stochastic induction of persister
cells by HipA through (p)ppGpp-mediated activation of mRNA endonucleases. Proc Natl Acad
Sci U S A. 2015;112(16):5171-6.

38. Ishihara H, Nakazaki M, Kanegae Y, Inukai K, Asano T, Katagiri H, et al. Effect of
mitochondrial and/or cytosolic glycerol 3-phosphate dehydrogenase overexpression on
glucose-stimulated insulin secretion from MIN6 and HIT cells. Diabetes. 1996;45(9):1238-44.

39. Talior I, Yarkoni M, Bashan N, Eldar-Finkelman H. Increased glucose uptake promotes
oxidative stress and PKC-delta activation in adipocytes of obese, insulin-resistant mice. Am J
Physiol Endocrinol Metab. 2003;285(2):E295-302.

40. Bone BS, B.; Olszowy, M. Reactive oxygen probes - a broad range of colors with easier
labeling and compatibility with fixation: novel CellROX reagents from Molecular Probes (P3295).
J Immunol. 2013;190(1 Supplement).

41. Zampieri M, Horl M, Hotz F, Muller NF, Sauer U. Regulatory mechanisms underlying
coordination of amino acid and glucose catabolism in Escherichia coli. Nat Commun.
2019;10(1):3354.

42. Kwan BW, Valenta JA, Benedik MJ, Wood TK. Arrested protein synthesis increases
persister-like cell formation. Antimicrob Agents Chemother. 2013;57(3):1468-73.

43. Dorr T, Vulic M, Lewis K. Ciprofloxacin causes persister formation by inducing the TisB
toxin in Escherichia coli. PLoS Biol. 2010;8(2):e1000317.

44. Aizenman E, Engelberg-Kulka H, Glaser G. An Escherichia coli chromosomal "addiction
module" regulated by guanosine [corrected] 3',5'-bispyrophosphate: a model for programmed
bacterial cell death. Proc Natl Acad Sci U S A. 1996;93(12):6059-63.

45. Bhaskar A, De Piano C, Gelman E, McKinney JD, Dhar N. Elucidating the role of (p)ppGpp
in mycobacterial persistence against antibiotics. IUBMB Life. 2018;70(9):836-44.

46. Albrethsen J, Agner J, Piersma SR, Hojrup P, Pham TV, Weldingh K, et al. Proteomic
profiling of Mycobacterium tuberculosis identifies nutrient-starvation-responsive toxin-
antitoxin systems. Mol Cell Proteomics. 2013;12(5):1180-91.

47.   McKenzie JL, Robson J, Berney M, Smith TC, Ruthe A, Gardner PP, et al. A VapBC toxin-
antitoxin module is a posttranscriptional regulator of metabolic flux in mycobacteria. J
Bacteriol. 2012;194(9):2189-204.
48.   Verstraeten N, Knapen WJ, Fauvart M, Michiels J. Membrane depolarization-triggered
responsive diversification leads to antibiotic tolerance. Microb Cell. 2015;2(8):299-301.
49.   Vilcheze C, Hartman T, Weinrick B, Jain P, Weisbrod TR, Leung LW, et al. Enhanced
respiration prevents drug tolerance and drug resistance in Mycobacterium tuberculosis. Proc
Natl Acad Sci U S A. 2017;114(17):4495-500.
50.   Bonev B, Hooper J, Parisot J. Principles of assessing bacterial susceptibility to antibiotics
using the agar diffusion method. J Antimicrob Chemother. 2008;61(6):1295-301.
51.   Laurell H, Iacovoni JS, Abot A, Svec D, Maoret JJ, Arnal JF, et al. Correction of RT-qPCR
data for genomic DNA-derived signals with ValidPrime. Nucleic Acids Res. 2012;40(7):e51.
52.   Dahl JL, Kraus CN, Boshoff HI, Doan B, Foley K, Avarbock D, et al. The role of RelMtb-
mediated adaptation to stationary phase in long-term persistence of Mycobacterium
tuberculosis in mice. Proc Natl Acad Sci U S A. 2003;100(17):10026-31.
652
653
654

 Figure legend

Figure 1: Time-kill curves and growth characteristics of *M. smegmatis* from nutrient rich and
nutrient starved conditions. **a**,**b**. Time-kill curves of MSM cultures grown in nutrient rich (NR) or
nutrient starved (NS) conditions treated with 5x MIC INH (**a**) or 5x MIC RIF (**b**). The solid lines
indicate experimentally observed time-kill curves. Data points are from three experimental
replicates, error bars were calculated by measuring standard deviation between replicates. The
dashed and dotted lines distinguish the slopes of the susceptible and tolerant subpopulations,
respectively. **c**. ScanLag analysis showing time of appearance (TOA) of cultures grown in
nutrient rich and nutrient starved conditions. Error bars within the violin plot are standard
deviation with confidence interval of 0.9. The dashed lines indicate the mean TOA from cultures
grown in nutrient rich or nutrient starved conditions. Mean TOA of nutrient starved and
nutrient rich conditions were compared with Students-t test.

Figure 2: PerSort isolation and characterization of subpopulations from ATc-induced MSM-
mEos2 cultures. **a**. Top: Population structure of ATc-induced (500ng/ml) MSM-mEos2 cells from
single-cell gates, represented by mEos2 fluorescence on x-axis and propidium iodide (PI, 5µl/ml)
fluorescence on y-axis. Polygons indicate the gates for non-culturable cells (PI(+)), dim cells, and
lit cells with the proportion of cells in that particular gate. Bottom: Cumulative distribution of
mEos2 florescence intensity in MSM-mEos2 cells, dashed lines indicate boundaries of sort gates
for dim, lit, and unselected PI(-) cells. **b**. Cultivability of PerSorted dim, lit and PI(-) cells from
ATc-induced MSM-mEos2 cultures. The "% cultivable" was calculated from the number of CFUs
from 200 sorted cells. Experiments were performed in triplicates and error bars represent the
standard deviation between replicates. Significance of cultivability difference between dim, lit
and non-culturable subpopulations was calculated with Students t-test. **c**. ScanLag analysis of
dim, unselected PI(-) and lit cells from PerSorted MSM-mEos2 cultures induced with ATc (500
ng/mL), the dashed lines in the violin plot indicate mean TOA of the sorted subpopulations.
Error bars within the violin plot are standard deviation with confidence interval of 0.9. **d,e**. %
survival of 5x MIC INH or 5x MIC RIF treatment of PerSorted dim and lit cells of MSM-mEos2
cultures induced with ATc (500 ng/ml), compared to % survival of whole populations (i.e.,
unsorted) of MSM-mEos2 induced with ATc (500 ng/ml) and grown in nutrient rich and starved
conditions. Experiments were performed in triplicates and error bars represent the standard
deviation between replicates. Significance of survival between dim and lit subpopulations was
calculated with Students-t test.

Figure 3: Regrowth dynamics of PerSorted dim and lit cells and population structure under
naïve versus nutrient starved conditions. **a.** Dim and lit cell proportions in cultures regrown
from PerSorted dim and lit populations obtained from ATc-induced MSM-mEos2 cultures. The
proportions of cells in each subpopulation were measured from the gating methods described
in Figure 2a. **b,c**. Lag phase and maximum growth rate were calculated from OD600 absorbance
measurements of dim and lit cells PerSorted into 7H9 media. Error bars were calculated by
measuring standard deviation in the lag phase and maximum growth rates between dim and lit

cells (n = 100,000). Significance of differences between dim and lit subpopulations were
calculated with Students-t test. NS, not significant. **d**. Population structure of ATc-induced
MSM-mEos2 cultures grown in naïve (green) and nutrient starved (grey) conditions. **e.** A model
describing the formation of dim cells under naïve and nutrient starved conditions in
mycobacterium. The probability of dim cell formation was calculated from their relative
proportion in the indicated culture conditions.
703
Figure 4: Single-cell gene expression and ROS levels from PerSorted single dim and lit cells. **a.**
Hierarchical clustering of top ranked features in PerSorted dim and lit single-cells. Blocks in the
cluster represent the statistically significant clusters (p-value = 0.01). **b**. qRT-PCR measurement
of 16s rRNA, 23s rRNA and Phospho-glucoisomerase (PGI, MSMEG_5541) levels in PerSorted
dim and lit cells (n = 300 cells). Error bars were calculated by measuring standard deviation of
gene expression. P-values were calculated with Student's t-test between the dim and lit cells. **c.**
Reactive oxygen Species (ROS) levels, measured with CellRox, of ATc-induced MSM-mEos2
cultures grown in nutrient rich conditions and PerSorted into dim and lit subpopulations. ROS
was quenched with N-acetyl cysteine and induced with MnTBAP and used as reference
controls. Significance of ROS levels between dim and lib subpopulations was calculated by K-S
test (K-S statistics of CellRox orange intensity between dim and lit population; K-S max
Difference: 56.1%, K-S max at Intensity: 14.8551, K-S probability: >99.9%).
716
Figure 5: Dim cell proportion and time-kill curves upon L-cysteine addition to nutrient starved
M. smegmatis. **a.** Proportion of dim cells in MSM-mEos2 cultures grown in nutrient starved
conditions with or without the addition of 4mM L-cysteine. Dim cell proportions in each
condition were calculated with respect to MSM-mEos2 cultures grown in naïve conditions. Error
bars were calculated by measuring standard deviation in dim cell proportions between 10
PerSort replicates. **b,c** Time-kill curves of MSM cultures grown in nutrient starved conditions
with 4mM L-cysteine and treated with 5x MIC INH (**b**) or 5x MIC RIF (**c**). Time-kill curve data for
nutrient rich and nutrient starved conditions (no L-cysteine) is from Figure 1. Data points are
averaged from three independent experiments and error bars represent the standard deviation
in percent survival between the replicates.
727
Figure 6. The persister subpopulation converges on a low O_2 metabolic state, which was
exploited by addition of L-cysteine to convert the subpopulation to drug susceptible cells. The
diagram depicts the processes driving dim (multidrug tolerant persisters) and lit (drug
susceptible) cell formation, as discovered through single-cell expression analysis. Evidence from
literature further connects these various mechanisms to the O_2 respiratory state of either dim
or lit cells. Red arrows indicate connections identified in this study, black arrows are previously
determined connections from literature (references indicated), and white boxes are proposed
mechanisms for persister cell formation.
736

737
738

a: Aizenman et al ; b: Verstraeten et al.; c: McKenzie et al.; d: Vilchèze et al.;

→ Active (this study); → Active (literature); ✗ Inactive (this study)

a INH treated

% Survival

- Nutrient rich (NR)
- Nutrient starved (NS)
- - - Susceptible sub-population
- ···· Tolerant sub-population

Time (hours)

b RIF treated

↔ Delayed killing

Time (hours)

c Time of appearance of CFUs

24 hrs + mins

~4 hrs
(p-value: <10⁻⁵)

NS
(128 CFUs)

NR
(110 CFUs)