Long-distance dispersal, ice sheet dynamics, and mountaintop isolation underlie the genetic

structure of glacier ice worms

Scott Hotaling[1], Daniel H. Shain[2], Shirley A. Lang[3], Robin K. Bagley[4], Lusha M. Tronstad[5],

David W. Weisrock[6], and Joanna L. Kelley[1]

Affiliations:

[1] School of Biological Sciences, Washington State University, Pullman, WA, USA

[2] Department of Biology, Rutgers University, Camden, NJ, USA

[3] Graduate School of Biomedical Sciences, Rowan University, Stratford, NJ, USA

[4] Department of Biology, University of Iowa, Iowa City, IA

[5] Wyoming Natural Diversity Database, University of Wyoming, Laramie, WY, USA

[6] Department of Biology, University of Kentucky, Lexington, KY, USA

Correspondence: Scott Hotaling, School of Biological Sciences, Washington State University,

Pullman, WA, 99164, USA; Email: scott.hotaling@wsu.edu; Phone: (828) 507-9950; ORCID:

0000-0002-5965-0986

Running title: Population genomics of ice worms

Keywords: RAD sequencing, Pacific Northwest, phylogeography, *Mesenchytraeus solifugus*,

global change biology, annelid

Abstract:

Disentangling the contemporary and historical factors underlying the spatial distributions of

species is a central goal of biogeography. For species with broad distributions but little capacity

to actively disperse, disconnected geographic distributions highlight the potential influence of

passive, long-distance dispersal (LDD) on their evolutionary histories. However, dispersal alone

cannot completely account for the biogeography of any species, and other factors–e.g., habitat

suitability, life history–must also be considered. North American ice worms (*Mesenchytraeus*

solifugus) are glacier-obligate annelids that inhabit coastal North American glaciers from Oregon

to Alaska. Previous studies identified a complex biogeographic history for ice worms, with evidence for genetic isolation, unexpectedly close relationships among geographically disjunct lineages, and evidence for contemporary migration across large (>1,500 km) areas of unsuitable habitat. In this study, we collected genome-scale sequence data for most of the known ice worm range. We found support for a deep divergence between populations along the Pacific Coast and the inland flanks of the Coast Mountains (mean F_{ST} = 0.60) as well as support for LDD from Alaska to Vancouver Island, perhaps mediated by migrating birds. Our results highlight the power of genomic data for disentangling complex biogeographic patterns, including the presence of LDD.

Introduction:

For more than a century, long-distance dispersal (LDD) among presumably isolated populations has intrigued biologists [1-4]. Historically considered rare and unpredictable, the idea that LDD can act as a general mechanism influencing the biogeography of presumably dispersal-limited, macroscopic organisms has gained traction in recent years, with examples accumulating for both plants [5] and invertebrates [6-9]. Many animal vectors play an integral role in plant and invertebrate LDD [e.g., 10, 11], however, in most cases, the resulting LDD is limited to < 10 km. For more extreme LDD events (e.g., > 100 km), the most common animal vector is likely migratory birds, as they seasonally move by the millions over broad spatial scales and geographic barriers, visiting similar habitats along the way [12, 13]. Through this mechanism, dispersal units (e.g., whole organisms, eggs, seeds, etc.) may be ingested and dispersed internally or by directly adhering to the bird's exterior [12]. Thus, as long as there is an opportunity for migratory birds and dispersal units to interact, the opportunity also exists for LDD. Physically quantifying LDD is difficult, however, because it requires real-time sampling and searching (internal and external) of migrating birds for hitchhiking dispersers. Moreover, because rare migratory events can affect species distributions [14] and influence genetic differentiation among populations [15], even thorough physical surveys of migratory birds that find no evidence for LDD cannot rule out its presence. Therefore, alternative approaches for detecting and characterizing LDD should be employed. Because population genomic tools are well-suited to detecting gene flow among populations [e.g., 16], these tools are also well-suited to the indirect detection of LDD, even in the absence of field observations.

Many mechanisms influence genetic relationships among taxa and a range of factors

should be considered when attempting to reconstruct biogeographic patterns. For instance, pulses

and contractions of glaciers and ice sheets have shaped the evolutionary histories of populations

and species throughout Earth's history [17-20]. These ice sheet dynamics typically affect

organisms as ice traverses the landscape, separating and reconnecting populations as it expands

and contracts. However, some species are directly tied to ice sheets [e.g., the meltwater stonefly,

17] and are therefore much more susceptible to ice sheet influence on their evolutionary

trajectories. Perhaps no species is more directly tied to ice sheets than the glacier ice worms,

Mesenchytraeus solifugus in North America [21] and *Sinenchytraeus glacialis* in Tibet [22]. The

geographic range of *M. solifugus* (hereafter "ice worm") follows a coastal arc from the Chugach

Mountains in southeast Alaska to the Cascade Volcanoes of Washington and Oregon [24]. Ice

worms can also not tolerate temperatures more than roughly ±7 °C from freezing and requires

glacier ice for survival and reproduction [23]. With such unique ecology and physiology, and a

dispersal-limited life history, the evolutionary history of ice worms since diverging from

conspecifics [25, 26] should be relatively simple, with gene flow occurring during glacial periods

and isolation (paired with genetic drift) driving divergence among mountaintop-isolated

populations during interglacial periods. Natural systems, however, are often more complex than

expected and indeed, the evolutionary history of ice worms challenges general expectations of

gene flow and evolutionary dynamics in mountain systems.

Previous genetic studies based on one or two genetic markers identified three ice worm

lineages: a "northern" clade in southern Alaska, a "central" clade in southeast Alaska and

northern British Columbia (BC), and a "southern" clade ranging over much of BC to southern

Oregon [23-25]. Surprisingly, phylogenetic evidence supported the northern and southern

lineages as being most closely related to one another despite the central clade separating them

geographically. The most curious aspect of ice worm biogeography, however, has been the

repeated discovery of closely related ice worms on glaciers several hundred to thousands of

kilometers south of their closest genetic relatives [23, 25; P. Wimberger, unpublished data].

These disjunct northern ice worms co-occurred with, but appeared genetically distinct from, their

conspecifics (either central or southern clade ice worms) on the same glaciers. Dial *et al.* [23]

laid out three possible explanations for this pattern: wind transport, passerine-mediated dispersal,

or a more extensive previous range of the northern clade. While wind transport seems unlikely,

the potential for passerine-mediated dispersal is reasonable, particularly in light of other

examples of bird-mediated LDD [see 7, 27]. The third scenario, a more extensive distribution of

the northern clade with holdover lineages inhabiting the same glacier as more recent colonizers

could indeed result in more than one distinct lineage on the same glacier. However, if this recent

colonization occurred at the Last Glacial Maximum when ice sheet extent last peaked in the

region, recently colonizing lineages should not remain closely related to the source without at

least some degree of contemporary gene flow.

In this study, we leveraged a modern population genomic toolkit to add new perspective

to the age-old challenge of identifying LDD in wild populations. We also provide new insight

into how multiple factors can interact to shape the evolutionary history of species. We

hypothesized that the biogeographic history of ice worms stemmed from a confluence of three

main factors: extreme LDD, glacier dynamics, and mountaintop isolation. To test this hypothesis,

we generated a genome-wide single nucleotide polymorphism (SNP) data set to answer three

more-specific questions: (1) How do the clades previously diagnosed from a small number of

markers hold up to genome-wide scrutiny? (2) What, if any, genomic evidence exists for LDD in

ice worms? (3) How do the evolutionary relationships among ice worm populations and genetic

clusters align with glacial history in the region [e.g., 28]? Beyond a refined view of ice worm

evolution, our study confirms that LDD does occur in ice worms, providing an example of LDD

in an annelid and a rare population genomic exploration of the process. Moreover, while

considerable evidence details the existence of refugia in the Pacific Northwest (PNW) during the

Pleistocene [29], few studies have explored how ice sheet dynamics influenced the evolutionary

history of species directly tied to them [e.g., 17]. Our results reveal the profound impact that a

putative ice ridge that formed along the crest of the Coast Mountains [28] may have had on ice

worm evolution, possibly precipitating an ongoing speciation event. Broadly, our findings

highlight the power for population genomics to capture contemporary evidence of LDD while

also providing novel evidence for reconstructing the glacial history of a region.

Table 1. Sampling information and summary statistics for all ice worm populations included in this study. Abbreviations: n = sample size, π = nucleotide diversity, Het = heterozygosity, F_{IS} = inbreeding coefficient, AK = Alaska, BC = British Columbia, VI = Vancouver Island. π, Het, and F_{IS} were calculated for variable sites only.

Population	Latitude, longitude	State/Prov.	Elev. (m)	n	π	Het	F_{IS}
Learnard (LEA)	60.806, -148.721	AK	624	3	0.114	0.100	0.026
Davidson (DAV)	59.067, -135.551	AK	986	15	0.078	0.065	0.037
Treaty (TRE)	56.586, -130.151	BC	1376	8	0.026	0.033	-0.012
Bear (BEA)	56.096, -129.681	BC	648	6	0.039	0.046	-0.013
William Brown (WIB)	54.611, -129.129	BC	1260	6	0.056	0.057	0.000
Jacobson (JAC)	52.050, -126.072	BC	1249	5	0.108	0.106	0.005
White Mantle (WHM)	50.795, -125.153	BC	1764	8	0.131	0.135	-0.005
Comox (COM)	49.545, -125.355	BC (VI)	1881	4	0.091	0.072	0.034
Mariner (MAR)	49.460, -125.764	BC (VI)	1754	4	0.098	0.084	0.024

Methods:

Sample collection, library preparation, and SNP calling

During the summer of 2009, ice worms were collected from nine glaciers across most of their range (Figs. 1, S1; Table 1). Samples were stored in > 80% EtOH until DNA was extracted from 59 ice worms using a Qiagen DNEasy Blood and Tissue Kit. Double-digest restriction-site associated DNA sequencing libraries were prepared following Peterson *et al.* [30] with restriction enzymes EcoRI and NlaIII. During library preparation, samples were divided into two groups and each sample was assigned a unique, variable-length barcode [31] which was incorporated during adapter ligation. Size selection for a 350 bp ± 35 bp window was performed with a Pippen Prep (Sage Science), and both sample groups were subsequently amplified using PCR primers containing a group-specific barcode. The 59-sample library was sequenced on one lane of an Illumina HiSeq4000 at the University of Illinois High-Throughput Sequencing and Genotyping Unit with single-end, 100 bp chemistry.

Raw reads were demultiplexed, quality-filtered, and RAD loci were assembled *de novo* using the *process_radtags* and *denovo_map* functions of the Stacks v1.46 pipeline [32]. We allowed a maximum distance between stacks of 2 and a minimum read depth of 10. Next, we applied a stringent filtering scheme to identify high-confidence SNPs that were shared among many individuals. We only included SNPs if they were present in ≥ 5 populations, genotyped in ≥ 50% of individuals per population, and were in Hardy-Weinberg equilibrium with a minor allele frequency of ≥ 0.025 overall. We further restricted analyses to one random SNP per locus for all analyses except fineRADstructure (see below). All post-Stacks filtering steps were

146   performed in PLINK v1.07 [33] and the commands used in this study are provided on GitHub

147   (https://github.com/scotthotaling/ice_worm_ddRAD).

Figure 1. Ice worm populations sampled for this study. Color-coding reflects the results of a
fineRADstructure coancestry analysis. Group I (circles) and II (square) populations were generally
defined by their presence to the east or west of a putative ice ridge that formed during the Pleistocene
~25,000 years ago [28] as well as their distance to the Pacific Ocean. The deep divergence between
groups I and II is clearly evident along with more recent differentiation within each group. One individual
from the Mariner Glacier (asterisk) is admixed between the Mariner/Comox (Vancouver Island) and
Learnard (southern Alaska) clusters, indicating recent LDD.

Population genetic and phylogenetic analyses

For each population, we calculated nucleotide diversity (π), heterozygosity (Het), and the

inbreeding coefficient (F_{IS}). We also calculated a pair-wise AMOVA F_{ST} for all population

combinations [34] – using the *populations* module in Stacks. To test for a signature of isolation-

by-distance [IBD, 35], we estimated Euclidean distances among sites with Google Earth and

tested the correlation between geographic distance and F_{ST} with four Mantel tests performed in

GenoDive v2.0b27 [36]. The first Mantel test included all nine populations and the second

excluded both Vancouver Island (VI) populations from the Mariner and Comox Glaciers to

assess whether unique histories for those populations were significantly altering results. The

third and fourth Mantel tests focused on signatures of IBD within groups "I" and "II" (see

Results). We also measured Euclidean distances for each population to the Pacific Ocean in

Google Earth Pro as an additional spatial comparison of groups I and II. We determined if mean

distances to the Pacific Ocean differed between the two groups with a one-way ANOVA.

Population structure was inferred in two ways: a maximum likelihood-based method

using ADMIXTURE 1.3.0 [37] and a discriminant analysis of principal components (DAPC)

with the R package *adegenet* [38]. ADMIXTURE analyses were performed with default settings,

a range of clusters (K) from 1-12, and 25 replicates per K with the current time as the random

seed. The cross-validation (CV) error for each K was plotted to identify the best-fit K (minimized

CV across the mean of all replicates for each K). After identifying the best-fit K, we considered

the replicate that minimized CV across all 25 replicates for all K's to be the best-fit solution

overall. However, because all runs did not converge on the same result, we also inspected best-fit

solutions for other replicates of $K = 7$ (the best-fit K overall) to clarify the distribution of best-fit

solutions. For DAPC, we first used the *find.clusters* function to identify the optimal K [i.e., the K

with the lowest Bayesian Information Criterion (BIC)]. Next, to avoid over-fitting of the model,

we retained the appropriate number of principal components (PCs) according to the α-score [PCs

retained = 6, Fig. S2; 38]. We performed a final DAPC analysis using the best-fit K and optimal

number of PCs identified in the previous two steps.

We extended our population structure analyses to infer both shared ancestry and

phylogenetic relationships in two ways: a nearest neighbor haplotype approach to infer

coancestry with fineRADstructure [39] and phylogenetic relationships inferred from singular

value decomposition estimates for quartets of tips using SVDQuartets [40] as implemented in

PAUP* v4.0a159 [41]. For fineRADstructure, we used 100,000 burn-in iterations followed by

100,000 iterations sampled every 1,000 steps for the Markov chain Monte Carlo clustering

algorithm. Next, we used 10,000 iterations of the tree-building algorithm to assess genetic

relationships among clusters. Since fineRADstructure is a haplotype-based approach, analyses

were performed using all SNPs for a given RAD locus (i.e., a haplotype) rather than randomly

selected single SNPs per locus. For SVDQuartets, we performed exhaustive sampling of all

possible quartets (every combination of four tips) and branch support was estimated with 100

nonparametric bootstrap replicates.

Table 2. Above the diagonal: Pair-wise AMOVA F_{ST} values for all localities populations included in this
study. Mean F_{ST} (bottom row) refers to the average pair-wise differentiation for columnar populations
versus all others. Below the diagonal (in gray): mean pair-wise shared loci for the fineRADstructure
coancestry analysis (see Fig. 1).

	LEA	DAV	TRE	BEA	WIB	JAC	WHM	COM	MAR
LEA	--	0.333	0.634	0.626	0.608	0.586	0.557	0.347	0.344
DAV	74.9	--	0.567	0.557	0.541	0.527	0.515	0.290	0.295
TRE	19.8	20.0	--	0.153	0.207	0.275	0.250	0.656	0.652
BEA	19.0	19.9	127.2	--	0.165	0.250	0.222	0.650	0.645
WIB	20.1	20.5	119.8	123.0	--	0.216	0.201	0.630	0.628
JAC	21.0	20.4	99.7	99.5	104.7	--	0.172	0.604	0.600
WHM	21.6	19.1	90.7	90.9	93.0	107.8	--	0.576	0.572
COM	84.8	86.1	21.8	21.3	23.0	22.9	22.1	--	0.160
MAR	164.5	82.9	21.2	21.3	22.3	21.8	21.4	262.3	--
Mean F_{ST}	0.504	0.453	0.424	0.408	0.399	0.404	0.383	0.489	0.487

Results:

We generated 343,875,880 reads with an average of 5,828,404 sequences per individual

(min. = 446,872 and max. = 40,982,490). Our total RAD data set included 360,534 unique loci.

After filtering, we retained 6,019 loci and 10,392 SNPs (mean = 1.73 SNPs per locus). This final

data set had genotype calls for ~65% of all SNPs. Nucleotide diversity (π) was highest in the

White Mantle and Learnard populations (0.131 and 0.114, respectively) and lowest in the Treaty,

Bear, and William Brown populations (0.026-0.056; Table 1). Heterozygosity followed the same

pattern as π (Table 1). The inbreeding coefficient (F_{IS}) was highest in the Davidson and Comox

populations (0.037 and 0.034, respectively) and lowest in Bear (-0.013) and Treaty (-0.012;

Table 1). Mean differentiation (F_{ST}) for all pairwise comparisons was 0.439. The Learnard

population from southern Alaska was, on average, the most differentiated from all others (mean

$F_{ST} = 0.504$) and White Mantle was the least differentiated (mean $F_{ST} = 0.383$; Table 2). We

detected no association between genetic and geographic distances in either study area-wide

Mantel tests (Mantel's *r*, all populations = -0.04, *P* = 0.42; Mantel's *r*, no VI populations = 0.15,

$P = 0.36$). There was, however, a signature of IBD within group II (Mantel's r, group I = 0.839,

$P = 0.025$) but not within group I (Mantel's r, group II = 0.839, $P = 0.082$).

Our DAPC analyses supported $K = 6$ as the optimal number of genetic clusters (Figs. 2A,C). For

ADMIXTURE, our results supported $K = 7$ as the best-fit to the data (Figs. 2B,C) and the

SVDQuartets phylogeny largely mirrored both lines of population structure evidence (Fig. 2D).

All analyses supported multiple independent genetic clusters of ice worms, many more than three

clades identified in previous studies [e.g., 25]. Our DAPC and ADMIXTURE results differed in

two ways: (1) DAPC analyses grouped the Treaty, Bear, and William Brown populations into

one cluster (as did SVDQuartets) whereas ADMIXTURE split William Brown into its own

cluster, and this accounted for the $K = 6$ versus $K = 7$ discrepancy between the two approaches.

(2) ADMIXTURE also split Learnard (southern Alaska) and the two Vancouver Island

populations – Comox and Mariner – into two clusters with one sample, MS5 (red asterisks in

Fig. 2), almost evenly split in terms of ancestry between the two. Conversely, DAPC split

Learnard and Comox+Mariner into two clusters with MS5 completely assigned to the Learnard

cluster, despite MS5 being sampled from the Mariner glacier far to the south (Figs. 1,2). Aside

from the best-fit solutions, we also observed a second common ADMIXTURE solution for $K = 7$

which differed from the best-fit solution in one way; rather than splitting Treaty+Bear and

William Brown into two clusters (Fig. 2B), Bear specimens were almost evenly admixed

between Treaty and William Brown (Fig. S3), its two closest neighbors to the north and south,

respectively. Finally, because our SNP filtering focused on overarching patterns in the data set,

and likely overlooked some degree of population-specific detail, our results are likely

conservative estimates of genetic structure in the group.

Figure 2. Population genetic structure of ice worms based upon (A) a discriminant analysis of principal components (DAPC) for *K* = 6 and (B) ADMIXTURE results for *K* = 7. (C) Comparisons of support for different values of *K* for DAPC (Bayesian information criterion, BIC; gray dashed line, left y-axis) and Admixture (cross-validation error, CV; dark line, right y-axis). The best-fit *K* (white square, ADMIXTURE; white circle, DAPC) corresponds to the *K* at which CV (ADMIXTURE) or BIC (DAPC) was minimized, respectively. For ADMIXTURE, vertical white bars represent the standard deviation of values for each *K* across 25 replicates. (D) An unrooted phylogeny of ice worms generated with SVDQuartets. Tip colorations reflect assignments in (A). Dark circles indicate nodes with > 95% bootstrap support. Specimens in group I (circles) and II (squares) are denoted with different symbols. Asterisks highlight a single specimen, MS5, which showed evidence of shared ancestry across geographically disjunct populations, indicating LDD.

Our fineRADstructure results mirrored those from DAPC, identifying the same six genetic clusters with MS5 assigned to the Learnard cluster rather than the Comox/Mariner cluster. Still, MS5 again exhibited substantial evidence for shared ancestry between the Learnard and Comox/Mariner genetic clusters (Fig. 1). On average, MS5 shared 411 loci with Learnard ice worms and 220 loci with ice worms from the Comox and Mariner ice worms (Fig. 1), a roughly 60/40 split between the two clusters. Notably, MS5 also exhibited the highest heterozygosity of any individual in the study (and this result did not stem from outsized

coverage, Fig. S4). Our fineRADstructure analysis also revealed the potential for an overarching,

deep divergence between two groups of ice worm populations (groups I and II; Fig. 1). This split

was corroborated by both SVDQuartets (Fig. 2D) and F_{ST} comparisons. Mean pairwise F_{ST}

among populations within groups I and II were 0.21 and 0.30, respectively. Across groups,

however, mean pairwise F_{ST} was 0.60. Mean distance to the Pacific Ocean also differed between

groups by 100.5 km (group I = 73.3 km, group II = 173.8 km; ANOVA, $P < 0.001$).

263   To be clear, the geographic groupings (I and II) described in this study are with respect to

the Coast Mountains, and specifically a putative ice ridge that formed along the crest of the range

during the Pleistocene [28]. All populations, regardless of grouping, followed the known pattern

of ice worms inhabiting relatively coastal glaciers (see Fig. S1). Interestingly, while the

geographic location of 8 of 9 populations fell on either side of the putative ice ridge, one

population did not: White Mantle. Our resolution of a deep, two-group split within ice worms

adds new clarification to the three-clade result identified by earlier studies [e.g., 23, 25], showing

that the previously recognized northern and central clades should likely be lumped together (now

group I).

Discussion:

274   Historical and contemporary factors, both biotic and abiotic, interact to shape the present-

day genetic structure of populations and species. Disentangling their varied contributions can be

difficult, however, particularly when evolutionary histories are muddled by unexpected events

(e.g., LDD of an organism with limited potential for active dispersal). The modern population

genomic toolkit provides historically unprecedented power to resolve biogeographical

complexity by allowing more quantitative perspectives of relatedness and greatly improved

resolution of genetic independence or similarity [42]. In this study, we used a population

genomic data set to refine understanding of the evolution history of the extremophile, glacier-

obligate ice worm, *M. solifugus*. Our results provide a clear genomic perspective of LDD,

showing unequivocally that migration has occurred between southern Alaska and the glaciers of

Vancouver Island ~1,900 km to the south and across the Pacific Ocean. We also provide an

independent line of biological evidence in support of the geological hypothesis that an ice ridge

formed along the crest of the Coast Mountains during the Pleistocene [28].

The recent biogeographic history of ice worms appears to have been shaped by three

main factors: i) ice sheet dynamics (likely during the Pleistocene), ii) mountaintop isolation from

conspecifics following the retreat of Pleistocene ice into higher elevations, and iii) LDD.

i.   Evidence for the first, overarching factor that has defined the recent evolution of ice

worms lies in our overwhelming support for divergence between two overarching

groups of ice worms (I and II). During the Pleistocene, ice was generated in the high

peaks of the Coast Mountains and subsequently flowed west to the Pacific and east to

the BC interior [28, 43, 44]. This potential western-eastern divergence in ice worms

aligns with this divide, suggesting that each group diverged in allopatry from their

conspecifics. At its maximum, the Cordilleran ice sheet was a ~2,000-3,000 m high

convex dish with gentle interior slopes that steepened at its periphery [28, 45]. Given

the sensitivity of ice worms to extreme cold [23], populations likely only persisted on

the margins of the ice sheet, as their present-day occurrences on the lower flanks of

higher elevation, low-latitude glaciers suggest. It is possible that, as suggested

previously [23, 25], the Boundary Ranges (the most northern subrange of the Coast

Mountains), are actually the biogeographic barrier that drove the observed deep

divergence in ice worms described in this study. However, without more fine-scale

population genomic sampling on both sides of the Coast Mountains (including the

Boundary Ranges), this nuance of ice worm biogeography will remain unclear. In the

same vein, the strong genetic similarity of White Mantle to populations east of the

proposed ice ridge (Fig. 1), despite falling on its western side indicates that either the

ice ridge actually formed more to the west than previously thought [28], the White

Mantle population has migrated west since the end of the Pleistocene, or the ice ridge

itself was not a barrier driving differentiation (as discussed above).

ii.   The Cordilleran ice sheet was seeded by alpine glaciers and throughout western North

America, the ice sheet's signature persists in mountainous regions. While the specific

dynamics of deglaciation on valley and drainage scales are unknown, a safe

assumption is that glaciers retreated from valleys into higher elevations, likely with

ice worm populations in tow. Increasing mountaintop isolation and subsequent

genetic drift likely precipitated more recent differentiation within groups I and II

318		since their initial split. Evidence for IBD within group II supports this hypothesis.
319		The lack of support for IBD in group I may simply reflect a smaller sample size or
320		LDD maintaining connections over larger spatial scales than a purely IBD model
321		would predict.
322	iii.	Despite limited sampling, we were able to identify one instance of recent LDD among
323		geographically disparate ice worm populations. Indeed, with a roughly 60/40 ancestry
324		split between the Mariner/Comox and Learnard clusters, one specimen (MS5), is
325		likely the progeny of recent hybridization between the two. This indicates that LDD is
326		both ongoing in ice worms and perhaps not particularly rare. The most plausible
327		mechanism for ice worm LDD is passive dispersal of mucous-coated ice worm
328		cocoons on southward migrating birds [e.g., 12]. Several passerines (e.g., Gray-
329		crowned rosy finches, *Leucosticte tephrocotis*) have been observed feeding on ice
330		worms [46, 47; S.H., personal observation] and the presence of an ice worm clitellum
331		[48] indicates that ice worms, like other *Mesenchytraeus* species, reproduce by egg-
332		laden cocoons [23]. The seemingly exclusive north-to-south pattern of ice worm LDD
333		also has temporal support from bird migratory behavior. Indeed, late autumn ice
334		worm reproduction [at the end of the productive season on mountain glaciers, 49]
335		likely occurs in concert with southward-migrating birds stopping to feed on glaciers
336		free of seasonal snow. In contrast, returning spring migrants pass over the same
337		glaciers when seasonal snowfall still covers overwintering ice worms [23], limiting
338		the potential for LDD in the reverse direction.
339		

One question remains, however, if birds are precipitating LDD in ice worms, why has it

only been observed for populations west of the Coast Mountains? This curiosity ties in to an

important question in North American biogeography: to what extent have ice sheets driven

present-day patterns of speciation and genetic differentiation among fauna of the northwest? For

ice worms, we hypothesize that ice worm populations comprising groups I and II have

accumulated some degree of reproductive isolation and are either already species-level lineages

or on their way to this conclusion. Thus, we predict a zygotic barrier may be limiting inter-group

migrants from leaving a genomic signature of their presence. It is also possible, and perhaps

likely, that LDD in ice worms is merely the result of vector migration patterns. For instance, *L.*

tephrocotis, like other songbirds [50], may preferentially follow coastlines during migration.

However, until more is known about the specific interactions of ice worms with various bird

species, and thus their potential to act as LDD vectors, relating bird migrations to ice worm

distributions will remain difficult. Beyond ice worms, ice sheets have been implicated as a key

driver of speciation in boreal birds [51] and phylogeographic structure of many taxa, from

nematodes to gray wolves [17, 29, 52-54], and our results clearly support these broader

implications for biodiversity accumulation and maintenance in North America.

Conclusions

In this study, we leveraged population genomic data to unravel the complex evolutionary

history of the North American ice worm, *M. solifugus.* Our results add new clarity to previous

perspectives on ice worm biogeography while also lending genomic support to the existence of

contemporary, passerine-mediated LDD in the group. While the phylogenetic data used in this

study (i.e., the lack of an outgroup) preclude us from diagnosing groups I and II as monophyletic,

given the results of previous studies [23-25], we predict that future efforts will diagnose them as

such, perhaps even representing two distinct species. Finally, our genomic data lend support to

the glaciological record in the region, adding a biological line of evidence to a postulated north-

south ice ridge that formed along the crest of the Coast Mountains during the Pleistocene [28].

This potential for genomics to inform the geological record is intriguing and ice worms, as a rare

glacier-obligate macroinvertebrate, may be an ideal taxon for similar studies in the future.

Acknowledgements: We thank Joe Giersch for map making assistance and the University of

Kentucky Center for Computational Sciences for high-performance computing resources. This

research was partially funded by NSF award #IOS-082050 to D.H.S.

Competing interests: We have no competing interests.

Author contributions: S.H., D.H.S., L.M.T, and D.W.W. conceived of and funded the study.

S.H., D.H.S., S.A.L, R.K.B., and D.W.W. collected the data. S.H. and J.L.K. analyzed the data

and wrote the manuscript with input from D.H.S., R.K.B., and D.W.W. All authors approved the

final version.

Data accessibility: Raw sequence data for this study has been submitted to GenBank under

BioProject #PRJNA479335 and code to reproduce the analyses is deposited on GitHub

(https://github.com/scotthotaling/ice_worm_ddRAD).

Funding: Field sampling for this study was supported by NSF award #IOS-0820505 to D.H.S.

References:

1.   Darwin C. 1859 *On the origin of species*, Routledge.

2.   Martin C.H., Turner B.J. 2018 Long-distance dispersal over land by fishes: extremely
rare ecological events become probable over millennial timescales. *Proc R Soc B* **285**(1881),
20172436.

3.   Trakhtenbrot A., Nathan R., Perry G., Richardson D.M. 2005 The importance of long‑
distance dispersal in biodiversity conservation. *Diversity and Distributions* **11**(2), 173-181.

4.   Helmuth B., Veit R., Holberton R. 1994 Long-distance dispersal of a subantarctic
brooding bivalve (Gaimardia trapesina) by kelp-rafting. *Marine biology* **120**(3), 421-426.

5.   Viana D.S., Gangoso L., Bouten W., Figuerola J. 2016 Overseas seed dispersal by
migratory birds. *Proc R Soc B* **283**(1822), 20152406.

6.   Gittenberger E., Groenenberg D.S., Kokshoorn B., Preece R.C. 2006 Biogeography:
molecular trails from hitch-hiking snails. *Nature* **439**(7075), 409.

7.   Miura O., Torchin M.E., Bermingham E., Jacobs D.K., Hechinger R.F. 2012 Flying
shells: historical dispersal of marine snails across Central America. *Proc R Soc B* **279**(1731),
1061-1067.

8.   Pearce T.A., Mulvihill R.S., Porter K.A. 2012 Land slugs (Gastropoda: Pulmonata) on
birds demonstrate dispersal potential. *Nautilus* **126**, 38-40.

9.   Suetsugu K., Funaki S., Takahashi A., Ito K., Yokoyama T. 2018 Potential role of bird
predation in the dispersal of otherwise flightless stick insects. *Ecology*.

10.   Campos-Arceiz A., Larrinaga A.R., Weerasinghe U.R., Takatsuki S., Pastorini J.,
Leimgruber P., Fernando P., Santamaría L. 2008 Behavior rather than diet mediates seasonal
differences in seed dispersal by Asian elephants. *Ecology* **89**(10), 2684-2691.

11.   Anderson J.T., Nuttle T., Rojas J.S.S., Pendergast T.H., Flecker A.S. 2011 Extremely
long-distance seed dispersal by an overfished Amazonian frugivore. *Proceedings of the Royal
Society of London B: Biological Sciences*, rspb20110155.

12.   Viana D.S., Santamaría L., Figuerola J. 2016 Migratory birds as global dispersal vectors.
Trends in ecology & evolution **31**(10), 763-775.

13.   Somveille M., Manica A., Butchart S.H., Rodrigues A.S. 2013 Mapping global diversity
patterns for migratory birds. *PloS one* **8**(8), e70907.

14.   Simpson G.G. 1940 Mammals and land bridges. *Journal of the Washington Academy of*
Sciences **30**(4), 137-163.

15.   Slatkin M. 1987 Gene flow and the geographic structure of natural populations. *Science*
236(4803), 787-792.

16.   Hohenlohe P.A., Day M.D., Amish S.J., Miller M.R., Kamps-Hughes N., Boyer M.C.,
Muhlfeld C.C., Allendorf F.W., Johnson E.A., Luikart G. 2013 Genomic patterns of
introgression in rainbow and westslope cutthroat trout illuminated by overlapping paired-end
RAD sequencing. *Molecular ecology* **22**(11), 3002-3013. (doi:10.1111/mec.12239).

17.   Hotaling S., Muhlfeld C.C., Giersch J.J., Ali O.A., Jordan S., Miller M.R., Luikart G.,
Weisrock D.W. 2018 Demographic modelling reveals a history of divergence with gene flow for
a glacially tied stonefly in a changing post‐Pleistocene landscape. *Journal of Biogeography*
45(2), 304-317.

18.   Jordan S., Giersch J.J., Muhlfeld C.C., Hotaling S., Fanning L., Tappenbeck T.H.,
Luikart G. 2016 Loss of genetic diversity and increased subdivision in an endemic alpine
stonefly threatened by climate change. *PloS one* **11**(6), e0157386.

19.   Antonelli A., Kissling W.D., Flantua S.G., Bermúdez M.A., Mulch A., Muellner-Riehl
A.N., Kreft H., Linder H.P., Badgley C., Fjeldså J. 2018 Geological and climatic influences on
mountain biodiversity. *Nature Geoscience* **11**(10), 718.

20.   Taberlet P., Fumagalli L., Wust-Saucy A., Cosson J. 1998 Comparative phylogeography
and postglacial colonization routes in Europe. *Molecular ecology* **7**(4), 453-464.

21.   Emery C. 1898 Diagnosi di un nuovi genere e nuova specie di annelidi della famiglia
degli Enchytraeidae. *Atti R Accademei Lincei* **5**, 100-111.

22.   Liang Y., Hsu C., Chang T. 1979 A new genus and species of Enchytraeidae from Tibet.
Acta Zootaxonomica Sinica **4**, 312-317.

23.   Dial R.J., Becker M., Hope A.G., Dial C.R., Thomas J., Slobodenko K.A., Golden T.S.,
Shain D.H. 2016 The role of temperature in the distribution of the glacier ice worm,
Mesenchytraeus solifugus (Annelida: Oligochaeta: Enchytraeidae). *Arctic, Antarctic, and Alpine*
Research **48**(1), 199-211.

24.   Hartzell P.L., Nghiem J.V., Richio K.J., Shain D.H. 2005 Distribution and phylogeny of
glacier ice worms (Mesenchytraeus solifugusandMesenchytraeus solifugusrainierensis).
Canadian Journal of Zoology **83**(9), 1206-1213. (doi:10.1139/z05-116).

25.   Dial C.R., Dial R.J., Saunders R., Lang S.A., Lee B., Wimberger P., Dinapoli M.S.,
Egiazarov A.S., Gipple S.L., Maghirang M.R., et al. 2012 Historical biogeography of the North
American glacier ice worm, Mesenchytraeus solifugus (Annelida: Oligochaeta: Enchytraeidae).
Mol Phylogenet Evol **63**(3), 577-584. (doi:10.1016/j.ympev.2012.01.008).

26.   Lang S.A., Saglam N., Kawash J., Shain D.H. 2017 Punctuated invasion of water, ice,
snow and terrestrial ecozones by segmented worms (Oligochaeta: Enchytraeidae:
Mesenchytraeus). *Proc R Soc B* **284**(1864), 20171081.

27.   Figuerola J., Green A.J., Michot T.C. 2004 Invertebrate eggs can fly: evidence of
waterfowl-mediated gene flow in aquatic invertebrates. *The American naturalist* **165**(2), 274-
280.

28.   Clague J.J., Ward B. 2011 Pleistocene glaciation of British Columbia. In *Developments*
in Quaternary Sciences (pp. 563-573, Elsevier.

29.   Shafer A.B., Cullingham C.I., Cote S.D., Coltman D.W. 2010 Of glaciers and refugia: a
decade of study sheds new light on the phylogeography of northwestern North America.
Molecular ecology **19**(21), 4589-4621. (doi:10.1111/j.1365-294X.2010.04828.x).

30.   Peterson B.K., Weber J.N., Kay E.H., Fisher H.S., Hoekstra H.E. 2012 Double digest
RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and non-
model species. *PloS one* **7**(5), e37135. (doi:10.1371/journal.pone.0037135).

31.   Burford Reiskind M., Coyle K., Daniels H., Labadie P., Reiskind M., Roberts N., Roberts
R., Schaff J., Vargo E. 2016 Development of a universal double‐digest RAD sequencing
approach for a group of nonmodel, ecologically and economically important insect and fish taxa.
Molecular ecology resources **16**(6), 1303-1314.

32.   Catchen J., Hohenlohe P.A., Bassham S., Amores A., Cresko W.A. 2013 Stacks: an
analysis tool set for population genomics. *Molecular ecology* **22**(11), 3124-3140.

33.   Purcell S., Neale B., Todd-Brown K., Thomas L., Ferreira M.A., Bender D., Maller J.,
Sklar P., De Bakker P.I., Daly M.J. 2007 PLINK: a tool set for whole-genome association and
population-based linkage analyses. *The American Journal of Human Genetics* **81**(3), 559-575.

34.   Weir B.S., Cockerham C. 1996 Genetic data analysis II: Methods for discrete population
genetic data. Sinauer Assoc. *Inc, Sunderland, MA, USA*.

35.   Wright S. 1943 Isolation by distance. *Genetics* **28**(2), 114-138.

36.   Meirmans P.G., Van Tienderen P.H. 2004 GENOTYPE and GENODIVE: two programs
for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Resources* **4**(4),
792-794.

37.   Alexander D.H., Novembre J., Lange K. 2009 Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res* **19**(9), 1655-1664. (doi:10.1101/gr.094052.109).

38.   Jombart T., Devillard S., Balloux F. 2010 Discriminant analysis of principal components:
a new method for the analysis of genetically structured populations. *BMC genetics* **11**(1), 94.

39.   Malinsky M., Trucchi E., Lawson D.J., Falush D. 2018 RADpainter and
fineRADstructure: population inference from RADseq data. *Molecular biology and evolution*
35(5), 1284-1290.

40.   Chifman J., Kubatko L. 2014 Quartet inference from SNP data under the coalescent
model. *Bioinformatics* **30**(23), 3317-3324.

41.   Swofford D.L. 1998 Phylogenetic analysis using parsimony.

42.   Allendorf F.W., Hohenlohe P.A., Luikart G. 2010 Genomics and the future of
conservation genetics. *Nat Rev Genet* **11**(10), 697-709. (doi:10.1038/nrg2844).

43.   Margold M., Jansson K.N., Kleman J., Stroeven A.P., Clague J.J. 2013 Retreat pattern of
the Cordilleran Ice Sheet in central British Columbia at the end of the last glaciation
reconstructed from glacial meltwater landforms. *Boreas* **42**(4), 830-847.

44.   Stumpf A.J., Broster B.E., Levson V.M. 2000 Multiphase flow of the late Wisconsinan
Cordilleran ice sheet in western Canada. *Geological Society of America Bulletin* **112**(12), 1850-
1863.

45.   Wilson J., Falconer G., Mathews W., Prest V. 1958 Glacial map of Canada. *Geological*
Association of Canada, Toronto, ON.

46.   Riedel J.L., Wilson S., Baccus W., Larrabee M., Fudge T., Fountain A. 2015 Glacier
status and contribution to streamflow in the Olympic Mountains, Washington, USA. *Journal of*
Glaciology **61**(225), 8-16.

47.   Rosvold J. 2016 Perennial ice and snow-covered land as important ecosystems for birds
and mammals. *Journal of biogeography* **43**(1), 3-12.

48.   Shain D.H., Carter M.R., Murray K.P., Maleski K.A., Smith N.R., McBride T.R.,
Michalewicz L.A., Saidel W.M. 2000 Morphologic characterization of the ice worm
Mesenchytraeus solifugus. *Journal of Morphology* **246**(3), 192-197.

49.   Hotaling S., Hood E., Hamilton T.L. 2017 Microbial ecology of mountain glacier
ecosystems: biodiversity, ecological connections and implications of a warming climate.
Environmental microbiology **19**(8), 2935-2948.

50.   Ruegg K.C., Smith T.B. 2002 Not as the crow flies: a historical explanation for circuitous
migration in Swainson's thrush (Catharus ustulatus). *Proceedings of the Royal Society of London*
B: Biological Sciences **269**(1498), 1375-1381.

51.   Weir J.T., Schluter D. 2004 Ice sheets promote speciation in boreal birds. *Proceedings of*
the Royal Society of London B: Biological Sciences **271**(1551), 1881-1887.

52.   Muñoz-Fuentes V., Darimont C.T., Wayne R.K., Paquet P.C., Leonard J.A. 2009
Ecological factors drive differentiation in wolves from British Columbia. *Journal of*
Biogeography **36**(8), 1516-1531.

53.   Burg T.M., Gaston A.J., Winker K., Friesen V.L. 2006 Effects of Pleistocene glaciations
on population structure of North American chestnut‐backed chickadees. *Molecular ecology*
15(9), 2409-2419.

54.   Koehler A.V., Hoberg E.P., Dokuchaev N.E., Tranbenkova N.A., Whitman J.S.,
Nagorsen D.W., Cook J.A. 2009 Phylogeography of a Holarctic nematode, Soboliphyme
baturini, among mustelids: climate change, episodic colonization, and diversification in a
complex host–parasite system. *Biological Journal of the Linnean Society* **96**(3), 651-663.
527