PETISCO is a novel protein complex required for 21U RNA biogenesis and embryonic

viability

Ricardo J. Cordeiro Rodrigues[1], António Miguel de Jesus Domingues[1], Svenja Hellmann[1],

Sabrina Dietz[2], Bruno F. M. de Albuquerque[1,3], Christian Renz[4], Helle D. Ulrich[4], Falk Butter[2],

René F. Ketting[1]

Institute of Molecular Biology (IMB), Ackermannweg 4, 55128, Mainz, Germany

[1] IMB, Biology of Non-coding RNA Group

[2] IMB, Quantitative Proteomics Group

[3]Graduate Program in Areas of Basic and Applied Biology, University of Porto, 4099-003

Porto, Portugal.

[4]IMB, Maintenance of Genome Stability Group

correspondence: r.ketting@imb.de

ORCID Ketting: 0000-0001-6161-5621

ORCID Rodrigues: 0000-0002-0541-2229

ORCID Domingues: 0000-0002-1803-1863

ORCID Renz: 0000-0003-3164-3597

ORCID Ulrich: 0000-0003-0431-2223

Abstract

Piwi proteins are important for germ cell development in almost all animals studied thus far.

These proteins are guided to specific targets, such as transposable elements, by small guide

RNAs, often referred to as piRNAs, or 21U RNAs in *C. elegans*. In this organism, even though

genetic screens have uncovered a number of potential 21U RNA biogenesis factors, little is

known about how these factors interact or what they do. Based on the previously identified

21U biogenesis factor PID-1, we here define a novel protein complex, PETISCO, that is

required for 21U RNA biogenesis. PETISCO contains both potential 5'-cap and 5'-phosphate

RNA binding domains, suggesting involvement in 5' end processing. We define the

interaction architecture of PETISCO and reveal a second function for PETISCO in embryonic

development. This essential function of PETISCO is not mediated by PID-1, but by TOST-1.

Vice versa, TOST-1 is not involved in 21U RNA biogenesis. Both PID-1 and TOST-1 are small,

intrinsically disordered proteins that interact directly with the PETISCO protein ERH-2

(enhancer of rudimentary homolog 2) using a conserved sequence motif. Finally, our data

suggest an important role for TOST-1:PETISCO in SL1 homeostasis in the early embryo. Our

work describes the first molecular platform for 21U RNA production in *C. elegans*, and

strengthens the view that 21U RNA biogenesis is built upon a much more widely used,

snRNA-related pathway.

Introduction

Germ cells in many organisms depend critically on the integrity of a small RNA-driven pathway known as the piRNA pathway [1] [2]. This pathway is characterized by members of the Piwi protein family that, bound to their small RNA cofactors (piRNAs), act in gene-regulatory and transposon silencing pathways [3,4]. In the absence of these pathways, germ cells do not form properly, ultimately leading to sterility. Even though this type of pathway is found in the germ cells of most, if not all animals, the mechanistic details of the Piwi pathways are remarkably different. In flies, for example, a piRNA amplification loop, driven by two Piwi paralogs, is coupled to the activity of a third, nuclear Piwi protein that drives transcriptional silencing [5] [4]. In other animals, such as the silk moth, the nuclear branch seems to be absent, while the Piwi amplification loop is present [6], and in mice a linear Piwi pathway drives nuclear accumulation of a Piwi protein [4]. While transposons represent major targets for these pathways, it is also clear that non-transposon targets are functionally relevant for these pathways [7].

In *C. elegans*, the main Piwi protein is named PRG-1 (piwi related gene 1) and the small RNA co-factors bound by PRG-1 are known as 21U RNAs [8-10]. This name stems from the fact that these small RNAs are 21 nucleotides long, and have a strong bias for a 5' uracil. Target RNA recognition by PRG-1:21U complexes does not depend on full-length base-pairing between the 21U RNA and the target RNA [11,12], and results in the recruitment of RNA-dependent RNA polymerase activity, that drives the synthesis of so called 22G RNAs [8,9]. These 22G RNAs, named after their predominant 22 nucleotide length and 5' G bias, are bound by argonaute proteins that are specific for nematodes (also referred to as WAGO-proteins), that ultimately drive the silencing of the 21U target.

In absence of PRG-1, the germline deteriorates over generations, eventually leading to sterility [13]. The defects that accumulate are most likely not genetic in nature, since the phenotype can be reverted by, for instance, starvation [13]. These observations have led to the suggestion that in *prg-1* mutants a form of stress accumulates and ultimately leads to germ cell dysfunction [14]. More acute fertility defects can be observed in *prg-1* mutants when the 22G RNA biogenesis machinery is reactivated in zygotes, after being defective in both of the parents. In this case, maternally provided PRG-1:21U complexes are required to prevent

immediate sterility, by preventing accumulation of 22G RNA populations that

inappropriately silence genes that should be expressed [15,16]. This demonstrates that PRG-1,

and its bound 21U RNA, has a critical function in maintaining a properly tuned 22G RNA

population in the germ cells. Nevertheless, the impact of PRG-1 on transposon silencing is

rather modest, as in *prg-1* mutants only activation of the Tc3 transposon has thus far been

demonstrated [8].

Interestingly, the silencing of a 21U target, at least a transgenic one, can become

independent of 21U RNAs themselves [5,17,18]. In this state, which has been named RNAe (for

<u>RNA</u>i induced <u>e</u>pigenetic silencing), the silencing has been completely taken over by a self-

sustaining 22G RNA response. This includes a nuclear component that changes the histone

methylation status of the targeted transgene, driven by the nuclear Argonaute protein

HRDE-1 [5,17-19]. Possibly, such an RNAe state may explain why transposons are not more

broadly upregulated in *prg-1* mutants, because in *prg-1;hrde-1* double mutants the Tc1

transposon is strongly activated [15]. How exactly RNAe is established is not clear, but once it

is, it can be extremely stable, for over tens or more of generations [5,17,18].

Given the important function of 21U RNAs driving a potentially very powerful silencing

response, the biogenesis of 21U RNAs is a critical process for the germ cells of *C. elegans*.

Nevertheless, very little is known about the mechanistic steps of this process. The large

majority of genes encoding 21U RNAs are found in two main clusters on chromosome IV,

and are characterized by a very specific sequence motif in their promoter that defines the 5'

end of the mature 21U RNA [20]. Transcription of these genes requires a protein named PRDE-

1 and the transcription factor SNPC-4 [21,22], the latter of which is also known to be involved in

transcription of other short structural RNAs, such as snRNAs and splice-leader RNAs [21]. An

evolutionary analysis of 21U RNA loci in diverse nematodes has revealed that 21U loci may

have evolved from snRNA loci [23]. These loci include both the strongly conserved U1, U2 loci,

as well as loci producing so-called splice leader RNAs (SL1 and SL2), that are trans-spliced to

the 5' ends of a large fraction of all mRNAs in *C. elegans* [24]. These observations raise the

possibility that also other aspects of the 21U RNA pathway may have mechanistic links to

snRNA biogenesis.

The 21U RNA precursor transcripts are short, around 27 nucleotides, and capped [25]. Even
though a biochemical reconstitution of the processing has not been achieved thus far, the
available data suggest the following order of steps in the maturation of the 21U precursor
transcripts into mature 21U RNAs [20,22,25-27]. First, the precursors are most likely processed at
the 5' end, resulting in de-capping and removal of two nucleotides. The enzymes involved
have not yet been identified, and whether this reaction is mediated by endo- or exo-
nucleolytic activities is not clear. This step is likely followed by loading of the 5'processed
precursor into PRG-1 and trimming of the 3' end by the 3'-5' exonuclease PARN-1 [28]. Finally,
the 3' end is 2'-O-methylated by HENN-1 [29-31]. Not much is known about other proteins
acting at these 21U maturation steps, even though a number of genes have been implicated
in this process [21,22,26,27 32].

Here, we follow up on our previous identification of PID-1 (piRNA induced silencing defective
1) as a protein essential for 21U RNA production [26]. Mutants lacking PID-1 produce very low
amounts of mature 21U RNAs, and the 21U-related molecules that remain tend to be
precursor transcripts, suggesting PID-1 acts at some step in 21U precursor processing. Other
factors potentially acting at this step of 21U biogenesis, TOFU-1, 2, 6 and 7, were identified
in a genome-wide RNAi screen [27]. How these factors interconnect, however, remained
unclear. We find, using immuno-precipitation (IP) and label free quantitative mass
spectrometry (IP-MS), that PID-1 interacts with two proteins that were identified in a
genome-wide RNAi screen for 21U RNA biogenesis factors: TOFU-6 and the unnamed protein
Y23H5A.3, henceforth referred to as PID-3. In addition, we identify two strongly conserved
proteins interacting with PID-1: IFE-3, a *C. elegans* eIF4E homolog, and ERH-2, one of the *C.*
elegans homologs of 'enhancer of rudimentary'. Enhancer-of-rudimentary homologs are
evolutionary very well conserved proteins, with homologs being present from plants to man.
Its mechanistic role is not very well established, but in *Schizosaccharomyces pombe* ERH1
drives the decay of meiotic transcripts, and interacts with the nuclear exosome and the
nuclear CCR4-NOT complex [33]. Since we always find PID-3, ERH-2, TOFU-6 and IFE-3 together
in a complex we named it PETISCO, for PID-3, ERH-2, TOFU-6, IFE-3 small RNA Complex. All
PETISCO proteins are required for 21U biogenesis. Additionally, PETISCO mutants display a
maternal effect lethal (Mel) phenotype, whereas *pid-1* and *prg-1* mutants are viable. We find
that this is caused by the fact that besides binding to PID-1, PETISCO can bind a protein with

similarities to PID-1. We named this protein TOST-1, for <u>T</u>wenty-<u>O</u>ne U pathway antagoni<u>st</u>.

Mutants for *tost-1* produce 21U RNAs, display enhanced 21U-driven silencing and have a

Mel phenotype. PID-1 and TOST-1 both interact with ERH-2 using a conserved motif, strongly

suggesting a mutually exclusive mode of binding to PETISCO, where PID-1 implicates it in

21U RNA biogenesis and TOST-1 in another pathway that is essential for embryonic viability.

Our data suggest that this pathway may be involved in providing the embryo with resources

required for trans-splicing, especially with the SL1 splice leader RNA.

With these findings, we pave the way for a better understanding of how 21U precursor

transcripts are processed. The identification of PETISCO, plus its implication in at least two

distinct processes provides the new insight that 21U RNA processing is closely related to a

more widely conserved process: that of snRNP biogenesis. Combined with the fact that 21U

RNA transcription also bears strong signatures of an snRNA-history, an image arises in which

the 21U RNA pathway may have evolved out of an already existing small non-coding RNA

network that became linked to an Argonaute-driven gene-silencing program.

Results

Identification of PID-1 interactors

To identify proteins interacting with the 21U biogenesis factor PID-1 [26] we performed IPs with PID-1 specific antibodies, followed by protein identification using mass spectrometry (IP-MS). As control, we precipitated PID-1 from two independent *pid-1* loss of function strains. Both experiments identified a rather restricted set of proteins (Figure 1a, S1a). TOFU-6 and PID-3 were identified as the most prominent PID-1 interactors, and for the latter we further validated the interaction with PID-1 through IP-Western blotting (Figure S1b). Given that these two proteins were identified in an RNAi-screen for 21U biogenesis factors [27], we considered these as functionally relevant PID-1 interactors. TOFU-6 is a protein with a Tudor domain, a potential eIF4E interaction motif and an RRM domain, whereas PID-3 has an RRM domain and a MID-domain (Figure S1c). Two other proteins were consistently identified as PID-1 interactors: IFE-3 and ERH-2 (Figure 1a, S1a). IFE-3 is one of the five *C. elegans* homologs of eIF4E. Previous work demonstrated that IFE-3 binds to the 7-methylguanylate (m7G) Cap [34]. Interestingly, 21U precursor transcripts appear to be not trans-spliced [25], implying that they have a m7G Cap structure at their 5' end. Hence, IFE-3 may bind to the 5' cap structure of the 21U precursor transcripts. ERH-2 is one of the two *C. elegans* homologs of a protein known as 'enhancer of rudimentary'. Homologs of this protein are strongly conserved from plants to mammals. Even though a protein structure for the human homolog has been described [35], its molecular function is still unclear. In Figure S1c we present a schematic of all these PID-1 interactors, with their identified domains.

We next probed the subcellular localization of PID-1 and its four identified interactors, by expressing them from transgenes generated through the so-called miniMos approach (see methods) [36]. For each gene we used its own promoter and 3' UTR sequences. All transgenes were able to rescue mutant phenotypes (not shown) indicating the expressed proteins are functional. All four interactors show a characteristic perinuclear, punctate localization in the germline syncytium, overlapping partially, but not fully with the P-granule marker PGL-1 (Figure 1b,c). IFE-3 has a clear granular expression in the primordial germ cells in the embryo (Figure S2b), whereas the remaining interactors show a dispersed cytoplasmic distribution across the entire early embryo (Figure S2c). These results show that all the identified PID-1

interactors are expressed concomitantly in early embryos, and in the germline, where they

are found in close proximity to P-granules.

PID-1 interactors mutually interact to form PETISCO

To probe to what extent the identified PID-1 interactors reciprocally interact, and to what

extent they in turn interact with additional proteins, we performed IP-MS on TOFU-6, IFE-3,

PID-3 and ERH-2 in young adult animals, using the tagged proteins expressed from the

above-described transgenes. Extracts from non-transgenic wild-type animals were used as

negative controls. As shown in Figure 2a-d, these experiments revealed that all four proteins

co-precipitate with each other. In addition, we also found an uncharacterised protein

(C35D10.13) which systematically co-precipitated with the PID-1 interactors, but was absent

from the PID-1 IPs. This factor, which we named TOST-1, will be further described below.

These interactions are RNAse resistant. RNase treatment did not disrupt PID-3 interactions

(Figure S3a), and had very little effect on IFE-3 partners, resulting only in the loss of PID-1

(Figure S3b). We summarize the network of PID-1 interactors in Figure 2e. Given the strong

reciprocal interactions between these proteins, it is likely that they form a discrete complex.

We named this complex PETISCO, for PID-3, ERH-2, TOFU-6, IFE-3 small RNA Complex.

As mentioned before, IFE-3 is one of the *C. elegans* eIF4E homologs. Surprisingly, in our

experiments we do not detect any of the known translation initiation factors to be

associated with IFE-3. The transgenically expressed IFE-3 does rescue the *ife-3* mutant

phenotype (not shown), implying that IFE-3 may not play a role in initiating translation. We

do detect a number of additional proteins bound to IFE-3 which provide clues to IFE-3

function. One such protein is IFET-1, a homolog of human EIF4E nuclear import factor 1 and

a negative regulator of translation [37]. We further consistently detect many, if not all

components of the SMN complex [38]. The SMN complex plays a major role in snRNP

biogenesis, implicating IFE-3 in this process as well.

PETISCO architecture

Having established the components of PETISCO, we next dissected the molecular

interactions within this complex. Using the yeast-two-hybrid (Y2H) system we scored

interactions between each possible pair of individual PETISCO subunits (Figure 3a, S4a). This

resulted in the following interactions that are most likely to be direct: IFE-3 interacts only

with TOFU-6. TOFU-6 in turn interacts with PID-3, which in turn also interacts with ERH-2.

Finally, PID-1 and TOST-1 were both found to interact only with ERH-2. Self-interaction was

detected for ERH-2, though under lower stringency conditions, indicating that ERH-2 may be

present as a dimer in PETISCO.

We also investigated which domains of the PETISCO subunits are responsible for the protein

interactions using the same Y2H set-up. The putative eIF4E interaction motif [39] [40] within the

C-terminus of TOFU-6 indeed is responsible for the interaction between TOFU-6 and IFE-3

(Figure 3b). No interactions were found for the Tudor domain of TOFU-6. The RRM domain

of TOFU-6 was found to bind to the RRM-domain of PID-3 (Figure 3c), and the same RRM

domain of PID-3 was found to also interact with ERH-2 (Figure 3d). Lower stringency and

control selections of the same experiments are shown in Figures S4b-d). This Y2H set-up did

not allow us to determine if the PID-3 RRM domain can sustain both the TOFU-6 and the

ERH-2 interaction simultaneously, however the fact that we find PID-3, TOFU-6 and ERH-2

strongly enriched in the IPs of one another, suggests that it can.

PETISCO subunits are required for 21U RNA biogenesis and are essential for

embryogenesis

Since at least two components of PETISCO, PID-1 and TOFU-6, play a major role in 21U RNA

biogenesis, we hypothesized that the other components are also part of this pathway. We

first tested whether they affect the silencing of a GFP sensor construct that reports on the

activity of the 21U pathway (21U sensor) [11]. We used a strain that contains this sensor plus a

pid-1(xf14) mutation that is rescued by transgenic expression of PID-1. In this strain, the

sensor is silenced (Figure 4a,b), although not completely. This partial 21U sensor-silencing

brings two advantages: first, the 21U sensor is not in an RNAe-state, in which it would no

longer be activated by loss of 21U RNAs; second, a semi-silenced state allows for scoring of

both increased and decreased sensor activity.

Animals from this sensitized sensor strain were subjected to RNAi targeting the various

PETISCO subunits and sensor activity was evaluated by microscopy and RT-qPCR (Figure

4a,b). As expected, RNAi against *pid-1* activated the sensor. Likewise, RNAi against *ife-3*, *pid-*

3, *tofu-6* and *erh-2* activated the sensor. These data show that PETISCO subunits, like the

PETISCO interactor PID-1, are required for a fully functional 21U RNA silencing pathway.

To extend the above observation to the overall 21U RNA population, we aimed to sequence

small RNAs from genetic mutants. For this, we created mutant alleles using CRISPR-Cas9 for

each of the PETISCO subunit genes. We were able to derive deletion alleles for *pid-3*, *erh-2*

and *ife-3* (Figure S5a), and isolated and sequenced small RNAs. We did not make mutants for

tofu-6, since RNAi on *tofu-6* was already shown to significantly reduce 21U RNA production

[27]. All experiments were done in triplicates. The results show that all tested PETISCO

subunits significantly affect 21U RNA biogenesis (Figure 4c). In particular, *pid-3(tm2417)* and

erh-2(xf168) mutants display a strong reduction in 21U RNA levels. The effect of *ife-3(xf102)*

mutation on 21U RNA accumulation is less pronounced, but still significant. Redundancy

between IFE-3 and other eIF4E homologs, could be a reason for this less pronounced effect

on 21U RNA levels, as we find IFE-1, a non-selective TMG/m7G binder [34], enriched in some of

our IP-MS experiments (Figure 2a-c, S3b). Other types of small RNAs, such as 22G, 26G and

miRNAs were mostly not affected by the tested mutations (Figure 4c and S5b-d). We note

that so-called type II 21U RNAs, which come from loci lacking the canonical Ruby motif and

are expressed at much lower levels [25], are only mildly, or not at all affected by loss of

PETISCO components.

Besides the defects in 21U biogenesis, the mutant alleles also display a so-called maternal

effect lethal (Mel) phenotype: homozygous mutant offspring from a heterozygous animal

develop into fertile adults, but the embryos display a fully penetrant arrest and never hatch

(Table 1). This phenotype has already been described for *tofu-6*, which is also known as *mel-*

47 [41]. The *ife-3* mutant displays a mixed phenotype, as previously described [42]: homozygous

mutant offspring from a heterozygous animal develop into adults that can either display a

masculinized germline (Mog) or a Mel phenotype. Interestingly, we note that *pid-1* mutants

also display a Mog phenotype, albeit at low frequency (Table 1 and Figure S6).

PID-1 and TOST-1 define distinct functions of PETISCO

The fact that PETISCO genes are essential for embryogenesis contrasts with the fact that loss

of 21U RNAs through other means, such as loss of PRG-1 or PID-1, does not result in a Mel

phenotype. Interestingly, RNAi-knockdown of *tost-1* resulted in an embryonic lethal

phenotype (not shown). In our IP-MS experiments, we consistently found this protein to

interact with the PETISCO subunits (Figure 2a-d), yet not with PID-1 (Figure 1a and S1a).

Furthermore, *tost-1(rnai)* displays enhanced, rather than disrupted silencing of the

sensitized 21U sensor (Figure 4a,b). This phenotype lead us to name this protein TOST-1,

short for twenty-one U antagonist.

We tested both PID-1 and TOST-1 for interactions with PETISCO subunits. We found that

both PID-1 and TOST-1 specifically interact only with ERH-2 (Figure 3a). While the overall

amino-acid sequences of PID-1 and TOST-1 do not display convincing homology (Figure S7a),

when aligned with PID-1 and TOST-1 homologs from other nematode species, a conserved

motif emerges $(_{[+][+]\Psi(T/S)}\mathbf{L}_{(N/S)[-]}\mathbf{RF}_{x\Psi xxx}\mathbf{G}_{(Y/F)}$ – Figure 5a). Strikingly, the *xf14* allele we

identified for *pid-1* in our previously described genetic screen caries a mutation of the fully

conserved arginine residue within this motif (R61C) [26]. When introduced into the Y2H

experiment, PID-1(R61C) did not interact with ERH-2, and the analogous mutation in TOST-1

also disrupted its ERH-2 interaction (Figure 5b). These data strongly suggest that both PID-1

and TOST-1 share a conserved short motif required for ERH-2 interaction.

To further understand TOST-1 function we generated loss of function alleles for *tost-1* using

CRISPR-Cas9 (Figure S5a). Loss of TOST-1 did not significantly affect 21U RNA biogenesis

(Figure 5c, S7b), but did result in a fully penetrant Mel phenotype (Table 1). Among the

alleles we generated, we found one allele, *tost-1(xf196)*, with a 58bp deletion which

removes the splice acceptor site of its third exon (Figure S5a), resulting in either a frame-

shift or a C-terminal truncation. *tost-1(xf196)* displays a temperature-sensitive (TS) effect: at

25°C the animals are fully Mel, while at 15°C the animals are viable. The fact that this allele

appears to be mostly functional at 15°C suggests that the N-terminal part of TOST-1 is the

critical part of this protein. The interaction motif we identified is still intact in *tost-1(xf196)*,

consistent with the idea that it is essential for TOST-1 function. This TS allele allowed us to

probe the Mel phenotype in more detail, through temperature-shift experiments (Figure

S7c). First, animals grown at the restrictive temperature were able to produce viable

offspring after shifting them to the permissive temperature. This shows that there are no

structural or developmental defects in the germline of these animals that prevent them from

producing live offspring. Second, the TS allele allows us to probe when TOST-1 function is
required. The time required for animals shifted to the permissive temperature to produce
viable embryos (8 hours) is significantly longer than the time it takes from fertilization to
egg-laying (approximately 200 min at 15^{o}C), implying that the embryos laid after 8 hours
were fertilized and raised at the permissive temperature. Yet, they are still arresting. This
observation places TOST-1 activity within the gonad. However, when animals are shifted
from permissive to restrictive temperature, the first arrested embryos are laid already after
two hours. This is very close to the time of residency within the uterus (approximately 150
min 25^{o}C), and hence these embryos had been fertilized very close to the time of the
temperature shift, in a gonad that was still functional. This suggests TOST-1 also has a
function within the embryo. This is consistent with PETISCO subunit expression within early
embryos (Figure S2c).

In conclusion, our data shows that both TOST-1 and PID-1 bind to the same PETISCO subunit,
while loss of these two proteins results in different phenotypes. The combined phenotypes
of both *pid-1* and *tost-1* mutants is found in mutants for the other PETISCO subunits,
strongly suggesting that PID-1 and TOST-1 define two distinct aspects of PETISCO function.

PETISCO interacts with SL1 RNA
Given the known links between the 21U RNA pathway and snRNA transcription, the fact that
we find many SMN-complex components in IFE-3 IPs, and that at least one SMN component
has a Mel phenotype (MEL-46), we hypothesized that the essential function of PETISCO is
linked to snRNA homeostasis. Consistent with this idea, we noticed a striking accumulation
of a 3' fragment of SL1 RNA in a number of the mutants we tested (Figure 6a,b, S8a). Such
accumulation was less pronounced, or absent for another small non-coding RNA that is
produced from the same gene cluster as SL1, 5S rRNA (Figure 6b,c). Importantly, this
enrichment of SL1 fragments was very clear in *tost-1* mutants, but absent in *pid-1* mutants.
Also *ife-3* mutants and *erh-2* mutants displayed a similar accumulation in all the replicates
that we sequenced. Mutants for *pid-3*, however, did not show the accumulation (Figure 6b).
For SL2 we only observe the enrichment in *ife-3* mutant libraries (Figure S8b), but we point
out that SL2 read counts are comparatively very low.

To further test a potential link between PETISCO and SL1 RNA, we probed whether SL1, or

SL1 fragments may be bound by PETISCO. To test this, we performed RIPseq experiments on

IFE-3 and PID-3. We performed these experiments in quadruplicate, and also performed

triplicate mock IPs from wild-type animals. Since full SL1 transcripts, but also 21U precursors

are capped, we treated the RNA with RppH to remove 5' caps before cloning. We detect

strong and significant enrichment of SL1, and to a lesser extent SL2 sequences in both PID-3

and IFE-3 IPs (Figure 6c, S8c). In contrast, transcripts from 21U loci, 5S rRNA and snRNAs are

not enriched, or are even depleted (Figure 6c, S8c). The read coverage over the SL1

transcript shows that two fragments are detected: one at the 5' and another at the 3' end

(Figure 6d, S8d). The 5' fragment is capped, as this fragment is absent from libraries from

non-RppH treated RNA (Figure 6d). We conclude that PETISCO binds to SL1 RNA and that loss

of PETISCO leads to an accumulation of SL1 RNA fragments.

Discussion

We have identified a protein complex, named PETISCO, that is involved in at least two different pathways. PID-1 guides PETISCO towards 21U RNA biogenesis, whereas TOST-1 brings PETISCO to a pathway that is essential for embryogenesis, but does not involve 21U RNAs. Here we discuss various aspects of this complex and present hypotheses for the different potential functions of PETISCO.

PID-3 and TOFU-6 define the core of PETISCO

Judging by the enrichments found in all IPs and the similarities in phenotypes of the corresponding mutants it seems likely that two proteins, PID-3 and TOFU-6 form the core of the PETISCO complex, and that the other identified factors may add as adaptors for specific functions of PETISCO. What could the core function(s) of this complex be? Based on protein identities and obtained results we consider the following, non-exclusive possibilities.

Based on the presence of two RRM domains, one in TOFU-6 and one in PID-3, this complex likely has RNA binding activities. Indeed, our RIPseq experiment identified 3' fragments of SL1 and SL2 bound to PETISCO. Interestingly, we found that both RRM domains act as protein-protein interaction modules. This does not mean that these two RRM domains cannot also be involved in RNA binding, as combined RNA- and protein-binding activities for RRM domains have been described before [43]. Possibly, the assembly of the RRM-RRM contact between PID-3 and TOFU-6 is facilitated by RNA. Even though we find that RNase treatment does not affect the pull-down efficiency between PID-3 and TOFU-6, the RNA involved may be inaccessible to RNases.

In addition to the RRM motifs, the MID-domain present in PID-3 also brings potential for protein-RNA-interactions. The MID-domain has thus far been described in context of argonaute proteins where it mediates interactions with the 5' end of the bound small RNA co-factor [44]. It is curious to find such a domain in a protein involved in generating small RNA co-factors for an argonaute protein. We hypothesize that the MID domain of PID-3 may help to stabilize a 5' processing intermediate. This would imply PETISCO as a 5'end processing platform for 21U RNAs. Biochemical reconstitution experiments will be required to test this.

The Tudor domain of TOFU-6 likely represents a protein-protein interaction unit. Tudor domains are known to interact with methylated arginines, and within perinuclear granules, where PETISCO is found, such methylated arginines are likely abundant [45]. Hence, the Tudor domain may be involved in sequestering PETISCO to these granules, possibly to PRG-1. PRG-1 itself has an N-terminal protein sequence that is highly suggestive of arginine methylation. In one of our IFE-3 pull-downs, we do enrich for PRG-1 (Figure 2a), supporting this possibility.

ERH-2: Bridge between core-PETISCO and PETISCO-guiding factors?

The presence of ERH-2 provides hints for PETISCO function. In *S. pombe*, Erh1 forms a tight complex with a protein named Mmi and together they are involved in nuclear mRNA degradation of meiotic transcripts, involving proteins such as ARS2 and the CCR4-NOT complex [33]. Hence, Erh1 in *S. pombe* appears to act as a bridge between an RNA binding protein (Mmi) and an RNA processing machinery. Our data strongly suggest that ERH-2 in *C. elegans* has a very similar function. On the one hand, it binds core PETISCO and on the other hand either PID-1 or TOST-1, which set the function of the complex (see below).

ERH-related proteins in other systems, including *S. pombe* and human cells, are nuclear proteins [33], while we find ERH-2 to be present mainly in perinuclear granules. Related to this it is interesting to note that *C. elegans* has a second ERH-like protein T21C9.4. Alignments show that this homolog is more closely related to *S. pombe* and human ERH1 (not shown), raising the possibility that T21C9.4 in *C. elegans* is nuclear and may be coupled to nuclear RNA processing.

Bi-functionalization of PETISCO through PID-1 and TOST-1

We show that PETISCO has at least two functions: 21U RNA biogenesis and another, essential function in embryogenesis, which is likely related to SL1/2 snRNP homeostasis. These two functions can be functionally separated through two different 'adapter'-proteins (PID-1 and TOST-1) that bind to PETISCO via ERH-2. The overall homology between PID-1 and TOST-1 is very low. Nevertheless, we identified a motif that is required for binding to ERH-2. Hence, both proteins likely bind to PETISCO in a mutually exclusive way. Indeed, no TOST-1 was detected in PID-1 IPs, and Zeng et al. (accompanying manuscript) did not find PID-1 in their TOST-1 IPs. An IP-MS experiment on PID-3 in absence of PID-1 (therefore enriching for

TOST-1:PETISCO) did not reveal much change compared to wild-type (not shown), suggesting

that stable interactions within PID-1:PETISCO and TOST-1:PETISCO are very similar and do

not provide further insights into the differential activities of PID-1 and TOST-1.

Potential roles for IFE-3

IFE-3 has been shown to bind m7G Cap structures, and to bind much less efficiently to the

typical TMG Cap structures found on the majority of trans-spliced mRNAs in *C. elegans* [34].

This activity of IFE-3 makes it a good candidate for binding 21U RNA precursors at a stage

before 5' processing. IFE-3 IP-MS experiments also enrich mildly for proteins involved in de-

capping (e.g. EDC-3, CAR-1, CGH-1 – Figure 2a). In relation to 21U RNA biogenesis, it is an

intriguing option that IFE-3 binds 21U precursor transcripts through their 5' cap structure,

after which they become de-capped by associated de-capping activities. IFE-3, however, is

not essential for 21U generation. This could be due to either functional redundancy with IFE-

1, which we regularly detect in our IP-MS experiments. Alternatively, IFE-3 has no direct role

in 21U RNA processing, and the small drop of 21U levels in *ife-3* mutants is an indirect effect,

for instance through effects on PETISCO overall stability.

We find that IFE-3 consistently comes down with many, if not all proteins found in the so-

called SMN complex. This complex is well known for its involvement in snRNP assembly [38]. In

C. elegans, the SMN complex has also been shown to be required for SL1 trans-splicing [46].

We note that a homolog for Gemin5, which has been shown to bind the m7G Cap structure

of pre-mature snRNAs in human cells [47], is not encoded by the *C. elegans* genome. Given the

cap-binding activity of IFE-3 and its association with the other SMN complex subunits, IFE-3

may fulfil this function in *C. elegans*. We note that like loss of IFE-3 [42], loss of the *C. elegans*

Gemin3 homolog, named MEL-46 [41], and the U2 snRNP-associated factor MOG-2 [48] also

result in Maternal effect lethality (Mel) and Masculinization of the germline (Mog)

phenotypes, further strengthening the links between IFE-3 and snRNP homeostasis. Given

that the SMN-complex proteins are not found in the IPs of any of the other PETISCO

subunits, the IFE-3-SMN interaction is likely to be physically separated from PETISCO. Finally,

it is interesting to note that *pid-1* mutants, but not *tost-1* mutants, also display a Mog

phenotype, albeit at a low frequency. Considering the interplay between PID-1 and TOST-1,

this could relate from excessive, or ectopic TOST-1:PETISCO in *pid-1* mutants affecting the

IFE-3-SMN interplay.

Potential mechanisms behind the Mel and 21U phenotypes of PETISCO mutants

Our data clearly show that PETISCO has at least two functions. Its role in 21U RNA biogenesis

is non-essential and is guided by PID-1. A second function, which is essential for early

development, is linked to PETISCO via TOST-1. While our current data do not provide

mechanistic details on molecular PETISCO function, they do provide interesting leads as to

what the essential function of PETISCO may be.

As mentioned above, PETISCO contains proteins with domains that interact with the 5' ends

of RNA, either capped or phosphorylated, raising the possibility PID-1-bound PETISCO plays a

role in 5' end processing of 21U precursor transcripts. In our RIP experiments, we did not

detect 21U RNA precursor molecules, suggesting that either levels are too low to be

detected in a RIP experiment, or that their processing is too fast. Given that RIP experiments

typically have high backgrounds, this may easily prevent detection of significant

enrichments. In view of the phenotypes and protein domains within PETISCO, we consider it

very likely that PETISCO does interact with 21U RNA precursors.

But what could the molecular function of PID-1-bound PETISCO be? In absence of PETISCO

21U RNA levels drop strongly, but we note that so-called type II 21U RNA levels are much

less affected (Figure S5b). These 21U RNA species do not derive from loci with the typical

Ruby-motif, and are expressed at much lower levels than type I 21U RNAs. We hypothesize

that PETISCO may function to specifically stimulate the processing of pre-cursors that come

from Ruby-motif-containing 21U loci, such that the PRG-1 RNP pool is dominated by PRG-1

protein bound to type I 21U RNAs.

TOST-1-bound PETISCO may have a similar function, only then coupled to the production of

an RNP that is essential for early development. Our RIPseq and mutant small RNAseq data

indicate that this could involve the SL1 snRNP. Our interpretation of these data is that

PETISCO binds full-length SL1 transcripts, but that these are partially degraded by nucleases

during the RIP-procedure, or within the animals when further processing of the complex is

stalled due to loss of PETISCO subunits such as TOST-1 or IFE-3. But how would this explain
the maternal effect lethality phenotype of PETISCO mutants? SL1 is a generally required
splice-leader RNA, and mutants that carry a large deletion covering the SL1 loci display a
direct embryonic lethal phenotype [49]. If PETISCO would have a core function in trans-splicing
one would expect a similarly severe zygotic phenotype. We hypothesize that PETISCO
specifically helps the accumulation of a large pool of SL1 snRNPs that will be maternally
loaded into the embryos. Later in development, when the demand for SL1 snRNPs may have
dropped, PETISCO may therefore not be required. This would create an interesting parallel
to 21U RNPs, that are also provided maternally [15].

Clearly, additional studies will be required to resolve the molecular function of PETISCO in
21U RNA processing and further test its role in SL1 homeostasis during early development.
However, our data firmly show that PETISCO is a 21U biogenesis complex, and in addition
has an essential role during early development. Our work shows that the processing of 21U
RNAs is likely derived from a much more widely conserved mechanism, and provides a new
and exciting view into how small RNA biogenesis can be intertwined with other gene-
regulatory mechanisms.

Figure Legends

Figure 1 – PID-1 interactors reside in P-Granules

a) Volcano plot representing label-free proteomic quantification of PID-1 IPs from non-
gravid adult extracts. IPs were performed and analyzed in quadruplicates. The x-axis
represents the median fold enrichment of individual proteins in wild type (WT) versus
pid-1(xf14) mutant strain. y-axis indicates -Log_{10}(p-value) of observed enrichments.
Dashed lines represent thresholds at p=0.05 and 2-fold enrichment. Blue data points
represent values out of scale. Red and Green data points represent values above and
below threshold, respectively.

b and c) Expression pattern and localization of tagged PETISCO components and P-
granule marker PGL-1. Endogenous promotors and 3'UTRs were used for PETISCO
proteins. Proteins and observable tags indicated in the panels. **b)** Immunostaining
images acquired with laser scanning confocal microscope. **c)** Live worm images
acquired under wide field fluorescent microscope. Scale bars represent 5 ɏm.
Contrast of images has been enhanced.

Figure 2 – PID-1 interactors form a novel protein complex, PETISCO

a-d) Volcano plot representing label-free proteomic quantification of quadruplicate **a)**
3xFLAG::mCherry::IFE-3;*ife-3(xf101)*; **b)** PID-3::mCherry::Myc;*pid-3(tm2417)*; **c)** TOFU-
6::GFP::HA;*tofu-6(it20)* and **d)** ERH-2::GFP::OLLAS;*erh-2(xf168)* IPs from non-gravid
adult extracts. The x-axis represents the median fold enrichment of individual
proteins in control (WT) versus transgenic strain. y-axis indicates -Log_{10}(p-value) of
observed enrichments. Dashed lines represent thresholds at p=0.05 and 2-fold
enrichment. Blue data points represent values out of scale. Red and Green data
points represent above and below threshold respectively.

e) Venn Diagram summarizing significant interactions in PETISCO protein IPs.
*represents protein found significantly enriched in only one experiment of
3xFLAG::mCherry::IFE-3;*ife-3(xf101)* IP.

Figure 3 – PETISCO Architecture

a-d) Yeast two-hybrid interaction assays of PETISCO subunits in low (TRP⁻LEU⁻HIS⁻) or high
stringency media (TRP⁻LEU⁻HIS⁻ADE⁻) as indicated. Interactions were screened in
both Y2H orientations. **a)** Full length proteins **b)** TOFU-6 and individual domains
tested for interaction with full length IFE-3 **c)** Interactions between PID-3 and TOFU-
6 **d)** Interaction between PID-3 and ERH-1. For details on domains and other
selection conditions see Figure S4.

Figure 4 – PETISCO is required for 21U RNA biogenesis

a) Wide field fluorescent microscopy of adult hermaphrodites carrying GFPH2B-21U
sensor transgene in a sensitized background. Worms were subjected to RNAi via
feeding (targets indicated in figure) from L1 larval stage to adulthood. Empty RNAi
Vector serves as negative control. Scale Bar 10 ɏm.

b) Quantitative RT-PCR of 21U sensor transgene in adult worm populations of A). Values
are obtained from experimental triplicates and technical duplicates, normalized to

pmp-3 mRNA levels. Significance was tested with Student's t-test: **p-value<0.01;
*p-value<0.05. Error bars represent standard deviation.

c) Global level of type I 21U RNA and miRNAs in the indicated strains. Values are in
reads per million (RPM). Individual data points of three independent replicates are
shown and horizontal bar represents the mean. Significance was tested with
Student's t-test and p-values are indicated in the graph.

Figure 5 – TOST-1 is essential but not required for 21U RNA biogenesis
a) Alignment of a short region of PID-1 and TOST-1 homologs from different nematodes.
Conserved arginine residue was found to be mutated in *pid-1(xf14)* as indicated.
Consensus sequence is presented below alignment. Alignment performed with
MUSCLE v3.8 [50] and representation with ESPrit v3.0 [51].

b) Y2H interaction assay of PID-1, TOST-1 and PID-1/TOST-1 carrying the corresponding
arginine mutation found in pid-1(xf14). High stringency plates (TRP⁻LEU⁻HIS⁻ADE⁻)
were used in the presented figure. For other conditions please see Figure S4d.

c) Global levels of type I 21U RNA and miRNAs in wild type (N2), *pid-1(xf35)* and *tost-*
1(xf194) gravid adult worms. Values are in reads per million (RPM). Individual data
points of three independent replicates are shown and horizontal bar represents the
mean. Significance was tested with Student's t-test and p-values are indicated in the
graph.

Figure 6 – PETISCO interacts with SL1 snRNA
a) Schematic representation of the Splicing Leader 1 RNA.
b) Global levels of SL1 RNA and 5S rRNA in wild type (N2), *pid-1(xf35)* and *tost-1(xf194)*
gravid adult worms and *pid-3(tm2417), erh-2(xf168)* and *ife-3(xf102)* non-gravid
adults. Values are in reads per million (RPM). Individual data points of three
independent replicates are shown and horizontal bar represents the total mean.
Significance was tested with Student's t-test and p-values are indicated in the graph.

c) Fold enrichments of SL1 RNA and 5S rRNA in Mock (N2), 3xFLAG::mCherry::IFE-3;*ife-*
3(xf101); and PID-3::mCherry::Myc;*pid-3(tm2417)* RIPs over paired inputs in non-
gravid adult worms. Top row displays non-treated and bottom row RppH treated
samples. Individual data points of three independent replicates are shown and
horizontal bar represents the mean. Significance was tested with Student's t-test and
p-values are indicated in the graph.

d) Coverage profile, normalized to paired input, of SL1, of the data displayed in **c**. Colors
under SL1 RNA correspond to scaled colors represented in **a.**

Figure 7 – Schematic representation of PETISCO function
A schematic of the proposed PETISCO activity, displaying its dual function. One in 21U
biogenesis and another potentially linking to splice leader homeostasis. The two different 5'
end binding domains may reflect stabilization of different RNA species, and may reflect 5'
end processing of transcripts bound by PETISCO. ERH-2 serves as an anchor for PID-1 or

TOST-1 driving PETISCO function towards 21U RNA biogenesis or SL1 homeostasis
respectively.
588
589 **Table 1 – PETISCO displays maternal effect lethality**
590 "n.a." not applicable; "n.d." not determined; "-" mild 21U RNA defect; "--" severe 21U RNA
591 defect; "+" no 21U RNA defect; "(TS)" temperature sensitive; "#" counts are for gonadal
592 arms (n=38) due to mixed phenotypes in individuals; "*" according to Goh et al. [27].
593

References

1. Ketting, R.F. The many faces of RNAi. *Dev Cell* **20**, 148-61 (2011).
2. Weick, E.M. & Miska, E.A. piRNAs: from biogenesis to function. *Development* **141**, 3458-71 (2014).
3. Luteijn, M.J. & Ketting, R.F. PIWI-interacting RNAs: from generation to transgenerational epigenetics. *Nat Rev Genet* **14**, 523-34 (2013).
4. Czech, B. & Hannon, G.J. One Loop to Rule Them All: The Ping-Pong Cycle and piRNA-Guided Silencing. *Trends Biochem Sci* **41**, 324-337 (2016).
5. Luteijn, M.J. et al. Extremely stable Piwi-induced gene silencing in Caenorhabditis elegans. *EMBO J* **31**, 3422-30 (2012).
6. Sakakibara, K. & Siomi, M.C. The PIWI-Interacting RNA Molecular Pathway: Insights From Cultured Silkworm Germline Cells. *Bioessays* **40**(2018).
7. Rojas-Rios, P. & Simonelig, M. piRNAs and PIWI proteins: regulators of gene expression in development and stem cells. *Development* **145**(2018).
8. Das, P.P. et al. Piwi and piRNAs act upstream of an endogenous siRNA pathway to suppress Tc3 transposon mobility in the Caenorhabditis elegans germline. *Mol Cell* **31**, 79-90 (2008).
9. Batista, P.J. et al. PRG-1 and 21U-RNAs interact to form the piRNA complex required for fertility in C. elegans. *Mol Cell* **31**, 67-78 (2008).
10. Wang, G. & Reinke, V. A C. elegans Piwi, PRG-1, regulates 21U-RNAs during spermatogenesis. *Curr Biol* **18**, 861-7 (2008).
11. Bagijn, M.P. et al. Function, targets, and evolution of Caenorhabditis elegans piRNAs. *Science* **337**, 574-578 (2012).
12. Lee, H.C. et al. C. elegans piRNAs mediate the genome-wide surveillance of germline transcripts. *Cell* **150**, 78-87 (2012).
13. Simon, M. et al. Reduced insulin/IGF-1 signaling restores germ cell immortality to Caenorhabditis elegans Piwi mutants. *Cell Rep* **7**, 762-73 (2014).
14. Heestand, B., Simon, M., Frenk, S., Titov, D. & Ahmed, S. Transgenerational Sterility of Piwi Mutants Represents a Dynamic Form of Adult Reproductive Diapause. *Cell Rep* **23**, 156-171 (2018).
15. de Albuquerque, B.F., Placentino, M. & Ketting, R.F. Maternal piRNAs Are Essential for Germline Development following De Novo Establishment of Endo-siRNAs in Caenorhabditis elegans. *Dev Cell* **34**, 448-56 (2015).
16. Phillips, C.M., Brown, K.C., Montgomery, B.E., Ruvkun, G. & Montgomery, T.A. piRNAs and piRNA-Dependent siRNAs Protect Conserved and Essential C. elegans Genes from Misrouting into the RNAi Pathway. *Dev Cell* **34**, 457-65 (2015).
17. Ashe, A. et al. piRNAs can trigger a multigenerational epigenetic memory in the germline of C. elegans. *Cell* **150**, 88-99 (2012).
18. Shirayama, M. et al. piRNAs initiate an epigenetic memory of nonself RNA in the C. elegans germline. *Cell* **150**, 65-77 (2012).
19. Buckley, B.A. et al. A nuclear Argonaute promotes multigenerational epigenetic inheritance and germline immortality. *Nature* **489**, 447-51 (2012).
20. Ruby, J.G. et al. Large-scale sequencing reveals 21U-RNAs and additional microRNAs and endogenous siRNAs in C. elegans. *Cell* **127**, 1193-207 (2006).
21. Kasper, D.M., Wang, G., Gardner, K.E., Johnstone, T.G. & Reinke, V. The C. elegans SNAPc component SNPC-4 coats piRNA domains and is globally required for piRNA abundance. *Dev Cell* **31**, 145-58 (2014).

22. Weick, E.M. et al. PRDE-1 is a nuclear factor essential for the biogenesis of Ruby
motif-dependent piRNAs in C. elegans. *Genes Dev* **28**, 783-96 (2014).
23. Beltran, T. et al. Evolutionary analysis im[licates RNA polymersase II pausing and
chromatin structure in nematode piRNA biogenesis. *BioRxiv*
https://doi.org/10.1101/281360 (2018).
24. Blumenthal, T. Trans-splicing and operons in C. elegans. *WormBook*, 1-11 (2012).
25. Gu, W. et al. CapSeq and CIP-TAP identify Pol II start sites and reveal capped small
RNAs as C. elegans piRNA precursors. *Cell* **151**, 1488-500 (2012).
26. de Albuquerque, B.F. et al. PID-1 is a novel factor that operates during 21U-RNA
biogenesis in Caenorhabditis elegans. *Genes Dev* **28**, 683-8 (2014).
27. Goh, W.S. et al. A genome-wide RNAi screen identifies factors required for distinct
stages of C. elegans piRNA biogenesis. *Genes Dev* **28**, 797-807 (2014).
28. Tang, W., Tu, S., Lee, H.C., Weng, Z. & Mello, C.C. The RNase PARN-1 Trims piRNA 3'
Ends to Promote Transcriptome Surveillance in C. elegans. *Cell* **164**, 974-84 (2016).
29. Kamminga, L.M. et al. Differential impact of the HEN1 homolog HENN-1 on 21U and
26G RNAs in the germline of Caenorhabditis elegans. *PLoS Genet* **8**, e1002702 (2012).
30. Billi, A.C. et al. The Caenorhabditis elegans HEN1 ortholog, HENN-1, methylates and
stabilizes select subclasses of germline small RNAs. *PLoS Genet* **8**, e1002617 (2012).
31. Montgomery, T.A. et al. PIWI associated siRNAs and piRNAs specifically require the
Caenorhabditis elegans HEN1 ortholog henn-1. *PLoS Genet* **8**, e1002616 (2012).
32. Cecere, G., Zheng, G.X., Mansisidor, A.R., Klymko, K.E. & Grishok, A. Promoters
recognized by forkhead proteins exist for individual 21U-RNAs. *Mol Cell* **47**, 734-45
(2012).
33. Sugiyama, T. et al. Enhancer of Rudimentary Cooperates with Conserved RNA-
Processing Factors to Promote Meiotic mRNA Decay and Facultative
Heterochromatin Assembly. *Mol Cell* **61**, 747-759 (2016).
34. Jankowska-Anyszka, M. et al. Multiple isoforms of eukaryotic protein synthesis
initiation factor 4E in Caenorhabditis elegans can distinguish between mono- and
trimethylated mRNA cap structures. *J Biol Chem* **273**, 10538-42 (1998).
35. Arai, R. et al. Crystal structure of an enhancer of rudimentary homolog (ERH) at 2.1
Angstroms resolution. *Protein Sci* **14**, 1888-93 (2005).
36. Frokjaer-Jensen, C. et al. Random and targeted transgene insertion in Caenorhabditis
elegans using a modified Mos1 transposon. *Nat Methods* **11**, 529-34 (2014).
37. Sengupta, M.S. et al. ifet-1 is a broad-scale translational repressor required for
normal P granule formation in C. elegans. *J Cell Sci* **126**, 850-9 (2013).
38. Cauchi, R.J. SMN and Gemins: 'we are family' ... or are we?: insights into the
partnership between Gemins and the spinal muscular atrophy disease protein SMN.
Bioessays **32**, 1077-89 (2010).
39. Mader, S., Lee, H., Pause, A. & Sonenberg, N. The translation initiation factor eIF-4E
binds to a common motif shared by the translation factor eIF-4 gamma and the
translational repressors 4E-binding proteins. *Mol Cell Biol* **15**, 4990-7 (1995).
40. Gruner, S. et al. The Structures of eIF4E-eIF4G Complexes Reveal an Extended
Interface to Regulate Translation Initiation. *Mol Cell* **64**, 467-479 (2016).
41. Minasaki, R., Puoti, A. & Streit, A. The DEAD-box protein MEL-46 is required in the
germ line of the nematode Caenorhabditis elegans. *BMC Dev Biol* **9**, 35 (2009).
42. Keiper, B.D. et al. Functional characterization of five eIF4E isoforms in Caenorhabditis
elegans. *J Biol Chem* **275**, 10590-6 (2000).

43. Maris, C., Dominguez, C. & Allain, F.H. The RNA recognition motif, a plastic RNA-
binding platform to regulate post-transcriptional gene expression. *FEBS J* **272**, 2118-
31 (2005).
44. Schirle, N.T. & MacRae, I.J. The crystal structure of human Argonaute2. *Science* **336**,
1037-40 (2012).
45. Siomi, M.C., Mannen, T. & Siomi, H. How does the royal family of Tudor rule the
PIWI-interacting RNA pathway? *Genes Dev* **24**, 636-46 (2010).
46. Philippe, L. et al. An in vivo genetic screen for genes involved in spliced leader trans-
splicing indicates a crucial role for continuous de novo spliced leader RNP assembly.
Nucleic Acids Res **45**, 8474-8483 (2017).
47. Xu, C. et al. Structural insights into Gemin5-guided selection of pre-snRNAs for snRNP
assembly. *Genes Dev* **30**, 2376-2390 (2016).
48. Zanetti, S., Meola, M., Bochud, A. & Puoti, A. Role of the C. elegans U2 snRNP protein
MOG-2 in sex determination, meiosis, and splice site selection. *Dev Biol* **354**, 232-41
(2011).
49. Ferguson, K.C., Heid, P.J. & Rothman, J.H. The SL1 trans-spliced leader RNA performs
an essential embryonic function in Caenorhabditis elegans that can also be supplied
by SL2 RNA. *Genes Dev* **10**, 1543-56 (1996).
50. Edgar, R.C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res* **32**, 1792-7 (2004).
51. Robert, X. & Gouet, P. Deciphering key features in protein structures with the new
ENDscript server. *Nucleic Acids Res* **42**, W320-4 (2014).
710

Gene	Allele	Mutation Type	21U RNA Presence	Terminal Phenotypes	Ratio (%)
ife-3	*xf101*	Start Codon Loss	n.d.	Mog/Mel	n.d.
	xf102	Frameshift	-	Mog/Mel	n.d.
	RNAi	n.a.	n.d.	Mog/Mel	47/53#
pid-3	*tm2417*	Frameshift	--	Mel	100
	xf149	Frameshift	n.d.	Mel	100
	xf151	Inframe Deletion	n.d.	Mel	100
	xf153	Frameshift	n.d.	Mel	100
tofu-6	*it20*	Nonsense	n.d.	Mel	100
	yt2	Nonsense	n.d.	Mel	100
	RNAi	n.a.	--*	Mel	n.d.
tost-1	*xf191*	Frameshift	n.d.	Viable	n.d.
	xf194	Start Codon Loss	+	Mel	100
	xf196	Splice Site Loss	n.d.	(TS) Mel	100 (25ºC)
pid-1	*xf14*	Missense	--	n.d.	n.d.
	xf35	Frameshift	--	Mog	<1
	xf36	Frameshift	--	n.d.	n.d.
erh-2	*xf168*	Stop Codon Loss	--	Mel	100

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6