Increased association between Epstein-Barr virus EBNA2 from type 2 strains and the

transcriptional repressor BS69 restricts B cell growth

Rajesh Ponnusamy[1], Ritika Khatri[1], Paulo B. Correia[2], Erika Mancini[1], Paul J. Farrell[2] and

Michelle J. West[1*]

1. School of Life Sciences, University of Sussex, Falmer, Brighton, BN1 9QG, UK.

2. Section of Virology, Imperial College London, Norfolk Place, London W2 1PG, UK.

* To whom correspondence should be addressed. Tel: +44 (0)1273 678404; Fax: +44 (0)1273

877586; Email: m.j.west@sussex.ac.uk

Abstract

Natural variation separates Epstein-Barr virus (EBV) into type 1 and type 2 strains. Type 2 EBV is less transforming *in vitro* due to sequence differences in the EBV transcription factor EBNA2. This correlates with reduced activation of the EBV oncogene LMP1 and some cell genes. Transcriptional activation by type 1 EBNA2 can be suppressed through the binding of two PXLXP motifs in its transactivation domain (TAD) to the dimeric coiled-coil MYND domain (CC-MYND) of the BS69 repressor protein (ZMYND11). We identified a third conserved PXLXP motif in type 2 EBNA2. We found that type 2 EBNA2 peptides containing this motif bound $BS69_{CC-MYND}$ efficiently and that the type 2 $EBNA2_{TAD}$ bound an additional $BS69_{CC-MYND}$ molecule. Full-length type 2 EBNA2 also bound BS69 more efficiently in pull-down assays. Molecular weight analysis and low-resolution structures obtained using small-angle X-ray scattering showed that three $BS69_{CC-MYND}$ dimers bound two molecules of type 2 $EBNA2_{TAD}$, in line with the dimeric state of full-length EBNA2 *in vivo*. Importantly, mutation of the third BS69 binding motif in type 2 EBNA2 improved B-cell growth maintenance. Our data indicate that increased association with BS69 restricts growth promotion by EBNA2 and may contribute to reduced B-cell transformation by type 2 EBV.

Author summary

Epstein-Barr virus (EBV) drives the development of many human cancers worldwide including specific types of lymphoma and carcinoma. EBV infects B lymphocytes and immortalises them, thus contributing to lymphoma development. The virus promotes B lymphocyte growth and survival by altering the level at which hundreds of genes are expressed. The EBV protein EBNA2 is known to activate many growth-promoting genes. Natural variation in the sequence of EBNA2 defines the two main EBV strains: type 1 and type 2. Type 2 strains immortalise B lymphocytes less efficiency and activate some growth genes poorly, although the mechanism of this difference is unclear. We now show that sequence variation in type 2 EBNA2 creates a third site of interaction for the repressor protein (BS69, ZMYND11). We have characterised the complex formed between type 2 EBNA2 and BS69 and show that three dimers of BS69 form a bridged complex with two molecules of type 2 EBNA2. We demonstrate that mutation of the additional BS69 interaction site in type 2 EBNA2 improves its growth-promoting function. Our results therefore provide a molecular explanation for the different B lymphocyte growth promoting activities of type 1 and type 2 EBV. This aids our understanding of immortalisation by EBV.

Introduction

Epstein-Barr virus (EBV) is a ubiquitous γ-herpesvirus that immortalises human B lymphocytes to establish a lifelong persistent infection that is usually harmless. Delayed primary EBV infection can however give rise to infectious mononucleosis. EBV is also associated with the development of malignancies that include Burkitt's (BL), Hodgkin's, diffuse large B cell and post-transplant lymphoma and nasopharyngeal or gastric carcinoma. EBV expresses nine latent proteins in *in vitro* infected lymphoblastoid cell lines (LCLs), including 6 Epstein-Barr nuclear antigens (EBNA1, 2, 3A, 3B, 3C and leader protein) and 3 latent membrane proteins (LMP1, 2A, 2B). The EBNA2 transcription factor is one of five of these latent genes essential for B cell transformation (1). EBNA2 functions as the master regulator of EBV latent gene transcription and activates numerous cell genes that control B cell growth and survival (2). It cannot however bind to DNA directly and hijacks cell DNA binding proteins e.g. RBPJ (RBPJκ, CBF1) and EBF1 to target viral and cell gene regulatory elements (2, 3). Although EBNA2 binding sites are close to gene promoters in the viral genome, in the B cell genome they are mostly found at enhancer elements and EBNA2 has been shown to promote enhancer-promoter interactions (4-6). EBNA2 activates transcription through interactions between its acidic transactivation domain (TAD) and histone acetyl transferases, ATP-dependent remodellers and components of the preinitiation complex (7-13).

EBV genome sequences worldwide separate into two main strains (type 1 and type 2) based on differences in the EBNA2 and EBNA3A, 3B and 3C genes (14-18). Type 2 strains are less efficient at immortalising resting B cells *in vitro* than type 1 strains (19). This phenotype is determined by sequence variation in EBNA2 since complementation of an EBNA2 defective virus with type 1 EBNA2 but not type 2 EBNA2 supports efficient primary B cell

immortalisation (1). Consistent with its reduced primary B cell transforming function, type 2

EBNA2 cannot complement loss of type 1 EBNA2 function to maintain the growth of

lymphoblastoid cell lines (20). Amino acids responsible for the differences in B cell growth

maintenance between type 1 and type 2 EBNA2 were mapped to the C-terminal region of

EBNA2 (20). Surprisingly, a single amino acid (aspartate 442 of type 1 EBNA2) in the TAD

appears to be a key determinant of B cell growth maintenance by type 1 EBNA2. Replacing

the serine that occurs at the equivalent position in type 2 EBNA2 (amino acid 409 of type 2

EBNA2) with aspartate (mutant S442D) confers efficient growth maintenance function (21).

Type 2 EBNA2 has reduced ability to activate expression of the EBV oncogene LMP1 and a

small number of cellular genes e.g. CXCR7 (22). These differences in gene activation could

underlie the reduced B cell growth maintenance and transforming function of type 2 EBNA2,

although the mechanism involved and the role played by the single aspartate residue is unclear.

Part of the mechanism may involve (or result in) reduced binding of type 2 EBNA2 to the

LMP1 promoter and cell gene regulatory elements (21). EBNA2 binding sites at genes

activated less efficiently by type 2 EBNA2 are enriched for composite binding motifs for ETS

and IRF transcription factors (ETS and IRF composite element; EICE), implicating ETS/IRF

family members in the gene specificity of the observed effects (21).

Despite a clear deficiency in the immortalising and B cell growth maintenance properties of

type 2 EBNA2 *in vitro*, no specific differences in disease association have been reported to

date for type 1 and type 2 EBV. Interestingly, although the outgrowth of immortalised cells is

less efficient and much slower in primary B cell cultures infected with type 2 EBV (19, 20),

the LCLs that are eventually established from type 2 viruses proliferate at similar rates to type

1 LCLs. Type 1 and 2 LCLs also show equivalent expression of LMP1 and CXCR7 (20). Over

extended periods of time, it is therefore possible to select for immortalised cells infected with

type 2 EBV that have the required levels of expression of these genes to support their long term

proliferation. *In vivo* other factors may create an environment that helps support B cell

immortalisation by type 2 EBV.

New research also suggests that type 2 EBV may use alternative approaches to persist *in vivo*.

Type 2 EBV has the unique capacity to infect T cells in culture and is detected in T cells from

healthy infants from Kenya, indicating that T cell infection may form part of a natural type 2

EBV infection (23, 24). Recent work also showed that a type 2 EBV strain was able to infect

both B cells and T cells in humanised mice (25). Mice infected with type 2 EBV developed

tumours that resembled the diffuse large B-cell lymphomas that also developed in mice infected

with type 1 EBV, confirming the tumorigenic potential of a persistent type 2 EBV infection,

once established (25).

BS69 (ZMYND11) is a multi-domain chromatin-associated repressor protein that suppresses

transcription elongation, regulates pre-mRNA processing and has tumour suppressor function

(26, 27). The BS69 gene undergoes chromosomal translocation in minimally differentiated

myeloid leukaemia leading to the expression of a BS69-MBTD1 fusion protein (28). BS69

contains three histone reader domains in its N-terminal region; a plant homeodomain, a

bromodomain and a PWWP domain. The tandemly-arranged bromodomain and PWWP

domain bind to histone H3 or the variant histone H3.3 when trimethylated on lysine K36 (27,

29). BS69 also contains a coiled-coil (CC) dimerisation domain adjacent to a MYND domain

in its C terminus. BS69 binds to a number of chromatin modifying enzymes (BRG1, HDAC1,

EZH2) and transcription factors (adenovirus E1a, c-Myb, ETS2, E2F6, and the Myc-associated

MGA protein) and inhibits transcription factor activation function (30-34). The BS69 MYND

domain binds to E1a and MGA through a PXLXP motif (34).

BS69 has also been shown to interact with the TAD of type 1 EBNA2 through two PXLXP

motifs and to restrict EBNA2 transcriptional activation function (34, 35). The structure of the

dimeric CC-MYND domain of BS69 bound to two peptides encompassing sequences from one

of the EBNA2 PXLXP motifs (motif 1) has been solved (35). Based on this three-dimensional

structure, a BS69 dimer was predicted to interact with the two adjacent PXLXP motifs in type

1 EBNA2. Interestingly the amino acid implicated in the type-specific differences in growth

maintenance observed for type 1 and type 2 EBNA2 (amino acid 442 in type 1 EBNA2) (21)

lies immediately adjacent to the second BS69 binding motif.

We set out to determine whether sequence differences between type 1 and type 2 EBNA2 affect

BS69 binding. We hypothesised that type 2 EBNA2 would show increased binding to BS69

and that this would impair its gene activation and growth maintenance function. We initially

examined the impact of type-specific differences in EBNA2 amino acid 442 on BS69 binding.

We also identified a third PXLXP BS69 binding motif in type 2 EBNA2, so we examined

whether the presence of this extra motif resulted in the interaction of additional molecules of

BS69 with type 2 EBNA2. We found that amino acid 442 did not affect BS69 binding, but

influenced the conformation of the TAD, potentially affecting binding of other transcriptional

regulators. We also demonstrated that the third PXLXP motif in type 2 EBNA2 was responsible

for the binding of an additional BS69 dimer. Importantly, mutation of the third PXLXP BS69

binding motif in full length type 2 EBNA2 restored B cell growth maintenance function

indicating that increased BS69 binding is responsible for impaired type 2 EBNA2 function.

Results

Sequence differences near type 2 EBNA2 BS69 binding motif 2 do not increase BS69 binding.

Previous studies demonstrated that differences in a single amino acid in the TAD between type

1 and type 2 EBNA2 determined the ability of EBNA2 to maintain the growth of an EBV-

infected LCL (21). This amino acid (located at position 442 in the EBNA2 sequence from the

prototypical type 1 B95-8 strain of EBV) is conserved as aspartate in type 1 strains and as

serine (at the corresponding position of 409) in type 2 strains. In type 1 EBNA2, aspartate 442

is located immediately adjacent to a previously identified binding motif for the cell

transcriptional repressor BS69 (motif 2) that fits the PXLXP consensus (PILFP$_{437-441}$)(34). In

the TAD of type 2 EBNA2, the PXLXP motif is conserved (PFLFP$_{404-408}$) and is flanked by

serine 409 (Figure 1A).

We hypothesised that the impaired gene activation and growth maintenance properties of type

2 EBNA2 may be the result of increased binding to BS69 as a result of the aspartate to serine

amino acid difference adjacent to BS69 binding motif 2. We therefore tested whether a BS69

binding motif 2 peptide from type 2 EBNA2 showed enhanced binding to BS69 compared to

a motif 2 peptide from type 1 EBNA2. We used isothermal titration calorimetry (ITC) to

determine the affinity of peptide binding to the C-terminal region of BS69 (amino acid 480-

602) comprising the CC-MYND domain that we expressed and purified from *E.coli* (Figure

1B and 1C). In contrast to our hypothesis, we found that the type 2 EBNA2 motif 2 peptide

bound to BS69$_{CC-MYND}$ with reduced affinity (K_D=176 μM) compared to the corresponding

peptide from type 1 EBNA 2 (K_D=47.7 μM) (Figure 1B and 1C and Supplementary Table S1).

The affinity of binding of the type 1 EBNA 2 motif 2 peptide to BS69$_{CC-MYND}$ was very similar

to the previously reported K_D of 35 μM (35). The difference in binding between type 1 and

type 2 EBNA2 motif 2 peptides could be influenced by both the aspartate to serine change and

differences in two other amino acids present in the sequence (Figure 1B and 1C).

An additional BS69 PXLXP binding motif previously identified in type 1 EBNA2 (motif 1)

located N-terminal to motif 2 is also present in type 2 EBNA 2 (Figure 1A). The $BS69_{CC\text{-}MYND}$

dimer binds a type 1 EBNA2 polypeptide containing both motif 1 and motif 2 with high affinity

and the structure of BS69 dimer could accommodate binding to both motifs simultaneously

(35). We therefore tested whether sequence differences in type 2 EBNA2 (including the

aspartate to serine change) affected the binding of a region of EBNA2 containing both motif 1

and motif 2 to BS69. Type 1 $EBNA2_{381\text{-}445}$ and type 2 $EBNA2_{348\text{-}412}$ were expressed and

purified from *E.coli* and their interaction with $BS69_{CC\text{-}MYND}$ examined using ITC. Consistent

with previous reports (35) we found that type 1 $EBNA2_{381\text{-}445}$ bound to $BS69_{CC\text{-}MYND}$ with high

affinity (K_D=0.95 μM) likely due to the high avidity of interaction with two binding sites

(Figure 1D). In the context of this larger region of EBNA2 we found very little difference in

the affinity of type 2 EBNA2 binding to $BS69_{CC\text{-}MYND}$ (K_D=1.21 μM) (Figure 1E). In addition

to measuring binding affinities, ITC data can also be used to calculate binding stoichiometry

(n) which can be visualised as the molar ratio at the mid (inflection) point of the sigmoidal

binding curve. We titrated EBNA2 polypeptides into a cell containing $BS69_{CC\text{-}MYND,}$ so the

stoichiometry values we obtained indicate the molar ratio at which the EBNA2 polypeptide

saturates the available sites in $BS69_{CC\text{-}MYND}$ monomers. Consistent with the presence of two

BS69 binding sites in the EBNA2 polypeptides, we obtained n values of 0.42 and 0.33 for type

1 and type 2 EBNA2: $BS69_{CC\text{-}MYND}$ binding, respectively (Figure 1D and 1E). These

approximate to the expected molar ratio of 0.5 taking into consideration some margin of error

in n value determination by ITC, which is heavily influenced by the accuracy of protein

concentrations and the proportion of 'active' protein in the sample.

We conclude that the aspartate 442 to serine amino acid difference between type 1 and type 2

EBNA2 does not affect the binding of BS69 to the TAD of type 2 EBNA2 in these assays. Our

data also indicate that additional sequence differences in and around BS69 binding motifs 1

and 2 in type 2 EBNA2 do not influence the binding of the $BS69_{CC-MYND}$ dimer to this region

of the protein.

Type 2 EBNA2 contains a third BS69 binding site

During the course of our study we also identified a third potential BS69 binding site in the type

2 EBNA2 TAD (Figure 2A). In type 2 EBNA2, a sequence that is an exact match to the PXLXP

BS69 consensus binding motif is present C-terminal to motif 2 ($PTLEP_{414-418}$). In type 1

EBNA2 the corresponding region has an isoleucine in place of the leucine residue ($PSIDP_{447-}$

$_{451}$). To determine whether these regions of EBNA2 also interact with BS69, we performed

ITC experiments using type 1 and type 2 EBNA2 peptides (Figure 2B and 2C). We were not

able to detect any binding of the type 1 EBNA2 peptide encompassing this region (T1

$EBNA2_{445-455}$) to $BS69_{CC-MYND}$, underscoring the importance of the central leucine in the

PXLXP motif for the BS69 interaction (Figure 2B). In contrast, a peptide from the

corresponding region of type 2 EBNA2 (T2 $EBNA2_{412-422}$) interacted with BS69 with a

K_D=219 µM (Figure 2C). The affinity of interaction with this new motif (that we named motif

3) is weaker than the interaction we observed for type 1 or type 2 EBNA2 motif 2 (Figure 1).

To determine the impact of motif 3 on the interaction of type 2 EBNA2 with $BS69_{CC-MYND}$ in

the presence of the two other BS69 binding motifs, we expressed and purified a larger type 2

EBNA2 polypeptide containing motif 1, 2 and 3 (T2 $EBNA2_{348-422}$) for use in ITC. For

comparison, we also analysed the binding of the corresponding larger region of type 1 EBNA2

(T1 EBNA2$_{381-455}$). We found that inclusion of the additional C-terminal amino acids had little

impact on the affinity of binding of type 1 EBNA2 to BS69$_{CC-MYND}$ or the stoichiometry of

binding (compare Figure 2D and Figure 1D) (Supplementary Table S2). In contrast, for type 2

EBNA2, we observed a change in the stoichiometry of binding from 0.33 when motif 1 and 2

were present (T2 EBNA2$_{348-412)}$ to 0.15 when motif 1, 2 and 3 were present (T2 EBNA2$_{348-422}$)

(compare Figure 2E and 1E). This is consistent with the presence of an additional BS69 binding

site and indicates the interaction of T2 EBNA2$_{348-422}$ with an additional BS69$_{CC-MYND}$ molecule.

Perhaps surprisingly, we did not observe an increase in the affinity of binding of the longer

type 2 EBNA2 polypeptide to BS69$_{CC-MYND}$ (compare Figure 2E and 1E). Nonetheless the

recruitment of more BS69 to type 2 EBNA2 could be physiologically relevant for the function

of type 2 EBNA2 as a transcriptional activator. To confirm that the observed change in binding

stoichiometry was due to the presence of motif 3 in the type 2 EBNA2 polypeptide, we analysed

the binding of T2 EBNA2$_{348-422}$ with motif 3 mutated from PTLEP to ATAEA (T2 EBNA2$_{348-}$

$_{422}$ motif 3 mt). We found that mutation of motif 3 altered the stoichiometry of binding to

BS69$_{CC-MYND}$ from 0.15 to 0.30, consistent with the loss of a BS69 binding motif (Figure 2F).

This is similar to the value obtained for the type 2 EBNA2 polypeptide containing only motif

1 and motif 2 (T2 EBNA2$_{348-412}$)(Figure 1E). We conclude that type 2 EBNA2 contains an

additional binding site for BS69 that is not present in type 1 EBNA2.

Three BS69 CC-MYND dimers bridge two molecules of type 2 EBNA2

To further examine whether type 2 EBNA2 could form higher-order complexes with the BS69

CC-MYND domain that are larger than type 1 EBNA2, we examined the properties of BS69-

EBNA2 complexes using size exclusion chromatography (SEC). Consistent with complex

formation, when pre-incubated with BS69$_{CC-MYND,}$ both T1 EBNA2$_{381-455}$ and T2 EBNA2$_{348-422}$

polypeptides migrated through the size exclusion column faster and eluted at a lower elution

volume compared to the migration of each component individually (Figure 3A). In line with

the binding of additional BS69$_{CC-MYND}$ molecules to T2 EBNA2$_{348-422}$ and the formation of

higher molecular weight complexes, we found that type 2 EBNA2 complexes eluted at a lower

volume than type 1 EBNA2 complexes (Figure 3A). SDS-PAGE of SEC column fractions

confirmed the presence of BS69$_{CC-MYND}$ and EBNA2 in the higher molecular weight complexes

(Figure 3B). Note that both type 1 and type 2 EBNA2 polypeptides migrate anomalously on

SDS-PAGE gels and not at their predicted molecular weights (MW) of 7.9 and 8.1 kDa

respectively, likely due to their high proline content (Figure 3B). They are however pure and

resolve as single species on gel filtration columns (Figure 3A).

Because migration in SEC is influenced by both size and shape and BS69$_{CC-MYND}$ has an

elongated structure due to the CC domain, we were unable to determine the MW of BS69-

EBNA2 complexes accurately using SEC. In order to obtain more accurate MW information

that would allow us to determine the number of molecules of BS69$_{CC-MYND}$ and EBNA2 present

in type 1 and type 2 complexes, we used SEC with multi-angle light scattering (SEC-MALS)

(Table 1). SEC-MALS gave MWs for T1 EBNA2$_{381-455}$ and T2 EBNA2$_{348-422}$ that matched the

theoretical MW of their monomeric forms and gave a MW for BS69$_{CC-MYND}$ consistent with its

dimeric state (Table 1). For the T1 EBNA2$_{381-455}$-BS69$_{CC-MYND}$ complex, SEC-MALS gave a

MW of 62.3 kDa. Given that there are two binding sites for BS69 in the T1 EBNA2$_{381-455}$

polypeptide, this figure most closely matches the MW of a complex containing two type 1

EBNA2 polypeptides and two BS69$_{CC-MYND}$ dimers (theoretical MW of 76.7 kDa) rather than

a single type 1 EBNA2 polypeptide with a one dimer of EBNA2 BS69$_{CC-MYND}$ (theoretical MW

of 38.3 kDa) (Table 1). For the BS69$_{CC-MYND}$-T2 EBNA2$_{348-422}$ complex, SEC-MALs gave a

MW of 135 kDa consistent with the larger complex size observed in SEC (Table 1 and Figure

3A). Given the presence of three BS69 binding motifs in type 2 EBNA2, this MW most closely

matches that of a complex containing three $BS69_{CC-MYND}$ dimers and two type 1 EBNA2

polypeptides (theoretical MW of 107.5 kDa) (Table 1).

Because of the discrepancies in the theoretical and experimentally determined MWs for

$BS69_{CC-MYND}$-EBNA2 complexes, we also used small-angle-X-ray scattering (SAXS) to obtain

information on the shape and size of these complexes in solution. Initially we used SEC-SAXS

to analyse each polypeptide individually. We used a Kratky representation to visualize features

of the scattering profiles obtained for T1 $EBNA2_{381-45,}$ T2 $EBNA2_{348-422}$ and $BS69_{CC-MYND}$

individually to identify the folding state of the polypeptides in solution. The absence of a bell-

shaped curve with a well-defined maximum for both EBNA2 polypeptides indicates that they

are natively unfolded in solution (Supplementary Figure S1). The bell-shaped curve obtained

for the $BS69_{CC-MYND}$ dimer indicates that it is folded in solution as expected from the crystal

structure (35). Three-dimensional models were created for the individual polypeptides by *ab

initio* shape determination. For $BS69_{CC-MYND}$ a solution structure consistent with the coiled-

coil dimer structure determined by X-ray crystallography was obtained (35)(Supplementary

Figure S2). For the EBNA2 polypeptides, solution structures consistent with flexible unfolded

peptide chains were obtained (Supplementary Figure S2). SAXS analysis of $BS69_{CC-MYND}$ pre-

mixed with either type 1 or type 2 EBNA2 polypeptides gave a larger Porod volume (directly

related to MW) compared to the individual proteins, consistent with complex formation

(Supplementary Table S3). An *ab initio* dummy atom model was generated for the type 1

EBNA2-$BS69_{cc-MYND}$ complex and this fitted well to the experimental SAXS data (χ^2 of 1.4)

(Figure 4A). The three-dimensional model generated by *ab initio* shape determination for the

type 1 EBNA2-$BS69_{cc-MYND}$ complex indicated that the complex has a large elongated shape

with a volume of 166 nm^3 and a maximum dimension (Dmax) of 138 Å (Figure 4B). This three-

dimensional model could accomodate two BS69$_{CC-MYND}$ dimer structures that were manually

docked into the SAXS envelope. The additional space at the bottom of the model was allocated

to the model solution structures of two type 1 EBNA2 polypeptides (Figure 4B). This docked

structural model for the type 1 EBNA2-BS69$_{CC-MYND}$ complex was then fitted to the

experimental scattering data and gave a reasonable χ^2 value of 2.56. For comparison a structural

model where only one BS69$_{CC-MYND}$ dimer and a single type 1 EBNA2 polypeptide were

docked into the SAXS envelope was created but this alternative model gave a worse fit to the

experimental data (Supplementary Figure S3). An *ab initio* dummy atom model was then

generated for the type 2 EBNA2-BS69$_{CC-MYND}$ complex and this fitted well to the experimental

SAXS data (χ^2 of 1.0) (Figure 4C). The three-dimensional model created by *ab initio* shape

determination for the type 2 EBNA2 complex had a larger volume (239 nm^3) and maximum

dimension (145 Å) than the type 1 EBNA2 complex model (Figure 4D). The type 2 EBNA2

model could accommodate the docking of three BS69$_{CC-MYND}$ dimer structures along with two

type 2 EBNA2 polypeptides and this structural model gave a good fit to the experimental data

(χ^2 of 1.44) (Figure 4C and D). In comparison a docked model containing two BS69$_{CC-MYND}$

dimers and a single type 2 EBNA2 polypeptide gave a worse fit to the experimental data

(Supplementary Figure S3).

Taken together our data indicate that BS69 forms higher order complexes with EBNA2 that

involve the interaction of each MYND domain of the BS69 dimer with binding sites in two

separate EBNA2 molecules. Rather than an *in vitro* artefact, this intermolecular 'bridging'

interaction is consistent with the fact that EBNA2 forms dimers *in vivo*. Although the N-

terminal regions of EBNA2 that mediate dimerisation (36) are absent in the EBNA2

polypeptides we examined in our interaction studies, our data indicate that BS69 may have the

capacity to stabilise or enhance dimerisation between two EBNA2 molecules held together

through their N-termini. Importantly, using multiple independent techniques, we also

demonstrate that type 2 EBNA 2 interacts with an additional BS69$_{CC-MYND}$ dimer.

The serine to aspartate change in type 2 EBNA2 alters its binding characteristics

During the course of our ITC experiments we observed that binding data obtained using the

longer EBNA2 polypeptides (T1 EBNA2$_{381-455}$ and T2 EBNA2$_{348-422}$) showed some deviation

from curves fitted using the single binding event ('one set of sites') model (where binding to

multiple sites cannot be detected as separate heat change events) (Figure 2D and 2E). This

suggested that the mode of binding of these polypeptides to BS69$_{CC-MYND}$ could involve more

than one distinguishable binding event. To determine whether this was the case, we performed

ITC experiments using an increased number of smaller injections of the EBNA2 polypeptide

to obtain more data points for curve fitting (Figure 5). For T1 EBNA2$_{381-455}$ the binding data

did not fit well to curves generated using an alternative two binding event ('two sets of sites')

model (χ^2/degrees of freedom=0.56) (Figure 5A and Supplementary Table S2). This indicates

that the deviation of T1 EBNA2$_{381-455}$-BS69$_{CC-MYND}$ binding data from fitted curves at low

molar ratios was unlikely to be the result of a separate binding event (Figure 2D and 5A). In

contrast, for T2 EBNA2$_{348-422}$ the binding profiles obtained fitted well to curves generated using

the two binding event model (χ^2/degrees of freedom=0.23) (Figure 5B and Supplementary

Table S2). This enabled the affinity of the two separate binding events to be determined, which

were both in the nanomolar range (K$_{D1}$=0.009 μM and K$_{D2}$=0.091 μM). These data indicate

that this region of type 2 EBNA2 may adopt a different conformation to type 1 EBNA2 when

binding to BS69$_{CC-MYND.}$

In further experiments we addressed the impact of changing the serine at position 409 in type

2 EBNA2 to the aspartate present at the equivalent position (aspartate 442) in type 1 EBNA2on

BS69 binding in the context of the longer type 2 EBNA2 polypeptide containing three BS69

binding sites. To do this we expressed and purified a type 2 EBNA2 polypeptide with an S409D

mutation (T2 EBNA2$_{348-422}$ SD mutant). Interestingly, we found that the SD substitution

enhanced the detection of the second binding event on interaction with BS69$_{CC-MYND}$ (Figure

5C). The second binding event for T2 EBNA2$_{348-422}$ SD was associated with a larger change in

enthalpy (ΔH -7.65 kcal/mol) than the second binding event detected for T2 EBNA2$_{348-422}$ (ΔH

-1.59 kcal/mol) (Supplementary Table S2). Importantly we found that the affinities of the two

binding events remained largely unaffected (Figure 5B and C), consistent with our earlier

observations that the presence of serine at position 409 does not affect the ability of a type 2

polypeptide containing motif 1 and motif 2 to bind BS69 (Figure 1). To determine whether the

impact of the SD mutation was dependent on the presence of BS69 binding motif 3, we also

produced a polypeptide containing the SD and motif 3 mutation (EBNA2$_{348-422}$ SD + m3 mt).

We found that two binding events were still clearly detectable on interaction of this double

mutant type 2 EBNA2 polypeptide with BS69$_{CC-MYND}$ and that the enthalpy change of the

second binding event was similar to that of the single SD mutant (ΔH -9.62 kcal/mol)

(Supplementary Table S2) indicating that the impact of the SD change is still evident. The

affinity of the second binding event (K_{D2}) was however reduced by approximately 2-fold for

EBNA2$_{348-422}$ SD + m3 mt compared to the EBNA2$_{348-422}$ SD mutant (Figure 5C and D). These

data indicate that motif 3 contributes to the second binding event. We conclude that the SD

mutation previously shown to enhance the growth maintenance properties of type 2 EBNA2

(21) does not affect BS69 binding but likely alters the conformation of the type 2 EBNA2 TAD.

This may therefore impact on the binding of other transcriptional regulators that influence type

2 EBNA2 function.

Full-length type 2 EBNA2 binds BS69$_{CC-MYND}$ *more efficiently in pull-down assays*

To confirm our *in vitro* observations that a type 2 EBNA2 polypeptide binds an additional

BS69 dimer, we examined the interaction of BS69$_{CC-MYND}$ with full-length EBNA2 proteins

stably expressed in B cells. Lysates from cells expressing type 1 or type 2 EBNA2 or the type

2 SD mutant were incubated with recombinant GST-BS69$_{CC-MYND}$ immobilised on glutathione

beads for increasing times and the amount of EBNA2 precipitated determined by Western

Blotting (Figure 6). Consistent with the presence of an additional BS69 binding site in type 2

EBNA2, we found that GST-BS69$_{CC-MYND}$ pulled down type 2 EBNA2 more efficiently than

type 1 EBNA2 at short incubation times (Figure 6). In agreement with our *in vitro* observations

using the type 2 EBNA2 SD mutant, we found that this protein interacted with BS69$_{CC-MYND}$

with the same efficiency as type 2 EBNA2 (Figure 6). After 30 minutes incubation, GST-

BS69$_{CC-MYND}$ became saturated with EBNA2 and differences in association were no longer

evident. A control GST fusion protein (GST-Rab11) did not precipitate EBNA2, confirming

the specificity of the interactions. These data therefore confirm the increased association of

BS69$_{CC-MYND}$ with type 2 EBNA2.

Mutation of BS69 binding motif 3 in type 2 EBNA2 increases its growth maintenance function

To determine whether the presence of the additional BS69 binding motif in type 2 EBNA2

(motif 3) had functional consequences for the activity of type 2 EBNA2, we examined the

ability of a type 2 EBNA2 motif 3 mutant to maintain B cell growth. We utilised a previously

described assay using an EBV-infected LCL (EREB2.5) in which the activity of a type 1

estrogen receptor-EBNA2 fusion protein can be switched off by estrogen withdrawal (37). Loss

of EBNA2 activity leads to growth arrest, but transfection of a stably-maintained plasmid

expressing type 1 EBNA2 into these cells supports their survival (20). In contrast, the

expression of type 2 EBNA2 cannot maintain the growth of these cells (20). We found that

mutation of BS69 binding motif 3 produced a type 2 EBNA2 protein that was able to support

the recovery of these cells from the loss of type 1 EBNA2 activity, with cells recovering well

2-4 weeks following estrogen withdrawal (Figure 7A). The type 2 EBNA2 motif 3 mutant

behaved similarly to the type 2 EBNA2 SD mutant that was previously shown to support B cell

growth in this assay (21). We also examined the ability of the SD and motif 3 double mutant

in this assay and found that it showed a slightly increased ability to support B cells growth

(Figure 7A). In our hands, expression of type 1 EBNA2 supported initial growth in this assay

better than any type 2 mutants, with the mutants supporting growth recovery from 2 weeks

(Figure 7A). We confirmed that all EBNA2 proteins were expressed at similar levels (Figure

7B). We conclude that the presence of the additional BS69 binding motif in type 2 EBNA2

impairs the ability of type 2 EBNA2 to maintain B cell growth.

A BS69 isoform containing the MYND domain is expressed in type 1 and type 2 EBV-infected

cells.

BS69 functions as a negative regulator of EBNA2 transcription activity in reporter assays (34,

35), but previous studies have reported that BS69 expression is downregulated on infection of

resting B cells by EBV and is low in the resulting immortalised LCLs (35). Transcriptional

repression of BS69 by EBNA2 was implicated in BS69 downregulation indicating that EBNA2

may act to restrict expression of its own negative regulator (35). The cell lines examined in this

previous study all harboured type 1 EBV or type 1 EBNA2, so we next addressed whether

BS69 was expressed at similar levels in cells infected with type 1 and type 2 EBV. We

examined BS69 protein levels in type 1 and type 2 LCLs using an anti-BS69 antibody raised

against a region within the MYND domain of BS69. We found that BS69 was expressed at

similar levels in type 1 and type 2 LCLs, but surprisingly levels in LCLs were similar to those

in an EBV negative B cell line (AK31) (Figure 8A). We expanded our analysis to include

additional EBV negative B cell lines (BJAB and DG75), EBV infected cell lines displaying the

EBNA1 only latency I pattern of EBV gene expression (Akata and Mutu I), an additional type

1 LCL (IB4) and a BL cell line expressing all EBV latent proteins including EBNA2 (Mutu

III) (both latency III cell lines) (Figure 8B). We found no correlation between BS69 expression

and EBV infection or EBNA2 expression (Figure 8B). BS69 did not therefore appear to be

downregulated as a result of EBNA2 expression. We also examined BS69 expression over the

course of a primary B cell infection and found that BS69 was not downregulated as previously

reported (Figure 8C).

We therefore explored the possibility that we were detecting a different isoform of BS69.

Alternative splicing has been reported to give rise to different BS69 isoforms and four have

been experimentally verified (33) (Figure 9A). The canonical isoform (isoform 1, UniProt

identifier: Q15326-1) contains 15 exons and encodes a protein of 71 kD (602 amino acids).

Isoform 2 (Q15326-2) lacks amino acids 93-146 encoding the PHD domain (exon 4) and

encodes a protein of 64.4 kD. Isoform 3 (Q15326-3) lacks amino acids 563-602 encoding the

MYND domain (exon 15) and has a unique C-terminus encoded by an extended exon 14

sequence. Isoform 3 encodes a protein of 66.6 kD. Isoform 4 (Q15326-3) lacks exon 4 and

exon 15 (and thus both the PHD and MYND domains) and encodes a protein of 60 kD. These

isoforms were previously described as full length (FL), ΔPHD, ΔMD and ΔPHD, ΔMD

respectively (33), but the exon numbering used in this previous study differed. The BS69

protein detected in Figure 8A and B has a molecular weight of approximately 64 kD consistent

with that expected for isoform 2. This was the only protein detected by this antibody (against

the MYND domain), indicating that isoform 1 was not expressed in the cell lines examined.

Since the antibody we used would not detect BS69 isoforms 3 and 4, we could not exclude the

possibility that one or more of these isoforms was also expressed and that an alternative BS69

isoform was detected previously (35). In line with this possibility, we noted that the QPCR

analysis carried out by Harter *et al* used primers located in exon 4, which is absent from isoform

2.

No detail was provided on the anti-BS69 antibody used previously (35) and we were not able

to find another antibody that detected isoform 3 and 4 in Western blotting. We therefore took

a non-quantitative PCR approach to screen for different BS69 isoforms using cDNA prepared

from LCLs and from B cells during a primary EBV infection. PCR using a forward primer in

exon 3 and a reverse primer in exon 13 amplified two products indicating the presence of at

least two different isoforms, one containing exon 4 (1139 bps) and one lacking exon 4 (977

bps) (Figure 9B). This would be consistent with the presence of isoform 3 (which contains

exon 4) and isoform 2 (which lacks exon 4 and was detected by Western blotting (Figure 8)).

PCR products were sequenced to confirm their identity (data not shown). However, since

isoform 4 also lacks exon 4, this PCR analysis cannot rule out the additional presence of

isoform 4. Since in isoforms 3 and 4 exon 15 is replaced by a short unique 3' sequence from

exon 14, we designed reverse PCR primers in this unique 3' region. PCR using these primers

amplified only one product of 1578 bps consistent with presence of exon 4 and the unique 3'

region (isoform 3) (Figure 9B). The identity of this PCR product was again confirmed by

sequencing (data not shown). Importantly, we did not detect a smaller product (1416 bps) that

would indicate the presence of isoform 4 (lacks exon 4 and exon 15). Our data therefore

indicate that LCLs infected with either type 1 or type 2 EBV express both isoform 2 and

isoform 3 of BS69.

To quantitatively examine whether either BS69 isoform 2 or isoform 3 were downregulated on

EBV infection and in cells expressing EBNA2 as previously described (35), we used QPCR to

analyse BS69 mRNA expression in primary B cells infected by EBV and in a panel of EBV

negative and positive B cell lines. QPCR using primers that spanned exon 14 and exon 15

(present in isoform 1 and 2) detected variable levels of BS69 across the cell lines examined,

with no obvious correlation with EBV positivity or EBNA2 expression (present in latency III

EBV infected cell lines). This is consistent with the variability in BS69 protein expression

detected in Western blot analysis of isoform 2 expression (Figure 8). Although, in one

experiment (#2) primary B cells expressed high levels of BS69 isoform 2 mRNA that were

reduced on EBV infection, the second primary infection experiment did not reproduce this

observation. In fact, primary infection #2 was the same infection analysed by Western blotting

in Figure 8C so this change in BS69 RNA expression did not result in decreased expression of

BS69 isoform 2 protein. It is most likely therefore that BS69 isoform 2 expression varies in an

EBV and EBNA2 independent manner. Analysis of BS69 mRNA expression using QPCR

primers that would specifically amplify BS69 isoforms containing the long form of exon 14

(isoforms 3 and 4) also detected variable expression of BS69 that did not correlate with EBV

positivity or EBNA2 expression indicating that isoform 3 expression is also EBV independent

(Figure 9D).

We conclude that B cells infected with type 1 or type 2 EBV do not consistently display reduced

expression of any detectable isoform of BS69 compared to uninfected B cells. Since BS69

isoform 2 contains the MYND domain that binds EBNA2 (that is absent in isoform 3), the

continued expression of isoform 2 in EBV infected B cells would be expected to restrict the

gene activation function of EBNA2.

Inhibition of BS69 function increases EBNA2 transactivation activity

To determine whether inhibition of BS69 function increased EBNA2 transactivation function,

we carried out EBNA2 transactivation assays in an EBV negative B cell line (BJAB) in which

we overexpressed isoform 3 of BS69 lacking the MYND domain (ΔMYND) (but containing

the coiled-coil dimerisation domain). This form of BS69 has been proposed to act as a dominant

negative inhibitor of the MYND-domain dependent functions of BS69 (33). We performed

transactivation assays using EBNA2-GAL4-DNA binding domain (DBD) fusion proteins and

a Firefly luciferase reporter plasmid containing a synthetic promoter with 4 GAL4 binding

sites. Plasmids expressing GAL4-DBD fusion proteins containing regions of type 1 EBNA2

(334-487) and type 2 EBNA2 (301-454) encompassing all BS69 binding motifs were

transfected into BJAB cells in the presence or absence of plasmids expressing either full length

BS69 (isoform 1) or isoform 3 (ΔMYND). Consistent with previous reports, we found that

overexpression of full length BS69 inhibited transactivation by type 1 EBNA2 (34, 35) (Figure

10). BS69 also inhibited transactivation by type 2 EBNA2 (Figure 10). Consistent with its

function as a dominant negative inhibitor, we found that expression of BS69 ΔMYND

increased transactivation by both type 1 and type 2 EBNA2 (Figure 10). To determine whether

this was a non-specific or EBNA2-dependent effect, we expressed BS69 ΔMYND in the

absence of any GAL4-DBD-EBNA2 expressing constructs. In the absence of EBNA2 fusion

protein expression, BS69 ΔMYND had no effect on the activity of the GAL4 reporter (Figure

10). These data therefore demonstrate that inhibition of the MYND-domain dependent function

of BS69 in B cells relieves repression of EBNA2 transactivation. These data support our

hypothesis that the expression of MYND-domain containing BS69 isoforms in B cells impedes

EBNA2 gene activation function.

Taken together our *in vitro* and cell-based assays suggest that during initial B cell infection the

increased association of BS69 with type 2 EBNA2 may impede key gene activation events that

are required for the efficient outgrowth of immortalised cell lines.

Discussion

Type 2 EBV strains have reduced B cell transformation capacity and type 2 EBNA2 activates some viral and cell genes less efficiently than type 1 EBNA2, a feature that may underlie the impaired transformation phenotype. We have identified an additional binding site for the transcriptional repressor BS69 in the EBNA2 protein encoded by type 2 strains of EBV and show that mutation of this additional binding site improves the B cell growth maintenance properties of type 2 EBNA2. Our data therefore implicate increased BS69 association in the impaired function of type 2 EBNA2.

Type 2 EBV transforms resting B cells more slowly and results in the outgrowth of less immortalised cell clones than type 1 EBV (1, 19). Although early type 2 EBV transformants show reduced cell growth, the immortalised LCLs that eventually arise from a type 2 EBV infection grow similarly to those infected with type 1 EBV. Type 2 LCLs also maintain similar levels of expression of key EBNA2 target genes (20). This indicates that the impaired function of type 2 EBNA2 restricts an early stage in the B cell transformation process *in vitro*. Indeed two EBNA2 target genes that are only weakly activated by type 2 EBNA2 compared to type 1 EBNA2 (22), the viral oncogene LMP1 and the cell gene CXCR7, display slower and weaker induction during primary infection with type 2 EBV (20). Although it was previously reported that BS69 is downregulated on EBV infection, we found that there is continued expression of BS69 isoform 2 in EBV-infected cells. Since this isoform contains the MYND domain that mediates the BS69-EBNA2 interaction the expression of BS69 would be expected to restrict EBNA2 activation function. Consistent with this prediction we found that the expression of a dominant negative form of BS69 (isoform 3) lacking the MYND domain enhances EBNA2 activation function in B cells. Our data are consistent with a model where BS69 acts as a

restriction factor for both type 1 and type EBNA2 but the association of type 2 EBNA2 with

more molecules of the BS69 repressor protein further restricts the activation of growth and

survival genes important in early transformation. Why a small number of specific genes are

activated less well by type 2 EBNA2 is as yet not fully clear, but sequences resembling EICEs

(bound by ETS and IRF transcription factors) are found at EBNA2 binding sites in the LMP1

promoter and binding sites closest to the cell genes that show reduced activation by type 2

EBNA2. Interestingly BS69 binding to PXLXP motifs in ETS2 has been shown to inhibit its

transactivation activity (32). Since the ETS family member PU.1 is known to bind to the

putative EICE in the LMP1 promoter and plays a role in LMP1 promoter activation, it is

possible that type 2 EBNA2 functions less well in the context of PU.1 binding sites.

Interestingly, PU.1 also contains PXLXP motifs that would be predicted to bind BS69, so

enhanced tertiary complex formation between type 2 EBNA2, BS69 and PU.1 at the regulatory

elements of specific genes may function to stabilise BS69 binding and further restrict gene

activation by type 2 EBNA2.

Our data also provide important new molecular information on the nature of the complexes

formed between EBNA2 and BS69 that may applicable to the way BS69 interacts with other

cellular and viral transcription factors via its MYND domain. Single and multiple PXLXP

BS69 binding motifs have been identified in the cell and viral binding partners of BS69, but

the elucidation of the structure of the dimeric coiled-coil-MYND domain of BS69 led to a

model that proposed that a BS69 dimer bound to the two adjacent PXLXP motifs (motif 1 and

motif 2) in the same type 1 EBNA2 molecule (35). However, although the authors found that

BS69 bound with increased affinity when two PXLXP motifs were present in EBNA2

polypeptides, the three-dimensional structure obtained comprised a $BS69_{CC-MYND}$ dimer bound

to two separate type 1 EBNA2 motif 1 peptides. Although the binding of motif 1 and motif 2

could be accommodated in the $BS69_{CC-MYND}$ structure if the intervening 52 amino acids were

looped out, the formation of this complex has not been formally demonstrated (35). Our SEC-

MALS and SAXS analysis provides the first evidence that the $BS69_{CC-MYND}$ dimer

preferentially forms an intermolecular bridge between PXLXP motifs located on different

EBNA2 molecules. This mode of binding is consistent with the fact that EBNA2 is a dimeric

protein, with dimerisation mediated by the N-terminal END domain comprising amino acids

1-58 (36). Additional self-associating regions have also been mapped elsewhere in EBNA2

and include amino acids 97–121 and 122-344 (38, 39), although no molecular information is

available on how these regions may contribute to dimerisation. Our data indicate that BS69

binding to sites in the C-terminal transactivation domain may contribute to the formation or

stabilisation of EBNA2 dimers. Interestingly, although SEC analysis clearly demonstrated

complex formation between both type 1 and type 2 EBNA 2 polypeptides and $BS69_{CC-MYND,}$

the elution profiles of both complexes were broad. The type 1 EBNA2-$BS69_{CC-MYND}$ elution

profile had a clear shoulder indicating the presence of smaller MW complexes (Figure 3A).

This would explain why the average MW determined by SEC-MALs was smaller than expected

for a complex that contained two molecules of type 1 EBNA2 and two $BS69_{CC-MYND}$ dimers.

It is possible that in solution *in vitro* there is a mixed population of dimeric type 1 EBNA2 and

monomeric type 1 EBNA2 complexes (where a single EBNA2 polypeptide is bound by one

$BS69_{CC-MYND}$ dimer as previously proposed). We were not able to investigate this further using

SAXS as this 'shoulder' was not clearly defined, so SAXS analysis for both type 1 and type 2

EBNA2-BS69 complexes focused on the major elution peak of the large complex. Given that

full length EBNA2 expressed in EBV-infected cells is a dimer, complexes involving two

EBNA2 molecules are more likely to be physiologically relevant.

Surprisingly, in our GAL4-EBNA2 fusion protein assays we did not see weaker transactivation

by the type 2 EBNA2 fusion protein compared to the type 1 EBNA2 fusion protein as reported

previously (21). We used a longer region of EBNA2 compared to this previous study that

encompassed all three BS69 binding sites for type 2 EBNA2 and the corresponding region of

type 2 EBNA2 (with only two functional BS69 binding sites). Previously GAL4-EBNA2

fusion protein constructs were used that expressed a type 1 EBNA2 protein containing only

BS69 binding motif 2 or the corresponding region of type 2 EBNA2 that contained BS69

binding motif 2 and 3 (21). It is not completely clear why the increased association of BS69

with type 2 EBNA2 is not associated with weaker transactivation in our assays in the context

of a longer region of EBNA2, but it could point to the importance of the dimerisation that

occurs in the context of the full-length protein in the assembly of larger BS69-EBNA2

complexes.

When considering the nature of assembly of BS69-EBNA2 complexes, it is likely that binding

to motif 1 (which in type 1 EBNA2 has the highest affinity for $BS69_{CC\text{-}MYND}$) would drive the

initial interaction between EBNA2 and BS69 and binding to motif 1 probably constitutes the

first binding event that can be distinguished in our ITC analysis using an increased number of

injections. For type 2 EBNA2, since both motif 2 and 3 bind BS69 with similar affinity, binding

to both of these motifs probably occurs with similar kinetics and is detectable as a single second

binding event by ITC. Given the fact that $BS69_{CC\text{-}MYND}$ dimers are predicted in the solution

structure of the BS69-EBNA2 complex to be located side by side along a dimeric EBNA2

molecule, it is possible that interactions between BS69 coiled-coil dimers play a role in

stabilising the oligomeric complex.

Our initial interest in examining type-specific binding of EBNA2 to BS69 centred around the

influence of a serine residue in the TAD of type 2 EBNA2 that plays a key role in restricting B

cell growth maintenance by type 2 EBNA2 (21). Although this residue is located immediately

adjacent to BS69 binding motif 2 in type 2 EBNA2, we found that it did not increase BS69

binding (as might have been expected) when binding was compared to the corresponding

region of type 1 EBNA2 where there is an aspartate residue in its place. It does not appear

therefore that the influence of serine 409 on growth maintenance is mediated through

alterations in BS69 binding affinity. Our ITC analysis however did find that a serine to

aspartate change at this position in type 2 EBNA2 altered the nature of BS69 binding indicating

that it may induce a conformational change in this region of EBNA2. This could result in

differences in the binding of other transcription regulators to the type 2 EBNA2 TAD compared

to the type 1 EBNA2 TAD. Possibilities could include increased binding of a repressor or co-

repressor to the type 2 EBNA2 TAD or decreased binding of an activator or co-activator.

BS69 may have a wider role in regulating B cell transformation and the growth of EBV-

infected cells in addition to its modulation of EBNA2 transactivation. BS69 localised to the

cell membrane has also been implicated as an adaptor in signalling mediated by the EBV

oncogene LMP1. The MYND domain of BS69 was reported to bridge an interaction between

the carboxy terminal cytoplasmic domain of LMP1 and the TRAF6 signalling protein to

activate the JNK signalling pathway (40). Conversely, BS69 has also been implicated as a

negative regulator of LMP1-mediated NF-κB signalling by decreasing the association between

C-terminal activation region (CTAR) 2 of LMP1 and the signalling adaptor TRADD (41) and

by binding to CTAR1 and bringing in the negative regulator of NF-κB signalling, TRAF3 (42).

Although further work appears to be required to fully understand the role of BS69 in LMP1

signalling and the relative proportions of nuclear and membrane-associated BS69, it is possible

that BS69 is a key modulator of growth promoting events in EBV-infected cells. In this context,

our work now sheds new light on how transformation by type 2 strains of EBV may be

specifically curbed as a result of sequence variation that results in the creation of an additional

binding site for BS69.

Materials and Methods

Cell lines

All cell lines were passaged twice weekly in RPMI 1640 media (Invitrogen) supplemented with 10% Fetal Bovine serum (Gibco), 1 U/ml penicillin G, 1 µg/ml streptomycin sulphate and 292 µg/ml L-glutamine at 37°C in 5% CO_2. DG75 (43) and AK31 (44) are EBV negative BL cell lines and BJAB (45) is an EBV negative B cell lymphoma line. Akata (46) and Mutu I are EBV positive latency I BL cell lines and Mutu III is a cell line derived from Mutu I cells that drifted in culture to express all EBV latent proteins (latency III) (47). All LCLs also display the latency III pattern of EBV gene expression and were described previously (48); IB4, spLCL, LCL3, C2 + Obaji, JAC-B2, BM + Akata LCLs are infected with type 1 EBV and C2 + BL16, WEI-B1, Jijoye and AFB1 LCLs are infected with type 2 EBV. The ER-EB 2.5 LCL, expressing a conditionally active oestrogen receptor (ER)-EBNA2 fusion protein, was provided by Prof B. Kempkes and was cultured in the presence of β-estradiol (37). The Daudi:pHEBo-MT:E2T1, Daudi:pHEBo-MT:E2T2 and Daudi:pHEBo-MT:E2T2 S442D cell lines were generated and cultured as described previously (21). B cell infection samples were described previously (49).

Plasmids

Constructs expressing N-terminal 6 x histidine tagged EBNA2 polypeptides were generated using the Sequence and Ligation Independent Cloning (SLIC) technique using type 1 EBNA2 (B95-8), type 2 EBNA2 (AG876) and type 2 EBNA2 SD (serine to aspartate at position 409) pSG5 expression plasmids as templates to amplify the regions of interest. DNA was PCR amplified using primers containing 20-30 bp of additional sequence from the regions 5' and 3' to the multiple cloning site of pET47b+. pET47b+ was digested using SmaI and HindIII and the PCR product and double-digested vector were then partially digested using the 3' to 5' exonuclease activity of T4 DNA Polymerase in the absence of dNTPs to generate long

complementary 5' overhangs. The PCR products and pET47b+ were then annealed on ice. The

ligated DNA fragments obtained contain four nicks that are repaired by *E. coli* after

transformation. For type 1 EBNA2, constructs encoded amino acids 381-445 or 381-455 to

generate pET47b+ T1 EBNA2$_{381-445}$ and pET47b+ T1 EBNA2$_{381-455.}$ Type 2 EBNA2 constructs

encoded amino acids 348-412 or 348-422 to generate pET47b+ T2 EBNA2$_{348-412,}$ pET47b+ T2

EBNA2$_{348-422}$ and pET47b+ T2 EBNA2$_{348-422}$ SD.

The BS69 CC-MYND domain (amino acids 480-602) was amplified from pCI-BS69

containing the full length human BS69 sequence (gift from Dr Stéphane Ansieau) and cloned

using SLIC into the SmaI and HindIII sites of pET49b+ to generate a construct expressing an

N-terminal GST-6x Histidine tag BS69$_{CC-MYND}$ fusion protein.

To create GAL4-DNA-binding domain-EBNA2 TAD fusion protein expressing constructs

pBlueScript plasmids carrying EBNA2 sequences were used as the template to PCR amplify

the type 1 EBNA2 TAD (amino acids 426-463) and the type 2 EBNA2 TAD (amino acids 334-

487) using *Taq* DNA polymerase. Primers contained *Bam*HI or *Not*I restriction sites at their 5'

ends. PCR products were first cloned into pCR2.1 using the TA cloning kit (Invitrogen)

according to the manufacturer's instructions. The pCR2.1 vector carrying the cloned PCR

product was then digested with *Bam*HI and *Not*I and the EBNA2 TAD fragment then cloned

into the *Bam*HI and *Not*I sites of pcDNA3.1-GAL4-DBD.

Site-directed mutagenesis

Reverse PCR with the Q5® Site-Directed Mutagenesis kit (NEB) was used to introduce

mutations into BS69 binding motif 3 in the pET47b+ T2 EBNA2$_{348-422}$ construct. This resulted

in a change from PTLEP to ATAEA and generated pET47b+ T2 EBNA2$_{348-422}$ motif 3 mt. The

same primers were used to mutate motif 3 in the context of the SD mutation to generate

pET47b+ T2 EBNA2$_{348-422}$ SD + motif 3 mt.

Growth maintenance assay

The EREB2.5 growth assay was performed as described previously (20). Briefly, 5 μg of OriP-

p294 plasmids expressing type 1 EBNA2, type 2 EBNA2 or type 2 EBNA2 mutants were

transfected into 5×10^6 EREB 2.5 cells resuspended in 110 μl of buffer T using Neon

transfection with 1 pulse of 1300 V for 30 msec. Following transfection, cell suspensions were

added to 2 ml of media supplemented with 10% FBS and antibiotics but without β-oestradiol

and incubated overnight in 12-well plates. The following day each transfected sample was

made up to 10 mls with media and divided into 5 x 2 ml aliquots in a 12-well plate. Samples

were harvested for cell counting and protein analysis at time points up to 4 weeks.

Reporter assays

Cells were diluted 1:3 in fresh culture medium one day before transfection. 2×10^6 BJAB cells

were used for each individual transfection. Cells were pelleted by centrifugation at 335*g* for 5

minutes at 4ºC and washed twice with pre-warmed PBS. Cells were resuspended in 100 μl of

Neon resuspension solution R (Invitrogen). Cell suspensions were then mixed with plasmid

DNA (2-12 μg in TE buffer). Cells were co-transfected with 300 ng of either type 1 GAL4-

DBD:EBNA2 (aa 334-487) or type 2 GAL4-DBD:EBNA2 (301-454) constructs, 500 ng of

pFRLuc (Agilent technologies), 10 ng of pRL-CMV (Promega) and 1 μg of BS69 (pCI-BS69)

or BS69 ΔMYND (pCI-BS69-ΔMYND) expressing plasmids (gift from Dr Stéphane Ansieau).

The DNA and cell mixture was transferred to a 100 μl Neon transfection pipette tip

(Invitrogen). Cells were electroporated using Neon transfection protocol 14 (1200 V of pulse

voltage, 20 ms of pulse width and 2 pulse number) and then transferred into 2 ml of pre-warmed

media in a 6-well plate and incubated at 37ºC for 24 hr.

Cell pellets were then lysed using 100 μl of 1X Passive Lysis buffer (Promega). Two freeze-

thaw cycles were performed to achieve efficient lysis (20 sec on dry ice and thawing at room

temperature for 2 min). Cell debris was removed by centrifugation at 25,000g, for 1 min at 4ºC

and the clear supernatant was then transferred to a fresh tube. 20 μl of lysate was assayed for

firefly and Renilla luciferase activity using 20 μl of each dual luciferase assay kit reagent

(Promega) and a microplate luminometer (LUMIstar Omega, BMG Labtech).

Recombinant protein production

pGEX4T1-BS69 (kindly provided by Dr Stéphane Ansieau) was used to express a GST-BS69

fusion protein containing amino acids 452-602 of BS69 encompassing the CC-MYND

domain (numbered according to the canonical isoform) (34). pGEX4T1-RAB11B expressing

GST-tagged RAB11B (gift from Prof Gill Elliott) was used to produce a negative control

protein for the GST pull-down assays.

For production of BS69$_{CC-MYND}$ and EBNA2 polypeptides, the relevant plasmids were

transformed into the Rosetta 2 (DE3) pLysS E. Coli strain and protein expression induced by

adding 0.4 mM of isopropyl βD-1-thiogalactopyranoside (IPTG) to 3 litre cultures at an

OD$_{600nm}$ of 0.6. The bacteria were then grown at 20°C overnight before harvesting for protein

purification. Cell pellets from a 3 litre induced culture were lysed for 30 minutes on ice with

constant stirring in 100 ml of lysis buffer (25 mM Tris-HCL pH 7.5, 500 mM NaCl, 5%

Glycerol). The lysis buffer was supplemented with 0.25 mg/ml lysozyme, 2 mM MgCl$_2$, 1 mM

TCEP (tris(2-carboxyethyl)phosphine), two protease-inhibitor complete tablets (Roche) and

DNase and 0.2 mg/ml of DNase I. Lysates were then sonicated at 37% amplitude for 5 minutes

with 10 seconds pulses using a Vibra-cell sonicator (SONICS). The cell debris was pelleted at

15000 rpm for 45 minutes at 4°C (Biofuge Stratos, Heraeus). Beads from 3 ml of HisPur™

Cobalt Resin slurry (Thermo Scientific) were added to the cleared lysate along with 2 mM

imidazole and the sample incubated for 1 hour and 30 minutes at 4°C with rolling. Samples

were decanted into a centrifuge column (Thermo Scientific Pierce) and washed with buffer

(25mM Tris-HCl, 500mM NaCl and 1mM TCEP, 2 mM of imidazole, pH 7.5). The protein

was eluted from the beads using buffer containing increasing concentrations of imidazole (5

mM, 10 mM, 20 mM, 50 mM, 100 mM, 200 mM, 300 mM and 500 mM). Fraction samples

were analysed by SDS-PAGE and the fractions containing recombinant protein were pooled

and incubated with 3C protease (200 μl of 2 mg/ml) in the presence of 2 mM DTT overnight

at 4°C to cleave the Histidine tag. His-tag cleaved proteins were then separated from the 3C

protease by passing the protein sample through a GSTrap HP column (Amersham) using a

peristaltic pump to capture the GST-tagged 3C protease. Untagged recombinant proteins were

then concentrated and injected into a HiLoad™ 16/600 Superdex™ 75g Column (GE

Healthcare) pre-equilibrated in 25 mM Tris-HCL, 200 mM NaCl, 1 mM TCEP, pH 7.5 purified

at 0.5 ml/min. Protein fractions containing purified protein were then pooled, concentrated and

stored at -80°C until required. Approximately 1 mg of purified EBNA2 polypeptides or 4 mg

of BS69$_{CC-MYND}$ was obtained from 1 litre of culture.

Isothermal titration calorimetry

Four commercially synthesized peptides (Peptide Synthetics) were used for ITC. These

included BS69 binding motif 2 of type 1 EBNA2 (435-445) or type 2 EBNA2 (402-412) and

putative BS69 binding motif 3 of type 1 EBNA2 (445-455) or type 2 EBNA2 (412-422).

Frozen protein was quickly thawed using running water and dialysed overnight at 4°C using

Slide-A-Lyzer® MINI Dialysis Units (Thermo Scientific) against ITC buffer (20mM Tris-HCl,

100mM NaCl and 1mM TCEP , pH 7.5). The next day, protein samples were centrifuged at

13000 rpm for 10 minutes at 4ºC and the concentration was determined by NanoDrop

spectroscopy (NanoDrop Technologies) with their respective molecular weights and extinction

coefficients. EBNA2 peptides (1mM) and polypeptides (type 1, 0.3 mM and type 2, 0.6 mM)

were titrated against $BS69_{CC-MYND}$ (0.1mM) at 25°C using a MicroCalTM iTC200 instrument

(Malvern). For peptides, 13 x 3.0 µl injections were used for titration. For EBNA2 polypeptides

19 x 2.0 µl or 29 x 1.3 µl injections were used for titration. ITC data were corrected for non-

specific heat and analysed using MicroCal Origin® 7.0. The experiments were performed in

triplicate alongside a control experiment with no $BS69_{CC-MYND}$ (buffer only in the cell). All

polypeptides were used within 24 hours of dialysis into ITC buffer.

Size Exclusion Chromatography

An S200 10/300 GL gel filtration column (GE Healthcare) was equilibrated with buffer

containing 20mM Tris-HCl, 100mM NaCl and 1mM TCEP, pH 7.5. Individual EBNA2 or

$BS69_{CC-MYND}$ polypeptides or complexes were applied to the column and analysed at a flow

rate of 0.5ml/min. The eluted fractions were then analysed by SDS-PAGE and Quick

Coomassie staining.

Size Exclusion Chromatography-multi-angle light scattering

EBNA2-BS69 complexes were prepared by pre-incubating proteins in a 1:3 molar ratio for at

least 30 mins at 4°C. Purified samples (45 µl) at a concentration of 5 mg/ml were loaded onto

a Shodex KW403-4F column at 25°C pre-equilibrated in 20 mM Tris-HCl, 100mM NaCl and

1mM TCEP, pH 7.5. Elution fractions were monitored using a DAWN HELEOS II MALS

detector followed by a refractive index detector Optilab T-rEX (Wyatt Technology). Molecular

masses of each individual peak were determined using ASTRA 6 software (Wyatt

Technology). For normalization of the light scattering and data quality, BSA was used as a

calibration standard.

Small-angle X-ray scattering

Synchrotron radiation X-ray scattering data from solutions of individual proteins or complexes

prepared as for SEC-MALS were collected on beamline B21 at Diamond Light Source (Didcot,

United Kingdom), with an inline HPLC system. X-ray scattering patterns were recorded on a

Pilatus detector after injection of 45 μl of protein sample (5-10 mg/ml) in a Superdex 200

3.2/300 column equilibrated in 20mM Tris-HCl, 100mM NaCl, 2% Sucrose and 1mM TCEP,

pH 7.5. Samples were analysed at 20°C using a flow-rate of 0.25 ml/min. Initial data processing

(background subtraction and radius of gyration Rg calculation) was performed using ScÅtter

(v3.0 by Robert P. Rambo; Diamond Light Source). *Ab initio* beads model for the complex

were prepared using DAMMIF (50). 23 independent dummy atom models were obtained by

running the program in 'slow' mode. DAMAVER was then used to align and average the

models (51). The *ab initio* generated beads models were refined using DAMMIN and compared

to the experimental scattering data to derive χ^2 values (52). The goodness-of-fit χ^2 values for the

docked structure compared to the experimental scattering data were determined with FoXS

(53).

GST pull-down assays

Nuclear extracts were prepared from control or EBNA2 expressing Daudi cell lines. EBNA2

expression in Daudi:pHEBo-MT:EBNA2 cells was induced with 5 μM $CdCl_2$ for 24 hours. At

least $4x10^7$ cells were then harvested and resuspended in 1 ml of buffer A (10 mM HEPES pH

7.9, 1.5 mM MgCl2, 10 mM KCl, 0.5 mM 1,4-dithiothreitol (DTT) (Sigma), 1 mM PMSF

(Sigma) and 1x complete protease inhibitor cocktail (Roche)). Cells were pelleted by

centrifugation at 1000*g* for 5 min at 4ºC and lysed in 100 μl of buffer A supplemented with

0.1% (v/v) NP-40 and incubated on ice for 5 min. Cell lysates were centrifuged at 2700*g* for

30 sec at 4ºC and the nuclei resuspended in 50 μl of buffer B (20 mM HEPES pH 7.9, 420 mM

NaCl, 1.5 mM MgCl2, 0.2 mM EDTA, 1 mM PMSF, 25% (v/v) glycerol, 1 mM DTT and 1x

complete protease inhibitor cocktail) at 4ºC for 20 min with rotation. Samples were finally

centrifuged at 11,600g for 10 min at 4ºC and the supernatants/nuclear extracts were transferred

to fresh eppendorf tubes and the protein concentration was determined before storage at -80ºC.

Lysates containing GST-tagged BS69$_{CC-MYND}$ or GST-RAB11B were prepared from 100 ml

cultures of E. Coli BL21 (DE3). Transformed cells were cultured at 30ºC until they reached an

OD$_{600nm}$ of 0.6 and protein expression was induced at 25ºC with 0.5 mM IPTG for 3-4 h. Cells

were pelleted at 2,800g for 20 min at 4ºC and then resuspended in 10 ml of Lysis buffer (20

mM Tris-Cl pH 8.0, 150 mM NaCl and 1 mM DTT) supplemented with 120 μl of lysozyme

(10 mg/ml) and lysates incubated on ice for 20 min. Lysates were sonicated at high speed for

3 x 15 sec pulses in ice water using an Ultrasonic XL2020 Processor (Heat Systems) and cell

debris pelleted at 17,900g for 30 min at 4ºC. Lysates were stored at -80ºC until required.

For pull-down assays, 50 μl of 50% Glutathione-Sepharose 4B Bead slurry (GE Healthcare)

was washed three times in ice-cold binding buffer (20 mM Tris-Cl pH 8.0, 150 mM NaCl, 1

mM DTT and 0.1 mg/ml BSA). Beads were pelleted by centrifugation at 25,000g for 1 min

and 100 μl of bacterial lysate containing the GST-tagged protein was incubated with the washed

beads for 1 h at 4ºC with rotation. Glutathione-Sepharose Beads bound to the GST-tagged

protein were then washed with ice-cold binding buffer six times and pelleted by centrifugation

at 25,000g for 1 min. Loaded GST- BS69$_{CC-MYND}$ beads were then incubated with nuclear

extracts containing EBNA2 at 4ºC for different times (5, 10 and 30 minutes). Loaded GST-

RAB11B beads were incubated with lysates for 30 minutes. Beads were then washed six times

with ice-cold binding buffer and pelleted by centrifugation at 25,000g for 1 min. Beads were

then resuspended in 25 μl of 2x SDS sample buffer (120 mM Tris-Cl pH 6.8, 4% (w/v) SDS,

2% (v/v) β-mercaptoethanol, 20% (v/v) glycerol and 0.01% (w/v) bromophenol blue) and

incubated at 95ºC for 5 min and analysed for EBNA2 levels by SDS-PAGE and Western

blotting.

SDS-PAGE and Western blotting

SDS-PAGE and Western blotting was carried out as described previously (54, 55) using the

anti-EBNA2 monoclonal antibody PE2 (gift from Prof M. Rowe) anti-actin 1/5000 (A-2066,

Sigma) and anti-BS69 1/1000 (ab190890, Abcam). Western blot visualisation and signal

quantification was carried out using a Li-COR Imager. Gels were stained using Quick

Coomassie stain (Generon Ltd).

PCR and QPCR

RNA was extracted from cells using Trireagent (Sigma), further purified using the RNeasy kit

(Promega) and cDNA synthesised using random primers and the ImProm II reverse

transcription kit (Promega). Standard PCR reactions were performed with Phusion DNA

polymerase (New England Biolabs) using the relevant BS69 primers listed in Supplementary

Table S1. Quantitative PCR was performed in duplicate using the standard curve absolute

quantification method on an Applied Biosystems 7500 real-time PCR machine as described

previously (56) using the relevant primers listed in Supplementary Table S1. The efficiency of

all primers was determined prior to use and in each experiment and all had amplification

efficiencies within the recommended range (90–105%).

Data Availability

SAXS data have been deposited in the small angle scattering biological data bank (SASBDB)

(www.sasbdb.org) under accession numbers SASDEF6, SASDEG6, SASDEH6, SASDEJ6,

SASDEK6;

https://www.sasbdb.org/data/SASDEF6/5llm1wasc4/

https://www.sasbdb.org/data/SASDEG6/4yff1ro01u/

https://www.sasbdb.org/data/SASDEH6/tpia09j2m0/

https://www.sasbdb.org/data/SASDEJ6/werm0avk55/

https://www.sasbdb.org/data/SASDEK6/07t9uflv7k/

Acknowledgements

We thank Dr Stéphane Ansieau and Prof Gill Elliott for providing plasmids. We thank

Diamond Light Source for beamtime (proposal mx14891) and the staff of beamline B21 for

assistance with data collection.

 Figure Legends

Figure 1. Isothermal titration calorimetry analysis of the interaction of type 1 and type 2

EBNA2 peptides and polypeptides containing BS69 binding motif 2 with BS69$_{CC-MYND}$.

(A) Amino acid sequence of the regions of type 1 (B95-8) and type 2 (AG876) EBNA2

containing BS69 binding motifs 1 and 2 (underlined). Asterisks show the positions of the P, L

and P amino acids in the PXLXP binding motif. Aspartate 442 in type 1 EBNA2 and the

corresponding serine 409 in type 2 EBNA2 adjacent to motif 2 are shown in red. (B) Isothermal

titration calorimetry (ITC) analysis of type 1 EBNA2 motif 2 peptide binding to BS69$_{CC-MYND}$.

The upper panel shows heat peak data as Differential Power (DP) versus time and the lower

panel shows ΔH (derived from integration of the heat peak intensities) plotted against the

BS69$_{CC-MYND}$/EBNA2 molar ratio (based on monomer concentrations). Titrations were

performed using a series of 13 2.0 µl injections of 1 mM EBNA2 peptide and 0.1 mM BS69$_{CC-}$

$_{MYND}$ in the cell. The solid line shows the best fit using a one-site (one event) binding model

with the n value fixed to 1. The dissociation constant (K$_D$) displayed shows the mean ± standard

deviation obtained from 3 independent experiments (C) ITC analysis of the binding of the type

1 EBNA2 polypeptide T1 EBNA2$_{381-445}$ (0.6 mM) to BS69$_{CC-MYND}$ using 19 2.0 µl injections

of 0.1 mM EBNA2 polypeptide. The n value indicates the stoichiometry of BS69$_{CC-}$

$_{MYND}$/EBNA2 binding calculated for monomeric proteins and shows the mean ± standard

deviation from 3 independent experiments. (D) ITC analysis of the binding of the type 2

EBNA2 polypeptide T2 EBNA2$_{348-412}$ (0.3 mM) to BS69$_{CC-MYND}$ (0.1 mM) as in C.

Figure 2. Isothermal titration calorimetry analysis of the interaction of type 1 and type 2

EBNA2 peptides and polypeptides containing BS69 binding motif 3 with BS69$_{CC-MYND}$.

(A) Amino acid sequence of the regions of type 1 (B95-8) and type 2 (AG876) EBNA2

containing BS69 binding motifs 1, 2 and 3 (underlined). Asterisks show the positions of the P,

L and P amino acids in the PXLXP binding motif. (B) Isothermal titration calorimetry (ITC)

analysis of type 1 EBNA2 motif 3 peptide binding to $BS69_{CC-MYND}$ (no binding was detected

so the K_D could not be determined (n.d.). (C) Isothermal titration calorimetry (ITC) analysis of

type 2 EBNA2 motif 3 peptide binding to $BS69_{CC-MYND}$ using 19 injections of 2.0 μl EBNA2

peptide. Data are displayed and analysed as in Figure 1. (C) ITC analysis of the binding of type

2 EBNA2 motif 3 peptide binding to $BS69_{CC-MYND}$ as in B. (D) ITC analysis of the binding of

the type 1 EBNA2 polypeptide $EBNA2_{381-455}$ to $BS69_{CC-MYND.}$ (E) ITC analysis of the binding

of the type 2 EBNA2 polypeptide T2 $EBNA2_{348-422}$ to $BS69_{CC-MYND}$. (F) ITC analysis of the

binding of the type 2 EBNA2 polypeptide T2 $EBNA2_{348-422}$ with BS69 binding motif 3 mutated

from PTLEP to ATAEA.

Figure 3. Solution state analysis of EBNA2 and $BS69_{CC-MYND}$ complexes. (A) Size exclusion

chromatography of individual type 1 and type 2 EBNA2 polypeptides, $BS69_{CC-MYND}$ and

$EBNA2-BS69_{CC-MYND}$ complexes. Absorbance was normalized to the type2 EBNA2-BS69

complex (highest absorbance) using UNICORN software. (B) Samples from the indicated

fractions were analysed by SDS-PAGE followed by Coomassie staining.

Figure 4. Solution structure of $EBNA2-BS69_{CC-MYND}$ complexes determined by SAXS. (A)

SAXS scattering data for type 1 $EBNA2_{381-455}-BS69_{CC-MYND}$ (black dots) fitted to the *ab initio*

DAMMIN dummy atom model (red line). SAXS scattering data fitted to the docked structural

complex shown in B (green) with the χ^2 determined by FoXS. Plots show relative log intensity

vs scattering vector (q). (B) Solution structure of type 1 $EBNA2_{381-455}-BS69_{CC-MYND.}$ The

SAXS envelopes (grey mesh) were generated by averaging 23 *ab-initio* models using the

DAMMIF programme. The crystal structures of two $BS69_{CC-MYND}$ dimers (PDB ID: 5HDA)

are shown in cyan and were manually docked into the SAXS envelope along with the *ab-initio*

dummy atom SAXS solution structures of two type 1 $EBNA2_{381-455}$ polypeptides (salmon). The

maximum dimension (D_{max}) and volume were calculated using ScÅtter. (C) SAXS scattering

data for type 2 $EBNA2_{348-422}$-$BS69_{CC-MYND}$ (black dots) fitted to the *ab initio* DAMMIN

dummy atom model (red line). SAXS scattering data fitted to the docked structural complex

shown in D (green) with the χ^2 determined by FoXS. (D) Solution structure of type 2

$EBNA2_{348-422}$-$BS69_{CC-MYND}$ obtained as described in (B). The crystal structures of three

$BS69_{CC-MYND}$ dimers (PDB ID: 5HDA) are shown in cyan and were manually docked into the

SAXS envelope along with the *ab-initio* dummy atom SAXS solution structures of two type 2

$EBNA2_{348-422}$ polypeptides (orange).

Figure 5. Isotheral titration calorimetry analysis of the interaction of type 1 and type 2

$EBNA2$ polypeptides with $BS69_{CC-MYND}$ using an increased number of injections. (A) ITC

analysis of the binding of the type 1 EBNA2 polypeptide $EBNA2_{381-455}$ to $BS69_{CC-MYND.}$

Titrations were performed using a series of 29 injections of 1.3 µl 0.3 mM EBNA2 polypeptide

and 0.1 mM $BS69_{CC-MYND}$ in the cell. The solid line shows the best fit using a two-site (two

event) binding model. The mean dissociation constant (K_D) ± standard deviation from 3

independent experiments is shown for each binding event. (B) ITC analysis of the binding of

the type 2 EBNA2 polypeptide T2 $EBNA2_{348-422}$ to $BS69_{CC-MYND}$ using 29 injections and fitting

using a two-site (two event) binding model as in A. (C) ITC analysis of the binding of the type

2 EBNA2 polypeptide T2 $EBNA2_{348-422}$ SD containing the serine 412 to aspartate mutation

using 29 injections and fitting using a two-site (two event) binding model. (D) ITC analysis of

the binding of the type 2 EBNA2 polypeptide T2 $EBNA2_{348-422}$ with BS69 binding motif 3

mutated from PTLEP to ATAEA using 29 injections and fitting using a two-site (two event)

binding model.

Figure 6. GST pulldown assay using GST-BS69$_{CC-MYND}$ and lysates from B cell lines

expressing type 1 and type 2 EBNA2. Nuclear extracts of Daudi:pHEBo-MT:EBNA-2 cells

expressing type 1 (T1), type 2 (T2) or type 2 SD EBNA2 proteins were incubated for 5, 15 or

30 min at 4ºC with glutathione beads which had been loaded with bacterial lysates expressing

the GST- BS69$_{CC-MYND}$. GST-RAB11B was used as a negative control and was incubated for

30 min with the nuclear extracts. Following washing, beads were resuspended in protein sample

buffer and analysed by SDS-PAGE and Western blotting for EBNA2 using the PE2 anti-

EBNA2 antibody. The EBNA2 proteins expressed by these cell lines display almost the same

size (72 kD) since the number of polyproline residues had been equalised in all the EBNA2

alleles.

Figure 7. LCL growth maintenance assay using type 2 EBNA2 mutants. (A) ER-EB 2.5

cells conditionally expressing an estrogen receptor type 1 EBNA2 fusion protein were cultured

in medium containing β-estradiol prior to resuspension in medium without β-estradiol. Cells

were transfected with OriP-based plasmids (p294) expressing full length type 1 or type 2

EBNA2 or the type 2 SD, motif 3 or SD and motif 3 mutant EBNA2 proteins and cultured in

medium without β-estradiol. Live cells (that excluded Trypan Blue) were counted 1, 2, 3 and

4 weeks post-transfection. Data from a representative experiment of 4 independent repeats is

shown. Error bars show the mean ± standard deviation of duplicate cell counts for each sample.

(B) Western blot analysis of EBNA2 expression in protein extracts from the transfected

EREB2.5 cells. Cells were harvested 2 days after transfection. EBNA2 was detected using the

PE2 monoclonal antibody and blots were probed for actin as a loading control.

Figure 8. Western blot analysis of BS69 expression in EBV infected B cells. (A) Western

blot analysis of BS69 expression in the EBV negative BL cell line AK31 and type 1 and type

2 EBV immortalised LCLs (that display the latency III pattern of gene expression associated

with expression of all EBV latent proteins). BS69 was detected using an antibody that

recognises a sequence in the MYND domain encoded by exon 15 (ab190890). EBNA2 was

detected using the PE2 monoclonal antibody. Type 2 EBV has a lower molecular weight due a

difference in the number of proline repeat residues present. Blots were probed for actin as a

loading control. (B) Western blot analysis of BS69 expression in EBV negative (BJAB, DG75,

AK31) and EBV positive latency I (Akata and Mutu I) and latency III B cell lines (Mutu III

(BL) and IB4 (LCL) as in A. (C) Western blot analysis of BS69 expression on primary B cell

infection. UI indicates uninfected primary B cells. Samples were taken at the indicated day

post-infection and analysed as in A.

1002

Figure 9. PCR analysis of the BS69 isoforms expressed in EBV infected B cells. (A)

Diagram of the four experimentally verified BS69 isoforms. The position of the start codon

(ATG) and stop codon (red asterisk) is indicated. Numbering of exons is *as per* the canonical

isoform (isoform 1). Exon lengths are not to scale. Exon 4 is shown in red, Exon 15 is shown

in green and exon 14 is shown in blue. The 3' part of exon 14 present in isoforms 3 and 4 is

shown in blue hatched lines. A sequence within exon 15 is recognised by the antibody

(ab190890) used in Figure 8 as indicated on the diagram. Approximate locations of primers

used for conventional PCR and QPCR are indicated by black and red arrows respectively. (B)

Agarose gel analysis of PCR products generated using primers located in specific BS69 exons.

The upper panel shows the PCR products amplified from cDNA samples from type 1 and type

2 LCLs and the primary infection samples shown in C using a forward primer located in exon

3 and a reverse primer located in exon 13. Transcripts containing exon 4 (isoform 1 and 3) will

generate a 1139 bp product and those lacking exon 4 (isoforms 2 and 4) will produce a 977 bp

product. The lower panel shows PCR carried out using the exon 3 forward primer and a reverse

primer in the 3' region of exon 14 that is uniquely present in differentially spliced BS69

transcripts lacking exon 15 (isoforms 3 and 4). Transcripts containing exon 4 and lacking exon

15 (isoform 3) will generate a 1578 bp product and those lacking exon 4 and exon 15 will

produce a 1416 bp product (isoform 4). (C) QPCR analysis of cDNA from a panel of EBV

negative and positive B cell lines and primary EBV infections using primers that amplify across

the exon 14 and exon 15 junction. For EBV infections, samples from two different experiments

(#1 and #2) were used. Primary B cells (uninfected, UI) were infected with EBV and samples

harvested after 2 days (+ EBV). Results show the mean ± standard deviation of QPCR

replicates from a representative experiment. BS69 relative quantities were normalized to β2

microglobulin. (D) QPCR analysis as in (C) using primers within exon 14. The reverse primer

is located in the 3' part of exon 14 only present in BS69 isoform 3 and 4.

1028

Figure 10. Expression of a dominant negative form of BS69 increases EBNA2

transactivation. Transactivation assays in BJAB cells using EBNA2-GAL4-DNA binding

domain fusion proteins and a GAL4 reporter plasmid. Cells were cotransfected with 300 ng of

either type 1 GAL4-DBD:EBNA2 (aa 334-487) or type 2 GAL4-DBD:EBNA2 (301-454)

constructs, 500 ng of pFRLuc (Gal4 firefly luciferase reporter), 10 ng of pRL-CMV and 1 μg

of BS69 (pCI-BS69) or BS69 ΔMYND (pCI-BS69-ΔMYND) expressing plasmids. For each

sample, firefly luciferase values were normalised for transfection efficiency using Renilla

luciferase values. Results are presented as luciferase activity relative to the pFR-Luc reporter

plasmid plus empty vector (pcDNA3.1-GAL4-DBD). BS69 ΔMYND was also transfected in

the absence of EBNA2 expressing constructs. Results show the mean of two independent

experiments ± standard deviation.

1040

Table 1. Size exclusion chromatography and multiangle might scattering (MALS) determination of the molecular weight of individual type 1 and type 2 EBNA2 polypeptides, BS69$_{CC-MYND}$ and EBNA2-BS69 complexes. Theoretical and experimentally determined molecular weights are shown. For EBNA2-BS69 complexes, the theoretical molecular weights of complexes containing different numbers of EBNA2 and BS69 molecules are shown in parentheses. The most likely solution state based the experimentally determined molecular weight is indicated for each sample.

1048

Supporting Information Legends

Supplementary Figure S1. Normalised (dimensionless) Kratky plots generated using ScÅtter (v3.0 by Robert P. Rambo; Diamond Light Source). $I(q)/I(0)*(q*Rg)^2$ vs q*Rg. Scattering intensity I (q), scattering vector (q), radius of gyration (Rg).

1053

Supplementary Figure S2. Solution structures of BS69$_{CC-MYND}$ and type 1 and type 2 EBNA2 polypeptides determined by SAXS. (A-C) The SAXS envelopes (grey mesh) were generated by averaging 20 *ab-initio* models using the DAMMIF programme and further refined with DAMMIN to produce refined dummy atom models (magenta mesh). The maximum dimension (D$_{max}$) and volume were calculated using the ScÅtter programme. In (A) the BS69$_{CC-MYND}$ dimer structure (cyan; PDB ID: 5HDA) was manually docked into the envelope. (D-F) SAXS scattering data (black dots) fitted to the *ab initio* DAMMIN dummy atom (red line). χ^2 values for fitting are shown.

1062

Supplementary Figure S3. Alternative models and their respective goodness-of-fit to the

experimental SAXS data. (A) One BS69$_{CC-MYND}$ dimer (cyan; PDB ID: 5HDA) and the *ab*

initio model of one type 1 EBNA2$_{381-455}$ polypeptide (salmon) were manually docked into the

ab initio envelope (grey mesh) of the type 1 EBNA2 BS69 complex. (B) SAXS scattering data

were fitted to the docked structural complex shown in A and gave a χ^2 of 13.33 using FoXS.

Graphs show relative log intensity vs scattering vector (q) (upper panel) and the deviation

(residual) of the model from the experimental data (lower panel). The hydration parameter (C$_2$)

was fixed to 0 to prevent the hydration shell increasing to beyond the maximum limit of 4 to

attempt to fit the structure into the envelope. (C) The structural model shown in Figure 4C (two

BS69$_{CC-MYND}$ dimers and two type 1 EBNA2$_{381-455}$ polypeptides) was refitted to the SAXS

envelope using FoXS with the C$_2$ value set to 0 for comparison. This gave a similar χ^2 (2.59)

to that shown in Figure 4A indicating a much better fit to the scattering data. (D) Two BS69$_{CC-}$

$_{MYND}$BS69$_{CC-MYND}$ dimers (cyan; PDB ID: 5HDA) and the *ab initio* model of one type 2

EBNA2$_{348-422}$ polypeptide (orange) were manually docked into the *ab initio* envelope (grey mesh). (E)

SAXS scattering data were fitted to the docked structural complex shown in D and gave a χ^2 of 4.65

using FoXS. Graphs show relative log intensity vs scattering vector (q) (upper panel) and the deviation

(residual) of the model from the experimental data (lower panel). The hydration parameter (C$_2$) was

fixed to 0 to prevent the hydration shell increasing to high levels to attempt to fit the structure into the

envelope. (E) The structural model shown in Figure 4D (three BS69$_{CC-MYND}$ dimers and two type 2

EBNA2$_{348-422}$ polypeptides) was refitted to the SAXS envelope using FoXS with the C$_2$ value set to 0

for comparison. This gave a similar χ^2 (1.47) to that shown in Figure 4B indicating a much better fit to

the scattering data.

1085

1086 **Supplementary Table S1**. Primer sequences.

1087

1088   **Supplementary Table S2.** Data obtained from Isothermal calorimetry analysis of EBNA2

1089   peptides and polypeptides binding to $BS69_{CC-MYND}$ (n.d. indicates binding not detected). Data

1090   show the mean ± standard deviation for three independent experiments. For peptides n values

1091   were fixed to 1. Data from type 2 EBNA2 peptides and polypeptides are in shaded columns.

1092

1093   **Supplementary Table S3.** SAXS data for BS69 and EBNA2 polypeptides and complexes.

1094

1095

1096

References

1.   Cohen JI, Wang F, Mannick J, Kieff E. Epstein-Barr virus nuclear protein 2 is a key
determinant of lymphocyte transformation. Proc Natl Acad Sci U S A. 1989
Dec;86(23):9558-62. PubMed PMID: 2556717.
2.   Kempkes B, Ling PD. EBNA2 and Its Coactivator EBNA-LP. Curr Top Microbiol
Immunol. 2015;391:35-59. PubMed PMID: 26428371.
3.   Glaser LV, Rieger S, Thumann S, Beer S, Kuklik-Roos C, Martin DE, et al. EBF1
binds to EBNA2 and promotes the assembly of EBNA2 chromatin complexes in B cells.
PLoS Pathog. 2017 Oct;13(10):e1006664. PubMed PMID: 28968461. Pubmed Central
PMCID: 5638620.
4.   West MJ. Chromatin reorganisation in Epstein-Barr virus-infected cells and its role in
cancer development. Current opinion in virology. 2017 Sep 11;26:149-55. PubMed PMID:
28910751.
5.   Zhao B, Zou J, Wang H, Johannsen E, Peng CW, Quackenbush J, et al. Epstein-Barr
virus exploits intrinsic B-lymphocyte transcription programs to achieve immortal cell growth.
Proc Natl Acad Sci U S A. 2011 Sep 6;108(36):14902-7. PubMed PMID: 21746931. Pubmed
Central PMCID: 3169132. Epub 2011/07/13. eng.
6.   Wood CD, Veenstra H, Khasnis S, Gunnell A, Webb HM, Shannon-Lowe C, et al.
MYC activation and BCL2L11 silencing by a tumour virus through the large-scale
reconfiguration of enhancer-promoter hubs. eLife. 2016 Aug 04;5. PubMed PMID:
27490482. Pubmed Central PMCID: 5005034.
7.   Wang L, Grossman SR, Kieff E. Epstein-Barr virus nuclear protein 2 interacts with
p300, CBP, and PCAF histone acetyltransferases in activation of the LMP1 promoter. Proc
Natl Acad Sci U S A. 2000 Jan 4;97(1):430-5. PubMed PMID: 10618435.
8.   Wu DY, Kalpana GV, Goff SP, Schubach WH. Epstein-Barr virus nuclear protein 2
(EBNA2) binds to a component of the human SNF-SWI complex, hSNF5/Ini1. J Virol. 1996
Sep;70(9):6020-8. PubMed PMID: 8709224.
9.   Wu DY, Krumm A, Schubach WH. Promoter-specific targeting of human SWI-SNF
complex by Epstein-Barr virus nuclear protein 2. J Virol. 2000 Oct;74(19):8893-903.
PubMed PMID: 10982332.
10.   Tong X, Drapkin R, Reinberg D, Kieff E. The 62- and 80-kDa subunits of
transcription factor IIH mediate the interaction with Epstein-Barr virus nuclear protein 2.
Proc Natl Acad Sci U S A. 1995 Apr 11;92(8):3259-63. PubMed PMID: 7724549.
11.   Tong X, Drapkin R, Yalamanchili R, Mosialos G, Kieff E. The Epstein-Barr virus
nuclear protein 2 acidic domain forms a complex with a novel cellular coactivator that can
interact with TFIIE. Mol Cell Biol. 1995 Sep;15(9):4735-44. PubMed PMID: 7651391.
12.   Tong X, Wang F, Thut CJ, Kieff E. The Epstein-Barr virus nuclear protein 2 acidic
domain can interact with TFIIB, TAF40, and RPA70 but not with TATA-binding protein. J
Virol. 1995 Jan;69(1):585-8. PubMed PMID: 7983760.
13.   Cohen JI, Kieff E. An Epstein-Barr virus nuclear protein 2 domain essential for
transformation is a direct transcriptional activator. J Virol. 1991 Nov;65(11):5880-5. PubMed
PMID: 1656076.
14.   Sample J, Young L, Martin B, Chatman T, Kieff E, Rickinson A. Epstein-Barr virus
types 1 and 2 differ in their EBNA-3A, EBNA-3B, and EBNA-3C genes. J Virol. 1990
Sep;64(9):4084-92. PubMed PMID: 2166806.
15.   Dambaugh T, Hennessy K, Chamnankit L, Kieff E. U2 region of Epstein-Barr virus
DNA may encode Epstein-Barr nuclear antigen 2. Proc Natl Acad Sci U S A. 1984
Dec;81(23):7632-6. PubMed PMID: 6209719. Pubmed Central PMCID: 392202.

16.   Rowe M, Young LS, Cadwallader K, Petti L, Kieff E, Rickinson AB. Distinction
between Epstein-Barr virus type A (EBNA 2A) and type B (EBNA 2B) isolates extends to
the EBNA 3 family of nuclear proteins. J Virol. 1989 Mar;63(3):1031-9. PubMed PMID:
2536817. Pubmed Central PMCID: 247795.
17.   Palser AL, Grayson NE, White RE, Corton C, Correia S, Ba Abdullah MM, et al.
Genome diversity of Epstein-Barr virus from multiple tumor types and normal infection. J
Virol. 2015 May;89(10):5222-37. PubMed PMID: 25787276. Pubmed Central PMCID:
4442510.
18.   Correia S, Bridges R, Wegner F, Venturini C, Palser A, Middeldorp JM, et al.
Sequence variation of Epstein-Barr virus: viral types, geography, codon usage and diseases. J
Virol. 2018 Aug 15. PubMed PMID: 30111570.
19.   Rickinson AB, Young LS, Rowe M. Influence of the Epstein-Barr virus nuclear
antigen EBNA 2 on the growth phenotype of virus-transformed B cells. J Virol. 1987
May;61(5):1310-7. PubMed PMID: 3033261. Pubmed Central PMCID: 254104.
20.   Cancian L, Bosshard R, Lucchesi W, Karstegl CE, Farrell PJ. C-terminal region of
EBNA-2 determines the superior transforming ability of type 1 Epstein-Barr virus by
enhanced gene regulation of LMP-1 and CXCR7. PLoS Pathog. 2011 Jul;7(7):e1002164.
PubMed PMID: 21857817. Pubmed Central PMCID: 3145799.
21.   Tzellos S, Correia PB, Karstegl CE, Cancian L, Cano-Flanagan J, McClellan MJ, et
al. A Single Amino Acid in EBNA-2 Determines Superior B Lymphoblastoid Cell Line
Growth Maintenance by Epstein-Barr Virus Type 1 EBNA-2. J Virol. 2014 Aug
15;88(16):8743-53. PubMed PMID: 24850736.
22.   Lucchesi W, Brady G, Dittrich-Breiholz O, Kracht M, Russ R, Farrell PJ. Differential
gene regulation by Epstein-Barr virus type 1 and type 2 EBNA2. J Virol. 2008
Aug;82(15):7456-66. PubMed PMID: 18480445. Pubmed Central PMCID: 2493322. Epub
2008/05/16. eng.
23.   Coleman CB, Daud, II, Ogolla SO, Ritchie JA, Smith NA, Sumba PO, et al. Epstein-
Barr Virus Type 2 Infects T Cells in Healthy Kenyan Children. J Infect Dis. 2017 Sep
15;216(6):670-7. PubMed PMID: 28934430. Pubmed Central PMCID: 5853903.
24.   Coleman CB, Wohlford EM, Smith NA, King CA, Ritchie JA, Baresel PC, et al.
Epstein-Barr virus type 2 latently infects T cells, inducing an atypical activation characterized
by expression of lymphotactic cytokines. J Virol. 2015 Feb;89(4):2301-12. PubMed PMID:
25505080. Pubmed Central PMCID: 4338898.
25.   Coleman CB, Lang J, Sweet LA, Smith NA, Freed BM, Pan Z, et al. Epstein-Barr
virus type-2 infects T-cells and induces B-cell lymphomagenesis in humanized mice. J Virol.
2018 Aug 8. PubMed PMID: 30089703.
26.   Wen H, Li Y, Xi Y, Jiang S, Stratton S, Peng D, et al. ZMYND11 links histone
H3.3K36me3 to transcription elongation and tumour suppression. Nature. 2014 Apr
10;508(7495):263-8. PubMed PMID: 24590075. Pubmed Central PMCID: 4142212.
27.   Guo R, Zheng L, Park JW, Lv R, Chen H, Jiao F, et al. BS69/ZMYND11 reads and
connects histone H3.3 lysine 36 trimethylation-decorated chromatin to regulated pre-mRNA
processing. Mol Cell. 2014 Oct 23;56(2):298-310. PubMed PMID: 25263594. Pubmed
Central PMCID: 4363072.
28.   de Rooij JD, van den Heuvel-Eibrink MM, Kollen WJ, Sonneveld E, Kaspers GJ,
Beverloo HB, et al. Recurrent translocation t(10;17)(p15;q21) in minimally differentiated
acute myeloid leukemia results in ZMYND11/MBTD1 fusion. Genes, chromosomes &
cancer. 2016 Mar;55(3):237-41. PubMed PMID: 26608508.
29.   Wang J, Qin S, Li F, Li S, Zhang W, Peng J, et al. Crystal structure of human BS69
Bromo-ZnF-PWWP reveals its role in H3K36me3 nucleosome binding. Cell Res. 2014
Jul;24(7):890-3. PubMed PMID: 24675531. Pubmed Central PMCID: 4085756.

30.   Ladendorff NE, Wu S, Lipsick JS. BS69, an adenovirus E1A-associated protein,
inhibits the transcriptional activity of c-Myb. Oncogene. 2001 Jan 4;20(1):125-32. PubMed
PMID: 11244510.

31.   Masselink H, Vastenhouw N, Bernards R. B-myb rescues ras-induced premature
senescence, which requires its transactivation domain. Cancer Lett. 2001 Sep 28;171(1):87-
101. PubMed PMID: 11485831.

32.   Wei G, Schaffner AE, Baker KM, Mansky KC, Ostrowski MC. Ets-2 interacts with
co-repressor BS69 to repress target gene expression. Anticancer research. 2003 May-
Jun;23(3A):2173-8. PubMed PMID: 12894593.

33.   Velasco G, Grkovic S, Ansieau S. New insights into BS69 functions. J Biol Chem.
2006 Jun 16;281(24):16546-50. PubMed PMID: 16565076.

34.   Ansieau S, Leutz A. The conserved Mynd domain of BS69 binds cellular and
oncoviral proteins through a common PXLXP motif. J Biol Chem. 2002 Feb 15;277(7):4906-
10. PubMed PMID: 11733528.

35.   Harter MR, Liu CD, Shen CL, Gonzalez-Hurtado E, Zhang ZM, Xu M, et al.
BS69/ZMYND11 C-Terminal Domains Bind and Inhibit EBNA2. PLoS Pathog. 2016
Feb;12(2):e1005414. PubMed PMID: 26845565. Pubmed Central PMCID: 4742278.

36.   Friberg A, Thumann S, Hennig J, Zou P, Nossner E, Ling PD, et al. The EBNA-2 N-
Terminal Transactivation Domain Folds into a Dimeric Structure Required for Target Gene
Activation. PLoS Pathog. 2015 May;11(5):e1004910. PubMed PMID: 26024477. Pubmed
Central PMCID: 4449002.

37.   Kempkes B, Spitkovsky D, Jansen-Durr P, Ellwart JW, Kremmer E, Delecluse HJ, et
al. B-cell proliferation and induction of early G1-regulating proteins by Epstein-Barr virus
mutants conditional for EBNA2. Embo J. 1995 Jan 3;14(1):88-96. PubMed PMID: 7828599.

38.   Tsui S, Schubach WH. Epstein-Barr virus nuclear protein 2A forms oligomers in vitro
and in vivo through a region required for B-cell transformation. J Virol. 1994 Jul;68(7):4287-
94. PubMed PMID: 8207803. Pubmed Central PMCID: 236351.

39.   Harada S, Yalamanchili R, Kieff E. Epstein-Barr virus nuclear protein 2 has at least
two N-terminal domains that mediate self-association. J Virol. 2001 Mar;75(5):2482-7.
PubMed PMID: 11160754. Pubmed Central PMCID: 114834.

40.   Wan J, Zhang W, Wu L, Bai T, Zhang M, Lo KW, et al. BS69, a specific adaptor in
the latent membrane protein 1-mediated c-Jun N-terminal kinase pathway. Mol Cell Biol.
2006 Jan;26(2):448-56. PubMed PMID: 16382137. Pubmed Central PMCID: 1346911.

41.   Ikeda O, Sekine Y, Mizushima A, Oritani K, Yasui T, Fujimuro M, et al. BS69
negatively regulates the canonical NF-kappaB activation induced by Epstein-Barr virus-
derived LMP1. FEBS Lett. 2009 May 19;583(10):1567-74. PubMed PMID: 19379743.

42.   Ikeda O, Miyasaka Y, Yoshida R, Mizushima A, Oritani K, Sekine Y, et al. BS69
cooperates with TRAF3 in the regulation of Epstein-Barr virus-derived LMP1/CTAR1-
induced NF-kappaB activation. FEBS Lett. 2010 Mar 5;584(5):865-72. PubMed PMID:
20138174.

43.   Ben-Bassat H, Goldblum N, Mitrani S, Goldblum T, Yoffey JM, Cohen MM, et al.
Establishment in continuous culture of a new type of lymphocyte from a "Burkitt like"
malignant lymphoma (line D.G.-75). Int J Cancer. 1977 Jan;19(1):27-33. PubMed PMID:
188769. Epub 1977/01/01. eng.

44.   Jenkins PJ, Binne UK, Farrell PJ. Histone acetylation and reactivation of Epstein-Barr
virus from latency. J Virol. 2000 Jan;74(2):710-20. PubMed PMID: 10623733. Pubmed
Central PMCID: 111591. Epub 2000/01/07. eng.

45.   Klein G. Studies on Epstein-Barr Virus Genome and Ebv-Determined Nuclear
Antigen in Human Malignant Disease. Cold Spring Harbor symposia on quantitative biology.
1974;39:783-90. PubMed PMID: WOS:A1974AB46200012. English.

46.   Takada K. Cross-linking of cell surface immunoglobulins induces Epstein-Barr virus
in Burkitt lymphoma lines. Int J Cancer. 1984 Jan 15;33(1):27-32. PubMed PMID: 6319296.
47.   Gregory CD, Rowe M, Rickinson AB. Different Epstein-Barr virus-B cell interactions
in phenotypically distinct clones of a Burkitt's lymphoma cell line. J Gen Virol. 1990 Jul;71 (
Pt 7):1481-95. PubMed PMID: 2165133. Epub 1990/07/01. eng.
48.   Spender LC, Lucchesi W, Bodelon G, Bilancio A, Karstegl CE, Asano T, et al. Cell
target genes of Epstein-Barr virus transcription factor EBNA-2: induction of the p55alpha
regulatory subunit of PI3-kinase and its role in survival of EREB2.5 cells. J Gen Virol. 2006
Oct;87(Pt 10):2859-67. PubMed PMID: 16963743. Epub 2006/09/12. eng.
49.   Brocard M, Khasnis S, Wood CD, Shannon-Lowe C, West MJ. Pumilio directs
deadenylation-associated translational repression of the cyclin-dependent kinase 1 activator
RGC-32. Nucleic Acids Res. 2018 Jan 27. PubMed PMID: 29385536. Pubmed Central
PMCID: 5909466.
50.   Franke D, Svergun DI. DAMMIF, a program for rapid ab-initio shape determination
in small-angle scattering. J Appl Crystallogr. 2009 Apr;42:342-6. PubMed PMID:
WOS:000264292800030. English.
51.   Volkov VV, Svergun DI. Uniqueness of ab initio shape determination in small-angle
scattering. J Appl Crystallogr. 2003 Jun;36:860-4. PubMed PMID: WOS:000182284400105.
English.
52.   Svergun DI. Restoring low resolution structure of biological macromolecules from
solution scattering using simulated annealing (vol 76, pg 2879, 1999). Biophys J. 1999
Nov;77(5):2896-. PubMed PMID: WOS:000083554900051. English.
53.   Schneidman-Duhovny D, Hammel M, Sali A. FoXS: a web server for rapid
computation and fitting of SAXS profiles. Nucleic Acids Research. 2010 Jul;38:W540-W4.
PubMed PMID: WOS:000284148900087. English.
54.   West MJ, Webb HM, Sinclair AJ, Woolfson DN. Biophysical and mutational analysis
of the putative bZIP domain of Epstein-Barr virus EBNA 3C. J Virol. 2004 Sep;78(17):9431-
45. PubMed PMID: 15308737.
55.   Bark-Jones SJ, Webb HM, West MJ. EBV EBNA 2 stimulates CDK9-dependent
transcription and RNA polymerase II phosphorylation on serine 5. Oncogene. 2006 Mar
16;25(12):1775-85. PubMed PMID: 16314842. Epub 2005/11/30. eng.
56.   McClellan MJ, Wood CD, Ojeniyi O, Cooper TJ, Kanhere A, Arvey A, et al.
Modulation of enhancer looping and differential gene targeting by Epstein-Barr virus
transcription factors directs cellular reprogramming. PLoS Pathog. 2013 Sep;9(9):e1003636.
PubMed PMID: 24068937. Pubmed Central PMCID: 3771879. Epub 2013/09/27. eng.

1280

1281

Figure 1

Main Figures combined file