**Exome sequencing identifies high-impact trait-associated alleles enriched in Finns**

Locke, Adam E[1,2,3,*]; Meltz Steinberg, Karyn[2,4,*]; Chiang, Charleston WK[5,6,*]; Service, Susan K[5,*]; Havulinna, Aki S[7,8]; Stell, Laurel[9]; Pirinen, Matti[7,10,11]; Abel, Haley J[2,12]; Chiang, Colby C[2]; Fulton, Robert S[2]; Jackson, Anne U[3]; Kang, Chul Joo[2]; Kanchi, Krishna L[2]; Koboldt, Daniel C[2,13,14]; Larson, David E[2,12]; Nelson, Joanne[2]; Nicholas, Thomas J[2,15]; Pietilä, Arto[8]; Ramensky, Vasily[5,16]; Ray, Debashree[3,17]; Scott, Laura J[3]; Stringham, Heather M[3]; Vangipurapu, Jagadish[18]; Welch, Ryan[3]; Yajnik, Pranav[3]; Yin, Xianyong[3]; Eriksson, Johan G[19,20,21]; Ala-Korpela, Mika[22,23,24,25,26,27]; Järvelin, Marjo-Riitta[28,29,30,31,32]; Männikkö, Minna[29,33]; Laivuori, Hannele[7,34,35]; FinnGen Project; Dutcher, Susan K[2,12]; Stitziel, Nathan O[2,36]; Wilson, Richard K[2,13,14]; Hall, Ira M[1,2]; Sabatti, Chiara[9,37]; Palotie, Aarno[7,38,39]; Salomaa, Veikko[8]; Laakso, Markku[18,40]; Ripatti, Samuli[7,10,39]; Boehnke, Michael[3,†]; Freimer, Nelson B[5,†]

[1]Department of Medicine, Washington University School of Medicine, St. Louis, MO
[2]McDonnell Genome Institute, Washington University School of Medicine, St. Louis, MO
[3]Department of Biostatistics and Center for Statistical Genetics, University of Michigan School of Public Health, Ann Arbor, MI
[4]Department of Pediatrics, Washington University School of Medicine, St. Louis, MO
[5]Center for Neurobehavioral Genetics, Jane and Terry Semel Institute for Neuroscience and Human Behavior, University of California Los Angeles, Los Angeles, CA
[6]Center for Genetic Epidemiology, Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA
[7]Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland
[8]National Institute for Health and Welfare, Helsinki, Finland
[9]Department of Biomedical Data Science, Stanford University, Stanford, CA
[10]Department of Public Health, University of Helsinki, Helsinki, Finland;
[11]Helsinki Institute for Information Technology HIIT and Department of Mathematics and Statistics, University of Helsinki, Helsinki, Finland
[12]Department of Genetics, Washington University School of Medicine, St. Louis, MO
[13]The Institute for Genomic Medicine, Nationwide Children's Hospital, Columbus, OH
[14]Department of Pediatrics, The Ohio State University College of Medicine, Columbus, OH
[15]USTAR Center for Genetic Discovery and Department of Human Genetics, University of Utah, Salt Lake City, UT
[16]Federal State Institution "National Medical Research Center for Preventive Medicine" of the Ministry of Healthcare of the Russian Federation, Moscow, Russia
[17]Departments of Epidemiology and Biostatistics, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD
[18]Institute of Clinical Medicine, Internal Medicine, University of Eastern Finland, Kuopio, Finland
[19]Department of Public Health Solutions, National Institute for Health and Welfare, Helsinki, Finland
[20]Folkhälsan Research Center, Helsinki, Finland

[21]Department of General Practice and Primary Health Care, University of Helsinki,
Helsinki and Helsinki University Hospital, Helsinki, Finland
[22]Systems Epidemiology, Baker Heart and Diabetes Institute, Melbourne, Victoria,
Australia
[23]Computational Medicine, Faculty of Medicine, University of Oulu and Biocenter Oulu,
University of Oulu, Oulu, Finland
[24]NMR Metabolomics Laboratory, School of Pharmacy, University of Eastern Finland,
Kuopio, Finland
[25]Population Health Science, Bristol Medical School, University of Bristol, Bristol, UK
[26]Medical Research Council Integrative Epidemiology Unit at the University of Bristol,
Bristol, UK
[27]Department of Epidemiology and Preventive Medicine, School of Public Health and
Preventive Medicine, Faculty of Medicine, Nursing and Health Sciences, The Alfred
Hospital, Monash University, Melbourne, Victoria, Australia
[28]Biocenter Oulu, University of Oulu, Oulu, Finland
[29]Center for Life Course Health Research, Faculty of Medicine, University of Oulu, Oulu,
Finland
[30]Unit of Primary Health Care, Oulu University Hospital, Oulu, Finland
[31]Department of Epidemiology and Biostatistics, MRC-PHE Centre for Environment and
Health, School of Public Health, Imperial College London, London, UK
[32]Department of Life Sciences, College of Health and Life Sciences, Brunel University
London, Uxbridge, UK
[33]Northern Finland Birth Cohorts, Faculty of Medicine, University of Oulu, Oulu,
Finland
[34]Medical and Clinical Genetics, University of Helsinki and Helsinki University Hospital,
Helsinki, Finland
[35]Department of Obstetrics and Gynecology, Tampere University Hospital and University
of Tampere, Faculty of Medicine and Life Sciences, Tampere, Finland
[36]Cardiovascular Division, Department of Medicine, Washington University School of
Medicine, St. Louis, MO
[37]Department of Statistics, Stanford University, Stanford, CA
[38]Analytical and Translational Genetics Unit (ATGU), Psychiatric &
Neurodevelopmental Genetics Unit, Departments of Psychiatry and Neurology,
Massachusetts General Hospital, Boston, MA
[39]Broad Institute of MIT and Harvard, Cambridge, MA
[40]Department of Medicine, Kuopio University Hospital, Kuopio, Finland
82
[*]These authors contributed equally to this work.
[†]These authors jointly supervised this work.

**ABSTRACT**

As yet undiscovered rare variants are hypothesized to substantially influence an

individual's risk for common diseases and traits, but sequencing studies aiming to

identify such variants have generally been underpowered. In isolated populations that

have expanded rapidly after a population bottleneck, deleterious alleles that passed

through the bottleneck may be maintained at much higher frequencies than in other

populations. In an exome sequencing study of nearly 20,000 cohort participants from

northern and eastern Finnish populations that exemplify this phenomenon, most novel

trait-associated deleterious variants are seen only in Finland or display frequencies more

than 20 times higher than in other European populations. These enriched alleles underlie

34 novel associations with 21 disease-related quantitative traits and demonstrate a

geographical clustering equivalent to that of Mendelian disease mutations characteristic

of the Finnish population. Sequencing studies in populations without this unique history

would require hundreds of thousands to millions of participants for comparable power for

these variants.

**INTRODUCTION**

Genotyping studies of common genetic variants (defined here as minor allele frequency

[MAF]>1%) have identified tens of thousands of genome-wide significant associations

with common diseases and disease-related quantitative traits[1]. For most traits, however,

these associations account for only a modest fraction of trait heritability, and the

mechanisms through which associated variants contribute to biological processes remain

mostly unknown. These observations have led to the expectation that rare variants

109 (defined here as MAF≤1%) which are not well-tagged by the single-nucleotide

110 polymorphisms (SNPs) on genome-wide genotyping arrays are probably responsible for

111 much of the heritability that remains unexplained[2]. Additionally, because purifying

112 selection acts to remove deleterious alleles from the population, most variants that exert a

113 sizable effect on complex traits, and that likely offer the best prospect for revealing

114 biological mechanisms, should be particularly rare.

116 Rare variants are unevenly distributed between populations and difficult to represent

117 effectively on commercial genotyping arrays, as evidenced by relatively sparse

118 association findings even from large array-based studies of coding variants[3-6].

119 Discovering rare variant associations will therefore almost certainly require exome or

120 genome sequencing of very large numbers of individuals. However, the sample size

121 required to reliably identify rare-variant associations remains uncertain; most sequencing

122 studies to date have identified few novel associations, and theoretical analyses confirm

123 that they have been underpowered to do so[7]. These analyses also suggest that power to

124 detect rare variant associations varies enormously between populations that have

125 expanded in isolation from recent bottlenecks compared to those that have not.

127 In isolated populations that expand rapidly following a bottleneck, alleles that pass

128 through the bottleneck often rise to a much higher frequency than in other populations[8-10].

129 If the bottleneck was recent, even deleterious alleles under negative selection may remain

130 relatively frequent in these populations, resulting in increased power to detect association

131 with disease-related traits. The Finnish population exemplifies this type of history. It

grew from bottlenecks occurring 2,000-4,000 years ago in the founding of the early-

settlement regions of southern and western Finland; internal migration in the 15[th] and 16[th]

centuries to the late-settlement regions of northern and eastern Finland created additional

bottlenecks[11]. The subsequent rapid growth of the Finnish population (to ~5.5 million,

larger than any other human isolate) generated sizable geographic sub-isolates in late-

settlement regions.

Geneticists have long noted that the bottlenecks that were so prominent in Finland's

recent history caused 36 Mendelian disorders to be much more common in Finland than

in other European countries, while several other disorders are much less common, a

phenomenon termed "the Finnish Disease Heritage"[12]. The identification of mutations for

35 of these disorders has confirmed that they mostly concentrate in late settlement

regions[12]. Additional studies demonstrated, in these regions, an overall enrichment of

deleterious variants more extreme compared to other isolates or to Finland generally[13-15].

We reasoned that this enrichment would enable exome sequencing studies of late-

settlement Finland to be better powered than studies in other populations to

systematically investigate the impact of low-frequency variants on disease-related

quantitative traits. Based on this expectation, we formed such a sample ("FinMetSeq")

from two Finnish population-based cohort studies: FINRISK and METSIM (see

Methods).

Using >1.4 M variants identified and genotyped by successful exome sequencing of

19,292 FinMetSeq participants, we conducted single-variant association analysis with 64

clinically relevant quantitative traits[16,17]. We identified 43 novel associations with

deleterious variants in 25 traits: 19 associations (11 traits) in FinMetSeq and 24

associations (20 traits) in a combined analysis of FinMetSeq with an additional 24,776

Finns from three cohorts for which imputed array-based genome-wide genotype data

were available. Nineteen of the 26 variants underlying these 43 novel associations were

unique to Finland or enriched >20-fold in FinMetSeq compared to non-Finnish

Europeans (NFE).

We demonstrate that (1) a well-powered exome sequencing study can identify numerous

rare alleles, each of which has a substantial effect on one or more traits in the individuals

who carry them, and (2) exome sequencing in a population that has expanded after recent

population bottlenecks is an extraordinarily efficient strategy to discover such effects. As

most of the novel putatively deleterious trait-associated variants that we identified are

unique to or highly enriched in Finland, similarly powered studies of these variants in

non-Finnish populations might require hundreds of thousands or even millions of

participants. Additionally, the geographical clustering of these enriched alleles, like the

Finnish Disease Heritage mutations, demonstrates that the distribution of trait-associated

rare alleles may vary significantly between locales within a country.

**RESULTS**

**Genetic variation**

We attempted to sequence the protein-coding regions of 23,585 genes covering 39 MB of

genomic sequence in 20,316 FinMetSeq participants. After extensive quality control, we

included in downstream analysis 19,292 individuals sequenced to 47x mean depth

(Methods, **Supplementary Table 1**). We identified 1,318,781 single nucleotide variants

(SNVs) and 92,776 insertion/deletion (indel) variants, with a mean of 20,989 SNVs and

604 indel variants per individual. The majority (87.5%) of SNVs identified were rare

(MAF<1%); 40.5% were singletons (**Table 1**). Each participant carried 15 singleton

variants on average, 17 rare (MAF≤1%) protein truncating variants (PTVs; annotated as

stop gain, essential splice site, start loss, or frameshift) alleles, and 171 common

(MAF>1%) PTVs (**Supplementary Table 2**). Frameshift indels accounted for the largest

proportion of PTVs (31% of rare, 42% of common), while stop gain variants were the

most frequent type of protein truncating SNVs (29% of rare, 20% of common).

We compared variant allele frequencies in FinMetSeq to those of NFE control exomes

from the Genome Aggregation Database (gnomAD v2.1, **Extended Data Fig. 1**). As in

previous smaller-scale comparisons of Finnish and NFE exomes, in FinMetSeq we

observe a depletion of the rarest alleles (singletons and doubletons) and a relative excess

of more common variants (minor allele count, MAC ≥5) compared to NFE for all classes

of variants. This effect is particularly marked for alleles predicted to be deleterious

(**Extended Data Fig. 2**).

**Single-variant association analyses**

We tested for association between genetic variants in FinMetSeq and 64 clinically

relevant quantitative traits measured in members of both FINRISK and METSIM

(**Supplementary Table 3**). We adjusted lipid and blood pressure traits for lipid lowering

and antihypertensive medication use, respectively, adjusted all traits for covariates using

linear regression (**Supplementary Table 4**), and inverse normalized trait residuals to

generate normally distributed traits for genetic association analysis that assumed an

additive model (Methods). Based on common variants, 62 of 64 traits exhibited

significant heritability (P<0.05; $h^2$ range 5.0-52.5%; **Fig. 1A**, **Supplementary Table 5**),

and there was substantial correlation between traits, phenotypically and genetically (**Fig.**

**1B**)**.**

We tested the 64 traits for single-variant associations with the 362,996 to 602,080 genetic

variants with MAC ≥3 among the 3,558 to 19,291 individuals measured for each trait

(**Supplementary Tables 3 & 4**). Association results are available for download and can

be explored interactively with PheWeb (http://pheweb.sph.umich.edu/FinMetSeq/) and

via the Type 2 Diabetes Knowledge Portal (www.type2diabetesgenetics.org). We

identified 1,249 trait-variant associations ($P<5\times10^{-7}$) at 531 variants (**Supplementary**

**Table 6**), with 53 of 64 traits associated with at least one variant (**Fig. 2A**). All 1,249

associations remained significant after multiple testing adjustment across the exome and

across the 64 traits with a hierarchical procedure setting average FDR at 5% (Methods).

Using the hierarchical FDR procedure, we detected an additional 287 trait-variant

associations at these 531 variants (**Supplementary Table 7**). These additional

associations reflect the high correlation between a subset of lipid traits, e.g. high-density

lipoprotein cholesterol (HDL-C) and apolipoprotein A1 (ApoA1). Given the diversity of

traits assessed in these cohorts, these associations may shed additional light on the

biology of measures that have been less frequently assayed in large GWAS, such as

intermediate density lipoproteins (IDL) and very-low-density lipoprotein (VLDL)

particles. Of the 531 associated variants, 59 (11%) were rare (MAF≤1%); by annotation, 200 (38%) were coding, 108 (20%) missense, and 11 (2%) protein truncating. Furthermore, minor alleles at >10-fold increased frequency in FinMetSeq compared to NFE are substantially more likely to be associated with a trait compared to variants with similar or lower MAF in FinMetSeq compared to NFE (OR=4.92, P=2.6×10$^{-5}$; **Extended Data Fig. 3**).

We clumped associated variants within 1 Mb and with r$^2$>0.5 into a single locus, irrespective of the associated traits (Methods). After clumping, the 531 associated variants represented 262 distinct loci (597 trait-locus pairs, **Supplementary Table 6**); 158 of the 262 loci (60%) consisted of a single trait-associated variant. As expected, the number of associated loci per trait was positively correlated with trait heritability (r=0.38, P=8.8×10$^{-4}$). Height was a noticeable outlier, with relatively few associations considering its high estimated heritability (**Fig. 2B**).

The majority of variants and loci (61%) were associated with a single trait; 4% were associated with ≥10 traits. Overlapping associations (**Fig. 2C**) strongly reflect the relationships exhibited by both trait and genetic correlations (**Fig. 1B**). For example, rs113298164, a missense variant in *LIPC* (p.Thr405Met), is associated with 11 traits, including cholesterols, fatty acids, apolipoproteins, and cholines. Similarly, the estimated genetic correlation of trait pairs is a strong predictor of the probability for a trait pair to share associated loci (**Fig. 2D**).

To determine which of the 1,249 single-variant associations were distinct from known

GWAS associations for the same traits, we repeated association analysis for each trait

conditional on published associated variants ($P<10^{-7}$) for the corresponding trait in the

EBI GWAS Catalog (December 2016 release). Of the 1,249 trait-variant associations,

478 (at 213 of 531 variants) remained significant ($P<5\times10^{-7}$) after conditional analysis,

representing 126 of the original 262 loci, including at least one conditionally significant

locus for each of 48 traits (**Supplementary Table 8**). The conditionally-associated

variants were more often rare (24% vs. 11%), more likely to alter or truncate the resulting

protein (31% vs. 22%), and more frequently >10x enriched in FinMetSeq relative to NFE

(19% vs. 10%) compared to the full set of associated variants.

**Gene-based association analyses**

To identify genes associated with the 64 traits, we performed aggregate tests of protein

coding variants, grouping variants using three different masks. Mask 1 comprised PTVs

of any frequency; Masks 2 and 3 also included missense variants with MAF<0.1% or

0.5% predicted to be deleterious by five algorithms (Methods). We identified 54 gene-

based associations with $P<3.88\times10^{-6}$ (adjusting for testing a maximum of 12,890 genes

containing at least two qualifying variants) and with multi-trait FDR<0.05, analogous to

the threshold used for single-variant association testing (Methods). Fifteen of these

associations required $\geq2$ variants to achieve significance (i.e. the association was not

driven by a single strongly associated variant; **Supplementary Table 9**). Extremely rare

(MAC$\leq$3) PTVs drove the association of eight traits with *APOB* (**Extended Data Fig. 4**).

We found a novel association between two very rare stop gain variants in *SECTM1* and

HDL2 cholesterol (P=7.2×10$^{-7}$, **Extended Data Fig. 5**). *SECTM1* encodes an interferon-

induced transmembrane protein that is negatively regulated by bacterial

lipopolysaccharide (LPS)[18]. The association could reflect the role of HDL particles in

binding and neutralizing LPS in infections and sepsis[19].

**Replication and follow-up of single-variant associations in three additional Finnish**

**cohorts: Identification of novel coding, deleterious variant associations**

We attempted to replicate the 478 single-variant associations from FinMetSeq

(unconditional and conditional P≤5×10$^{-7}$) and to follow-up the 2,120 suggestive but sub-

threshold associations from FinMetSeq (unconditional 5×10$^{-7}$<P≤5×10$^{-5}$, conditional

P≤5×10$^{-5}$) in 24,776 participants from three Finnish cohort studies for which varying

subsets of the 64 FinMetSeq traits were available: FINRISK[20,21] participants not

sequenced in FinMetSeq (n=18,215), the Northern Finland Birth Cohort 1966[22]

(n=5,139), and the Helsinki Birth Cohort[23] (n=1,412). For each of the three cohorts, we

carried out genotype imputation using the Finnish-specific SISu v2 reference panel

(http://www.sisuproject.fi), which is comprised of 5,380 haplotypes from whole-genome

based sequencing and 10,184 haplotypes from whole-exome based sequencing in coding

regions, and then used the same single-variant association analysis strategy employed in

FinMetSeq. We then carried out meta-analysis of the three imputation-based studies to

test for replication of associated FinMetSeq variants ("replication analysis") and four-

study meta-analysis with FinMetSeq to follow-up suggestive associations ("combined

analysis"; Methods).

We obtained data for 448 of the 478 significant variant-trait associations (191 of the 213

requested variants). Of the 448 associations for which we had replication data, 439

(98.0%) had the same direction of effect in replication analysis as in FinMetSeq; 392 of

the 448 replicated at P<0.05 (87.5%; **Supplementary Table 10**). We also obtained data

to follow up 1,417 of the 2,120 sub-threshold associations (1,014 of the 1,554 requested

variants); >60% of the variants that we could not follow up were very rare in FinMetSeq

and were not present in the SISu reference panel. Of the 1,417 sub-threshold trait-variant

associations, 431 reached $P<5\times10^{-7}$ in the combined analysis (**Supplementary Table**

**11**).

Among the significant results from FinMetSeq or combined analysis, 43 associations

were with 26 predicted deleterious variants that conditional analysis and literature review

suggest are novel (**Table 2**). Nineteen such associations, at 15 deleterious coding

variants, were significant in FinMetSeq (**Table 2; Supplementary Table 10**). Twelve of

these associations replicated (P<0.05) in the replication analysis and remained significant

in the combined analysis; for the other seven associations we either did not have

replication data (six associations) or did not replicate but had very low power (<5%) in

the replication analysis (one association). Four of the 15 variants were PTVs; 11 were

missense variants predicted to be deleterious by at least one of five prediction algorithms.

Another 24 associations, with 16 variants (two PTVs and 14 missense variants predicted

to be deleterious), only reached significance in the combined analysis (**Table 2;**

**Supplementary Table 11**). Five variants with significant associations in FinMetSeq

alone were associated with additional traits in combined analysis (**Table 2**).

Of the 43 associations shown in **Table 2**, 34 were with 19 variants either seen only in

Finland or enriched by >20-fold in FinMetSeq compared to NFE (13 of 15 variants in

FinMetSeq and 11 of 16 variants in combined analysis with five variants overlapping).

Identifying associations for these 19 variants would have required much larger samples in

NFE populations than in FinMetSeq (**Fig. 3A & B**). We provide brief summaries relating

each of these highly enriched associations to known biology and prior genetic evidence

relating to the respective genes in **Supplementary Information.** We highlight a few of

the most striking findings, below.

*Anthropometric traits.* As a group these are among the most extensively investigated

quantitative traits, with thousands of common variant associations reported, most of very

small effect[24-28]. We identified several rare, large effect variants for these traits, including

a predicted damaging missense variant (rs200373343, p.Arg94Cys) in *THBS4* 45X more

frequent in FinMetSeq than in NFE and associated in the combined analysis with a mean

decrease in body weight of 5.9 kg (**Table 2**). *THBS4* encodes thrombospondin 4, a

matricellular protein found in blood vessel walls and highly expressed in heart and

adipose tissue[29]. *THBS4* is involved in local signaling in the developing and adult nervous

system, and may function in regulating vascular inflammation[30]. Coding variants in

*THBS4* and other thrombospondin genes have been implicated in increased risk for heart

disease[31-33].

We identified a predicted damaging missense variant (rs2273607, p.Val104Met) in *DLK1*

that is 177X more frequent in FinMetSeq than in NFE and is associated in the combined

analysis with a mean decrease in height of 1.3 cm (**Table 2**). *DLK1* encodes Delta-Like

Notch Ligand 1, an epidermal growth factor that interacts with fibronectin and inhibits

adipocyte differentiation. Uniparental disomy of *DLK1* causes Temple Syndrome and

Kagami-Ogata Syndrome, characterized by pre- and postnatal growth restriction,

hypotonia, joint laxity, motor delay, and early onset of puberty[34-36]. Paternally-inherited

common variants near *DLK1* have been associated with child and adolescent obesity,

type 1 diabetes, age at menarche, and central precocious puberty in girls[37-39].

Homozygous null mutations in the mouse ortholog Dlk-1 lead to embryos with reduced

size, skeletal length, and lean mass[40], while in Darwin's finches, SNVs at this locus have

a strong effect on beak size[41].

*HDL-C.* Two novel variants with large effects on HDL-C in FinMetSeq are absent in

NFE. The predicted deleterious missense variant rs750623950 (p.Arg112Trp) in

*CD300LG* is associated in FinMetSeq with a mean increase in HDL-C of 0.95 mmol/l,

and also associated with HDL2-C and ApoA1 (**Table 2**). *CD300LG* encodes a type I cell

surface glycoprotein. A missense variant in *ABCA1* (rs765246726, p.Cys2107Arg) is

associated in FinMetSeq with a mean reduction in HDL-C of 0.64 mmol/l (**Table 2**).

Fifteen more variants (including ten which are absent in NFE) contributed to a strong

*ABCA1* gene-based association signal (P=$2.2 \times 10^{-13}$; **Supplementary Table 9, Extended**

**Data Fig. 6**). *ABCA1* encodes the cholesterol efflux regulatory protein, which regulates

cholesterol and phospholipid metabolism. Individuals who are homozygotes or

compound heterozygotes for any of several *ABCA1* mutations produce very little HDL-C

and experience the manifestations of severe hypercholesterolemia.

*Amino Acids.* A stop gain variant (rs780671030, p.Arg722X) in *ALDH1L1* is associated

in FinMetSeq with a mean reduction in serum glycine levels of 0.03 mmol/l but is not

observed in NFE (**Table 2**); this effect may increase risk for several cardiometabolic

disorders[42,43]. *ALDH1L1* encodes 10-formyltetrahydrofolate dehydrogenase, which

competes with the enzyme serine hydroxymethyltransferase to alter the ratio of serine to

glycine in the cytosol. Although rs780671030 was the strongest associated variant, gene-

based association tests suggest that additional PTVs and missense variants in *ALDH1L1*

also alter glycine levels (P=$1.4\times10^{-20}$, **Extended Data Fig. 7**, **Supplementary Table 9**).

*Ketone bodies.* A predicted damaging missense variant (rs201013770, p.Phe517Ser) in

*ACSS1* is associated in the combined analysis with mean increased serum acetate level of

0.005 mmol/l but is not observed in NFE (**Table 2**). *ACSS1* encodes an acyl-coenzyme A

synthetase and plays a role in the conversion of acetate to acetyl-CoA. In rodents,

increased acetate levels lead to obesity, insulin resistance, and metabolic syndrome,

mediated by activation of the parasympathetic nervous system[44].

**Associated variants and disease endpoints**

Newly available GWAS data from the FinnGen project[45] enabled us to test the hypothesis

that deleterious variants responsible for our novel quantitative trait associations (**Table 2**)

could also contribute to disease endpoints related to these traits. FinnGen has particularly

rich data on such endpoints as the samples are largely drawn from Finnish hospital

biobanks. In total, we examined 22 disease endpoint phenotypes for all 25 available

variants in **Table 2**. Three variants were associated with disease endpoints in FinnGen at

a Bonferroni-corrected threshold of $P<0.05/(22\times25)=9.0\times10^{-5}$ (**Supplementary Table**

**12**).

A predicted damaging missense variant (17:39135270:A/G; p.Ser32Pro) in *KRT40* which

is not observed in NFE and associated in FinMetSeq with a mean elevation in HDL-C of

1.07 mmol/l (**Table 2**), is associated in FinnGen with increased risk for pancreatitis.

While this is the first disease association reported for this gene, the type I keratin family,

of which *KRT40* is a member, is believed to play an important role in regulating exocrine

pancreas homoeostasis[46]. A 29 bp deletion on chromosome 1 causes a frameshift in

*FAM151A* which is 6.7X more frequent in FinMetSeq than NFE and associated in

FinMetSeq with both decreased total cholesterol in IDL and decreased IDL particle

concentration (**Table 2**), is associated in FinnGen with decreased risk of myocardial

infarction. The interpretation of this association is complicated by the fact that the variant

is also present in an overlapping transcript (*ACOT11*), a gene that plays a role in fatty

acid metabolism and lies <1 MB from a well-known cardioprotective variant in *PCSK9*.

Finally, a predicted damaging missense variant (rs77273740; p.Arg65Trp) in *DBH* that is

23.8X more frequent in FinMetSeq than in NFE and is associated with a mean decrease

of 1 mmHg in diastolic blood pressure in our combined analysis (**Table 2**), is associated

in FinnGen with decreased risk for hypertension. Distinct loci in this gene have

previously been shown with mean arterial pressure and this variant was included in a

gene-based association with mean arterial pressure[5,6].

**Replication outside of Finland: UK Biobank**

To begin to assess the generalizability outside of Finland of the novel associations that

we detected, we attempted to replicate associations from our combined Finnish analyses

in the UK Biobank (UKBB), a European sample that is approximately ten-fold larger.

Across eight anthropometric and blood pressure traits for which UKBB data are publicly

available, our Finnish combined analysis had identified 31 trait-variant associations

reaching $P<5 \times 10^{-7}$. More than a quarter of these variants (8 of 31) were not present in the

UKBB database. Of the remaining 23 associations, 20 were to variants that were common

in FinMetSeq (MAF> 1%) and had a comparable frequency in UKBB; 15 (75%) of these

variants showed association in UKBB at $P<0.05/23=2.2 \times 10^{-3}$ (Bonferroni correction for

23 tests). Of the three rare variants in this analysis, all of which were enriched at >10x

frequency in FinMetSeq compared to UKBB, none showed association in UKBB

(**Supplementary Table 13**). Even after adjusting for winner's curse[47] and with a sample

size of 340,000-360,000, we had <50% power to detect all three of these associations in

UKBB (**Supplementary Table 13**). This comparison supports the argument that

extremely large samples will be needed in most other populations to achieve the power

for rare variant association studies that we have observed in Finland.

**Geographical clustering of associated variants**

Given the concentration within sub-regions of northern and eastern Finland of most

Finnish Disease Heritage mutations[48], we hypothesized that the distribution of rare trait-

associated variants discovered through FinMetSeq might also display geographical

clustering. In support of this hypothesis, principal component analysis revealed broad-

scale population structure within the late-settlement region among 14,874 unrelated

FinMetSeq participants whose parental birthplaces are known (**Extended Data Fig. 8**).

Consistent with our hypothesis, parental birthplaces were significantly more

geographically clustered for carriers of PTVs and missense alleles than for carriers of

synonymous alleles, even after adjusting for MAC (**Supplementary Tables 14A, 14B**).

To enable finer scale analysis of the distribution of variants within late-settlement

Finland, we delineated geographically distinct population clusters using patterns of

haplotype sharing among 2,644 unrelated individuals with both parents known to be born

in the same municipality (Methods, **Extended Data Fig. 9**). Taking the cluster that is

most genetically similar to early-settlement Finland as a reference, we compared variant

counts for different functional classes and frequencies between this reference cluster and

each of the other 12 clusters that contained ≥100 individuals (**Fig. 4, Supplementary**

**Tables 15, 16**). In the two clusters that represent the most heavily bottlenecked late-

settlement regions (Lapland and Northern Ostrobothnia), we observed a marked deficit of

singletons and significant enrichment of variants at intermediate frequency compared to

other clusters. This pattern is most significant for missense variants, which are present in

the exome in large numbers; PTVs show consistently greater enrichment, but with less

statistical significance likely due to very small counts (**Fig. 4**). Two clusters in which we

observed marked enrichment of singletons, Surrendered Karelia and South Ostrobothnia,

showed the highest levels of genetic diversity across the frequency spectrum, likely

reflecting relatively recent gene flow into these regions from neighboring countries

(Russia and Sweden, respectively, **Fig. 4**).

We observed particularly strong geographical clustering among variants >10X enriched

in FinMetSeq compared to NFE (**Fig. 5A, Extended Data Fig. 10, Supplementary**

**Table 17**). We further characterized geographical clustering for FinMetSeq-enriched

trait-associated variants, by comparing the mean distances between birthplaces for

parents of minor allele carriers to those of non-carriers (**Supplementary Table 18**). Most

such variants were highly localized. For example, for variant rs780671030 in *ALDH1L1,*

which may be unique to Finns, the mean distance between parental birthplaces is 135 km

for carriers and 250 km for non-carriers ($P<1.0\times10^{-7}$, **Fig. 5B**). In contrast, few of the

variants that displayed sub-threshold association in FinMetSeq but that showed

significant associations in the combined analysis were significantly geographically

clustered within Finland (**Supplementary Table 18**).

Finally, we compared the geographic clustering of FinMetSeq-enriched trait-associated

variants to that of 35 Finnish Disease Heritage mutations carried by ≥3 FinMetSeq

individuals with known parental birthplaces. FinMetSeq carriers of monogenic Finnish

Disease Heritage mutations and FinMetSeq carriers of trait-associated variants identified

in FinMetSeq displayed a comparable degree of geographic clustering.  This clustering

was dramatically greater than that observed for the non-carriers of both sets of variants

(**Fig. 5C**), suggesting that rare variants associated with complex traits may be much more

unevenly distributed geographically than has been appreciated to date.

**DISCUSSION**

We have demonstrated that a well-powered exome sequencing study of deeply

phenotyped individuals can identify numerous rare variants associated with medically

relevant quantitative traits. The variants that we identified may provide a useful starting

point for studies aimed at uncovering biological mechanisms or fostering clinical

translation. For example, further investigation of the p.Arg722X variant in *ALDH1L1*

associated with reduced serum glycine could help elucidate the role of this gene in

astrocyte function, a topic of growing interest in neurobiology. Glycine is a key

inhibitory neurotransmitter localized to astrocytes[49], while the mouse ortholog, *Aldh1l1,*

is the primary marker for astrocytes in experimental research, since it is strongly

expressed in astrocytes, but not in neurons[50].

The substantial power of this study for discovering rare variant associations derives from

the occurrence, in the recently expanded and heavily bottlenecked populations of

northern and eastern Finland, of a large pool of deleterious variants that appear unique to

Finland or at frequencies orders of magnitude greater than in NFE. This observation

motivates a strategy for scaling up the discovery of rare variant associations by

prioritizing the sequencing of populations beyond Finland that have expanded in isolation

from recent bottlenecks. Examples of other such populations include those of

Ashkenazim[51], Iceland[52], Quebec[53], highland regions of Latin America[54], and

geographically isolated regions of larger European countries such as Sardinia[55] and

Crete[10]. In each of these populations, genetic drift has almost certainly caused a different

set of alleles to pass through the corresponding population-specific bottlenecks, enriching

some variants while depleting others. The numerous rare-variant associations that could

be identified by sequencing available, phenotyped samples across multiple population

isolates could rapidly increase our understanding of the genetic architecture of complex

traits. One caveat is that the extended LD blocks that are typical in such populations may

make it difficult to identify the causative variant from among multiple deleterious

variants within an association region[56].

Recent studies have suggested a continuity between the genetic architectures of complex

traits and disorders classically considered monogenic[57,58]. Our results offer strong support

for this continuity, not only in identifying numerous deleterious variants with large

effects on quantitative traits, but in demonstrating that such variants show geographical

clustering comparable to that of the mutations responsible for the Finnish Disease

Heritage.

The use of a Finland-specific genotype reference panel[59] to impute FinMetSeq variants

into array-genotyped samples from three other Finnish cohorts enabled us to identify

many additional novel associations. This result suggests that using sequence data from a

subset of individuals in each population to impute variants in array-genotyped samples

from the same population is a cost-effective strategy for detecting rare-variant

associations. However, the clustering in FinMetSeq of deleterious trait-associated

variants within limited geographical regions and our inability to follow-up >700 sub-

threshold associations from FinMetSeq for which the associated variants were not present

in the Finnish imputation reference panel, emphasize the importance of extensively

representing regional subpopulations when designing such reference panels, to account

for fine-scale population structure.

To fully realize the value of large-scale sequencing studies in population isolates, it will

be necessary to increase the richness of phenotypes available in sequenced cohorts from

these populations. For example, we associated <100 of the >24,000 deleterious, highly

enriched variants identified in FinMetSeq with one of the 64 cardiometabolic-related

quantitative traits studied here. In Finland, the national health care system and the

population's willingness to participate in biomedical research mean that extensive

medical records and population registries are available for mining additional phenotype

data, and create an opportunity for callback by genotype for further phenotyping and

collection of biological samples[60]. Notably, the associations we identified to disease

endpoints in FinnGen give a hint of the discoveries that will be possible when that

database reaches its full size of 500,000 participants. The insights gained from such

efforts will accelerate the implementation of precision health, informing projects in

larger, more heterogeneous populations which are still at an early stage[61].

**References**

1    MacArthur, J. *et al.* The new NHGRI-EBI Catalog of published genome-wide
     association studies (GWAS Catalog). *Nucleic Acids Res* **45**, D896-D901,
     doi:10.1093/nar/gkw1133 (2017).

2    Manolio, T. A. *et al.* Finding the missing heritability of complex diseases. *Nature*
     **461**, 747-753, doi:10.1038/nature08494 (2009).

3    Mahajan, A. *et al.* Refining the accuracy of validated target identification through
     coding variant fine-mapping in type 2 diabetes. *Nature genetics* **50**, 559-571,
     doi:10.1038/s41588-018-0084-1 (2018).

4    Turcot, V. *et al.* Protein-altering variants associated with body mass index
     implicate pathways that control energy intake and expenditure in obesity. *Nature
     genetics* **50**, 26-41, doi:10.1038/s41588-017-0011-x (2018).

5    Liu, C. *et al.* Meta-analysis identifies common and rare variants influencing blood
     pressure and overlapping with metabolic trait loci. *Nature genetics* **48**, 1162-
     1170, doi:10.1038/ng.3660 (2016).

6    Surendran, P. *et al.* Trans-ancestry meta-analyses identify rare and common
     variants associated with blood pressure and hypertension. *Nature genetics* **48**,
     1151-1161, doi:10.1038/ng.3654 (2016).

7    Zuk, O. *et al.* Searching for missing heritability: designing rare variant association
     studies. *Proc Natl Acad Sci U S A* **111**, E455-464, doi:10.1073/pnas.1322563111
     (2014).

8    Xue, Y. *et al.* Enrichment of low-frequency functional variants revealed by
     whole-genome sequencing of multiple isolated European populations. *Nature
     communications* **8**, 15927, doi:10.1038/ncomms15927 (2017).

9    Ganna, A. *et al.* Quantifying the Impact of Rare and Ultra-rare Coding Variation
     across the Phenotypic Spectrum. *American journal of human genetics* **102**, 1204-
     1211, doi:10.1016/j.ajhg.2018.05.002 (2018).

10   Southam, L. *et al.* Whole genome sequencing and imputation in isolated
     populations identify genetic associations with medically-relevant complex traits.
     *Nature communications* **8**, 15606, doi:10.1038/ncomms15606 (2017).

11   Jakkula, E. *et al.* The genome-wide patterns of variation expose significant
     substructure in a founder population. *Am J Hum Genet* **83**, 787-794,
     doi:10.1016/j.ajhg.2008.11.005 (2008).

12   Polvi, A. *et al.* The Finnish disease heritage database (FinDis) update-a database
     for the genes mutated in the Finnish disease heritage brought to the next-
     generation sequencing era. *Hum Mutat* **34**, 1458-1466, doi:10.1002/humu.22389
     (2013).

13   Manning, A. *et al.* A Low-Frequency Inactivating AKT2 Variant Enriched in the
     Finnish Population Is Associated With Fasting Insulin Levels and Type 2
     Diabetes Risk. *Diabetes* **66**, 2019-2032, doi:10.2337/db16-1329 (2017).

14   Lim, E. T. *et al.* Distribution and medical impact of loss-of-function variants in
     the Finnish founder population. *PLoS genetics* **10**, e1004494,
     doi:10.1371/journal.pgen.1004494 (2014).

15   Service, S. K. *et al.* Re-sequencing expands our understanding of the phenotypic
     impact of variants at GWAS loci. *PLoS genetics* **10**, e1004147,
     doi:10.1371/journal.pgen.1004147 (2014).

16   Wurtz, P. *et al.* Metabolite profiling and cardiovascular event risk: a prospective
study of 3 population-based cohorts. *Circulation* **131**, 774-785,
doi:10.1161/CIRCULATIONAHA.114.013116 (2015).
17   Laakso, M. *et al.* The Metabolic Syndrome in Men study: a resource for studies of
metabolic and cardiovascular diseases. *Journal of lipid research* **58**, 481-493,
doi:10.1194/jlr.O072629 (2017).
18   Huyton, T., Gottmann, W., Bade-Doding, C., Paine, A. & Blasczyk, R. The T/NK
cell co-stimulatory molecule SECTM1 is an IFN "early response gene" that is
negatively regulated by LPS in human monocytic cells. *Biochim Biophys Acta*
**1810**, 1294-1301, doi:10.1016/j.bbagen.2011.06.020 (2011).
19   Pirillo, A., Catapano, A. L. & Norata, G. D. HDL in infectious diseases and
sepsis. *Handb Exp Pharmacol* **224**, 483-508, doi:10.1007/978-3-319-09665-0_15
(2015).
20   Borodulin, K. *et al.* Forty-year trends in cardiovascular risk factors in Finland.
*Eur J Public Health* **25**, 539-546, doi:10.1093/eurpub/cku174 (2015).
21   Abraham, G. *et al.* Genomic prediction of coronary heart disease. *Eur Heart J* **37**,
3267-3278, doi:10.1093/eurheartj/ehw450 (2016).
22   Sabatti, C. *et al.* Genome-wide association analysis of metabolic traits in a birth
cohort from a founder population. *Nature genetics* **41**, 35-46, doi:10.1038/ng.271
(2009).
23   Pulizzi, N. *et al.* Interaction between prenatal growth and high-risk genotypes in
the development of type 2 diabetes. *Diabetologia* **52**, 825-829,
doi:10.1007/s00125-009-1291-1 (2009).
24   Locke, A. E. *et al.* Genetic studies of body mass index yield new insights for
obesity biology. *Nature* **518**, 197-206, doi:10.1038/nature14177 (2015).
25   Shungin, D. *et al.* New genetic loci link adipose and insulin biology to body fat
distribution. *Nature* **518**, 187-196, doi:10.1038/nature14132 (2015).
26   Wood, A. R. *et al.* Defining the role of common variation in the genomic and
biological architecture of adult human height. *Nature genetics* **46**, 1173-1186,
doi:10.1038/ng.3097 (2014).
27   Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and
body mass index in approximately 700000 individuals of European ancestry. *Hum*
*Mol Genet*, doi:10.1093/hmg/ddy271 (2018).
28   Pulit, S. L. *et al.* Meta-analysis of genome-wide association studies for body fat
distribution in 694,649 individuals of European ancestry. *Hum Mol Genet*,
doi:10.1093/hmg/ddy327 (2018).
29   Fagerberg, L. *et al.* Analysis of the human tissue-specific expression by genome-
wide integration of transcriptomics and antibody-based proteomics. *Mol Cell*
*Proteomics* **13**, 397-406, doi:10.1074/mcp.M113.035600 (2014).
30   Corsetti, J. P. *et al.* Thrombospondin-4 polymorphism (A387P) predicts
cardiovascular risk in postinfarction patients with high HDL cholesterol and C-
reactive protein levels. *Thromb Haemost* **106**, 1170-1178, doi:10.1160/TH11-03-
0206 (2011).
31   Cui, J. *et al.* Thrombospondin-4 1186G>C (A387P) is a sex-dependent risk factor
for myocardial infarction: a large replication study with increased sample size

from the same population. *Am Heart J* **152**, 543 e541-545,
doi:10.1016/j.ahj.2006.06.002 (2006).

32    Wessel, J., Topol, E. J., Ji, M., Meyer, J. & McCarthy, J. J. Replication of the
association between the thrombospondin-4 A387P polymorphism and myocardial
infarction. *Am Heart J* **147**, 905-909, doi:10.1016/j.ahj.2003.12.013 (2004).

33    Zhang, X. J. *et al.* Association between single nucleotide polymorphisms in
thrombospondins genes and coronary artery disease: A meta-analysis. *Thromb*
*Res* **136**, 45-51, doi:10.1016/j.thromres.2015.04.019 (2015).

34    Beygo, J. *et al.* New insights into the imprinted MEG8-DMR in 14q32 and
clinical and molecular description of novel patients with Temple syndrome. *Eur J*
*Hum Genet* **25**, 935-945, doi:10.1038/ejhg.2017.91 (2017).

35    Prats-Puig, A. *et al.* The placental imprinted DLK1-DIO3 domain: a new link to
prenatal and postnatal growth in humans. *Am J Obstet Gynecol* **217**, 350 e351-350
e313, doi:10.1016/j.ajog.2017.05.002 (2017).

36    Rosenfeld, J. A. *et al.* Clinical features associated with copy number variations of
the 14q32 imprinted gene cluster. *Am J Med Genet A* **167A**, 345-353 (2015).

37    Wallace, C. *et al.* The imprinted DLK1-MEG3 gene region on chromosome
14q32.2 alters susceptibility to type 1 diabetes. *Nature genetics* **42**, 68-71,
doi:10.1038/ng.493 (2010).

38    Day, F. R. *et al.* Genomic analyses identify hundreds of variants associated with
age at menarche and support a role for puberty timing in cancer risk. *Nature*
*genetics* **49**, 834-841, doi:10.1038/ng.3841 (2017).

39    Perry, J. R. *et al.* Parent-of-origin-specific allelic associations among 106
genomic loci for age at menarche. *Nature* **514**, 92-97, doi:10.1038/nature13545
(2014).

40    Cleaton, M. A. *et al.* Fetus-derived DLK1 is required for maternal metabolic
adaptations to pregnancy and is associated with fetal growth restriction. *Nature*
*genetics* **48**, 1473-1480, doi:10.1038/ng.3699 (2016).

41    Chaves, J. A. *et al.* Genomic variation at the tips of the adaptive radiation of
Darwin's finches. *Mol Ecol* **25**, 5282-5295, doi:10.1111/mec.13743 (2016).

42    Ding, Y. *et al.* Plasma Glycine and Risk of Acute Myocardial Infarction in
Patients With Suspected Stable Angina Pectoris. *J Am Heart Assoc* **5**,
doi:10.1161/JAHA.115.002621 (2015).

43    Wittemans, L. B. L. *et al.* Assessing the causal association of glycine with risk of
cardio-metabolic diseases. *Nature communications* **10**, 1060, doi:10.1038/s41467-
019-08936-1 (2019).

44    Perry, R. J. *et al.* Acetate mediates a microbiome-brain-beta-cell axis to promote
metabolic syndrome. *Nature* **534**, 213-217, doi:10.1038/nature18309 (2016).

45    Tabbassum, R. *et al.* Genetics of human plasma lipidome: Understanding lipid
metabolism and its link to diseases beyond traditional lipids. *bioRxiv*,
doi:10.1101/457960 (2018).

46    Casanova, M. L. *et al.* Exocrine pancreatic disorders in transsgenic mice
expressing human keratin 8. *J Clin Invest* **103**, 1587-1595, doi:10.1172/JCI5343
(1999).

47 Palmer, C. & Pe'er, I. Statistical correction of the Winner's Curse explains
replication variability in quantitative trait genome-wide association studies. *PLoS*
*genetics* **13**, e1006916, doi:10.1371/journal.pgen.1006916 (2017).
48 Norio, R. Finnish Disease Heritage I: characteristics, causes, background. *Hum*
*Genet* **112**, 441-456, doi:10.1007/s00439-002-0875-3 (2003).
49 Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**,
999-1014 e1022, doi:10.1016/j.cell.2018.06.021 (2018).
50 Chai, H. *et al.* Neural Circuit-Specialized Astrocytes: Transcriptomic, Proteomic,
Morphological, and Functional Evidence. *Neuron* **95**, 531-549 e539,
doi:10.1016/j.neuron.2017.06.029 (2017).
51 Rivas, M. A. *et al.* Insights into the genetic epidemiology of Crohn's and rare
diseases in the Ashkenazi Jewish population. *PLoS genetics* **14**, e1007329,
doi:10.1371/journal.pgen.1007329 (2018).
52 Gudbjartsson, D. F. *et al.* Large-scale whole-genome sequencing of the Icelandic
population. *Nature genetics* **47**, 435-444, doi:10.1038/ng.3247 (2015).
53 Peischl, S. *et al.* Relaxed Selection During a Recent Human Expansion. *Genetics*
**208**, 763-777, doi:10.1534/genetics.117.300551 (2018).
54 Service, S. *et al.* Magnitude and distribution of linkage disequilibrium in
population isolates and implications for genome-wide association studies. *Nature*
*genetics* **38**, 556-560, doi:10.1038/ng1770 (2006).
55 Chiang, C. W. K. *et al.* Genomic history of the Sardinian population. *Nature*
*genetics*, doi:10.1038/s41588-018-0215-8 (2018).
56 Peltonen, L., Palotie, A. & Lange, K. Use of population isolates for mapping
complex traits. *Nature reviews. Genetics* **1**, 182-190, doi:10.1038/35042049
(2000).
57 Bastarache, L. *et al.* Phenotype risk scores identify patients with unrecognized
Mendelian disease patterns. *Science* **359**, 1233-1239, doi:10.1126/science.aal4043
(2018).
58 Niemi, M. E. K. *et al.* Common genetic variants contribute to risk of rare severe
neurodevelopmental disorders. *Nature*, doi:10.1038/s41586-018-0566-4 (2018).
59 Surakka, I. S., A.-P.; Ruotsalainen, S.E.; Durbin, R.; Salomaa, V.; Daly, M.;
Palotie, A.; Ripatti, S. The rate of false polymorphisms introduced when imputing
genotypes from global imputation panels. *bioRxiv*, doi:10.1101/080770 (2016).
60 Latva-Rasku, A. *et al.* A Partial Loss-of-Function Variant in AKT2 Is Associated
With Reduced Insulin-Mediated Glucose Uptake in Multiple Insulin-Sensitive
Tissues: A Genotype-Based Callback Positron Emission Tomography Study.
*Diabetes* **67**, 334-342, doi:10.2337/db17-1142 (2018).
61 Collins, F. S. & Varmus, H. A new initiative on precision medicine. *N Engl J Med*
**372**, 793-795, doi:10.1056/NEJMp1500523 (2015).
714

**Acknowledgements**

Thanks to Terri Teshiba for coordinating ethical permissions and samples. Thanks to Sini

Kerminen, Daniel Lawson, and George Busby for discussions and providing scripts to

run fineSTRUCTURE. SR was supported by the Academy of Finland Center of

Excellence in Complex Disease Genetics (Grant No 312062), Academy of Finland (No.

285380), the Finnish Foundation for Cardiovascular Research, the Sigrid Juselius

Foundation, Biocentrum Helsinki and University of Helsinki HiLIFE Fellow grant. VR

acknowledges support by RFBR, research project No. 18-04-00789 A. VS was supported

by the Finnish Foundation for Cardiovascular Research. CS and LS received funding

from HG006695, HL113315, and MH105578. MAK is supported by a Senior Research

Fellowship from the National Health and Medical Research Council (NHMRC) of

Australia (APP1158958). He also works in a unit that is supported by the University of

Bristol and UK Medical Research Council (MC_UU_12013/1). The Baker Institute is

supported in part by the Victorian Government's Operational Infrastructure Support

Program. AUJ, DR, LJS, HMS, RW, PY, XY, and MB received funding from

DK062370. SKS, CWKC, and NBF received funding from HL113315 and NS062691.

The METSIM study was supported by grants from Academy of Finland (No. 321428),

the Sigrid Juselius Foundation, the Finnish Foundation for Cardiovascular Research,

Kuopio University Hospital, and Centre of Excellence of Cardiovascular and Metabolic

Diseases supported by the Academy of Finland (ML). Sequencing was funded by

5U54HG003079, and AEL, KMS, HJB, CCC, CJK, KLK, DCK, DEL, JN, TJN, SKD,

NOS, IMH, and RKW were funded by 5U54HG003079 and 5UM1HG008853-03.

**Author Contributions**

AEL, LJS, RKW, AaP, VS, ML, SR, MB, and NBF designed the study. AEL, KMS,

HJA, RSF, DCK, DEL, JN, TJN, and JV produced and quality-controlled the sequence

data. AEL, AUJ, ArP, HMS, MAK, VS, and ML produced and quality-controlled the

clinical data. AEL, KMS, CWKC, SKS, ASH, LS, MP, CCC, AUJ, CJK, KK, VR, DR,

JV, RW, PY, and XY analyzed data. JGE, MAK, MRJ, and MM provided replication

data. HL, SKD, NOS, IMH, CS, SR, MB, and NBF supervised experiments and analyses.

AEL, KMS, CWKC, SKS, CS, MB and NBF wrote the paper. AEL, KMS, CWKC, and

SKS contributed equally to this work. NBF and MB jointly supervised this work.

Competing interests statements:

VS has participated in a conference trip sponsored by Novo Nordisk and received a

honorarium from the same source for participating in an advisory board meeting. He also

has ongoing research collaboration with Bayer Ltd.

HL is a member of the Nordic Expert group unconditionally supported by Gedeon

Richter Nordics and has received an honorarium from Orion.

Correspondence and requests for materials should be addressed to

nfreimer@mednet.ucla.edu or boehnke@umich.edu.

Data Availability: The sequence data can be accessed through dbGaP using the following

study numbers: FINRISK: phs000756, METSIM: phs000752. Association results can be

accessed at http://pheweb.sph.umich.edu/FinMetSeq/.

**Table 1.** Sequence variants identified using whole-exome sequencing of 19,292 FinMetSeq participants. Percentages are the percent of all variants in the given category to either have MAF <1% or to be singleton variants.

| Variant type | All variants | MAF<1% | Singleton variants |
|---|---|---|---|
| SNV | 1,318,781 | 87.5% | 40.5% |
| Insertion/Deletion | 92,776 | 87.0% | 43.1% |
| Predicted LoF | 33,156 | 96.4% | 55.0% |
| Non-synonymous | 353,228 | 92.2% | 46.4% |

| Variant Annotation | All variants | MAF<1% | Singleton variants |
|---|---|---|---|
| Splice Acceptor | 3,180 | 95.4% | 50.8% |
| Splice Donor | 3,795 | 96.2% | 53.3% |
| Stop Gain | 11,382 | 97.3% | 54.3% |
| Frameshift | 12,845 | 96.6% | 58.2% |
| Stop Loss | 621 | 88.1% | 48.1% |
| Initiator Codon/Start Loss | 1,333 | 93.6% | 49.1% |
| Inframe Insertion | 1,673 | 90.3% | 44.5% |
| Inframe Deletion | 4,936 | 92.9% | 46.8% |
| Missense | 353,228 | 92.3% | 46.4% |
| Splice Region | 40,248 | 87.1% | 41.2% |
| Incomplete Terminal Codon | 16 | 81.3% | 50.0% |
| Stop Retained | 217 | 86.2% | 42.4% |
| Synonymous | 180,104 | 85.7% | 40.0% |
| Coding Sequence | 78 | 88.5% | 41.0% |
| Mature miRNA | 239 | 92.9% | 48.5% |
| 5' UTR | 35,572 | 87.8% | 38.2% |
| 3' UTR | 66,539 | 86.2% | 38.6% |
| Non-coding Exonic | 82,126 | 85.8% | 37.8% |
| Intronic | 601,362 | 85.1% | 37.4% |
| Upstream | 8,820 | 86.5% | 38.3% |
| Downstream | 3,050 | 84.6% | 38.3% |
| Intergenic | 193 | 85.5% | 31.1% |

Variant annotation refers to the "most deleterious" annotation for a given variant across all Ensembl (v88) transcripts, following the order defined by VEP (https://useast.ensembl.org/info/genome/variation/prediction/predicted_data.html).

**Table 2**. Associations with predicted deleterious variants that conditional analysis and literature review suggest are novel. These associations reach exome-wide significance in FinMetSeq alone or in a combined analysis of FinMetSeq with three replication cohorts.

| Chr:Pos (GRCh37) | Gene | Anno: Prediction^ | FMS MAF | NFE MAF# | MAF Ratio (95% CI) | Trait | FMS P | FMS Beta | Repl. or combined P** | Repl. or combined Beta | Mean in carriers \| non-carriers |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1:55076137 | FAM151A | FS:PTV | .099 | .0147 | 6.7 (5.6-7.8) | Total Chol. in IDL | $5.4\times10^{-16}$ | -0.187 | **$2.1\times10^{-17}$** | -0.191 | .84 \| .87 mmol/l |
| | | | | | | IDL Particle Conc. | $8.9\times10^{-14}$ | -0.172 | **$1.9\times10^{-16}$** | -0.185 | .130 \| .134 umol/l |
| 2:120848049 | EPB41L5 | MS:T;B;B;N;D | .085 | .044 | 1.9 (0.9-3) | eGFR* | $1.7\times10^{-6}$ | -0.093 | $4.8\times10^{-12}$ | -0.107 | 88.6 \| 89.9 SI |
| | | | | | | Creatinine* | $2.5\times10^{-6}$ | 0.091 | $2.5\times10^{-12}$ | 0.098 | 81.62 \| 80.64 umol/l |
| 3:125831672 | ALDH1L1 | SG:PTV | .0026 | 0 | ∞ | Glycine | $1.8\times10^{-8}$ | -0.873 | **$4.5\times10^{-4}$** | -0.827 | .24 \| .27 mmol/l |
| 4:13612630 | BOD1L1 | MS:D;D;D;N;D | .0001 | 0 | ∞ | WHR adj. BMI | $4.7\times10^{-7}$ | -2.501 | NA | NA | .88 \| .93 |
| 5:79336091 | THBS4 | MS:D;D;D;D;D | .0045 | .0001 | 45 (41.9-48.1) | Weight* | $6.7\times10^{-7}$ | -0.377 | $3.2\times10^{-7}$ | -0.252 | 74.6 \| 80.5 kg |
| 5:140181423 | PCDHA3 | FS:PTV | .0001 | NA | NA | WHR adj. BMI | $2.7\times10^{-7}$ | 2.559 | NA | NA | 1.14 \| .93 |
| 9:107548661 | ABCA1 | MS:D;D;D;D;D | .00023 | 0 | ∞ | Serum HDL Chol. | $4.8\times10^{-10}$ | -2.046 | NA | NA | .80 \| 1.44 mmol/l |
| 9:136501728 | DBH | MS:D;D;P;N;N | .05 | .0021 | 23.8 (22.4-25) | Diastolic BP* | $1.5\times10^{-6}$ | -0.115 | $2.8\times10^{-12}$ | -0.11 | 83.1 \| 84.1 mmHg |
| 11:47282929 | NR1H3 | MS:D;P;P;D;D | .0042 | .00003 | 140 (132.8-147.2) | Serum HDL Chol. | $1.4\times10^{-7}$ | 0.425 | **$6.7\times10^{-7}$** | 0.435 | 1.59 \| 1.44 mmol/l |
| | | | | | | HDL2 Chol.* | $3.2\times10^{-6}$ | 0.473 | $1.3\times10^{-8}$ | 0.458 | 1.07 \| .92 mmol/l |
| | | | | | | VLDL Chol.* | $4.0\times10^{-6}$ | -0.469 | $3.1\times10^{-7}$ | -0.412 | .75 \| .91 mmol/l |
| 11:116692293 | APOA4 | MS:T;D;P;N;N | .0096 | .012 | 0.8 (-0.4-2) | Serum HDL Chol.* | $2.2\times10^{-5}$ | 0.225 | $1.5\times10^{-7}$ | 0.196 | 1.51 \| 1.44 mmol/l |
| 11:117352857 | DSCAML1 | MS:T;B;B;.;D | .016 | .0002 | 80 (77.8-82.2) | VLDL Chol. | $4.1\times10^{-8}$ | 0.299 | **$2.0\times10^{-3}$** | 0.162 | 1.01 \| .90 mmol/l |
| 14:101198426 | DLK1 | MS:T;B;B;N;D | .023 | .00013 | 177 (174.3-179.6) | Height* | $2.7\times10^{-5}$ | -0.149 | $1.2\times10^{-10}$ | -0.163 | 170.7 \| 172.0 cm |
| 16:55862682 | CES1 | MS:D;D;D;D;D | .0018 | .00003 | 60 (52.8-67.2) | Serum HDL Chol. | $1.1\times10^{-10}$ | 0.771 | **$3.8\times10^{-6}$** | 0.793 | 1.77 \| 1.44 mmol/l |
| | | | | | | Serum ApoA1* | $1.9\times10^{-6}$ | 0.668 | $4.0\times10^{-9}$ | 0.718 | 1.65 \| 1.47 g/l |
| 16:56996009 | CETP | SD:PTV | .0017 | .00003 | 56.7 (49.4-63.9) | Serum ApoA1 | $2.6\times10^{-8}$ | 0.834 | **$1.8\times10^{-4}$** | 1.034 | 1.70 \| 1.47 g/l |
| | | | | | | Serum HDL Chol. | $1.1\times10^{-14}$ | 0.946 | **$8.8\times10^{-21}$** | 1.217 | 1.84 \| 1.44 mmol/l |
| 16:68013570 | DPEP3 | MS:T;B;B;N;D | .0099 | .00044 | 22.5 (20.8-24.2) | Serum HDL Chol. | $1.6\times10^{-7}$ | -0.295 | **$7.2\times10^{-15}$** | -0.373 | 1.33 \| 1.44 mmol/l |
| | | | | | | Serum ApoA1* | $5.2\times10^{-6}$ | -0.294 | $4.0\times10^{-7}$ | -0.253 | 1.40 \| 1.47 g/l |
| 16:68732169 | CDH3 | MS:D;D;D;D;D | .0044 | .00064 | 6.9 (5.2-8.5) | Pyruvate* | $3.7\times10^{-5}$ | 0.417 | $6.6\times10^{-10}$ | 0.471 | .08 \| .07 mmol/l |
| 17:6599157 | SLC13A5 | MS:D;D;D;D;D | .00091 | 0 | ∞ | Citrate | $1.3\times10^{-9}$ | 1.294 | **$9.5\times10^{-12}$** | 1.309 | .14 \| .11 mmol/l |
| 17:7129898 | DVL2 | MS:D;D;D;D;D | .02 | .02 | 1 (-0.2-2.1) | Valine* | $4.2\times10^{-5}$ | -0.239 | $5.7\times10^{-9}$ | -0.232 | .210 \| .217 mmol/l |
| 17:39135270 | KRT40 | MS:D;P;P;N;D | .00013 | 0 | ∞ | Serum HDL Chol. | $3.2\times10^{-8}$ | 2.416 | NA | NA | 2.51 \| 1.44 mmol/l |
| 17:41062979 | G6PC | MS:T;P;P;D;D | .025 | 0 | ∞ | Total MUFA | $4.4\times10^{-7}$ | 0.275 | $3.5\times10^{-1}$ | 0.067 | 3.88 \| 3.62 mmol/l |
| | | | | | | Glycerol* | $5.8\times10^{-6}$ | 0.218 | $4.1\times10^{-7}$ | 0.183 | .092 \| .088 mmol/l |
| | | | | | | Plasma CRP* | $1.6\times10^{-5}$ | 0.175 | $4.0\times10^{-9}$ | 0.185 | 2.47 \| 2.17 mg/l |
| | | | | | | Triglycerides* | $1.0\times10^{-6}$ | 0.23 | $1.3\times10^{-7}$ | 0.197 | 1.60 \| 1.46 mmol/l |
| 17:41926216 | CD300LG | MS:T;D;P;N;N | .00034 | 0 | ∞ | Serum HDL Chol. | $4.8\times10^{-14}$ | 2.061 | **$4.9\times10^{-2}$** | 0.801 | 2.39 \| 1.44 mmol/l |
| | | | | | | HDL2 Chol. | $1.3\times10^{-7}$ | 2.154 | NA | NA | 1.88 \| .92 mmol/l |
| | | | | | | Serum ApoA1 | $8.1\times10^{-8}$ | 1.694 | NA | NA | 2.04 \| 1.47 g/l |
| 18:47091686 | LIPG | SA:PTV | .0025 | 0 | ∞ | HDL2 Chol.* | $1.2\times10^{-5}$ | 0.579 | $5.6\times10^{-10}$ | 0.624 | 1.13 \| .92 mmol/l |
| | | | | | | Phosphocholines* | $3.1\times10^{-6}$ | 0.624 | $1.1\times10^{-8}$ | 0.578 | 2.44 \| 2.20 mmol/l |
| | | | | | | Phosphoglycerides* | $9.0\times10^{-6}$ | 0.594 | $1.1\times10^{-7}$ | 0.538 | 2.50 \| 2.25 mmol/l |

| Position | Gene | Annotation^ | MAF | NFE MAF# | MAF Ratio | Trait | P-value | Beta | Rep. P-value** | Rep. Beta | Means |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 19:10683762 | *AP1M2* | MS:D;D;D;D;D | .015 | .00009 | 167 (162.7-170.7) | Serum ApoB | $5.8\times10^{-8}$ | -0.282 | **$1.5\times10^{-3}$** | -0.199 | .96 \| 1.02 g/l |
| | | | | | | Total Chol. in IDL* | $1.1\times10^{-6}$ | -0.289 | $6.9\times10^{-14}$ | -0.319 | .81 \| .87 mmol/l |
| | | | | | | IDL Particle Conc.* | $2.1\times10^{-6}$ | -0.281 | $8.5\times10^{-14}$ | -0.318 | .125 \| .133 umol/l |
| | | | | | | Remnant Chol.* | $8.0\times10^{-6}$ | -0.268 | $2.7\times10^{-12}$ | -0.301 | 1.65 \| 1.77 mmol/l |
| 19:11350904 | *ANGPTL8* | SG:PTV | .0025 | 0 | ∞ | HDL2 Chol.* | $3.4\times10^{-6}$ | 0.564 | $1.1\times10^{-8}$ | 0.574 | 1.06 \| .92 mmol/l |
| 19:49318380 | *HSD17B14* | MS:D;D;D;D;D | .046 | .05 | 0.9 (-0.2-2) | Valine* | $3.4\times10^{-5}$ | -0.152 | $2.1\times10^{-7}$ | -0.144 | .212 \| .217 mmol/l |
| 20:24994201 | *ACSS1* | MS:D;D;D;D;D | .0026 | 0 | ∞ | Acetate* | $1.3\times10^{-5}$ | 0.626 | $2.1\times10^{-12}$ | 0.631 | .046 \| .041 mmol/l |

^Annotations are from VEP: FS=Frameshift; SG=Stop Gain; SD=Splice Donor; SA=Splice Acceptor; MS=Missense. All but MS are PTV. Predictions for missense variants are presented for five different prediction algorithms, each separated by semi-colon: SIFT (D=Damaging; T=Tolerated); PolyPhen2 - human diversity (D=Probably Damaging; P=Possibly Damaging; B=Benign); PolyPhen2 - hum variation (D=Probably Damaging; P=Possibly Damaging; B=Benign); Mutation Taster (A=Automatic Disease Causing; D=Disease Causing; N=Polymorphism; P=Automatic Polymorphism); and LRT (D=Deleterious; N=Neutral; U=Unknown).

# Non-Finnish European (NFE) MAF are taken from exomes of gnomAD v2.1 control individuals that were not from Estonia or Sweden. A variant with frequency 0 indicates that the variant was present in some subset of gnomAD, but was not found in NFE controls. NA indicates the variant was not present in gnomAD.

Minor Allele Frequency Ratio (MAF Ratio) is MAF in FinMetSeq/MAF in gnomAD NFE.

*Indicates an association only reaching significance in meta-analysis combining FinMetSeq and the three replication cohorts. If unlabeled, the association was significant in FinMetSeq alone.

** Replication P-values <0.05 are highlighted in bold.

**METHODS**

**METSIM and FINRISK studies: designs, phenotypes, and sequenced participants**

**METSIM** is a single-site study comprised of 10,197 men randomly selected from the

population register of Kuopio, Eastern Finland, aged 45 to 73 years at initial examination

from 2005 to 2010[17,62]. The goal of METSIM is to investigate genetic and non-genetic

factors associated with Type 2 Diabetes (T2D), cardiovascular disease (CVD), insulin

resistance, and related traits. The METSIM study protocol includes collection of data on

CVD history and risk factors, measurements of height, weight, waist, hip, blood pressure,

and collection of a blood sample for laboratory measurements and DNA extraction.

Diagnoses of myocardial infarction[63], stroke[64], and peripheral vascular disease were

verified based on medical records at baseline. We attempted exome sequencing of all

METSIM study participants.

**FINRISK** is a series of health examination surveys carried out by the National Institute

for Health and Welfare (formerly National Public Health Institute) of Finland every five

years beginning in 1972[65]. The surveys are based on random population samples from

five (six in 2002) geographical regions of Finland. Participants were selected by 10-year

age group, sex, and study area. Survey sample sizes have varied from 7,000 to 13,000

individuals and participation rates from 60% to 90%. The age-range was 25 to 64 years

until 1992 and 25 to 74 years since 1997. The survey includes a self-administered

questionnaire, a standardized clinical examination carried out by specifically trained

study nurses, and collection of a blood sample for laboratory measurements and DNA

extraction[66]. For exome sequencing, we chose 10,192 participants from the 1992, 1997,

2002, and 2007 FINRISK surveys from northeastern Finland (former provinces of North

Karelia, Oulu, and Lapland). This selection was based on the hypothesis that the rapid

growth in isolation of the populations of this region from severe bottlenecks in the 16[th]-

17[th] centuries might have resulted in deleterious variants rising to a much higher

frequency than in other populations.

METSIM participants fasted for more than 10 hours prior to blood draw. FINRISK

participants were instructed to fast for four hours before the scheduled examination and

to avoid heavy meals earlier in the day; duration of fasting was recorded. Laboratory

measurements in METSIM included an oral glucose tolerance test with samples at 0, 30,

and 120 minutes (only fasting measurements in known diabetics) for glucose, insulin,

proinsulin, and free fatty acids, as well as fasting laboratory measurements including

lipids, lipoproteins, apolipoproteins, adiponectin, and hs-CRP. Most of these

measurements were carried out in FINRISK non-fasting samples; two-hour oral glucose

tolerance tests with glucose and insulin measured at 0 and 120 minutes were carried out

in subsets of FINRISK 1992, 2002 and 2007 participants. Both studies include proton

NMR metabolomics measurements of lipoprotein subclasses, their lipid concentrations

and composition, apolipoprotein A-I and B, multiple cholesterol and triglyceride

measures, albumin, various fatty acids, and numerous low-molecular-weight metabolites,

including amino acids, glycolysis related measures and ketone bodies[67,68].

METSIM and FINRISK participants are followed up regularly via record linkage using

the Finnish health registries, allowing for near complete follow-up of multiple disease

diagnoses; participants may also be called back in person for additional studies.

Participants in both studies provided informed consent, and all study protocols were

approved by the Ethics Committees at the participating institutions (FINRISK cohorts

1992 & 1997: National Public Health Institute of Finland; FINRISK 2002, 2007, & 2012:

Ethical Review Board of the Hospital District of Helsinki and Uusimaa; METSIM:

Research Ethics Committee, Hospital District of Northern Savo IRB #1).

**Selection of traits, harmonization, exclusions, covariate adjustment, and**

**transformation**

Of the 257 quantitative metabolic and cardiovascular traits measured in both METSIM

and FINRISK, we selected 64 traits for association analysis in the current study based on

clinical relevance for cardiovascular and metabolic health (**Supplementary Tables 3, 4**).

*Exclusions*

We excluded 126 individuals, 92 with type 1 diabetes and 34 women who were pregnant

at the time of phenotyping, from all analyses, and 3,088 individuals with T2D from

analyses of glycemic traits. For traits influenced by food consumption (amino acids, fatty

acids, LDL cholesterol, total triglycerides, and glycemic traits), we excluded individuals

not fasting for at least 8 hours after their last meal. A complete list of exclusions can be

found in **Supplementary Table 4**.

*Trait adjustments*

For individuals on antihypertensive medication at the time of testing, we added 15 and 10

mmHg to the measured values of systolic and diastolic blood pressures, respectively[69,70].

For individuals on lipid regulating medications, we multiplied the measured total

cholesterol by 1.25 [71]. For FINRISK participants, we calculated LDL cholesterol (LDL-

C) levels using the Friedewald equation (LDL-C = adjusted total cholesterol – HDL-C –

(triglycerides/2.2)) and excluded individuals with elevated triglycerides (>2.5mmol/l);

LDL-C was measured directly in METSIM participants and for participants on lipid

regulating medication, values were divided by 0.7 [72]. All trait adjustments are listed in

**Supplementary Table 4**.

*Trait transformations and adjustment for covariates*

We prepared quantitative traits for association analysis separately for METSIM and

FINRISK participants by linear regression on trait-specific covariates; skewed variables

were log transformed prior to linear regression analysis. Both before and after log

transformation, we examined all variables for normality and for outliers. Log

transformation was adequate in all cases to approximate normality for initial covariate

adjustment. Outliers, of which there were no more than 5 individuals with values >5SD

for any trait prior to adjustment and inverse normalization, were not removed. Covariates

for these regression analyses always included covariates age and age$^2$ for METSIM and

sex, age, age$^2$, and cohort year for FINRISK. Trait transformations and trait-specific

covariates are listed in **Supplementary Table 4**. Several traits were adjusted for sex

hormone treatment, which included women on contraceptives or hormone replacement

therapy. We transformed residuals from these initial regression analyses to normality

using inverse normal scores.

**Exome sequencing**

We carried out exome sequencing in two phases.

Phase 1 We quantified the 10,379 Phase 1 DNA samples with the Quant-iT PicoGreen

dsDNA reagent on a Varioskan Microplate Reader (ThermoFisher Scientific) and

randomly parsed samples with adequate DNA (>250ng) into cohort specific files. We

then re-arrayed samples using the BioMicroLab XL20 (USA Scientific) to ensure equal

numbers of METSIM and FINRISK samples on each 96-well plate, alternating samples

between studies in consecutive positions within and across plates, to reduce opportunities

for between-study batch effects.

We constructed dual indexed libraries using 100-250ng of genomic DNA and the KAPA

HTP Library Kit (KAPA Biosystems) on the SciClone NGS instrument (Perkin Elmer).

The target insert size was 250 bp. We then pooled twelve libraries prior to hybridization

with the SeqCap EZ HGSC VCRome (Roche) reagent that targets 45.1 Mb (23,585 genes

and 189,028 non-overlapping exons) of the human genome. Each library pool contained

samples from both cohorts and was comprised of 300-400 ng of each individual library

for a total library input of 3.6-4.8 μg into the initial hybridization. We estimated the

concentration of each captured library pool by qPCR (Kapa Biosystems) to produce

appropriate cluster counts for the HiSeq2000 platform (Illumina). We then generated

2x100bp paired-end sequence data yielding ~6 Gb per sample to achieve a coverage

depth of ≥20x for ≥70% of targeted bases for every sample.

<u>Phase 2</u> We quantified, prepared, pooled, and captured the 9,937 Phase 2 samples just as

in Phase 1. Here we generated 2×125 bp paired-end sequencing reads on the HiSeq2500

1T to again achieve a coverage depth of ≥20x for ≥70% of targeted bases for every

sample.

*Contamination detection, sequence alignment, sample QC, and variant calling*

We aligned sequence reads to human genome reference build 37 using bwa-mem

(v0.7.7), marked duplicates with picard MarkDuplicates (v1.113;

http://broadinstitute.github.io/picard), and realigned indels with the GATK[73]

IndelRealigner (v2.4). We used BamUtil clipOverlap (v1.0.11;

http://genome.sph.umich.edu/wiki/BamUtil:_clipOverlap) to mark overlapping bases

from paired reads resulting from short insert fragments.

For each sample, we required ≥70% of target bases sequenced at ≥20x depth, and SNV

genotype array concordance >90% if SNV array data were available. We used

verifyBamID[74] (v1.1.1) to detect and exclude samples with estimated contamination

>3%. Where available, we also used existing genotype data with verifyBamID to detect

and exclude sample swaps. Of 20,316 samples attempted, 13 failed sequencing, 35 were

sample swaps, 760 either had low genotype concordance, sex mismatch, or estimated

contamination >3%, and four had discrepancies between reported and genotype-estimated

relationships (**Supplementary Table 1**).

We called SNVs and short indels using the recommended best practices for cohort-level

calling with GATK[73] (v3.3). For each individual, we called bases and identified variant

sites for all targeted exome bases and 500 bp of sequence up and downstream of each

target region using HaplotypeCaller, resulting in calling substantial numbers of non-

exonic variants. We merged these calls in batches of 200 individuals using

CombineGVCFs and recalled genotypes for all individuals at all variable sites with

GenotypeGVCFs.

After merging genotypes for the 19,378 samples that passed preliminary QC checks, we

filtered SNVs and indels separately using the recommended best practices for Variant

Quality Score Recalibration (VQSR). We used the set of true positive variants provided

in the GATK resource bundle (v2.5; build37) for training the VQSR model after

restricting to sites in targeted exome regions. After assessment with VQSR, we retained

variants for which we identified ≥99% of true positive sites used in the training model

(i.e. truth sensitivity) for both SNVs and indels.

Following initial variant filtering, we decomposed multi-allelic variants into bi-allelic

variants, left-aligned indels, and dropped redundant variants using vt[75] (version 0.5). We

filtered variants with >2% missing calls and/or Hardy-Weinberg p-value<$10^{-6}$. We

applied an additional filter removing variants with an overall allele balance (AB; alternate

AC/sum of total AC) <30% in genotyped samples. We then excluded 86 individuals with

>2% missing variant calls yielding a final analysis set of 19,292 individuals.

**Array genotypes, genotype imputation, and integrated exome+imputation panel**

METSIM participants were previously genotyped with the Illumina OmniExpress array; genotyping and quality control are described elsewhere[76]. FINRISK participants were previously genotyped in several batches on different arrays[21]. We lacked genotype array data for 1,488 participants (57 METSIM, 1,431 FINRISK). From the available genotype array data, we generated three datasets: 1) a merged array-based genotype call set of all variants present in ≥90% of array-genotyped individuals across both cohorts; 2) a merged Haplotype Reference Consortium (HRC) v1.1 imputed data set of the array-based genotypes; 3) an integrated data set containing genotyped and imputed array-based variants and exome sequence variants (HRC+exome). Where there was overlap between the sequence and imputed genotypes, we kept the sequence-based genotypes. We excluded the 1,488 individuals with no array data from the HRC+exome panel.

We prepared array genotypes for imputation using the Imputation Preparation and Checking tool (http://www.well.ox.ac.uk/~wrayner/tools/HRC-1000G-check-bim.v4.2.5.zip) and used the Michigan Imputation Server[77] (www.imputationserver.sph.umich.edu) to impute genotypes using the HRC (v1.1) reference panel[78]. METSIM samples were imputed in a single batch. FINRISK samples were imputed in batches based on the genotyping array and/or center where genotypes were generated.

**Annotation**

We annotated the final set of variants passing QC using Ensembl's variant effect

predictor (VEP v76)[79] and Combined Annotation-Dependent Depletion[80] (CADD v1.2).

We employed five *in silico* algorithms implemented in dbNSFP v2.4

(https://sites.google.com/site/jpopgen/dbNSFP) to predict the functional impact of

missense variants: PolyPhen2 HumDiv and HumVar[81], LRT[82], MutationTaster[83], and

SIFT[84].

**Association testing**

*Single variants*

We carried out single-variant association tests for transformed trait residuals with

genotype dosages for variants with MAC≥3 assuming an additive genetic model. We

used the EMMAX[85] linear mixed model approach, as implemented in EPACTS (v3.3.0;

http://genome.sph.umich.edu/wiki/EPACTS), to account for relatedness between

individuals. We used genotypes for sequenced variants with MAF≥1% to construct the

genetic relationship matrix (GRM).

*Conditioning on associated variants from prior GWAS*

To differentiate association signals identified in this study from known association

signals, for each trait we performed exome-wide association analysis conditioning on

variants previously associated ($P<10^{-7}$) with that trait. We compiled a list of known

variants for each trait from the EBI GWAS catalog

(https://www.ebi.ac.uk/gwas/downloads; December 4, 2016 version), from recent papers,

and from manuscripts in preparation of which we were aware[76,86-88]. The keywords from

the GWAS catalog we used to assign known variants to each of our traits are listed in

**Supplementary Table 19**. We also manually curated the associations from Inouye et

al.[89] and Kettunen et al.[86] to attribute "blood metabolite" associations to the specific

associated metabolites.

Using the combined HRC+exome panel (see above), we pruned each trait-specific list of

associated variants ("GWAS variants") based on linkage disequilibrium (LD) ($r^2$>0.95).

Twenty-three GWAS variants were not present in the HRC+exome panel. For 17 of these

23 variants, we identified a proxy ($r^2$>0.80) variant instead; we excluded the remaining

six variants from conditional analysis. The list of variants included in conditional analysis

for each trait is included in **Supplementary Table 20**. We extracted genotypes for

variants used in conditional analysis from the integrated HRC+exome panel and

converted dosages to alternate allele counts by rounding to the nearest integer (0, 1, or 2).

We imputed missing genotypes for the 1,488 individuals without array data using the

mean genotype dosage for purposes of conditional analysis.

For conditional analysis for each trait, we ran association analysis using the same linear

mixed model approach as in unconditional analysis but including the complete set of

pruned GWAS variants as covariates in the association test. Following conditional

association, we further determined novelty based on absence of the variant from OMIM

descriptions, ClinVar, and extensive literature searches.

*Defining loci*

The set of >1.4M variants tested for association includes variants in LD. To identify the

number of distinct associations for each trait, we performed LD clumping using Swiss

(https://github.com/welchr/swiss). We selected the subset of variants with (1)

unconditional $P<5\times10^{-7}$ or (2) both unconditional and conditional $P<5\times10^{-5}$ for at least

one trait. For each variant in this subset, we provided Swiss with the minimum

unconditional p-value across all traits. The clumping procedure starts with the variant

with the smallest p-value (index variant), and merges into one locus all variants within ±1

Mb that have $r^2>0.5$ with the index variant. The procedure is repeated iteratively until no

variants remain. Trait associations with variants in the same locus are considered to

represent the same signal and trait associations with variants in different loci to be

distinct signals.

1000

1001    *Calculating effects and variance explained of individual variants*

1002    For novel variants highlighted in **Table 2** we evaluated the effect of each variant on the

1003    trait values. We did this by calculating the mean trait value in carriers and non-carriers,

1004    assuming no homozygous carriers. Differences noted are the difference in the two means.

1005

1006    Given that the effect estimates from our association tests are standardized, we calculated

1007    variance explained for a given variant with the equation $2f(1-f)\hat{\beta}^2$, where f is the minor

1008    allele frequency and $\hat{\beta}$ is the estimated effect size. The variance explained is included in

1009    **Supplementary Table 8**.

1010

*Gene-based testing*

We carried out gene-based association tests using the mixed model implementation of

SKAT-O[90], which tests for the optimal mixture of burden and dispersion-style multi-

marker tests while adjusting for relatedness between individuals using the same GRM

calculated for the single-variant tests. EMMAX and the mixed model version of SKAT-O

(mmskat) are implemented in EPACTS.

1017

We implemented gene-based tests using three different, but nested, sets of variants

(variant "masks"):

(1) PTVs at any allele frequency with VEP annotations: frameshift_variant,

initiator_codon_variant, splice_acceptor_variant, splice_donor_variant, stop_lost,

stop_gained;

(2) PTVs included in (1) plus missense variants with MAF<0.1% scored as "damaging"

or "deleterious" by all five functional prediction algorithms;

(3) PTVs included in (1) plus missense variants with MAF<0.5% scored as "damaging"

or "deleterious" by all five functional prediction algorithms.

1027

For each trait and mask, we only tested genes with at least two qualifying variants. Each

mask contained a different number of genes with at least two qualifying variants: up to

7,996, 12,795, and 12,890 for the three masks, respectively. The exact number of genes

tested varied by trait due to sample size. We first used a Bonferroni-corrected exome-

wide threshold for 12,890 genes, which corresponds to a threshold of $P<3.88\times10^{-6}$.

Analogous to single-variant association, we passed genes meeting this association

threshold forward for additional consideration with hierarchical FDR correction as

described below.

1036

**Hierarchical FDR correction for testing multiple traits and variants**

In controlling for multiple testing our goal was to make sure that, by looking across 64

traits, we did not increase the proportion of falsely discovered variants. To accomplish

this, we adopted a FDR controlling procedure described in Peterson et al.[91], which uses a

hierarchical strategy to increase power while controlling type I error (**Supplementary**

**Methods**). This procedure has two stages. Stage 1 identifies the set of R variants that are

associated with at least one trait, controlling genome-wide FDR across all variants at

0.05. Stage 2 identifies all the traits that are associated with the discovered variants in a

manner that guarantees an average FDR<0.05.

1046

In Stage 1 we restricted ourselves to the R=531 variants that have an unconditional

association $P<5\times10^{-7}$ with at least one trait. For these, we calculated a p-value for the

hypothesis of no association between the variant and any of the 64 traits using Simes'

rule[92], a combination rule that is robust to dependence between phenotypes. To account

for the fact that we did an initial selection of these R variants from the total number of

variants tested (T), we passed the Simes p-values to a Benjamini-Hochberg (BH)

procedure that controls FDR at target level 0.05×R/T, a modification[93] which guarantees

that the FDR in the set of S variants discovered to be associated with at least one trait is

less than 0.05.

1056

1057     In stage 2, to determine which traits are associated with the set of the S selected variants

1058     we apply the Benjamini and Bogomolov[94] procedure. This procedure applies a

1059     multiplicity correction variant by variant, passing the 64 trait association p-values from

1060     each of the S selected variants and all 64 traits to a BH procedure that controls FDR at

1061     target level $0.05 \times S/T$.

1062

1063     We applied this hierarchical correction to two different sets of results: the set of single-

1064     variant unconditional results and the set of gene-based test results. The gene-based tests

1065     used a threshold of $P<3.88 \times 10^{-6}$ to identify the R nominally significant genes in the first

1066     stage of the hierarchical procedure.

1067

1068     **Genotype validation**

1069     We validated exome sequence-based genotype calls using Sanger sequencing for

1070     METSIM carriers of 13 trait-associated very rare variants with MAF<0.1% in seven

1071     genes. All but one of 108 (99.1%) non-reference genotypes validated were concordant.

1072

**Association replication in additional Finnish cohorts**

We performed replication analysis of significant single-variant associations ($P<5 \times 10^{-7}$)

and follow-up analysis of suggestive single-variant associations ($P<5 \times 10^{-5}$) in up to

24,776 individuals from three GWAS cohort studies: Northern Finland Birth Cohort 1966

(NFBC1966), the Helsinki Birth Cohort Study (HBCS), and FINRISK study participants

not included in the exome sequencing portion of FinMetSeq.

1079

1080    A detailed description of the NFBC1966 study has been published previously and

1081    additional information is available at: http://www.oulu.fi/nfbc/node/18091 [22]. The data

1082    used here, including clinical measurements and blood samples for genetic and NMR

1083    metabolite analyses, were collected at the 31-year follow-up in 1997. NFBC1966 samples

1084    (n=5,139) were genotyped on the Illumina 370k array.

1085

1086    The HBCS includes participants born in Helsinki from 1934-1944 and has been described

1087    elsewhere[23]; a basic description is available at https://thl.fi/fi/web/thlfi-en/research-and-

1088    expertwork/projects-and-programmes/helsinki-birth-cohort-study-hbcs-idefix.       HBCS

1089    samples (n=1,412) were genotyped on the Illumina 610k array.

1090

1091    The FINRISK cohort was described in detail above, and participants (replication

1092    n=18,125) were genotyped in several batches on the Illumina 610k, CoreExome, or

1093    OmniExpress arrays[20,21].

1094

For each replication cohort, prior to phasing we performed genotype quality control

batch-wise using standard quality thresholds for both sample-wise and variant-wise

filtering: call rate>95%, HWE>$10^{-6}$, MAF>5%. We pre-phased array genotypes with

Eagle[95] (v2.3) and imputed genotypes genome-wide with IMPUTE[96] (v2.3.1) using the

SISu v2 reference panel consisting of 2,690 sequenced Finnish genomes and 5,092

sequenced Finnish exomes. Following imputation, we assessed imputation quality by

confirming sex, comparing sample allele frequencies with reference population estimates,

and examining imputation quality (INFO score) distributions. We excluded any variant

with INFO<0.7 within a given batch from all replication/follow-up analyses.

1104

For each of the three cohorts, we matched, harmonized, covariate adjusted, and

transformed available phenotypes as described above for FinMetSeq. We used the same

covariates as for FINRISK. For each cohort, we ran single-variant association using the

EMMAX linear mixed model implemented in EPACTS after generating kinship matrices

from LD-pruned (command: plink --indep-pairwise 50 5 0.2) directly genotyped variants

with MAF>5%.

1111

**Association to disease endpoints in FinnGen**

From a list of >1,100 disease endpoints available for analysis in the FinnGen project, we

selected 22 we considered most likely to be related to the quantitative traits analyzed in

FinMetSeq. As described in detail in Tabassum et al.[45], variant associations with disease

endpoints in FinnGen biobank participants were tested using SPAtest R package and

adjusting for age, sex, and first 10 PCs in up to ~97,000 individuals.

1118

**Association replication in UK Biobank**

For the eight traits analyzed in FinMetSeq that were also available in the current UKBB

release (height, weight, BMI, hip circumference, waist circumference, fat percentage,

systolic blood pressure, and diastolic blood pressure), we extracted trait-variant

association statistics for variants reaching $P<5\times10^{-7}$ in the FinMetSeq combined analysis

from the analysis of unrelated white British individuals generated by the Neale lab

([http://www.nealelab.is/uk-biobank](http://www.nealelab.is/uk-biobank)). Seven of the eight traits had at least one associated

variant and 23 of the total of 31 variants were available in UKBB. A comparison of

association results is in **Supplementary Table 13**.

1128

**Population genetic analyses**

*Identifying unrelated individuals*

To identify a set of nearly independent common autosomal SNVs, we removed SNVs

with MAF<5% and pruned the remaining SNVs in windows of 50 SNVs, in steps of 5

SNVs, such that no pair of SNVs had $r^2$>0.2. We used the resulting 26,036 SNVs to

estimate pairwise relationships among the 19,292 exome-sequenced individuals using

KING[97]. We then removed one individual from each of the 4,418 pairs inferred by KING

to have a relationship of 3rd degree or closer, resulting in a set of 14,874 (nearly)

unrelated individuals for population genetic analyses.

1138

*Identifying sub-population clusters in FinMetSeq*

We first combined exome sequence variants and a genome-wide set of 220,798 SNVs

from GWAS arrays to provide a genome-wide backbone to aid in phasing and computing

haplotype sharing. After removing variants with MAC<3, variants in known regions of

long range LD[98] and variants with HWE<$10^{-4}$, we phased the remaining 764,696 variants

using SHAPEIT[99] (version 2, r837). To assess the substructure in our dataset while

minimizing the effect of mixing due to recent population mobility, we focused on the

2,644 unrelated individuals born by 1955 whose parents were both born in the same

municipality (irrespective of the birth location of the individual).

1148

We identified sub-populations of the 2,644 individuals using ChromoPainter (version 2)

and fineSTRUCTURE[100] (version 2.0.8). We first used ChromoPainter to generate a

pairwise co-ancestry matrix, which represents each individual's DNA as a count of

haplotype blocks copied from every other individual in the dataset. Following previous

practices[101], for computational efficiency, we estimated and fixed the switch and global

emission rates as input for ChromoPainter on a subset of the data; cluster inference is

known to be robust to up to 10-fold deviations of the estimated switch and emission

rates[102]. For further computational speedup, we generated an initial clustering by

applying a normal mixture model clustering[103] (mclust package in R, version 5.1) to the

top ten principal components of the coancestry matrix and used this initial cluster

solution as seed to the fineSTRUCTURE analysis. We conducted 1 million Markov chain

Monte Carlo (MCMC) iterations retaining one sample for every 1,000 iterations after

discarding 3 million iterations as burn-in. After MCMC, we used fineSTRUCTURE to

perform *post-hoc* refinement of cluster membership; we started with the MCMC sample

with the highest posterior probability and reassigned membership, taking into account the

cluster membership at each of the recorded MCMC samples[102].

1165

In total, we ran five MCMC chains using fineSTRUCTURE, retaining the configuration

with highest posterior probability for further analysis. We confirmed convergence of the

fineSTRUCTURE MCMC runs by calculating Geweke's convergence diagnostic using

the coda package (version 0.18) in R to compare the number of inferred clusters in the

first 10% and last 50% of the MCMC chain, and visual inspections of the general

consistency of cluster memberships between independent MCMC chains. In total, we

inferred 245 sub-population clusters among the 2,644 individuals.

1173

We inspected the initial clustering solution from fineSTRUCTURE by examining for

each individual the estimated proportion of their haplotype length derived from each of

the inferred clusters using non-negative least squares[102,104]. This approach showed many

individuals derived a substantial proportion of their haplotype length not from the cluster

initially assigned by fineSTRUCTURE, but instead from a different but related sub-

cluster on the fineSTRUCTURE hierarchical clustering tree, suggesting redundancy in

fineSTRUCTURE-inferred clusters. We therefore combined related clusters by

successively merging pairs of clusters that resulted in the smallest decrease in the

posterior probability of the fineSTRUCTURE hierarchical clustering tree. At each merge,

we reorganized individuals into merged cluster memberships and re-estimated the

haplotype-sharing profile for each individual. We iteratively merged the hierarchical tree

until ≥90% of individuals were assigned to the cluster where they also derive the highest

proportion of haplotype sharing, resulting in 16 clusters for the 2,644 reference

individuals, each named based on the most common parental birthplaces of its members

(**Supplementary Table 15**).

1189

*Enrichment of predicted functionally deleterious alleles in Finland*

We assessed enrichment of predicted functionally deleterious alleles in Finland by

comparing the 14,874 nearly unrelated (pairwise kinship coefficient <0.0448) FinMetSeq

individuals to the 14,944 NFE control exomes in gnomAD, excluding from the NFE

individuals from the neighboring countries of Estonia and Sweden in which substantial

numbers of Finns reside. We analyzed sites with base quality score >10, mapping quality

score >20, and coverage equal to or greater than that found in ≥80% of variable sites

(17.73X in FinMetSeq, 32.27X in gnomAD), resulting in ~38.6 Mbp for comparisons.

We considered only the two most common alleles at each site. We contrasted the

proportional site frequency spectra for FinMetSeq and NFE for five functional variant

categories (PTVs, missense, synonymous, UTR, and intronic variants) after accounting

for sample size differences between datasets by down-sampling both datasets to 18,000

chromosomes.

1203

We also assessed the enrichment of functional alleles within subpopulations of the

FinMetSeq dataset. Of the 16 sub-population clusters identified by fineSTRUCTURE, we

used as the reference population a cluster for which the highest proportion of the parents

of its members were from the southwestern, "early-settlement" part of Finland (NSv3,

**Supplementary Table 15**). Twelve of the remaining 15 clusters also have >100 members

and were used in subsequent analyses (**Supplementary Table 15**). We then compared

the ratio of the site frequency spectra to the reference for PTVs, missense, and

synonymous variants, again down-sampling both datasets to 200 haploid chromosomes to

account for sample size differences. For a given comparison, we computed statistical

evidence for enrichment or depletion at a given allele count bin by exact binomial test

against a null of equal number of variants found in both the test and reference cluster.

1215

*Geographical clustering of predicted functionally deleterious alleles*

We first generated a distance matrix tabulating the pairwise geographical distance in

kilometers between the birthplaces of all available parents of unrelated sequenced

individuals. For each variant of interest, we computed for the minor allele carriers in

FinMetSeq the mean distance among all parent pairs. For example, for a variant with

three carriers with information for five (of the possible six) parents, we computed the

mean for all (5-choose-2 = 10) distances. We evaluated statistical significance of

geographical clustering by comparing the mean distance to the means for up to

10,000,000 sets of randomly drawn non-carrier individuals matched by cohort status and

number of parents with birthplace information available.

1226

To assess whether PTVs or missense variants may be more geographically clustered than

synonymous variants, we first identified a set of near-independent variants ($r^2 > 0.02$) with

MAC$\geq$3 and MAF$\leq$5% among the 14,874 unrelated individuals. This set included 4,312

PTVs, 91,851 missense variants, and 49,842 synonymous variants. For each variant, we

computed the mean pairwise geographical distance in kilometers between the birthplaces

across all pairs of the available parents of carriers of the minor allele and regressed this

mean distance on variant class (PTVs, missense, or synonymous) and MAC, MAC$^2$, and

MAC$^3$ (**Supplementary Table 14**).

1235

We also assessed whether variants showing stronger enrichment (compared to NFE) are

more likely to be geographically clustered. Starting with the three functional classes of

variants identified above, we further restricted analysis to those variants found in

gnomAD so we could calculate the enrichment in frequency over gnomAD NFE. We

included 1,540 PTVs, 46,953 missense, and 28,912 synonymous variants in this analysis

after pruning variants for LD with PLINK. As above, we computed the mean pairwise

distances among parents of carriers of the minor allele and regressed mean distance on

the logarithm of enrichment and MAC, $MAC^2$, and $MAC^3$ (**Supplementary Table 17**). In

both analyses, we first assessed a model with the interaction terms but reported only the

model without interactions if the interactions were not significant.

1246

*Heritability estimates and genetic correlations*

We used genome-wide array genotype data on the 13,326 unrelated individuals for whom

both exome sequence and array data were available to estimate heritability and genetic

correlations for the 64 traits. We constructed a GRM with PLINK[105] (v.1.90b,

https://www.cog-genomics.org/plink2) by applying additional filters for MAF>1% and

genotype missingness rate <2% to the set of previously-used genotyped SNVs, leaving

205,149 SNVs for GRM calculation. We used the exact mixed model approach of

biMM[106] (v.1.0.0, http://www.helsinki.fi/~mjxpirin/download.html) to estimate the

heritability of our 64 traits and the genetic correlation of the 2,016 trait pairs.

**Methods References**
62    Stancáková, A. *et al.* Changes in insulin sensitivity and insulin release in relation
to glycemia and glucose tolerance in 6,414 Finnish men. *Diabetes* **58**, 1212-1221,
doi:10.2337/db08-1607 (2009).
63    Thygesen, K. *et al.* Third universal definition of myocardial infarction. *J. Am.*
*Coll. Cardiol.* **60**, 1581-1598, doi:10.1016/j.jacc.2012.08.001 (2012).
64    European Stroke Initiative Executive, C. *et al.* European Stroke Initiative
Recommendations for Stroke Management-update 2003. *Cerebrovasc. Dis.* **16**,
311-337 (2003).
65    Borodulin, K. *et al.* Cohort Profile: The National FINRISK Study. *Int J*
*Epidemiol*, doi:10.1093/ije/dyx239 (2017).
66    Borodulin, K. *et al.* Forty-year trends in cardiovascular risk factors in Finland.
*Eur. J. Public Health* **25**, 539-546, doi:10.1093/eurpub/cku174 (2015).

67 Soininen, P., Kangas, A. J., Würtz, P., Suna, T. & Ala-Korpela, M. Quantitative
serum nuclear magnetic resonance metabolomics in cardiovascular epidemiology
and genetics. *Circ. Cardiovasc. Genet.* **8**, 192-206,
doi:10.1161/CIRCGENETICS.114.000216 (2015).
68 Wurtz, P. *et al.* Quantitative Serum Nuclear Magnetic Resonance Metabolomics
in Large-Scale Epidemiology: A Primer on -Omic Technologies. *American*
*journal of epidemiology* **186**, 1084-1096, doi:10.1093/aje/kwx016 (2017).
69 Wu, J. *et al.* A summary of the effects of antihypertensive medications on
measured blood pressure. *Am J Hypertens* **18**, 935-942,
doi:10.1016/j.amjhyper.2005.01.011 (2005).
70 Tobin, M. D., Sheehan, N. A., Scurrah, K. J. & Burton, P. R. Adjusting for
treatment effects in studies of quantitative traits: antihypertensive therapy and
systolic blood pressure. *Statistics in medicine* **24**, 2911-2935,
doi:10.1002/sim.2165 (2005).
71 Liu, D. J. *et al.* Exome-wide association study of plasma lipids in >300,000
individuals. *Nature genetics* **49**, 1758-1766, doi:10.1038/ng.3977 (2017).
72 Friedewald, W. T., Levy, R. I. & Fredrickson, D. S. Estimation of the
concentration of low-density lipoprotein cholesterol in plasma, without use of the
preparative ultracentrifuge. *Clin Chem* **18**, 499-502 (1972).
73 DePristo, M. A. *et al.* A framework for variation discovery and genotyping using
next-generation DNA sequencing data. *Nat. Genet.* **43**, 491-498,
doi:10.1038/ng.806 (2011).
74 Jun, G. *et al.* Detecting and estimating contamination of human DNA samples in
sequencing and array-based genotype data. *Am. J. Hum. Genet.* **91**, 839-848,
doi:10.1016/j.ajhg.2012.09.004 (2012).
75 Tan, A., Abecasis, G. R. & Kang, H. M. Unified representation of genetic
variants. *Bioinformatics* **31**, 2202-2204, doi:10.1093/bioinformatics/btv112
(2015).
76 Davis, J. P. *et al.* Common, low-frequency, and rare genetic variants associated
with lipoprotein subclasses and triglyceride measures in Finnish men from the
METSIM study. *PLoS genetics* **13**, e1007079, doi:10.1371/journal.pgen.1007079
(2017).
77 Das, S. *et al.* Next-generation genotype imputation service and methods. *Nature*
*genetics* **48**, 1284-1287, doi:10.1038/ng.3656 (2016).
78 McCarthy, S. *et al.* A reference panel of 64,976 haplotypes for genotype
imputation. *Nature genetics* **48**, 1279-1283, doi:10.1038/ng.3643 (2016).
79 McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122,
doi:10.1186/s13059-016-0974-4 (2016).
80 Kircher, M. *et al.* A general framework for estimating the relative pathogenicity
of human genetic variants. *Nat. Genet.* **46**, 310-315, doi:10.1038/ng.2892 (2014).
81 Adzhubei, I. A. *et al.* A method and server for predicting damaging missense
mutations. *Nature methods* **7**, 248-249, doi:10.1038/nmeth0410-248 (2010).
82 Chun, S. & Fay, J. C. Identification of deleterious mutations within three human
genomes. *Genome research* **19**, 1553-1561, doi:10.1101/gr.092619.109 (2009).

83    Schwarz, J. M., Cooper, D. N., Schuelke, M. & Seelow, D. MutationTaster2:
1314           mutation prediction for the deep-sequencing age. *Nature methods* **11**, 361-362,
1315           doi:10.1038/nmeth.2890 (2014).

1316    84    Kumar, P., Henikoff, S. & Ng, P. C. Predicting the effects of coding non-
1317           synonymous variants on protein function using the SIFT algorithm. *Nature*
1318           *protocols* **4**, 1073-1081, doi:10.1038/nprot.2009.86 (2009).

1319    85    Kang, H. M. *et al.* Variance component model to account for sample structure in
1320           genome-wide association studies. *Nature genetics* **42**, 348-354,
1321           doi:10.1038/ng.548 (2010).

1322    86    Kettunen, J. *et al.* Genome-wide study for circulating metabolites identifies 62
1323           loci and reveals novel systemic effects of LPA. *Nature communications* **7**, 11122,
1324           doi:10.1038/ncomms11122 (2016).

1325    87    Kettunen, J. *et al.* Genome-wide association study identifies multiple loci
1326           influencing human serum metabolite levels. *Nature genetics* **44**, 269-276,
1327           doi:10.1038/ng.1073 (2012).

1328    88    Teslovich, T. M. *et al.* Identification of seven novel loci associated with amino
1329           acid levels using single-variant and gene-based tests in 8545 Finnish men from
1330           the METSIM study. *Hum Mol Genet* **27**, 1664-1674, doi:10.1093/hmg/ddy067
1331           (2018).

1332    89    Inouye, M. *et al.* Novel Loci for metabolic networks and multi-tissue expression
1333           studies reveal genes for atherosclerosis. *PLoS Genet.* **8**, e1002907,
1334           doi:10.1371/journal.pgen.1002907 (2012).

1335    90    Lee, S. *et al.* Optimal unified approach for rare-variant association testing with
1336           application to small-sample case-control whole-exome sequencing studies.
1337           *American journal of human genetics* **91**, 224-237, doi:10.1016/j.ajhg.2012.06.007
1338           (2012).

1339    91    Peterson, C. B., Bogomolov, M., Benjamini, Y. & Sabatti, C. Many Phenotypes
1340           Without Many False Discoveries: Error Controlling Strategies for Multitrait
1341           Association Studies. *Genet. Epidemiol.* **40**, 45-56, doi:10.1002/gepi.21942 (2016).

1342    92    Simes, R. J. An Improved Bonferroni Procedure for Multiple Tests of
1343           Significance. *Biometrika* **73**, 751-754, doi:Doi 10.2307/2336545 (1986).

1344    93    Brzyski, D. *et al.* Controlling the Rate of GWAS False Discoveries. *Genetics* **205**,
1345           61-75, doi:10.1534/genetics.116.193987 (2017).

1346    94    Benjamini, Y. & Bogomolov, M. Selective inference on multiple families of
1347           hypotheses. *J. R. Stat. Soc. Series B Stat. Methodol.* **76**, 297-318,
1348           doi:10.1111/rssb.12028 (2014).

1349    95    Loh, P. R. *et al.* Reference-based phasing using the Haplotype Reference
1350           Consortium panel. *Nature genetics* **48**, 1443-1448, doi:10.1038/ng.3679 (2016).

1351    96    Howie, B. N., Donnelly, P. & Marchini, J. A flexible and accurate genotype
1352           imputation method for the next generation of genome-wide association studies.
1353           *PLoS genetics* **5**, e1000529, doi:10.1371/journal.pgen.1000529 (2009).

1354    97    Manichaikul, A. *et al.* Robust relationship inference in genome-wide association
1355           studies. *Bioinformatics* **26**, 2867-2873, doi:10.1093/bioinformatics/btq559 (2010).

1356    98    Price, A. L. *et al.* Long-range LD can confound genome scans in admixed
1357           populations. *American journal of human genetics* **83**, 132-135; author reply 135-
1358           139, doi:10.1016/j.ajhg.2008.06.005 (2008).

1359    99    Delaneau, O., Zagury, J. F. & Marchini, J. Improved whole-chromosome phasing
1360           for disease and population genetic studies. *Nat Methods* **10**, 5-6,
1361           doi:10.1038/nmeth.2307 (2013).
1362    100   Lawson, D. J., Hellenthal, G., Myers, S. & Falush, D. Inference of population
1363           structure using dense haplotype data. *PLoS genetics* **8**, e1002453,
1364           doi:10.1371/journal.pgen.1002453 (2012).
1365    101   Kerminen, S. *et al.* Fine-Scale Genetic Structure in Finland. *G3* **7**, 3459-3468,
1366           doi:10.1534/g3.117.300217 (2017).
1367    102   Leslie, S. *et al.* The fine-scale genetic structure of the British population. *Nature*
1368           **519**, 309-314, doi:10.1038/nature14230 (2015).
1369    103   Fraley, C. & Raftery, A. E. Model-based clustering, discriminant analysis, and
1370           density estimation. *J Am Stat Assoc* **97**, 611-631, doi:Doi
1371           10.1198/016214502760047131 (2002).
1372    104   Busby, G. B. *et al.* Admixture into and within sub-Saharan Africa. *Elife* **5**,
1373           doi:10.7554/eLife.15266 (2016).
1374    105   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger
1375           and richer datasets. *Gigascience* **4**, 7, doi:10.1186/s13742-015-0047-8 (2015).
1376    106   Pirinen, M. *et al.* biMM: efficient estimation of genetic variances and covariances
1377           for cohorts with high-dimensional phenotype measurements. *Bioinformatics* **33**,
1378           2405-2407, doi:10.1093/bioinformatics/btx166 (2017).
1379

**Figure Legends**

**Figure 1. Characterization of traits by heritability, Pearson correlation, and genetic correlation.** Traits in both figures are in the same order, clockwise in A, and left to right and top to bottom in B, and following the trait group color key.

A) Estimated heritability ($h_x^2$) for each of the 64 traits included in association analysis. Heritability is based on ~205,000 common variants from GWAS arrays available in 13,342 unrelated individuals. Height has the highest heritability estimate at 52.5%. Estimates of trait heritability for metabolic measures are somewhat lower than previous reports (Kettunen, 2012) because estimates are from population-level data as opposed to twin studies and heritability was estimated from covariate adjusted and inverse normal transformed residuals, rather than raw trait values. Trait abbreviations are listed in Supplementary Table 3. All traits are significantly heritable except for 2hr-FFA (Fatty Acid) and His (Amino Acid), see Supplementary Table 5 for estimates, SEs, and P-values.

B) Heatmap of: 1) absolute Pearson correlations of standardized trait values in upper triangle, and 2) absolute values of the genetic correlation, $\rho_G(x,y)$, in lower triangle, where $\rho_G(x,y)$ is the estimated genetic correlation of traits $x$ and $y$. Values below the diagonal in gray had non-estimable genetic correlations.

**Figure 2. Characterization of discovered associations.**

A) Number of genomic loci associated with each trait. Each bar is subdivided into common (MAF>1%, dark blue) and rare (MAF<1%, light blue). Traits are sorted by group as in Figure 1.

B) Relationship between estimated heritability and number of genomic loci detected for each trait. Each trait is colored by trait group following the trait group color key. Vertical bars indicated ±2 standard errors of the heritability estimate. The gray line shows the linear regression fit, shown to indicate the general trend.

C) Heatmap of shared genomic associations by pairs of traits. For traits $x$ and $y$, the color in row $x$ and column $y$ reflects the number of loci associated with both traits divided by the number of loci associated with trait $x$. Traits are presented in the same order as in 2A, and the side and top color bars reflect the trait groups.

D) Relationship between estimated genetic correlation and extent of sharing of genetic associations. For each pair of traits, the extent of locus sharing is defined as the number of loci associated with both traits divided by the total number of loci associated with either trait. The bar within each box is the median, the box represents

the inter-quartile range, whiskers extend up to 1.5x the interquartile range, and outliers are presented as individual points. Analysis using the absolute value of the Pearson correlation of the residual series results in a very similar pattern.

**Figure 3. Allelic enrichment in the Finnish population and its effect on genetic discovery.**
A) Relationship between MAF and estimated effect size for associations discovered in FinMetSeq exomes alone. Each variant reaching significance in FinMetSeq is plotted. Those associations highlighted in Table 2 are represented with a dark blue point (FinMetSeq MAF) and a corresponding brown point reflecting the NFE MAF (gnomAD). The purple lines indicate the 80% power curves for significance at $5 \times 10^{-7}$ for sample sizes of 10,000 and 20,000. The right end of the power curve for N=20,000 terminates at MAF = 0.007. Plots show the dramatic increase in power due to higher relative frequency in Finland.

B) Relationship between MAF and estimated effect size for associations discovered in the combined analysis. Same plot as in A, highlighting the variants in Table 2 only reaching significance in the combined analysis.

**Figure 4. Regional variation in allele frequencies by functional annotation.** Enrichment of functional allelic class in sub-populations (regions) of Northern and Eastern Finland. For each minor allele count bin, we computed the ratio of number of variants found in each subpopulation to an internal reference subpopulation (NSv3), after down-sampling the frequency spectra of all populations to 200 chromosomes. Pink cells represent an enrichment (ratio >1), blue cells represent a depletion (ratio <1). The 12 sub-populations with sample size >100 are shown. The results are consistent with multiple independent bottlenecks followed by subsequent drift in Northern and Eastern Finland, particularly for populations in Lapland and Northern Ostrobothnia. Abbreviations for regions: Kainuu (Kai), Lapland (Lap1, Lap2), Northern Karelia (NKa1, NKa2, NKa3, NKa4), Northern Ostrobothnia (NOs1, NOs2, NOs3, NOs4), Northern Savonia (NSv1, NSv2, NSv3), Southern Ostrobothnia (SOs), and Surrendered Karelia (SuK). For more detailed information on region definitions see Supplementary Table 15. Confidence intervals on the enrichment ratios, and their P-values, are presented in Supplementary Table 16.

**Figure 5. Geographical clustering of associated variants.**
A) Geographical clustering of PTVs as a function of MAC and frequency enrichment over NFE from gnomAD. For each PTV ($r^2 \leq 0.02$, MAC$\geq$3, MAF$\leq$0.05) we computed the mean distance between birth places of available

parents of all carriers of the minor allele. We compared the frequency of the minor allele in FinMetSeq to gnomAD NFE. Blue and pink colors denote the frequency is lower or higher in FinMetSeq than in gnomAD NFE, respectively. The size of the point is proportional to the logarithm of the frequency ratio difference. In general, we observe that variants enriched in FinMetSeq are more geographically clustered.

B) Example of geographical clustering for a trait associated variant. The birth locations of all parents of carriers (orange) and a matching number of parents of non-carriers (blue) of the minor allele for variant chr3:125831672 (rs780671030, p.Arg722X) in *ALDH1L1* are displayed on a map of Finland. This variant is associated with serum glycine levels in FinMetSeq and has a frequency of 0 in NFE samples from gnomAD. The parents of carriers are born on average 135 km apart, the parents of non-carriers on average 250 km apart ($P<10^{-7}$ by permutation).

C) Comparison of geographical clustering between Finnish Disease Heritage (FDH) mutations and trait-associated variants that are >10x more frequent in FinMetSeq than in NFE. The degree of geographical clustering (based on parental birthplace) is comparable between carriers of those variants that showed significant associations in FinMetSeq alone (FMS) and carriers of FDH mutations, and greater than that seen in carriers of variants that showed significant association only in the combined analysis (FMS+Replication). For all variants, carriers of the minor allele displayed greater clustering than non-carriers. The bar within each box is the median, the box represents the inter-quartile range, whiskers extend up to 1.5x the interquartile range, and outliers are presented as individual points.

**Figure Legends (Extended Data Figures)**

**Extended Data Fig. 1. Comparison of allele frequencies of variants in FinMetSeq and NFE from gnomAD.** The comparison of allele frequencies shows the excess of variants at higher frequency in Finland as a result of the multiple bottlenecks experienced in Finnish population history.

**Extended Data Fig. 2. Proportional site frequency spectra between FinMetSeq and gnomAD NFE by variant annotation class**. In general, we find a depletion of the variants in the rarest frequency class, as well as enrichment of variants in the intermediate to common frequency range. The site frequency spectra were down-sampled to 18,000 chromosomes for each dataset.

**Extended Data Fig. 3. Comparison of MAFs for trait-associated variants in FinMetSeq and NFE gnomAD.** Plotted in gray background is a 2-D histogram of variants with non-zero allele frequencies in both gnomAD and FinMetSeq but no trait associations. Variants significantly associated with at least one trait are colored and scaled proportionately to the association p-value, with more significant associations having a larger symbol. Variants >10X enriched in FinMetSeq compared to NFE are pink, those <10X enriched are in blue. The dashed line is the line of equal frequency. Variants unique to Finns and absent in gnomAD are not plotted.

**Extended Data Fig. 4. Gene-based association of extremely rare variants in *APOB* with serum total cholesterol**. The upper panel shows the distribution of the covariate adjusted and inverse-normal transformed phenotype. The lower panel displays the association statistics for each variant included in the gene-based test along with the trait value for minor allele carriers of each variant (orange triangles). SV.P is the P-value from the analysis of each variant in a single-variant analysis.

**Extended Data Fig. 5. Gene-based association of rare variants in *SECTM1* with HDL2 cholesterol**. The upper panel shows the distribution of the covariate adjusted and inverse-normal transformed phenotype. The lower panel displays the association statistics for each variant included in the gene-based test, along with the trait value for minor allele carriers of each variant (orange triangles). SV.P is the P-value from the analysis of each variant in a single-variant analysis.

**Extended Data Fig. 6. Gene-based association of extremely rare variants in *ABCA1* with serum HDL**

**cholesterol**. The upper panel shows the distribution of the covariate adjusted and inverse-normal transformed phenotype. The lower panel displays the association statistics for each variant included in the gene-based test, along with the trait value for minor allele carriers of each variant (orange triangles). SV.P is the P-value from the analysis of each variant in a single-variant analysis.

**Extended Data Fig. 7. Gene-based association of extremely rare variants in *ALDH1L1* with glycine levels**. The upper panel shows the distribution of the covariate adjusted and inverse-normal transformed phenotype. The lower panel displays the association statistics for each variant included in the gene-based test, along with the trait value for minor allele carriers of each variant (orange triangles). SV.P is the P-value from the analysis of each variant in a single-variant analysis.

**Extended Data Fig. 8. Population structure of the FinMetSeq dataset, by region**. Population structure, by region, from principal components analysis of exome sequencing variant data (MAF > 1%), for 14,874 unrelated individuals whose parental birthplaces were known. Color indicates individuals with both parents born in the same region; gray indicates individuals with different parental birth regions, or missing information for one parent. Abbreviations for the regions: Usm, Uusimaa; Swf, Southwest Finland; Stk, Satakunta; Khm, Kanta-Hame; Prk, Pirkanmaa; Phm, Paijat-Hame; Kyl, Kymenlaakso; SKa, Southern Karelia; Nka, Northern Karelia; SSv, Southern Savonia; NSv, Northern Savonia; Ctf, Central Finland; SOs, Southern Ostrobothnia; Osb, Ostrobothnia; COs, Central Ostrobothnia; NOs, Northern Ostrobothnia; Kai, Kainuu; Lap, Lapland; x, split parental birthplaces. Large solid circles represent the center of each region. A map of Finland with regions labeled is supplied for reference.

**Extended Data Fig. 9. Hierarchical clustering tree produced by fineSTRUCTURE**. We identified 16 subpopulations within the FinMetSeq dataset by applying a haplotype-based clustering algorithm, fineSTRUCTURE, on 2,644 unrelated individuals born by 1955 whose parents were both born in the same municipality (Methods). Each subpopulation is named based on the most common parental birth location among its members, with the following abbreviations: NKa, North Karelia; NSv, North Savonia; SOs, South Ostrobothnia; NOs, North Ostrobothnia; Kai, Kainuu; Lap, Lapland; SuK, Surrendered Karelia. A map of Finland with regions labeled is supplied for reference. If multiple subpopulations share the same location label, the subpopulation is further distinguished with a numeral. NSv3 is used as an internal reference in enrichment analysis. See **Supplementary Table 15** for more detailed demographic descriptions of each subpopulation.

**Extended Data Fig. 10. Geographical clustering of missense and synonymous variants as a function of minor allele count and frequency enrichment over gnomAD NFE**. This represents the same analysis as Figure 5A, but for missense and synonymous variants rather than PTVs. Similar to PTVs, missense and synonymous variants that show greater enrichment in FinMetSeq are more likely to be geographically clustered. Blue and pink colors denote the frequency is lower or higher in FinMetSeq than in gnomAD NFE, respectively. The size of the point is proportional to the logarithm of the frequency ratio difference.

Figure 1. Characterization of traits by heritability, Pearson correlation, and genetic correlation. Traits in both figures are in the same order, clockwise in A, and left to right and top to bottom in B, and following the trait group color key.

A) Estimated heritability ($h_x^2$) for each of the 64 traits included in association analysis. Heritability is based on ~205,000 common variants from GWAS arrays available in 13,342 unrelated individuals. Height has the highest heritability estimate at 52.5%. Estimates of trait heritability for metabolic measures are somewhat lower than previous reports (Kettunen, 2012) because estimates are from population-level data as opposed to twin studies and heritability was estimated from covariate adjusted and inverse normal transformed residuals, rather than raw trait values. Trait abbreviations are listed in Supplementary Table 3. All traits are significantly heritable except for 2hr-FFA (Fatty Acid) and His (Amino Acid), see Supplementary Table 5 for estimates, SEs, and P-values.

B) Heatmap of: 1) absolute Pearson correlations of standardized trait values in upper triangle, and 2) absolute values of the genetic correlation, ρg(x,y), in lower triangle, where ρg(x,y) is the estimated genetic correlation of traits x and y. Values below the diagonal in gray had non-estimable genetic correlations.

Figure 2. Characterization of discovered associations.

A) Number of genomic loci associated with each trait. Each bar is subdivided into common (MAF>1%, dark blue) and rare (MAF<1%, light blue). Traits are sorted by group as in Figure 1.

B) Relationship between estimated heritability and number of genomic loci detected for each trait. Each trait is colored by trait group following the trait group color key. Vertical bars indicated ±2 standard errors of the heritability estimate. The gray line shows the linear regression fit, shown to indicate the general trend.

C) Heatmap of shared genomic associations by pairs of traits. For traits x and y, the color in row x and column y reflects the number of loci associated with both traits divided by the number of loci associated with trait x. Traits are presented in the same order as in 2A, and the side and top color bars reflect the trait groups.

D) Relationship between estimated genetic correlation and extent of sharing of genetic associations. For each pair of traits, the extent of locus sharing is defined as the number of loci associated with both traits divided by the total number of loci associated with either trait. The bar within each box is the median, the box represents the inter-quartile range, whiskers extend up to 1.5x the interquartile range, and outliers are presented as individual points. Analysis using the absolute value of the Pearson correlation of the residual series results in a very similar pattern.

Figure 3. Allelic enrichment in the Finnish population and its effect on genetic discovery.

A) Relationship between MAF and estimated effect size for associations discovered in FinMetSeq exomes alone. Each variant reaching significance in FinMetSeq is plotted. Those associations highlighted in Table 2 are represented with a dark blue point (FinMetSeq MAF) and a corresponding brown point reflecting the NFE MAF (gnomAD). The purple lines indicate the 80% power curves for significance at 5x10$^{-7}$ for sample sizes of 10,000 and 20,000. The right end of the power curve for N=20,000 terminates at MAF = 0.007. Plots show the dramatic increase in power due to higher relative frequency in Finland.

B) Relationship between MAF and estimated effect size for associations discovered in the combined analysis. Same plot as in A, highlighting the variants in Table 2 only reaching significance in the combined analysis.

Figure 4. Regional variation in allele frequencies by functional annotation. Enrichment of functional allelic class in sub-populations (regions) of Northern and Eastern Finland. For each minor allele count bin, we computed the ratio of number of variants found in each subpopulation to an internal reference subpopulation (NSv3), after down-sampling the frequency spectra of all populations to 200 chromosomes. Pink cells represent an enrichment (ratio >1), blue cells represent a depletion (ratio <1). The 12 sub-populations with sample size >100 are shown. The results are consistent with multiple independent bottlenecks followed by subsequent drift in Northern and Eastern Finland, particularly for populations in Lapland and Northern Ostrobothnia. Abbreviations for regions: Kainuu (Kai), Lapland (Lap1, Lap2), Northern Karelia (NKa1, NKa2, NKa3, NKa4), Northern Ostrobothnia (NOs1, NOs2, NOs3, NOs4), Northern Savonia (NSv1, NSv2, NSv3), Southern Ostrobothnia (SOs), and Surrendered Karelia (SuK). For more detailed information on region definitions see Supplementary Table 15. Confidence intervals on the enrichment ratios, and their P-values, are presented in Supplementary Table 16.

**Figure 5. Geographical clustering of associated variants.**

A) Geographical clustering of PTVs as a function of MAC and frequency enrichment over NFE from gnomAD. For each PTV (r2≤0.02, MAC≥3, MAF≤0.05) we computed the mean distance between birth places of available parents of all carriers of the minor allele. We compared the frequency of the minor allele in FinMetSeq to gnomAD NFE. Blue and pink colors denote the frequency is lower or higher in FinMetSeq than in gnomAD NFE, respectively. The size of the point is proportional to the logarithm of the frequency ratio difference. In general, we observe that variants enriched in FinMetSeq are more geographically clustered.

B) Example of geographical clustering for a trait associated variant. The birth locations of all parents of carriers (orange) and a matching number of parents of non-carriers (blue) of the minor allele for variant chr3:125831672 (rs780671030, p.Arg722X) in *ALDH1L1* are displayed on a map of Finland. This variant is associated with serum glycine levels in FinMetSeq and has a frequency of 0 in NFE samples from gnomAD. The parents of carriers are born on average 135 km apart, the parents of non-carriers on average 250 km apart (P<10⁻⁷ by permutation).

C) Comparison of geographical clustering between Finnish Disease Heritage (FDH) mutations and trait-associated variants that are >10x more frequent in FinMetSeq than in NFE. The degree of geographical clustering (based on parental birthplace) is comparable between carriers of those variants that showed significant associations in FinMetSeq alone (FMS) and carriers of FDH mutations, and greater than that seen in carriers of variants that showed significant association only in the combined analysis (FMS+Replication). For all variants, carriers of the minor allele displayed greater clustering than non-carriers. The bar within each box is the median, the box represents the inter-quartile range, whiskers extend up to 1.5x the interquartile range, and outliers are presented as individual points.