Mechanism and resistance for antimycobacterial activity of a fluoroquinophenoxazine

compound

Pamela K. Garcia[1,2], Thirunavukkarasu Annamalai[1,2], Wenjie Wang[1,2], Raven Bell[1], Duc Le[1],

Paula Martin Pancorbo[1], Sabah Sikandar[1], Ahmed Seddek[1,2], Xufen Yu[3], Dianqing Sun[3], Anne-

Catrin Uhlemann[4], Purushottam B. Tiwari[5], Fenfei Leng[1,2], Yuk-Ching Tse-Dinh[1,2*]

[1] Department of Chemistry & Biochemistry, Florida International University, Miami, Florida,

United States of America

[2] Biomolecular Sciences Institute, Florida International University, Miami, Florida, United States

of America

[3] Department of Pharmaceutical Sciences, The Daniel K. Inouye College of Pharmacy,

University of Hawaii at Hilo, 34 Rainbow Drive, Hilo, Hawaii, United States of America

[4] Division of Infectious Diseases, Columbia University Medical Center, New York, New York,

United States of America

[5] Department of Oncology, School of Medicine, Lombardi Comprehensive Cancer Center

Georgetown University Medical Center, Washington DC, United States of America

* Corresponding author, Email: ytsedinh@fiu.edu (YT)

Abstract

We have previously reported the inhibition of bacterial topoisomerase I activity by a fluoroquinophenoxazine compound (FP-11g) with a 6-bipiperidinyl lipophilic side chain that exhibited promising antituberculosis activity (MIC = 2.5 µM against *Mycobacterium tuberculosis*, SI = 9.8). Here, we found that the compound is bactericidal towards *Mycobacterium smegmatis*, resulting in greater than 5 Log_{10} reduction in colony-forming units [cfu]/mL following a 10 h incubation at 1.25 µM (4X MIC) concentration. Growth inhibition (MIC = 50 µM) and reduction in cfu could also be observed against a clinical isolate of *Mycobacterium abscessus*. Stepwise isolation of resistant mutants of *M. smegmatis* was conducted to explore the mechanism of resistance. Mutations in the resistant isolates were identified by direct comparison of whole-genome sequencing data from mutant and wild-type isolates. These include mutations in genes likely to affect the entry and retention of the compound. FP-11g inhibits *Mtb* topoisomerase I and *Mtb* gyrase with IC_{50} of 0.24 and 31.5 µM, respectively. Biophysical analysis showed that FP-11g binds DNA as an intercalator but the IC_{50} for inhibition of *Mtb* topoisomerase I activity is >10 fold lower than the compound concentrations required for producing negatively supercoiled DNA during ligation of nicked circular DNA. Thus, the DNA-binding property of FP-11g may contribute to its antimycobacterial mechanism, but that alone cannot account for the observed inhibition of M*tb* topoisomerase I.

Introduction

Tuberculosis (TB) is a devastating disease caused by *Mycobacterium tuberculosis*

infection. The World Health Organization (WHO) reported that TB is the leading cause of death

worldwide from a single infectious agent. The 2018 Global Tuberculosis Report estimated that

TB caused ~1.6 million deaths in 2017, with up to 10 million newly diagnosed TB cases

annually [1]. Significantly, drug-resistant TB poses a great threat to public health. In 2017, there

were 558,000 new cases of drug resistant TB, including multidrug resistant TB (MDR-TB).

resistant to rifampicin and isoniazid (two first-line anti-TB drugs) and rifampicin-resistant TB

(RR-TB) [1]. These TB patients are more likely to have poor treatment outcomes. Therefore,

there is an urgent need for potent new TB drugs with novel modes of action [2]. Resistance to

current antibiotics also makes it difficult to treat infections caused by nontuberculous

mycobacteria (NTM). It is often difficult to distinguish between *M. tuberculosis* and NTM as

the cause of lung infections, so there is a medical need for new antibiotics that have broad

antimycobacterial activity.

DNA topoisomerases are essential enzymes for maintaining optimal DNA topology to

facilitate vital functions including replication, transcription, recombination and DNA repair [3].

Topoisomerases belong to type I or type II families corresponding to cleavage of a single or

double strand of DNA being utilized for catalysis [4] with subfamilies based on sequence

homology and mechanism [5]. Every bacterium has at least one type IA topoisomerase for

overcoming topological barriers that require cutting and rejoining of a single strand of DNA [3,

6]. Topoisomerase I catalytic activity is essential for the viability of *Mycobacterium smegmatis*

(*Msm*) and *M. tuberculosis* (*Mtb*) [7, 8], because these organisms only have one type IA

topoisomerase. *Mtb* topoisomerase I (MtbTopI) is a validated anti-TB target; [7, 9, 10] for

identification of inhibitors that can potentially be developed into leads for new TB therapy.

We have previously shown [11] that a fluoroquinophenoxazine compound (FP-11g,

shown in Scheme 1) with a 6-bipiperidinyl lipophilic side chain inhibited the catalytic activity of

Escherichia coli topoisomerase I (IC_{50} = 0.48 µM), and showed promising antituberculosis

activity (MIC = 2.5 µM against *Mtb* H37Rv, SI = 9.8 for Vero cell cytotoxicity). In this study,

we followed up on the antimycobacterial activity of this compound to further explore its

therapeutic potential. The interactions of this compound with MtbTopI and DNA were

characterized in biophysical assays. Mechanism of resistance to its antimycobacterial activity

was studied with *M. smegmatis*. This commonly used model mycobacterial organism is non-

pathogenic and fast growing, but still shares many features with pathogenic mycobacteria.

Materials and Methods

Synthesis of FP-11g

FP-11g (Scheme 1) was synthesized using our previously reported procedure [11]. The structure

was confirmed by ^{1}H NMR, and high-resolution mass spectrometry (HRMS). Purity was

determined to be >99% by reverse phase C18-HPLC.

MIC (minimum inhibitory concentration) against *M. smegmatis* and *M. abscessus*

M. smegmatis mc2 155 (WT from ATCC) and FP-11g resistant mutants derived from WT were

cultured in 5 ml Middlebrook 7H9 broth containing 0.2% glycerol, 0.05% Tween 80 with or

without a supplement of 10% albumin, dextrose, sodium chloride (ADN) at 37°C with shaking.

Stationary phase bacteria cultures were adjusted to an optical density (OD_{600}) of 0.1 and

subsequently diluted 1:10 using growth media. Fifty microliters (~10^5 cfu) of the diluted culture

were transferred to the individual wells of a clear round-bottom 96-well plate containing 50 µl of

serially diluted compounds. The 96-well plate was then incubated at 37°C with shaking. After 48

hours of incubation, resazurin (final concentration 0.002%) was added to the individual wells

and the fluorescence reading at 560/590 nm was taken with a BioTek Synergy plate reader after

approximately 5h of incubation at 37°C.

A clinical isolate of *M. abscessus* bacterium (isolated at the Columbia University Medical

Center) was cultured in Middlebrook 7H9 ADN broth till it reached an optical density (OD_{600}) of

1.0. The culture was stored at -80°C as 1 ml aliquots containing 15% glycerol. These frozen

aliquots were then used as the inoculum for *M. abscessus* MIC assays. Prior to conducting each

MIC assay, an aliquot of frozen *M. abscessus* was thawed and diluted 1:100 in Middlebrook 7H9

broth. After dilution, the bacterial cells (~10^5 cfu) were added to the wells of a 96-well plate

containing the serially diluted compounds as described for *M. smegmatis* cells and incubated at

37°C with no shaking for 48 h. Subsequently, resazurin (final concentration 0.005%) was added

to each well and fluorescence reading at 560/590 nm (excitation/emission) was taken after 24 h

of incubation at 37°C. MIC was determined as compound concentration that showed at least 90%

reduction in fluorescence when compared to the control (no inhibitor) wells. Ciprofloxacin and

Clarithromycin were included in the assay as positive controls. On the day of each MIC assay,

thawed inoculum of *M. abscessus* was also serially diluted and spread plated on LB agar plates

to confirm for the inoculum load of ~10^5 cfu/well. MIC determination for each bacteria was

repeated at least three times.

Survival assays

The bactericidal effect of FP-11g compound was evaluated in 96-well plates using a protocol

similar to the MIC assay. In brief, *M. smegmatis* was incubated at 37°C with shaking in the

presence of 1X, 2X, 4X and 8X MIC of FP-11g for 6, 10, 24 and 48 hours; *M. abscessus* was

incubated at 37°C without shaking in the presence of 1X and 2X MIC of FP-11g for 24, 48 and

72 hours. At each time point 20 µl from the treatment wells were serially diluted (10 fold),

spread on LB agar plates and incubated at 37°C for 4 or 7 days for counting the viable colonies

of *M. smegmatis* and *M. abscessus* respectively. Ten microliters from the treatment wells were

also enriched in 5-ml of Middlebrook 7H9 broth if there are no viable colonies from a particular

treatment-time combination on the LB agar plates. The survival percentage was calculated by

dividing the number of viable colonies at each time point by the initial viable count prior to the

treatment (time 0). Survival assays were repeated three times.

Isolation of resistant mutants

M. smegmatis mc2 155 was exposed stepwise to increasing concentrations of FP-11g to isolate

mutant strains with different levels of resistance for FP-11g by a slightly modified protocol as

described by Fujimoto-Nakamura et al [12]. Overnight culture of *M. smegmatis* was adjusted

to an optical density (OD_{600}) of 0.1 and a volume of 100 µl ($2x10^6$ cfu) was spread on 7H9 ADN

agar containing 2.5 µM (8X MIC) FP-11g for isolation of resistant mutants. Colonies appearing

on these plates were scored as resistant mutants only if they grew again on 7H9 ADN broth and

agar plates containing 8X MIC of FP-11g. Mutation frequency was calculated based on these

resistant colonies. Only one resistant mutant was isolated at 8X MIC of FP-11g. This resistant

mutant was next exposed to 16X MIC of FP-11g in 7H9 ADN broth to further isolate resistant

colonies PGM1, PGM2 shown in Table 1. As our initial resistant mutation isolation at 8X MIC

of FP-11g resulted in only one mutant, a second round of stepwise mutation was initiated with

4X MIC to higher concentrations of FP-11g (PGM3 – PGM6 shown in Table 1).

Whole genome sequencing

M. smegmatis WT and mutant strains were cultured in 5 ml of Middlebrook 7H9 ADN broth and

2 ml was spun down for genomic DNA extraction. DNA extraction was performed using the

BACTOZOL™ Bacterial DNA isolation kit (Molecular Research Center) according to

manufacturer's instructions**.** DNA quality was evaluated through the UV absorbance 260/280

ratio (>1.8) and DNA quantity was measured using a fluorescence-based Qubit® dsDNA BR

(Broad-Range) Assay Kit (Thermo Fisher). DNA concentration was adjusted to 0.2 ng/μl with

10 mM Tris-HCl pH 8.0. After concentration adjustment, all the genomic DNA samples were

quantified once again to confirm the concentration. Library was prepared with the Nextera XT

DNA Library Prep Kit, 96 Indexes (FC-131-1096). After clean-up of the libraries, random

samples were selected for the Bio-analyzer analysis to verify the average size of the fragments.

Cleaned up libraries were normalized manually and pooled together. The pooled libraries were

then denatured with NaOH (final concentration 0.1 N) and the PhiX Control v3 library added.

This mix was loaded in the pre-thawed reagent cartridge associated with the MiSeq Reagent kit

v3 and sequenced using Illumina MiSeq® Next Generation Sequencer. FASTQ files generated

from the sequencing run was further analyzed in CLC genomics workbench 10 software

(QIAGEN). Sequence of *M. smegmatis* mc2 155 WT strain from our laboratory was used as

reference to compare all the mutant sequences and detect variations (Single Nucleotide

Variations, Deletions and Insertions).

Enzyme assays

MtbTopI relaxation inhibition assay - MtbTopI was expressed in *E. coli* T7 Express crystal strain

(New England Biolabs) and purified as previously described [13,14]. The relaxation activity of

MtbTopI was assayed in a buffer containing 10 mM Tris-HCl, pH 8.0, 50 mM NaCl, 0.1 mg/ml

gelatin, and 0.5 mM $MgCl_2$. Serial dilutions of FP-11g dissolved in DMSO was mixed with 10 µl

of the reaction buffer containing 12.5 ng of enzyme before the addition of 9 µl of reaction buffer

containing 160 ng of supercoiled pBAD/Thio plasmid DNA purified by cesium chloride gradient

as the substrate. The reactions were terminated following an incubation of the mixtures at 37°C

for 30 min by the addition of 4 µl of stop solution (50% glycerol, 50 mM EDTA, and 0.5%

(vol/vol) bromophenol blue) and the mixtures were analyzed by agarose gel electrophoresis with

TAE buffer (40 mM Tris acetate and 1 mM EDTA, pH 8.2). The gels were stained in ethidium

bromide and photographed under UV light.

DNA gyrase supercoiling inhibition assay – *Mtb* DNA gyrase was purchased from Inspiralis. The

supercoiling assays were carried out by mixing serial dilutions of FP-11g in 0.5 µl DMSO with 1

U of the enzyme in reaction buffer of 50 mM HEPES pH 7.9, 100 mM potassium glutamate, 6

mM magnesium acetate, 4 mM DTT, 1 mM ATP, 2 mM spermidine, 0.05 mg/ml BSA, followed

by the addition of 300 ng of relaxed covalently closed circular DNA, for a final reaction mixture

volume of 20 µl. The samples were incubated at 37°C for 30 min before the reactions were

terminated and analyzed by agarose gel electrophoresis.

Surface Plasmon Resonance

Biacore T200 SPR instrument was used to conduct all experiments. MtbTopI was used as ligand

to immobilize on a CM5 sensor surface in 10 mM sodium acetate buffer at pH 5.5, using

standard amine coupling chemistry, to a level of ~13400 RU. PBS-P (20 mM phosphate buffer,

pH 7.4, 2.7 mM KCl, 137 mM NaCl, 0.05% v/v surfactant P20) was used as the immobilization

running buffer. FP-11g was used as an analyte to inject over the ligand immobilized surface in

various concentrations in the presence of PBS-P buffer supplemented with 10% DMSO. The

flow rate of all analyte solutions was maintained at 50 μl/min. One 20 s pulse of 1M NaCl

solution was injected for surface regeneration. The contact and dissociation times used were 60s

and 300s, respectively. SPR sensorgrams were both reference and bulk subtracted. All

experiments were conducted at 25°C.

Measurement of visible absorbance and fluorescence emission spectra

Visible absorbance spectra of free and bound FP-11g were recorded in a Cary Bio50 UV-VIS

spectrophotometer. Fluorescence emission spectra were measured by using a Cary fluorescence

spectrophotometer.

DNA UV Melting studies

DNA UV melting curves were determined using a Cary 100 UV-Vis spectrophotometer

equipped with a thermoelectric temperature-controller. The salmon testes (ST) DNA in the

presence of different concentrations of FP-11g in 1×BPE buffer (6 mM Na_2HPO_4, 2 mM

NaH_2PO_4, and 1 mM EDTA, pH 7) was used for UV melting studies. Samples were heated at a

rate of 1°C min^{-1}, while continuously monitoring the absorbance at 260 nm. Primary data were

transferred to the graphic program Origin (MicroCal, Inc., Northampton, MA) for plotting and

analysis.

Dialysis experiments

Dialysis assays were carried out according to the previously published procedure [15]. Briefly, a

volume of 0.3 ml of 75 µM (bp) of ST DNA was pipetted into a 0.3 ml disposable dialyzer. The

dialysis units were then placed into a beaker with 100 ml of 1×BPE buffer (6 mM Na_2HPO_4, 2

mM NaH_2PO_4, and 1 mM EDTA, pH 7) containing 1 µM of FP-11g. The dialysis was allowed to

equilibrate with continuous stirring for 72 hours at room temperature (24°C). After the dialysis,

the free, bound, and total concentrations of FP-11g were determined spectrophotometrically.

These values were used to determine the DNA binding constant of FP-11g.

DNA ligation assays

DNA ligation assays were carried out in 1×DNA T4 DNA ligase buffer (50 mM Tris-HCl (pH

7.5), 10 mM $MgCl_2$, 1 mM ATP, and 10 mM DTT) using 1,200 units of T4 DNA ligase (New

England Biolabs) in 100 µl of solution containing the nicked plasmid pAB1 [16] in the presence

of different concentrations of FP-11g. After incubation at 37°C for 30 min, the ligation reactions

were stopped by extraction with 100 µl of phenol. The DNA samples were analyzed by

electrophoresis in a 1% agarose gel in TAE. After electrophoresis, the agarose gel was stained

with ethidium bromide and photographed using a Kodak imaging system.

Results

Growth inhibition of *M. smegmatis* and *M. abscessus*

The MIC of FP-11g for *M. smegmatis* mc2 155 was found to be 0.31 µM in multiple

measurements. The MIC value is the same in 7H9 media with or without ADN supplement.

There is also no effect on the MIC from shaking during the incubation. For the clinical *M.*

abscessus isolate, the MIC was 50 μM, compared to Clarithromycin MIC of 0.7-1.56 μg/ml (1-2

μM) and Ciprofloxacin MIC of 12.5 μg /ml (38 μM), IC_{50} for 50% growth inhibition of the *M.*

abscessus strain was estimated to be 3-6 μM.

Bactericidal activity of FP-11g

FP-11g showed strong bactericidal activity against *M. smegmatis*. When *M. smegmatis* cells

(~1x10^5 cells in each well in 96-well plate) were exposed to 2X MIC of FP-11g, the cell viability

was diminished by ~ 4.5 Log after 48 hours of incubation (Fig 1A). For concentrations of 4X

MIC and 8X MIC, 24 hours were sufficient to diminish cell viability by greater than 5 Log, with

no viable cells remaining at 48 hours.

Treatment of *M. abscessus* with 2X MIC of FP-11g resulted in >3 Log loss of cell

viability (Fig 1B). The *M. abscessus* strain grows slower than *M. smegmatis*, so treatment was

extended to 72 hours. Because of the higher MIC value for *M. abscessus*, FP-11g concentrations

could only be tested at up to 4X MIC in the survival assay.

Fig 1. Bactericidal effect of FP-11g on *M. smegmatis* mc2 155 and *M. abscessus*

(A) The bactericidal effect of FP-11g on *M. smegmatis* were determined using 1X (0.31 μM), 2X

(0.62 μM), 4X (1.25 μM) and 8X (2.5 μM) MIC at different time points. The downward arrow

indicates that no viable colonies were detected following treatment with 4X and 8X MIC at time

points beyond 10 hours. **(B)** Survival of clinical isolate of *M. abscessus* following treatment

with FP-11g at 50 and 100 μM. Percent survival is calculated by the ratio of cfu from treated

culture versus cfu from culture prior to addition of FP-11g. The error bars represent the standard

deviation of results of experiments repeated three times.

Isolation and verification of resistant mutants

253   The mutation frequency for resistance to FP-11g at 8X (MIC) was estimated to be 5×10^{-7}. Six

254   *M. smegmatis* strains (PGM1 – PGM6, Table 1) isolated from stepwise increase of FP-11g

255   concentrations were verified to be resistant to the compound by determination of MIC. An

256   increase in the resistance to Moxifloxacin was also observed but the fold-increase in

257   Moxifloxacin was significantly lower than the fold-increase for FP-11g MIC.

Mutations identified in WGS

The mutations found in each of the resistant strains are listed in Table 2. Mutations in ten

different genes associated with deletions, insertions or SNV (Single Nucleotide Variations) were

detected in these mutants (Table S1). All of the resistant strains have SNV on *MSMEG_0965*

gene (*mspA*), which codes for the major porin in *M. smegmatis.* This porin is important for the

permeation of nutrients and antibiotics inside the cell. Previous studies showed that the deletion

of this gene in *M. smegmatis* resulted in a reduced permeability to drugs such as β-lactams,

fluoroquinolones and chloramphenicol [17-20]. The *mspA* gene is present in fast-growing

mycobacteria only, and is not found in *M. tuberculosis* [21, 22].

The second most frequent mutation (in 3 out of 6 mutants) detected corresponds to

MSMEG_5623 and *MSMEG_6430. MSMEG_5623* gene codes for an L-carnitine dehydratase

homolog of unknown function in *M. smegmatis,* with no homolog in *M. tuberculosis* according

to Tuberculist (http://svitsrv8.epfl.ch/tuberculist/)*.* However, using the blast tool it was

determined that this L-carnitine dehydratase homolog from *M. smegmatis* has 32% identity when

aligned with the L-carnitine dehydratase from *M. tuberculosis* (*Rv3272),* including Asp33

(conserved amino acid) that corresponds to the amino acid mutated in FP-11g resistant strains.

MSMEG_6430, with a Thr to Lys substitution mutation detected in two mutants, is classified as a

membrane protein that may have diverse functions in the cells, such as transportation of

molecules through the membrane or serving as receptors for chemical signals [23, 24]. An Ile to

Ser substitution mutation was also detected in *MSMEG_2820* which encodes for an unknown

integral membrane protein. The mutations in diverse membrane proteins detected here may

affect the transport of FP-11g across the membrane.

The remainder of detected mutations occurred at lower frequencies. *MSMEG_1513* (Ser

to Cys substitution) codes for a hypothetical protein with no homolog in *M. tuberculosis* and has

been classified as an oxidoreductase. *MSMEG_0241* (homolog of *Rv0202c*) with frame shift

mutation detected encodes for MmpL11 (mycobacteria membrane protein large), a protein that

belongs to a family of transporters and contribute to the cell wall biosynthesis in mycobacteria

[25]. MmpL proteins family has been associated with drug resistance in *M. abscessus* and *M.*

tuberculosis [26]. Frame shift mutation was also detected in *MSMEG_0240* (homolog of

Rv0201c), which does not have a known function.

Comparison of inhibition of MtbTopI versus DNA gyrase

The IC_{50} of FP-11g for inhibition of *E. coli* topoisomerase I relaxation activity has been reported

as 0.48 µM [11]. In comparison, the IC_{50}s for inhibition of human topoisomerase I relaxation

activity and topoisomerase IIα decatenation activity were found to be both around 3.9 µM in the

previous report [11]. Here we determined the IC_{50} for inhibition of MtbTopI and *Mtb* DNA

gyrase by FP-11g (Fig 2). The IC_{50} for MtbTopI (0.24 µM) is ~100-fold lower than the IC_{50} for

inhibition of *Mtb* DNA gyrase (31 µM).

Fig 2. Comparison of inhibition of MtbTopI and DNA gyrase activities by FP-11g

(**A**) Inhibition of MtbTopI relaxation of negatively supercoiled DNA by increasing concentrations

of FP-11g. (**B**) Inhibition of *Mtb* DNA gyrase supercoiling of relaxed DNA requires higher

concentrations of FP-11g.

FP-11g can interact directly with MtbTopI

To detect direct enzyme-inhibitor interaction, MtbTopI was immobilized on the CM5 chip

surface and allowed to interact with FP-11g in the analyte solution. The SPR sensorgram (Fig 3)

qualitatively showed that MtbTopI interacts directly with FP-11g.

Fig 3. Qualitative evaluation of the binding of FP-11g with MtbTopI using SPR

SPR sensorgrams showing the direct binding of FP-11g with the immobilized MtbTopI. FP-11g

was injected at 2.5 µM (green), 5 µM (blue), and 10 µM (red) in duplicate.

FP-11g binds to DNA with intercalation

Since FP-11g has a planar tetracyclic ring scaffold (Scheme 1), it may bind to DNA by

intercalation. Fig 4 shows the visible absorption spectra (Fig 4A) and fluorescence emission (Fig

4B) spectra of free and DNA-bound FP-11g. Binding of the compound to DNA results in a red

shift of the maximum absorbance from 386 nm to 406 nm. Interestingly, upon binding to DNA,

FP-11g has a pronounced induced visible band around 330 nm with a maximum absorbance at

334 nm. Binding of FP-11g to DNA also causes a red shift to the fluorescence emission spectrum

of FP-11g (Fig 4B). The fluorescence intensity is slightly enhanced upon binding to DNA. Table

3 summarizes the optical properties of free and DNA-bound FP-11g.

Fig 4C shows the DNA UV melting profiles in which different concentrations of FP-11g

were added to the same concentration of ST DNA. In the absence of FP-11g, the UV melting

temperature of ST DNA (Tm) was determined to be 70.8°C. In the presence of saturated FP-11g

(40 µM), the DNA UV melting temperature was increased to 81.9°C, indicating that FP-11g

tightly binds to ST DNA. In this study, we also carried out a dialysis binding assay in which 75

µM (bp) ST DNA was extensively dialyzed against a large excess of 1 µM of FP-11g. Fig 4D

shows our results. After 72 hours of dialysis, the concentration of FP-11g outside the dialysis bag

was not significantly changed. In contrast, the concentration of FP-11g inside the dialysis bag

where the ST DNA was located was increased to 25 µM. The DNA binding constant of FP-11g

was estimated to be 1×10^6 M^{-1} in 1XBPE.

Fig 4. Binding of FP-11g to salmon testes DNA in 1×BPE buffer (6 mM Na$_2$HPO$_4$, 2 mM

NaH$_2$PO$_4$, and 1 mM EDTA, pH 7). (A) Visible absorption spectra of free (black line) and DNA

bound (red line). **(B)** Fluorescence emission spectra of free (black line) and DNA bound (red line).

The fluorescence emission spectra were recorded with λ_{em} = nm. **(C)** DNA UV melting of salmon

testes DNA in the presence of various concentrations of FP-11g. The FP-11g concentrations are 0,

2, 5, 7.5, 10, 15, 20, and 40 µM from left to right. **(D)** The DNA dialysis assay was performed as

described in the Methods. Visible absorption spectra of FP-11g inside the dialysis bag (red line)

and outside dialysis bag (black line).

341  Fig 5 shows the results of ligation of nicked circular DNA in the presence of different

342 concentrations of FP-11g by T4 DNA ligase. Change in DNA linking number was observed at 2

343 μM. Concentrations of FP-11g at 5 and 10 μM drove the plasmid DNA template into (-)

344 supercoiled DNA. This result suggests that FP-11g can act as a DNA intercalator at these higher

345 concentrations. The DNA intercalation of FP-11g at the observed IC_{50}s could be the basis for the

346 inhibition of DNA gyrase and human topoisomerases [11], but could not be the sole basis of

347 inhibition of bacterial topoisomerase I observed at 10-fold lower IC_{50}s.

349 **Fig 5. Products of DNA ligation in the presence of FP-11g.** DNA ligation of nicked pAB1

350 plasmid DNA by T4 DNA ligase in the presence of indicated concentrations of FP-11g were

351 performed as described under Methods.

Discussion

354 In previous studies, the FP-11g compound has been proposed as an antimycobacterial agent

355 active against *M. tuberculosis* (MIC = 2.5 μM) [11]. Here we demonstrate its activity against the

356 pathogenic NTM *M. abscessus*. Compounds active against both species would be useful in

357 clinical settings since in some cases the symptoms and clinical manifestations of infections

358 caused by these two organisms are difficult to differentiate, consequently empiric treatment

359 effective for both organisms is required. Additionally, patients suffering from cystic fibrosis or

360 other immunosuppressive condition can be infected by different mycobacteria at the same time.

361 In fact, co-infections with *M. abscessus* and *M. tuberculosis* have been reported [27, 28]. The

362 strong bactericidal effect of FP-11g against *M. smegmatis* supports investigation of this

363 compound and its derivatives for its potential use as a broad antimycobacterial agent. The

364   finding of the growth inhibitory effect of FP-11g on *M. abscessus* is encouraging because of the

365   lack of response of NTM clinical strains and subspecies of *M. abscessus* to nearly all of the

366   current antibiotics.

367   No specific mutations found in FP-11g *M. smegmatis* resistant mutants could be

definitively associated with the proposed mechanism of action of this compound, but all of them

may play a role in drug resistance. Expression of *M. smegmatis* MspA porin in *M. tuberculosis*

promotes not only cell growth but also antibiotic susceptibility [29]. To date, porin genes have

not yet been identified in *M. tuberculosis*, even though some studies suggest the presence of

these structures in these organisms [22]. The mutation in MspA porin is likely to be the first

mechanism of resistance developed by *M. smegmatis*, and hence the common mutation was

found. Changes in the porin may interrupt the drug transportation into the cell and support cell

survival. Since this porin is present in fast growing mycobacteria, it would be of significance in

the treatment of atypical pathogenic mycobacteria including *M. abscessus* studied here [30].

377   Characterization of integral membrane proteins has been always a challenge due to the

difficulty for the expression, solubility and crystallization of these proteins. Most of the

information has been obtained through computational approaches, which predict the structural

conformation of a protein based on the primary sequence and classify the different types of

transmembrane proteins [31, 32]. Transmembrane proteins could be associated with

mechanisms of resistance to drugs that are not targeting them directly. Integral component of

membrane are channels through the cellular membrane that transport amino acids, lipids,

coenzymes, carbohydrates, nucleotides and other metabolites [33]. Mutations in integral

membrane proteins detected here may affect drug transportation through the membrane.

Variation in channels that transport molecules through the cell wall will affect the intake of

molecules including drugs and metabolites. These integral components of membrane could also

be non-characterized efflux pumps that may be extruding the drug from the organism and

support the drug resistance.

In this study we detected the MmpL11 mutation in two resistant mutants. Previous

studies with *M. tuberculosis* strains harboring mutations in MmpL proteins showed that all of the

mutant strains retained general drug susceptibility to diverse antibacterial agents. This suggests

that MmpL proteins do not play a direct role in drug resistance [34]. Nonetheless, recent studies

have shown that the extended RND permease superfamily in *M. tuberculosis* include both RND

multidrug efflux transporters and members of the MmpL family [35]. In *M. abscessus*, MmpL

has been associated with drug resistance through efflux pumps [36]. Additionally, mutations on

MmpL11 proteins has been associated with impairment of biofilm formation in *M. smegmatis*.

The absence of this gene has generated a reduced permeability to antimicrobial agents [37, 38],

which is evidence that this mutation may play an important role in FP-11g resistance.

Remarkably, we did not detect mutations in the *topA* gene in the resistant mutant isolates.

This might be due to FP-11g acting as an unconventional catalytic inhibitor. These types of

inhibitors are described as compounds that bind to the DNA as well as to the topoisomerase

enzyme to inhibit the topoisomerase activity [39]. Mutations that affect the essential activity of

topoisomerase I could compromise cell growth to a significant extent and would less likely be

selected for resistance than the mutations detected here that can limit compound transport into

the cell. It is also possible that due to its interaction with DNA, FP-11g inhibits multiple DNA

binding proteins in its mode of action, including DNA gyrase as observed here. Further studies

are required to identify other analogs of FP-11g that can be more selective in targeting

topoisomerase I activity in its antimycobacterial mechanism of action. This would require

enhancing the interaction of the inhibitor with the MtbTopI enzyme, and reducing the

dependence of inhibition on DNA binding.

Acknowledgments

414 We thank Dr. DeEtta Mills and Christina Burns from the International Forensic Research

415 Institute at FIU for support in WGS.

Funding

This research was supported by NIH grant R01 AI069313 (to YT). Experimental SPR

sensorgrams were measured using Biacore T200 SPR instrument available in Biacore Molecular

Interaction Shared Resource (BMISR) facility at Georgetown University. The BMISR is

supported by NIH grant P30CA51008. REU participant RB was supported by the NSF-REU

Site Grant CHE1560375 to FIU. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

References

1. World Health Organization, W.H.O., *Global tuberculosis report.* 2018.

2. Marais, B.J., *The global tuberculosis situation and the inexorable rise of drug-resistant disease.* Adv Drug Deliv Rev, 2016. **102**: p. 3-9.

3. Wang, J.C., *Cellular roles of DNA topoisomerases: a molecular perspective.* Nat Rev Mol Cell Biol, 2002. **3**(6): p. 430-40.

4. Corbett, K.D. and J.M. Berger, *Structure, molecular mechanisms, and evolutionary relationships in DNA topoisomerases.* Annu Rev Biophys Biomol Struct, 2004. **33**: p. 95-118.

5. Schoeffler, A.J. and J.M. Berger, *DNA topoisomerases: harnessing and constraining energy to govern chromosome topology.* Q Rev Biophys, 2008. **41**(1): p. 41-101.

6. Forterre, P. and D. Gadelle, *Phylogenomics of DNA topoisomerases: their origin and putative roles in the emergence of modern organisms.* Nucleic Acids Res, 2009. **37**(3): p. 679-92.

7. Ahmed, W., et al., *Conditional silencing of topoisomerase I gene of Mycobacterium tuberculosis validates its essentiality for cell survival.* FEMS microbiology letters, 2014. **353**(2): p. 116-123.

8. Ahmed, W., et al., *Reduction in DNA topoisomerase I level affects growth, phenotype and nucleoid architecture of Mycobacterium smegmatis.* Microbiology (Reading, England), 2015. **161**(Pt 2): p. 341-353.

9. Nagaraja, V., et al., *DNA topoisomerase I and DNA gyrase as targets for TB therapy.* Drug Discov Today, 2017. **22**(3): p. 510-518.

447   10.   Ravishankar, S., et al., *Genetic and chemical validation identifies Mycobacterium*

448   *tuberculosis topoisomerase I as an attractive anti-tubercular target.* Tuberculosis

449   (Edinb), 2015. **95**(5): p. 589-98.

450   11.   Yu, X., et al., *Synthesis, evaluation, and CoMFA study of fluoroquinophenoxazine*

451   *derivatives as bacterial topoisomerase IA inhibitors.* Eur J Med Chem, 2017. **125**: p. 515-

452   527.

453   12.   Fujimoto-Nakamura, M., et al., *Accumulation of mutations in both gyrB and parE genes*

454   *is associated with high-level resistance to novobiocin in Staphylococcus aureus.*

455   Antimicrob Agents Chemother, 2005. **49**(9): p. 3810-5.

456   13.   Annamalai, T., et al., *Analysis of DNA relaxation and cleavage activities of recombinant*

457   *Mycobacterium tuberculosis DNA topoisomerase I from a new expression and*

458   *purification protocol.* BMC Biochem, 2009. **10**: p. 18.

459   14.   Cao, Y., et al., *Investigating mycobacterial topoisomerase I mechanism from the analysis*

460   *of metal and DNA substrate interactions at the active site. Nucleic Acids Res.* 2018.

461   **46***:*7296-7308.

462   15.   Ren, J. and J.B. Chaires, *Sequence and structural selectivity of nucleic acid binding*

463   *ligands.* Biochemistry, 1999. **38**(49): p. 16067-75.

464   16.   Gu, M., et al., *Fluorescently labeled circular DNA molecules for DNA topology and*

465   *topoisomerases.* Sci Rep, 2016. **6**: p. 36006.

466   17.   Stephan, J., et al., *The growth rate of Mycobacterium smegmatis depends on sufficient*

467   *porin-mediated influx of nutrients.* Mol Microbiol, 2005. **58**(3): p. 714-30.

468   18.   Stahl, C., et al., *MspA provides the main hydrophilic pathway through the cell wall of*

469   *Mycobacterium smegmatis.* Mol Microbiol, 2001. **40**(2): p. 451-64.

470   19.   Stephan, J., et al., *Multidrug resistance of a porin deletion mutant of Mycobacterium*

471   *smegmatis.* Antimicrob Agents Chemother, 2004. **48**(11): p. 4163-70.

472   20.   Danilchanka, O., M. Pavlenok, and M. Niederweis, *Role of porins for uptake of*

473   *antibiotics by Mycobacterium smegmatis.* Antimicrob Agents Chemother, 2008. **52**(9): p.

474   3127-34.

475   21.   Niederweis, M., et al., *Cloning of the mspA gene encoding a porin from Mycobacterium*

476   *smegmatis.* Mol Microbiol, 1999. **33**(5): p. 933-45.

477   22.   Kartmann, B., S. Stenger, and M. Niederweis, *Porins in the cell wall of Mycobacterium*

478   *tuberculosis.* J Bacteriol, 1999. **181**(20): p. 6543-6.

479   23.   Adams, R., et al., *Binding sites in membrane proteins--diversity, druggability and*

480   *prospects.* Eur J Cell Biol, 2012. **91**(4): p. 326-39.

481   24.   Overington, J.P., B. Al-Lazikani, and A.L. Hopkins, *How many drug targets are there?*

482   Nat Rev Drug Discov, 2006. **5**(12): p. 993-6.

483   25.   Delmar, J.A., et al., *Structural Basis for the Regulation of the MmpL Transporters of*

484   *Mycobacterium tuberculosis.* J Biol Chem, 2015. **290**(47): p. 28559-74.

485   26.   Nessar, R., et al., *Mycobacterium abscessus: a new antibiotic nightmare.* J Antimicrob

486   Chemother, 2012. **67**(4): p. 810-8.

487   27.   Ishiekwene, C., et al., *Case report on pulmonary disease due to coinfection of*

488   *Mycobacterium tuberculosis and Mycobacterium abscessus: Difficulty in diagnosis.*

489   Respir Med Case Rep, 2017. **20**: p. 123-124.

490   28.   Sohn, S., et al., *Mixed Infection of Mycobacterium abscessus subsp. abscessus and*

491   *Mycobacterium tuberculosis in the Lung.* Korean J Thorac Cardiovasc Surg, 2017. **50**(1):

492   p. 50-53.

29.   Mailaender, C., et al., *The MspA porin promotes growth and increases antibiotic*

susceptibility of both Mycobacterium bovis BCG and Mycobacterium tuberculosis.

Microbiology, 2004. **150**(Pt 4): p. 853-64.

30.   Cavalli, Z., et al., *High incidence of non-tuberculous mycobacteria-positive cultures*

among adolescent with cystic fibrosis. J Cyst Fibros, 2017. **16**(5): p. 579-584.

31.   Korepanova, A., et al., *Cloning and expression of multiple integral membrane proteins*

from Mycobacterium tuberculosis in Escherichia coli. Protein Sci, 2005. **14**(1): p. 148-

58.

32.   Ding, C., et al., *Identification of mycobacterial membrane proteins and their types using*

over-represented tripeptide compositions. J Proteomics, 2012. **77**: p. 321-8.

33.   He, Z. and J. De Buck, *Cell wall proteome analysis of Mycobacterium smegmatis strain*

MC2 155. BMC Microbiol, 2010. **10**: p. 121.

34.   Domenech, P., M.B. Reed, and C.E. Barry, 3rd, *Contribution of the Mycobacterium*

tuberculosis MmpL protein family to virulence and drug resistance. Infect Immun, 2005.

73(6): p. 3492-501.

35.   Sandhu, P. and Y. Akhter, *The internal gene duplication and interrupted coding*

sequences in the MmpL genes of Mycobacterium tuberculosis: Towards understanding

the multidrug transport in an evolutionary perspective. Int J Med Microbiol, 2015.

305(3): p. 413-23.

36.   Strnad, L. and K.L. Winthrop, *Treatment of Mycobacterium abscessus Complex.* Semin

Respir Crit Care Med, 2018. **39**(3): p. 362-376.

37.   Pacheco, S.A., et al., *MmpL11 protein transports mycolic acid-containing lipids to the*

515   *mycobacterial cell wall and contributes to biofilm formation in Mycobacterium*

516   *smegmatis.* J Biol Chem, 2013. **288**(33): p. 24213-22.

517   38.   Purdy, G.E., M. Niederweis, and D.G. Russell, *Decreased outer membrane permeability*

518   *protects mycobacteria from killing by ubiquitin-derived peptides.* Mol Microbiol, 2009.

519   **73**(5): p. 844-57.

520   39.   Akerman, K.J., et al., *Gold(III) macrocycles: nucleotide-specific unconventional catalytic*

521   *inhibitors of human topoisomerase I.* J Am Chem Soc, 2014. **136**(15): p. 5670-82.

524 **Table 1.** FP-11g resistant mutants, resistance levels and cross-resistance to Moxifloxacin

M. smegmatis Strain	FP-11g concentration for mutant isolation X MIC (0.31 µM)	FP-11g MIC (µM)	Fold-increase in FP-11g MIC	Moxifloxacin MIC (µg/ml)	Fold-increase in Moxifloxacin MIC
WT	N/A	0.31	N/A	0.08-0.16	N/A
PGM1	8X → 16X	2.5	8X	0.16	2X
PGM2	8X → 16X	5	16X	0.32	4X
PGM3	4X → 10X	5	16X	0.32	4X
PGM4	4X → 13X	10	32X	0.32	4X
PGM5	4X → 16X	5	16X	0.32	4X
PGM6	4X → 16X	5	16X	0.16	2X

534   **Table 2.** Mutations identified in each FP-11g resistant mutant

Strain ID	Region	Gene	Type	Nucleotide		Amino acid		Functional annotation in *M. smegmatis*
				Ref	Allele	Ref	Allele	
PGM1	270553^270554	MSMEG_0241	INS	-	C	L	fs	MmpL11 protein. Function unknown
	1039173	MSMEG_0965	SNV	T	C	L	P	Porin MspA
	6498888^6498889	MSMEG_6430	INS	-	C	S	fs	Hypothetical protein. Function unknown. Integral component of membrane
PGM2	270553^270554	MSMEG_0241	INS	-	C	L	fs	MmpL11 protein. Function unknown
	1039173	MSMEG_0965	SNV	T	C	L	P	Porin MspA
	4715711^4715712	MSMEG_4629	INS	-	T			pseudogene
	6498888^6498889	MSMEG_6430	INS	-	C	S	fs	Hypothetical protein. Function unknown. Integral component of membrane
	6498891^6498892	MSMEG_6430	INS	-	C	D	fs	Hypothetical protein. Function unknown. Integral component of membrane
	6498895	MSMEG_6430	SNV	G	C	S	C	Hypothetical protein. Function unknown. Integral component of membrane
	6498901	MSMEG_6430	SNV	G	T	T	K	Hypothetical protein. Function unknown. Integral component of membrane
PGM3	1013796	MSMEG_0933	DEL	C	-	R	fs	Conserved hypothetical protein. Function unknown
	1039098	MSMEG_0965	SNV	C	T	T	I	Porin MspA
	2883459	MSMEG_2820	SNV	T	G	I	S	Hypothetical protein. Function unknown. Integral component of membrane
	5707525	MSMEG_5623	SNV	C	A	D	Y	L-carnitine dehydratase. Function unknown
PGM4	1039098	MSMEG_0965	SNV	C	T	T	I	Porin MspA
	1604708	MSMEG_1513	SNV	G	C	S	C	Conserved hypothetical protein. Function unknown. oxidoreductase activity
PGM5	269640^269641	MSMEG_0240	INS	-	A	M	fs	Conserved hypothetical protein. Function unknown
	1039098	MSMEG_0965	SNV	C	T	T	I	Porin MspA
	1604708	MSMEG_1513	SNV	G	C	S	C	Conserved hypothetical protein. Function unknown. oxidoreductase activity
	5707525	MSMEG_5623	SNV	C	A	D	Y	L-carnitine dehydratase. Function unknown
PGM6	1039098	MSMEG_0965	SNV	C	T	T	I	Porin MspA
	3612348	MSMEG_3552	SNV	C	G	E	Q	Conserved hypothetical protein. Function unknown
	5707525	MSMEG_5623	SNV	C	A	D	Y	L-carnitine dehydratase. Function unknown

	6498888^6498889	MSMEG_6430	INS	-	C	S	fs	Hypothetical protein. Function unknown. Integral component of membrane
	6498891^6498892	MSMEG_6430	INS	-	C	D	fs	Hypothetical protein. Function unknown. Integral component of membrane
	6498895	MSMEG_6430	SNV	G	C	S	C	Hypothetical protein. Function unknown. Integral component of membrane
	6498898^6498899	MSMEG_6430	INS	-	T	P	fs	Hypothetical protein. Function unknown. Integral component of membrane
	6498901	MSMEG_6430	SNV	G	T	T	K	Hypothetical protein. Function unknown. Integral component of membrane

535   Ref: reference; SNV: single nucleotide variation; INS: insertion; DEL: deletion; fs: frame shift

537   **Table 3.** Optical properties of FP-11g in the presence and absence of ST DNA in 1×BPE buffer
538

FP-11g	λ_{max} (nm)	ε_{334} $M^{-1}cm^{-1}$	ε_{386} $M^{-1}cm^{-1}$	ε_{406} $M^{-1}cm^{-1}$	$\Delta T_m{}^a$ (°C)	Relative fluorescence[b]
Free	386	4,646	6,033	5,587	N/A	1.0
Bound	406	12,321	5,284	4,418	11.1	1.5
	334					

539   [a]$\Delta T_m = T_m - T_m{}^0$ where T_m and $T_m{}^0$ represent the DNA UV melting temperature in the presence

540   and absence of 40 µM of FP-11g (Fig. 4C).

541   [b]At $\lambda_{em} = 550$ nm with $\lambda_{ex} = 386$ nm. The fluorescence intensity is relative to that of free FP-11g.

Scheme 1. Structure of FP-11g

Scheme 1

A

Survival (%) vs Time (h)

Legend: 2.5 μM, 1.25 μM, 0.625 μM, 0.31 μM

B

Survival (%) vs Time (h)

Legend: 100 μM, 50 μM

Fig 1

A

0   0.06   0.12   0.24   0.48   0.96   μM FP-11g

Nicked + Relaxed
Partially Relaxed
Supercoiled

1   2   3   4   5   6   7

B

0   3.9   7.8   15.6   31.5   62.5   125   μM FP-11g

Nicked + Relaxed
Partially Relaxed
Supercoiled

1   2   3   4   5   6   7   8

Fig 2

Fig 3

Fig 4

Fig 5