Protein composition of the occlusion bodies of *Epinotia aporema* granulovirus

Tomás Masson, María Laura Fabre, María Leticia Ferrelli, Matías Luis Pidre, Víctor Romanowski.

1 Instituto de Biotecnología y Biología Molecular (IBBM, UNLP-CONICET), Facultad de Ciencias Exactas, Universidad Nacional de La Plata, La Plata, Buenos Aires, Argentina.

Abstract

Within family *Baculoviridae*, members of the *Betabaculovirus* genus are employed as biocontrol agents against lepidopteran pests, either alone or in combination with selected members of the *Alphabaculovirus* genus. *Epinotia aporema* granulovirus (EpapGV) is a fast killing betabaculovirus that infects the bean shoot borer (*E. aporema*) and is a promising biopesticide. Because occlusion bodies (OBs) play a key role in baculovirus horizontal transmission, we investigated the composition of EpapGV OBs. Using mass spectrometry-based proteomics we could identify 56 proteins that are included in the OBs during the final stages of larval infection. Our data provides experimental validation of several annotated hypothetical coding sequences. Proteogenomic mapping against genomic sequence detected a previously unannotated ac*110*-like core gene and a putative translation fusion product of ORFs *epap48* and *epap49*. Comparative studies of the proteomes available for the family *Baculoviridae* highlight the conservation of core gene products as parts of the occluded virion. Two proteins specific for betabaculoviruses (Epap48 and Epap95) are incorporated into OBs. Moreover, quantification based on emPAI values showed that Epap95 is one of the most abundant components of EpapGV OBs.

Introduction

The family *Baculoviridae* comprises a diverse group of large double stranded DNA

viruses that infect larvae of the insect orders *Lepidoptera*, *Hymenoptera* and *Diptera*

[1]. Baculovirus have a rod shaped, enveloped virion with a circular genome ranging

from 80 to 180 kbp [2]. Virions are found in the environment embedded in a

proteinaceous matrix that forms occlusion bodies (OBs), a phenotype that is resistant

to desiccation and UV radiation. OBs on leaves that are consumed by foraging larvae

reach the midgut and, after being dissolved at high pH, release the occlusion derived

viruses (ODVs), which initiate infection of the epithelial cells. These infected cells

produce budded viruses (BVs) that disseminate the infection systemically [3]. Based on

OBs morphology, baculoviruses were first classified in two groups:

Nucleopolyhedrovirus (NPV) and Granulovirus (GV) [1]. Later, they were taxonomically

divided into four genera: *Alphabaculovirus* (lepidopteran-specific NPV),

Betabaculovirus (lepidopteran-specific GV), *Gammabaculovirus* (hymenopteran-

specific NPV) and *Deltabaculovirus* (dipteran-specific NPV) [1].

Among different entomopathogenic viruses, the baculoviruses have received most of

the attention due to their narrow host range which makes them safe pesticides. The

majority of commercial products are based on virus isolates that belong to the genera

Alphabaculovirus and *Betabaculovirus* [4]. The bean shoot borer (*Epinotia aporema*) is

an oligophagous pest that attacks soybean crops [5]. A poliorganotropic fast killing

betabaculovirus for this species, *Epinotia aporema* granulovirus (EpapGV), has been

discovered and sequenced by our group [6, 7]. In order to improve our understanding

of the infectious process we set out to analyze the protein content of EpapGV OB using

a proteomic approach.

Mass spectrometry-based (MS) proteomics represents a powerful technique to

interrogate the structural landscape of viral particles [8]. In addition to direct protein

identification, spectral data derived from proteomic experiments can be used to

identify novel features within genomic and transcriptomic datasets. This

proteogenomic methodology is independent of reference annotation, thus providing

excellent means for the refinement of gene models and the discovery of novel protein

coding sequences [9].

Virion proteomics has been applied to study several DNA virus families (*Ascoviridae*

[10], *Herpesviridae* [11], *Iridoviridae* [12], *Nudiviridae* [13] and *Poxviridae* [14]). In

relation with the present study, eight ODV proteomes of baculoviruses have been

analyzed (AcMNPV [15], AgMNPV [16], ChchNPV [17], HearNPV [18], MabrNPV [19],

ClanGV [20], PiraGV [21] and CuniNPV [22]). These datasets point at a complex virion

comprising a large number of proteins involved in virion morphogenesis, OBs

formation and infection of insect midgut epithelial cells.

In this work we examined the protein content of EpapGV OBs using MS-based shotgun

proteomics in order to describe the composition of this virion phenotype. A total of 56

viral proteins from EpapGV OBs were identified, the majority of which are conserved

components among the members of the family *Baculoviridae* and few are

betabaculovirus-specific proteins. Comparative proteomics of baculovirus showed a

set of core gene products present in the majority of proteomes analyzed.

Materials and Methods

Larvae and virus

Larvae of the bean shoot borer (*Epinotia aporema*), were collected from field in the

experimental station of the Instituto Nacional de Tecnología Agropecuaria (INTA) and

reared in our laboratory with an artificial diet and controlled light cycle (16 hours of

light). The strain used in this study was EpapGV (Refseq ID NC_018875), collected in

Oliveros (Santa Fe, Argentina) [6].

Occlusion bodies (OBs) production and purification

Fourth instar *E. aporema* larvae were infected *per os* using artificial diet contaminated

with a solution containing EpapGV OBs. Dying larvae with signs of infection were

stored and processed as described previously [7]. Briefly, infected larvae were stored

in distilled water and later homogenized in a Dounce homogenizer. The resulting

solution was filtered through three layers of cheesecloth to eliminate insoluble insect

debris. This extract was clarified by three steps of centrifugation at 10000 x g for 10

minutes followed by a wash with 0.05% v/v SDS solution. Clarified solution was

subjected to ultracentrifugation in a continuous 30-60% w/w sucrose gradient (50000 x

g, one hour, 4°C, Beckman SW 41 Ti rotor). The whitish/opalescent band corresponding

to OBs was collected, diluted 10-fold in distilled water and pelleted by centrifugation at

14000 x g for 10 minutes. The final pellet was resuspended in distilled water and

stored frozen at -20°C. Two biological independent samples were processed. Total

protein mass in the sample was quantified using the Bradford assay [23].

Mass spectrometry analysis

Protein digestion and analysis were performed at the Proteomics Core Facility

CEQUIBIEM, at the University of Buenos Aires/CONICET (National Research Council) as

follows: protein samples were reduced with 10 mM dithiothreitol in 50 mM

ammonium bicarbonate pH 8 (45 min, 56°C) and carbamidomethylated with 20 mM

iodoacetamide in the same solvent (40 min, room temperature, in darkness). This

protein solution was precipitated with 0.2 volumes of 100% w/v trichloroacetic acid

(Sigma) at -20 °C for at least two hours and centrifuged at 12000 x g for 10 min (4°C).

The pellet was washed twice with ice-cold acetone and dried at room temperature.

Proteins were resuspended in 50 mM ammonium bicarbonate pH 8 and digested with

trypsin (Promega V5111). The resulting peptides were desalted with ZipTip C18

columns (Millipore).

The digests were analyzed by nanoLC-MS/MS in a Thermo Scientific Q Exactive Mass

Spectrometer coupled to a nanoHPLC EASY-nLC 1000 (Thermo Scientific). For the LC-

MS/MS analysis, approximately 1 µg of peptides was loaded onto the column and

eluted for 120 minutes using a reverse phase column (C18, 2 µm x 10 nm particle size,

50 µm x 150 mm) Easy-Spray Column PepMap RSLC (P/N ES801) suitable for separating

complex mixtures of peptides with a high degree of resolution. The flow rate used for

the nano-column was 300 nL min^{-1} and the solvent range from 7% B (5 min) to 35% B

(120 min). Solvent A was 0.1% formic acid in water whereas B was 0.1% formic acid in

acetonitrile. The injection volume was 2 µL. The MS equipment has a high collision

dissociation cell (HCD) for fragmentation and an Orbitrap analyzer (Thermo Scientific,

Q-Exactive). A voltage of 3.5 kV was used for Electro Spray Ionization (Thermo

Scientific, EASY-SPRAY).

XCalibur 3.0.63 (Thermo Scientific) software was used for data acquisition and

equipment configuration that allows peptide identification at the same time of their

chromatographic separation. Full-scan mass spectra were acquired in the Orbitrap

analyzer. The scanned mass range was 400-1800 m/z, at a resolution of 70000 at 400

m/z and the 12 most intense ions in each cycle, were sequentially isolated, fragmented

by HCD and measured in the Orbitrap analyzer. Peptides with a charge of +1 or with

unassigned charge state were excluded from fragmentation for MS2.

Analysis of MS data

Q Exactive raw data was processed using Proteome Discoverer^{TM} software (version

2.1.1.21, Thermo Scientific) and searched against EpapGV protein database

downloaded from NCBI (accession number NC_018875, National Center for

Biotechnology Information; www.ncbi.nlm.nih.gov) digested with trypsin with a

maximum of one missed cleavage per peptide. Proteome Discoverer^{TM} searches were

performed with a precursor mass tolerance of 10 ppm and product ion tolerance of

0.05 Da. Static modifications were set to carbamidomethylation of Cys, and dynamic

modifications were set to oxidation of Met and N-terminal acetylation. Protein hits

were filtered for high confidence peptide matches with a maximum protein and

peptide false discovery rate of 1% calculated using a reverse database strategy. The

exponentially modified protein abundance index (emPAI) was calculated automatically

by Proteome Discoverer^{TM} software and used to estimate the relative abundance of

identified proteins within the sample.

Non annotated peptides search

The complete genome sequence of EpapGV was translated *in silico* in all six frames

using the Mascot search software. Spectral data was searched and all peptides hits

were filtered to discard matches in previously annotated ORFs. The remaining peptides

were mapped to the corresponding genomic position. Search for putative unannotated

ORFs was done extending peptide hits until a stop codon was found at C-terminus, and

a start or stop codon for the N-terminus. Homologous sequences were searched using

the TBLASTN tool against all baculovirus genomes.

Orthologs clustering

A database comprising all the ODV proteins detected in baculoviruses was generated

using previous proteomic data sets [15-22]. The software BLASTP [24] and HHMER [25]

were used to identify groups of orthologous proteins (orthogroups) among different

proteomes by reciprocal best hits.

Results

Structural components of the EpapGV OB

We determined the composition of purified EpapGV OBs employing a shotgun

proteomic approach. The peptide mixture was separated by liquid chromatography

and analyzed with tandem mass spectrometry (LC-MS/MS). This approach was used to

avoid protein loss associated with SDS-PAGE gel extraction. We detected 56 proteins in

our purified EpapGV OBs samples. Genes encoding these proteins comprise 43.93% of

EpapGV total number of annotated ORFs (Fig 1), showing that a large part of the viral

genome codes for structural proteins. A total of 10 proteins (Epap10, Epap62, Epap71,

Epap123, LEF6, P6.9, Hel-1, P18, DNA Polymerase and DNA Ligase) were detected with

only a single peptide, which might be related to low molar proportions of these

polypeptides in the sample (Table 1). As additional evidence for the identification of

these proteins, we checked the presence of several ion products belonging to the

theoretical *b* and *y* spectral ions series for these peptides. The full list of proteins is

shown in Table 1. In our samples we were unable to detect PIF3 and desmoplakin, two

core gene products which have been confirmed in other virions by MS and western

blot [15]. This could be attributed to proteolytic degradation, low protein level or

deficient ionization of these proteins in our samples.

Fig 1. Genomic localization of ORFs coding for proteins found in EpapGV OBs

Proteins identified from spectral data (green arrows) were mapped to their respective

genomic coordinate (grey line). The novel identified Ac110-like peptide is highlighted

in red script. The baculovirus collinearity region is shown in black.

180   **Table 1. Proteins detected in EpapGV OBs**

ORF	Protein	NCBI Protein Id.	Size (aa)	% Coverage	# Peptides	emPAI	% emPAI
1	Granulin	YP_006908509.1	248	60,48	25	1995261	857918

5	P78/83	YP_006908513.1	137	16,06	2	0,33	0,14
6	PK1	YP_006908514.1	276	30,43	9	3,44	1,48
10	Epap10	YP_006908518.1	90	23,33	1	0,33	0,14
14	EFP	YP_006908522.1	541	5,18	1	0,08	0,03
21	PEP2	YP_006908529.1	142	45,07	5	4,62	1,99
22	PEP/P10	YP_006908530.1	308	48,70	8	9	3,87
25	PEP1	YP_006908533.1	178	37,64	6	5,31	2,28
27	**PIF5**	YP_006908535.1	354	31,07	7	5,58	2,40
28	**Ac142**	YP_006908536.1	457	44,86	18	4,84	2,08
29	**ODV-E18**	YP_006908537.1	88	45,45	6	176,83	76,03
31	v-Cath	YP_006908539.1	329	5,77	2	0,29	0,12
39	ODV-E66	YP_006908547.1	654	33,64	14	5,31	2,28
40	Epap40	YP_006908548.1	102	75,49	6	16,78	7,22
47	**PIF2**	YP_006908555.1	374	20,05	7	0,69	0,30
48	Epap48	YP_006908556.1	446	37,67	13	2,54	1,09
49	Epap49	YP_006908557.1	1465	29,83	35	2,88	1,24
52	v-Ubi	YP_006908560.1	93	33,33	3	1,37	0,59
53	**ODV-EC43**	YP_006908561.1	348	47,41	12	6,20	2,67
58	SOD	YP_006908566.1	183	55,19	9	24,12	10,37
59	**PIF0**	YP_006908567.1	653	11,02	5	0,47	0,20
62	Epap62	YP_006908570.1	106	10,38	1	0,39	0,17
66	P24	YP_006908574.1	165	27,88	3	0,87	0,37
69	**PIF1**	YP_006908577.1	538	7,99	3	0,52	0,22
71	Epap71	YP_006908579.1	104	6,73	1	0,47	0,20
74	LEF6	YP_006908582.1	82	12,19	1	0,47	0,20
78	**P48/45**	YP_006908586.1	380	6,58	2	0,10	0,04
79	P12	YP_006908587.1	115	38,26	3	4,62	1,99
80	**P40**	YP_006908588.1	373	25,20	9	1,68	0,72
81	**P6.9**	YP_006908589.1	56	14,29	1	2,16	0,93
83	**38K**	YP_006908591.1	295	26,44	5	1,45	0,62
84	**PIF4**	YP_006908592.1	162	20,99	3	0,67	0,29
85	**Hel-1**	YP_006908593.1	1085	1,29	1	0	0,00
86	**ODV-E25**	YP_006908594.1	213	43,19	7	9	3,87
87	**P18**	YP_006908595.1	158	5,06	1	0	0.00
88	**P33**	YP_006908596.1	254	36,61	7	1,98	0,85
90	ChaB	YP_006908598.1	75	45,33	2	2,98	1,28
92	**VP39**	YP_006908600.1	293	75,43	21	232,27	100,00
93	**ODV-EC27**	YP_006908601.1	284	24,65	8	3,39	1,46
94	BRO	YP_006908602.1	359	5,85	2	0,23	0,10
95	Epap95	YP_006908603.1	73	69,86	5	176,83	76,03
96	**PIF8**	YP_006908604.1	567	14,10	7	1,22	0,52
98	**Ac81**	YP_006908606.1	191	23,04	5	1,61	0,69
99	**GP41**	YP_006908607.1	286	74,82	23	87,59	37,66
100	**Ac78**	YP_006908608.1	88	44,32	2	16,78	7,22
101	**VLF-1**	YP_006908609.1	368	35,87	11	3,06	1,32
103	Ac75	YP_006908611.1	149	56,38	9	7,11	3,06
106	**DNA Pol**	YP_006908614.1	1068	2,72	1	0	0.00
109	**PIF6**	YP_006908617.1	148	41,22	5	2,59	1,11
113	FP25K	YP_006908621.1	146	19,86	3	0,23	0,10
115	DNA Lig	YP_006908623.1	534	4,87	1	0,08	0,03
123	Epap123	YP_006908630.1	102	6,87	1	0,47	0,20

126	Epap126	YP_006908633.1	347	45,82	11	3,49	1,50
127	Epap127	YP_006908634.1	69	30,43	2	1,15	0,49
129	**VP1054**	YP_006908636.1	344	11,05	2	0,1	0,04
133	ME53	YP_006908640.1	373	9,91	2	0,22	0,09
-	**Ac110**	This study	47	29.79	1	-	-

% Coverage: percentage of the protein sequence covered by identified peptides.

emPAI: exponentially modified Protein Abundance Index.

ORF: Open Reading Frame, as numbered in EpapGV genome map published by Ferrelli *et al* [7].

Peptides: number of individual peptides identified for each protein

Core gene products are in bold characters.

In addition to identifying the components of the OBs, we estimated the relative

abundance of each protein; to this end we calculated the emPAI value proposed by

Ishihama *et al* [26] for each protein. The emPAI value for the major capsid protein

VP39 was used to normalize protein abundance (Table 1). Taking a cutoff value of at

least 10 % VP39 emPAI, the most abundant proteins are GP41, Granulin, ODV-E18, SOD

and Epap95, together with VP39. The major capsid component VP39, the tegument

protein GP41 and the major component of OB matrix granulin were expected to be

among the most abundant proteins due to their known structural function. ODV-E18

ortholog in AcMNPV is an essential protein for BV production that also localizes to the

ODV membrane [27]. Cu-Zn superoxide dismutase (SOD) activity in virion preparations

of Chlorovirus PBCV-1 has been recently associated with reactive oxygen species

reduction during the early stages of virus infection [28]. Finally, Epap95 a protein with

orthologs in all the members of the genus *Betabaculovirus*, has been consistently

detected in the granuloviruses of *Clostera anachoreta* (ClanGV) and *Pieris rapae*

(PiraGV) as a component within ODVs.

Betabaculovirus gene content remains poorly characterized, with a large number of

hypothetical genes predicted by phylogenomics methods. Some of these genes lack

bona fide experimental evidence to confirm the actual existence of their putative

protein products. Our proteomic data confirmed the presence of translation products

for 10 hypothetical proteins annotated in the genome of EpapGV, namely, Epap10,

Epap40, Epap48, Epap49, Epap62, Epap71, Epap95, Epap123, Epap126 and Epap127.

Short peptides encoded in EpapGV genome that do not belong to annotated ORFs

To identify possible unannotated proteins, we searched our spectral data against a

theoretical database comprising all translation products predicted in the six reading

frames of EpapGV genome sequence (we included all possible ORFs, without

introducing a minimal size criterion). We detected seven peptides that mapped to the

EpapGV genome but did not belong to the set of annotated ORFs [7]. Their sequence

and genomic location are detailed in S1 Appendix. One of these peptides matches an

unannotated 47 amino acids long ORF overlapping *epap51* but in the opposite

orientation. We further examined the presence of this novel ORF in other members of

the family *Baculoviridae* and found that it is an ortholog of the core gene *ac110* [29].

This gene has been described as the *per os* infectivity factor 7 (*pif7*) and its product has

only been detected in the proteome of HearNPV ODV [18] and EpapGV (this study).

Genomic localization and orientation of this *ac110*-like gene is conserved within

Betabaculovirus, providing additional evidence about its evolutionary conservation.

The remaining six peptides overlap with annotated ORFs (*chitinase*, *dna ligase* and

granulin) or intergenic regions. Two peptides were found between ORFs *epap48* and

epap49 and one peptide between *epap61* and *epap62*. TBLASTN was used to find

putative homologous unannotated peptides in other baculovirus genomes. Only the

peptides overlapping with *chitinase* and *granulin* are conserved in homologous *loci* in

GV and NPV genomes (S1 Appendix).

Remarkably, peptides between *epap48* and *epap49* almost cover the entire 145 bp

intergenic sequence (Fig 2). *Epap48* encodes a 446 amino acid long protein that is

conserved in the betabaculoviruses. The putative translation product of *epap49* is a

large protein composed of 1465 residues with no orthologs detected in other

baculoviruses. Mapped peptides are located in the same reading frame as the

translation product of *epap49*, but no methionine codon has been found in frame (Fig

2). One hypothesis that could explain the presence of these peptides is that Epap48

and Epap49 may be expressed as a fusion protein due to a putative +1 frameshifting

event near the C-terminus of Epap48; further experimental validation of this potential

fusion protein will be needed to confirm this hypothesis.

Fig 2. Putative fusion protein Epap48-Epap49

(A) Genomic locus containing *epap48* (green) and *epap49* (blue) genes. The two

peptides detected by MS inside the intergenic region are depicted in purple. (B)

Genome sequence and translation frame for each gene. Start and stop codons are

shown in red (nucleotide numbers are those from NCBI accession number

NC_018875).

Conservation of structural proteins in the family Baculoviridae

The reports of ODV proteomes belonging to several baculoviruses were used to

evaluate the conservation of the viral particle composition in this family. To date, eight

proteomic studies were carried out on ODV, including five members of the genus

Alphabaculovirus (AcMNPV, AgMNPV, ChchNPV, MabrNPV and HearNPV), two of the

genus *Betabaculovirus* (ClanGV and PiraGV) and one of *Deltabaculovirus* (CuniNPV).

Our study expands this data with the proteins present in the EpapGV OBs. Sequences

of proteins detected in occluded virions of baculovirus were used to construct groups

of orthologous proteins (S1 Table). For each of these orthogroups we scored the

number of proteomes in which they are present as a measure of their conservation.

We assigned them a class based in the phylogenetic conservation of their coding

sequence (core, lepidopteran-specific, genus-specific and virus specific) (Fig 3A). Most

conserved protein groups (present in a larger number of proteomes) are enriched in

core and lepidopteran-specific gene products. In contrast, proteins specific to a small

set of proteomes are encoded by genus-specific and virus-specific genes.

The betabaculovirus proteomes (EpapGV, ClanGV and PiraGV) were compared using a

Venn diagram (Fig 3B). From the proteins present in all three viruses, BRO, Epap48,

Epap95 and Epap126 are the only orthogroups without functional characterization.

Interestingly, Epap95 is one of the most abundant proteins according to emPAI values.

Additionally, Epap126 is shared between group II alphabaculoviruses and

betabaculoviruses (except for ClanGV) (S1 Table). On the other hand, Epap10, Epap49

and Epap62 are proteins unique to EpapGV OBs. Remarkably, *epap10* orthologs are

encoded only in five alphabaculoviruses that infect insects of the family *Tortricidae*,

Choristoneura fumiferana NPV, *Choristoneura occidentalis* NPV, *Choristoneura*

rosaceana NPV, *Cryptophlebia peltastica* NPV and *Epiphyas postvittana* NPV. This could

be the product of an ancestral horizontal gene transfer between alphabaculoviruses

and betabaculoviruses that coinfected the same host, based on gene conservation

evidence.

Fig 3. Baculovirus proteome comparison. (A) Protein conservation in baculovirus

occluded virion proteomes. Protein orthogroups derived from proteomics datasets

were scored according to the number of proteomes in which they were detected.

Gene class distribution (core genes, lepidopteran baculovirus-specific, genus-specific,

specie-specific) of orthologs groups within *Baculoviridae* is highlighted in different

colours. (B) Proteins detected in proteomes of betabaculoviruses were grouped in sets

of orthologs and represented using a Venn diagram. A set of 34 proteins is present in

all three viruses [20, 21, this study]; core gene products are highlighted in bold. Two of

these protein clusters being specific of *Betabaculovirus* (Epap48 and Epap95).

291   Discussion

Occluded virions are responsible for baculovirus primary infection. Proteome of OBs is

related with oral infectivity, providing relevant information about conserved

components potentially associated with midgut infection. Until now, the proteomes of

ClanGV and PiraGV ODVs have been interrogated using MS-based techniques [20, 21].

These viral species are phylogenetically distant to EpapGV [7]. We explored possible

divergence in protein composition employing a bottom-up proteomic approach to

characterize the protein content of EpapGV OBs. A diagram of the EpapGV virion

particle summarizing qualitative and semi-quantitative composition is shown in Fig 4.

Virion components can be grouped in five classes based in their localization: 18

nucleocapsid proteins, 15 ODV envelope proteins, 5 occlusion matrix proteins, 1

tegument protein and 17 proteins of undefined localization. Comparisons across virion

proteomes available for members of the family *Baculoviridae* highlighted the

conservation of several structural components forming the mature virion. On the other

hand, comparative genomics highlight the conservation of a collinear genomic region

for lepidopteran-infecting baculoviruses [30]. Combined genomics and proteomics

information suggests that this *locus*, compared to the rest of genome, is densely

populated by protein coding sequences corresponding predominantly to structural

polypeptides (Fig 1 and S1 Fig). Intriguingly, the product of the core gene desmoplakin

could not be detected in the betabaculovirus structural proteomes but it has been

reported for alphabaculoviruses; we do not know if this is related to a different

localization of this protein within betabaculovirus or due to technical reasons. In

AcMNPV, desmoplakin has been implied in the segregation of nucleocapsids destined

to build BV (which are ubiquitinated) and ODV (non ubiquitinated) [31].

Fig 4. Schematic model of EpapGV OB. Qualitative and semi-quantitative virion model

based on our proteomic data and localization described in published reports. Protein

levels were estimated using the emPAI value and expressed as relative value with

respect to the major capsid protein VP39.

It has been previously reported that proteins related to viral DNA metabolism and DNA

binding capacity may be retained in the virion [15]. In the present study we could

identify the DNA polymerase, DNA ligase and Helicase-1 in OBs. For other

baculoviruses IE1, Alk-Exo, LEF1 and LEF3 have been detected also. This reinforces the

idea that the viral DNA is associated with various proteins (in addition to the major

condensing protein P6.9) inside the viral capsid.

The envelope that surrounds the ODV morphotype is especially adapted for primary

infection of the insect midgut and presents a complex complement of proteins. These

can be classified in two functional groups, those required for virion envelopment and

those related with oral infectivity. Envelope morphogenesis begins with the formation

of intranuclear microvesicles (IMV) derived from the inner nuclear membrane and the

association with viral capsids. The ODV membrane proteins Ac75 and P18 are

necessary for the generation of these IMV [32, 33]. Subsequently, envelopment of

assembled nucleocapsids requires the ODV proteins Ac78, Ac81, Ac142, ODV-E25,

ODV-EC43, P33 and P48 to form mature OBs [34-40] . On the other hand, several ODV

membrane proteins are members of the PIF complex (PIF0, PIF1, PIF2, PIF3, PIF4, PIF5,

PIF6 and PIF8); this molecular complex is the main effector of oral infection in the

insect midgut. These proteins are encoded by core genes conserved in all the members

of the *Baculoviridae* family [41].

The biological relevance of the betabaculovirus-specific orthogroups Epap48 and

Epap95 within OBs is currently unknown. Moreover, the high content of Epap95 in the

OBs may also be biologically relevant. On the other hand, Epap126 orthogroup is

present in group II alphabaculoviruses and betabaculoviruses, which represents a

conserved protein potentially involved in oral infection.

Baculovirus genomes are densely populated with coding sequences (overlapping in

several cases) and contain short intergenic regions [2]. A recent study has described

the transcriptional landscape of baculovirus infection, demonstrating the existence of

several polycistronic and overlapping viral transcripts [42]. Together with other

technologies, proteogenomic mapping is a valuable tool to improve the annotation of

these complex coding regions. This approach has been used in proteome research for

several virus families [43], but was applied only for one baculovirus, AgMNPV [44]. We

identified seven peptides that do not map to previously annotated coding regions. One

of these peptides turned to be an ortholog of *ac110*; this ORFs overlapped with the

coding sequence of *epap51*. Moreover, the presence of peptides derived from

alternative frames inside the coding regions of *granulin* and *chitinase* raises the

question about the underlying complexity of baculovirus transcription and translation

processes.

Surprisingly, two unmapped peptides were found to be encoded in the intergenic

region of *epap48* and *epap49* and suggest the presence of a putative fusion product

between these proteins. Epap49 is the largest protein annotated in EpapGV genome; it

is 1465 amino acids long. As previously reported, it was difficult to annotate this as a

hypothetical protein due to its atypically large size, absence of homologous proteins in

Genbank and lack of known promoter motifs [7]. Also, it was noted that large proteins

were coded in similar locations in the genomes of ChocGV and HearGV, 1144 and

1279 amino acids long, respectively [45, 46]. In the case of ChocGV it was not

annotated in the genome [45] and in HearGV it was found to be a fusion of ORFs

homologous to XecnGV 47 and 48 [46]. In this study, we found evidence that Epap49 is

actually translated.

371 Conclusion

The protein composition of EpapGV OBs was interrogated using an MS-based

proteomic approach. A total of 56 proteins have been detected in the EpapGV

occluded virion, suggesting the presence of a highly conserved protein profile in

baculoviral OBs. We identified Epap95, a betabaculovirs-specific protein, as a highly

abundant capsid component. This protein represents an interesting candidate for

further functional studies to explore its role in betabaculovirus pathogenesis. Through

proteogenomic search we could detect a non-annotated coding region with a high

degree of sequence identity to *ac110*. In addition, our data strongly suggest the

translation of a putative fusion protein involving Epap48 and Epap49. Our study

highlight the usefulness of MS proteomics to characterize the protein complement of

the viral particle and the possibility to improve genome annotation through

experimental evidence for translation of predicted coding regions.

Acknowledgements

The authors thank Dr. Silvia Margarita Moreno and Dr. María Pía Valacco, from the

Centro de Estudios Químicos y Biológicos por Espectrometría de Masas (CEQUIBIEM-

CONICET-FCEN-UBA), for their help with sample preparation protocols, data

acquisition and subsequent analysis. This work was supported by grants from the

Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) and UNLP to Víctor

Romanowski.

References

1. Harrison RL, Herniou EA, Jehle JA, Theilmann DA, Burand JP, Becnel JJ, et al. ICTV Virus Taxonomy Profile: Baculoviridae. Journal of General Virology. 2018;99(9):1185-1186. pmid: 29947603.

2. van Oers MM, Vlak JM. Baculovirus genomics. Curr Drug Targets. 2007;8(10), 1051–68. pmid: 17979665

3. Passarelli AL. Barriers to success: how baculoviruses establish efficient systemic infections. Virology. 2011;411(2):383-92. pmid: 21300392

4. Haase S, Sciocco-Cap A, Romanowski V. Baculovirus insecticides in Latin America: historical overview, current status and future perspectives. Viruses. 2015;7:2230–67. pmid: 25941826

5. Sánchez NE, Pereyra PC. Neotropical soybean budborer, Crocidosema aporema (Walsingham, 1914) (Lepidoptera: Tortricidae). In J. L. Capinera (Ed.), Encyclopedia of Entomology. Dordrecht: Springer Netherlands. 2008. pp. 2587–2588. doi: https://doi.org/10.1007/978-1-4020-6359-6_2186

6. Sciocco-Cap A, Parola AD, Goldberg AV, Ghiringhelli PD, Romanowski V. Characterization of a granulovirus isolated from Epinotia aporema Wals. (Lepidoptera: Tortricidae) larvae. Appl Environ Microbiol. 2001;67(8):3702-6. pmid: 11472950

7. Ferrelli ML, Salvador R, Biedma M, Berretta M, Haase S, Sciocco-Cap, A, et al. Genome of Epinotia aporema granulovirus (EpapGV), a polyorganotropic fast killing betabaculovirus with a novel thymidylate kinase gene. BMC Genomics. 2012;13(1):548. pmid: 23051685

8. Greco TM, Diner BA, Cristea IM. The impact of mass spectrometry–based proteomics on fundamental discoveries in virology. Annu Rev Virol. 2014;1(1):581-604. pmid: 26958735

9. Nesvizhskii AI. Proteogenomics: concepts, applications and computational strategies. Nat Methods. 2014;11(11):1114–1125. pmid: 25357241

10. Tan Y, Bideshi DK, Johnson JJ, Bigot Y, Federici BA. Proteomic analysis of the Spodoptera frugiperda ascovirus 1a virion reveals 21 proteins. J Gen Virol. 2009;90:359–365. pmid: 19141444

11. Vidick S, Leroy B, Palmeira L, Machiels B, Mast J, François S,et al. Proteomic characterization of murid herpesvirus 4 extracellular virions. PLoS ONE. 2013;8(12):e83842. pmid: 24386290

12. Ince IA, Boeren SA, Van Oers MM, Vervoort JJM, Vlak JM. Proteomic analysis of Chilo iridescent virus. Virology. 2010;405(1):253–258. pmid: 20598335

13. Bézier A, Harichaux G, Musset K, Labas V, Herniou EA. Qualitative proteomic analysis of Tipula oleracea nudivirus occlusion bodies. J Gen Virol. 2017;98(2):284–295. pmid: 28284235

14. Doellinger J, Schaade L, Nitsche A. Comparison of the Cowpox Virus and Vaccinia Virus mature virion proteome: analysis of the species- and strain-specific proteome. PLOS ONE. 2015;10(11):e0141527. pmid: 26556597

15. Braunagel SC, Russell WK, Rosas-Acosta G, Russell DH, Summers MD. Determination of the protein composition of the occlusion-derived virus of Autographa californica nucleopolyhedrovirus. Proc Natl Acad Sci USA. 2003;100(17):9797-802. pmid: 12904572

16. Braconi CT, Ardisson-Araújo DM, Paes Leme AF, Oliveira JV, Pauletti BA, Garcia-Maruniak A, et al. Proteomic analyses of baculovirus Anticarsia gemmatalis multiple nucleopolyhedrovirus budded and occluded virus. J Gen Virol. 2014;95:980–989. pmid: 24443474

17. Xu F, Ince IA, Boeren S, Vlak JM, Van Oers MM. Protein composition of the

occlusion derived virus of Chrysodeixis chalcites nucleopolyhedrovirus. Virus Res. 2011;158(1-2):1–7. pmid: 21354223

18. Hou D, Zhang L, Deng F, Fang W, Wang R, Liu X, et al. Comparative proteomics reveal fundamental structural and functional differences between the two progeny phenotypes of a baculovirus. J Virol. 2013;87(2):829–39. pmid: 23115289

19. Hou D, Chen X, Zhang LK. Proteomic analysis of Mamestra Brassicae Nucleopolyhedrovirus progeny virions from two different hosts. PLOS ONE. 2016; 11(4):e0153365. pmid: 27058368

20. Zhang, X., Yin, X., Liang, Z., & Shao, X. Proteomic analysis of the occlusion-derived virus of Clostera anachoreta granulovirus. J Gen Virol. 2015;96(8):2394–2404. pmid: 25872743

21. Wang XF, Zhang BQ, Xu HJ, Cui YJ, Xu YP, Zhang MJ, et al. ODV-associated proteins of the Pieris rapae granulovirus. J Proteome Res. 2011;10(6):2817–2827. pmid: 21517121

22. Perera O, Green TB, Stevens SM Jr, White S, Becnel JJ. Proteins associated with Culex nigripalpus nucleopolyhedrovirus occluded virions. J Virol. 2007;81(9):4585-90. pmid: 17301145

23. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. 1976;72(1–2):248–54. pmid: 942051

24. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J. Mol. Biol. 1990;215(3):403-410. pmid: 2231712

25. Eddy SR. Accelerated Profile HMM Searches. PLoS Comput Biol. 2011;7(10):e1002195. pmid: 22039361

26. Ishihama Y, Oda Y, Tabata T, Sato T, Nagasu T, Rappsilber J, et al. Exponentially modified protein abundance index (emPAI) for estimation of absolute protein amount in proteomics by the number of sequenced peptides per protein. Mol Cell Proteomics. 2005;4(9):1265–1272. pmid: 15958392

27. McCarthy CB, Theilmann DA. AcMNPV ac143 (odv-e18) is essential for mediating budded virus production and is the 30th baculovirus core gene. Virology. 2008;375(1):277-291. pmid: 18328526

28. Kang M, Duncan GA, Kuszynski C, Oyler G, Zheng J, Becker DF, et al. Chlorovirus PBCV-1 encodes an active copper-zinc superoxide dismutase. J Virol. 2014;88(21):12541-12550. pmid: 25142578

29. Javed MA, Biswas S, Willis LG, Harris S, Pritchard C, van Oers MM, et al. Autographa californica multiple nucleopolyhedrovirus AC83 is a per os infectivity factor (PIF) protein required for occlusion-derived virus (ODV) and budded virus nucleocapsid assembly as well as assembly of the PIF complex in ODV envelopes. J Virol. 2017;91(5):e02115-16. pmid: 28031365

30. Zhu Z, Yin F, Liu X, Hou D, Wang J, Zhang L, et al. Genome sequence and analysis of Buzura suppressaria nucleopolyhedrovirus: a group II Alphabaculovirus. PloS One. 2014;9(1):e86450. pmid: 24475121

31. Biswas S, Willis LG, Fang M, Nie Y, Theilmann DA. Autographa californica nucleopolyhedrovirus AC141 (Exon0), a potential E3 ubiquitin ligase, interacts with viral ubiquitin and AC66 to facilitate nucleocapsid egress. J Virol. 2018;92(3):e01713-17 pmid: 29142135

32. Shi A, Hu Z, Zuo Y, Wang Y, Wu W, Yuan M, et al. Autographa californica nucleopolyhedrovirus ac75 is required for the nuclear egress of nucleocapsids and intranuclear microvesicle formation. Journal of Virology. 2017;92(4):e01509-17. pmid: 29212928

33. Yuan M, Huang Z, Wei D, Hu Z, Yang K, Pang Y. Identification of Autographa californica nucleopolyhedrovirus ac93 as a core gene and its requirement for

494 intranuclear microvesicle formation and nuclear egress of nucleocapsids. J Virol.
495 2011;85(22):11664–74. pmid: 21880748

34. Tao XY, Choi JY, Kim WJ, Lee JH, Liu Q, Kim SE, et al. The Autographa californica multiple nucleopolyhedrovirus ORF78 is essential for budded virus production and general occlusion body formation. J Virol. 2013;87(15):8441–8450. pmid: 23698311

35. Dong F, Wang J, Deng R, Wang X. Autographa californica multiple nucleopolyhedrovirus gene ac81 is required for nucleocapsid envelopment. Virus Res. 2016;221:47–57. pmid: 27212683

36. McCarthy CB, Dai X, Donly C, Theilmann DA. Autographa californica multiple nucleopolyhedrovirus ac142, a core gene that is essential for BV production and ODV envelopment. Virology. 2008;372(2):325–339. pmid: 18045640

37. Chen L, Hu X, Xiang X, Yu S, Yang R, Wu X. Autographa californica multiple nucleopolyhedrovirus odv-e25 (Ac94) is required for budded virus infectivity and occlusion-derived virus formation. Arch Virol. 2012;157(4):617–625. pmid: 22218963

38. Alfonso V, Maroniche GA, Reca SR, López MG, del Vas M, Taboga O. AcMNPV Core gene ac109 is required for budded virion transport to the nucleus and for occlusion of viral Prpgeny. PLoS ONE. 2012;7(9):e46146. pmid: 23049963

39. Wu W, Passarelli AL. Autographa californica multiple nucleopolyhedrovirus Ac92 (ORF92, P33) is required for budded virus production and multiply enveloped occlusion-derived virus formation. J Virol. 2010;84(23):12351–61. pmid: 20861245

40. Yuan M, Wu W, Liu C, Wang Y, Hu Z, Yang K, et al. A highly conserved baculovirus gene p48 (ac103) is essential for BV production and ODV envelopment. Virology. 2008;379(1):87–96. pmid: 18656219

41. Boogaard B, van Oers MM, van Lent JWM. An advanced view on baculovirus per os infectivity factors. Insects. 2018;9(3):E84 pmid: 30018247

42. Moldován N, Tombácz D, Szűcs A, Csabai Z, Balázs Z, Kis E, et al. Third-generation sequencing reveals extensive polycistronism and transcriptional overlapping in a baculovirus. Sci. Rep. 2018;8(1):8604. pmid: 29872099

43. Leroy B, Gillet L, Vanderplasschen A, Wattiez R. Structural proteomics of herpesviruses. Viruses. 2016;8(2):50. pmid: 26907323

44. Brito AF, Braconi CT, Weidmann M, Dilcher M, Alves JMP, Gruber A, et al. The Pangenome of the Anticarsia gemmatalis Multiple Nucleopolyhedrovirus (AgMNPV). Genome Biol Evol. 2015;8(1):94–108. pmid: 26615220

45. Escasa SR, Lauzon HAM, Mathur AC, Krell PJ, Arif BM. Sequence analysis of the Choristoneura occidentalis granulovirus genome. J Gen Virol. 2006;87(7):1917–33. pmid:16760394.

46. Harrison RL, Popham HJR. Genomic sequence analysis of a granulovirus isolated from the Old World bollworm, Helicoverpa armigera. Virus Genes. 2008;36(3):565–81. pmid:18418706.

536 **Supplementary information**

537 **S1 Appendix.** EpapGV unannotated peptides detected by MS.

538 **S1 Figure.** Parity plot of sequences coding for structural proteins present in AcMNPV,

539 ChchNPV and PiraGV against EpapGV.

540 **S1 Table.** Proteomic profiles of baculovirus occluded virions proteomes.

Figure

Figure

Figure

Figure