Stress activity is not predictive of coping style in North American red squirrels

Sarah E. Westrick[1*], Freya van Kesteren[1], Rupert Palme[2], Rudy Boonstra[3], Jeffery E. Lane[4], Stan

Boutin[5], Andrew G. McAdam[6], Ben Dantzer[1,7]

[1]Department of Psychology, University of Michigan, Ann Arbor, MI, USA

[2]Department of Biomedical Sciences, University of Veterinary Medicine, Vienna, Austria

[3]Department of Biological Sciences, University of Toronto Scarborough, Toronto, ON, Canada

[4]Department of Biology, University of Saskatchewan, Saskatoon, SK, Canada

[5]Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

[6]Department of Integrative Biology, University of Guelph, Guelph, ON, Canada

[7]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA

*Corresponding author: westse@umich.edu, 623-687-8615

Keywords: coping styles, glucocorticoids, North American red squirrels, behavioral syndromes

Acknowledgments

We thank Agnes MacDonald and her family for long-term access to her trapline, and the

Champagne and Aishihik First Nations for allowing us to conduct our work within their

traditional territory. We thank Adi Boon, Amanda Kelley, and Ryan W. Taylor for collecting

much of the behavioral data and all the volunteers, field assistants, and graduate students for

their assistance in data collection. This work was supported by American Society of

Mammalogists to SEW; University of Michigan to SEW and BD; National Science Foundation

(IOS-1749627) to BD; and Natural Sciences and Engineering Research Council to SB, AGM, JL,

and RB. This is publication XX of the Kluane Red Squirrel Project.

Abstract

Individuals vary in their behavioral and physiological responses to environmental changes. These

behavioral responses are often described as 'coping styles' along a proactive-reactive continuum.

Studies in laboratory populations often, but not always, find that behavioral responses and

physiological responses to stressors covary, where more proactive (more aggressive and active)

individuals have a lower physiological stress response, specifically as measured by

hypothalamic-pituitary-adrenal (HPA) axis activity. These studies support the possibility of

hormonal pleiotropy underlying the presentation of behaviors that make up the proactive-reactive

phenotype. However, recent research in wild populations is equivocal, with some studies

reporting the same pattern as found in many controlled laboratory studies, whereas others do not.

We tested the hypothesis that physiological and behavioral stress responses are correlated in wild

adult North American red squirrels (*Tamiasciurus hudsonicus*). We used fecal cortisol

metabolites (FCMs) as a non-invasive, integrated estimate of circulating glucocorticoids for our

measurement of HPA axis activity. We found that FCM concentrations were not correlated with

three measures of behavioral coping styles (activity, aggression, and docility) among individuals.

This does not support the hypothesis that hormonal pleiotropy underlies a proactive-reactive

continuum of coping styles. Instead, our results support the "two-tier" hypothesis that behavioral

and physiological stress responses are independent and uncorrelated traits among individuals in

wild populations that experience naturally varying environments rather than controlled

environments. If also found in other studies, this may alter our predictions about the evolutionary

consequences of behavioral and endocrine coping styles in free-living animals.

Significance Statement

Individuals vary in how they respond to stressors through behavior and physiology, but we find the two responses are independent in wild animals. Many laboratory studies find links between the behavioral and physiological stress responses, however studies conducted with wild populations are less conclusive. In wild North American red squirrels, independence between the physiological response and behavioral response may allow adaptive responses to a changing environment without pleiotropic constraint.

Introduction

Organisms can respond to fluctuating environmental challenges and aversive stimuli both through behavioral responses and physiological stress responses. Laboratory studies often find these responses to be associated with one another (but see Steimer and Driscoll 2003; Koolhaas et al. 2007). In behavioral ecology and behavioral neuroscience, 'coping styles' have been recognized as one method of categorizing behavioral reactions to environmental challenges and stressors. Coping styles refer to a consistent set of behavioral responses to a stressor (Gosling 2001; Réale et al. 2007; Koolhaas et al. 2010; Stamps and Groothuis 2010). Furthermore, the suite of behaviors that make up an individual's coping style is theorized to be mediated by hormones that exert pleiotropic actions (Koolhaas et al. 1999; McGlothlin and Ketterson 2008).

This unidimensional model has been repeatedly supported by studies describing how the hypothalamic-pituitary-adrenal (HPA) axis mediates coping styles (Koolhaas et al. 1999). Many of these studies have used selected lines, or have been done under controlled conditions in the laboratory. The conclusion from this model is that the behavioral stress response and physiological stress response run along the same axis. This hypothesis suggests a unidimensional

response along a proactive-reactive continuum, where 'proactive' individuals are highly

aggressive, highly active, and exhibit lower HPA axis reactivity and activity compared to

'reactive' individuals (Koolhaas et al. 1999; Cockrem 2007; Carere et al. 2010). The vast

majority of these studies have been conducted using laboratory animals or wild animals selected

for specific behavioral phenotypes, producing individuals at the extremes of this behavioral

continuum. For example, in wild Great Tits (*Parus major*) lines selected for divergent

personality types show the predicted unidimensional relationship between behavioral and stress

responses in that more proactive birds exhibited lower HPA axis reactivity in response to capture

and restraint (Baugh et al. 2012), and lower baseline corticosterone metabolites (Stöwe et al.

2010).

As more empirical studies are testing these models, the results from studies in the wild

have been equivocal. Whereas there is some support for the unidimensional model in wild

animals (see Table 1), recent studies that have used this coping style paradigm to test the

relationship between behavior and HPA axis reactivity or activity in free-living animals have

found that the proactive-reactive continuum is not predictive of the physiological stress response

(Garamszegi et al. 2012; Ferrari et al. 2013; Dosmann et al. 2015). For example, though

laboratory selection line results are consistent with predictions of the unidimensional model,

when testing Great Tits in the laboratory with natural, non-selected variation in exploratory

behavior, the relationship no longer holds (Baugh et al. 2012).

Discrepancies in the lab between recent observed relationships (Koolhaas et al. 2007) and

the simple unidimensional model (Koolhaas et al. 1999), have recently led to the development of

a 'two-tier' coping style model. This two-tier model proposes that individuals in a population can

vary independently in both behavioral responses and physiological responses to environmental

challenges (Koolhaas et al. 2010). This model of coping styles reframed the original model to

establish behavioral coping strategies on a continuum independent of physiological coping

strategies (Koolhaas et al. 2010). The distinction between the unidimensional and two-tier coping

style models is significant in assessing the ecological and evolutionary consequences of variation

in response to stressors. If the phenotypic correlation between behavioral and physiological

stress responses (assumed by the unidimensional coping styles model) reflects an underlying

correlation mediated by the effects of a hormone, this may present a limitation in the ability of

populations to adapt to changing environmental stressors (Sih et al. 2004; Dantzer and Swanson

2017). Alternatively, if the two-tier model of coping style is supported, and there are two

separate axes for behavioral and physiological stress responses, this suggests the potential for

each trait to be an independent target of selection, potentially facilitating rapid adaptation to new

environmental challenges (McGlothlin and Ketterson 2008; Ketterson et al. 2009). Exploring

how coping styles relate to the physiological stress response in wild populations allows us to test

across the entire spectrum of naturally occurring individual variation in behavioral coping styles,

thus informing our perspective on how these mechanisms function in wild populations (Réale et

al. 2007; Ferrari et al. 2013).

We investigated the relationship between three behavioral traits and one measure of HPA

axis activity (concentrations of fecal cortisol metabolites, FCM) in a natural population of North

American red squirrels (*Tamiasciurus hudsonicus*, hereafter, 'red squirrels'). Previous studies in

this species showed that there was a repeatable, correlated suite of behavioral traits, specifically

aggression, activity, and docility, across the adulthood of an individual (Boon et al. 2007; Taylor

et al. 2012). These suites of behavioral traits can also be placed along the proactive-reactive

continuum as coping styles, with the more active, aggressive, and less docile individuals at the

proactive end of the continuum. Differences in coping styles in red squirrels have clear

environment-dependent fitness correlates (Boon et al. 2007, 2008; Taylor et al. 2014), and

variation in heritable coping styles among individuals (Taylor et al. 2012) in this population may

be maintained through fluctuating selection caused by changing environmental conditions

(Taylor et al. 2014).

We used fecal samples as a non-invasive proxy for HPA axis activity and reactivity,

which is unaffected by trapping-induced stress (Dantzer et al. 2010). In red squirrels, FCM is

representative of the circulating plasma cortisol over the past ~12 hours, with a 10.9 ± 2.3 hours

lag time to peak excretion following experimental administration of cortisol (Dantzer et al.

2010). Influences of the circadian rhythm on circulating cortisol are not detected in fecal samples

collected throughout the day (Dantzer et al. 2010). Additionally, glucocorticoid concentrations in

fecal samples have been shown to be representative of HPA activity and reactivity (Sheriff et al.

2011; Palme 2019).

It is important to note that glucocorticoids are metabolic hormones and only one

mediator of the reactive physiological stress response of an individual (Romero et al. 2009).

However evidence for both the unidimensional and two-tier models of coping styles specifically

connect glucocorticoids with the behavioral response (Table 1), in addition to catecholamines

(reviewed in Koolhaas et al. 1999, 2010). While this is not a perfect measure of the overall

physiological stress response of an individual, glucocorticoids are an important physiological

mediator of the multifaceted stress response (Sapolsky et al. 2000; Romero et al. 2009).

Glucocorticoids are secreted to mobilize energy in response to a stressor in the environment, but

also exert pleiotropic effects (Sapolsky et al. 2000). For example, fluctuating baseline

glucocorticoids act as a mediator of future reproductive investment in European Starlings

(*Sturnus vulgaris*) by preparing individuals for energetically expensive reproductive seasons

(Love et al. 2014).

To test the unidimensional and two-tier models of the overall stress response, we

measured FCM as a non-invasive marker of HPA axis activity and the behavior of individuals

using three behavioral assays (open-field trial, mirror-image stimulation trial, and handling

docility assay) to measure coping style. We then compared the FCM concentrations to the

behavioral coping style of individual squirrels. A relationship between FCMs and behavioral

coping style across individuals would support the unidimensional model, whereas a lack of

relationship would support the two-tier model.

Methods

Study species

North American red squirrels are a sexually monomorphic species of arboreal squirrels

(Boutin and Larsen 1993). Females and males are both territorial of their food-cache (located on

the center of their territory) year-round (Dantzer et al. 2012; Siracusa et al. 2017). Red squirrels

in the region of our study rely on seeds produced by white spruce (*Picea glauca*) trees as their

primary food source (Fletcher et al. 2010). Squirrel population density is closely associated with

mast seeding of the white spruce, or episodes of booms and busts in food availability (McAdam

and Boutin 2003; Fletcher et al. 2010; Dantzer et al., 2013). Red squirrels have one litter per

year, with the exception of mast years when autumn spruce seed is superabundant (Boutin et al.

2006; McAdam et al. 2007).

Our study was conducted as a part of the Kluane Red Squirrel Project, a long-term study

of wild population of red squirrels within Champagne and Aishihik First Nation traditional

territory along the Alaska Highway in the southwest Yukon, Canada (61°N, 138°W). Each

squirrel was tagged with a unique set of alphanumeric stamped ear tags (National Band and Tag

Company, Newport, KY, USA). At each live-trapping (Tomahawk Live Trap, Tomahawk, WI,

USA) event, body mass and reproductive status of the squirrel were recorded. Female

reproductive status was determined through changes in body mass, by nipple condition, and by

abdominal palpations of developing fetuses in females. Male reproductive status was determined

by palpating for the presence of testicles either in the scrotum (breeding) or abdomen (non-

breeding). A more detailed description of the population and general methods can be found in

McAdam et al. (2007).

The local population of red squirrels was broken down into three study populations in

different locations: two were control populations (referred to hereafter as 'control grids') and one

was provided with supplemental food between 2004 and 2017, such that squirrel density was

increased (Dantzer et al., 2013; hereafter referred to as 'high-density grid'). Squirrels on the

high-density grid were provided with 1 kg of peanut butter (no sugar or salt added)

approximately every six weeks from October to May (Dantzer et al. 2012). We included these

squirrels to increase our sample size, and included study grid type as a covariate in all models to

control for variation between the grids. Additionally, high conspecific competition is a

significant environmental factor influencing the physiological stress response of red squirrels

(Dantzer et al. 2013) and so was important to include as a covariate in our statistical models

(Table 2). All work was conducted under the animal ethics approvals from Michigan State

University (AUF#04/08-046-00) and University of Guelph (AUP#09R006).

Behavioral trials

Squirrels were subjected to two behavioral trials to measure 'activity' and 'aggression':

an open-field (OF) trial, and a mirror image stimulation (MIS) trial (Boon et al. 2007; Taylor et

al. 2012). These behavioral trials were conducted for other projects and were not evenly

distributed across years. We performed trials in 2005, 2008, 2009, 2010, and 2012 (see Table 2

for a breakdown of sample sizes). During these years, additional trials were conducted on

squirrels in this population for multiple studies (see Boon et al. 2007; Boon et al. 2008; Kelley et

al. 2015; Taylor et al. 2012, 2014), but for the purposes of this analyses, we only included trials

for which we also had FCM concentration data for that individual. All squirrels were mature

adults (>1 year old) at the time of the trial.

To measure an individual's coping style, we conducted OF and MIS trials during the

same trapping event, with the OF trial completed first followed by the MIS trial. Squirrels

habituate to these tests (Archer 1973; Boon et al. 2008; Martin and Réale 2008), but the behavior

of individual squirrels over time is known to be repeatable (Boon et al. 2007, 2008; Taylor et al.

2014). For simplicity, we thus used only the results of each individual's first test as a measure of

its activity and aggression. We transferred focal squirrels from a live trap into the arena using a

canvas handling bag. The same portable testing arena was used for both trials, and consisted of a

60 x 80 x 50 cm white corrugated plastic box with a clear acrylic lid (Taylor 2012). Four blind

holes made with black PVC caps in the bottom of arena allowed the squirrel to explore possible

'escape routes'. We exposed a 45 x 30 cm mirror fixed to one end of the arena after the OF trial

to begin the MIS trial. A digital video camera recorded behavior in the arena. We performed all

behavioral trials on the territory of the focal individual. Between trials, we cleaned the arena

using 70% isopropyl alcohol.

To quantify behavior from the videos, we used manual scoring methods with an ethogram

developed and used in previous red squirrel studies (Boon et al. 2007, 2008; Taylor et al. 2012,

2014; Kelley et al. 2015; Supplementary Material Table S1). Because these videos were

collected and scored across multiple years, observers used different software programs

depending on what program was available the year they were scored. Trials conducted in 2005

were scored using The Observer Video-Pro 5.0 (Noldus Information Technology, Wageningen,

The Netherlands). Trials conducted from 2008-2010 were scored using JWatcher Video 1.0

(Blumstein and Daniel 2007). Trials conducted in 2012 were scored using Cowlog software

(Hänninen and Pastell 2009). Regardless of the software used, the ethogram and the overall

method of scoring the videos remained consistent. Because this is a manual process and the

software simply records keystrokes indicating behaviors observed, it is not likely that the

software used impacted the score. Furthermore, a previous study using some of our dataset

showed high inter-observer reliability for the behavioral measures we recorded from these videos

(Taylor et al., 2012), so it again seems unlikely that the software used would influence the

behavioral data we extracted from the videos. During the OF trial, we recorded the mutually

exclusive behaviors of time spent walking, sniffing, chewing, rearing, grooming, and being still.

Additionally, we recorded the number of jumps and head-dips in the false holes. During the MIS

trial, we recorded the amount of time spent in the third of the arena closest to the mirror, and the

amount of time spent in the third of the arena farthest from the mirror. We also recorded the

number of aggressive contacts with the mirror (attacks), the latency until the first attack, and the

latency until the first approach towards the mirror. A detailed description of the video scoring

methods can be found in Boon et al. (2008). Following Taylor et al. (2012), behaviors with an

inter-observer reliability of greater than 0.7 were used in analyses (see Supplementary Material

Table S1 and S2 for a list of behaviors used in the analyses).

As an additional behavioral measurement, we also measured 'docility' of individual

squirrels. In 2008 and 2009, docility measurements were collected on many squirrels for other

studies (see Boon et al. 2007; Taylor et al. 2012), but for the purposes of this study, we focused

only on trials that were conducted during a trapping event where fecal samples were also

collected and subsequently analyzed (n = 168 trapping events). We quantified docility as the

squirrel's response to handling (for examples in other species, see Carere and Oers 2004; Martin

and Réale 2008; Montiglio et al. 2012). We transferred squirrels from the trap into a canvas

handling bag and placed the squirrel on a flat surface. We measured docility during handling by

counting the number of seconds out of 30 seconds in which the squirrel was not struggling. A

squirrel that spent most of the time immobile during the test was considered docile, a trait

previously demonstrated to be repeatable (Boon et al. 2007; Taylor et al., 2012) and heritable

(Taylor et al., 2012) in this population. This test was conducted an average of 8 (min = 1, max =

42) times on 79 individual squirrels caught between 2008 and 2009. Docility scores were z-

scored for analyses. See Table 2 for detailed sample sizes.

Fecal cortisol metabolites

From 2005 to 2014, we opportunistically collected a total of 703 fecal samples during

routine trapping of squirrels with peanut butter for measurement of FCM concentrations

corresponding to individuals with behavioral data (see Dantzer et al. 2010). Fecal samples were

collected from under live-traps within two hours of trapping and placed in 1.5 mL vials stored in

a -20 °C freezer within five hours of collection. Urine contaminated feces were excluded. All

fecal samples were lyophilized for 14-16 h before being pulverized in liquid nitrogen using a

mortar and pestle. Using 0.05 g of dry ground feces, steroid metabolites were extracted by

adding 1 mL of 80% methanol and vortexing samples at 1450 RPM for 30 min, and then

centrifuging for 15 min at 2500 g (Dantzer et al. 2010; Palme et al. 2013). The resulting

supernatant was stored at -20 °C for analysis via glucocorticoid metabolite assay using a 5α-

pregnane-3β,11β,21-triol-20-one antibody enzyme immunoassay (EIA; see Touma et al. 2003).

A detailed validation and description of steroid extraction and EIA with red squirrel fecal

samples can be found in Dantzer et al. (2010).

Fecal samples were analyzed across multiple assays and in two different labs (n = 355 at

University of Toronto Scarborough and n = 348 at University of Michigan) but using the same

protocol. We confirmed that our measures of FCM concentrations were highly repeatable across

assays or labs through the following. First, a separate group of fecal samples (n = 128 samples)

were analyzed in both labs and the optical density of these samples were closely correlated

(Pearson correlation = 0.88). This indicates that the data were comparable, but we also included a

covariate in our statistical models for where the data were analyzed (see below). Second, using

pooled samples that were run repeatedly on different plates (n = 115), we found that the

estimates of optical density for these pool samples were highly repeatable ($R = 0.85$, 95% CI =

0.54-0.93). Finally, using a linear mixed-effects model, we partitioned the variance in the optical

density recorded for the pooled samples that were run across these different plates. We found

that most of the variance was due to the sample itself (85.1%) with relatively little of it being

explained by intra-assay variation as all samples were run in duplicate (4.9%) or by inter-assay

variation (9.9%). Together, this indicates that our measures of FCM concentrations should be

comparable across assays and across labs. See Table 2 for a representation of how sample sizes

were broken down in each dataset.

Statistical methods

All statistical analyses were conducted in R version 3.4.3 (R Core Team 2016). For the

OF and MIS trials, we used two principal components analyses to reduce the redundancy among

behavioral measurements and calculate composite behavioral scores for each trial, as we have

done previously in this system (Boon et al. 2007, 2008; Taylor et al. 2012, 2014; Kelley et al.

2015). To conduct the principal components analyses with correlation matrices, we used the R

package 'ade4' version 1.7-10 (Dray and Dufour 2007). By reducing the multiple behaviors

observed down to one metric for each trial, we were able to assess the primary variation among

individuals along those axes. All further analyses used the scores calculated from the principal

component loadings (Supplementary Material Table S2) for each trial. From the OF trial, we

interpreted the first principal component as a measure of overall 'activity', as it has previously

been interpreted in this population (Boon et al. 2007, 2008; Taylor et al. 2012, 2014). In our data

set, the first component explained 64% of the variation in behavior across OF trials. From the

MIS trial, we interpreted the first principal component as a measure of 'aggression', as it has also

previously been interpreted (Boon et al. 2007, 2008; Taylor et al. 2012, 2014). In our data set,

the first component explained 60% of the variation in behavior across MIS trials.

Because we were interested in an estimate of the covariance of FCM concentrations and

personality, we used a multivariate framework to conservatively address how the two types of

stress responses interact, as explained in Houslay and Wilson (2017). In this study, we were

interested in how FCM concentrations and behavioral traits co-varied among individuals. To

investigate this in a multivariate framework, we used a Bayesian generalized linear mixed effects

multivariate model based on a Markov chain Monte Carlo algorithm with the R package

'MCMCglmm' version 2.25 (Hadfield 2010) to assess the relationship between FCM

concentrations and behavior. All fecal cortisol metabolite concentration data were ln-transformed

to improve normality of residuals.

For docility analyses, we used measurements of docility paired with the FCM

concentrations of that trapping event. Using a bivariate generalized linear mixed-effects model,

we asked whether individuals with higher mean FCM concentrations have higher mean docility

scores (among-individual covariance), and whether individual observations of FCM

concentration and docility relative to the individual's mean concentrations were correlated

(within-individual covariance). Within-individual covariance indicates how the FCM

concentrations and docility scores of one individual covary across multiple observations for that

individual; in essence if we have multiple unique measurements of FCM concentration and

docility from one individual, does the docility score predict FCM concentrations on that day? In

contrast, among-individual covariance measures the relationship between FCM concentration

and docility across individuals in the population; in other words, does an individual's average

docility score predict its average FCM concentration? The bivariate model for docility and FCM

concentration included fixed effects of study grid (control or high-density), sex, Julian date of

trapping event (continuous), trial number, breeding status (breeding/non-breeding/lactating), and

a variable to indicate where the fecal sample was processed (UT Scarborough/UM). Docility

measurements were taken across multiple trapping events for a squirrel, therefore we included

trial number to control for any variation caused by habituation to the process of being trapped

and restrained in the bag (Boon et al., 2007; Taylor et al., 2012). We specified in the model to

estimate the fixed effects of Julian date of trapping event and the location of fecal sample

processing for only FCM concentration. These fixed effects were included because they have

previously been shown to influence FCM concentration in red squirrels (Dantzer et al. 2010,

2013), and thus were included to control for variation among these variables. Although the

correlation between UT Scarborough and UM samples was high (0.88), we included location of

fecal sample processing to account for any minor variation between the locations.

Because we only had one measurement of aggression and activity per individual, we were

unable to estimate the within-individual covariance between FCM concentration and the

activity/aggression of that individual. Thus, the bivariate models for activity and aggression were

structured to control for only one individual activity and aggression score per individual, and

only estimate among-individual covariance. The models for activity and aggression included the

same fixed effects as the model for docility, except trial number was not included. All trials were

the first trial in the lifetime of that individual, so there is no variance in novelty of the arena

across individuals. Again, we estimated fixed effects of the Julian date of trapping event and

location of fecal sample processing for only FCM concentration. With this model structure, we

were able to more precisely control for variation in FCM concentration due to reproductive

condition and time of year.

We fit all bivariate MCMCglmm models with uninformative priors (as in Houslay and

Wilson 2017) for 2,100,000 iterations with the first 100,000 discarded, and 1 out of every 1,000

of the remaining iterations used for parameter estimations. Credible intervals (95%) around the

correlation were based on the MCMC chain iterations. To confirm convergence using a

combination of methods, as suggested in a comparative review (Cowles and Carlin 1996), we ran

all MCMCglmm models three times for comparison using the Gelman-Rubin statistic (Gelman

and Rubin 1992), and we also ran the Geweke diagnostic (Geweke 1992). All models passed

both diagnostics for convergence.

Data availability

The datasets analyzed during the current study available from the corresponding author on

reasonable request.

Results

Our results indicate that docility, activity, and aggression did not co-vary with FCM

concentrations among individuals. Using a bivariate generalized linear mixed effects model

approach, the within-individual covariance indicated that an individual's FCM concentrations

did not correlate with docility ($r = 0.020$, CI = [-0.14, 0.28]). Our models also indicated that

among individuals, FCM concentrations did not correlate with docility ($r = 0.14$, CI = [-0.64,

0.83], Figure 1), activity (r = 0.15, CI = [-0.17, 0.47], Figure 2) or aggression (r = 0.29, CI = [-

0.098, 0.56], Figure 2) (Table 3). Regardless of the statistical significance of these relationships,

the direction of the observed effect was opposite to the predicted relationship between behavior

and FCM concentrations. The direction of these correlations suggest more active and more

aggressive squirrels may have higher FCM concentrations, but this is not conclusive due to

confidence intervals overlapping zero.

Discussion

We demonstrated that the behavioral coping style (represented by three behaviors) and

one measure of the physiological stress response (FCM concentrations) did not co-vary in a free-

ranging mammal. Independent variation between the behavioral and physiological stress

responses supports the two-tier model of coping strategies proposed by Koolhaas et al. (2010).

This model proposes that within a species, individuals can exhibit a consistent behavioral

response anywhere along the proactive-reactive continuum but independent of their

physiological stress response, which can range from a low to a high HPA axis activity. Contrary

to many studies (Raulo and Dantzer, 2018), proactive, or highly active/aggressive red squirrels

did not always exhibit lower HPA axis activity than reactive individuals. In fact, the parameter

estimates were in the opposite direction from those predicted by the unidimensional model.

Specifically, the unidimensional model predicts that a more active behavioral stress response and

HPA axis activity should be negatively correlated and we found that they were instead positively

correlated, though again these confidence intervals overlapped zero. Although we did find that

the three behavioral measures were phenotypically correlated (see Supplementary Material),

physiological stress, as measured by FCM concentration, does not appear to be the pleiotropic

mechanism causing this covariation.

Previous studies that have found behavior and HPA axis reactivity are linked have used a

different statistical framework than our study. Thus, it is possible our results are simply the

outcome of using a more conservative statistical test. However, our results were robust across

statistical techniques as we also ran the same models using a different statistical technique (linear

mixed effects models) that has been used in previous studies (e.g. Lendvai et al. 2011; Montiglio

et al. 2012). These results from the linear mixed effects models (Supplementary Material Table

S5) and those from the bivariate models (presented above) both support the hypothesis that

behavior and physiology are independent in our study.

Our study contributes to an emerging trend of a lack of a strong relationship between

behavioral and physiological stress responses in wild and laboratory animals (reviewed by Raulo

and Dantzer 2018). For example, wild alpine marmots (*Marmota marmota*) exhibit a lack of

among-individual correlation between activity and plasma cortisol concentrations, as well as

between docility and plasma cortisol (Ferrari et al. 2013). Likewise, docility and exploration

were not correlated with a change in plasma cortisol in response to a stressor in plateau pika

(*Ochotona curzoniae*: Qu et al. 2018). Additional studies measuring fecal glucocorticoid

metabolites demonstrate that HPA axis activity does not correlate with shy-bold behavioral types

in wild flycatchers (*Ficedula albicollis*; Garamszegi et al. 2012), or with exploration/activity in

Belding's ground squirrels (*Spermophilus beldingi*; Dosmann et al. 2015). In captivity, Holstein

Friesian heifer calves (*Bos taurus*) HPA axis reactivity to ACTH is not correlated with their

response to novelty (Van Reenen et al. 2005).

The unidimensional model posits that both HPA axis activity and reactivity should be

lower in proactive animals (Koolhaas et al. 1999). However, it should be noted that

measurements of fecal glucocorticoid metabolites in red squirrels may not allow for direct

measurement of the reactivity of the HPA axis, which may correlate more strongly with

behavioral stress responses compared to basal regulation (Baugh et al. 2012). Although a study

on free-ranging eastern chipmunks (*Tamias striatus*) showed evidence supporting covariance of

behavioral response and physiological stress response from fecal samples, the study used only

one metric of physiological stress (coefficient of variation of fecal glucocorticoid metabolites)

per individual (Montiglio et al. 2012). This statistical method was limiting in that it did not

consider the uncertainty around each individual's measure of HPA axis activity. Our research

expands upon the chipmunk study by using more conservative statistical methods, which were

not widely used until recently (Houslay and Wilson 2017), in addition to linear models and

multiple behavioral assays to establish coping styles. Using both of these approaches, we showed

that the behavioral coping style (comprised of three correlated behaviors) does not covary with

one measure of physiological stress. We acknowledge that the studies included in Table 1 are

across multiple taxa, behaviors, and HPA axis activity measurements. This likely contributes to

the equivocal nature of support for these models in wild animals. Though we focused only on

wild animals in our brief review (Table 1), empirical studies using laboratory animals also

include variation in measurements. Due to the large variability across studies in measurements of

HPA axis activity and behavior, perhaps a less generalized model of the relationship between

behavior and physiological stress may be more predictive for future studies than our current

models.

Our study was conducted using adult red squirrels. In this population, high juvenile

mortality results in a high opportunity for selection during the first year of life (McAdam et al.

2007). Due to these strong selective pressures, we must consider the possibility that selection

may have already eroded the (co)variance of physiological and behavioral stress responses in

surviving adults. For example, perhaps juveniles with high covariance of the physiological and

behavioral stress response were unable to adaptively respond to environmental conditions,

whereas juveniles with low covariance were able to adaptively respond to conditions with the

two stress responses decoupled. Additionally, these selective pressures fluctuate across years

because red squirrels rely on a masting food source (white spruce) that goes through episodes of

booms and busts in production of reproductive cones (McAdam and Boutin 2003; Fletcher et al.

2010). Following the masting of spruce trees, squirrel populations increase in density, which may

generate density-dependent selection on juvenile traits (Dantzer et al. 2013; Fisher et al. 2017).

These fluctuations in selection may maintain genetic variation in behavioral traits in this

population (Taylor et al. 2014), and, if a pleiotropic hormone was the mechanism underlying

these behavioral correlations, it could limit an adaptive behavioral response to this fluctuating

selection if the selective forces on hormone levels and the behavior push in opposite directions

(Ketterson and Nolan Jr. 1999; McGlothlin and Ketterson 2008). For example, if a high activity

is beneficial in one environmental condition but high HPA axis reactivity is not, a strong

correlation between the two traits would constrain an individual's ability to show an adaptive

behavioral response to the current environmental conditions. Recent work, however, suggests a

hormonal pleiotropic relationship is likely not powerful enough to constrain independent

evolution of two traits (Dantzer and Swanson 2017). Alternatively, if the hormone does not show

a pleiotropic relationship with behavior and selection for both is working in the same direction,

then the absence of a correlation could slow their adaptive response relative to a situation with a

positive pleiotropic relationship between the two traits (Ketterson et al. 2009).

We also must consider the possibility that different behavioral traits are favored at

different life stages. Additional work in this study system has shown wider variation in these

behavioral traits among juvenile squirrels, with individuals at both extremes of the proactive-

reactive continuum, and individuals regress to the mean as they age (Kelley et al. 2015). This is a

potential limitation of our study. Studies using selection lines, and therefore individuals at

extremes in behavioral response, may therefore be more appropriate for making predictions

about juvenile red squirrels, though perhaps not appropriate for predictions about adults. In both

Great Tits and rainbow trout (*Oncorhynchus mykiss*), studies using exploration selection lines

have found evidence to support a correlation between behavioral and physiological stress

responses (Øverli et al. 2007; Baugh et al. 2012). This suggests the potential to detect a

relationship between behavioral and physiological stress responses in juvenile red squirrels,

when individuals are more widely dispersed along the proactive-reactive continuum. Future work

on the relationship between coping styles and physiological stress responses should investigate

the ontogeny of the relationship, and how it may change across life stages.

Our study helps establish a foundation to use in exploring the fitness consequences of

variability across two axes of the stress response, behavioral and physiological. Building upon

this current work, we have an opportunity to explore the mechanisms contributing to each axis of

variation independently. For instance, the maternal environment during ontogeny may influence

the development of the physiological stress axis, or contribute to the behavioral coping style

(reviewed in Meaney 2001). Furthermore, our study provides additional evidence supporting the

lack of direct phenotypic correlation between behavioral and physiological stress responses in

wild animals exhibiting natural variation in stress responses. Our study, in conjunction with

previous studies on these models in wild animals (Table 1), suggests a need for a more

generalizable model of the relationship between the behavioral and physiological stress

responses, perhaps taking into account the environmental conditions experienced by the species.

Further studies in wild animals are needed to explore the mechanisms underlying this variation

along the phenotypic landscape of the stress response and the adaptive value of such variation.

Studies on the relationship between the behavioral and physiological stress phenotypes in

wild animals in variable environments provide insight into the pleiotropic constraints on the

evolutionary paths these populations may take. Our study contributes to a growing body of work

in support of the two-tier model of coping styles and physiological stress reactivity and activity

in wild and laboratory populations. Specifically, our study demonstrated that the FCM

concentration of wild red squirrels is independent of an individual's activity, aggression, and

docility. Given that red squirrels in this region experience a fluctuating environment in terms of

competitors (Dantzer et al. 2013), food (Boutin et al. 2006), and predators (O'Donoghue et al.

1998; Studd et al. 2014) and also fluctuating selection on behavioral traits (Boon et al., 2007;

Taylor et al., 2014), having behavioral and physiological responses that are uncorrelated may be

beneficial for adapting to this environmental variability. If similar results are found in other

species, the lack of a phenotypic relationship between the behavioral and physiological stress

responses could have important evolutionary implications, particularly for those species living in

fluctuating environments.

Ethical approval: All applicable international, national, and/or institutional guidelines for the

care and use of animals were followed. All procedures performed in studies involving animals

were in accordance with the ethical standards of the institution or practice at which the studies

were conducted.

Conflict of interest: The authors declare that they have no conflict of interest.

References

Archer J (1973) Tests for emotionality in rats and mice: A review. Anim Behav 21:205–235. doi:

10.1016/S0003-3472(73)80065-X

Baugh AT, Schaper S V., Hau M, et al (2012) Corticosterone responses differ between lines of

great tits (*Parus major*) selected for divergent personalities. Gen Comp Endocrinol

175:488–494. doi: 10.1016/j.ygcen.2011.12.012

Blumstein DT, Daniel JC (2007) Quantifying behavior the JWatcher way. Sinaeur Associates,

Inc., Sunderland, MA

Boon AK, Reale D, Boutin S (2008) Personality, habitat use, and their consequences for survival

in North American red squirrels *Tamiasciurus hudsonicus*. Oikos 117:1321–1328. doi:

10.1111/j.2008.0030-1299.16567

Boon AK, Réale D, Boutin S (2007) The interaction between personality, offspring fitness and

food abundance in North American red squirrels. Ecol Lett 10:1094–1104. doi:

10.1111/j.1461-0248.2007.01106.x

Boutin S, Larsen KW (1993) Does food availability affect growth and survival of males and

females differently in a promiscuous small mammal, *Tamiasciurus hudsonicus*? J Anim

Ecol 62:364–370. doi: 10.2307/5367

Boutin S, Wauters LA, McAdam AG, et al (2006) Anticipatory reproduction and population

growth in seed predators. Science 314:1928–1930. doi: 10.1126/science.1135520

Carere C, Caramaschi D, Fawcett TW (2010) Covariation between personalities and individual

differences in coping with stress: Converging evidence and hypotheses. Curr Zool 56:728–

741

Carere C, Oers K Van (2004) Shy and bold great tits (*Parus major*): body temperature and breath

rate in response to handling stress. Physiol Behav 82:905–912. doi:

10.1016/j.physbeh.2004.07.009

Cockrem JF (2007) Stress, corticosterone responses and avian personalities. J Ornithol 148:169–

178. doi: 10.1007/s10336-007-0175-8

Cowles MK, Carlin BP (1996) Markov chain Monte Carlo convergence diagnostics: A

comparative review. J Am Stat Assoc 91:883–904. doi: 10.1080/01621459.1996.10476956

Dantzer B, Boutin S, Humphries MM, McAdam AG (2012) Behavioral responses of territorial

red squirrels to natural and experimental variation in population density. Behav Ecol

Sociobiol 66:865–878. doi: 10.1007/s00265-012-1335-2

Dantzer B, McAdam AG, Palme R, et al (2010) Fecal cortisol metabolite levels in free-ranging

North American red squirrels: Assay validation and the effects of reproductive condition.

Gen Comp Endocrinol 167:279–286. doi: 10.1016/j.ygcen.2010.03.024

Dantzer B, Newman AEM, Boonstra R, et al (2013) Density triggers maternal hormones that

increase adaptive offspring growth in a wild mammal. Science 340:1215–1217. doi:

10.1126/science.1235765

Dantzer B, Swanson EM (2017) Does hormonal pleiotropy shape the evolution of performance

and life history traits? Integr Comp Biol 57:372–384. doi: 10.1093/icb/icx064

Dosmann AJ, Brooks KC, Mateo JM (2015) Within-individual correlations reveal link between a

behavioral syndrome, condition, and cortisol in free-ranging Belding's ground squirrels.

Ethology 121:125–134. doi: 10.1111/eth.12320

Dray S, Dufour A-B (2007) The ade4 package: implementing the duality diagram for ecologists.

J Stat Softw 22:. doi: 10.18637/jss.v022.i04

Ferrari C, Pasquaretta C, Carere C, et al (2013) Testing for the presence of coping styles in a

wild mammal. Anim Behav 85:1385–1396. doi: 10.1016/j.anbehav.2013.03.030

Fisher DN, Boutin S, Dantzer B, et al. (2017) Multilevel and sex-specific selection on

546 competitive traits in North American red squirrels. Evolution 71:1841–1854. doi:

547 10.1111/evo.13270

548 Fletcher QE, Boutin S, Lane JE, et al (2010) The functional response of a hoarding seed predator

549 to mast seeding. Ecology 91:2673–2683. doi: 10.1890/09-1816.1

550 Garamszegi LZ, Rosivall B, Rettenbacher S, et al (2012) Corticosterone, avoidance of novelty,

551 risk-taking and aggression in a wild bird: No evidence for pleiotropic effects. Ethology

552 118:621–635. doi: 10.1111/j.1439-0310.2012.02049.x

553 Gelman A, Rubin DB (1992) Inference from iterative simulation using multiple sequences. Stat

554 Sci 7:457–472. doi: 10.1214/ss/1177011136

555 Geweke J (1992) Evaluating the accuracy of sampling-based approaches to the calculation of

556 posterior moments. In: Bernardo J, Berger A, Dawid A, Smith A (eds) Bayesian Statistics 4.

557 Oxford University Press, Oxford, pp 169–193

558 Gosling SD (2001) From mice to men: what can we learn about personality from animal

559 research? Psychol Bull 127:45–86. doi: 10.1037/0033-2909.127.1.45

560 Hadfield JD (2010) MCMC methods for multi-response generalized linear mixed models: The

561 MCMCglmm R package. J Stat Softw 33:. doi: 10.18637/jss.v033.i02

562 Hänninen L, Pastell M (2009) CowLog: Open-source software for coding behaviors from digital

563 video. Behav Res Methods 41:472–476. doi: 10.3758/BRM.41.2.472

564 Houslay TM, Wilson AJ (2017) Avoiding the misuse of BLUP in behavioural ecology. Behav

565 Ecol 28:948–952. doi: 10.1093/beheco/arx023

566 Kelley AD, Humphries MM, McAdam AG, Boutin S (2015) Changes in wild red squirrel

567 personality across ontogeny: activity and aggression regress towards the mean. Behaviour

568 152:1291–1306. doi: 10.1163/1568539X-00003279

569 Ketterson ED, Atwell JW, McGlothlin JW (2009) Phenotypic integration and independence:

Hormones, performance, and response to environmental change. Integr Comp Biol 49:365–379. doi: 10.1093/icb/icp057

Ketterson ED, Nolan, Jr. V (1999) Adaptation, exaptation, and constraint: A hormonal perspective. Am Nat 154:S4–S25. doi: 10.1086/303280

Koolhaas JM, De Boer SF, Buwalda B, Van Reenen K (2007) Individual variation in coping with stress: A multidimensional approach of ultimate and proximate mechanisms. Brain Behav Evol 70:218–226. doi: 10.1159/000105485

Koolhaas JM, de Boer SF, Coppens CM, Buwalda B (2010) Neuroendocrinology of coping styles: Towards understanding the biology of individual variation. Front Neuroendocrinol 31:307–321. doi: 10.1016/j.yfrne.2010.04.001

Koolhaas JM, Korte SM, De Boer SF, et al (1999) Coping styles in animals: Current status in behavior and stress-physiology. Neurosci Biobehav Rev 23:925–935. doi: 10.1016/S0149-7634(99)00026-3

Lendvai ÁZ, Bókony V, Chastel O (2011) Coping with novelty and stress in free-living house sparrows. J Exp Biol 214:821–828. doi: 10.1242/jeb.047712

Love OP, Madliger CL, Bourgeon S, et al. (2014) Evidence for baseline glucocorticoids as mediators of reproductive investment in a wild bird. Gen Comp Endocrinol 199:65–69. doi: 10.1016/j.ygcen.2014.01.001

Martin JGA, Réale D (2008) Temperament, risk assessment and habituation to novelty in eastern chipmunks, *Tamias striatus*. Anim Behav 75:309–318. doi: 10.1016/j.anbehav.2007.05.026

McAdam AG, Boutin S (2003) Variation in viability selection among cohorts of juvenile red squirrels (*Tamiasciurus hudsonicus*). Evolution 57:1689–1697. doi: 10.1111/j.0014-3820.2003.tb00374.x

McAdam AG, Boutin S, Sykes AK, Humphries MM (2007) Life histories of female red squirrels

and their contributions to population growth and lifetime fitness. Ecoscience 14:362. doi:

10.2980/1195-6860(2007)14[362:LHOFRS]2.0.CO;2

McGlothlin JW, Ketterson ED (2008) Hormone-mediated suites as adaptations and evolutionary

constraints. Philos Trans R Soc B Biol Sci 363:1611–1620. doi: 10.1098/rstb.2007.0002

Meaney MJ (2001) Maternal care, gene expression, and the transmission of individual

differences in stress reactivity across generations. Annu Rev Neurosci 24:1161–1192

Montiglio P-O, Garant D, Pelletier F, Réale D (2012) Personality differences are related to long-

term stress reactivity in a population of wild eastern chipmunks, *Tamias striatus*. Anim

Behav 84:1071–1079. doi: 10.1016/j.anbehav.2012.08.010

O'Donoghue M, Boutin S, Krebs CJ, et al (1998) Behavioural responses of coyotes and lynx to

the snowshoe hare cycle. Oikos 82:169. doi: 10.2307/3546927

Øverli Ø, Sørensen C, Pulman KGT, et al (2007) Evolutionary background for stress-coping

styles: Relationships between physiological, behavioral, and cognitive traits in non-

mammalian vertebrates. Neurosci Biobehav Rev 31:396–412. doi:

10.1016/j.neubiorev.2006.10.006

Palme R (2019) Non-invasive measurement of glucocorticoids: Advances and problems. Physiol

Behav 199:229–243. doi: 10.1016/j.physbeh.2018.11.021

Palme R, Touma C, Arias N, et al (2013) Steroid extraction: Get the best out of faecal samples.

Wiener Tierärztliche Monatsschrft 100:238–246

Qu J, Fletcher QE, Réale D, et al (2018) Independence between coping style and stress reactivity

in plateau pika. Physiol Behav 197:1–8. doi: 10.1016/j.physbeh.2018.09.007

Raulo A, Dantzer B (2018) Associations between glucocorticoids and sociality across a

continuum of vertebrate social behavior. Ecol Evol 1–20. doi: 10.1002/ece3.4059

Réale D, Reader SM, Sol D, et al (2007) Integrating animal temperament within ecology and

evolution. Biol Rev 82:291–318. doi: 10.1111/j.1469-185X.2007.00010.x

Romero LM, Dickens MJ, Cyr NE (2009) The reactive scope model - A new model integrating

homeostasis, allostasis, and stress. Horm Behav 55:375–389. doi:

10.1016/j.yhbeh.2008.12.009

Sapolsky RM, Romero LM, Munck AU (2000) How do glucocorticoids influence stress

responses? Integrating permissive, suppressive, stimulatory, and preparative actions. Endocr

Rev 21:55–89. doi: 10.1210/er.21.1.55

Sheriff MJ, Dantzer B, Delehanty B, et al (2011) Measuring stress in wildlife: Techniques for

quantifying glucocorticoids. Oecologia 166:869–887. doi: 10.1007/s00442-011-1943-y

Sih A, Bell AM, Johnson JC, Ziemba RE (2004) Behavioral syndromes: An integrative

overview. Q Rev Biol 79:241–277. doi: 10.1086/422893

Siracusa E, Morandini M, Boutin S, et al (2017) Red squirrel territorial vocalizations deter

intrusions by conspecific rivals. Behaviour 154:1259–1273. doi: 10.1163/1568539X-

00003467

Stamps J, Groothuis TGG (2010) The development of animal personality: Relevance, concepts

and perspectives. Biol Rev 85:301–325. doi: 10.1111/j.1469-185X.2009.00103.x

Steimer T, Driscoll P (2003) Divergent stress responses and coping styles in psychogenetically

selected Roman high-(RHA) and low-(RLA) avoidance rats: Behavioural, neuroendocrine

and developmental aspects. Stress 6:87–100. doi: 10.1080/1025389031000111320

Stöwe M, Rosivall B, Drent PJ, Möstl E (2010) Selection for fast and slow exploration affects

baseline and stress-induced corticosterone excretion in Great tit nestlings, Parus major.

Horm Behav 58:864–871. doi: 10.1016/j.yhbeh.2010.08.011

Studd EK, Boutin S, McAdam AG, et al (2014) Predators, energetics and fitness drive neonatal

reproductive failure in red squirrels. J Anim Ecol 84:249–259. doi: 10.1111/1365-

2656.12279

Taylor RW (2012) Quantitative genetics, selection, mate choice and red squirrel behavior in a

fluctuating environment. Michigan State University

Taylor RW, Boon AK, Dantzer B, et al. (2012) Low heritabilities, but genetic and maternal

correlations between red squirrel behaviours. J Evol Biol 25:614–624. doi: 10.1111/j.1420-

9101.2012.02456.x

Taylor RW, Boutin S, Humphries MM, McAdam AG (2014) Selection on female behaviour

fluctuates with offspring environment. J Evol Biol 27:2308–2321. doi: 10.1111/jeb.12495

Touma C, Sachser N, Möstl E, Palme R (2003) Effects of sex and time of day on metabolism and

excretion of corticosterone in urine and feces of mice. Gen Comp Endocrinol 130:267–278.

doi: 10.1016/S0016-6480(02)00620-2

Van Reenen CG, O'Connell NE, Van der Werf JTN, et al (2005) Responses of calves to acute

stress: Individual consistency and relations between behavioral and physiological measures.

Physiol Behav 85:557–570. doi: 10.1016/j.physbeh.2005.06.015

658 **Figure 1.** Stress reactivity and activity, as measured by average FCM concentration, is not

659 predicted by docility in North American red squirrels. Size of the points represents number of

660 fecal samples included for that individual.

661 **Figure 2.** Stress reactivity and activity, as measured by average FCM concentration, is not

662 predicted by activity or aggression in North American red squirrels. Activity and aggression are

663 from scores determined by the principal component analyses. Size of the points represents

664 number of fecal samples included for that individual.

7   **Table 1. Review of field studies testing for the covariance in behavioral coping styles and HPA activity**

8   This non-comprehensive table includes studies conducted with natural populations testing the unidimensional (Koolhaas et al. 1999b)

9   and two-tier models (Koolhaas et al. 2010) of the among individual relationship between behavioral coping styles ('Behavioral Trait') and

HPA axis activity ('Physiological Measurement'). The first section includes studies that support the main prediction from the unidimensional

model that more proactive individuals have lower HPA axis activity. The second section includes studies that are contrary to the main

prediction from the unidimensional model that more proactive individuals would have lower HPA axis activity. These studies do show that

behavior and HPA activity covary, but the relationship is in the opposite of the direction predicted by the unidimensional model with

proactive individuals having lower HPA axis activity. The third section includes studies that support the two-tier model that predicts that

behavior and HPA axis activity do not covary in either direction. Correlations or estimates are included if available in the corresponding

manuscript. Confidence/credible intervals are included if available; if not, p-values are included when available. If the confidence/credible

interval overlapped zero, we interpreted this as the behavior trait measured and the physiological measurement did not covary. Non-true

baseline samples were the first sample collected but involved some handling or trapping stress. (n = sample size of individuals, NS = non-

significant ($p > 0.05$), FCM = fecal cortisol metabolites, DEX = dexamethasone (synthetic corticosteroid that exerts negative feedback on the

0   HPA axis), ACTH = adrenocorticotropic hormone that increases adrenocortical activity)

Species	Behavioral Trait	Physiological Measurement	n	Correlation or Estimate [CI]
Evidence supporting the predictions of the unidimensional model that behavioral and physiological traits are negatively correlated				
Great Tits (*Parus major*) (Baugh et al. 2017)	Exploration	Blood corticosterone *(ACTH challenge induced)*	85	$R^2 = 0.051$, $p = 0.04$
Great Tits (*Parus major*) (Baugh, van Oers, Naguib, & Hau, 2013)	Exploration Exploration	Blood corticosterone *(90-min handling-restraint stress-induced)* Blood corticosterone *(area under the stress-induced curve)*	16 16	$b = -0.417$, $p = 0.015$ $b = -24.66$, $p = 0.007$
Belding's ground squirrels (*Urocitellus belingi*) (Clary et al. 2014)	Vigilance	FCM *(nominal baseline)*	12	$b = 2.109$ [0.17, 4.05]
Brook charr (*Salvelinus fontinalis*) (Farwell et al. 2014)	Activity	Whole-body cortisol *(baseline and handling/novel object stress-induced samples)*	66	NA*
House Sparrows (*Passer domesticus*) (Lendvai et al. 2011)	Hovering	Blood corticosterone *(30 min handling-restraint stress-induced)*	18	Pearson's r = -0.58, $p = 0.017$
Eastern chipmunks (*Tamias striatus*) (Montiglio et al. 2012)	Exploration	FCM *(coefficient of variation)*	58	$b = -13.68$ [-27.62, -4.45]
Plateau pika (*Ochotona curzoniae*) (Qu et al. 2018)	Shyness	Blood cortisol *(true baseline)*	292	posterior R = 0.45 [0.09, 0.66]
Evidence in the opposite direction of predictions of the unidimensional model				
Poeciliid fish (*Brachyrhaphis episcopi*) (Archard et al. 2012)	Exploration, Activity	Water-borne cortisol *(open-field stress-induced)*	96	Pearson's r = -0.29, $p = 0.005$
Great Tits (*Parus major*) (Baugh et al. 2013)	Exploration	Blood corticosterone *(true baseline)*	82	$b = 0.536$, $p = 0.003$
Alpine marmots (*Marmota marmota*) (Costantini et al. 2012)	Exploration, Activity	Blood cortisol *(non-true baseline)*	28	$b = 0.54$, Fischer *C*-statistic model selection
Graylag Geese (*Anser anser*) (Kralj-Fišer et al. 2009)	Aggression	FCM *(handling stress-induced)*	10	Spearman's r = 0.782, $p = 0.008$
Evidence supporting the predictions of the two-tier model that behavioral and physiological traits are not significantly correlated				

Species	Behavioral Trait	Physiological Measurement	n	Correlation or Estimate [CI]
Great Tits (*Parus major*)	Exploration	Blood corticosterone *(handling-restrain stress-induced)*	85	$R^2 = 0.03$, $p = 0.14$
(Baugh, Davidson, Hau, & van Oers, 2017)	Exploration	Blood corticosterone *(DEX challenge response)*	85	$R^2 = 0.02$, $p = 0.20$
Belding's ground squirrels (*Urocitellus belingi*)	Exploration	FCM *(non-true baseline)*	12	b = -0.76 [-1.12, 0.98]
(Clary et al., 2014)				
Alpine marmots (*Marmota marmota*)	Exploration, Activity	Blood cortisol *(pre-restraint to post-restraint response)*	28	NA**
(Costantini et al., 2012)				
Belding's ground squirrels (*Urocitellus belingi*)	Activity	FCM *(non-true baseline)*	157	posterior R = -0.007, [-0.123, 0.095]
(Dosmann et al. 2015)	Exploration	FCM *(non-true baseline)*	157	posterior R = -0.046, [-0.175, 0.036]
	Docility	FCM *(non-true baseline)*	157	posterior R = 0.016, [-0.077, 0.139]
Alpine marmots (*Marmota marmota*)	Activity	Blood cortisol *(non-true baseline)*	146	posterior R = 0.04, [-0.56, 0.71]
(Ferrari et al. 2013)	Impulsivity	Blood cortisol *(non-true baseline)*	146	posterior R = 0.08, [-0.68, 0.62]
	Docility	Blood cortisol *(non-true baseline)*	146	posterior R = 0.14, [-0.64, 0.63]
Collard Flycatchers (*Ficedula albicollis*)	Novel object avoidance	FCM *(non-true baseline)*	51	Pearson's r = -0.017, [-0.291, 0.260]
(Garamszegi et al. 2012)	Aggression	FCM *(non-true baseline)*	56	Pearson's r = -0.076, [-0.332, 0.191]
	Risk-taking	FCM *(non-true baseline)*	54	Pearson's r = 0.074, [-0.198, 0.335]
Nazca Boobies (*Sula granti*)	Gardening (non-social)	Blood corticosterone *(true baseline)*	222	b = 0.26, [-0.09, 0.61]
(Grace and Anderson 2014)	Shaking (non-social)	Blood corticosterone *(true baseline)*	222	b = -0.06, [-0.25, 0.13]
	Aggression (non-social)	Blood corticosterone *(true baseline)*	222	b = 0.19, [-0.07, 0.45]
	Gardening (non-social)	Blood corticosterone *(maximum value across 4 time points during capture-restraint)*	222	b = -0.30, [-0.98, 0.38]
	Shaking (non-social)	Blood corticosterone *(maximum value across 4 time points during capture-restraint)*	222	b = -0.08, [-0.44, 0.28]
	Aggression (non-social)	Blood corticosterone *(maximum value across 4 time points during capture-restraint)*	222	b = -0.07, [-0.35, 0.21]
	Gardening (non-social)	Blood corticosterone *(area under the curve across 4 time points during capture-restraint)*	222	b = -0.16, [-0.7, 0.38]
	Shaking (non-social)	Blood corticosterone *(area under the curve across 4 time points during capture-restraint)*	222	NA, CI in figure includes 0
	Aggression (non-social)	Blood corticosterone *(area under the curve across 4 time points during capture-restraint)*	222	NA, CI in figure includes 0
	Gardening (social)	Blood corticosterone *(true baseline)*	222	b = 0.20, [0.02, 0.38]
	Shaking (social)	Blood corticosterone *(true baseline)*	222	b = -0.04 [-0.18, 0.1]

Species	Behavioral Trait	Physiological Measurement	n	Correlation or Estimate [CI]
	Aggression (social)	Blood corticosterone *(true baseline)*	222	b = -0.02 [-0.11, 0.07]
	Gardening (social)	Blood corticosterone *(maximum value across 4 time points during capture-restraint)*	222	b = -0.03 [-0.23, 0.17]
	Shaking (social)	Blood corticosterone *(maximum value across 4 time points during capture-restraint)*	222	b = 0.04 [-0.2, 0.28]
	Aggression (social)	Blood corticosterone *(maximum value across 4 time points during capture-restraint)*	222	b = -0.01 [-0.02, 0]
	Gardening (social)	Blood corticosterone *(area under the curve across 4 time points during capture-restraint)*	222	b = 0.02 [-0.15, 0.19]
	Shaking (social)	Blood corticosterone *(area under the curve across 4 time points during capture-restraint)*	222	NA, CI in figure includes 0
	Aggression (social)	Blood corticosterone *(area under the curve across 4 time points during capture-restraint)*	222	NA, CI in figure includes 0
Graylag Geese (*Anser anser*) (Kralj-Fišer et al. 2009)	Aggression	FCM *(non-true baseline)*	10	Spearman's r = -0.503, NS
	Sociability	FCM *(non-true baseline)*	10	Spearman's r = -0.212, NS
	Sociability	FCM *(handling stress-induced)*	10	Spearman's r = -0.127, NS
House Sparrows (*Passer domesticus*) (Lendvai et al. 2011)	Exploration	Blood corticosterone *(30 min handling-restraint stress-induced)*	18	Pearson's r = -0.07, *p* = 0.797
	Shy-bold	Blood corticosterone *(30 min handling-restraint stress-induced)*	18	Pearson's r = -0.04, *p* = 0.866
	Exploration	Blood corticosterone *(true baseline)*	18	Pearson's r = -0.18, *p* = 0.487
	Shy-bold	Blood corticosterone *(true baseline)*	18	Pearson's r = -0.07, *p* = 0.801
	Hovering	Blood corticosterone *(true baseline)*	18	Pearson's r = -0.14, *p* = 0.573
Plateau pika (*Ochotona curzoniae*) (Qu et al. 2018)	Docility, in cage	Blood cortisol *(true baseline)*	292	posterior R = 0.25 [-0.02, 0.63]
	Docility, in bag	Blood cortisol *(true baseline)*	292	posterior R = -0.06 [-0.28, 0.31]
	Exploration	Blood cortisol *(true baseline)*	292	posterior R = -0.16 [-0.52, 0.05]
	Docility, in cage	Blood cortisol *(40 min post-capture change from baseline)*	292	posterior R = 0.22 [-0.06, 0.47]
	Docility, in bag	Blood cortisol *(40 min post-capture change from baseline)*	292	posterior R = 0.16 [-0.07, 0.48]
	Exploration	Blood cortisol *(40 min post-capture change from baseline)*	292	posterior R = 0.10 [-0.42, 0.24]
	Shyness	Blood cortisol *(40 min post-capture change from baseline)*	292	posterior R = -0.23 [-0.28, 0.22]

* Used AICc model selection approach. All top models included activity as predictor of cortisol.

Based on figure in text, the top model indicates more active individuals have less whole-body

cortisol.

** Used nested path model selection approach. Based on figure in text, the top model does not

include a relationship between change in cortisol and coping style.

Table 2. Bivariate model structures and sample sizes. This table breaks down the sample sizes and the variables (fixed and random

effects) considered in each bivariate mixed effects model. Response variables shown are for the bivariate models that tested the

association between behavioral traits (docility, activity, or aggression) and a measure of the physiological stress response (fecal

cortisol metabolites or FCM). Italics indicate fixed effects estimated for FCM concentration only.

Model	Response Variables	Fixed Effects	Random Effect	Year	# of individuals	# of fecal samples
1	docility, FCM	study grid (control or high-density) sex docility trial number breeding, non-breeding, or lactating *fecal processing lab* *Julian date of trapping event*	squirrel ID	2008:	32	36
				2009:	56	132
				total:	**79***	**168**
2	activity, FCM	study grid (control or high-density) sex breeding, non-breeding, or lactating *fecal processing lab* *Julian date of trapping event*	squirrel ID	2005:	2	9
				2008:	27	139
				2009:	15	117
				2010:	1	10
				2012:	28	260
				total:	**72***	**484**
3	aggression, FCM	study grid (control or high-density) sex breeding, non-breeding, or lactating *fecal processing lab* *Julian date of trapping event*	squirrel ID	2005:	2	9
				2008:	27	139
				2009:	15	117
				2010:	1	10
				2012:	28	260
				total:	**72***	**484**

695 *Some squirrels were sampled in multiple years.

Table 3. Multivariate results for relationships between FCM and behaviors

Results from our three bivariate generalized linear mixed-effects model models to examine the relationships between FCM and each of three behaviors individually (activity and aggression models: n = 484 fecal samples; docility model: n = 168 fecal samples). Among-individual variances are listed on the diagonal, covariances below and correlations above (with the lower and upper bounds of 95% CIs in parentheses).

	ln(FCM)	Docility
ln(FCM)	0.049	0.14
	[1.86-8, 0.l3]	[-0.63, 0.83]
Docility	0.012	0.40
	[-0.074, 0.11]	[0.15, 0.71]

	ln(FCM)	Activity
ln(FCM)	0.075	0.15
	[0.018, 0.14]	[-0.17, 0.47]
Activity	0.04	0.55
	[-0.034, 0.11]	[0.38, 0.73]

	ln(FCM)	Aggression
ln(FCM)	0.077	0.29
	[0.016, 0.14]	[-0.098, 0.56]
Aggression	0.061	1.00
	[-0.03, 0.16]	[0.68, 1.33]

704 **Figure 1.**

709   **Figure 2.**