Dynamics of age-related catastrophic mitotic failures and recovery in yeast

Authors: Matthew M. Crane[1]*, Adam E. Russell[1+], Brent J. Schafer[1+], Mung Gi Hong[1], Joslyn E. Goings[1], Kenneth L. Chen[1,2,3], Ben W. Blue[1] and Matt Kaeberlein[1]*

[1]Department of Pathology, University of Washington, Seattle, WA, USA.
[2]Department of Genome Sciences, University of Washington, Seattle, Washington, USA
[3]Medical Scientist Training Program, University of Washington, Seattle, Washington USA

[+]Authors contributed equally
*Correspondence to: mcrane2@uw.edu ; kaeber@uw.edu

Abstract: Genome instability is a hallmark of aging and contributes to age-related disorders such as progeria, cancer, and Alzheimer's disease. In particular, nuclear quality control mechanisms and cell cycle checkpoints have generally been studied in young cells and animals where they function optimally, and where genomic instability is low. Here, we use single cell imaging to study the consequences of increased genomic instability during aging, and identify striking age-associated genome missegregation events. During these events the majority of mother cell chromatin, and often both spindle poles, are mistakenly sent to the daughter cell. This breakdown in mitotic fidelity is accompanied by a transient cell cycle arrest that can persist for many hours, as cells engage a retrograde transport mechanism to return chromosomes to the mother cell. The repetitive ribosomal DNA (rDNA) has been previously identified as being highly vulnerable to age-related replication stress and genomic instability, and we present several lines of evidence supporting a model whereby expansion of rDNA during aging results in nucleolar breakdown and competition for limited nucleosomes, thereby increasing risk of catastrophic genome missegregation.

Main Text: Each cell cycle involves a delicate choreography of duplicating genetic material and cellular organelles, and apportioning them appropriately between mother and daughter cell. Failures of cell cycle regulation can result in severely compromised fitness or cells that respond improperly to environmental cues and emerge as cancerous precursors (Hanahan and Weinberg, 2011). In particular, aneuploidy (the gain or loss of partial or whole chromosomes) can be deleterious to fitness(Beach et al., 2017; Sunshine et al., 2016) and has been implicated in many different types of cancers(Gordon et al., 2012) as well as developmental diseases such as Down Syndrome(Nagaoka et al., 2012). Recent work has also documented extensive damage and genomic rearrangements that can occur from micronuclei or from telomeric crisis(Maciejowski et al., 2015; Zhang et al., 2015), and identified the rDNA sequences as particularly vulnerable to genomic damage(Flach et al., 2014; Xu et al., 2017). Mitotic processes and checkpoints of the budding yeast *Saccharomyces cerevisiae* have been extensively studied, but the vast majority of work has focused on logarithmically growing young cells(Beach et al., 2017; Dotiwala et al., 2007; London and Biggins, 2014; Palmer et al., 1989; Santaguida and Amon, 2015). By studying nuclear dynamics in yeast as they age, we have uncovered a cause of age-associated genomic instability and an active mechanism to maintain nuclear integrity and proper segregation of genetic material in aged cells.

We characterized the dynamics of genome replication and partitioning during replicative aging by imaging cells expressing fluorescently tagged histone 2B (Htb2:mCherry) in a microfluidic device over their entire lifespans(Crane et al., 2014). During each cell cycle, the amount of Htb2 in the mother cell nucleus increases by about two-fold and then drops by one-half, as the cell enters mitosis and chromosomes are segregated to the newly formed daughter. The vast majority of cell divisions in young cells follow this characteristic pattern (Figure 1A). As cells age, however, abnormal segregation events become common (Figure 1B, Videos 1-2, please ensure volume is on for all Videos to hear audio explanation). The single cell trace shown in Figure1C, for example, shows a cell undergoing multiple cell cycles with proper division until an abnormal segregation occurs in which the majority of detectable histones are sent to the daughter cell. These genome-level missegregation (GLM) events result in cell cycle arrest that can range from a few minutes (Figure 1C-top) to many hours (Figure1C-middle), before they are usually corrected by returning the aberrantly segregated genetic material to the mother cell. The range of arrest durations is broad, with most events resolved within an hour, but some lasting many hours (Figure 1 – figure supplement 1). If corrected by this REtrograde TRansport Nuclear (RETRN) process, mother cells are able to proceed through subsequent divisions, but if not, the mother cells will terminally exit the cell cycle and senesce (Figure 1C-bottom).

To further characterize GLM and RETRN dynamics during aging, cells were imaged over their entire replicative lifespan, with birth events, GLMs, and RETRN events assessed. The probability of a GLM increased dramatically at the end of life (Figure 1D), with approximately three quarters of mother cells experiencing one or more GLMs (Figure 1E). About 90% of GLMs were corrected through successful RETRN events, allowing individual mother cells to live approximately 30% longer on average than if all GLMs were terminal (Figure 1 – figure supplement 2). However, even when corrected, mother cells that undergo a GLM are more likely to die in the near future than cells of the same age that have not experienced such an event, and GLMs become increasingly predictive of impending mortality with increasing age (Figure 1 – figure supplement 2). To confirm that the histones do indeed co-localize with DNA during these events, we imaged old mothers and observed the dynamics of Htb2 in cells exposed to the DNA stain Hoechst 3342. As can be clearly seen (Figure 1F, Videos 3-4), both the DNA and histones move in concert during these events. Furthermore, we confirmed that histone levels reflect nuclear DNA abundance in single cells by staining with DAPI and comparing Htb2 levels with DNA content in old cells (Figure 1 – figure supplement 3). GLM and RETRN dynamics were not influenced by the fluorophore used or which histone is tagged, as the dynamics of both Htb2:mCherry and histone 2A tagged with GFP (Hta2:GFP) did not differ (Figure 1 – figure supplement 4). GLM and RETRN frequency is not an artifact of our imaging protocol, as modifying the excitation power or the cumulative excitation energy exposure had no effect on these observations (Figure 1 – figure supplement 5). For clarity, the strain containing Htb2:mCherry is referred to as wild-type hereafter.

In order to observe the nuclear periphery during GLM and RETRN events, we imaged aging cells expressing both Htb2:mCherry and Nup49:GFP, and compared normal divisions (Video 5) with RETRN events (Figure 1G-top, Video 6) and terminal GLMs (Figure 1G-bottom, Video 7). The dynamics of the histone missegregation and recovery can be clearly seen in these time-lapse series, and strikingly the mother cells retain an intact nuclear envelope during these events – even if they lose all the chromatin (Figure1G). Passage of the histones fully into the daughter cell is evident from cells co-expressing a bud neck marker (Myo1:GFP) along with Htb2:mCherry (Videos 1-2). Interestingly, during these events both spindle poles often enter the

daughter and move in concert with the tagged histones (Figure1H). Spindle missegregation and
RETRN was visualized through time-lapse imaging of aging cells expressing the spindle
component Spc72 tagged with GFP (Spc72:GFP, Video 8). In terminal GLMs without RETRN,
both spindle poles generally remain in the daughter cell (Figure 1H, Video 9). This can also be
observed in videos where tubulin is tagged with GFP (Tub1:GFP), and all of the detectable
nuclear microtubules enter the daughter cell during GLMs (Video 10-11).
98

Fig 1

99
Fig. 1. During replicative aging cells frequently undergo dramatic genomic missegregation events. **A)** Schematic
showing the process of a normal cell division where chromatin (red) doubles during S-phase and is divided between
mother and daughter during mitosis. **B)** Aging cells frequently experience Genome Level Missegregation (GLM)
events where most genomic material enters the daughter while the nuclear envelope stays behind. Usually this
missegregation is corrected (top, RETRN event), allowing mother cells to go on to divide and produce more

daughters. If not corrected and cytokinesis occurs (bottom), this becomes a terminal event wherein mother cells
replicatively senesce. **C)** Representative single cell traces of mother Htb2 levels showing missegregation (shaded)
and active retrograde correction events. Corrections can occur quickly (top), or can take hours to be completed
(middle). A GLM becomes terminal (bottom) if it is not corrected. (*) indicates the formation of new buds, and both
cells with RETRN events produce additional daughters. AU indicates arbitrary units. **D)** Missegregation
probabilities increase dramatically near the end of replicative lifespan. n=359 mother cells examined. **E)** Over their
entire replicative lifespan, individual mother cells have a greater than 70% chance of having one or more
missegregation events. **F)** Genomic DNA and histones co-localize during GLM events. Two cells expressing
Htb2:mCherry and stained with a live DNA dye Hoechst 3342. **G)** Time-lapse dynamics of a GLM with RETRN
correction (top, mother cell replicative age 14) and a terminal missegregation (bottom, mother cell replicative age
12) in cells co-expressing Htb2:mCherry and Nup49:GFP. During both GLMs the nuclear envelope is clearly visible
in both mother (M) and daughter (D) cells. See Videos 6 and 7. **H)** Time-lapse dynamics of a GLM with RETRN
correction (top, mother cell replicative age 13) and a terminal missegregation (bottom, mother cell replicative age
16) in cells expressing Htb2:mCherry and Spc72:GFP. Both spindle poles can be seen to enter the daughter (D)
during these events, and during the RETRN event a spindle pole returns to the mother (M). In the terminal
missegregation, the spindle pole fails to reenter the mother cell. See Videos 8 and 9. Times are indicated in
hours:mins from the start of the displayed time-lapse, not the start of the experiment. Arrows indicate mother cells
without visible chromatin.
123
To investigate the temporal and cell cycle dynamics of the genome missegregation and
RETRN events, we employed two cell cycle reporters, Whi5:GFP and Myo1:GFP(Di Talia et al.,
2007). Whi5 prevents exit from G1 when localized in the nucleus, and the transition of Whi5
from the nucleus to the cytoplasm signifies the end of G1 (Figure 2A) (Charvin et al., 2010).
Individual young cells proceed in a reliable fashion through the cell cycle, with Whi5 becoming
nuclear localized immediately after Htb2 levels fall and the cell enters telophase (FigureS6). By
aligning all annotated cell divisions without GLMs (Figure 2B), the temporal dynamics of
histone separation into the daughter cells are clear and immediately followed by the transition of
Whi5 from the cytoplasm into the nucleus. In contrast, when a GLM occurs, Htb2 levels fall
precipitously in mother cells, but the cell delays the cytoplasmic to nuclear transition of Whi5
until after the RETRN correction (Figure 2C). Because the Whi5 cytoplasm-to-nuclear transition
occurs upon activation of the mitotic exit network(Bean et al., 2006; Costanzo et al., 2004), this
delay demonstrates that the cell prolongs mitosis until the GLM is corrected.
In a complementary fashion, we confirmed that GLMs induced a delay in mitotic exit by
following Myo1:GFP localization and abundance. Myo1 is produced at the end of G1 as the cell
begins the formation of a new daughter(Bi and Park, 2012; Weiss, 2012). At the end of mitosis,
following successful partitioning of the genomic material, Myo1 is degraded during cytokinesis
(Figure 2A, Figure 2 – figure supplement 1). By manually aligning normal cell divisions, the
increase of Myo1 levels at the bud neck, and subsequent decrease during cytokinesis, can be
clearly seen (Figure 2D). When the Htb2 levels reach their lowest point, Myo1 levels are at a
maximum, but they immediately begin to fall as the cell eliminates the bud neck during
cytokinesis. In contrast, in divisions where a GLM occurs, the levels of Myo1 continue to
increase even after the Htb2 levels reach their lowest point (Figure 2E). Only following the
RETRN event does Myo1 begin to fall.
To determine whether GLMs result from improper spindle attachment and whether
RETRN events are caused by activation of the spindle assembly checkpoint, we deleted the gene
encoding the spindle assembly checkpoint component Mad3 (mammalian BubR1). This failed to
alter the age-related increase in missegregation, and older *mad3Δ* cells had the same
missegregation rates as wild type cells (Figure 2 – figure supplement 2). Taken together, these
data differentiate RETRN events and terminal GLMs from prior observations of nuclear

oscillations near the bud neck linked to alignment of the spindle poles between mother and
daughter(Palmer et al., 1989; Yang et al., 1997; Yeh et al., 2000, 1995). Similarly, nuclear
excursions where a single spindle-pole entered the daughter, but then returned to the mother has
been identified in DNA damage checkpoint mutants (Dotiwala et al., 2007); however, when cells
arrest as a result of the DNA damage response during metaphase, only the nucleolus enters the
daughter while all chromatin is retained by the mother cell (Witkin et al., 2012). A recent report
identified segregation of the nucleus and spindle poles into the daughter cell in five aging yeast
cells (of the 10 observed), which is likely to be the same phenotype detailed here(Neurohr et al.,
2018). They also identified elevated rates of chromosome I missegregation in old cells (in ten of
the forty cells observed). In prior studies, mutants that missegregate chromosomes or both
spindle poles to the daughter cell are unable to correct these events(Finley et al., 2008; Thrower
et al., 2003; Yeh et al., 2000, 1995); however, the RETRN events seen in aged mother cells
unambiguously delay mitotic exit and correct these failures by actively transferring chromatin,
microtubules, and/or spindle poles from the daughter cell to the mother cell.

Fig 2

168
Fig. 2. Characterization of missegregation and corrections during the cell cycle. **A)** Schematic showing the temporal
dynamics of the proteins used to characterize missegregation events. Whi5 exits the nucleus to initiate START and
move the cell into S phase. It re-enters the nucleus at the end of mitosis. Myo1 is produced at the end of G1 and
localizes to the bud neck until cytokinesis ends mitosis. **B)** Average Htb2 and Whi5 dynamics for cell cycles without
missegregation events (n=2,697). Whi5 begins to transition from cytoplasmic to nuclear when Htb2 levels reach the
lowest point as indicated by the arrow. **C)** Average traces from cell cycles with RETRN events. In these cases
(n=466), the Whi5 cytoplasmic to nuclear transition is delayed until the missegregation is corrected. **D)** Average

Htb2 and Myo1 dynamics for cell cycles without missegregations (n=3,146). Myo1 levels begin to fall immediately
after chromosome segregation, as shown by the arrow. **E**) In cell cycles with a RETRN event (n=459), even after
chromosomes enter the daughter (noted by the arrow), Myo1 levels continue to increase until the missegregation is
corrected, and then cytokinesis begins. N-values indicate the number of cell cycles analyzed, and all error bars are
standard error. Cells are aligned so that time t=0 indicates the beginning of mitotic exit and the M to G1 transition,
as determined by Whi5 and Myo1 dynamics.
182
Changes in nucleosome occupancy have been linked to age-related genomic
instability(Hu et al., 2014), DNA damage(Hauer et al., 2017), and tumorigenesis(Sharma et al.,
2010), and during the course of our studies we observed increasing Htb2 levels at late replicative
ages (Figure 3A, Video 12). Given that prior studies have reported both increasing and
decreasing histone levels with age(Feser et al., 2010; Janssens et al., 2015), we analyzed
expression of two histones tagged with GFP (Hta2, Htb2) in addition to Htb2:mCherry and
confirmed that this end of life increase is a general behavior of histones (Figure 3 – figure
supplement 1). Furthermore, all histones examined showed a dramatic increase in expression
between 5-10 divisions prior to death (Figure 3 – figure supplement 2).
We speculated that the elevated histones in aged cells may reflect an expansion of the
rDNA in the form of extrachromosomal rDNA circles (ERCs), which increase dramatically
during aging and have been proposed as a molecular mechanism of aging(Sinclair and Guarente,
1997) and genome instability (Ganley et al., 2009). As a result, rDNA copy number in old cells
has been shown to increase nearly twenty fold (Dang et al., 2009). To test whether histone levels
reflect ERC abundance, we determined the impact of reducing ERC formation by removing the
replication fork block protein Fob1; deletion of *FOB1* reduces rDNA recombination and ERC
abundance(Defossez et al., 1999; Kobayashi et al., 1998; Sinclair and Guarente, 1997) and
significantly extends lifespan (Figure 3 – figure supplement 3). In comparison to wild type cells,
the increase in histone levels with age is attenuated in *fob1*Δ cells (Figure 3B). Furthermore,
although wild-type and *fob1*Δ mother cells stained with DAPI after 16h of growth had
statistically indistinguishable replicative ages (Figure 3C, median RLS=13, p=0.42), wild-type
cells have significantly greater DNA levels (Figure 3D, p=0.0002). The lower DNA and histone
levels in old *fob1*Δ cells relative to age-matched wild-type is consistent with the model that
increasing histone levels are driven by ERC accumulation in aging yeast. If increasing histone
levels reflect ERC levels, then histone levels should act as a biomarker that predicts remaining
lifespan and could be associated with other age-associated failures. Indeed, within cells of the
same replicative age, we determined that histone abundance is negatively correlated with
remaining lifespan, and this correlation becomes stronger as cells age (Figure 3 – figure
supplement 3). Histone levels are less correlated with mortality in *fob1*Δ cells, supporting a
model where slowing ERC production reduces the underlying pathology linking histone
abundance to death.

Fig. 3. Age-related rDNA instability impairs Cdc14 release and anaphase entry. **A)** Time course of a single mother cell expressing Htb2:mCherry and Nup49:GFP shows the increase in histone levels during normal replicative aging, full movie in Video 12. The mother cell shown here has a replicative age of 16 at the timelapse start and 21 at the end. **B)** Old wild type cells have significantly greater histone levels compared to young wild type cells and accumulation of histones during aging is attenuated in cells lacking Fob1 (n=481) compared to wild type (wt, n=587). Significance of age and genotype (p<0.0001 in both cases) determined by repeated measures ANOVA. This links rDNA expansion in the form of ERCs with histone accumulation. **C)** Wild-type and *fob1Δ* cells were grown in the microfluidic device for 16h, and then fixed. At the time of fixation, both strains had similar replicative ages (p=0.42, two tailed t-test). **D)** Although both strains had similar replicative ages, the aged wild-type mother cells had increased DAPI staining levels relative to aged *fob1Δ* cells, indicating an increased level of DNA (p=0.0002, two tailed t-test). **E)** In young cells the nucleolus (Utp13:GFP) is adjacent to the genomic DNA (marked by Htb2). In aged cells, however, ERCs accumulate and are localized to the nucleolus which becomes colocalized with histones as the nucleolus fragments. Colocalization is particularly evident toward the end of a mother cells life (see Videos 13 and 14). **F)** The size of the nucleolus as measured by segmenting Utp13:GFP increases during replicative aging (n=93, p<0.0001, Mann-Kendall trend test).

The rDNA is localized to the nucleolus, a phase separated organelle inside the nucleus(Lindström et al., 2018). To examine the relationship between histone dynamics, ERCs, and nucleolar structure, we followed aging mother cells expressing Htb2:mCherry along with a GFP fusion to the nucleolar protein Utp13. In young cells, the nucleolus appears as a nuclear site adjacent to the majority of histones, while in old cells, a large quantity of histones is

localized within the nucleolus (Figure 3C, Videos 13-14). Furthermore, the size of the nucleolus
increases dramatically as cells age and often becomes fragmented into multiple foci (Figure 3D,
Videos 13-14), further indicating that the excess histones are localized to ERCs in old cells and
confirming earlier work linking ERCs to nucleolar fragmentation(Neurohr et al., 2018; Sinclair
et al., 1997; Sinclair and Guarente, 1997).
The protein Cdc14 is also localized to the nucleolus where it functions in the Regulator of
Nucleolar Silencing and Telophase Exit (RENT) complex to silence transcription (Clemente-
Blanco et al., 2011; Shou et al., 1999) and ensure proper segregation of chromosomes(D'Amours
et al., 2004; Rock and Amon, 2009; Sullivan et al., 2004). Recently Cdc14 was identified as the
limiting step in anaphase, and separately it was observed that compaction of rDNA within the
nucleolus interfered with proper release of Cdc14 from the nucleolus(de los Santos-Velázquez et
al., 2017; Roccuzzo et al., 2015). Given the increased nucleolar size and the increased nucleolar
histone levels during aging, we hypothesized that this might affect proper Cdc14 release from the
nucleolus during anaphase. To test this, we observed both Cdc14:GFP and Htb2:mCherry
dynamics in individual cells, and compared normal cell cycles with cell cycles where GLMs
occurred. Cdc14 exit from the nucleolus is controlled in two stages, first by the CDC Fourteen
Early Anaphase Release (FEAR) network and then the Mitotic Exit Network (MEN)(Rock and
Amon, 2009). By averaging the dynamics of cell cycles that behave normally, it is clear that
Cdc14 begins to exit the nucleolus prior to division of genomic material between mother and
daughter cells (Figure 4A, Video 15). In cell cycles that experience GLMs, however, Cdc14
remains localized to the nucleolus but is released immediately preceding a RETRN event (Figure
4B, Video 15). This agrees with prior work showing that Cdc14 release during anaphase is
required to generate pulling forces within the mother to counteract those in the daughter (Ross
and Cohen-Fix, 2004).
Cdc14 is specifically required for condensation and segregation of repetitive DNA
sequences including the rDNA and telomeres(D'Amours et al., 2004; Sullivan et al., 2004). In
order to further explore the consequences of failed Cdc14 release on anaphase dynamics during
GLM events, we directly observed Chromosome XII by targeting a LacI:GFP reporter to LacO
sites engineered on Chr XII(Ide et al., 2010). During GLMs where the majority of DNA enters
the daughter cell, both Chr XII chromatids remain behind in mother cell (Figure 4C, Video 16-
17). Furthermore, during these GLMs, the Chr XII sister chromatids appear as a single point,
only separating into two distinct foci following a RETRN event (Figure 4C, Video 16-17). This
suggests that delayed Cdc14 activation prevents separation of the rDNA and results in improper
condensation and segregation of Chr XII. During this process, as Chr XII remains localized to
the nucleolus in the mother cell, the remaining genomic content missegregates to the daughter.
We confirmed that this behavior is unique to Chr XII by imaging Chr IV and V during GLMs.
Both copies of Chr IV and Chr V are missegregated to the daughter cells with the rest of the
chromatin during GLMs (Figure 4D,E, Videos 18-19). Thus, we propose that age-associated
expansion of the rDNA and histone depletion lead to dysregulation of Cdc14 during the
metaphase to anaphase transition which results in improper genomic segregation. In both
terminal GLMs and RETRN events, Cdc14 eventually exits the nucleolus to trigger anaphase
(Figure 4 – figure supplement 1).

Fig 4

279
Figure 4. Altered Cdc14 dynamics during aging disrupts segregation of rDNA containing Chromosome XII. **A)**
Dynamics of Cdc14 and Htb2 in mother cells during normal divisions (n=2513). As the cell enters mitosis, Cdc14
leaves the nucleolus which triggers anaphase (indicated by the arrow). **B)** Dynamics of Cdc14 and Htb2 in mother
cells during RETRN events (n=298). Although genomic content has left the mother and entered the daughter, Cdc14
is still localized to the nucleolus. Shortly following Cdc14 release from the nucleolus (indicated by the arrows), the
cells experience a RETRN event. **C)** Direct observation of Chr XII using LacI:GFP and LacO repeats on Chr XII.
When the cell experiences a missegregation event, both chromatids of Chr XII remain behind in the mother until the
RETRN event. After this, a single green dot can be seen in both mother and daughter cells. Mother cell replicative
age equals 8 at the beginning of this timelapse. **D)** Direct observation of Chr IV using TetR:GFP and TetO repeats
on Chr IV. When the cell experiences a GLM, both chromatids of Chr IV move to the daughter along with the
majority of the chromatin. Following a RETRN event, single green dot can be seen in both mother and daughter

cells. Mother cell replicative age equals 5 at the beginning of this timelapse. **E)** Direct observation of Chr V using
TetR:GFP and TetO repeats on Chr V. When the cell experiences a GLM, both chromatids of Chr V move to the
daughter along with the majority of the chromatin. Following a RETRN event, single green dot can be seen in both
mother and daughter cells. Mother cell replicative age equals 19 at the beginning of this timelapse. The gray arrows
mark the location of the labelled chromosomes. Times are indicated in hours:mins. In all panels, n-values report the
number of individual cells analyzed.
297

During mitosis, cells must remove and then replace nucleosomes on each DNA strand.
Because all replicating DNA pulls histones from a common pool, a mismatch between local
histone supply and demand could lead to gaps in nucleosome occupancy, which have been
previously observed in aged cells(Hu et al., 2014). We hypothesized that expansion of the rDNA
due to increasing ERC levels during aging elevates mother cell demand for histones, which is
only partially compensated for by the observed increase in histone expression. By this model,
histone demand should be lower in old *fob1Δ* cells compared to age-matched wild-type cells (due
to fewer ERCs), which should result in a reduced probability of GLMs. Observations of GLMs
in a *fob1Δ* mutant support this model (Figure 5A). In particular, *fob1Δ* and wild-type cells
diverge most dramatically following replicative age 10 (Figure 5A), when the wild-type cells
begin to increase histone levels (Figure 3B).
To directly test the mechanistic link between histone competition and GLMs, we
genetically manipulated the supply of histones (Figure 5B, Figure 5 – figure supplement 1). To
increase histone levels, we first removed Hpc2, a component of the HIR complex which
represses histone transcription(Green et al., 2005). Deletion of *HPC2* results in reduced
frequency of GLMs in aging cells (Figure 5B). Likewise, deletion of *TOM1*, which encodes a
factor required for degradation of excess histones(Singh et al., 2009), also reduced GLMs
(Figure 5B). To reduce histone abundance, we deleted *SPT21*, which encodes a protein that
positively regulates histone expression(Dollard et al., 1994; Kurat et al., 2014) and whose loss
has been previously shown to reduce histone levels and increase rDNA instability(Eriksson et al.,
2012; Kobayashi and Sasaki, 2017). In contrast to deletion of either *HPC2* or *TOM1*, deletion of
SPT21 caused increased frequency of GLMs in aging mother cells. When averaged over the
entire lifespan, *spt21Δ* cells experienced significantly more GLMs than wild-type cells, while
hpc2Δ and *tom1Δ* cells had significantly fewer events (Figure 5C). These observations
demonstrate that altering histone abundance in aging cells is sufficient to modulate the frequency
of GLM events both upward and downward, and support a model whereby rDNA expansion
causes increasing competition for a common histone pool in aging cells which drives the
dramatic increase in GLMs during aging (Figure 5D).

Fig 5

Figure 5. Genomic missegregations increase with age as a result of competition for histones. **A)** Deleting *FOB1* reduces the accumulation of ERCs with age, and also reduces the probability that cells will experience a missegregation event, shown as P(missegregation) at different ages. **B)** Increasing histone supply (*hpc2Δ* & *tom1Δ*) reduces the fraction of cells that have missegregation events, while reducing histone supply (*spt21Δ*) increases the fraction of cells with missegregation events. Error bars are standard error. N-values report the number of cells. **C)** The mean number of genome level missegregations (GLM) events per mother cell by replicative age twenty. Error bars are 95% confidence intervals generated by bootstrapping with replacement. (*) Indicates samples with confidence intervals that do not overlap with wild-type. **D)** In young cells the common pool of histones is available for genomic DNA, but as cells age the demand for histones increases as ERCs compete with genomic DNA for the free pool of histones. Old *fob1Δ* cells have fewer ERCs which lowers the competition for histones. By increasing the common histone supply (*hpc2Δ* and *tom1Δ*) competition between genomic DNA and ERCs is reduced and the missegregation rate is reduced.

Because function declines in many different and subtle ways during aging, catastrophic failures and homeostatic systems like the those uncovered here may only be detected in aged organisms. Imaging of individual yeast cells through microfluidic trapping allowed us to observe GLMs that occur in most mother cells at some point during their lives. These events are exceedingly rare in young cells, are actively repaired through transient arrest of the cell cycle, and likely result from insufficient histone availability due to expansion of the nuclear genome via accumulation of rDNA. The rDNA has been linked to genomic instability in yeast(Ganley et al., 2009; Ide et al., 2010; Saka et al., 2013) and within cancer cell lines(Xu et al., 2017), and nucleolar size is anti-correlated with lifespan across organisms(Tiku et al., 2016). Nucleolar structure has also been linked to cancer pathogenesis(Lindström et al., 2018). Given the ubiquity of aneuploidy in age-related human cancers(Sansregret and Swanton, 2017), we speculate that mechanisms for responding to mitotic failures may be important to cope with age-associated

genomic instability in multicellular eukaryotes. Interestingly, although humans have only five acrocentric chromosomes with rDNA repeats, these are far more likely to result in developmental trisomy than other chromosomes(Nagaoka et al., 2012). Whether the mechanisms accounting for age-related increases in genomic instability uncovered here, and the RETRN process to repair these events, are conserved in higher eukaryotes requires further investigation. Such research into the behavior of natural cellular processes during aging could provide insights into age-related pathology and uncover potential targets for intervention.

Acknowledgements: We would particularly like to thank S. Biggins, B. Brewer and M. Raghuraman for constructive discussions. We also thank L. Veenhoff and Kaeberlein lab members for feedback and advice. Strains YSI129, AMY914 and AMY1081were generous gifts from Jessica Tyler and Adele Marston. This work was supported by NIH grants T32AG000057, R01AG056359, and P30AG013280.

References

Bakker E. 2016. Quantitative fluorescence microscopy methods for studying transcription with application to the yeast GAL1 promoter. The University of Edinburgh.

Bakker E, Swain PS, Crane MM. 2017. Morphologically constrained and data informed cell segmentation of budding yeast. *Bioinformatics*. doi:10.1093/bioinformatics/btx550

Beach RR, Ricci-Tam C, Brennan CM, Moomau CA, Hsu P-H, Hua B, Silberman RE, Springer M, Amon A. 2017. Aneuploidy Causes Non-genetic Individuality. *Cell* **169**:229–242.e21.

Bean JM, Siggia ED, Cross FR. 2006. Coherence and Timing of Cell Cycle Start Examined at Single-Cell Resolution. *Mol Cell* **21**:3–14.

Bi E, Park H-O. 2012. Cell Polarization and Cytokinesis in Budding Yeast. *Genetics* **191**:347–387.

Cai L, Dalal CK, Elowitz MB. 2008. Frequency-modulated nuclear localization bursts coordinate gene regulation. *Nature* **455**:485–490.

Charvin G, Oikonomou C, Siggia ED, Cross FR. 2010. Origin of irreversibility of cell cycle start in budding yeast. *PLoS Biol* **8**. doi:10.1371/journal.pbio.1000284

Clemente-Blanco A, Sen N, Mayan-Santos M, Sacristán MP, Graham B, Jarmuz A, Giess A, Webb E, Game L, Eick D, Bueno A, Merkenschlager M, Aragón L. 2011. Cdc14 phosphatase promotes segregation of telomeres through repression of RNA polymerase II transcription. *Nat Cell Biol* **13**:1450–1456.

Costanzo M, Nishikawa JL, Tang X, Millman JS, Schub O, Breitkreuz K, Dewar D, Rupes I, Andrews B, Tyers M. 2004. CDK activity antagonizes Whi5, an inhibitor of G1/S transcription in yeast. *Cell* **117**:899–913.

Crane MM, Clark IBN, Bakker E, Smith S, Swain PS. 2014. A microfluidic system for studying ageing and dynamic single-cell responses in budding yeast. *PLoS One* **9**:e100042.

D'Amours D, Stegmeier F, Amon A. 2004. Cdc14 and condensin control the dissolution of cohesin-independent chromosome linkages at repeated DNA. *Cell* **117**:455–469.

Dang W, Steffen KK, Perry R, Dorsey JA, Johnson FB, Shilatifard A, Kaeberlein M, Kennedy BK, Berger SL. 2009. Histone H4 lysine 16 acetylation regulates cellular lifespan. *Nature* **459**:802–807.

Defossez PA, Prusty R, Kaeberlein M, Lin SJ, Ferrigno P, Silver PA, Keil RL, Guarente L. 1999. Elimination of replication block protein Fob1 extends the life span of yeast mother cells. *Mol Cell* **3**:447–455.

de los Santos-Velázquez AI, de Oya IG, Manzano-López J, Monje-Casas F. 2017. Late rDNA Condensation Ensures Timely Cdc14 Release and Coordination of Mitotic Exit Signaling with Nucleolar Segregation. *Curr Biol* **27**:3248–3263.e5.

Di Talia S, Skotheim JM, Bean JM, Siggia ED, Cross FR. 2007. The effects of molecular noise and size

control on variability in the budding yeast cell cycle. *Nature* **448**:947–951.

Dollard C, Ricupero-Hovasse SL, Natsoulis G, Boeke JD, Winston F. 1994. SPT10 and SPT21 are
required for transcription of particular histone genes in Saccharomyces cerevisiae. *Mol Cell Biol*
14:5223–5228.

Dotiwala F, Haase J, Arbel-Eden A, Bloom K, Haber JE. 2007. The yeast DNA damage checkpoint
proteins control a cytoplasmic response to DNA damage. *Proc Natl Acad Sci U S A* **104**:11358–
11363.

Eriksson PR, Ganguli D, Nagarajavel V, Clark DJ. 2012. Regulation of histone gene expression in
budding yeast. *Genetics* **191**:7–20.

Fernius J, Marston AL. 2009. Establishment of Cohesion at the Pericentromere by the Ctf19 Kinetochore
Subcomplex and the Replication Fork-Associated Factor, Csm3. *PLoS Genet* **5**:e1000629.

Feser J, Truong D, Das C, Carson JJ, Kieft J, Harkness T, Tyler JK. 2010. Elevated histone expression
promotes life span extension. *Mol Cell* **39**:724–735.

Finley KR, Bouchonville KJ, Quick A, Berman J. 2008. Dynein-dependent nuclear dynamics affect
morphogenesis in Candida albicans by means of the Bub2p spindle checkpoint. *J Cell Sci* **121**:466–
476.

Flach J, Bakker ST, Mohrin M, Conroy PC, Pietras EM, Reynaud D, Alvarez S, Diolaiti ME, Ugarte F,
Forsberg EC, Le Beau MM, Stohr BA, Méndez J, Morrison CG, Passegué E. 2014. Replication
stress is a potent driver of functional decline in ageing haematopoietic stem cells. *Nature* **512**:198–
202.

Ganley ARD, Ide S, Saka K, Kobayashi T. 2009. The Effect of Replication Initiation on Gene
Amplification in the rDNA and Its Relationship to Aging. *Mol Cell* **35**:683–693.

Gordon DJ, Resio B, Pellman D. 2012. Causes and consequences of aneuploidy in cancer. *Nat Rev Genet*
13:189–203.

Granados AA, Crane MM, Montano-Gutierrez LF, Tanaka RJ, Voliotis M, Swain PS. 2017. Distributing
tasks via multiple input pathways increases cellular survival in stress. *Elife* **6**.
doi:10.7554/eLife.21415

Green EM, Antczak AJ, Bailey AO, Franco AA, Wu KJ, Yates JR 3rd, Kaufman PD. 2005. Replication-
independent histone deposition by the HIR complex and Asf1. *Curr Biol* **15**:2044–2049.

Hanahan D, Weinberg RA. 2011. Hallmarks of cancer: the next generation. *Cell* **144**:646–674.

Hauer MH, Seeber A, Singh V, Thierry R, Sack R, Amitai A, Kryzhanovska M, Eglinger J, Holcman D,
Owen-Hughes T, Gasser SM. 2017. Histone degradation in response to DNA damage enhances
chromatin dynamics and recombination rates. *Nat Struct Mol Biol* **24**:99–107.

Henderson KA, Hughes AL, Gottschling DE. 2014. Mother-daughter asymmetry of pH underlies aging
and rejuvenation in yeast. *Elife* **3**:e03504.

Huh W, Falvo JV, Gerke LC, Carroll AS, Howson RW, Weissman JS, O'Shea EK. 2003. Global analysis
of protein localization in budding yeast. *Nature* **425**:686–691.

Hu Z, Chen K, Xia Z, Chavez M, Pal S, Seol JH, Chen CC, Li W, Tyler JK. 2014. Nucleosome loss leads
to global transcriptional up-regulation and genomic instability during yeast aging. *Genes and*
Development **28**:396–408.

Ide S, Miyazaki T, Maki H, Kobayashi T. 2010. Abundance of ribosomal RNA gene copies maintains
genome integrity. *Science* **327**:693–696.

Janssens GE, Meinema AC, González J, Wolters JC, Schmidt A, Guryev V, Bischoff R, Wit EC,
Veenhoff LM, Heinemann M. 2015. Protein biogenesis machinery is a driver of replicative aging in
yeast. *Elife* **4**:e08527.

Jorgensen P, Edgington NP, Schneider BL, Rupes I, Tyers M, Futcher B. 2007. The size of the nucleus
increases as yeast cells grow. *Mol Biol Cell* **18**:3523–3532.

Kobayashi T, Heck DJ, Nomura M, Horiuchi T. 1998. Expansion and contraction of ribosomal DNA
repeats in Saccharomyces cerevisiae: requirement of replication fork blocking (Fob1) protein and the
role of RNA polymerase I. *Genes Dev* **12**:3821–3830.

Kobayashi T, Sasaki M. 2017. Ribosomal DNA stability is supported by many "buffer genes"-
introduction to the Yeast rDNA Stability Database. *FEMS Yeast Res* **17**. doi:10.1093/femsyr/fox001
Kruegel U, Robison B, Dange T, Kahlert G, Delaney JR, Kotireddy S, Tsuchiya M, Tsuchiyama S,
Murakami CJ, Schleit J, Sutphin G, Carr D, Tar K, Dittmar G, Kaeberlein M, Kennedy BK, Schmidt
M. 2011. Elevated proteasome capacity extends replicative lifespan in saccharomyces cerevisiae.
PLoS Genet **7**. doi:10.1371/journal.pgen.1002253
Kurat CF, Lambert J-P, Petschnigg J, Friesen H, Pawson T, Rosebrock A, Gingras A-C, Fillingham J,
Andrews B. 2014. Cell cycle-regulated oscillator coordinates core histone gene transcription through
histone acetylation. *Proc Natl Acad Sci U S A* **111**:14124–14129.
Lindström MS, Jurada D, Bursac S, Orsolic I, Bartek J, Volarevic S. 2018. Nucleolus as an emerging hub
in maintenance of genome stability and cancer pathogenesis. *Oncogene*. doi:10.1038/s41388-017-
0121-z
London N, Biggins S. 2014. Signalling dynamics in the spindle checkpoint response. *Nat Rev Mol Cell
Biol* **15**:736.
Maciejowski J, Li Y, Bosco N, Campbell PJ, De Lange T. 2015. Chromothripsis and Kataegis Induced by
Telomere Crisis. *Cell* **163**:1641–1654.
McCormick MA, Delaney JR, Tsuchiya M, Tsuchiyama S, Shemorry A, Sim S, Chou AC-Z, Ahmed U,
Carr D, Murakami CJ, Schleit J, Sutphin GL, Wasko BM, Bennett CF, Wang AM, Olsen B, Beyer
RP, Bammler TK, Prunkard D, Johnson SC, Pennypacker JK, An E, Anies A, Castanza AS, Choi E,
Dang N, Enerio S, Fletcher M, Fox L, Goswami S, Higgins SA, Holmberg MA, Hu D, Hui J, Jelic
M, Jeong K-S, Johnston E, Kerr EO, Kim J, Kim D, Kirkland K, Klum S, Kotireddy S, Liao E, Lim
M, Lin MS, Lo WC, Lockshon D, Miller HA, Moller RM, Muller B, Oakes J, Pak DN, Peng ZJ,
Pham KM, Pollard TG, Pradeep P, Pruett D, Rai D, Robison B, Rodriguez AA, Ros B, Sage M,
Singh MK, Smith ED, Snead K, Solanky A, Spector BL, Steffen KK, Tchao BN, Ting MK, Vander
Wende H, Wang D, Welton KL, Westman EA, Brem RB, Liu X-G, Suh Y, Zhou Z, Kaeberlein M,
Kennedy BK. 2015. A Comprehensive Analysis of Replicative Lifespan in 4,698 Single-Gene
Deletion Strains Uncovers Conserved Mechanisms of Aging. *Cell Metab* **22**:895–906.
Nagaoka SI, Hassold TJ, Hunt PA. 2012. Human aneuploidy: mechanisms and new insights into an age-
old problem. *Nat Rev Genet* **13**:493–504.
Neurohr GE, Terry RL, Sandikci A, Zou K, Li H, Amon A. 2018. Deregulation of the G1/S-phase
transition is the proximal cause of mortality in old yeast mother cells. *Genes Dev*.
doi:10.1101/gad.312140.118
Otsu N. 1979. A Threshold Selection Method from Gray-Level Histograms. *IEEE Trans Syst Man Cybern*
9:62–66.
Palmer RE, Koval M, Koshland D. 1989. The dynamics of chromosome movement in the budding yeast
Saccharomyces cerevisiae. *J Cell Biol* **109**:3355–3366.
Roccuzzo M, Visintin C, Tili F, Visintin R. 2015. FEAR-mediated activation of Cdc14 is the limiting step
for spindle elongation and anaphase progression. *Nat Cell Biol* **17**:251–261.
Rock JM, Amon A. 2009. The FEAR network. *Curr Biol* **19**:R1063–8.
Ross KE, Cohen-Fix O. 2004. A role for the FEAR pathway in nuclear positioning during anaphase. *Dev
Cell* **6**:729–735.
Saka K, Ide S, Ganley ARD, Kobayashi T. 2013. Cellular senescence in yeast is regulated by rDNA
noncoding transcription. *Curr Biol* **23**:1794–1798.
Sansregret L, Swanton C. 2017. The Role of Aneuploidy in Cancer Evolution. *Cold Spring Harb Perspect
Med* **7**. doi:10.1101/cshperspect.a028373
Santaguida S, Amon A. 2015. Short- and long-term effects of chromosome mis-segregation and
aneuploidy. *Nat Rev Mol Cell Biol* **16**:473–485.
Shaner NC, Steinbach PA, Tsien RY. 2005. A guide to choosing fluorescent proteins. *Nat Methods
2:905–909.
Sharma S, Kelly TK, Jones PA. 2010. Epigenetics in cancer. *Carcinogenesis* **31**:27–36.

Shou W, Seol JH, Shevchenko A, Baskerville C, Moazed D, Chen ZW, Jang J, Shevchenko A,
Charbonneau H, Deshaies RJ. 1999. Exit from mitosis is triggered by Tem1-dependent release of the
protein phosphatase Cdc14 from nucleolar RENT complex. *Cell* **97**:233–244.
Sinclair DA, Guarente L. 1997. Extrachromosomal rDNA circles--a cause of aging in yeast. *Cell*
91:1033–1042.
Sinclair DA, Mills K, Guarente L. 1997. Accelerated aging and nucleolar fragmentation in yeast sgs1
mutants. *Science* **277**:1313–1316.
Singh RK, Kabbaj M-HM, Paik J, Gunjan A. 2009. Histone levels are regulated by phosphorylation and
ubiquitylation-dependent proteolysis. *Nat Cell Biol* **11**:925–933.
Sullivan M, Higuchi T, Katis VL, Uhlmann F. 2004. Cdc14 phosphatase induces rDNA condensation and
resolves cohesin-independent cohesion during budding yeast anaphase. *Cell* **117**:471–482.
Sunshine AB, Ong GT, Nickerson DP, Carr D, Murakami CJ, Wasko BM, Shemorry A, Merz AJ,
Kaeberlein M, Dunham MJ. 2016. Aneuploidy shortens replicative lifespan in Saccharomyces
cerevisiae. *Aging Cell* **15**:317–324.
Thrower DA, Stemple J, Yeh E, Bloom K. 2003. Nuclear oscillations and nuclear filament formation
accompany single-strand annealing repair of a dicentric chromosome in Saccharomyces cerevisiae. *J
Cell Sci* **116**:561–569.
Tiku V, Jain C, Raz Y, Nakamura S, Heestand B, Liu W, Späth M, Suchiman HED, Müller R-U,
Slagboom PE, Partridge L, Antebi A. 2016. Small nucleoli are a cellular hallmark of longevity. *Nat
Commun* **8**:16083.
Weiss EL. 2012. Mitotic exit and separation of mother and daughter cells. *Genetics* **192**:1165–1202.
Winzeler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, Andre B, Bangham R, Benito R,
Boeke JD, Bussey H, Chu AM, Connelly C, Davis K, Dietrich F, Dow SW, El Bakkoury M, Foury
F, Friend SH, Gentalen E, Giaever G, Hegemann JH, Jones T, Laub M, Liao H, Liebundguth N,
Lockhart DJ, Lucau-Danila A, Lussier M, M'Rabet N, Menard P, Mittmann M, Pai C, Rebischung
C, Revuelta JL, Riles L, Roberts CJ, Ross-MacDonald P, Scherens B, Snyder M, Sookhai-Mahadeo
S, Storms RK, Véronneau S, Voet M, Volckaert G, Ward TR, Wysocki R, Yen GS, Yu K,
Zimmermann K, Philippsen P, Johnston M, Davis RW. 1999. Functional Characterization of the S.
cerevisiae Genome by Gene Deletion and Parallel Analysis. *Science* **285**:901–906.
Witkin KL, Chong Y, Shao S, Webster MT, Lahiri S, Walters AD, Lee B, Koh JLY, Prinz WA, Andrews
BJ, Cohen-Fix O. 2012. The budding yeast nuclear envelope adjacent to the nucleolus serves as a
membrane sink during mitotic delay. *Curr Biol* **22**:1128–1133.
Xu B, Li H, Perry JM, Singh VP, Unruh J, Yu Z, Zakari M, McDowell W, Li L, Gerton JL. 2017.
Ribosomal DNA copy number loss and sequence variation in cancer. *PLoS Genet* **13**:e1006771.
Yang SS, Yeh E, Salmon ED, Bloom K. 1997. Identification of a mid-anaphase checkpoint in budding
yeast. *J Cell Biol* **136**:345–354.
Yeh E, Skibbens RV, Cheng JW, Salmon ED, Bloom K. 1995. Spindle dynamics and cell cycle
regulation of dynein in the budding yeast, Saccharomyces cerevisiae. *J Cell Biol* **130**:687–700.
Yeh E, Yang C, Chin E, Maddox P, Salmon ED, Lew DJ, Bloom K. 2000. Dynamic positioning of
mitotic spindles in yeast: role of microtubule motors and cortical determinants. *Mol Biol Cell*
11:3949–3961.
Zhang C-Z, Spektor A, Cornils H, Francis JM, Jackson EK, Liu S, Meyerson M, Pellman D. 2015.
Chromothripsis from DNA damage in micronuclei. *Nature* **522**:179–184.

Supplementary Materials and Methods

Microfluidics

Cells were imaged using a PDMS microfluidic flow chamber modified from an earlier design (Crane et al., 2014) to increase retention over the replicative lifespan of the mother cells. The microfluidic device was composed of multiple chambers in the same fashion as (Granados et al., 2017), which allowed individual genotypes to be exposed to identical environments and imaged in the same experiment while being physically isolated. Cells were loaded according to previously published methods (Granados et al., 2017). A volumetric flow rate of 3-7 µL/min per chamber was used, with the flow rate starting low, and increasing during the experiment to improve mother cell retention and to ensure that cells do not aggregate, which can clog the device.

Microscopy

Cells were imaged using a Nikon Ti-2000 microscope with a 40X oil immersion objective with a 1.3 NA and using the Nikon Perfect Focus System. An enclosed incubation chamber was used to maintain a stable 30C environment for the duration of the experiment. Two Aladdin syringe pumps were used for media flow. An LED illumination system (Excelitas 110-LED) was used to provide consistent excitation energies, and to minimize the exposure, illumination was triggered by the camera. Images were acquired using a Hammamatsu Orca Flash 4.0 V2. The microscope was controlled by custom software written in Matlab® and Micromanager.

Images were corrected for illumination artifacts in two stages. First, to correct for individual differences in the pixel biases, 1,000 images were acquired with no illumination, and the individual pixel means were determined. Second, to correct for flatness of field, a fluorescent dye was added to a microfluidic device instead of using a slide with dye. Using a slide containing dye introduces a large amount of out-of-focus light, which results in an underestimation of the field curvature. In order to compensate for the microfluidic features, 1,000 images were acquired each with a small offset in the x and y positions. Images were then dilated, and the median value at each location was used. Thus, for each image, the camera bias for that pixel was subtracted, and then it was multiplied by a flatness of field correction factor.

Images were acquired at 5 min intervals for bright-field and fluorescent channels. The fluorescence excitation power was set to 25% for all imaging except the GFP tagged histones, where it was set to 12%. Fluorescence and brightfield light was activated during image acquisition and all other lights in the room were turned off. For bright-field, 3 z-sections were acquired with 2.5 µm intervals, exposure times of 30 ms and were used for automated segmentation and tracking. For the fluorescent channels, 3 z-sections were acquired with 1.5 µm spacing. GFP images were acquired using a Chroma ET49002 filter set, and mCherry images were acquired using a Chroma ET49306. GFP images were acquired using exposure times of 60ms for all proteins except Htb2 and Hta2 which were acquire using a 30ms exposure time. mCherry images were acquired using a 60 ms exposure time. These imaging conditions were found to work as a reasonable compromise between the desire for frequent, dense imaging to enable identification of missegregations and retrograde transport, while also minimizing phototoxicity. We performed control experiments to verify that these exposure conditions did not affect the rates of genomic missegregation or replicative lifespan (Figure 1 – figure supplement

589   5). Each strain was imaged in multiple independent experimental runs, each with approximately
590   equal numbers of cells.
591
Data processing and single cell scoring
Following data acquisition, cells were identified and tracked using previously published
software(Bakker et al., 2017). This identified the cell outline, and performed initial tracking of
the cells through time. To ensure that only young, healthy cells were assessed, we only used cells
that were identified in the first three hours of the experiment. Birth events for these cells were
then manually scored, and any errors in tracking were corrected. This was all done using the
bright-field images. Birth events were scored by multiple observers who were blinded. Because
individual cells can be lost from traps prior to death, it can be challenging to know whether using
censored (lost) cells is most appropriate. In the supplementary information, all data is presented
with and without censoring. In the main text, plots aligned based on increasing age used all cells
present at that age, even if they were later lost from the device. For plots aligned by death, only
cells that had either died or senesced (failed to initiate a new cell division but did not visibly lose
cell wall integrity during the experiment) were used. Because censoring in lifespan experiments
relies on the assumption that losses are unbiased, we provide replicative lifespan curves both
including and excluding censored cells for all strains. Censoring does not change the
interpretation or statistical outcome of any of the experiments presented here.
608
609   For each cell division, the mean histone levels over that cell cycle were used. For cell cycles that
610   last more than three hours, only the first three hours were used to determine the histone levels.
611   This ensured that cells which have a terminal missegregation at end of life do not show an
612   inaccurately low histone level for the last cell cycle.
613
Following manual scoring of birth events, the fluorescent channel containing the histones was
used to observe the missegregation dynamics. To ensure consistent scoring across experiments
and eliminate bias, information about the experiments was masked from the scorer until after the
data was evaluated. Following genome level missegregations, RETRN events were defined as
where the histone fluorescence decreased in the daughter cell while simultaneously increasing in
the mother cell. This prevented any changes in focus or gradual fluorescence increases due to
fluorophore maturation from being inadvertently scored as a RETRN event. RETRN events were
scored at the timepoint that the histones return to the mother cell. During cell cycles where cells
had multiple retrograde events during the same cell cycle, only the final RETRN event was
scored. Events were scored as terminal missegregation events if, prior to a correction, the
daughter cell visibly separated from the mother cell (indicating cytokinesis) or if the mother
died.
626
Fluorescence quantification
Quantification of the level of protein localized to either the nucleolus (Cdc14) or the nucleus
(Whi5), was done using a measure of how asymmetrically distributed the fluorescent signal was.
Specifically, we used average brightness of the top 2% of pixels, divided by the cell median. By
normalizing to median fluorescence, we corrected for any changes in fluorescence that could
occur as a result of photobleaching. This method has been used previously as an accurate
measure of the fraction of protein that is nuclear localized(Cai et al., 2008; Granados et al.,
2017). For Myo1 quantification, we used the mean fluorescence level along the periphery of the

cell as segmented. The nucleolar localized fluorescent protein (Utp13:GFP) was used to segment
the nucleolus and determine the nucleolar size. This was done by calculating a segmentation
threshold using Otsu's threshold(Otsu, 1979) for each cell applied to the maximum projection of
the fluorescent image stack. This approach was found to have good agreement with earlier
estimates of nucleolar size in young cells grown in glucose(Jorgensen et al., 2007).
640
Because we are utilizing an epi-fluorescent microscope and as a result of out-of-focus light,
mean fluorescence is not directly linked to protein concentration (Bakker, 2016). Because the
fluorescence levels change dramatically during aging, segmenting out the nucleus to determine
fluorescence levels could introduce significant biases as the cells age. Additionally, taking the
mean fluorescence level could introduce significant errors depending on the segmentation
accuracy. Instead, to normalize for changes, we sum a constant area within each cell. This
prevents segmentation (of either the nucleus or cell) from contributing to the determination of
histone amount. Pixels are sorted from high to low, and the top fraction corresponding to a circle
with diameter of 3.8 μm are used to calculate the mean.
650
Yeast Strains and Growth
The GFP strains were all acquired from the yeast GFP collection (Huh et al., 2003). The
Htb2:mCherry strain was created by mating and sporulation of the strain from (Granados et al.,
2017). This strain was then crossed with the relevant GFP strains (Nup49:GFP, Myo1:GFP,
Tub1:GFP, Spc72:GFP, Cdc14:GFP, Utp13:GFP) or deletion strains (*hpc2Δ*, *fob1Δ*, *spt21Δ*,
tom1Δ, *mad3Δ*) from the deletion collection (Winzeler et al., 1999) and then confirmed by
colony PCR. The LacI-GFP strain with 50 LacO repeats on ChrXII was obtained from (Ide et al.,
2010). The strains containing TetR-GFP and TetO repeats integrated into ChrIV or ChrV were
obtained from (Fernius and Marston, 2009). These were then crossed with the strain containing
Htb2:mCherry. Complete list of strains available in Table S1.
661
Prior to each experiment, single colonies were picked into SC media (Sunrise Biosciences) with
2% dextrose. Cells were grown overnight, and then diluted 1:200 in fresh media and grown for
5-6h. Prior to loading into the microfluidic device, 0.5 mL of SC 2% dextrose with 0.5% BSA
was added to each 5 mL culture to prevent the cells from adhering to the PDMS during loading.
During experiments, SC media with 2% dextrose and 0.1% BSA was used, and cells were
imaged for 72h.
668
Statistical Analysis
Error bars in the figures which contained bar plots were generated by bootstrapping with
replacement, and then determining the 95% confidence intervals. Error bars in figures with line
plots are standard error. Statistical significance for lifespan was determined using the log-rank
test. Log-rank test was performed with, and without, censored cells that were lost prior to
senescence or death. To compare distributions (such as numbers of missegregation events over
the lifespan), a two-tailed t-test assuming equal variance was used. Correlations between histones
or missegregation events and remaining replicative lifespan were calculated with the Spearman
correlation using the population of cells alive at each replicative age. Correlation between
Htb2:mCherry levels and DAPI staining were also done using the Spearman correlation.
679

Supplementary Text

Differences between censored and uncensored survival data

Frequently in experiments or clinical studies that involve the generation of survival curves, some samples will be removed from the population under observation. For example, a patient may leave a study not because of death, but because they move to a different country. This can be treated in a relatively straightforward manner statistically including these individuals in the analysis until the point that they are lost (or censored). This relies on the assumption that there is no bias in whether a sample is lost or retained. A recurring concern with microfluidic aging experiments involving yeast is whether there is a bias in how cells are lost or retained. This appears especially important when the mutation or transgene affects cell morphology or cell cycle, as this can result in a bias in which cells are lost from the traps. To reduce the likelihood that our observations were directly affected by loss rates, in the main text we have plotted all cells that were present at that replicative age for plots from birth. Thus, if a cell was lost at replicative age 20, it was included in the plots until age 19. In the supplementary materials we have also repeated every plot, but only using cells that died in the microfluidic device. Given that this is an altered population distribution and smaller number of cells, these plots are slightly different, but they do not affect the conclusions. Furthermore, in the plots where cells are aligned by birth rather than death, we are forced to only use cells that die in the device. For replicative lifespans shown in the supplementary, we include survival curves with and without censored cells.

Aligning cells from birth or from death

Cells can be aligned either by birth (counting up from replicative age = 0), or by death (counting back from death). Either processing makes some assumptions about how similar cells are to one another. If cells are most similar to each other when they are born, aligning by birth makes sense, and as the replicative age increases, the number of samples decreases because cells are removed by death or senescence. In contrast, assuming that cells are similar at death implies that the phenotype of interest is most similar as cells approach death. For example, the average time cells take to proceed through each cell cycle increases geometrically when cells are aligned by birth, but exponentially when aligned by death. In the supplementary figures, we show the population means aligned using both approaches.

Differences between Htb2:mCherry and Hta2:GFP

Although both of these strains were found to have similar numbers of missegregation events during their lifetimes, and similar fractions of these events were corrected (Figure 1 – figure supplement 4B,C), there are subtle differences between the strains. Most notably, the strain with Hta2:GFP had what we consider to be a normal lifespan for this background (median RLS= 21 for cells that died/senesced in the device, median RLS=24 including censored cells, Figure 1 – figure supplement 4D,E). The strain with Htb2:mCherry, however, had a somewhat shorter lifespan (median RLS = 16 for cells that died/senesced, median RLS=18 including censored cells, Figure 1 – figure supplement 4D,E). Removing FOB1, however, results in an increase of the replicative lifespan of this strain by ~30% (Figure 3 – figure supplement 3), which is in line with results from literature (McCormick et al., 2015). Furthermore, the increase in replicative lifespan as a result of increased histone transcription has been less thoroughly studied, but our results are in line with those previously reported by another group (Feser et al., 2010; Kruegel et

725 al., 2011). Thus, although there is an unexpected reduction in lifespan for the Htb2:mCherry
726 strain, we do not believe that it affects our results.
727
Likewise, as shown in the main text, we determined the correlation between missegregation
events and remaining lifespan at the single cell level. The correlation is between the binary
presence or absence of a missegregation event during a cell cycle and the remaining lifespan.
Strikingly, as shown in Figure 1 – figure supplement 2, for both strains, the correlation between
missegregation events and remaining replicative lifespan is the same for both Htb2:mCherry and
Hta2:GFP. This is in spite of the difference in absolute lifespan between the two strains.
734
735 Because GFP fluorescence is much more affected than mCherry fluorescence by changing pH
736 (Shaner et al., 2005), and the pH of the cytoplasm in aging yeast has previously been shown to
737 increase (Henderson et al., 2014), we chose to perform the majority of the experiments using
738 mCherry. This ensured that any changes in pH homeostasis during aging would not affect our
739 measurements of histone levels.
740
GFP Tagged Histones and Correlation with remaining RLS
As discussed in the main text, we imaged GFP tagged histones acquired from the GFP collection
(Hta2, Htb2). This allowed us to determine that the increase in histone levels was a general
aspect of aging physiology, not confined to specific histones. In order to determine whether
histone levels were accurate biomarkers that predicted remaining lifespan, we used the single cell
measurements collected for both strains (Figure 3 – figure supplement 1). At each replicative
age, cells still alive are used to determine the correlation between their remaining replicative
lifespan and the single cell histone levels. Thus, for prediction of remaining lifespan at
replicative age 5, only cells that bud more than 5 times are included. The negative correlation
between histone levels and remaining lifespan is similarly true for the GFP tagged histones
imaged: Hta2, Htb2 (Figure 3 – figure supplement 1A). Furthermore, all histones are similarly
predictive of remaining lifespan, and become increasingly predictive as cells age.
753
To provide a more complete picture of the data acquired, we plotted the GFP mean histone levels
where single cells are aligned based on their replicative ages (Figure 3 – figure supplement 1B).
As the population ages, the underlying distribution changes as cells die and are removed from the
pool. This creates some variability with the increasing replicative age of the population. When
individual cells are aligned by death, the population mean for histone levels at replicative ages
preceding death looks quite different (Figure 3 – figure supplement 1C). By aligning cells by
death, it is clear that histone levels begin to increase around 5-10 replications prior to death.
When including all cells, including those that were lost from the traps, cells expressing GFP
tagged histones have similar replicative lifespans (Figure 3 – figure supplement 1D,E).
763
Single Cell DNA Levels and FOB1
As discussed in the main text, we determined the correlation between histone levels and DNA by
comparing, in individual cells, levels of Htb2:mCherry with levels of DAPI. To do this, mother
cells were aged in the device for 16h, and then fixed with ethanol and stained with DAPI. To
determine whether the accumulation of ERCs with age leads to a detectable increase in DNA
content, we compared DAPI staining of wild-type and *fob1Δ*. We acquired age-matched mother
cells by using a multi-chamber device to simultaneously age wild-type and *fob1Δ* cells.

Comparing mothers that had grown in the device for 16h (median of 13 divisions and thus
middle aged), we confirmed that DNA levels increase in wild-type cells are significantly higher
compared with fob1Δ. The confirmed that the deletion of FOB1 decreased DNA content in aging
cells, and not just histone levels. By manually scoring the number of divisions each strain had up
until the point of ethanol fixation, we determined that there was no difference between strains in
the number of daughters (Figure 3C, p=0.42). Although the replicative ages for each strain were
statistically indistinguishable, the wild-type strain had a statistically significant increase in the
DAPI staining levels (Figure 3D, p=0.0002). Thus, *fob1Δ* cells accumulate both histones and
excess DNA at lower rates compared with wild-type cells. Furthermore, we compared Htb2 and
DAPI levels at the single cell level, and histone levels are highly predictive of DNA content at
the single cell level (Figure 1 – figure supplement 3).
782
<u>Manipulating histone demand (fob1Δ)</u>
To provide a more complete picture of how Htb2 levels changed during aging in *fob1Δ* cells
compared with wild-type, we performed the same procedure as with the GFP strains but using
only the cells that die in the device to complement the plots in the main text. These plots look
very similar to the plots using censored cells indicating that censoring doesn't alter the
population distribution. By aligning cells based on the replicative age, and plotting the mean
Htb2:mCherry levels of the population, it is clear that there is a dramatic difference between
wild-type and *fob1Δ* cells in Htb2 levels (Figure 3 – figure supplement 3A). This also reflects the
missegregation rate (Figure 3 – figure supplement 3B). Missegregations are less predictive of
death in *fob1Δ* cells than wild-type cells (Figure 3 – figure supplement 3C). When cells are
aligned by death, as opposed to birth, both wild-type and fob1Δ cells experience an increase in
histone levels, although *fob1Δ* cells are attenuated relative to wild-type (Figure 3 – figure
supplement 3D). Because *fob1Δ* cells live longer, it would be reasonable to expect that by the
end-of-life, the missegregation rate is closer between wild-type and *fob1Δ* cells than when
aligned by birth (Figure 3 – figure supplement 3E). Similar to previous reports (Defossez et al.,
1999), we find a significant increase the replicative lifespan of *fob1Δ* cells compared with wild-
type (Figure 3 – figure supplement 3F). Survival curves of only the cells which died in the
device, and thus were used in the analysis, showed similar lifespan differences (Figure 3 – figure
supplement 3G).
802
<u>Manipulating histone supply (*tom1Δ*, *hpc2Δ* and *spt21Δ*)</u>
To provide a more complete picture of how the manipulation of the histone supply affected
lifespan and Htb2 levels, we performed a similar procedure as previously described. Spt21
positively regulates the transcription of all histone genes, and Hpc2 negatively regulates histone
genes. The protein Tom1 is involved in the ubiquitination and degradation of excess histone
proteins. Thus by deleting *HPC2*, we upregulate histone transcription, deleting *SPT21* suppresses
histone transcription and deleting *TOM1* allows excess histone proteins to remain in the cell.
Similar to the figures in the main text, we used only cells that die in the device to plot
missegregation rates aligned by birth (Figure 5 – figure supplement 1A) and by death (Figure 5 –
figure supplement 1B). As previously identified, *hpc2Δ* and *tom1Δ* cells live significantly longer
than wild-type cells (Figure 5 – figure supplement 1C,D). Surprisingly, *spt21Δ* cells do not
appear shorter lived compared to wild-type (Figure 5 – figure supplement 1C,D). These trends
are maintained in survival curves of only cells that die in the device (Figure 5 – figure
supplement 1C).

Strain Name	Genotype	Source
BY4741	MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0	
BY4742	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0	
YSI129	ade2-1 ura3-1 his3-11 trp1-1 leu2-3,112 can1-100 fob1Δ::LEU2 his3-11::GFP-LacI-HIS3 LacO(50)-ADE2:445kb ChXII (110 rDNA copies)	Ide, et al. 2010
AMY914	MATa ura3 trp1 his3-11 MET-CDC20::URA pURA::tetR-GFP::LEU2 cenIV::tetOx448::URA3	Fernius, et al. 2009
AMY1081	MATa ade2-1 leu2-3 trp1-1 trp1 his3-11 can1-100 GAL psi+ MET-CDC20::URA pURA::tetR-GFP::LEU2 ura3::tetOx112::URA3	Fernius, et al. 2009
MC213	HTB2::mCherry-URA3 his3 leu2 met15 LYS2	this study
MC230	fob1Δ::KanMX his3 leu2 HTB2::mCherry-URA met15 LYS2	this study
MC237	HTA2::GFP-HIS3 his3 leu2 ura3 met15 LYS2	Yeast GFP Collection
MC239	HTB2::GFP-HIS3 his3 leu2 ura3 met15 LYS2	Yeast GFP Collection
MC245	NUP49::GFP-HIS3 his3 leu2 ura3 met15 LYS2	Yeast GFP Collection
MC247	TUB1::GFP-HIS3 his3 leu2 ura3 met15 LYS2	Yeast GFP Collection
MC250	HTB2::mCherry-URA3 WHI5::GFP-HIS3 leu2 met15 LYS2	this study
MC255	spt21Δ::KanMX his3 leu2 HTB2::mCherry-URA met15 LYS2	Yeast KO Collection
MC257	hpc2Δ::KanMX his3 leu2 HTB2::mCherry-URA met15 LYS2	Yeast KO Collection
MC258	NUP49::GFP-HIS3 his3 leu2 ura3 met15 LYS2 htb2::mCherry-URA	this study
MC263	HTB2::mCherry-URA3 his3 leu2 met15 lys2 hpc2Δ::KanMX	this study
MC264	HTB2::mCherry-URA3 TUB1::GFP-HIS2 leu2 met15 LYS2	this study
MC266	HTB2::mCherry-URA3 his2 leu2 MET15 lys2 spt21Δ::KanMX	this study
MC281	HTB2::mCherry-URA3 his3-11::GFP-LacI-HIS3 RDN1::LacO(50)-ADE2	this study
MC349	mad3Δ::KanMX his3 leu2 htb2::mCherry-URA met15 LYS2	this study
MC351	SPC72::GFP-HIS3 his3 leu2 ura3 met15 LYS2	Yeast GFP Collection
MC352	UTP13::GFP-HIS3 his3 leu2 ura3 met15 LYS2	Yeast GFP Collection
MC354	CDC14::GFP-HIS3 his3 leu2 ura3 met15 LYS2	Yeast GFP Collection
MC355	SPC72::GFP-HIS3 his3 leu2 ura3 met15 LYS2 HTB2:mCherry-URA	this study
MC360	CDC14::GFP-HIS3 his3 leu2 ura3 met15 LYS2 HTB2:mCherry-URA	this study
MC364	UTP13::GFP-HIS3 his3 leu2 ura3 met15 LYS2 HTB2:mCherry-URA	this study
MC367	mad3Δ::KanMX his3 leu2 ura3 met15 LYS2 HTB2:mCherry-URA	this study
MC372	tom1Δ::KanMX his3 leu2 ura3 met15 LYS2 HTB2:mCherry-URA	this study
MC394	ura3 his3-11 pURA::tetR-GFP::LEU2 cenIV::tetOx448::URA3 HTB2:mCherry-URA	this study
MC395	leu2-3 his3-11 pURA::tetR-GFP::LEU2 ura3::tetOx112::URA3 HTB2:mCherry-URA	this study

817   **Table S1.** List of all strains and genotypes used in this study.

818 **Supplementary Figures**

819
Figure 1 – figure supplement 1. Histogram showing the duration of genomic missegregation
events in wild-type cells. Only events which eventually resulted in a RETRN event are shown,
as that provides a clear end of the missegregation event. **A)** Many GLMs are corrected within an
hour, but some events can last several hours. This includes GLMs from cells of all replicative
ages. **B)** When events are separated based on the age of the mother cell at time of missegregation
event, there is no significant difference in the distribution of GLM durations (p=0.5, two-tailed t-
test). N-value are the number of individual missegregation events quantified.
827

828
Figure 1 – figure supplement 2. GLMs are predictive of death, and most GLMs are
corrected by RETRN events. A) At the single cell level, genome level missegregation (GLM)
events are correlated with impending mortality (anti-correlated with remaining lifespan, as
shown), and become more predictive (more strongly anti-correlated, as shown) with increasing
replicative age. The strength of the anti-correlation with age is similar regardless of the
fluorophore used or histone protein tagged. Dots show correlation between an individual histone
and the remaining lifespan at that replicative age. To show trends, these have been smoothed
with a moving average (solid line). **B)** When a GLM occurs, approximately 90% of the time it is
corrected through a RETRN event, and the remainder of the time it becomes a terminal
missegregation. Error bars are 95% confidence intervals generated by bootstrapping with
replacement. N-values were individual cells, and only cells that died or senesced in the device
were used for both plots. **C)** RETRN events significantly increase replicative life span.
Comparison of experimentally observed replicative lifespan (corrected GLMs) to expected
lifespan if all GLMs were terminal (not corrected) for wild type cells. To generate the GLM not
corrected lifespan, the first GLM event for each mother cell was assumed to be terminal, and the
lifespan was truncated accordingly. This plot uses only cells that die in the device. RETRN
events increase median lifespan by 33% (p<0.0001). **D)** Same is in panel S2C, but including cells
that are censored. The increase in median lifespan by RETRN events is 29% (p<0.0001).
847

848
Figure 1 – figure supplement 3. Htb2 levels are predictive of DNA content at the single cell
level. Wild-type cells were grown in the microfluidic device for 16h, and then fixed. Cells had a
median replicative age of 13 divisions. The level of Htb2:mCherry is highly correlated
($p<0.0001$) with the level of DAPI staining at the single cell level in individual cells. This is true
across the entire range of Htb2:mCherry fluorescence, as even cells that have the highest
Htb2:mCherry levels have correspondingly high DAPI staining. N=800 individual mother cells.

855
Figure 1 – figure supplement 4. GLMs increase at end of life regardless of histone tagged
and fluorophore used. **A**) Both Htb2:mCherry and Hta2:GFP strains experience a dramatic
increase in the probability of GLM events in the 5-6 divisions preceding death. Error bars are
standard error. **B**) There is no difference between Htb2:mCherry and Hta2:GFP with respect to
the fraction of cells that experience GLM events. Error bars are 95% confidence intervals from
bootstrapping over individual cells. **C**) The distribution of the number of GLM events that cells
experience over their lifetime is similar between the strains but statistically different (p=0.02
two-tailed t-test), with Htb2:mCherry cells experiencing, on average, more events over their
lifetime. **D**) Replicative lifespan curves for Htb2:mCherry including censored cells (pink) and
excluding (red). **E**) Replicative lifespan curves for Hta2:GFP including censored cells (light
blue) and excluding (blue).

867
Figure 1 – figure supplement 5. GLMs are not caused by imaging conditions. A) To
determine whether cells were affected by the cumulative exposure to fluorescence excitation
energy, we compared GLM rates in cells imaged over their entire lifespans (blue) with those only
imaged after they were already aged for 20 hours (red), corresponding to a median replicative
age of 16 generations. This is equivalent to approximately 75% of the median lifespan of this
strain. To compare these cells to the control, we only quantified GLMs that occurred after 20h.
Error bars are 95% confidence intervals generated by bootstrapping with replacement over all
cells. No difference was detected in cells imaged continuously over their entire lifespan or only
after 20 hours, indicating that there is no cumulative effect of the exposure to fluorescence
excitation light on the frequency of GLMs or the ability of cells to correct these through RETRN
events.
879

Figure 2 – figure supplement 1. Single cell traces of the Whi5 and Myo1 data. A) Schematic
showing the temporal dynamics of the proteins used to characterize missegregation events. Whi5
exits the nucleus to initiate START and move the cell into S phase. It re-enters the nucleus at the
end of mitosis. **B)** A representative trace of a single cell expressing Htb2:mCherry and
Whi5:GFP. Histone levels (pink) increase, and then fall during mitosis. Arrows indicate the
timepoint before Whi5 transitions from the cytoplasm to the nucleus. (*) indicate raw Htb2
measurements, which were smoothed using a moving average for legibility (solid line). **C)**
Schematic showing the temporal dynamics of Myo1. Myo1 is produced at the end of G1 and
localizes to the bud neck until cytokinesis ends mitosis. **D)** Representative single cell trace of a
cell containing Htb2:mCherry and Myo1:GFP. (*) indicate raw Htb2 or Myo1 measurements,
which were smoothed using a moving average for legibility (solid line). Arrows indicate the
lowest point of Htb2 levels following mitosis.

893
Figure 2 – figure supplement 2. Removing Mad3 (mammalian BubR1) fails to affect the
age-related increase in missegregation rate. **A)** Wild-type and *mad3Δ* cells experience a
similar increase in genome level missegregation (GLM) events with age when looking at all
cells. **B)** The increase in GLM rate is similar when only comparing cells that die or senesce in
the device. **C)** When only cells that die in the device are aligned by death, both *mad3Δ* and wild-
type cells experience a similar age-related increase in GLM rates that begins around 5 divisions
prior to death. **D)** At the single cell level, GLM events are equally strongly correlated with
impending mortality in both *mad3Δ* and wild-type cells. Dots show correlation between a GLM
and the remaining lifespan at that replicative age for that genotype. To show trends, these have
been smoothed with a moving average for legibility (solid line). **E)** RLS curve including
censored cells (used in A). **F)** RLS curve excluding censored cells (used in B-D). All error bars
are standard error.

906
Figure 3 – figure supplement 1. GFP tagged histone data. Histones tagged with GFP increase
during aging, and are predictive of mortality. **A**) At the single cell level, histone levels are
strongly correlated with impending mortality, regardless of the histone tagged. Histones become
increasingly negatively correlated with remaining replicative lifespan as the cells age. Dots show
correlation between an individual histone and the remaining lifespan at that replicative age. To
show trends, these have been smoothed with a moving average (solid line). **B**) Mean histone
levels at each division aligned based on replicative age for each cell. **C**) Mean histone levels at
each division aligned using individual cell death. This shows that, on average, all histones begin
to increase sometime around 5-10 divisions prior to death. **D**) The replicative lifespans of the
strains including censored cells. **E**) The replicative lifespan of only cells that die or senesce in
the device. These are the cells that are used in the analysis for A-C. N values indicate the number
of cells, and the error bars are standard error.

919
Figure 3 – figure supplement 3. Fob1 vs Wild Type data. A) Mean Htb2 levels of mother
cells aligned by birth and excluding cells that were censored (See Figure 3B) **B)** Mean
missegregation rates of wild-type and *fob1Δ* mother cells aligned by birth and excluding cells
that were censored (See Figure 4A). **C)** At the single cell level, missegregation events are less
strongly anti-correlated with remaining lifespan in *fob1Δ* compared with wild-type. The
correlation curve for *fob1Δ* cells appears parallel to wild-type, but with an offset that is roughly
proportional to the increase in replicative lifespan we observed for the *fob1Δ* cells. Excludes
censored cells. Dots show correlation between an individual histone and the remaining lifespan
at that replicative age. To show trends, these have been smoothed with a moving average (solid
line). **D)** Mean Htb2 levels of mother cells aligned by death instead of birth, excluding censored
cells. **E)** Probability of missegregation events for wild-type and *fob1Δ* cells, but aligned by death
instead of birth and excluding censored cells. **F)** Survival curve of wild-type and *fob1Δ* cells
containing only cells which are not lost and censored. These are the cells used in A-E. Survival
curves comparing wild-type cells and cells lacking *FOB1*. **G)** Survival curve including censored
cells lost during the experiment. Compared with wild-type cells, *fob1Δ* mutants live longer
(p<1E-10, log-rank test). These are the cells used in the main figures. N-values report the number
of cells, and error bars for C,D,F are standard error.

937
Figure 4 – figure supplement 1. Cdc14 single cell traces in cells with terminal GLM and
RETRN events. A) Schematic showing cell cycle dynamics. Cdc14 is localized to the nucleolus
during the majority of the cell cycle and exits it in two stages during mitosis. Cdc14 re-enters the
nucleolus at the end of mitosis. **B)** A representative trace of a single cell expressing
Htb2:mCherry and Cdc14:GFP showing normal divisions and RETRN events. Histone levels
(pink) increase, and then fall during mitosis. Arrows indicate the timepoints where Cdc14 is at a
minima in the nucleolus. In normal divisions this coincides with the Htb2 minima. (*) indicate
raw Htb2 and Cdc14 measurements, which were smoothed using a moving average and
Savitzky-Golay smoothing respectively for legibility (solid line). In the last two divisions, a
RETRN events can be observed. **C)** Representative trace of a single cell expressing
Htb2:mCherry and Cdc14:GFP showing a terminal missegregations. Cdc14 exits the nucleolus
nearly eight hours after the GLM, but there is no RETRN event. **D)** Representative trace of a
single cell expressing Htb2:mCherry and Cdc14:GFP showing a GLM with a RETRN event (4h)
and then a terminal missegregation.

952
Figure 5 – figure supplement 1. Manipulating histone supply in yeast affects genome level
missegregation (GLM) rates. **A**) By manipulating the supply of histones through deleting
SPT21, *TOM1* and *HPC2* we can affect the probability of GLMs occurring. **B**) Mean GLM rates
for wild-type (wt), *spt21Δ, hpc2Δ* and *tom1Δ* mother cells aligned by birth and excluding cells
that were censored (See Figure 4B). **C**) Survival curves of only cells that are not lost. These are
the cells used for A,B. **D)** Survival curves which account for cells which are censored based on
loss from the traps, for wt, *hpc2Δ* and *spt21Δ* and *tom1Δ*. These are the cells used in Figure 5 of
the main text. Wild-type and *hpc2Δ* have different lifespans (p<0.001, log-rank test), wild-type
and *tom1Δ* have different lifespans (p<0.001, log-rank test), and wild-type and *spt21Δ* do not
have different lifespans (p=0.1 log-rank test). N-values indicate cell numbers, and error bars for
A,B are standard error.
964
965

Video Legends

Video 1 – Normal divisions followed by RETRN events in a strain expressing Htb2:mCherry and Myo1:GFP. The mother cell undergoes four normal divisions, and on the fifth (at timepoint 7h:35min), it experiences a missegregation event. The bud neck is clearly maintained until the retrograde transport occurs at 12h. Following this event, the bud neck is quickly removed, and is completely gone by 12h:15min. The blue arrow points to the mother cell during timepoints where it is experiencing a missegregation event. Timestamp is Hours:Min

Video 2 – RETRN events in a strain expressing Htb2:mCherry and Myo1:GFP. At timepoint 1h:20min the mother cell experiences a missegregation event. The bud neck is clearly maintained until the retrograde transport occurs at 2h:40min. Following this event, the bud neck is quickly removed, and is completely gone by 2h:55min. The blue arrow points to the mother cell during timepoints where it is experiencing a missegregation event. Video is significantly slower than other Videos. Timestamp is Hours:Min.

Video 3 - DNA colocalizes with tagged histones through mitosis. Cells expressing Htb2:mCherry were stained with Hoechst 3342, a live DNA stain. The first part of the video shows an overlay of red (Htb2:mCherry) and blue (Hoechst 3342), and the second shows the channels separated. As can be clearly seen in a normal cell cycle, the histones colocalize with the DNA, and both increase or decrease in fluorescence in the mother cell simultaneously.

Video 4 - DNA colocalizes with tagged histones through mitosis, including during missegregation and RETRN events. Cells expressing Htb2:mCherry were stained with Hoechst 3342, a live DNA stain. The first part of the video shows an overlay of red (Htb2:mCherry) and blue (Hoechst 3342), and the second shows the channels separated. As can be clearly seen, the histones colocalize with the DNA, and both increase or decrease in fluorescence in the mother cell simultaneously.

Video 5 – Normal divisions in a strain expressing Htb2:mCherry and Nup49:GFP. This cell undergoes six divisions, with histone and nuclear envelope behavior that is characteristic of young, healthy cells. Timestamp is Hours:Min

Video 6 – A missegregation followed by RETRN event in a strain expressing Htb2:mCherry and Nup49:GFP. The initial missegregation can be seen at timepoint 3h:30min, and the RETRN event at 8h:30min. Following the RETRN event, the mother cell is able to bud again at 12h, but the nuclear morphology of the daughter (for example, at 16h) is significantly altered. The blue arrow points to the mother cell during timepoints where it is experiencing a missegregation event. Timestamp is Hours:Min

Video 7 – Terminal missegregations in a strain expressing Htb2:mCherry and Nup49:GFP. At 3h, the mother cell can be seen to undergo a missegregation event. At 15h:30min, the daughter cell buds and can be seen to undergo mitosis, indicating that the daughter cell has separated from the mother. The blue arrow points to the mother cell during timepoints where it is experiencing a missegregation event. The mother cell eventually dies at 40h. Timestamp is Hours:Min.

Video 8 - A normal cell division, followed by a missegregation and RETRN event in a cell expressing Htb2:mCherry and Spc72:GFP. The two green dots indicate the spindle poles, and at numerous timepoints both poles enter the daughter cell. The blue arrow points to the mother cell during timepoints where it is experiencing a missegregation event. Timestamp is Hours:Min

Video 9 - A terminal missegregation event in a cell expressing Htb2:mCherry and Spc72:GFP. The two green dots indicate the spindle poles, and both poles enter the daughter around 2h40m. The poles move around and are highly active, with one at times reentering the mother cell. Finally, at 5h:20m, the daughter cell is washed away indicating it has fully separated from the

mother and that this is a terminal missegregation. The blue arrow points to the mother cell during
timepoints where it is experiencing a missegregation event. Timestamp is Hours:Min.
Video 10 – Normal divisions followed RETRN events in a strain expressing Htb2:mCherry and
Tub1:GFP. The mother cell undergoes four divisions normally and on the fifth, at timepoint
7h:50min it experiences a missegregation event that is resolved by a RETRN event at
10h:30min. The blue arrow points to the mother cell during timepoints where it is experiencing a
missegregation event. Timestamp is Hours:Min
Video 11 – A terminal divisions followed RETRN events in a strain expressing Htb2:mCherry
and Tub1:GFP. At timepoint 3h:15min the mother experiences a missegregation event, and both
the chromatin and microtubules can be seen entering the daughter cell. At 7h:25 min the
daughter cell is washed away indicating the cell has completed cytokinesis. The blue arrow
points to the mother cell during timepoints where it is experiencing a missegregation event.
Timestamp is Hours:Min
Video 12 – Normal divisions, and an increase in histone levels in a strain expressing
Htb2:mCherry and Nup49:GFP. Over a period of 15h this cell grows normally and produces
daughters. The increase in Htb2:mCherry can be clearly seen over the course of this experiment.
Video 13 - Age related structural and morphological changes to the nucleolus. This strain is
expressing a nucleolar localized protein (Utp13:GFP) and Htb2:mCherry. In the young cell, the
green nucleolus is adjacent to the red genomic material. As the cell ages and accumulates ERCs,
the histone level begins to increase noticeably around 14h. This alters the nucleolar morphology
and size, and by 23h the nucleolar morphology is dramatically different and it surrounds a large
portion of the histones. Timestamp is Hours:Min.
Video 14 - Age related structural and morphological changes to the nucleolus. This strain is
expressing a nucleolar localized protein (Utp13:GFP) and Htb2:mCherry. In the young cell, the
green nucleolus is adjacent to the red genomic material. As the cell ages and accumulates ERCs,
the histone level begins to increase noticeably around 9h. This alters the nucleolar morphology
and size, and by 12h the nucleolar morphology is dramatically different and it surrounds a large
portion of the histones. Timestamp is Hours:Min.
Video 15 - A missegregation and RETRN event showing that the Cdc14 remains localized to the
nucleolus even when the cell experiences a missegregation event. The cell is expressing
Htb2:mCherry and Cdc14:GFP. The exit of Cdc14:GFP from the nucleolus at 2h coincides with
the RETRN event. The blue arrow points to the mother cell during timepoints where it is
experiencing a missegregation event. Timestamp is Hours:Min
Video 16 - ChrXII dynamics during a missegregation and RETRN event. Cell is expressing
Htb2:mCherry and LacI:GFP, and has LacO repeats inserted into ChrXII. The Chromosome XII
sister chromatids clearly remain behind despite the majority of the genome entering the daughter
cell. Simultaneous to the RETRN event, the sister chromatids separate and can be identified in
both mother and daughter cells. The blue arrow points to the mother cell during timepoints where
it is experiencing a missegregation event. Timestamp is Hours:Min
Video 17 - ChrXII dynamics during a missegregation and RETRN event. Cell is expressing
Htb2:mCherry and LacI:GFP, and has LacO repeats inserted into ChrXII. The Chromosome XII
sister chromatids clearly remain behind despite the majority of the genome entering the daughter
cell. Simultaneous to the RETRN event, the sister chromatids separate and can be identified in
both mother and daughter cells. The blue arrow points to the mother cell during timepoints where
it is experiencing a missegregation event. Timestamp is Hours:Min

Video 18 - ChrIV dynamics during a missegregation and RETRN event. Cell is expressing
Htb2:mCherry and TetR:GFP, and has TetO repeats inserted into ChrIV. The Chromosome IV
sister chromatids clearly move into the daughter with the majority of the genome. Simultaneous
to the RETRN event, the sister chromatids separate and can be identified in both mother and
daughter cells. The blue arrow points to the mother cell during timepoints where it is
experiencing a missegregation event. Timestamp is Hours:Min
Video 19 - ChrV dynamics during a missegregation and RETRN event. Cell is expressing
Htb2:mCherry and TetR:GFP, and has TetO repeats inserted into ChrV. The Chromosome V
sister chromatids clearly move into the daughter with the majority of the genome. Simultaneous
to the RETRN event, the sister chromatids separate and can be identified in both mother and
daughter cells. The blue arrow points to the mother cell during timepoints where it is
experiencing a missegregation event. Timestamp is Hours:Min
Video 20- Cdc14 dynamics during normal cell divisions. Cell is expressing Htb2:mCherry and
Cdc14:GFP. Timestamp is Hours:Min
Video 21 - Spindle pole dynamics during normal cell divisions. Cell is expressing Htb2:mCherry
and Spc72:GFP. Timestamp is Hours:Min
Video 22 - Microtubule dynamics during normal cell divisions. Cell is expressing Htb2:mCherry
and Tub1:GFP. Timestamp is Hours:Min
Video 23- Bud neck dynamics during normal cell divisions. Cell is expressing Htb2:mCherry
and Myo1:GFP. Timestamp is Hours:Min
1077
1078