1 **Contextual interference effect is independent of retroactive inhibition but variable practice**

2 **is not always beneficial**

Benjamin Thürer[1,2]*, Sarah Gedemer[2], Anne Focke[2], Thorsten Stein[2]

1: Brain Signaling Group, Department of Molecular Medicine, Institute of Basic Medical

Sciences, University of Oslo, 0317 Oslo, Norway

2: BioMotion Center, Institute of Sports and Sports Science, Karlsruhe Institute of Technology,

76131 Karlsruhe, Germany

*benjamin.thurer@medisin.uio.no

Abstract

Positive effects of variable practice conditions on subsequent motor memory consolidation and

generalization are widely accepted and described as the contextual interference effect (CIE).

However, the general benefits of CIE are low and these benefits might even depend on decreased

retest performances in the blocked-practicing control group, caused by retroactive inhibition. The

aim of this study was to investigate if CIE represents a true learning phenomenon or possibly

reflects confounding effects of retroactive inhibition. We tested 48 healthy human participants

adapting their reaching movements to three different force field magnitudes. Subjects practiced the

force fields in either a Blocked (B), Random (R), or Constant (C) schedule. In addition, subjects of

the Blocked group performed either a retest schedule that did (Blocked-Matched; BM) or did not

(Blocked-Unmatched; BU) control for retroactive inhibition. Results showed that retroactive

inhibition did not affect the results of the BU group much and that the Random group showed a

better consolidation performance compared to both Blocked groups. However, compared to the

Constant group, the Random group showed only slight benefits in its memory consolidation of the

mean performance across all force field magnitudes. This indicates that CIE reflects a true motor

learning phenomenon, which is independent of retroactive inhibition. However, random practice is

not always beneficial over constant practice.

Keywords: Motor memory consolidation, Force field adaptation, Sensorimotor learning, Motor

adaptation, Retrograde inhibition,

Introduction

It is widely accepted that variable practice conditions can be beneficial for motor memory

consolidation (Schmidt, 1975; Shea & Morgan, 1979). In particular, the contextual interference

effect (CIE) – originally formulated by Battig (1972) for verbal learning – describes an increased

retest and transfer performance (Shea & Morgan, 1979; Magill & Hall, 1990) due to highly-

interfering cognitive processes during random practice conditions (Kantak et al., 2010; Lage et al.,

2015). This effect seems to be more robust in basic research than in applied settings (Brady 2004,

Brady 2008). In addition, CIE is commonly examined by comparing a random with a blocked

practice schedule and test for their corresponding effects on posttest and transfer test performance.

In such a random practice schedule, different tasks (or parameters) change randomly from trial to

trial, whereas, in a blocked practice schedule, one specific task (or parameter constellation) is

practiced as a whole block first, before switching to the next task (or parameter constellation).

Although CIE seems to be robust, there is no widely accepted hypothesis that accounts for this

effect. Classical explanations include the elaboration hypothesis (Shea & Zimny, 1983), the

reconstruction hypothesis (Lee & Magill, 1983), and the retroactive inhibition hypothesis (Shea &

Titzer, 1993). Thus, it is still unsolved whether CIE stems from an increased memory consolidation

due to the random practice condition (e.g. elaboration or reconstruction hypothesis) or by a

decreased retention performance of the blocked practice condition (retroactive inhibition

hypothesis). This latter assumption – which is in the focus of this paper – derives from the

observation that subsequent learning of different tasks can lead to inhibition of a previous memory,

an effect called retroactive inhibition (see Robertson et al., 2004 for a review). Concerning CIE,

retroactive inhibition might lead to disadvantages for the blocked practicing subjects since their

previous memory might be inhibited due to the blocked practice schedule. Therefore, these subjects

might show the worst performance when recalling the first task and the best performance when

recalling the last task they have practiced. Previous work showed possible confounding effects of

this retroactive inhibition on the motor retrieval after blocked practice and, therefore, questioned

the validity of CIE (Del Rey et al., 1994, Shewokis et al., 1998). Furthermore, when retroactive

inhibition was eliminated by using a reminder trial, no differences in memory recall were observed

between random and blocked practice schedules (Shea & Titzer, 1993). Retroactive inhibition

harms CIE when the last practice and the first retention block are not congruent and controlling for

this confounding effect would need a Latin Square design to test all possible task orders in the

practice and in the retention session. For instance, testing for CIE using three different tasks would

lead to 36 orders (6 practice x 6 retest orders) and, thus, to a complex study design which has never

been used in previous CIE studies.

So far, CIE and retroactive inhibition were discussed in the context of skill learning, in

which most CIE studies were conducted. Skill learning is commonly defined as a "set of processes

associated with practice or experience leading to relatively permanent changes in the capacity for

skilled movement" (Schmidt et al., 2018, p. 283). In contrast, motor adaptation is interpreted as a

different type of motor learning, in which the motor system responds to changes in environmental

conditions and/or changes in the body to regain the former capacity for a skilled movement under

these new conditions (Krakauer & Manzoni, 2011). This study focuses on motor adaptation using

a force field paradigm (Shadmehr & Mussa-Ivaldi, 1994), for which CIE has been demonstrated in

previous studies from our laboratory (Thürer et al., 2017; Thürer & Weber et al., 2018). In these

studies, subjects had to adapt their reaching movements to different force field magnitudes either

in a blocked or random fashion. However, these former studies did not control for retroactive

inhibition and a constant group, practicing only the force field magnitude that needed to be recalled,

was not included.

Therefore, the first purpose of this study is to examine the contextual interference effect in

force field adaptation under consideration of possible confounding effects of retroactive inhibition.

The second purpose of this study is to examine if variable practice schedules (blocked and random)

outperform a constant practice schedule even if subjects have to adapt their reaching movements

only to a single force field.

Methods

Participants

This study tested 48 healthy right-handed participants (24 ± 4 years; 10 women) with no previous

experience at a robotic manipulandum. Handedness was tested by the Edinburgh inventory

(Oldfield, 1971) and participant's vision was normal or corrected to normal. The study was

approved by the Institutional Review Board. All participants were informed about the protocol and

gave their written informed consent before participating in the study.

Apparatus and experimental task

The experimental task was implemented at a robotic manipulandum (Kinarm End-Point Lab, BKIN

Technologies, Kingston, Canada) which can produce forces via a handle towards the participants'

hands. In addition, we used a virtual reality display that allowed the participants to see the visual

information in the horizontal plane (Fig 1A). Please note that vision of handle, hand, and arm was

occluded by the virtual reality display. Positions and forces of the robot handle were sampled at

1000 Hz.

We will briefly describe the experimental task which can be found elsewhere in more detail

(Thürer et al., 2017). Participants were seated in front of the manipulandum and the virtual reality

display was calibrated to the robot's handle. All participants performed center-out reaching

movements with their dominant (right) hand. While performing this task, the horizontal display

shows a white cursor which is controlled via the handling of the manipulandum (Fig. 1B). Every

trial started by holding the cursor in the center target on the screen and a "go" signal was given by

the highlighting of a target. From that "go" signal on, participants were allowed to start their

reaching movement without any pressure of time (no fast reaction times required). When

participants reached the target position, subjects were actively moved backwards to the center

position by the manipulandum. After a short pause in the center position of 800 ms, the next target

highlighted in a pseudo-randomized order. In total, six target positions were defined building a

circle with a diameter of 20 cm surrounding the center point. Pseudo-randomization facilitated that

in every block of six trials every target highlighted just once and that every participant had a

different target order so that no influence of target direction was given on the group level.

To provide similar movement times across trials and subjects, visual feedback was

implemented in every single trial. The feedback was given via a change in the target color after

reaching it. Target color switched to red if the movement was too fast (< 450 ms), blue if it was

too slow (> 550 ms), and green otherwise.

To induce motor adaptation and subsequent memory consolidation, we implemented

velocity-dependent counter-clockwise directed force fields at the robotic manipulandum. These

force fields perturbed the participants' movements and typically degraded their initial motor

performance leading to curved hand trajectories (Fig. 1C). In order to investigate practice schedules

with different amounts of variabilities, three separate force field viscosities were implemented with

each viscosity inducing a force field magnitude of either 8, 15, or 22 Ns/m. Therefore, each force

field magnitude represented an object with different physical properties. The absolute maximum

perpendicular displacement between the participant's hand path and a direct line joining center

point and target quantified the motor error (Fig. 1C).

Figure 1: Apparatus and task. **A**: Robotic manipulandum with the virtual reality display. Subjects

hold the robotic handle but look into the virtual reality add-on in front of them. In addition, a fabric,

not shown here, was attached to the reality add-on and to the subjects' shoulders to prevented

further visual input of the arms. The virtual cursor is controlled via the robotic handle. **B**:

Experimental task at the robotic manipulandum. The subjects see the cursor and targets on the

screen in the horizontal plane. **C**: the maximum perpendicular displacement (PDmax) was used to

quantify the motor performance. Dashed arrows indicate the force field. **D**: Experimental protocol

over two consecutive days. Red colors indicate that both Blocked groups differed in their practice

schedule on day 2 whereas the Random and Constant groups differed in their practice schedules on

day 1. Triangles, rectangles, and stars symbolize different force field magnitudes.

Experimental procedure

Participants were equally distributed into 4 groups (Blocked-Matched, BM; Blocked-Unmatched,

BU; Constant, C; Random, R; each $n = 12$). The groups differed only in their task protocol during

Practice and during Retest. The study took place on two consecutive days with 24 h between the

two test sessions (Fig. 1D).

On day 1, all participants received instructions about the behavioral task and performed 144

familiarization trials under null field conditions (motors of the robot were turned off) with two

breaks of 30 s after every 48th trials. Then, participants performed a baseline measurement

consisting 30 null field trials. After that, all participants performed 540 force field trials during

Practice, with a different force field schedule according to their group allocation. To avoid fatigue,

participants had a 30 s break after each 60th trial. The participants performed all trials on day 1

with their dominant right hand.

The practice schedule was identical between the two Blocked groups (BM, BU) but

different for the Random and Constant groups. Participants of the Blocked groups performed the

three force field magnitudes (8, 15, 22 Ns/m) in a blocked order. Therefore, all trials of one specific

magnitude were practiced first, before switching to the next magnitude. This resulted in three

blocks, each containing 180 trials of one specific force field magnitude. The Random group

performed a highly-variable practice schedule so that the three force field magnitudes changed on

a single-trial level. For the Constant group, each participant practiced only one specific force field

magnitude (15 Ns/m) and, thus, encountered no force field variability at all. The force field

magnitude (for C) and the magnitude order (for BM, BU, and R) was counter-balanced across

participants so that the mean force field magnitude was 15 Ns/m on the group level. In addition,

for the Blocked and Random groups, the mean force field magnitude across the whole Practice

session was 15 Ns/m for each single participant.

164   On day 2, all participants performed a Posttest and Transfer test. To quantify Posttest

165   performance, all participants performed 18 force field trials divided into three blocks with each

166   block representing one force field magnitude. Then, participants performed 60 trials of a constant

167   force field magnitude with their non-dominant left hand (Transfer test) to investigate long-term

168   effects on the contralateral hand indicated by a previous study from our group (Thürer & Weber et

169   al., 2018).

170   The order of force field magnitudes on day 2 differed between groups. For the Blocked-

Matched group, the magnitudes in Posttest were in a reversed order compared to Practice and, thus,

"matched" in terms of a reduced effect of retroactive inhibition on the first block (Fig. 1D). For

instance, when for a specific participant the Practice order was 15, 8, 22 Ns/m, the Posttest order

was set to 22, 8, 15 Ns/m. For the Blocked-Unmatched group, however, the order of force field

magnitudes was the same for the Practice and the Posttest session. For instance, when for a specific

participant the Practice order was 15, 8, 22 Ns/m, the Posttest order was set to 15, 8, 22 Ns/m. The

order of magnitudes on day 2 was similar for the Random and for the Constant group. Both groups

started the Posttest with the force field magnitude (for C) or with the mean force field magnitude

(for R) of the Practice session. Accordingly, all participants of the Random group were tested on

15 Ns/m in the first Posttest block because it was shown that participants adapt to the mean force

field magnitude (Scheidt et al., 2001). Regarding the Constant group, the first block's magnitude

was different between participants for each single participant had a different magnitude during

Practice due to counterbalancing. Both groups (R, C) were counterbalanced for the order of the

remaining two force field magnitudes so that, still, the mean across groups for the second and third

block of the Posttest was at 15 Ns/m. According to the Posttest, all participants performed the

Transfer test on the left hand at a specific constant force field magnitude, which was the same as

the first magnitude in the Posttest (Fig. 1D).

Statistics

For the statistical analyses, mean performance for the first and the last 6 trials of the Practice session

(Practice FT, Practice LT) was computed. Posttest performance was computed by the mean of the

first, middle, and last 6 Posttest trials (Posttest FT, MT, LT) and the mean across all 18 Posttest

trials (Posttest ALL). Contralateral Transfer performance was quantified by the initial 6 Transfer

trials (Transfer FT) and the whole 60 Transfer trials (Transfer ALL).

To test for the possible influence of retroactive inhibition on CIE, we performed mixed-

model ANOVAs with the factors time (Practice LT, Posttest FT; Practice LT, Posttest ALL) and

group (BM, BU). For a possible effect on the generalization from one hand to the other, the factor

time was adjusted accordingly (Practice LT, Transfer FT; Practice LT, Transfer ALL). In addition,

we investigated if random practice even outperforms constant practice by using standard Fischer t-

tests between groups (R, C). We calculated differences between each force field magnitude of the

Posttest (Posttest FT, MT, LT, ALL) and the last trials of the Practice session (Practice LT)

accordingly.

It is widely accepted that p-values alone are not a good marker for potential results in

research (e.g. Nuzzo, 2014). Therefore, besides using classical inferential statistics, we provide

effect sizes (partial eta squared, $pEta^2$; Cohen's d, d) and additional Bayesian statistics. Bayesian

statistics are provided in the supplementary material and confirm the results and interpretations of

the classical inferential statistics.

All parameters were tested for normal distribution and homogeneity of variances using

Shapiro-Wilk and Levene test. Statistical analyses were performed using MATLAB R2015b

(Mathworks, Natick, USA) and JASP 0.8.6 (Team JASP). Threshold for statistical significance

was set to $p = 0.05$. Multiple comparisons were corrected by the False Discovery Rate (FDR,

Benjamini & Hochberg, 1995) and in case of multiple comparisons, p-values in this study represent

the FDR corrected p-value (Benjamini & Yekutieli, 2001).

Results

CIE is unaffected by the different retest schedules of the Blocked groups

The progress in motor performance for both Blocked groups is depicted in Fig. 2. First, we tested

if motor adaptation during Practice differed between groups. Both groups (BM, BU) adapted

successfully to the force field schedule during Practice (time: $F(1,22) = 104.09$, $p < .001$, $pEta^2 =$

0.83) and showed no differences in their adaptation (group: $F(1,22) = 0.79$, $p = .385$, $pEta^2 = 0.03$;

time*group: $F(1,22) = 1.67$, $p = .210$, $pEta^2 = 0.07$, for mixed-model ANOVAs with factors time

(Practice FT, Practice LT) and group (BM, BU)).

Consolidation of motor memory (from Practice to Posttest) did not differ between Blocked

groups regarding their recall of the first force field magnitude ($F(1,22) = 0.47$, $p = .498$, $pEta^2 =$

0.02) or regarding all force field magnitudes ($F(1,22) = 0.06$, $p = .808$, $pEta^2 < 0.01$, for uncorrected

time*group interactions with factors time (Practice LT, Posttest FT; Practice LT, Posttest ALL)

and group (BM, BU)). Although descriptive statistics indicate slight benefits for the BM group in

recalling the very first force field magnitude during Posttest (Fig 2A, Fig 3A), this is not supported

by additional post-hoc statistics ($t(22) = -0.94$, $p = .358$, $d = -0.38$, for uncorrected independent t-

test between groups' Posttest FT performance). However, memory consolidation was significantly

stronger for the Random group compared to the BM group ($F(1,22) = 5.65$, $p = .029$, $pEta^2 = 0.20$)

and descriptively stronger to the BU group ($F(1,22) = 5.49$, $p = .054$, $pEta^2 = 0.20$, for FDR

corrected time*group interactions with factors time (Practice LT, Posttest ALL) and group (BM,

R; BU, R)), which confirms the contextual interference effect.

We further investigated if retroactive inhibition affected the generalization from the

dominant hand (Practice) to the non-dominant (Transfer). No differences between Blocked groups

($F(1,22) = 0.15$, $p = .701$, $pEta^2 < 0.01$, for time*group interaction with factors time (Practice LT,

Transfer FT) and group (BM, BU)) were observed and the Random group performed similar to

both Blocked groups (BM vs. R: $F(1,22) = 1.18$, $p = .568$, $pEta^2 = 0.05$; BU vs. R: $F(1,22) = 0.34$,

$p = .580$, $pEta^2 = 0.02$ for FDR corrected time*group interactions with factors time (Practice LT,

Transfer FT) and group (BM, R; BU, R)). However, further adaptation (quantified by Transfer LT

- Transfer FT) was observed to be faster for the Random group (Fig 3B), with this effect becoming

more pronounced if adaptation during Practice (Practice LT - Practice FT) is compared with

adaptation during Transfer (BM vs. R: $F(1,22) = 3.39$, $p = .079$, $pEta^2 = 0.13$; BU vs. R: $F(1,22) =$

8.05, $p = .020$, $pEta^2 = 0.27$, for FDR corrected time*group interaction with factors time (Practice

LT − Practice FT, Transfer LT − Transfer FT) and group (BM, R; BU, R)). This indicates that,

although motor memory generalization from Practice to Transfer did not differ between groups,

adaptation within the Transfer test appears to be faster for the Random compared to both Blocked

groups.

Figure 2: Descriptive results. **A**: Progress of the mean motor error + SEM for Practice, Posttest, and Transfer test of the Blocked-Matched (BM) and the Blocked-Unmatched (BU) group. **B**: Progress of the mean motor error for the Random (R) and the Constant (C) group.

Random practice improves mean memory consolidation of multiple force field magnitudes

The second aim of this study was to examine if constant practice leads to better memory consolidation of only one force field magnitude than random practice and if random practice outperforms constant practice in the recall of multiple force field magnitudes. Our results show that memory consolidation of the Constant and of the Random group did not differ for each single force field magnitude (quantified by differences between Posttest FT/MT/LT and Practice LT. Posttest FT: $t(22) = 0.89$, $p = .386$, $d = 0.36$; Posttest MT: $U = 99$, $p = .171$; Posttest LT $t(22) = 1.97$, $p = .122$, $d = 0.80$, for FDR corrected t- and U-tests between groups (C, R)). However, the Random

group showed a better mean memory consolidation across all force field magnitudes (Posttest ALL:

$t(22) = 3.23$, $p = .016$, $d = 1.32$, FDR corrected), with this effect most pronounced predicting high

effect sizes using Bayesian statistics (see Supplementary Figure S2). It is important to mention here

that this consolidation effect occurred due to performance differences at the end of Practice, for

there is no group difference regarding absolute Posttest values (Posttest FT: $U = 46$, $p = .143$, $d =$

-0.36; Posttest ALL: $t(22) = -0.41$, $p = .684$, $d = -0.17$, for t- and U-tests between groups (C, R)).

Further, we were interested in the mechanisms during Practice, which facilitate Posttest

performance. Uncorrected correlation analyses indicated that Posttest performances of all groups

were associated with at least two different mechanisms, namely practice success ($r = 0.56$, $p <$

.001) and practice variability ($r = 0.58$, $p < .001$, for Pearson correlation of all participants (BM,

BU, C, R) between Practice LT and Posttest ALL and between the individual's standard deviation

of the whole practice session and Posttest ALL). However, since the Random group revealed a

lower practice success ($t(22) = -2.89$, $p = .009$, $d = -1.18$) and a higher practice variability ($t(22) =$

-4.22, $p < .001$, $d = -1.72$, for t-tests between groups (R, C)) compared to the Constant group,

correlation coefficients might be compromised due to the inclusion of the Random group (Fig. 3C).

A deeper investigation of the data shows that Pearson coefficients of the Random group were

indeed higher compared to coefficients of the other groups but differed significantly only for the

practice variability (practice success: $z = -1.28$, $p = .201$; practice variability: $z = -2.63$, $p = .018$,

for the uncorrected differences between groups (R, [BM BU C]) after r-to-z transformation). This

indicates that variability during Practice was more beneficial - in terms of an improved motor

memory consolidation - for the Random group than for the other groups.

Figure 3: Deeper investigation of the behavioral results. **A**: Overview of the mean motor error +

SEM computed across trials and subjects for each group over the whole experiment. **B**: Differences

for Practice LT - Practice FT and Transfer LT - Transfer FT as an indication for the range of motor

adaptation. **C**: Correlation analysis for practice success (motor error of Practice LT; left) and

practice variability (standard deviation of the whole practice session; right) each associated with

Posttest ALL. Lines indicate linear fits for the Random (blue) and all the other groups (black).

Discussion

Our results showed no differences between the two Blocked groups although the Posttest schedule

of one group (BM) did and the other schedule (BU) did not control for retroactive inhibition.

Compared to the Random group both Blocked groups showed a limited memory consolidation,

which depicts that retroactive inhibition does not account for CIE in motor adaptation tasks.

Comparisons between Random and Constant groups showed a similar memory consolidation for

each single force field magnitude. However, the Random group outperformed the Constant group

in its mean memory consolidation across all three force field magnitudes.

Retroactive inhibition does not affect the contextual interference effect in motor adaptation

The experimental procedure of the BM group controlled for possible confounding effects of

retroactive inhibition. Nevertheless, BM performed similar to BU and its memory consolidation

was hampered compared to the Random group. These findings contradict previous skill learning

studies (Shea & Titzer, 1993, Del Rey et al., 1994, Shewokis et al., 1998), assuming retroactive

inhibition as the underlying mechanism for the contextual interference effect. Although retroactive

inhibition seemed to decrease the Posttest performance in the BU group (Fig. 3A), this effect was

too small to explain the benefits after random practice.

These benefits for the Random group were also observed when testing for the generalization

of memory to the contralateral hand. This finding concurs with the literature, which frequently

showed CIE for transfer tests in skill learning tasks (e.g. Shea & Morgan, 1979, Brady, 2004,

Wright et al., 2015) and reproduces earlier findings from our lab using a motor adaptation task

(Thürer & Weber et al. 2018). Our results indicate that participants of the Random group were not

only able to consolidate better in the meantime between sessions, leading to similar initial

performances in the Post- and Transfer test, they also were able to adapt faster towards the force

field condition with their left hand. This positive effect of variability on subsequent motor

adaptation is in line with a previous study, demonstrating that participants revealing a highly

variable baseline period adapt faster during the practice period (Wu et al., 2014). It is assumed that

this positive effect occurred due to noise in the motor planning system but not due to noise in the

motor execution system (Dhawale et al., 2017). That leads to the suggestion that the nervous

system, at least in some way, uses the knowledge of uncertainty of measured and/or predicted

feedback (Izawa & Shadmehr, 2008) to improve motor adaptation (Wei & Körding, 2010).

However, future work is needed to investigate this more deeply.

Correlation results are in accordance with the suggested role of motor variability. We were

able to show that both, motor success and motor variability are positively associated with the

Posttest performance. This indicates that positive effects of random practice depend more on the

increased motor variability and less on motor success, with the inverse proportion taking place for

the other groups. From a practical perspective, it is not clear if a perfect tradeoff between motor

success and motor variability would lead to absolute better performances after random compared

to blocked or constant practice. However, in a previous study we used a random design with lower

inter-trial variabilities and were able to show better absolute performance values for Random

compared to Blocked groups throughout the whole transfer test (Thürer & Weber et al., 2018). This

indicates that a good tradeoff is beneficial for long-term practice outcomes.

Contextual interference improves only the mean memory consolidation of multiple force field

magnitudes

Although CIE reflects a widely accepted phenomenon and seems to be unaffected by retroactive

inhibition in motor adaptation tasks, it is not clear whether random practice is always better than

constant practice. Our results showed that consolidation benefits of random compared to constant

practice in motor adaptation occur only if multiple force field magnitudes are retested. This

indicates that memory consolidation of a single task might not be improved by a highly variable

practice schedule. This concurs with the especial skill effect for skill learning (Breslin et al. 2010)

but contradicts previous work regarding random practice (Shea & Kohl, 1991).

This finding is also in line with our observation of a positive association between motor

success and Posttest performance. Our results indicate that benefits due to higher practice success

in constant practice equals the benefits of practice variability in random practice. However, since

participants of the Constant group did only train a single force field magnitude, mean consolidation

of multiple force field magnitudes was hampered compared to the Random group.

It is important to note that absolute values of Posttest performance did not differ

significantly between groups. Nevertheless, derived from a practical perspective, Random practice

might be the better choice since it leads to similar results than constant practice but has the

opportunity to increase mean memory consolidation of multiple force field magnitudes and to

increase generalization in terms of faster re-adaptation on the contralateral hand (Fig. 3B).

Limitations

This study showed some minor limitations which we would like to address. The Constant group

trained the same amount of trials as the other groups but each subject of only one force field

magnitude. Therefore, this group was able to draw on a greater practice experience for one specific

magnitude compared to the other groups. We cannot state how much this affected the results but

from a practical perspective it was important to have the same amount of practice time for each

group.

The force field magnitudes might have been too different and, thus, induced a too high

practice variability in the Random group. This might be the reason why we were not able to show

absolute Posttest and Transfer test performance benefits for the Random group. In a previous study

with a lower amount of variability, we were able to show these absolute benefits after Random

practice in the transfer test on the contralateral hand (Thürer &Weber et al., 2018).

The order of Post- and Transfer tests was not counter-balanced. Therefore, similar group

performances in the first Transfer trials might be caused by the 18 Posttest trials. However, we

were previously able to show that contralateral transfer from the dominant to the non-dominant

hand after random practice is almost independent of the Posttest performance (Thürer & Weber et

al., 2018) and, therefore, suggest that this had only minor effects on our results.

In this study, we investigated motor adaptation and not skill acquisition and, therefore, our

interpretations cannot be generalized to skill learning tasks. However, from a theoretical point of

view, confounding effects of retroactive inhibition should be more prone to happen in motor

adaptation than in skill acquisition, due to a bigger potential overlap of the underlying neural

structures.

Conclusion

In this study, we were able to show that the contextual interference effect represents a valid learning

phenomenon that is not affected by retroactive inhibition. Furthermore, we were able to show that

benefits of random practice are more related to the memory consolidation of multiple tasks or task

parameters and not on the consolidation of only one single task. The higher variability during

practice seems to be beneficial for this effect whereas motor success seems to facilitate the benefits

of constant practice. However, it remains unsolved how the motor system uses variability to

improve subsequent motor memory consolidation

Funding

This work was supported by the Graduate Funding from the German States. We acknowledge

support by Deutsche Forschungsgemeinschaft and Open Access Publishing Fund of Karlsruhe

Institute of Technology.

Acknowledgment

We like to thank Alexander Wolpert for his technical assistance and Ernst Hossner for the fruitful

discussions and his valuable input.

Literature

Battig WF (1972). Intratask interference as a source of facilitation in transfer and retention. In: *Topics in learning and performance* (Thompson RF and Voss JF, eds), pp131-159. New York: Academic Press.

Benjamini Y, Hochberg Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B* 57, 289-300.

Benjamini Y, Yekutieli D (2001). The control of the false discovery rate in multiple testing under dependency. *Ann Stat* 29, 1165-1188.

Brady F (2004). Contextual interference: a meta-analytic study. *Percept Mot Skills* 99(1), 116-126.

Brady, F. (2008). The contextual interference effect and sport skills. *Perceptual and Motor Skills*, 106(2), 461–472.

Brashers-Krug T, Shadmehr R, Bizzi E (1996). Consolidation in human motor memory. *Nature 382*(6588), 252.

Breslin G, Hodges NJ, Kennedy R, Hanlon M, Williams AM (2010). An especial skill: Support for a learned parameters hypothesis. *Acta Psychologica* 134(1), 55–60. https://doi.org/10.1016/J.ACTPSY.2009.12.004

Cohen J (1988). *Statistical power analysis for the behavioral sciences*. New York: Academic Press.

Del Rey P, Liu X, Simpson KJ (1994). Does retroactive inhibition influence contextual interference effects?. *Res Q Exerc Sport*, 65(2), 120-126.

Dhawale AK, Smith MA, Ölveczky BP (2017). The role of variability in motor learning. *Annu Rev Neurosci* 40, 479-98.

Izawa J, Shadmehr R (2008). On-line processing of uncertain information in visuomotor control. *J Neurosci* 28, 11360.

JASP Team (2018). JASP (Version 0.8.6) [Computer software].

Kantak SS, Sullivan KJ, Fisher BE, Knowlton BJ, Winstein CJ (2010). Neural substrates of motor

memory consolidation depend on practice structure. *Nat Neurosci 13*, 923-925.

Krakauer JW, Mazzoni P (2011). Human sensorimotor learning: adaptation, skill, and beyond.

Curr Opin Neurobiol 21: 636–644.

Lage GM, Ugrinowitsch H, Apolinário-Souza T, Vieira MM, Albuquerque MR, Benda RN (2015).

Repetition and variation in motor practice: A review of neural correlates. *Neurosci*

Biobehav Rev, 57, 132–141. https://doi.org/10.1016/j.neubiorev.2015.08.012

Lee TD, Magill RA (1983). The Locus of Contextual Interference in Motor-Skill Acquisition. *J*

Exp Psychol Learn Mem Cogn 9, 730-746.

Magill RA, Hall KG (1990). A review of the contextual interference effect in motor skill

acquisition. *Hum Mov Sci 9*, 241-289.

Nuzzo R (2014). Scientific method: statistical errors. *Nature* 506(7487), 150.

Oldfield RC (1971). The assessment and analysis of handedness: the Edinburgh inventory.

Neuropsychologia 9(1), 97-113.

Robertson EM, Pascual-Leone A, Miall RC (2004). Current concepts in procedural consolidation.

Nat Rev Neurosci 5(7), 576-582.

Rouder JN, Morey RD, Speckman PL, Province JM (2012). Default Bayes factors for ANOVA

designs. *J Math Psychol* 56(5), 356–374. https://doi.org/10.1016/j.jmp.2012.08.001

Scheidt RA, Dingwell JB, Mussa-Ivaldi FA (2001). Learning to move amid uncertainty. *J*

Neurophysiol 86, 971-985.

Schmidt, R. A. (1975). A schema theory of discrete motor skill learning. *Psychological Review*,

82, 225–260

Schmidt RA, Lee TD (2011). *Motor control and learning: A behavioral emphasis*. Champaign, IL:

Human Kinetics.

Schmidt, R.A., Lee, T.D., Winstein, C.J., Wulf, G. & Zelaznik, H.N. (2018). *Motor Control and*

Learning: A Behavioral Emphasis (6th ed.). Champaign, IL: Human Kinetics.

Shea CH, Kohl RM (1991). Composition of practice: Influence on the retention of motor skills.

Res Q Exerc Sport 62(2), 187-195.

Shea JB, Morgan RL (1979). Contextual interference effects on the acquisition, retention, and

transfer of a motor skill. *J Exp Psychol Hum Learn 5*, 179-187.

Shea JB, Titzer RC (1993). The influence of reminder trials on contextual interference effects. *J*

Mot Behav 25(4), 264-274.

Shea JB, Zimny ST (1983). Context effects in memory and learning movement information. In:

Memory and control of action (Magill RA, ed.) pp. 345–366. North-Holland.

Shewokis PA, Del Rey P, Simpson KJ (1998). A test of retroactive inhibition as an explanation of

contextual interference. *Res Q Exerc Sport 69*(1), 70-74.

Thürer B, Stockinger C, Focke A, Putze F, Schultz T, Stein T (2016). Increased gamma band power

during movement planning coincides with motor memory retrieval. *NeuroImage 125*, 172-

181.

Thürer B, Stockinger C, Putze F, Schultz T, Stein T (2017). Mechanisms within the Parietal Cortex

Correlate with the Benefits of Random Practice in Motor Adaptation. *Front Hum Neurosci*

11, 1–11. https://doi.org/10.3389/fnhum.2017.00403

Thürer B, Weber FD, Born J, Stein T (2018). Variable training but not sleep improves consolidation

of motor adaptation. *Sci Rep* 8(1), 15977. https://doi.org/10.1038/s41598-018-34225-w

Wei K, Körding K (2010). Uncertainty of feedback and state estimation determines the speed of

motor adaptation. *Front Comp Neurosci* 4, 11. https://doi.org/10.3389/fncom.2010.00011

Wolpert DM, Ghahramani Z, Jordan MI (1995). An internal model for sensorimotor integration.

Science 269(5232), 1880-1882.

Wright D, Verwey W, Buchanen J, Chen J, Rhee J, Immink M (2015). Consolidating behavioral

and neurophysiologic findings to explain the influence of contextual interference during

motor sequence learning. *Psychon Bull Rev* 1-21.

Wu HG, Miyamoto YR, Gonzalez Castro LN, Ölveczky BP, Smith MA (2014). Temporal structure

of motor variability is dynamically regulated and predicts motor learning ability. *Nat*

Neurosci 17, 312-21.