Meiotic MCM proteins promote and inhibit crossovers during meiotic recombination

Michaelyn Hartmann[*], Kathryn P. Kohl[†], Jeff Sekelsky[*,‡,§], Talia Hatkevich[*,**]

[*]*Curriculum in Genetics and Molecular Biology, University of North Carolina, Chapel Hill,*

North Carolina 27599

[†]*Department of Biology, Winthrop University, 701 Oakland Avenue, Rock Hill, SC 29733*

[‡]*Department of Biology, University of North Carolina, Chapel Hill, North Carolina 27599*

[§]*Integrative Program in Biological and Genome Sciences, University of North Carolina, Chapel*

Hill, North Carolina 27599

[**]*Corresponding author*

Running title: Genetic analysis of *Drosophila* mei-MCMs

Five key words or phrases: *Drosophila*, meiotic recombination, meiosis, crossover, mei-MCM

Corresponding author*:* CB #3280, 303 Fordham Hall, Curriculum of Genetics and Molecular

Biology, University of North Carolina, Chapel Hill, NC 27599-3280.

E-mail: hatkevic@email.unc.edu

Abstract

Crossover formation as a result of meiotic recombination is vital for proper segregation

of homologous chromosomes at the end of meiosis I. In many organisms, crossovers are

generated through two crossover pathways: Class I and Class II. To ensure accurate crossover

formation, meiosis-specific protein complexes regulate the degree in which each pathway is

used. One such complex is the mei-MCM complex, which contains MCM (mini-chromosome

maintenance) and MCM-like proteins REC (ortholog of Mcm8), MEI-217, and MEI-218,

collectively called the mei-MCM complex. The mei-MCM complex genetically promotes Class I

crossovers and inhibits Class II crossovers in *Drosophila*, but it is unclear how individual mei-

MCM proteins contribute to crossover regulation. In this study, we perform genetic analyses to

understand how specific regions and motifs of mei-MCM proteins contribute to Class I and II

crossover formation and distribution. Our analyses show that the long, disordered N-terminus of

MEI-218 is dispensable for crossover formation, and that mutations that disrupt REC's Walker A

and B motifs differentially affect Class I and Class II crossover formation. In Rec Walker A

mutants, Class I crossovers exhibit no change, but Class II crossovers are increased. However, in

rec Walker B mutants, Class I crossovers are severely impaired, and Class II crossovers are

increased. These results suggest that REC may form multiple complexes that exhibit differential

REC-dependent ATP binding and hydrolyzing requirements. These results provide genetic

insight into the mechanisms through which mei-MCM proteins promote Class I crossovers and

inhibit Class II crossovers.

Introduction

To reestablish the diploid genome upon sexual fertilization, the genome of progenitor germ cells must be successfully reduced by half through meiosis. Accurate reduction of the genome at the end of meiosis I requires crossover formation between homologous chromosomes during meiotic recombination. Meiotic recombination is initiated by the formation of multiple double-strand breaks (DSBs); the majority of meiotic DSBs are repaired as noncrossovers, while a selected subset are repaired as crossovers between homologs (reviewed in Lake and Hawley 2012).

Two distinct types of meiotic crossovers have been described: Class I and Class II. First defined in budding yeast (de Los Santos *et al.* 2003), Class I and Class II crossovers exist in most sexually reproducing organisms, but the relative proportions of each crossover type vary among organisms (Hollingsworth and Brill 2004). In *Drosophila*, most – if not all – crossovers are generated through the Class I pathway (Hatkevich *et al.* 2017), as shown through their dependence on the putative catalytic unit of the Class I meiotic resolvase MEI-9 (Sekelsky *et al.* 1995; Yıldız *et al.* 2002) and their display of crossover interference (Hatkevich *et al.* 2017). Most crossovers in *Drosophila* are also dependent upon a group of MCM- or MCM-like proteins, called the mei-MCM complex (Baker and Carpenter 1972; Grell 1978; Liu *et al.* 2000; Kohl *et al.* 2012).

The mei-MCM complex consists of REC (the *Drosophila* ortholog of MCM8), MEI-217, and MEI-218. REC appears to be a bona fide MCM protein, based on conservation of both the N-terminal MCM domain and the C-terminal AAA+ ATPase domain, which includes Walker A and B boxes that bind and hydrolyze ATP (Figure 1A). In contrast, MEI-217 and MEI-218 are highly divergent MCM-like proteins, and together resemble one full MCM protein. MEI-217 is

structurally similar to the MCM N-terminal domain, though this similarity is not detected in

BLAST or conserved domain searches (Kohl *et al.* 2012). The carboxy-terminus of MEI-218 has

a domain related to the AAA+ ATPase domain, but key residues are not conserved, including the

Walker A and B motifs that are critical for binding and hydrolyzing ATP, respectively (Iyer *et al.*

2004) (Figure 1B). Because key residues in the Walker A and B motifs are not conversed,

MEI-218 may not exhibit ATPase activity or it may exhibit partial function. In addition,

MEI-218 has a long N-terminal extension that is poorly conserved and predicted to be

disordered. The function of this region is unknown, but gene swap studies suggest that it may

contribute to differences in the recombination landscape among *Drosophila* species (Brand *et al.*

2018). For further analysis and details regarding the evolution of the mei-MCM complex, see

Supplemental Figures S1-S3.

While most crossovers are generated through the Class I pathway in wild-type

Drosophila and are mei-MCM dependent, mutants that lack the Bloom syndrome helicase (Blm)

generate only Class II crossovers based on their independence of MEI-9 and lack of patterning

(*e.g.*, interference) that is associated with Class I crossovers (Hatkevich *et al.* 2017). Blm is an

ATP-dependent 3'-5' helicase that exhibits vital anti-crossover functions in both meiotic and

somatic DSB repair (reviewed in Hatkevich and Sekelsky 2017). Interestingly, mutations in *mei-*

MCM and *Blm* genes genetically interact. In *Blm* mutants, crossovers are reduced by 30% but in

a *Blm rec* double mutant, crossovers are significantly increased compared to wild-type (Kohl *et*

al. 2012). This suggests that the mei-MCMs may function to inhibit crossovers within the Class

II pathway, in addition to their role promoting crossovers in the Class I pathway.

While the mei-MCMs function as a complex, little is known about how individual

mei-MCMs contribute to Class I and II crossover regulation. Here, we investigate specific

features of MEI-218 and REC to understand better how these proteins contribute to meiotic

recombination. We find that the N-terminus of MEI-218 is dispensable for crossover formation

and general crossover distribution. By mutating key residues in REC's Walker A and B motifs

(rec^{KA} and rec^{DA}, respectively), we found that rec^{KA} mutants exhibit no Class I crossover defect,

while Class II crossovers are significantly increased. Surprisingly, rec^{DA} mutants exhibit a severe

decrease in Class I crossovers and a significant increase in Class II crossovers. Our results

suggest that the mei-MCMs function in multiple roles and may complex in a variety of

configurations to properly regulate crossover formation.

Materials and Methods

Drosophila stocks. Flies were maintained on standard medium at 25°C. Some mutant alleles

have been previously described, including *mei-9ᵃ* (Baker and Carpenter 1972; Yıldız *et al.* 2004),

mei-218¹ and *mei-218⁶* (Baker and Carpenter 1972; Mckim *et al.* 1996), Blm^{N1} and Blm^{D2}

(Mcvey *et al.* 2007), *rec¹* and *rec²* (Grell 1978; Matsubayashi and Yamamoto 2003; Blanton *et*

al. 2005). The maternal-effect lethality in Blm^{N1}/Blm^{D2} mutants was overcome by the

UAS::GAL4 rescue system previously described (Kohl *et al.* 2012).

Generating mei-218 transgenic alleles. The transgenes for *mei-218$^{\Delta N}$* and *mei-218^{FL}* were

constructed by cloning cDNA for *mei-218* into *P{attBUASpW}* (AddGene). Full-length *mei-218*

included codons 1-1186; the *mei-218$^{\Delta N}$* transgene included codons 527-1186. Transgenics were

made by integrating into a phiC31 landing site in 2A on the *X* chromosome.

Generating **rec^{KA}** ***and*** **rec^{DA}** ***mutants.*** Annealed oligonucleotides were inserted into *Bbs*I-

digested   pU6-BbsI-chiRNA   plasmid   (Addgene). rec^{KA}:

CTTCGCCGAGAAGGGATAGTAAAC; rec^{DA}: CTTCGTTGCAGTGCCTACAATCAG.

Resulting plasmids were co-injected with repair template plasmid, consisting of synthesized

gBlocks (IDT DNA) cloned into pBlueScript plasmid (sequences available on request). Injected

larvae were raised to adulthood, and their male progeny were crossed to *TM3*/*TM6B* females

(Bloomington Stock Center) to generate stocks, after which DNA was extracted for screening

through PCR and restriction digest.

Nondisjunction assay. *X*-chromosome nondisjunction (NDJ) was assayed by mating virgin

females to *y cv v f / T(1:Y)B^S* males. Each cross was set up as a single experiment with 20-50

separate vials. The progeny of each vial were counted separately. Viable nondisjunction progeny

are *XXY* females with Bar eyes and *XO* males with Bar$^+$ eyes and the phenotypes from *y cv v f*

chromosome. Total (adjusted) represents the total with inviable exceptional progeny accounted

for (*XXX* and *YO*). NDJ rates and statistical comparisons were done as in Zeng et al. 2010.

Crossover distribution assay. Crossover distribution on chromosome *2*L was scored by crossing

virgin *net dpp^{d-ho} dp b pr cn / +* female flies with mutant background of interest to *net dpp^{d-ho} dp*

b pr cn homozygous males. Each cross was set up as a single experiment with at least 25

separate vials scored. The first set of vials was flipped after three days of mating into vials of a

new batch, although these were counted as one experiment. Batch effects for recombination

assays have not been observed in repeated studies for multiple genotypes used in this study

(Figure S4). These include wild-type (unpublished data), *Blm* (unpublished data), *rec* (Blanton *et*

al. 2005; Kohl *et al.* 2012), *mei-9* (Sekelsky *et al.* 1995), and *mei-9; rec* (Blanton *et al.* 2005).

All progeny were scored for parental and recombinant phenotypes. Crossover numbers in flies

are shown as cM where cM = (number of crossovers / total number of flies) * 100. Chi-squared

tests with Bonferroni correction were performed for each interval. For total cM, Fisher's Exact

Test was used to compare total crossovers to total number of flies. Crossover distribution is

represented as cM/Mb where Mb is length of the interval without transposable elements (TEs)

because crossovers rarely occur within TEs (Miller *et al.* 2016).

Protein structure and alignment. Structural domains of proteins were determined by using

PHYRE 2. All of the MCM regions identified correspond to the protein data bank ID #c2vl6C

and the AAA ATPase domains identified correspond to protein data bank ID #d1g8pa.

Alignment of the Walker A and Walker B motifs (Kohl *et al.* 2012) was done using MEGA 5

and aligned with the ClustalW program. Identical and conserved residues are shaded based on

groups of amino acids with similar chemical properties.

Data availability. All data necessary for confirming the conclusions in this paper are included in

this article and in supplemental figures and tables. *Drosophila* stocks and plasmids described in

this study are available upon request. We have uploaded Supplemental Material to Figshare.

Figure S1 illustrates distribution of Msh4, Msh5, Mcm8, Mcm9, MEI-217, and MEI-218 in

Diptera. Figure S2 illustrates the structure of MEI-217 and MEI-218 in Diptera. Figure S3 shows

sequence alignment of MEI-218. Figure S4 compares crossover frequencies in different batches

of the same genotype. Figure S5 details the cross scheme of *mei-218* transgene experiments.

Table S1 includes analysis of genetic interval differences between *WT* and *mei-218^{FL}*. Table S1

includes analysis of genetic interval differences between *mei-218^{FL}* and *mei-218$^{\Delta N}$*. Table S2

includes complete data set for calculating nondisjunction of *WT, rec$^-$/rec$^+$*, and *rec^{DA}/+*. Table S3

includes all data sets for meiotic crossovers for all genotypes discussed.

Results and Discussion

The N-terminus of MEI-218 is dispensable for crossover formation

MCMDC2 is a distantly-related member of the MCM family of proteins that is unique in

that the ATPase domain is predicted to be incapable of binding or hydrolyzing ATP. Orthologs

in Dipteran insects are further distinguished by having the N-terminal and ATPase-like domains

encoded in separate open reading frames. The two polypeptides, MEI-217 and MEI-218 interact

physically, at least in *Drosophila melanogaster*, presumably reconstituting a single MCM-like

protein. MEI-218 is also distinguished by possessing an N-terminal extension of variable length

in different species. *Drosophila melanogaster* MEI-218 can be divided into three distinct regions

(Figure 1A): an N-terminal tail (residues 1-500), a central acidic region (residues 500-800) and

the C-terminal ATPase-related region (residues 850-1116) (Kohl *et al.* 2012; Brand *et al.* 2018).

The N-terminal and middle regions are predicted to be disordered (Kohl *et al.* 2012) and are

poorly conserved (Figure S3). Results obtained through gene swap experiments suggest that the

N-terminal tail and central region regulate crossover number and distribution within *Drosophila*

species (Brand *et al.* 2018).

To genetically examine the function of the N-terminus of MEI-218, we compared

functions of a transgene that expresses a truncated form of MEI-218 that lacks the N-terminal

526 amino acids (*mei-218$^{\Delta N}$*) to a matched full-length transgene (*mei-218^{FL}*) (Figure 2A). Due to

the relatively high conservation among *Drosophila* species, the middle region of mei-218 was

retained for this experiment (Figure S3). Using the *UAS*/*GAL4* system (Duffy 2002), we

expressed both constructs in *mei-218* null mutants using the germline-specific *nanos* promoter

and measured crossovers along five adjacent intervals that span most of *2*L and part of *2R*

(Figure S4; for simplicity, we refer to this chromosomal region as *2*L.)

In wild-type females, the genetic length of *2*L is 45.8 cM (Hatkevich *et al.* 2017) (Figure

2B), whereas *mei-218* mutants exhibit a severe decrease in crossovers, with genetic length of

2.92 cM (Kohl *et al.* 2012). Expression of *mei-218^{FL}* in *mei-218* mutants (*mei-218^{FL}*) fully

rescues the crossover defect, exhibiting a genetic length of 54.1 cM. Unexpectedly, expression of

mei-218$^{\Delta N}$ in *mei-218* mutants (*mei-218$^{\Delta N}$*) restored crossing over to the same level as in

mei-218; mei-218^{FL} (55.9 cM; *n.s. p* = 0.61).

Brand *et al.* (2018) expressed *Drosophila mauritiana* MEI-217 and MEI-218 in

Drosophila melanogaster and found that crossovers were increased in proximal and distal

regions, resulting in an overall change in crossover distribution. We examined crossover

distribution in *mei-218; mei-218^{FL}* and *mei-218; mei-218$^{\Delta N}$* (Figure 2C). Overall, distributions

are similar, with both genotypes exhibiting a strong inhibition of crossovers near the centromere

(referred to as the centromere effect; Beadle 1932) and the majority of the crossovers placed in

the medial-distal regions (Figure 2C).

We conclude that the N-terminal tail of MEI-218 is dispensable for both crossover

formation and overall distribution on chromosome *2*L. This conclusion is supported by the

observation that of 16 sequenced mutations in *Drosophila melanogaster mei-218*, 14 are

nonsense or frameshift, and the only two missense mutations alter residues in the C-terminus

(amino acids 845 and 1107) (Collins *et al.* 2012).

The reasons why the MCM domains have been separated into MEI-217 and MEI-218

polypeptides and why MEI-218 has an N-terminal extension are unknown, but this structure has

been maintained for more than 250 million years of Dipteran evolution (Supplemental Figures

S2). Interestingly, MEI-218 is expressed moderately highly in testes (Thurmond *et al.* 2018)

even though males do not experience meiotic recombination. The predominant or exclusive

transcript in males does not encode MEI-217 (Thurmond *et al.* 2018), the seemingly obligate

partner for MEI-218 the female meiotic recombination. Males that lack *mei-218* are viable,

fertile, and do not exhibit elevated nondisjunction (Baker and Carpenter 1972; Mckim *et al.*

1996). For these reasons, we speculate that an unknown function of MEI-218 (independent of

MEI-217) in the male germline explains why its overall structure has been evolutionarily

maintained.

REC ATPase motifs are required for crossover formation

Of the three known mei-MCM subunits, only REC harbors well-conserved Walker A and

B motifs, suggesting that REC has ATP binding and hydrolysis activity (Kohl *et al.* 2012). It is

unknown whether the mei-MCM complex utilizes REC's putative ATPase activity for its

function *in vivo*. To test this, we used CRIPSR/Cas9 to introduce into *rec* mutations predicted to

disrupt Walker A and B motif functions (Figure 3A). The Walker A mutation (rec^{KA}) results in

substitution of a conserved lysine residue with alanine; this mutation in other AAA+ ATPases,

including replicative MCMs, prevents binding of ATP (Bell and Botchan 2013). The Walker B

mutation (rec^{DA}) results in substitution of a conserved aspartic acid with alanine; in MCMs and

other AAA+ ATPases, this mutation destroys the ability to coordinate Mg^{++} for ATP hydrolysis

(Bochman *et al.* 2008).

We assayed crossover frequency along *2*L in rec^{KA} and rec^{DA} mutants (Figure 3B).

Surprisingly rec^{KA} ATP binding mutants exhibit a genetic length of 44.9 cM, which is not

significantly different from wild-type ($p = 0.4016$), suggesting that ATP binding by REC is not

required for crossover formation. Conversely, there is a severe reduction in crossovers in rec^{DA}

mutants, with a genetic length of 1.6 cM ($p < 0.0001$), suggesting that REC's ability to hydrolyze

ATP is required for crossover formation.

Because the genetic length of rec^{DA} is significantly lower than rec null mutants (Figure

3B, $p < 0.0001$), we hypothesized that rec^{DA} is an antimorphic mutation. To test this, we

examined crossover levels and X chromosome nondisjunction (NDJ) in rec^{DA}/rec^{+} (Figure 3B

and 3C, respectively). The genetic length of 2L in $rec^{DA/+}$ is slightly lower than wild-type, but

not significantly different (43.9 cM and 45.8 cM, respectively; $p = 0.35$). For X-NDJ, both

wild-type and rec^{-}/rec^{+} mutants exhibit rates below 0.5%, while rec^{DA}/rec^{+} mutants exhibit a

significant increase to 1.4% NDJ ($p < 0.0001$). These data support the conclusion that rec^{DA} is

weakly antimorphic and suggest that rec^{DA} results in an inactive mei-MCM complex that is

antagonistic to the wild-type complex. In light of these interpretations, we propose that the

mei-MCM complex binds to recombination sites independent of REC binding to ATP, and that

REC-dependent ATP hydrolysis is required for the removal of the mei-MCM complex from

these sites.

The phenotypes of rec^{KA} and rec^{DA} mutants suggest that REC's ability to hydrolyze ATP

is required for crossover formation, whereas its ATP binding capability is dispensable. The

disparate requirements for REC's ATP binding and hydrolysis are similar to those of other

ATPase-dependent complexes. Rad51 paralogs, which form multi-protein complexes and contain

Walker A and B motifs, are proposed to exhibit ATPase activity in *trans* between adjacent

subunits, each of which contributes a Walker A or Walker B motif to the active site (Wu *et al.*

2004; Wu *et al.* 2005; Wiese *et al.* 2006). Because neither MEI-217 nor MEI-218 possess an

ATPase domain that harbors conserved key enzymatic residues (Figure 1B) (Kohl *et al.* 2012),

we propose that ATPase activity of the mei-MCM complex requires REC for ATP hydrolysis

and an unknown mei-MCM protein for ATP binding. Alternatively, because REC is highly

diverged, its Walker A and B motifs may function non-canonically. Biochemical studies are

needed to test these hypotheses, but these may require identification of the putative missing

subunit.

REC-dependent ATP hydrolysis is required for MEI-9-dependent crossovers

To gain insight into the crossover pathways that are used in rec^{KA} and rec^{DA} mutants, we

examined whether these crossovers require the Class I endonuclease/resolvase. In *Drosophila,*

the catalytic subunit of the putative Class I meiosis-specific endonuclease is MEI-9 (Sekelsky *et*

al. 1995; Yıldız *et al.* 2002; Hatkevich *et al.* 2017). The *2*L genetic length within a *mei-9* mutant

is 2.75 cM (Figure 4), demonstrating that at least 90% of crossovers are dependent upon MEI-9.

However, the genetic length in *mei-9*; *rec* mutants is not significantly different than that of *rec*

null single mutants (4.11 cM *vs* 4.66 cM, *p* = 0.64) suggesting that in the absence of REC, the

resulting crossovers are likely independent of MEI-9. Similarly, it has been shown previously

that *mei-218 mei-9* double mutants do not have reduced crossovers compared to *mei-218* single

mutants (Sekelsky *et al.* 1995), indicating that crossovers generated in the absence of the mei-

MCM complex are MEI-9-independent.

Because rec^{KA} mutants exhibit the same distribution and number of crossovers as

wild-type (Figure 3B), we hypothesized that rec^{KA} crossovers are dependent on MEI-9. To test

this, we examined genetic length across *2*L in *mei-9; rec^{KA}* double mutants (Figure 4). Mutants

for *mei-9; rec^{KA}* exhibit a genetic length of 2.72 cM, which is significantly decreased compared

to the rec^{KA} single mutant (*p* < 0.0001), but not significantly different from *mei-9* single mutants

(*p* = 0.94), showing that crossovers in rec^{KA} are indeed dependent upon MEI-9 nuclease. In

contrast, we predicted that crossovers in rec^{DA} will be independent of MEI-9, similar to

crossovers generated in *rec* null mutants. We observe that *mei-9; rec^{DA}* double mutants exhibit a

genetic length of 1.1 cM, which is significantly lower than that of *mei-9* single mutants ($p <$

0.001). Importantly, crossing over in the *mei-9; rec^{DA}* double mutant is not significantly different

than in rec^{DA} single mutants ($p = 0.23$), demonstrating that crossovers in rec^{DA} are independent of

MEI-9 (Figure 4).

From these data we conclude that the crossovers in rec^{KA} mutants arise through the

normal, MEI-9-dependent pathway, whereas mitotic nucleases generate the residual crossovers

in rec^{DA} mutants. These data show that REC^{KA} functions normally in the Class I pathway, but

this pathway is nonfunctional in *rec* null and rec^{DA} mutants. We suggest that the REC's ability to

hydrolyze, but not bind, ATP is required for the formation of Class I crossovers.

REC ATPase motifs are required to prevent Class II crossovers

In wild-type *Drosophila*, most or all crossovers are generated through the Class I

pathway (Hatkevich *et al.* 2017), and these crossovers are dependent upon the mei-MCM

complex (Kohl *et al.* 2012). However, in *Blm* mutants, crossovers are generated exclusively

through the Class II pathway (Hatkevich *et al.* 2017). In *Drosophila Blm* mutants, meiotic

crossovers are decreased by 30%, suggesting that the Class II pathway is less efficient at

generating crossovers than the Class I pathway, even though what may be the primary anti-

crossover protein, Blm helicase, is absent. It has previously been shown that loss of Blm

suppresses the high nondisjunction of *mei-218* and *rec* mutants (Kohl *et al.* 2012). However, in

Blm rec double mutants, crossovers are increased significantly compared to *Blm* single mutants

(Kohl *et al.* 2012), suggesting that REC and/or the mei-MCM complex has an anti-crossover role

in *Blm* mutants, and therefore in the Class II crossover pathway.

To further understand the role of REC in the Class II pathway, we investigated whether

REC's predicted ATP binding or hydrolysis function is required for its Class II anti-crossover

function. To do this, we measured the crossovers across *2*L in rec^{KA} and rec^{DA} in the background

of *Blm* mutants. If REC ATP binding or hydrolysis is required for an anti-crossover role in Class

II, then the genetic length of *Blm* rec^{KA} or *Blm* rec^{DA} double mutants will be similar to that of *Blm*

rec double mutants. Conversely, if REC ATP binding or hydrolysis is not required, then double

mutants will exhibit genetic lengths similar to that of *Blm* single mutants.

Interestingly, *Blm* rec^{KA} mutants exhibit a genetic length of 43.3 cM, which is not

significantly different than *Blm rec* mutants ($p = 0.10$) but significantly higher than *Blm* single

mutants ($p < 0.0001$; Figure 5A). Similarly, *Blm* rec^{DA} double mutants have a recombination rate

of 53.4 cM, which not significantly different from *Blm rec* double mutants ($p = 0.52$), but

significantly higher than *Blm* single mutants ($p < 0.0001$). These results suggest that REC's

predicted abilities to bind and hydrolyze ATP are both required for the inhibition of crossovers at

REC-associated Class II recombination sites. Therefore, it appears that REC forms different

complexes within the Class II pathway and Class I pathway. It is unknown whether this Class II

REC-associated complex requires the other mei-MCM proteins, and additional genetic studies

will be valuable to discern this.

In summary, the mei-MCMs are a family of diverged proteins that help to establish the

recombination landscape in *Drosophila melanogaster* by promoting Class I crossovers and

inhibiting Class II crossovers. Results obtained in this study have further elucidated meiotic

recombination roles for two mei-MCM proteins, MEI-218 and REC. While the N-terminus of

MEI-218 is dispensable for crossover formation (Figure 2), REC's predicted ability to bind and

hydrolyze ATP exhibit differential requirements for regulating Class I and Class II crossover

formation. From our genetic analyses, we suggest that the Walker B motif of REC, but not the

Walker A motif, is required for promoting the formation Class I, MEI-9 dependent crossovers

(Figures 3 and 4). The weakly antimorphic phenotype of rec^{DA} demonstrates that an impaired

REC Walker B mutant renders a poisonous complex – a complex in which we propose cannot be

released from recombination sites. Both Walker A and Walker B motifs block crossovers in the

Class II pathway, suggesting that REC forms different complexes to execute its pro- and anti-

crossover functions. Biochemical and cytological studies are needed to support or refute these

hypotheses.

Acknowledgements

We thank Juan Carvajal Garcia, Carolyn Turcotte, and anonymous reviewers for thoughtful

comments. This work was supported in part by a grant from the National Institute of General

Medical Sciences to J.S. under award 1R35 GM-118127. K.P.K. was supported in part by NIH

grant P20GM103499. T.H. was supported in part by NIH grants 5T32GM007092 and

1F31AG055157.

 Figure Legends

 Figure 1. MCM protein structure and alignments. (A) Structural domains of *Drosophila*

 melanogaster REC, MEI-217, MEI-218 and *Mus musculus* MCMDC2. Structural domains

 identified using PHYRE 2 (Kohl *et al.* 2012). "MCM domain" corresponds to protein data bank

 ID #c2vl6C and the AAA ATPase domains identified correspond to protein data bank ID

 #d1g8pa. The X on *Dm* MEI-218 and *Mm* MCMDC2 represents predicted inactive AAA ATPase

 domains. (B) Consensus sequence for Walker A motif (Walker *et al.* 1982), and consensus

 sequence for Walker B motif (Forsburg 2004). Identical or conserved amino acids are denoted

 with black background. Arrows denote the conserved catalytic residues.

 Figure 2. The role of MEI-218 N-terminus in crossover formation and distribution. (A)

 Schematic of transgenes for full length *mei-218* and N-terminal deleted *mei-218*, in which the

 first 526 amino acids are absent. (B) Map units of WT (Hatkevich *et al.* 2017), *mei-218* (Kohl *et*

 al. 2012)*, mei-218^{FL}* and *mei-218$^{\Delta N}$*. Map units represented as centimorgans (cM). Error bars

 indicate 95% confidence intervals. *n.s.* = not significant ($p = 0.61$). (C) Crossover distribution

 (solid lines) of *mei-218^{FL}* and *mei-218$^{\Delta N}$* represented as cM/Mb. Mb is measured distance of

 defined interval, excluding centromere, pericentromeric heterochromatin and transposable

 elements. Dotted lines represent mean crossover density across *2*L. Figure S5 details the cross

 scheme of *mei-218* transgene experiments. Refer to tables S1and S3 for complete data sets.

 Figure 3. REC ATPase binding and hydrolysis requirements for crossover formation. (A)

 Schematic representation of the mutated residues in *rec^{KA}* and *rec^{DA}*. (B) Map units of *WT*

 (Hatkevich *et al.* 2017), *rec^{1}/rec^{2}, rec^{KA},* and *rec^{DA}, rec^{DA}/rec^{+}*. Map units represented as

 centimorgans (cM). Error bars show 95% confidence intervals. (C) Percent nondisjunction of

WT, rec¹/rec⁺, and recᴰᴬ/rec⁺. (D) Model of possible complex depicting the functional Walker B

motif of REC protein interacting with a Walker A motif on a potential partner. $* \, p < 0.05$; $*** p$

< 0.0001. Refer to tables S2 and S3 for complete data sets.

Figure 4. MEI-9-dependent crossovers in *recᴷᴬ* and *recᴰᴬ* mutants. Map units of *WT*

(Hatkevich *et al.* 2017), *rec, mei-9, mei-9;rec, recᴷᴬ, mei-9;recᴷᴬ, recᴰᴬ*, and *mei-9;recᴰᴬ*. Map

units represented as centimorgans (cM). Error bars show 95% confidence intervals. $* \, p < 0.05$ $**$

$p < 0.001$ $*** p < 0.0001$; (*mei-9* vs *mei-9; recᴷᴬ* $p = 0.94$) (*recᴰᴬ* vs *mei-9; recᴰᴬ* $p = 0.23$).

Refer to Table S3 for complete data set.

Figure 5. Requirements of REC ATPase activity in Blm function. Map units of *WT*

(Hatkevich *et al.* 2017), *Blm* (Kohl *et al.* 2012)*, rec, Blm rec* (Kohl *et al.* 2012)*, Blm recᴷᴬ,* and

Blm recᴰᴬ. Map units represented as centimorgans (cM). Error bars show 95% confidence

intervals. Refer to Table S3 for complete data set. $*** \, p < 0.0001$. (*Blm rec* vs *Blm recᴷᴬ* $p =$

0.10) (*Blm rec* vs *Blm recᴰᴬ* $p = 0.52$).

References

Baker, B. S., and A. T. C. Carpenter, 1972 Genetic analysis of sex chromosomal meiotic mutants
in *Drosophila melanogaster*. Genetics **71:** 255-286.
Beadle, G. W., 1932 A possible influence of the spindle fibre on crossing-over in *Drosophila*.
Proc Natl Acad Sci U S A **18:** 160-165.
Bell, S. D., and M. R. Botchan, 2013 The minichromosome maintenance replicative helicase.
Cold Spring Harb Perspect Biol **5:** a012807.
Blanton, H. L., S. J. Radford, S. Mcmahan, H. M. Kearney, J. G. Ibrahim *et al.*, 2005 REC,
Drosophila MCM8, drives formation of meiotic crossovers. PLoS Genet **1:** e40.
Bochman, M. L., S. P. Bell and A. Schwacha, 2008 Subunit organization of Mcm2-7 and the
unequal role of active sites in ATP hydrolysis and viability. Mol Cell Biol **28:** 5865-
5873.

Brand, C. L., M. V. Cattani, S. B. Kingan, E. L. Landeen and D. C. Presgraves, 2018 Molecular
evolution at a meiosis gene mediates species differences in the rate and patterning of
recombination. Curr Biol **28:** 1289-1295 e1284.
Collins, K. A., J. G. Callicoat, C. M. Lake, C. M. Mcclurken, K. P. Kohl *et al.*, 2012 A germline
clone screen on the *X* chromosome reveals novel meiotic mutants in *Drosophila*
melanogaster. G3 (Bethesda) **2:** 1369-1377.
De Los Santos, T., N. Hunter, C. Lee, B. Larkin, J. Loidl *et al.*, 2003 The Mus81/Mms4
endonuclease acts independently of double-Holliday junction resolution to promote a
distinct subset of crossovers during meiosis in budding yeast. Genetics **164:** 81-94.
Duffy, J. B., 2002 GAL4 system in *Drosophila*: a fly geneticist's Swiss army knife. Genesis **34:**
1-15.
Forsburg, S. L., 2004 Eukaryotic MCM proteins: beyond replication initiation. Microbiol Mol
Biol Rev **68:** 109-131.
Grell, R. F., 1978 Time of recombination in the *Drosophila melanogaster* oocyte: evidence from
a temperature-sensitive recombination-deficient mutant. Proc Natl Acad Sci U S A **75:**
3351-3354.
Hatkevich, T., K. P. Kohl, S. Mcmahan, M. A. Hartmann, A. M. Williams *et al.*, 2017 Bloom
syndrome helicase promotes meiotic crossover patterning and homolog disjunction. Curr
Biol **27:** 96-102.
Hatkevich, T., and J. Sekelsky, 2017 Bloom syndrome helicase in meiosis: Pro-crossover
functions of an anti-crossover protein. Bioessays **39**.
Hollingsworth, N. M., and S. J. Brill, 2004 The Mus81 solution to resolution: generating meiotic
crossovers without Holliday junctions. Genes Dev **18:** 117-125.
Iyer, L. M., D. D. Leipe, E. V. Koonin and L. Aravind, 2004 Evolutionary history and higher
order classification of AAA+ ATPases. J Struct Biol **146:** 11-31.
Kohl, K. P., C. D. Jones and J. Sekelsky, 2012 Evolution of an MCM complex in flies that
promotes meiotic crossovers by blocking BLM helicase. Science **338:** 1363-1365.
Lake, C. M., and R. S. Hawley, 2012 The molecular control of meiotic chromosomal behavior:
events in early meiotic prophase in Drosophila oocytes. Annu Rev Physiol **74:** 425-451.
Liu, H., J. K. Jang, J. Graham, K. Nycz and K. S. Mckim, 2000 Two genes required for meiotic
recombination in Drosophila are expressed from a dicistronic message. Genetics **154:**
1735-1746.
Matsubayashi, H., and M. T. Yamamoto, 2003 REC, a new member of the MCM-related protein
family, is required for meiotic recombination in Drosophila. Genes Genet Syst **78:** 363-
371.
Mckim, K. S., J. B. Dahmus and R. S. Hawley, 1996 Cloning of the *Drosophila melanogaster*
meiotic recombination gene *mei-218*: a genetic and molecular analysis of interval 15E.
Genetics **144:** 215-228.
Mcvey, M., S. L. Andersen, Y. Broze and J. Sekelsky, 2007 Multiple functions of Drosophila
BLM helicase in maintenance of genome stability. Genetics **176:** 1979-1992.
Miller, D. E., C. B. Smith, N. Yeganeh Kazemi, A. J. Cockrell, A. V. Arvanitakis *et al.*, 2016
Whole-genome analysis of individual meiotic events in *Drosophila melanogaster* reveals
that noncrossover gene conversions are insensitive to interference and the centromere
effect. Genetics **203:** 159-171.

Sekelsky, J. J., K. S. Mckim, G. M. Chin and R. S. Hawley, 1995 The Drosophila meiotic
recombination gene *mei-9* encodes a homologue of the yeast excision repair protein
Rad1. Genetics **141:** 619-627.
Thurmond, J., J. L. Goodman, V. B. Strelets, H. Attrill, L. S. Gramates *et al.*, 2018 FlyBase 2.0:
The next Generation. Nucleic Acids Res **47:** D759-765.
Walker, J. E., M. Saraste, M. J. Runswick and N. J. Gay, 1982 Distantly related sequences in the
alpha- and beta-subunits of ATP synthase, myosin, kinases and other ATP-requiring
enzymes and a common nucleotide binding fold. EMBO J **1:** 945-951.
Wiese, C., J. M. Hinz, R. S. Tebbs, P. B. Nham, S. S. Urbin *et al.*, 2006 Disparate requirements
for the Walker A and B ATPase motifs of human RAD51D in homologous
recombination. Nucleic Acids Res **34:** 2833-2843.
Wu, Y., Y. He, I. A. Moya, X. Qian and Y. Luo, 2004 Crystal structure of archaeal recombinase
RADA: a snapshot of its extended conformation. Mol Cell **15:** 423-435.
Wu, Y., X. Qian, Y. He, I. A. Moya and Y. Luo, 2005 Crystal structure of an ATPase-active
form of Rad51 homolog from Methanococcus voltae. Insights into potassium
dependence. J Biol Chem **280:** 722-728.
Yıldız, Ö., H. Kearney, B. C. Kramer and J. J. Sekelsky, 2004 Mutational analysis of the
Drosophila DNA repair and recombination gene *mei-9*. Genetics **167:** 263-273.
Yıldız, Ö., S. Majumder, B. C. Kramer and J. Sekelsky, 2002 Drosophila MUS312 interacts with
the nucleotide excision repair endonuclease MEI-9 to generate meiotic crossovers. Mol
Cell **10:** 1503-1509.
437

Author Summary

Crossover formation between homologs is essential for accurate segregation at the end of
meiosis I. Crossovers are typically formed through two pathways: Class I and Class II. The mei-
MCMs are a class of proteins that promote Class I crossover formation and prohibit Class II
crossovers in Drosophila. Although the mei-MCMs are conserved, little is known about their
function. Here, we investigate the roles of two mei-MCMs, REC and MEI-218, in Class I and
Class II crossover formation in Drosophila. From results in this study, we generate novel,
testable hypotheses to further elucidate the meiotic function of the mei-MCM proteins.