1

4 **Behavioral evolution drives hindbrain diversification among**

5 **Lake Malawi cichlid fish**

6 **Authors:** Ryan A. York[1,2*], Allie Byrne[1], Kawther Abdhilleh[3], Chinar Patil[3], J. Todd Streelman[3],

7 Thomas E. Finger[4,5], Russell D. Fernald[1,6]

Affiliations:

[1]Department of Biology, Stanford University, Stanford, California 94305, USA

[2]Department of Neurobiology, Stanford University, Stanford, California 94305, USA

[3]School of Biological Sciences and Institute of Bioengineering and Bioscience, Georgia Institute

of Technology, Atlanta, Georgia 30332, USA

[4]Department of Cell and Developmental Biology, University of Colorado Denver, Aurora, CO

80045

[5]Rocky Mountain Taste and Smell Center, University of Colorado Denver, Aurora, CO 80045

[6]Neuroscience Institute, Stanford University, Stanford, California 94305, USA

[*]Correspondence: ryanyork@stanford.edu

Abstract

The evolutionary diversification of animal behavior is often associated with changes in

the structure and function of nervous systems. Such evolutionary changes arise either through

alterations of individual neural components ("mosaically") or through scaling of the whole brain

("concertedly"). Here we show that the evolution of a specific courtship behavior in Malawi

cichlid fish, the construction of mating nests known as bowers, is associated with rapid,

extensive, and specific diversification of orosensory, gustatory centers in the hindbrain. We find

that hindbrain volume varies significantly between species that build pit (depression) compared

to castle (mound) type bowers and that hindbrain features evolve rapidly and independently of

phylogeny among castle-building species. Using immediate early gene expression, we confirmed

a functional role for hindbrain structures during bower building. Comparisons of bower building

species in neighboring Lake Tanganyika show patterns of neural diversification parallel to those

in Lake Malawi. Our results suggest that mosaic brain evolution via alterations to individual

brain structures is more extensive and predictable than previously appreciated.

Keywords: Brain evolution, courtship behavior, mosaic evolution, hindbrain, vagal lobe, cichlid

fish, Lake Malawi

Introduction

Animal behaviors vary widely, as do their neural phenotypes[1]. Evolutionary

neuroscience identifies how the brain diversifies over time and space in response to selective

pressures [2]. A key goal of evolutionary neuroscience has been to identify whether brain

structures evolve independently ("mosaically") or in tandem with each other as they reflect key

life history traits, especially behavior [3-6]. While a number of studies have linked variation in

brain structure with other traits across evolutionary time [2, 7-9], it remains unclear whether or

not this variation is predictable. Specifically, when similar behavioral traits evolve among two or

more species, do their neural bases evolve correspondingly? If parallel brain evolution is

predictable then it may be possible to understand general principles of neural organization and

function across animals. This would expand our ability to manipulate brain function, but if this is

not true, new strategies will be needed to reveal the mechanisms of brain evolution.

Fishes, as both the most speciose (50% of extant vertebrates) and most varied vertebrate

radiation [10] offer opportunities to answer these questions. Fish species live in diverse

ecological, sensory, and social environments and have evolved elaborate variations in neural

structure and function from a common basic ground plan [11] making rapid and variable

diversification of brain structures a broad and general feature of their evolution [10].

The cichlid fishes of Lake Malawi, Africa offer a particularly striking model of these

patterns of diversification. Although geologically young (less than 5 million years old), Lake

Malawi contains at least 850 species of cichlids [12] that, based on molecular phylogenetic

analyses, can be sorted into six diverse clades [13]: sand-dwelling shallow benthic species (287

species), deep benthic species (150 species), rock-dwelling 'mbuna' (328 species), the deep-

water pelagic genus *Diplotaxodon* (19 species), the pelagic genus *Rhamphochromis* (14 species),

and the sand-breeding pelagic 'utaka' species (55 species) [14]. These clades exhibit substantial

variation in habitat use, visual sensitivity, diet, behavior, and coloration as well as craniofacial

morphology and tooth shape, presumably arising through repeated divergence in macrohabitat,

trophic specializations, diet, and coloration [15-17].

We studied the behavior, brain structures and bower construction of ~200 sand-dwelling

shallow living benthic species with males that build species-specific mating nests, known as

bowers, to court females in competition with conspecifics [17,18]. There are two basic bower

types: 1) pits or depressions in the sand, and 2) castles, where sand is heaped into a volcano

structure with showing species-specific differences in size and shape [17,18]. Pits and castles are

constructed via differential scooping and spitting of sand. Pit and castle type bower building is

innate and appears to be rapidly and relatively free from phylogenetic constraint; multiple sand-

dwelling genera contain both pit and castle-building species [17]. Furthermore, phylogenomic

analyses suggest evidence of the repeated evolution of bower types in Malawi cichlid phylogeny

[17].

How is construction of different bower types reflected in brain anatomy? Do differences

in brain structure evolve in tandem with bower building and, if so, how are they organized and

functionally related to this behavior? Here we address these questions using neuroanatomical,

phylogenetic, functional, and behavioral analyses and compare patterns of diversification in Lake

Malawi to other East African cichlid radiations.

Results

Bower type predicts hindbrain volume

We compiled measurements of six brain regions (telencephalon, hypothalamus,

cerebellum, optic tectum, olfactory bulb, and hindbrain) for 37 bower building species from a

phenotypic data set comprising brain measurements for 189 cichlid species from East Africa and

Madagascar [19]. For each brain region, we calculated a volume estimate using an ellipsoid

model [20] and controlled for size differences between species by normalizing the results using

mean standard length for each species (Supplementary material). Pairwise comparisons of

normalized volumetric measures for each region revealed a significant difference in hindbrain

volume between pit digging and castle building species (Kruskal-Wallis test; H = 13.87, 1 d.f., p

= 0.00019) but not for any other brain structure (Table 1). Plotting structural volume against

standard length reveals strong evidence for allometric scaling of the pit and castle hindbrain

(Figure 1A). An analysis of variance (ANOVA) comparing the slopes of standard length

compared to those of hindbrain volume reveals that pit and castle species significantly differ in

this relationship (F = 25.04, 1 d.f., $p = 1.82 \times 10^{-05}$) as also confirmed by post hoc Tukey test (p

<0.05).

Hindbrain volume varies independent of phylogeny

If hindbrain volume evolved mosaically, as suggested by our allometric analyses, then

pairwise tests of phenotypic variation controlling for phylogeny should reveal a lack of similarity

among closely related species in the size of this brain region. Phylogenetic ANOVA using a

genome-wide maximum likelihood phylogeny [21] revealed significant variation between pit and

castle species independent of relatedness (F = 13.18, 1 d.f., $p < 0.05$). Plotting the distribution of

volumetric measures along the full tree demonstrates this pattern (Figure 1B). We next calculated

more explicit measures of the phylogenetic signal using Pagel's λ (Pagel 1999) and Blomberg's

K [22]. Pagel's λ ranges from 0 to 1 with values closer to 1 representing stronger phylogenetic

signal while values of Blomberg's $K < 1$ suggest less phylogenetic signal in the trait. Both

measures indicated a lack of phylogenetic signal among the species tested (λ = 0.25; K = 0.29),

suggesting that hindbrain volume may be rapidly evolving but not correlated to phylogeny [22].

133   Since recently evolved clades such as Lake Malawi cichlids are prone to processes such

as introgression and incomplete lineage sorting that make resolving phylogenetic relationships

difficult [12], we re-performed the phylogenetic ANOVAs and tests of phylogenetic on non-

overlapping genomic windows each containing 10,000 SNPs (1,029 windows; Figure S1A). We

reasoned that if the patterns revealed by the genome-wide tests held in a substantial number of

these windows then the observed results should be robust to alternative phylogenetic scenarios

among the species sampled. Applying phylogenetic ANOVAs across the windows we found that

all tested yielded a p-value less than 0.05 with the median of 0.006 (Figure S1B). Similarly,

median values of phylogenetic signal were less than 1 across all windows (median λ = 0.011;

median K = 0.728; Figures S1C-D). Of note, the distributions of K and λ are multimodal,

suggesting that multiple phylogenetic patterns may still exist within the genomes of these

species. This is reflective of the genomic complexity found among Malawi cichlids mentioned

above and may have been captured by the relatively large window size used for these tests.

Nonetheless, taken together these results support a model in which hindbrain volume, at least

among the species tested, is associated with a deficit of phylogenetic signal.

Castle building is associated with increased rates of hindbrain diversification

149   We analyzed rates of trait diversification between rock and sand Malawi cichlid species

with a whole-genome phylogeny using Bayesian Analysis of Macroevolutionary Mixtures

(BAMM) v.2 [23, showing that hindbrain volume has increasingly diversified since the split

between rock and sand lineages around 800,000 years ago (Figure 1D). Separating the overall

phylogenetic tree into rock and sand clades revealed extremely different rates of diversification

between these two groups with the sand group (including bower builders) displaying a 7.4-fold

faster rate of phenotypic diversification (Figure 1E-F).

Given this rapid evolution, we hypothesized that hindbrain diversification rates would be

greatest in more recently diverging clades. Consistent with this hypothesis, the model with the

best rate shift configuration in BAMM included two significant shifts in younger bower-building

subclades within the sand-dwelling lineage (Posterior probability = 31.58; Figure S2). We further

found that three of the four best shifts accounting for the majority of posterior probabilities

sampled within the 95% credible shift set significant shifts on at least one of these branches

(Figure S2). These subclades were enriched for castle building species, including one that was

entirely composed of castle-builders, suggesting that the evolution of castle building is

associated with increased diversification rates and trait values for hindbrain size. The

phylogenetic analyses conducted here, and previously [17], suggest that the construction of pit

bowers may be ancestral from which castle building may have arisen multiple times. We then

hypothesized that hindbrain volumes among pit digging species should be more similar to those

of the rock lineage while castle building species should differ significantly. Indeed, hindbrain

volumes of pit digging and rock dwelling species are statistically indistinguishable (Kruskal-

Wallis test; H = 0.66, 1 d.f., p = 0.42) while volumes of castle building and rock dwelling

species differ substantially (Kruskal-Wallis test; H = 16.74, 1 d.f., p = 4.28 x 10^{-5}). Taken

together these results demonstrate that hindbrain volume is evolutionary labile and appears to

have rapidly diversified in concert with bower type, suggesting extensive mosaic evolution of

this structure within the sand-dwelling lineage.

Variation in hindbrain architecture and connectivity among pit and castle species

To identify whether or not variation in hindbrain volume is also associated with

cytoarchitecture and/or connectivity in that structure, we compared the neuroanatomy of two

approximately size matched species - *Copadichromis virginalis* (CV; pit-digging) and *Mchenga*

conophoros (MC; castle building, smaller hindbrain compared to TB) - and a castle builder with

a comparatively elaborated hindbrain, *Tramitichromis brevis* (TB; castle building, larger

hindbrain compared to MC). Diversification of the hindbrain in other fish species is associated

with expansion of the vagal lobe in the dorsal medulla, a key center for taste sensation and

processing that receives projections from sensory structures involved in gustation [24]. We found

that CV, MC, and TB displayed three grades of anatomical complexity in the morphology of

both these nerves (vagal nerve complex) and in associated vagal sensory structures of the

brainstem (oropharynx). The simplest morphology (Figure 2A-B), in *Copadichromis virginalis*

(CV), was similar to that of most teleost genera examined, with a smooth epithelium to the

posterior oropharynx with no specialized secondary structures such as masticatory pads or palatal

organ. Correspondingly, the vagus nerve was relatively small, with a single main root, and the

brainstem vagal complex was relatively small compared to the other species. In contrast, the oral

apparatus and vagal complex of *Mchenga conophoros* (MC) was more highly developed. A

distinct palatal organ (PO) was associated with the gill arches located most caudally and the oral

palatal surface morphology was more convoluted (Figure 2C-D). Concomitantly, the vagal nerve

was more developed showing multiple roots at the point of entrance to the brainstem and the

central vagal complex was larger than in *C. virginalis*. More complex still was the oropharynx of

Tramitichromis brevis. The posterior palate housed a large palatal organ whose epithelium was

thrown into rough protruding papillae (Figure S3A-S3C). The surface protrusions exhibited

numerous taste buds at their apex as revealed by staining for calretinin. The vagal nerve complex

was elaborated with distinct rootlets penetrating each of the branchial arches (Figure 3b). The

medulla was dominated by a large vagal lobe protruding from the dorsomedial surface.

201   Comparing sections through the vagal lobe complex of CV and MC showed differences

in elaboration of the central vagal complex corresponding to the differences in oral anatomy

(Figure 4, S3-4). In the caudal half of the medulla, the vagal lobe of CV showed a small, albeit

clearly laminated vagal lobe. Numerous densely-packed small cells lay along the medial,

ventricular face of the lobe at all levels. At the caudal end of the molecular layer of the nucleus

medialis, two relatively superficial layers of small neurons were evident. Two or more relatively

poorly defined layers of cells lay between these superficial layers and the periventricular layer.

Although the overall size of the vagal lobe of MC was larger and more highly structured than

that of CV, the essential laminated structure was similar. The vagal lobe of TB was larger still. In

both TB and MC, a layer of densely packed small neurons lined the ventricle while densely

packed and slightly clumped aggregates of small neurons dominated the superficial layers. As in

CV, several layers of neurons with no apparent organization formed the central portion of the

lobes.

214   The central projections and motor neurons of the vagus nerve were identified in all three

species by post-mortem application of diI to the vagus nerve roots just distal to their entrance

into the brainstem (see Methods). As seen in Figure S4, the pattern of termination of the vagus

nerve in all three species appears similar. In all species, the sensory root of the nerve entered the

lobe ventrolaterally and divided into a superficial and deep root terminating respectively in the

superficial half of the lobe and in the deeper one-third of the lobe with an intervening zone of

neuropil relatively free of vagal endings. This terminal-free zone was, however, penetrated by

numerous fascicles of the nerve running from the more superficial root into the deeper terminal

layers.

The motoneurons of the vagus nerve lay in a periventricular position ventral to the vagal

lobe proper, as in most other species of fish (e.g. catfish) [24]. This differs from goldfish and

common carp where vagal motor neurons lie along the innermost layer within the vagal lobe itself

[24, 25].

The palatal apparatus and attendant vagal lobe are proportionately larger in TB than in

the other 2 species examined. Nonetheless, the basic projections of the vagus nerve, the nature of

the vagal lobe lamination and the position of the vagal motoneurons appear similar to those of

the other species. In TB a distinct glossopharyngeal nerve root was identifiable (Figure S4N)

which terminated in a glossopharyngeal region (Figure S4N-O) midway between the facial lobe

(unlabeled by DiI positive fibers in Fig. S4M) and the more clearly laminated vagal lobe (Figure

S4P-S). Although this glossopharyngeal region appeared continuous with both the facial and

vagal lobes, it is clearly distinct from the latter in terms of cytological organization as in some

cyprinid species [24, 25]. The vagal lobe has superficial and deep terminal zones similar to those

in the other species described, whereas the glossopharyngeal zone does not. The terminal zone in

the glossopharyngeal zone was continuous only with the dorsal terminal zone of the vagal lobe;

no deep terminal zone was apparent. Separate glossopharyngeal and vagal motor neuron pools

were evident below the ventrolateral edge of the ventricle, each pool being situated beneath the

respective terminal field zone within the overlying lobes.

The vagal lobe is functionally involved in bower building and feeding

We used *cFos* mRNA expression as a proxy for neural activity to assess which vagal lobe

neurons are activated during bower building. Comparing the castle-building species *M.*

conophoros (MC) and the pit-digging species *C. virginalis* (CV), we measured *cFos* expression

in the vagal lobe during bower construction and compared it to control males who were housed

without sand or female conspecifics (Figure S5A-F). Cells in the vagal lobe of bower building

MC males showed greater *cFos* expression than control (2.601-fold difference, $p = 0.019$,

bootstrap 1-way ANOVA, n/group > 3) (Figure 4A). The level of *cFos* expression in the vagal

lobe of CV also increased significantly (2.517-fold difference, $p = 0.034$). Likewise, *cFos*

expression in the vagal lobe increased during feeding in both MC (2.254-fold difference, $p = $

0.046) and CV (2.438-fold difference, $p = 0.033$) (Figure 4B).

To measure vagal lobe activation during bower building we used *in situ* hybridization of

cFos and immunohistochemical labeling of phosphorylated ribosomal protein S6 (pS6). Labeling

of pS6 indicates recent neural activity and displays punctate staining, allowing for granular

quantification in individual neurons [26]. Staining of pS6 revealed that the number of VL

neurons activated during bower building differed significantly between CV and MC. As with

cFos, pS6 cell labeling was robust throughout the vagal lobe of both CV and MC after bower

building (Figure S6A-H). Quantification of pS6 labeling in the vagal lobe showed significant

differences during bower building in both species (CV: 4.12 fold difference, $p = 0.015$; MC:

13.84 fold difference, $p = 0.022$). Moreover, we found that MC bower building males displayed

an almost four-fold greater abundance of pS6 positive neurons compared to CV (3.68 fold

increase, $p = 0.04$) (Figure 4C).

Hindbrain volume has evolved in parallel with bower building in Lake Tanganyika

To ask whether parallel diversification of hindbrain volumes may have occurred in non-

Malawi cichlids, we measured variation in hindbrain volumes in the cichlid radiations of Lakes

Victoria and Tanganyika. We found that the Lake Victoria species sampled had hindbrain

volumes similar to those of the rock-dwelling cichlids of Lake Malawi (Figure 5A) whereas

normalized hindbrain volumes of Lake Tanganyika species were distributed similar to those of

Malawi species (Kruskal-Wallis test, H= 0.043, 1 d.f., $p = 0.84$), providing evidence of increased

hindbrain diversification within Tanganyika cichlids. This observation was notable given that

Lake Tanganyika has a number of reported bower building species, in contrast with Lake

Victoria where few, if any, are known. This suggests that hindbrain diversification has occurred

where cichlids have evolved this specific behavior (Malawi and Tanganyika), but not in lakes

where bower-building does not occur (Lake Victoria) . Accordingly, differences in the hindbrain

volumes of Tanganyikan bower building species compared to non-bower builders were

comparable to those found between the Malawi rock and sand lineages (Figure 5A).

Phylogenetic visualization revealed that hindbrain diversification within Lake

Tanganyika was largely limited to the Ectodini tribe that has a number of bower building species

(Figure 5B). We found that in this phylogenetic grouping of bower builders, in contrast to Lake

Malawi cichlids, there was little hindbrain evolution independent of the Lake Tanganyika

phylogeny (ANOVA (F = 9.79, 1 d.f., p = 0.12). Similarly, we identified significant amounts of

phylogenetic signal in hindbrain volume (K = 0.53, p = 0.035; λ = 0.65, p = 0.048).

Analyses of diversification rates using BAMM showed hindbrain diversification has

increased within the Lake Tanganyika Ectodini tribe as compared with non-Ectodini species

(Figure 5C-E). Our data support a model of parallel diversification of hindbrain volume

associated with bower building in both Lake Malawi and Lake Tanganyika, but with

substantially different evolutionary histories associated with the trait in each lake.

Discussion

We show here that hindbrain volume increases in tandem with the behavioral evolution

of castle-type bower building, both within the dozens of species sampled from Lake Malawi here

and also as compared to species in Lake Tanganyika but not Victoria, which seems to contain no

known builders. Despite previous results identifying mosaic evolution of the brain, whether this

may occur repeatedly given common ecological, evolutionary, or phylogenetic pressures is

unclear. Our findings suggest that in certain scenarios brain evolution may proceed in a

predictable manner, as previously proposed for systems such as bony fish [10] and vocal learning

birds and mammals [8], but has rarely has this been tested explicitly in a controlled phylogenetic

context. However, resolving phylogenetic relationships among Malawi cichlid species is difficult

given their close genetic relationships and complex histories of gene flow and incomplete lineage

sorting [14]. Therefore, future work attempting to disentangle further the differential roles of

ecology, behavior, and evolution on brain morphology in Lake Malawi should include wider,

more targeted sampling from the phylogeny in addition to comprehensive collection of

demographic traits for each species. Sequencing more Malawi species would allow useful

phylogenomic analyses of genetic association and identification of the roles of gene flow and/or

incomplete lineage sorting in the context of brain evolution.

Our results also indicate that variation in hindbrain size is associated specifically with

diversification of the vagal lobe, a key gustatory region of the fish hindbrain. The vagal lobe

differs significantly between pit and castle species in size and structure, but not connectivity.

This contrasts starkly with observations in cyprinid fish, another teleost clade with extensive

vagal lobe diversification. Among typical food-sorting cyprinids that have a palatal organ, such

as goldfish and carp, the vagal lobe is highly laminated with clear motor and sensory zones that

extend dorsally in parallel along the deep and superficial layers of the lobe [24, 25, 27].

Similarly, the vagal lobe of *Heterotis niloticus,* an unrelated fish of the order *Osteoglossiformes,*

has a laminated vagal lobe with sensory layers overlying motor layers [28]. In fish species that

lack an elaborate palatal organ, such as catfish and zebrafish, the vagal lobe is cytologically

simpler with less obvious lamination and a motor nucleus which is restricted to a sub-ventricular

area with little penetration into the lobe itself [29]. Malawi cichlid hindbrain diversification is an

intermediate between the vagal lobes of food sorting and non-food sorting cyprinids. While the

vagal motor nuclei of pit and castle species is similar in location to that of catfish, histology and

DiI labeling indicate that castle building species tend to show increases in vagal lobe size,

lamination, and manner of termination of primary vagal sensory fibers. These patterns of vagal

lobe elaboration are supported by the presence of palatal organs in just the castle building species

sampled. Furthermore, our studies of immediate early gene expression and ps6 abundance during

behavior reveal not only that the vagal lobe is involved in both bower building and feeding, but

that there are detectable species differences in neuronal activity during bower building. Such

differences may merely reflect differences in the degree to which oral manipulation of substrate

is necessary for the different types of bower: hole versus structure.

Evidently, the largest differences in hindbrain structure and function among bower

builders are determined mostly by activity (via behavioral state) and size, but only moderately by

changes in connectivity and histological structure. Any functional differences in the vagal lobe

associated with pit and castle bower types, then, likely arise through both variations in

developmental patterning and, in adulthood, modulation of vagal lobe function. It does not

appear that bower building is associated with the evolution of novel, behavior-specific hindbrain

circuits. Support for this comes from previous work showing that ecologically-relevant

differences in forebrain size among rock- and sand-dwelling Malawi cichlids arise from a

common blueprint that is differentially modified by patterning genes [30]. Similarly, the

hindbrains of the sand-dwelling cichlids analyzed here appear to have a relatively conserved

brain bauplan at base, but with substantial elaboration in size and modulation among species.

The high degree of relatedness and recent evolution of Malawi cichlid species likely constrains

the phenotypic possibilities available for the evolution of brain and behavior. The significant

increase in diversification rate of the hindbrain among castle-builder clades in both Malawi and

Tanganyika suggests that behavioral evolution in these lakes is likely supported by neural

variation producing bigger brain structures that generate more activity.

The degree of phenotypic predictability displayed by the hindbrain in Malawi cichlids

suggests that, given a common phylogenomic context, diversifying species may present some

degree of commonality in the evolution of neural and behavioral traits in response to similar

pressures. Of particular interest will be the study of how evolution acts on conserved and novel

genes and genetic networks to regulate the brain and behavior across evolutionary distances,

potentially revealing common principles of the evolution of behavior.

Methods

Fish were bred, housed, and maintained at Stanford University following established

Stanford University IACUC protocols.

Volumetric analyses were performed using measurements from a phenotypic data set

comprising brain measurements for 189 cichlid species from East Africa and Madagascar [19].

Phylogenetic comparisons were conducted in R [38]. The packages SNPhylo [31] and ape [32]

were used to construct ultrametric phylogenies for the Lake Malawi species from whole genome

SNP data [21]. A previously published phylogeny was used to infer phylogenetic relationships of

the Lake Victoria and Tanganyika species [35]. Phylogenetic ANOVAs were performed with the

package geiger [38] while phylogenetic signal and trait diversification rates were inferred from

the packages phyloSignal [40] and BAMMtools [23], respectively.

Oropharyngeal anatomy was assayed by scanning electron microscopy of the oral cavities

of *C. virginalis* (CV), *M. conophoros* (MC), and *T. brevis* (TB) using a Hitachi S-3400N VP

scanning electron microscope. Taste buds and innervation patterns were identified by

immunocytochemical labeling using antisera against human calretinin (CR 7697; AB_2619710)

and acetylated tubulin (Sigma T7451; AB_609894).

DiI tracing was performed on CV, MC, and TB by placing DiI crystals on either the

vagus nerve root or the surface of the vagal lobe. After a diffusion period of 1-6 months the

brains were sections at 75-100 um on a vibratome and then imaged.

Behaviorally-induced neural activity was assessed by running CV and MC males through

one of three behavioral paradigms: bower building, feeding, and control. For ISH we used RT-

PCR to amplify a portion of coding sequence from *cFos* (NM_001286320), and subcloned

products into pCR-TOPO4 (Life Technologies). *cFos* forward, 5'-AAT TGG ATC CAA GCC

CAG ATC TTC AGT GG-3'; *cFos* reverse, 5'-AAT TGA ATT CAT AGC CCT GTG ATC

GGC AC-3'. Abundance of pS6 was inferred by immunohistochemical labelling with with rabbit

anti Phospho-S6 Ser244 Ser247 (ThermoFIsher Scientific 44-923G, RRID AB_2533798). ISH

and immunohistochemical labeling were imaged in FIJI and quantified with custom scripts in R.

Additional information can be found in the supplemental experimental methods.

Vertebrate animal use

All animal experiments were approved by the Stanford University Animal Care and Use

Committee (protocol number APLAC-28757).

Data availability

All sequencing data in support of the findings of this study have been deposited in the Short

Read Archive under accessions SRR6314224, SRR6314225, SRR6314226, SRR6314228,

SRR6314230, SRR6322515. All morphometric, phylogenetic, and histological data are available

as electronic supplements to this manuscript.

Declaration of interests

The authors declare no competing interests.

Acknowledgements

We thank Beau Alward, Austin Hilliard, Patrick McGrath, and Hunter Fraser for helpful

comments on earlier drafts of the manuscript. We thank Robert Huber for sharing raw brain

measurement data and Catherine Wagner for the Tanganyika and Victoria species phylogeny.

This research was supported by awards from the NIH to RDF and JTS (R01GM101095), RDF (

NS034950), from the Stanford Bio-X Bowes Fellowship (RAY) and a Stanford CEHG

Fellowship (RAY). The authors have no competing interests.

Contributions

RY conceived of the study, designed the study, carried out behavioral and molecular

experiments, performed all phylogenetic and statistical analyses, and wrote the manuscript. AB

performed behavioral assays and *in situ* hybridization and immunohistochemical staining

protocols. KA, CP, and JTS collected and processed tissue samples for next generation

sequencing and identified genetic variants for phylogeny construction. TF performed the

neuroanatomical comparisons of Malawi cichlid hindbrain structure and helped draft the

manuscript. RDF provided funding, aided in design and coordination of the study and helped

write the manuscript.

Fig 1. Evolution and diversification of hindbrain volume among Malawi cichlids.

(A) Scatterplot comparing standard length and hindbrain volume for castle building species (green) and pit digging species (yellow). **(B)** Maximum likelihood phylogeny for 19 Lake Malawi sand and rock species. Hindbrain volumes normalized by standard length are represented by dots (castle = green, pit = yellow, rock = purple; for species labels and node support see Fig. S2). **(C)** Representative photos of whole brain and hindbrain (indicated by arrows) size for the castle species *Tramitichromis brevis* and the pit species *Copadichromis virginalis*. **(D-F)** Normalized hindbrain volume diversification rates for rock and sand **(D)**, sand alone **(E)**, and rock alone **(F)**.

Fig 2. Oral cavity and brain morphology of *Mchenga conophoros* and *Copadichromis*
***virginalis.* (A)** The dorsal oral cavity of the pit-digger *Copadichromis virginalis* is comparatively
simple and lacks a distinct palatal organ. **(B)** The dorsal oral cavity of the castle-builder *Mchenga*
conophoros displays a more convoluted structure and caudal palatal organ labeled 'PO'. **(C)**
Photograph of *in situ* caudal brain structures of *C. virginalis*. The approximate location of the vagal
lobe is outlined and labeled 'VL' and the cerebellum is labeled 'Cblm'. **(D)** Photograph of *in situ*
caudal brain structures of *M. conophoros*. Like in **(C)** the approximate location of the vagal lobe
is outlined and labeled 'VL' and the cerebellum is labeled 'Cblm'.

Fig 3. Vagus nerve termination. Composite micrographs comparing the laminar pattern of termination of vagus nerves in the vagal lobes of *C. virginalis* and *M. conophoros*. The sensory roots of the vagus nerve, labeled by DiI and rendered in magenta, terminate similarly in the two species: in superficial and deep layers leaving the central region void of terminals but crossed by fascicles running from the superficial root into the deeper layers. Green =Nissl staining inverted and rendered in green as if a fluorescent Nissl stain; magenta = diI label; image superimposed on the image of Nissl staining from another specimen.

512   **Fig 4. Molecular correlates of vagal lobe activity associated with bower building. (A)** Barplot

513   comparing *cFos* mRNA staining in MC across control, feeding, and building conditions across the

514   VL. *$P < 0.05$. **(B)** Barplot comparing *cFos* mRNA staining in CV across control, feeding, and

515   building conditions across the VL. *$P < 0.05$. **(C)** Barplot comparing pS6 positive cell abundance

516   during bower building in *C. virginalis* and *M. conophoros*. *$P < 0.05$.

Fig 5. Evolution and diversification of hindbrain volume across East African Cichlids.

(A) Violin plots of normalized hindbrain volumes for Malawi rock-dwelling species (n=47),

Lake Tanganyika non-bower building species (n=14), Lake Victoria species (n=55), Malawi

sand-dwelling species (n=37), and Tanganyika bower building species (n=12). Lakes are

represented by cartoon outlines accompanying their respective violin plots. **(B)** Maximum

likelihood phylogeny of Lake Tanganyika species sampled with accompanying normalized

hindbrain volumes. Red represents the Ectodini tribe, yellow represents non-Ectodini, stars

indicate known bower building species (for species and node labels see Fig. S10) **(C-E)**

Normalized hindbrain volume diversification rates for Ectodini and non-Ectodini **(C)**, Ectodini

(D), and non-Ectodini **(E)**. **(F)** Boxplot comparing per-species normalized hindbrain volume

diversification rates between Ectodini and non-Ectodini tribes.

Structure	Chi-squared statistic (H)	*p*-value
Hindbrain	13.874	1.9×10^{-4}***
Telencephalon	0.060	0.807
Cerebellum	0.046	0.830
Olfactory bulb	0.054	0.817
Hypothalamus	0.001	0.975
Optic tectum	0.790	0.374

Table 1. Allometric comparisons of brain structure volume

Results from comparisons of pit and castle species' brain structure volumes (normalized by standard length) using Kruskal-Wallis test. *** indicates $p < 0.001$.

References

1. Zupanc, G. n. K. H. 2004. Behavioral neurobiology : an integrative approach. Oxford University Press, Oxford ; New York.

2. Striedter, G. F. 2005. Principles of brain evolution. Sinauer Associates, Sunderland, Mass.

3. Finlay, B. L. and R. B. Darlington. 1995. Linked regularities in the development and evolution of mammalian brains. Science 268:1578-1584.

4. Barton, R. A. and P. H. Harvey. 2000. Mosaic evolution of brain structure in mammals. Nature 405:1055-1058.

5. Yopak, K. E., T. J. Lisney, R. B. Darlington, S. P. Collin, J. C. Montgomery, and B. L. Finlay. 2010. A conserved pattern of brain scaling from sharks to primates. Proc Natl Acad Sci U S A 107:12946-12951.

6. Hager, R., L. Lu, G. D. Rosen, and R. W. Williams. 2012. Genetic architecture supports mosaic brain evolution and independent brain-body size regulation. Nat Commun 3:1079.

7. Krebs, J. R., D. F. Sherry, S. D. Healy, V. H. Perry, and A. L. Vaccarino. 1989. Hippocampal specialization of food-storing birds. Proc Natl Acad Sci U S A 86:1388-1392.

8. Lefebvre, L., S. M. Reader, and D. Sol. 2004. Brains, innovations and evolution in birds and primates. Brain Behav Evol 63:233-246.

9. Dunbar, R. I. and S. Shultz. 2007. Understanding primate brain evolution. Philos Trans R Soc Lond B Biol Sci 362:649-658.

10. Kotrschal, K., M. J. Van Staaden, and R. Huber. 1998. Fish brains: evolution and environmental relationships. Rev Fish Biol Fisher 8:373-408.

11. Nieuwenhuys, R., H. J. t. Donkelaar, and C. Nicholson. 1998. The central nervous system of vertebrates. Springer, Berlin ; New York.

12. Brawand, D., C. E. Wagner, Y. I. Li, M. Malinsky, I. Keller, S. Fan, O. Simakov, A. Y. Ng, Z. W. Lim, E. Bezault, J. Turner-Maier, J. Johnson, R. Alcazar, H. J. Noh, P. Russell, B. Aken, J. Alfoldi, C. Amemiya, N. Azzouzi, J. F. Baroiller, F. Barloy-Hubler, A. Berlin, R. Bloomquist, K. L. Carleton, M. A. Conte, H. D'Cotta, O. Eshel, L. Gaffney, F. Galibert, H. F. Gante, S. Gnerre, L. Greuter, R. Guyon, N. S. Haddad, W. Haerty, R. M. Harris, H. A. Hofmann, T. Hourlier, G. Hulata, D. B. Jaffe, M. Lara, A. P. Lee, I.

608   MacCallum, S. Mwaiko, M. Nikaido, H. Nishihara, C. Ozouf-Costaz, D. J. Penman, D.

609   Przybylski, M. Rakotomanga, S. C. P. Renn, F. J. Ribeiro, M. Ron, W. Salzburger, L.

610   Sanchez-Pulido, M. E. Santos, S. Searle, T. Sharpe, R. Swofford, F. J. Tan, L. Williams,

611   S. Young, S. Yin, N. Okada, T. D. Kocher, E. A. Miska, E. S. Lander, B. Venkatesh, R.

612   D. Fernald, A. Meyer, C. P. Ponting, J. T. Streelman, K. Lindblad-Toh, O. Seehausen,

613   and F. Di Palma. 2014. The genomic substrate for adaptive radiation in African cichlid

614   fish. Nature 513:375-381.

615 13. Moran, P., I. Kornfield, and P. N. Reinthal. 1994. Molecular Systematics and Radiation

616   of the Haplochromine Cichlids (Teleostei, Perciformes) of Lake Malawi. Copeia:274-

617   288.

618 14. Malinsky, M., Svardal, H., Tyers, A.M., Miska, E.A., Genner, M.J., Turner, G.F., Durbin,

619   R. 2017. Whole Genome Sequences Of Malawi Cichlids Reveal Multiple Radiations

620   Interconnected By Gene Flow. bioRxiv.

621 15. Kocher, T. D. 2004. Adaptive evolution and explosive speciation: the cichlid fish model.

622   Nat Rev Genet 5:288-298.

623 16. Streelman, J. T. and P. D. Danley. 2003. The stages of vertebrate evolutionary radiation.

624   Trends in Ecology & Evolution 18:126-131.

625 17. York, R. A., Patil, C., Hulsey, C. D., Anoruo, O., Streelman, J. T., Fernald, R.D. 2015.

626   Evolution of bower building in Lake Malawi cichlid fish: phylogeny, morphology, and

627   behavior. Frontiers in Ecology and Evolution.

628 18. Keenleyside, M., ed. 1991. Cichlid Fishes: Behavior, Ecology, Evolution. Chapman and

629   Hall, London.

630 19. van Staaden, M. J., Huber, R., Kaufman, L. S., & Liem, K. F. 1995. Brain evolution in

631   cichlids of African Great Lakes: brain and body size, general patterns, and evolutionary

632   trends. ZACS 98:165-178.

633 20. Huber, R., M. J. van Staaden, L. S. Kaufman, and K. F. Liem. 1997. Microhabitat use,

634   trophic patterns, and the evolution of brain structure in African cichlids. Brain Behav

635   Evol 50:167-182.

636 21. York, R.A., Patil, C., Abdilleh, K., Johnson, Z.V., Conte, M.A., Genner, M.J., McGrath,

637   P.T., Fraser, H.B., Fernald, R.D., and Streelman, J.T. 2018. Behavior-dependent *cis-*

638   regulation reveals genes and pathways associated with bower building in cichlid fishes.

639   Proc Natl Acad Sci U S A.

640 22. Blomberg, S. P., T. Garland, Jr., and A. R. Ives. 2003. Testing for phylogenetic signal in

641   comparative data: behavioral traits are more labile. Evolution 57:717-745.

642 23. Rabosky, D. L. 2014. Automatic detection of key innovations, rate shifts, and diversity-

643   dependence on phylogenetic trees. PLoS One 9:e89543.

644 24. Morita, Y. and T. E. Finger. 1985. Topographic and laminar organization of the vagal

645   gustatory system in the goldfish, Carassius auratus. J Comp Neurol 238:187-201.

646 25. Morita, Y., T. Murakami, and H. Ito. 1983. Cytoarchitecture and topographic projections

647   of the gustatory centers in a teleost, Carassius carassius. J Comp Neurol 218:378-394.

648 26. Knight, Z. A., K. Tan, K. Birsoy, S. Schmidt, J. L. Garrison, R. W. Wysocki, A.

649   Emiliano, M. I. Ekstrand, and J. M. Friedman. 2012. Molecular profiling of activated

650   neurons by phosphorylated ribosome capture. Cell 151:1126-1137.

651 27. Herrick, C. J. 1905. The central gustatory paths in the brains of bony fishes. J Comp

652   Neurol Psycho 15:375-456.

653 28. Braford, M.R. 1986. De gustibus non est disputandem: a spiral center for taste in the

654   brain of the teleost fish, Heterotis niloticus. Science 232:489-491.

655 29. Finger, T. E. 2008. Sorting food from stones: the vagal taste system in Goldfish,

656   Carassius auratus. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 194:135-

657   143.

658 30. Sylvester, J. B., C. A. Rich, C. Yi, J. N. Peres, C. Houart, and J. T. Streelman. 2013.

659   Competing signals drive telencephalon diversity. Nat Commun 4:1745.

660 31. Lee, T. H., H. Guo, X. Wang, C. Kim, and A. H. Paterson. 2014. SNPhylo: a pipeline to

661   construct a phylogenetic tree from huge SNP data. BMC Genomics 15:162.

662 32. Paradis, E., Claude, J., and Strimmer, K. 2004. APE: analyses of phylogenetics and

663   evolution in R language. Bioinformatics 20:289-290.

664 33. Ivory, S. J., M. W. Blome, J. W. King, M. M. McGlue, J. E. Cole, and A. S. Cohen. 2016.

665   Environmental change explains cichlid adaptive radiation at Lake Malawi over the past

666   1.2 million years. Proc Natl Acad Sci U S A 113:11895-11900.

34.   Sturmbauer, C., S. Baric, W. Salzburger, L. Ruber, and E. Verheyen. 2001. Lake level

fluctuations synchronize genetic divergences of cichlid fishes in African lakes. Mol Biol

Evol 18:144-154

35.   Wagner, C. E., L. J. Harmon, and O. Seehausen. 2012. Ecological opportunity and sexual

selection together predict adaptive radiation. Nature 487:366-369.

36.   Konings, A. 2015. Tanganyika Cichlids in their Natural Habitat: 3rd Edition. Hollywoord

Import & Export Inc., Florida.

37.   Konings, A. 2007. Malawi Cichlids in their Natural Habitat, 4th Edition. Cichlid Press.

38.   Harmon, L. J., J. T. Weir, C. D. Brock, R. E. Glor, and W. Challenger. 2008. GEIGER:

investigating evolutionary radiations. Bioinformatics 24:129-131.

39.   Keck, F., Rimet, F., Bouchez, A., and Franc, A. 2016. phylosignal: an R package to

measure, test, and explore the phylogenetic signal. Ecology and Evolution 6(9):2774-

2780.

40.   Danecek, P., Auton, A., Abecasis, G. Albers, C.A., Banks, E., DePristo, M.A.,

Handsaker, R., Lunter, G., Marth, G., Sherry, S.T., et al. 2011. The variant call format

and VCFtools. Bioinformatics 27(15):2156-2158.

41.   Browning, S.R., and Browning, B.L. 2007. Rapid and accurate haplotype phasing and

missing data inference for whole genome association studies by use of localized

haplotype clustering. Am J Hum Genet 81:1084-1097.

42.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. 2012. Fiji: an open-source

platform for biological-image analysis. Nat Methods 9:676-682.

43.   Schneider, C.A., Rasband, W.S., and Eliceriri, K.W. 2016. NIH Image to ImageJ: 25

years of image analysis. Nat Methods 9(7):671-675.

44.   Mckaye, K.R., Stauffer, J.R., Turnger, G.F., Konings, A., & Sato, T. 2001. Fishes, as

well as birds, build bowers. J Aquaricult Aquat Sci 9:121-129.

45.   Konings, A. 2015. Tanganyika Cichlids in their Natural Habitat: 3rd Edition, (Florida:

Hollywoord Import & Export Inc.).

46.   Stauffer, J.R., and Konings, A. 2006. Review of Copadichromis (Teleostei: Cichlidate)

with the description of a new genus and six new species. Ichthyol. Explor. Freshwaters

7(1):9-42.

47.   Kellog, K.A., Markert, J.A., Stauffer, J.R., and Kocher, T.D. (1998). Intraspecific brood

mixing and reduced polyandry in a maternal mouth-brooding cichlid. Behavioral Ecology

9(5), 309-312.

48.   Kidd, M.R., Kidd, C.E., and Kocher, T.D. (2006). Axes of differentiation in the bower-

building cichlids of Lake Malawi. Mol Ecol 15(2), 459-478.