Evolutionary Models of Amino Acid Substitutions Based on the Tertiary Structure of their Neighborhoods.

Elias Primetis[1,2], Spyridon Chavlis[2], and Pavlos Pavlidis[3]

[1]University of Crete, Department of Biology
[2]Foundation for Research and Technology, Hellas, Institute of Molecular Biology and Biotechnology, 100 Nikolaou Plastira str., Vassilika Vouton, Heraklion, Crete GR 700 13, Greece
[3]Foundation for Research and Technology, Hellas, Institute of Computer Science, 100 Nikolaou Plastira str., Vassilika Vouton, Heraklion, Crete GR 700 13, Greece

Abstract

Intra-protein residual vicinities depend on the involved amino acids. Energetically favorable vicinities (or interactions) have been preserved, while unfavorable vicinities have been eliminated during evolution. We describe, statistically, the interactions between amino acids using resolved protein structures. Based on the frequency of amino acid interactions, we have devised an amino acid substitution model that implements the following idea: amino acids that have similar neighbors in the protein tertiary structure can replace each other, while substitution is more difficult between amino acids that prefer different spatial neighbors. Using known tertiary structures for α-helical membrane (HM) proteins, we build evolutionary substitution matrices. We constructed maximum likelihood phylogenies using our amino acid substitution matrices and compared them to widely-used methods. Our results suggest that amino acid substitutions are associated with the spatial neighborhoods of amino acid residuals.

1 Introduction

Structure and functionality of a protein are closely related (Worth et al., 2009). Its amino acid sequence as well as cofactors, ligands and other parts of the same or other proteins form a complex network of interactions that is the basis of the unique physicochemical properties of each protein related to its function (Worth et al., 2009). During the last decade, a multitude of tertiary protein structures have been determined by using techniques such as crystallography, NMR, electron microscopy and hybrid methods (Egli, 2010). Consequently, the number of the available tertiary structures in the RCSB-PDB database (Bernstein et al., 1977) amino acid is rapidly increasing.

Computational methods facilitate structural, functional and evolutionary characterization of the proteins (Nath Jha et al., 2011). Protein function is tightly linked to its tertiary structure, and consequently to the vicinities, or interactions, amino acid residues have formed. For example, globular and membrane

proteins are characterized by totally different physicochemical environments. Membrane proteins show a limited interaction with water molecules, and, in contrast, they are able to interact with the lipid bilayer. Thus, their trans-membrane region adopts a single type of secondary structure – either a helix or a beta-sheet. Largely, the secondary structure is defined by non-neighboring (on the sequence level) amino acid residue interactions (Nath Jha et al., 2011).

Recently, based on the ideas of (Nath Jha et al., 2011), we developed PrInS (freely available from `http://pop-gen.eu/wordpress/software/prins-protein-residues-interaction-statistics` (Protein Interaction Statistics; Pavlidis et al., unpublished) an open-source software to score proteins based on the frequency of their residue interactions. PrInS uses protein structures stored in Protein Data Bank (PDB) to construct a statistical model of intra-protein amino acid residue interactions for a certain class of proteins (e.g., membrane proteins). PrInS scores every amino acid a proportionally to the number of 'unexpected residues' that interact with it. The term 'unexpected residues' means residues that they are rarely found to be in the vicinity of a if we consider all protein structures of a given dataset. Therefore, it is able to pinpoint residues characterized by a large number of 'less frequent' interactions, and therefore they may represent functional areas of the proteins or targets of natural selection based on the following assumption: even though a large number of unlikely interactions characterizes these amino acids, nature has preserved them. Here, we use PrInS to describe statistically the intra-protein amino acid interactions and consequently to construct an amino acid substitution matrix endowed with the principle that amino acids with similar (residual) neighborhoods can substitute each other during evolution.

Protein evolution comprises two major principles. The first principle suggests that protein structure is more conserved than the sequence (Siltberg-Liberles et al., 2011). The second principle suggests that the physicochemical properties of amino acids constrain the structure, the function and the evolution of proteins (Hatton and Warr, 2015). Protein evolutionary rate is strongly correlated with fractional residue burial(Siltberg-Liberles et al., 2011). This is due to the fact that the core of a protein is mostly formed by buried residues, which often play a crucial role in the stability of the folded structure (Franzosa and Xia, 2008). The three-dimensional structure of the protein determines its evolutionary rate, since most mutations in the core of a protein tend to destabilize the protein.

Within protein families the backbone changes are infrequent, thus, preserving the folding properties over relatively long evolutionary distances, while substitutions are found often at the side chains. For example, for proteins with binding function, the binding interface is under functional constraint and may evolve the slowest, with differences in rate between affinity-determining and specificity-determining residues (Siltberg-Liberles et al., 2011). In addition, the secondary structural elements of a protein evolve at different rates. Beta sheets evolve more slowly than helical regions and random coils evolve the fastest (Siltberg-Liberles et al., 2011). Secondary structure changes may eventually occur due to varying helix/sheet propensity. Some of these changes in secondary structural composition may be evolutionary

neutral, whereas some structural transition may involve negative or positive selection. In the latter case, a new mutationally accessible fold may enable the development of a new favorable function that was not possible within the previous fold (Siltberg-Liberles et al., 2011).

In the context of folding, the thermodynamic stability of the proteins with a stable unique tertiary structure is important. Thermodynamic stability is maintained throughout evolution despite the desta­bilizing effect of the non-synonymous mutations, which are often removed from the populations as a result of negative selection. The protein structure is important because it acts as a scaffold for properly orientating functional residues, such as a binding interface and a catalytic residue. As a result, the selec­tive pressure for particular sequences (and not structures) over longer evolutionary periods is decreased, generating a neutral network of sequences interconnected via mutational changes (Siltberg-Liberles et al., 2011).

Choi and Kim (Choi and Kim, 2006), based on the most common structural ancestor (CSA), showed that not all present-day proteins evolved from one single set of proteins in the last common ancestral organism, but new common ancestral protein were born at different evolutionary times. These proteins are not traceable to one or two ancestral proteins, but they follow the rules of the "multiple birth model" for the evolution of protein sequence families (Choi and Kim, 2006).

In this study we used the scoring matrices that are obtained from PrInS algorithm to examine the evo­lution of proteins and we focused on the evolution of α-helical membrane proteins. The main hypothesis is that protein evolution is related to the three dimensional neighborhoods of amino acids. Specifically, amino acids that have the same amino acid neighbourhoods can substitute each other during evolution.

Methods

Dataset retrieval and name conversion

In eukaryotes, α-helical proteins exist mostly in the plasma membrane or sometimes in the outer cell membrane. In prokaryotes, they are present in their inner membranes. We used 82 α-helical membrane proteins, which were also scrutinized previously by Nath Jha et al. (2011) to describe the statistical properties of amino acid interactions within α-helical membrane proteins. First, for each of the protein in the dataset, we downloaded multiple sequence alignments of homologous protein sequences from the UCSC Genome Browser (Kent et al., 2002) for 16 primate and 3 non-primate mammalian species. Second, three-dimensional structures were downloaded from the Protein Data Bank (PDB) database.

The PrInS software

We applied PrInS on the three dimensional structures downloaded from PDB. PrInS constructs a scoring matrix M as follows: If the tertiary structure of a protein is represented by P and the total number of

amino acid residues is l, then residues $1 \leq k, m \leq l$ interact if and only if:

$$A_{km} = \begin{cases} 1, & \text{if } d(C_\alpha^k - C_\alpha^m) \leq 6.5\text{Å} \\ 0, & \text{otherwise} \end{cases} \tag{1}$$

where $d(C_\alpha^k - C_\alpha^m)$ denotes the distance between the C_α atoms of the k and m amino acids. In other words, the k^{th} and the m^{th} residues of a protein interact if and only if the Euclidean distance between their C_α atoms is less than 6.5Å and they are not located on adjacent positions on the amino acid chain.

Defining the environment of each amino acid

According to Nath Jha et al. (2011), amino acids of a HM protein can be classified in three environments: The first environment comprises amino acids at different distances from the lipid bilayer. The second environment classifies the helices (and thus, their amino acids) on the basis of their inter-helical interactions inside the membrane. Finally, the third environment classifies amino acid residues based on the number of interactions they make. In our study we have used the third type of environments (residue-contact-based interaction) based on the results from Nath Jha et al. (2011), who demonstrated that the residue-contact-based environment description of residue interaction is more accurate for the α-helical proteins they studied. Thus, every amino acid residue P_i in protein P can be assigned to the pair (A, K), where A indexes the amino acid type of the k^{th} residue (e.g. Alanine) and K represent the environment of the k^{th} residue. Based on the results of Nath Jha et al. (2011), we used the number of non-covalent contacts each amino acid makes to define its environment. Thus, environment I comprises all amino acids with 1-5 contacts, environment II, amino acids with exactly 6 contacts and environment III amino acids with more than 6 contacts.

Initially, PrInS was used to construct a matrix M, a 60×60 matrix (or equivalently nine 20×20 scoring matrices), that scores amino acid interactions in all environment pairs. For example, $M_{i,j}, i, j \leq 20$ describe score interactions between amino acids of the environment I. In M the pairs of amino acids with the lower scores are those that interact frequently. In general, the interaction between amino acid A ($1 \leq A \leq 20$) and amino acid B ($1 \leq B \leq 20$) that belong to the environments K ($1 \leq K \leq 3$) and Q ($1 \leq Q \leq 3$), respectively, is given by:

$$M_{ij} = -ln\left(\frac{n_{A,K-B,Q}}{g \times (S_A/S) \times (S_B/S) \times E_{K,Q}}\right) \tag{2}$$

.

The coordinates i and j are given by $i = 20(K - 1) + A$ and $j = 20(Q - 1) + B$, respectively. The parameter $g = 2$ if the contacting pair comprises different amino acids in the same environment ($A \neq B, K = Q$), and $g = 1$, otherwise. S_A and S_B are the total number of amino acids A and B in

the dataset, respectively. S is the total number of amino acids in the dataset and $E_{K,Q}$ equals to the number of interactions between environment K and Q. Finally, $n_{A,K-B,Q}$ is the number of interactions between amino acids A in environment K and amino acid B in environment Q. It is evident that $M_{i,j}$ assumes large (positive) values when the observed number of interactions $n_{A,K-B,Q}$, between A and B, is much lower than the interactions expected based on the total frequency of the amino acids A and B in the dataset and the number of interactions between the K and Q environments. In other words the value of M_{ij} is large for rare interactions.

Distance matrix

The M scoring matrix was split in nine 20×20 matrices and distance metrics (Euclidean, Manhattan and Pearson Squared) between the elements of each matrix (amino acids) were calculated. Let G_d represent the average 20×20 distance matrix for the distance metrics d ($d \in$ {Euclidean, Manhattan, Pearson Squared}). We name these matrices **n**eighborhood **E**uclidean-**d**istance based (nEd), nMd and nPd, respectively. For comparison purposes, we created an additional 20×20 distance matrix based on the Hamming distance between a pair of codons showing the minimum number of nucleotide changes required to transform one amino acid to another. Smaller distance values indicate higher similarity between the respective amino acids. Distance matrices have been visualized as heatmaps (**Supplementary Files S1a, S1c and S1d**). Two amino acids that have similar neighborhoods, *i.e.* similar amino acid neighbors in the tertiary structure will, thus, have a small distance (high similarity) between them. Consequently, according to our hypothesis, they will substitute each other during evolution at a higher rate.

Substitution rate matrix

Each of the four distance matrices were transformed into rate matrices D_r by using a similar procedure as in Dayhoff et al. (1978). In particular, the following procedure was followed:

1. Find the maximum value of the distance matrix D_g, D_{max}.

2. Subtract each matrix element from the D_{max} and divide by D_{max}. This will transform D_g to an amino acid "similarity" symmetric matrix, where 1 denotes maximum 'similarity' between two amino acids.

3. Similar to the substitution matrices BLOSUM62 and PAM120, one element of the matrix is defined as a reference. All other elements are scaled proportionally to the reference.

4. The diagonal of the matrix is defined as the negative sum of all other elements of the respective row.

If an amino acid pair is characterized by a large (relative) substitution rate, then they can substitute each

other during evolution at a high rate. The substitution rate matrix that is derived with the aforementioned algorithm is symmetric.

Evaluation of PrInS ability to predict amino acid substitutions

For every multiple alignment, a substitution matrix S was created from the alignment, by counting the occurrences of the two most common amino acids for every alignment site, and assuming that a substitution between these two amino acids has occurred during evolution. To evaluate the relation of our neighborhood-based substitution rate matrices D_r and the multiple sequence alignments, we calculated the Pearson correlation coefficients between S and D_r (for each distance metric). High Pearson correlation coefficient values suggest that the amino acid substitutions in the alignments are concordant with the neighborhood-based substitution rates distances.

Moreover, we manually analyzed the amino acid pairs with the lowest distance values in the Euclidean, Manhattan and Pearson distance matrices with the tool 'Common Substitution Tool' in the 'Amino Acid Explorer' (`https://www.ncbi.nlm.nih.gov/Class/Structure/aa/aa_explorer.cgi`) of NCBI (Bulka et al., 2006). Common Substitution Tool sorts an amino acid list from the most common to the least common substitutions, given an amino acid as input, based on the BLOSUM62 substitution matrix (Henikoff and Henikoff, 1992).

Overall and Site Likelihoods

Likelihood is a function of the parameters of a statistical model for given data. Since substitution rate matrices are part of the model, we compared the per site and the overall likelihood values obtained by our substitution matrices and those obtained by using BLOSUM62 and PAM120. Both the phylogeny and the equilibrium frequencies of the amino acids were considered known in this analysis. Thus, by keeping the remaining parameters the same for both models, we aimed to assess which rate matrix can better explain the observed multiple alignments. As overall likelihood, we defined the total likelihood of the alignment, while a site likelihood is the likelihood of a single position in the amino acid alignment. The phylogenetic tree was retrieved from the UCSC Genome Browser (Kent et al., 2002), and the equilibrium frequencies of amino acids were obtained from the equilibrium frequencies in the BLOSUM62 model. We used the same phylogenetic tree and equilibrium frequencies for all comparisons.

Results

Evaluation of PrInS Ability to Predict Amino Acid Substitutions

The substitution matrix counts the amino acid substitutions that occurred in the 224 downloaded multiple alignments using the two most frequent amino acid residues at each alignment site. For each amino acid in

this matrix, we evaluated its correlation (Pearson correlation coefficient) with the same amino acid in the distance matrices. Thus, if, for a given amino acid, both matrices suggest similar substitution preferences, then neighborhood-based substitution models are concordant with the observed substitutions. Since we used distance matrix in this analysis, negative correlation coefficient indicate concordance. Results suggest that there is concordance between the substitution matrix and our neighborhood-based distance matrices for most of the amino acids. Figure 1 shows the Pearson correlation coefficients for amino acids using the substitution matrix and the *nEd*. The correlation plots between the substitution matrix and other distances are illustrated in the Supplementary material (Figures S2a-S2c).

Pearson Correlation Coefficient: Consensus–Euclidean

Figure 1: Pearson Correlation Coefficient between the substitution and *nEd*. Concordance is indicated with negative values, while discordance with positive values.

As it is shown in Figure 1, most of the amino acids are concordant when we compare the Euclidean distance matrix and the substitution matrix. For example, both the uncorrelated and discordant amino acids were glycine, proline, arginine, serine, tryptophan and tyrosine.

In the Supplementary Figure S2b (substitution vs Manhattan distance), the discordant amino acids were the five out of six amino acids of Figure 1. Similar results are shown in Supplementary Figure S2c, where the substitution matrix is compared against the squared Pearson distance matrix. Finally, there are no discordant amino acids in the correlation between the Consensus matrix and the Genetic Code matrix (Supplementary Figure S2a).

Furthermore, the amino acid pairs with the lowest values in the distance matrices are at the top of the amino acid list that the Common Substitution Tool of NCBI Amino Acid Explorer reports. This means that amino acids with similar three dimensional neighborhoods tend to substitute each other. For example, in Euclidean distance matrix, the Leukine-Isoleukine pair has the lowest distance value and according to the substitution list of Amino Acid Explorer, Isoleukine is the most common substitution

of Leukine. Figure 2 shows the first five most common amino acid substitutions for Leucine according to 'Common Substitution Tool' (Figure 2a) and according to the Euclidean distance matrix (Figure 2b). Evidently, the Euclidean distance matrix can predict the most common substitutions of Leucine, but in slightly different order. For example, Common Substitution Tool reports the following order of amino acids: Isoleucine, Methionine, Valine, Phenylalanine and Alanine. According to the Euclidean distance matrix the order is: Isoleucine, Phenylalanine, Valine, Tryptophan and Tyrosine. Methionine is the sixth most common substitution of Leucine, while Alanine is the tenth most common substitution of Leucine according to the Euclidean distance matrix. On the other hand, Tryptophan and Tyrosine are the seventh and the eighth most common substitutions of Leucine according to the Common Substitution Tool. Neglecting the order of the first five substitutions, drawing five amino acids and observing three common is marginally significant (p-value = 0.07, right tail of hypergeometric distribution). Differences between two reports are possibly due to the fact that Euclidean distance matrix is solely based on the structural information of a protein family, while the Common Substitution Tool is based on BLOSUM62 substitution matrix that was created using alignments of multiple protein families.

(a)

(b)

Figure 2: The first five most common substitutions of Leucine according to (a) Common Substitution Tool and (b) Euclidean Distance Matrix. The same color code is used in both figures. Dark green shows the first most common substitution, while the red shows the fifth most common substitution. Methionine and alanine are not found among the first five most common substitutions of Leucine in Euclidean distance matrix, while Tryptophan and Tyrosine are not found among the first most common substitutions in Common Substitution Tool.

Pairwise comparison of proteins in different species

Using the Euclidean-based distance matrix, we scored the differences of each protein sequence to its human homologue. Results are presented as a heatmap and are clustered hierarchically based on their distance

from the human homologue (Figure 3). Darker gray tone in Figure 3 denotes high similarity between human and other species homologues, whereas lighter tones suggest lower similarity. As expected, proteins from species that are evolutionarily distant from humans are more dissimilar to human homologues. This is especially true for a group of proteins clustered together using the Euclidean distance-based substitution matrix (Figure 3 proteins labeled with three dashes).

Figure 3: An illustration of the scoring of multiple alignments with the Euclidean distance matrix as a heatmap. Each protein (rows) from human (first column) is compared against the homologous proteins from another species (columns) using pairwise amino acid sequence alignments. We used only the alignment sites with amino acids A and B, where $A \neq B$ and $A \neq -$ and $B \neq -$. The total score (a cell in the heatmap) is calculated as the sum of all site scores. Proteins form two clusters based on the hierarchical tree on the left side of the figure. Proteins of the bottom cluster (also denoted with three dashes are further analyzed using Gene Ontology terms.

The group of proteins clustered together by the Euclidean-based matrix (marked in the lower part of Figure 3) was further scrutinized by Gene Ontology (GO) terms. We used gProfiler (Reimand et al., 2007), to obtain the related GO terms for this set of proteins. These proteins are characterized by a distinct function compared to proteins found in other clusters. On the one hand, the proteins that are located in this cluster play a crucial role in the intestinal absorption of phytosterol and in cholisterol and lipid transportation. On the other hand, the remaining proteins, which are not found in this discrete cluster, are responsible for the homeostatic and transducer/receptor mechanisms of the cells.

Overall and Site Likelihoods

By converting PrInS distance matrices into rate matrices, it was possible to compute the overall likelihood and site likelihoods for every protein alignment. The resulting matrices from the nEd, nMd and $nPSm$ are named as nEs, nMs and $nPSs$, respectively. For most of the proteins, 170 out of 223, the likelihoods computed by BLOSUM62 and PAM120 were better than our neighborhood Euclidean-based substitution matrices. This result is expected since BLOSUM62 and PAM120 substitution matrices were constructed based on alignment files, whereas the presented substitution matrices are alignment-unaware. For the remaining 54 protein alignments, our models and more specifically the Euclidean-based rate matrix resulted in higher likelihoods.

For every protein, we calculated the individual site likelihood for each amino acid site in the multiple sequence alignment using either the BLOSUM62 (or PAM120) or the nEs matrix. Results are illustrated in Figure 4) for the protein NCKX1, where the likelihood difference between BLOSUM68 and nEs is plotted for every amino acid. Thus, negative values are related to sites where the likelihood is greater for the nEs matrix, whereas positive values show sites where BLOSUM62 results in greater likelihoods. For the protein NCKX1, 240 sites are scored with a higher likelihood using the Euclidean substitution matrix derived from PrInS (negative values), whereas 390 sites are scored with a higher likelihood using BLOSUM62 (positive values). The overall likelihood difference for this protein is positive (264.4) indicating that overall BLOSUM62 results in higher likelihood scores. NCKX1 was randomly selected to illustrate the site likelihood differences between the two approaches. It is a critical component of the visual transduction cascade, controlling the calcium concentration of outer segments during light and darkness McKiernan and Friedlander (1999).

Comparison between the average substitution likelihoods for different rate matrices

The 20 amino acids form $\binom{20}{2} = 190$ (unordered) pairs. For each of them, we have calculated the average differences between the BLOSUM62-based and the nEs-based likelihoods. For the calculations, we considered only the sites that consist of two amino acid states. Since all the parameters of the evolutionary model, but the substitution rate matrix are fixed (phylogenetic tree, equilibrium frequencies), then the matrix that results in the highest average likelihood for a certain amino acid pair describes the preferential model for this amino acid pair. Surprisingly, even though most of the protein alignments are scored with a higher likelihood using BLOSUM62, 111 (out of 190) amino acid pairs are characterized by negative average likelihood difference. This indicates that the nEs-based likelihood is greater than the BLOSUM62-based likelihood. The average likelihood is greater for the BLOSUM62 for only 50 pairs, whereas it cannot be assessed for 29 pairs because no occurrences of these 29 pairs were present in the

Figure 4: The differences between the BLOSUM62-based and the Euclidean-based likelihoods for each site of the NCKX1 protein is shown. On the x axis, the protein sites are shown, while on the y axis the differences between the likelihoods are depicted. Positive differences indicate higher likelihoods for the BLOSUM62 rate matrix, whereas negative differences indicate higher likelihoods for the Euclidean rate matrix.

alignment datasets. To scrutinize further this result, we plotted the average likelihood difference as a function of the occurrence frequency of the amino acid pair in the alignment (Figure 5). In Figure 5, it is apparent that the average likelihood difference is positively correlated with the frequency of occurrences of the amino acid pairs ($r = 0.621$, r is the correlation coefficient, CI: (0.55, 0.67), CI denotes the 95% confidence intervals). In other words, the more frequent a substitution is in the alignments, the higher the difference of likelihoods, favoring the BLOSUM62 versus the nEs.

The heatmap in Figure 6 shows all amino acid pairs and the likelihood differences between the BLO-SUM62 and the nEs matrix for all amino acid pairs. Cells denoted by a '-' are characterized by a higher likelihood for the Euclidean-based matrix, whereas a '+' denotes cells with higher BLOSUM62-based likelihood. The darker the color of the cell the higher the difference between the two likelihoods. Cells with an 'o' illustrate pairs where no site with this pair of amino acids was found in any of the alignments. The greatest value for a '-' cell is represented for the 'Arginine-Glycine' pair. This means that the nEs matrix is a preferential model for the specific amino acid pair. On the other hand, the pair with the highest '+' value is the 'Isoleucine-Valine' pair.

Figure 5: The average likelihood difference between the BLOSUM62-based and the Euclidean-based approaches that have been used to calculate likelihoods. As the figure indicates, for the majority of amino acid pairs the likelihood difference is close to 0, *i.e.*, both the Euclidean-based calculations and the BLOSUM62-based calculations result in similar outcomes. Positive values are fewer than negative values (see text), however the magnitude is much greater for positive values than for negative, indicating that BLOSUM62 results in much greater likelihoods than the Euclidean-based approach. Furthermore, as the occurence frequency of the amino acid increases the difference of the likelihood increases as well, suggesting that BLOSUM62 outperforms the Euclidean-based approach for the amino acid pairs that occur frequently, either within the same protein or in different proteins.

Figure 6: Comparison of likelihoods between BLOSUM62 and nEs for all amino acid pairs found in the protein alignments. Boxes with a '-' pinpoint to the specific amino acid pairs where nEs calculates a higher site likelihood. Contrarily, a '+' indicates amino acid pairs that the BLOSUM62 approach results in greater likelihood. Finally, 'o' cells depict the amino acid pairs that were not found in the multiple alignments, thus no comparison was possible. The darker the tone of the cell the greater the difference in the likelihood between the BLOSUM62 and the nEs approach.

Comparison to Grantham's and Sneath's distance matrices

Grantham's distance Grantham (1974) between two amino acids depends on three properties: composition (defined as the atomic weight ratio of noncarbon elements in end groups or rings to carbons in the side chain), polarity and molecular volume. Based on these three properties, distance $D_{G[i,j]}$, between amino acids i and j, is defined as:

$$D_{G[i,j]} = [\alpha(c_i - c_j)^2 + \beta(p_i - p_j)^2 + \gamma(v_i - v_j)^2]^{1/2} \tag{3}$$

where c is the composition, p the polarity and v the molecular volume. The three components of the distance (c, p, v) are not independent. Thus, the constants α, β, γ serve as normalizing factors and they can be calculated as a function of c, v, p Grantham (1974). Grantham (1974) provides the distances of Equation 3 for all amino acid pairs, thus a distance matrix D. Furthermore, Grantham demonstrated that the relative substitution frequency of amino acid pairs and their distances $D_{G[i,j]}$'s are correlated, underlying the physicochemical basis of amino acid substitution.

Similarly to Grantham's amino acid distances, Sneath's index Sneath (1966) takes into account 134 categories of activity and structure. The dissimilarity index $D_{S[i,j]}$ is the percentage of the sum of all properties not shared between two amino acids.

The amino acid neighborhood-based distance is correlated with both the Grantham's and Sneath's distances Figure 7, highlighting the fact that amino acid neighborhoods capture information related to the physicochemical properties of amino acids and also their relative substitution frequency.

Figure 7: Scatterplots between amino acid distances for different distance methods. (A) Euclidean-based distance versus Sneath's index ($r \approx 0.5$). (B) Sneath's index versus Grantham's distance ($r \approx 0.5$). (C) Euclidean-based distance versus Grantham's distance ($r \approx 0.66$).

Discussion

We studied the spatial properties of amino acid substitutions. We investigated if amino acid substitutions can be predicted from their neighborhood tertiary structure. We described statistically the amino acid residual neighborhoods using the software PrInS. Then, we converted PrInS output files to amino acid distance matrices and substitution rate matrices and evaluated their ability to model evolutionary changes.

Correlation analysis between the observed substitution frequencies (from multiple sequence alignments) and the distance matrices indicated that residual neighborhoods capture evolutionary information and thus they can be useful in modeling evolution. In other words, substitutions in multiple sequence alignments can be predicted from the amino acid neighborhoods: residuals with similar neighbors can substitute each other, with a higher rate, during evolution. More specifically, from the three distance matrices (Euclidean, Manhattan, and Pearson Squared) that were compared against the substitution matrix, only five amino acids were discordant in at least two comparisons. These amino acids were proline, arginine, serine, tryptophan and tyrosine. From the substitution matrix, only a few substitutions involve tryptophan and tyrosine, whereas there is a multitude of substitutions involving proline, arginine and serine. In contrast, based on distance matrices, in the columns of tryptophan and tyrosine there are small distance values, while in the proline, arginine and serine columns distances are large. This can explain the discordance of these amino acids in these two matrices, because a large number of substitutions is associated with small distance values for an amino acid.

In addition, the Common Substitution Tool of NCBI Amino Acid Explorer Bulka et al. (2006) enhances our main result: Amino acid pairs with the lower distance values in the distance matrices, especially using the nEd, are found to substitute each other more frequently. Amino Acid Explorer returned a list of the amino acids from the most common to the the less common substitution for each amino acid. There are some differences in the order of the amino acids between the Common Substitution Tool and the distance matrices (e.g Euclidean). These differences are possibly based on the fact that the generation of the nEd matrix is based solely on the structural information of a protein family (α-helical membrane proteins), while the Common Substitution Tool is based on BLOSUM62 substitution matrix that was created from alignments from multiple protein families. Thus, a plausible explanation is that the amino acid neighborhood approach we followed cannot capture adequately well substitutions of certain amino acids. Another plausible explanation is that we followed a linear approach to convert the amino acid distance matrix to the substitution rate matrix. Thus, if the distance between a pair of amino acids A_1, B_1 is d_1, whereas the distance of amino acids A_2 and B_2 is d_2, then the relation between the substitution rates of A_1, B_1 and A_2, B_2 will be a linear function of the d_1 and d_2. A non linear function between the distances and the substitution rates will perhaps result in more accurate substitution rates.

An interesting observation springs from Figure 5 where amino acid substitutions that occur very

frequently in binary sites of alignments (*i.e.*sites with two states) are better predicted with BLOSUM62 than nEs matrix. The relation between the average likelihood difference between BLOSUM62 and nEs is positively correlated with the occurrence frequency of amino acid substitution. Thus, the neighborhood based approach cannot predict well substitutions that occur frequently in an alignment. A plausible solution would be to modify nEs taking into account the frequency of the involved amino acids (which is presumably related to the substitution frequency between them).

The neighborhood distance matrices are highly correlated with the Grantham's and Sneath's distance matrices, illustrating that the formation of amino acid neighborhoods is determined by structural constraints. Grantham Grantham (1974) demonstrated that amino acid substitution is related to their structural properties. Thus, a plausible explanation of the amino acid substitution prediction ability of nEd approach is that neighborhood properties are defined by the structural properties of amino acids which also affect the substitutions between amino acids.

References

Frances C Bernstein, Thomas F Koetzle, Graheme JB Williams, Edgar F Meyer, Michael D Brice, John R Rodgers, Olga Kennard, Takehiko Shimanouchi, and Mitsuo Tasumi. The protein data bank. *The FEBS Journal*, 80(2):319–324, 1977.

Blazej Bulka, Stephen J. Freeland, and others. An interactive visualization tool to explore the biophysical properties of amino acids and their contribution to substitution matrices. *BMC bioinformatics*, 7(1): 329, 2006. URL https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-7-329.

In-Geol Choi and Sung-Hou Kim. Evolution of protein structural classes and protein sequence families. 103(38):14056–14061, 2006. URL http://www.pnas.org/content/103/38/14056.short.

MO Dayhoff, RM Schwartz, and BC Orcutt. 22 a model of evolutionary change in proteins. In *Atlas of protein sequence and structure*, volume 5, pages 345–352. National Biomedical Research Foundation Silver Spring, MD, 1978.

Martin Egli. Diffraction techniques in structural biology. In Serge L. Beaucage, Donald E. Bergstrom, Piet Herdewijn, and Akira Matsuda, editors, *Current Protocols in Nucleic Acid Chemistry*, pages 7.13.1–7.13.35. John Wiley & Sons, Inc., 2010. ISBN 978-0-471-14270-6. URL http://doi.wiley.com/10.1002/0471142700.nc0713s41. DOI: 10.1002/0471142700.nc0713s41.

Eric Franzosa and Yu Xia. Structural perspectives on protein evolution. *Annu Rep Comput Chem*, 4(1): 3–21, 2008.

R Grantham. Amino acid difference formula to help explain protein evolution. *Science*, 185(4154): 862–864, 1974.

Leslie Hatton and Gregory Warr. Protein structure and evolution: are they constrained globally by a principle derived from information theory? 10(5):e0125663, 2015. URL http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0125663.

Steven Henikoff and Jorja G. Henikoff. Amino acid substitution matrices from protein blocks. *Proceedings of the National Academy of Sciences*, 89(22):10915–10919, 1992. URL http://www.pnas.org/content/89/22/10915.short.

W. James Kent, Charles W. Sugnet, Terrence S. Furey, Krishna M. Roskin, Tom H. Pringle, Alan M. Zahler, and David Haussler. The human genome browser at UCSC. 12(6):996–1006, 2002. URL http://genome.cshlp.org/content/12/6/996.short.

Colleen J. McKiernan and Martin Friedlander. The retinal rod na+/ca2+, k+ exchanger contains a noncleaved signal sequence required for translocation of the n terminus. 274(53):38177–38182, 1999. URL http://www.jbc.org/content/274/53/38177.short.

A. Nath Jha, S. Vishveshwara, and J. R. Banavar. Amino acid interaction preferences in helical membrane proteins. 24(8):579–588, 2011. ISSN 1741-0126, 1741-0134. doi: 10.1093/protein/gzr022. URL https://academic.oup.com/peds/article-lookup/doi/10.1093/protein/gzr022.

Jri Reimand, Meelis Kull, Hedi Peterson, Jaanus Hansen, and Jaak Vilo. g:profilera web-based toolset for functional profiling of gene lists from large-scale experiments. 35:W193–W200, 2007. ISSN 1362-4962, 0305-1048. doi: 10.1093/nar/gkm226. URL https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkm226.

Jessica Siltberg-Liberles, Johan A. Grahnen, and David A. Liberles. The evolution of protein structures and structural ensembles under functional constraint. 2(4):748–762, 2011. ISSN 2073-4425. doi: 10.3390/genes2040748. URL http://www.mdpi.com/2073-4425/2/4/748/.

PHA Sneath. Relations between chemical structure and biological activity in peptides. *Journal of theoretical biology*, 12(2):157–195, 1966.

Catherine L. Worth, Sungsam Gong, and Tom L. Blundell. Structural and functional constraints in the evolution of protein families. 2009. ISSN 1471-0072, 1471-0080. doi: 10.1038/nrm2762. URL http://www.nature.com/doifinder/10.1038/nrm2762.