Differential susceptibility to reduced dopamine in SNc versus VTA in a mouse Tyrosine

hydroxylase KO model of Parkinson's Disease

M. Koltun, K. Cichewicz[a], J.T. Gibbs[b], M. Darvas[b], J. Hirsh*

Department of Biology, University of Virginia, Charlottesville, VA 22903, USA

[a]Present address: Center for Neuroscience, UC Davis, Davis, CA, USA

[b]Department of Pathology, University of Washington School of Medicine, University of

Washington, Box357470, Seattle, WA 98195, USA

[*]: Corresponding author

Abstract

Parkinson's Disease (PD), is a neurodegenerative disorder affecting both cognitive and motor functions. It is characterized by decreased brain dopamine (DA) and a selective and progressive loss of dopaminergic neurons in the substantia nigra pars compacta (SNc), whereas dopaminergic neurons in the ventral tegmental area (VTA) show reduced vulnerability. The majority of animal models of PD are genetic lesion or neurotoxin exposure models that lead to death of dopaminergic neurons. Here we use a DAT:TH KO mouse model that by inactivation of the tyrosine hydroxylase (*Th*) gene in dopamine transporter-expressing neurons, causes selective depletion of striatal dopamine without affecting DA neuron survival. We analyzed transcriptome responses to decreased DA in both pre- and post-synaptic dopaminergic brain regions of DAT:TH KO animals. None of the post-synaptic regions analyzed - Dorsal Striatum, Ventral Striatum and Prefrontal Cortex - show an overall change in transcriptome as a function of DA deficiency. This suggests that the broad striatal transcriptional changes in neurodegeneration-based PD models are not direct effects of DA depletion, but are rather a result of DA neuronal death. However, we find a number of dopaminergic genes differentially expressed in SNc, and to a lesser extent in VTA, as a function of DA deficiency, providing evidence for a DA-dependent feedback loop. Of particular interest, expression of Nr4a2, a crucial transcription factor maintaining DA neuron identity, is significantly decreased in SNc, but not VTA, of DAT:TH KO mice. Unexpectedly, we find that half of DAT:TH KO animals show near-normal levels of *Th* transcription, and also display an overall transcript expression pattern more similar to wild type animals than to the remaining DAT:TH KO mice. These animals could represent recruitment of a pathway attempting to compensate for dopamine loss.

Keywords

ventral tegmental area, substantia nigra, dopamine, Parkinson's disease, tyrosine hydroxylase

Introduction

Midbrain dopamine neurons in the ventral tegmental area (VTA) and substantia nigra pars

compacta (SNc) are the major source of DA in the brain through their projections to the striatum,

prefrontal cortex (PFC) and other limbic areas including the amygdala and hippocampus

(Björklund and Dunnett 2007). Via these projections to the striatum and PFC, DA neurons play

crucial roles in reward, motor and cognitive processes (Wise 2004). SNc and VTA DA neurons

have different major projection fields: SNc DA neurons project primarily to the dorsal striatum

whereas VTA DA neurons project to the ventral striatum and to the PFC, hippocampus and

amygdala. Loss-of-function studies using neurotoxins and genetic approaches indicate that VTA

and SNc DA neurons have specific involvement in reward, motor, memory and learning

processes (Darvas et al. from 2008-2014, Baunez and Robbins 1999; Chudasama and Robbins

2006; Da Cunha et al. 2002; De Leonibus et al. 2007; Mura and Feldon 2003). Parkinson's

disease (PD) is characterized by early degeneration of SNc neurons and the presence of alpha-

synuclein inclusion bodies (Lewy bodies), yet only much later is there involvement of the VTA

in PD. Clinical observations in PD patients suggest that loss of SNc DA neurons contribute to

both motor and cognitive symptoms of PD (Cools et al. 2001, 2003; Sawamoto et al. 2008).

The earliest changes induced by alpha synuclein accumulation are predegenerative loss of DA

neurotransmission, followed by axonopathy and subsequent loss of DA neurons (Luk et al. 2012;

Lundblad et al. 2012). Early changes in midbrain neurons resulting from loss of DA may be

involved in the chain of pathologic processes leading to degeneration of DA neurons. We have

generated a genetic model that allows investigation of the effects of DA loss on midbrain gene

expression without degeneration of midbrain DA neurons. Our model lacks tyrosine hydroxylase

(TH) selectively in DA-transporter (DAT) expressing neurons, resulting from a conditional

(floxed) *Th* deletion caused by expression of Cre recombinase in DAT neurons (Henschen,

Palmiter, and Darvas 2013; Jackson et al. 2012; Zhuang et al. 2005). Although these animals

have severe loss of striatal DA and learning and motor deficits, they retain basic locomotor

function, and are normally motivated to drink and eat on their own, with normal viability

(Darvas, Henschen, et al. 2014; Garrett Morgan et al. 2015; Henschen et al. 2013).

We use this model, referred to as DAT:TH KO, to investigate transcriptome changes in response

to DA loss. Exquisite mechanisms maintain DA tone during normal physiology, with regulation

at nearly all steps of DA synthesis, release and reception of the DA signal. These steps include

regulation of synthesis and feedback inhibition of the rate limiting DA biosynthetic enzyme TH,

regulation of size and content of synaptic vesicles, regulation of DA reuptake via DAT,

regulation of D2-like DA presynaptic autoreceptors, and postsynaptic DA receptors. In addition,

there can be direct or indirect effects of classical transmitters that act to modulate DA neurons

(reviewed in Sulzer et al., 2016). With this degree of regulation in normal DA physiology, it is

not surprising that in pathological conditions that cause loss of DA neurons brain physiology is

altered in a compensatory manner. Both in Parkinson's Disease and in Parkinson's Disease

models resulting from agents leading to death of DA neurons, motor symptoms occur only after a

majority of DA neurons are lost (Ballanger et al. 2016; Golden et al. 2013).

This compensation for DA neuron loss is not well understood (Bezard et al. 2001; Bezard, Gross,

and Brotchie 2003; Bezard and Gross 1998; Blesa et al. 2012; Hornykiewicz 1975; Perez et al.

2008). Although many of these reports indicate compensatory modulation of the DA system,

other transmitter systems may also be involved, including striatal preproenkephalin (Bezard et al.

2001), altered cortical-basal ganglia-thalamocortical circuitry (Bezard et al. 2003; Melief et al.

2018), alterations of the serotonergic system, in addition to altered DA signaling and altered

extracortical circuits (Ballanger et al. 2016).

Here we analyze transcriptome responses to DA loss in pre- and postsynaptic brain regions, using

DAT:CRE driven inactivation of *Th*. We find a significant difference in responses to DA loss

between SNc and VTA and a surprising diversity of responses that may represent a compensatory

response to DA loss.

Results

A novel *Th* partial transcript can be detected in DAT:TH KO presynaptic libraries

We used the DAT:TH KO model to test for effects of striatal DA deficiency on the transcriptome

in the regions studied. We expected that the lack of DA will affect preferentially DA producing

and DA receiving parts of the brain. Therefore, we studied transcripts from five distinct brain

regions, two of which are presynaptic: *Substantia nigra pars Compacta* (SNc) and Ventral

tegmental area (VTA); and three postsynaptic: dorsal striatum (DS), ventral striatum (VS) and

prefrontal cortex (PFC), from six wild type and six DAT:TH KO male mice. Samples from each

brain region were sequenced with similar depth of 42.2 ± 4.75 million uniquely mapped reads

per library. The distributions of number of reads mapped per gene in all brain regions followed

the same pattern, with the majority of genes represented by over 1000 unique reads (Figure

S1A). In all the following analyses we filtered out weakly expressed genes as represented by

fewer than 30 reads total in all samples from a given brain region. In the final dataset, a majority

of genes (14,157) was detectable in the samples from all brain regions, but there are 33 to 142

genes that are found in only one region (Figure S1B).

To validate an effective *Th* knockout in DAT expressing neurons of the DAT:TH KO animals,

we examined the expression level of *Th* in the SNc and VTA samples. Unexpectedly, we

detected a high level of *Th* mRNA in SNc and VTA from three DAT:TH KO mice which we

designate as 'DAT:TH KO$^{\text{High}}$' versus 'DAT:TH KO$^{\text{Low}}$' (Figure 1A). Additionally, SNc and

VTA samples from one WT animal have remarkably low levels of *Th* expression. To determine

if this low level of *Th* mRNA correlates with the DAT:TH KO^{Low} samples, we mark samples

from this WT animal with minus sign ("-") in this and subsequent figures.

One could imagine that inefficient removal of the *Th* exon 1 by Cre recombinase by low level of

Cre expression, could explain the observed variation in levels of TH transcripts over the

DAT:TH KO samples. However, we detect very few exon 1 reads in all of the DAT:TH KO

samples (Figure 1B), even in the DAT:TH KO samples with high *Th* expression level (DAT:TH

KO^{High}) which are otherwise of normal downstream structure (Figure 1C).

We used two analytic methods, PCA and hierarchical cluster analysis, to determine whether the

SNc transcriptomes of WT, DAT:TH KO^{High} and DAT:TH KO^{Low} show divergence in their

overall transcriptomes. The PCA and the hierarchical cluster analysis from SNc show that

DAT:TH KO^{High} have overall expression pattern more similar to wild type than to the DAT:TH

KO samples with low level of *Th*. The WT and DAT:TH KO^{High} samples from SNc are localized

closer to the upper-left corner of the PC1 vs PC2 plot, whereas DAT:TH KO samples are

grouped near lower-right corner (Figure 2A, a cluster of samples with low level of *Th* is outlined

with dashed line). This separation is even more visible on plot summarizing the hierarchical

cluster analysis. DAT:TH KO samples with low level of *Th* creates one cluster with the

aforementioned WT sample with the lowest level of Th ("WT-"), whereas DAT:TH KO samples

with high level of Th are grouped with the remainder of the WT samples on the other branch of

dendrogram (Figure 2B).

Lack of striatal dopamine does not alter the overall pattern of gene expression

Upon finding variation between the DAT:TH KO samples, we examined general differences

between brain regions and genotypes by two techniques, principal component analysis (PCA)

(Figure 3A) and hierarchical cluster analysis (Figure 3B). The first principal component in PCA

explains over the half of the variability, and clearly separates pre- and postsynaptic regions. The

second principal component separates SNc, VTA, DS and VS from PFC. On the contrary, WT vs

DAT:TH KO genotype is not recognizable from this analysis, indicating that the lack of DA does

not trigger global transcriptome changes. All samples from both presynaptic regions, SNc and

VTA, are closely localized, independent of genotype, reflecting similarity in their gene

expression profiles. The libraries from DS and VS create another sparse group, and the PFC

samples form an independent group, reflecting their separate origin. However, one sample from

DAT:TH KO PFC (Figure 3A, datapoint indicated by a hollow square) did not localize closely to

the PFC group. The library used to generate this datapoint had only 20% of the expected read

number, with 30% of reads representing poly-G, suggesting a problem with library preparation

or sequencing. None of the other libraries showed these issues, so this library was eliminated

from further consideration. Similar strong grouping by a region of origin is visible in a heatmap

representing results of hierarchical cluster analysis (Figure 3B). Two main branches of the

dendrogram separate pre- and postsynaptic samples, with a further division into samples from

PFC and striatum. As with the PCA, the DAT:TH KO and WT samples do not show clustering.

Moreover, we do not detect any strong batch affect and samples from three different sequencing

batches are clustered in alternating order (color coded in the leaves of the dendrogram in Figure

3B).

Lack of DA results in small transcriptomic effects in postsynaptic brain regions

To address whether DA deficiency affects gene expression in the DS, VS and PFC, we compared

transcriptomes from the DAT:TH KO^{Low} and WT animals using a DESeq2 statistical analysis

based on a negative binomial distribution model. We analyzed the expression level of 14,648,

14,694 and 14,692 genes per brain region, respectively. DS samples have the strongest observed

response to the decreased DA level among the postsynaptic brain parts, with 7 differentially

expressed (DE) genes at a false discovery rate (FDR) < 0.05. We detected only 3 downregulated:

Nefm, *Pdyn* and *Sntg2* (Figure 4A); and only 4 upregulated genes in DS: *Anxa3*, *Fam43a*,

Slc7a8, and *Wdfy* (Figure 4B). Only one of those genes, *Sntg2*, is also differentially expressed in

the ventral striatum of DAT:TH KO^{Low} animals. Additionally, *Gpx6* has a significantly higher

level of expression and *Shisa8* is significantly downregulated in DAT:TH KO^{Low} VS samples.

Finally, we do not detect any DE genes in DAT:TH KO PFC samples.

DAT:TH KO^{Low} show reduced expression of DA related genes in SNc

We next asked whether DA deficiency leads to differential gene expression in the SNc.

Comparing expression level of 15,282 genes from the DAT:TH KO^{Low} and WT animals, we find

34 that are DE at a false discovery rate (FDR) < 0.05 (Figure 5A). The DAT:TH KO^{High} libraries

were not used during the detection of the DE genes since earlier analyses show that these

libraries most resemble WT. However, when we examined the expression level of the DE genes

from the DAT:TH KO^{High} with the DAT:TH KO^{Low} and the WT SNc libraries, DAT:TH KO^{High}

cluster with the WT samples (Figure 5B), indicating that TH expression level is a better predictor

of clustering than genotype. When we directly compared the DAT:TH KO^{High} and the WT SNc

samples, none of the previously detected downregulated genes except for *Lcn2* and *Lrg1* had

significant DE.

All, but one of the DE genes are downregulated in the DAT:TH KO^{Low} SNc samples (Figure

5A). The upregulated gene, 2900052N01Rik, encodes a long intergenic noncoding RNA with no

known function. For most genes we observed a ~10-fold expression level decrease (with median

log2 FoldChange of -3.47 and mean log2 FoldChange of -3.2) in in the absence of DA. Among

the downregulated genes are several transcription factors involved in DA neuron specification:

Foxa2, *Gata3*, *Nr4a2*, *Pax5*, *Pitx3*, *En1* and *En2* (Figure 6A). Consistent with decreased

expression of the crucial dopaminergic transcriptional regulators, there is a reduced level of

classical dopaminergic markers: *Dao*; *Ddc*; *Ret*; *Gch1*; *Slc6a3* (codes DAT); *Slc18a2* (VMAT2)

(representative expression plots are collected on Figure 5B). Supplemental file 1: Table S1

reports the complete results of this analysis, providing full information about DE genes, with the

values of log2FoldChange, log2foldchange Standard Error, Wald statistic, p-value and FDR

adjusted p-value.

VTA and SNc Transcriptomes Respond Distinctly to DA Deficiency

Comparing expression levels of 15,187 genes detected in VTA in the DAT:TH KO^{Low} and WT

samples, we detected 89 DE genes at FDR < 0.05. In contrast to the general directionality of the

DE observed in SNc, the majority of DE genes were upregulated in DAT:TH KO^{Low} VTA samples

(55 upregulated vs 34 downregulated genes; Figure 7A). Moreover, the magnitude of observed

changes is smaller than in SNc. The majority of the DE is down- or upregulated less than two-

fold (60 out of 89 DE genes, with median absolute value of log2 Fold Change of 0.7 and mean of

1.09). However, the hierarchical cluster analysis based on the DE genes shows again that all

DAT:TH KO^{Low} samples are clustered together, whereas DAT:TH KO^{High} are more similar to WT.

Moreover, the "WT-" sample with a relatively low level of the *Th* expression is clustered with

the DAT:TH KO^{Low} libraries, showing an intermediate pattern of DE (Figure 7B). Furthermore,

when we compared DE across the entire transcriptome, none of the DE genes detected between

DAT:TH KO^{Low} and all WT samples have a statistically significant difference in expression level

between DAT:TH KO^{High} and WT VTA.

In addition to *Th*, only five DE genes from DAT:TH KO^{Low} SNc were also significantly

downregulated in VTA samples: *Chrnb3*, *Cox20*, *En1*, *Pitx3* and *Slc6a3* (Figure 8A). Among the

rest of the downregulated genes in VTA only additional two are known transcription factors:

Foxa1 and *Pbx3*. According to this, no additional classical dopaminergic markers were

downregulated in VTA.

To further examine potential differences between the dopamine dependency in SNc and VTA we

looked at the Gene Ontology (GO) terms enriched in detected DE genes, with top enriched

Biological Process (BP) GO terms are collected in Table 1. The full results of the GO enrichment

analyses are collected in the Additional file 2 Table S2. As expected, the top ten most enriched

terms in genes downregulated in SNc are connected to DA metabolism and to dopaminergic

neurons and their functions (20 out of 34 enriched categories). Results of the enrichment analysis

from the genes downregulated in the VTA are not as explicit. Only 5 categories were

significantly enriched after correction using Benjamini–Hochberg procedure. While none of

them directly connect to DA, one term, functionally related to DA, the GO:0007626 (locomotory

behavior), is also enriched in DE genes in SNc. The other two related categories are connected to

mitochondria. In contrast to the downregulated, the genes upregulated in VTA show high level of

enrichment in 85 BP categories. This difference could stem from the greater number of genes

with increased level of expression in DAT:TH KO^{Low} VTA samples. However, it could also

suggest stronger and active upregulation of particular pathways. The top ten enriched terms

revolve around synaptic transmission and glutamatergic signaling. Other terms of particular

interest point to the regulation of calcium and GABA-ergic signaling. These results suggest that

while the decrease level of DA in SNc may cause loss of dopaminergic identity, the neurons in

VTA seem to have more synaptic plasticity and react by remodeling of synaptic signaling.

Biological Process GO term enriched in DE genes in SNc

ID	Description	Count	p-value	p.adjust	Genes
GO:0009713	catechol-containing compound biosynthetic process	7	1.97E-14	1.12E-11	Ddc/Slc6a3/Gch1/Gata3/ Nr4a2/Dao/Th
GO:0042423	catecholamine biosynthetic process	7	1.97E-14	1.12E-11	Ddc/Slc6a3/Gch1/Gata3/ Nr4a2/Dao/Th
GO:0071542	dopaminergic neuron differentiation	6	5.81E-11	9.45E-09	En1/Pitx3/Nr4a2/Foxa2/ Pax5/En2
GO:0009636	response to toxic substance	10	2.28E-10	2.88E-08	Ddc/Slc6a3/Gch1/ Slc18a2/Lcn2/Nr4a2/ Th/ Gucy2c/ Chrnb3/Chrna6
GO:0098700	neurotransmitter loading into synaptic vesicle	4	3.94E-09	4.08E-07	Ddc/Slc18a2/Slc10a4/Th
GO:0007626	locomotory behavior	8	1.96E-08	1.86E-06	En1/Slc6a3/Sncg/Pitx3/ Slc18a2/Nr4a2/Foxa2/Th
GO:1901617	organic hydroxy compound biosynthetic process	7	5.64E-08	4.94E-06	Ddc/Slc6a3/Gch1/Gata3/ Nr4a2/Dao/Th
GO:0097164	ammonium ion metabolic process	6	3.94E-07	2.99E-05	Ddc/Slc6a3/Gch1/Nr4a2/ Dao/Th
GO:0001505	regulation of neurotransmitter levels	8	4.71E-07	3.35E-05	Ddc/Slc6a3/Colq/Sncg/ Gch1/Th/Chrnb3/Chrna6

Biological Process GO term enriched in downregulated genes in VTA

ID	Description	Count	p-value	p.adjust	Genes
GO:0007005	mitochondrion organization	8	3.07E-07	0.000205	Cox20/Dcn/Prdx3/ Ndufaf4/Tomm5/Ndufab1/ Higd1a/Apoo

ID	Description	Count	p-value	p.adjust	Genes
GO:0033108	mitochondrial respiratory chain complex assembly	3	1.81E-05	0.006042	*Cox20/Ndufaf4/ Ndufab1*
GO:0007626	locomotory behavior	5	0.000154	0.034539	*En1/Slc6a3/Pitx3/Pbx3/Th*
GO:0046500	S-adenosylmethionine metabolic process	2	0.000235	0.039340	*Mat2a/Mat2a*
GO:0051591	response to cAMP	3	0.000328	0.043891	*Slc6a3/Mat2a/Mat2a*

Biological Process GO term enriched in upregulated genes in VTA

ID	Description	Count	p-value	p.adjust	Genes
GO:0097060	synaptic membrane	14	1.37E-10	2.49E-08	*Epha4/Grm1/Atp2b1/Adcy8/ Adgrb1/Kcnb1/Lrrc7/Gabbr2/ Grin3a/Rims3/Cacna1c/Prkcg/ Gabrg3/Iqsec2*
GO:0099699	integral component of synaptic membrane	10	5.02E-09	4.56E-07	*Epha4/Grm1/Atp2b1/Adcy8/ Adgrb1/Lrrc7/Gabbr2/Grin3a/ Cacna1c/Gabrg3*
GO:0099240	intrinsic component of synaptic membrane	10	1.09E-08	6.60E-07	*Epha4/Grm1/Atp2b1/Adcy8/ Adgrb1/Lrrc7/Gabbr2/Grin3a/ Cacna1c/Gabrg3*
GO:0098984	neuron to neuron synapse	1q	1.97E-08	7.44E-07	*Epha4/Grm1/Ptk2b/Adcy8/ Adgrb1/Syngap1/Lrrc7/Grin3a /Cacna1c/Prkcg/Iqsec2*
GO:0098978	glutamatergic synapse	13	2.04E-08	7.44E-07	*Epha4/Grm1/Atp2b1/Ptk2b/ Adcy8/Adgrb1/Syngap1/Lrrc7 /Gabbr2/Grin3a/Rims3/ Cacna1c/Iqsec2*
GO:0050804	modulation of chemical synaptic transmission	13	1.16E-09	7.75E-07	*Epha4/Grm1/Jph4/Ptk2b/ Adcy8/Adgrb1/Syngap1/ Cacna1b/Kcnb1/Grin3a/Rims3 /Prkcg/Iqsec2*
GO:0099177	regulation of trans-synaptic signaling	13	1.19E-09	7.75E-07	*Epha4/Grm1/Jph4/Ptk2b/ Adcy8/Adgrb1/Syngap1/ Cacna1b/Kcnb1/Grin3a/Rims3 /Prkcg/Iqsec2*
GO:0014069	postsynaptic density	11	1.06E-07	2.98E-06	*Epha4/Grm1/Ptk2b/Adgrb1/ Syngap1/Lrrc7/Grin3a/ Cacna1c/Prkcg/Prkcg/Iqsec2*
GO:0032279	asymmetric synapse	10	1.15E-07	2.98E-06	*Epha4/Grm1/Ptk2b/Adgrb1/ Syngap1/Lrrc7/Grin3a/ Cacna1c/Prkcg/Iqsec2*
GO:0099055	integral component of postsynaptic membrane	7	2.16E-06	3.93E-05	*Epha4/Grm1/Adgrb1/Lrrc7/ Gabbr2/Grin3a/Cacna1c*
GO:0097060	synaptic membrane	14	1.37E-10	2.49E-08	*Epha4/Grm1/Atp2b1/Adcy8/ Adgrb1/Kcnb1/Lrrc7/Gabbr2/ Grin3a/Rims3/Cacna1c/Prkcg/ Gabrg3/Iqsec2*

Discussion

The majority of animal models of PD are based on neurotoxins or neuropharmacological agents targeting mitochondria, in particular the complex I of respiratory chain (Gubellini and Kachidian 2015). Although mitochondrial disfunction is well described in the pathogenesis of PD (Dawson and Dawson 2003; Gubellini et al. 2010), the neurodegeneration induced in those models is more rapid and acute than observed in PD. The strong cellular and behavioral phenotypes are observable in animals just weeks after a treatment with 6-OHDA or MPTP (Jackson-Lewis et al. 1995; Ungerstedt 1968; Ungerstedt and Arbuthnott 1970), whereas PD develops slowly, and the first locomotor symptoms could occur years after initiation of neurodegeneration (Halliday et al. 2008; Postuma et al. 2012). In this study we used a DAT:TH KO mouse model, generated by DA transporter driven CRE (DAT-CRE) to inactivate the *Th* gene, with loxP sites flanking *Th* exon 1, which encodes the *Th* translation startsite. This approach results in animals that lack 95% of a normal level of DA in the striatum, however the number of DA neurons is otherwise unaffected (Darvas, Henschen, et al. 2014; Garrett Morgan et al. 2015). This allows observation of transcriptome changes in response to DA loss, especially in SNc and VTA, that could have been missed in models based on the loss of DA neurons. Furthermore, as the DAT:TH KO model used here does not induce neuronal death, we could separate effects of the DA deficiency and a decreased amount of DA projections into striatum that are difficult to discriminate in the lesion-based models. Previously published comparisons between these two approaches show that more severe symptoms of PD, such as bradyphrenia, cognitive rigidity and spatial learning deficits, are dependent on both loss of DA and DA neuronal loss in striatum (Darvas, Henschen, et al. 2014). However, cognitive flexibility, working memory and cue-dependent associative learning can be

244 caused solely by decrease in striatal DA, independent of depletion of nigrostriatal DA neurons

245 (Garrett Morgan et al. 2015).

246 Based on the previously described distinct phenotype of DAT:TH KO animals, we hypothesized

247 that potential transcriptome changes might affect only subpopulations of cells that are the most

248 sensitive to dysregulation of DA homeostasis. We focused on the two midbrain regions, SNc and

249 VTA, as they are the major source of DA in the brain, and the three postsynaptic regions, VS, DS

250 and PFC that are innervated by VTA and SNc neuronal projections. This approach increased

251 sensitivity of our RNA-Seq analysis, allowing us to assess genes expressed specifically in these

252 brain regions, that could be overlooked in whole brain samples.

253 Surprisingly, we detected near normal levels of *Th* transcripts in half of the DAT:TH KO

254 samples (Fig. 1A). However, almost no reads were mapped to the FLOXed *Th* exon 1 in either of

255 the DAT:TH KO samples, eliminating the possibility that this results from inefficient Cre

256 recombinase function (Fig. 1B and C). The few *Th* exon 1 reads in DAT:TH KO samples could

257 come from DA neurons that do not express DAT (Darvas, Henschen, et al. 2014).

258 The unusual *Th* transcript that is detectable in the DAT:TH KO[High] samples lacks the first exon

259 that encodes the normal translation startsite, yet the remaining transcript has the same structure

260 as native *Th* mRNA. The normal N-terminal domain, encoded by exon 1, is involved in

261 regulation of TH enzymatic activity, with several phosphorylatable serines involved in TH

262 activation (Tekin et al. 2014), though deletion of this region can stabilize the TH protein

263 (Nakashima et al. 2005). To date, only one isoform of *Th* was characterized in mouse, however,

264 there are four alternative splicing forms of *TH* in humans that differ in structure at the 5' end of

265 the transcript, resulting in proteins with different N-terminal regulatory segments (Tekin et al.

266 2014). The *Th* transcripts detected in DAT:TH KO[High] animals could potentially represent a

novel mouse *Th* isoform that is differentially regulated relative to wildtype *Th*. The regional PCA

and the hierarchical cluster analysis based on SNc samples show that DAT:TH KO^{High} cluster

more strongly with WTs than with DAT:TH KO^{Low} animals (Fig. 2), supporting this hypothesis.

Therefore, we decided to treat the samples from those animals as a separate group, that could

represent potential DA-compensatory mechanism. Interestingly, when we compared the

expression levels of the genes, detected as differentially expressed between DAT:TH KO^{Low} and

WT, the samples from DAT:TH KO^{High} resemble WT animals. In the hierarchical cluster

analysis (Fig. 5 and 7) all DAT:TH KO^{High} samples clustered between WT samples.

Additionally, WT samples with lower level of Th, and thus presumably lower level of DA, are

clustered closer to the DAT:TH KO^{Low} animals. Those results support our hypothesis that the

alternative Th transcripts from DAT:TH KO^{High} animals could rescue the DAT:TH KO^{Low}

phenotype. However, further studies will be needed to confirm if DAT:TH KO^{High} animals

exhibit more wildtype-like behavioral phenotype and whether they express normal levels of

striatal DA.

The PCA and hierarchical cluster analysis of all samples clearly reflect differences in

transcriptomes of pre- and postsynaptic brain areas (Fig. 3A and B). The samples of those two

origins are strongly separated on the PC1 axis, whereas the PC2 additionally creates cluster of

the PFC datapoints. However, we could not observe the clear separation of DAT:TH KO samples

in any of tested regions based on the two first PC nor in the results of hierarchical cluster

analysis. These finding suggest that previously described broad transcriptional changes in

striatum in neurodegeneration-based PD models are induced by decrease in the number of DA

neurons, and not only by the decrease in DA level (Chin et al. 2008; Patel et al. 2008; Pattarini et

al. 2008; Singh et al. 2010).

In our study we detected nine genes DE in the postsynaptic regions, seven in DS and three in VS,

with one gene downregulated in both parts of striatum. We confirmed the dopaminergic

regulation of dynorphin expression in DS described previously in DA deficient models based on

both complete inactivation of *Th* and 6-OHDA-lesions (Sivam 1996; Zhou and Palmiter 1995).

Dynorphin belongs to an endogenous opioid peptide family, involved in regulation of DA

release, motor behavior, and learning (Carey et al. 2009; Gerfen and Steiner 1998; Reid et al.

1990; You, Herrera-marschitz, and Terenius 1999). The transcriptional activity of *Pdyn*, the gene

encoding prodynorphin, is dependent on signaling through the DA D1 receptor, and works in a

regulatory loop suppressing excessive activation of the striatal neurons (Gerfen and Steiner 1998;

Xu et al. 1994). We detected a decreased expression level of a related protein, syntrophin gamma

2, encoded by *Sntg2*, in both DS and VS DAT:TH KO^{Low} samples. Syntrophin gamma 2 belongs

to the family of cytoplasmic peripheral membrane proteins that binds to the C-terminal of

dynorphin and act as a adaptor proteins by its PDZ domain (Alessi et al. 2006; Piluso et al.

2000). However, target proteins associated with dynorphin-syntrophin scaffold complex are

currently unknown.

Subsequently, we focused on the SNc and VTA, as inactivation of Th directly affect neurons

from those areas.

Looking more closely at differences between animals with expected low level of *Th* mRNA and

WT controls, we detect 33 strongly downregulated genes in SNc (Fig. 5). These genes include

dopaminergic markers, such as *Aldh1a7*, *Ddc*, *En1*, *En2*, *Foxa2*, *Nr4a2* (*Nurr1*), *Pitx3*, *Slc18a2*

(VMAT2), *Slc6a3* (DAT) and *Th* (Domanskyi et al. 2014; Jacobs et al. 2011; Tekin et al. 2014).

Significant 1-2 log decreases in expression of those genes could be potentially explained by a

depletion of DA neurons in the SNc in DAT:TH KO animals. However we did not detect a

downregulation of antiapoptotic genes or an upregulation of canonical genes connected to cell

death, genes that are usually differentially expressed in PD and PD models, such as genes from

Bcl2 family, *Mapk8ip3*, *Lrpprc* and *Nfrkb* or *Casp8* (Elmore 2007; Simunovic et al. 2009). The

lack of evidence for cell death is consistent with the previous descriptions of this DAT:TH KO

model (Darvas, Henschen, et al. 2014; Garrett Morgan et al. 2015). Nevertheless,

downregulation of the DA related genes in SNc of DAT:TH KO[Low] animals is in accordance with

changes in PD and the lesion-based PD animal models. However, here for the first time we show

that decreased level of expression of dopaminergic genes in SNc and VTA could be a direct

effect of decreased DA as opposed to a result of death of dopaminergic neurons.

Additionally, we detected downregulation of genes associated with PD, such as the

neuroprotective gene *Fgf20* (Gao et al. 2008; Sleeman, Boshoff, and Duty 2012; Van Der Walt et

al. 2004); and *Gch1*, of which variants are connected with decreased risk of PD (Kalia and Lang

2015). We also found that two different subunits of Nicotinic Acetylcholine Receptor (*Chrna6*,

Chrnb3) are also downregulated in DAT:TH KO SNc. Nicotinic receptors involved in regulation

of DA release and are proposed as potential targets for PD treatment (Koranda et al. 2014; Leino

et al. 2018; Quik and Wonnacott 2011). Among the DE genes in SNc are also ones previously

unconnected to PD, yet of particular interest: *Dao*, D-amino acid oxidase, that may interact with

Aldh1a7 in alternative pathway of GABA synthesis, that was also implicated in alternative

pathway for DA biosynthesis (Kawazoe et al. 2007; J.-I. Kim et al. 2015); *Slc10a4*, a relatively

less known vesicular monoaminergic and cholinergic associated transporter, that plays a role in

the regulation of DA homeostasis (Larhammar et al. 2015; Melief et al. 2016). Another

potentially important transcription factor that is involved in the development of DA neurons,

Pax5 (Simon et al. 2003), is also significantly downregulated in DAT:TH KO[Low] SNc. That

suggests, that some of the observed changes could be connected to decreased striatal DA level

during development. The downregulation of the postmitotic selector *Gata3*, involved in

differentiation of SNc and VTA GABAergic neurons indicate developmental effect of DAT:TH

KO as well (Haugas et al. 2016; Moriguchi 2006).

The results of differential gene expression analysis of DAT:TH KO^Low and WT VTA samples

highlight the important differences between VTA and SNc. We not only detect more DE genes,

but moreover the majority of them are upregulated in the absence of DA. This distinct response

could reflect a differential variability of DA neurons in SNc and VTA in the PD (Brichta and

Greengard 2014), as only four DA genes have significantly lower level in DAT:TH KO^Low VTA

(*Th*, *Slc6a3*, *Pitx3* and *En1*; **Error! Reference source not found.**). Among the genes

downregulated in SNc, but not in VTA, *Nr4a2*, gene encoding the highly conserved transcription

factor Nurr1, is of particular interest. In PD patient brains, Nurr1 is downregulated in regions

affected by α-synuclein inclusions and variants of *Nr4a2* are associated with cases of PD

(Grimes et al. 2006; Le et al. 2003; Xu et al. 2002; Zheng, Heydari, and Simon 2003). Nurr1 is

expressed in the early postmitotic neurons, is necessary for a proper differentiation and is

maintained in midbrain DA neurons in the adult brain (Perlmann and Wallén-Mackenzie 2004;

Zetterström et al. 1997). Nurr1 null mice die within 24h after birth and conditional knockout of

Nr4a2 in DA neurons in adult animals caused decreased expression of DA markers, including *Th*

and *Slc6a3*, followed by progressive loss of DA (Castillo et al. 1998; Kadkhodaei et al. 2009,

2013). Six additional DA-related transcription factors are downregulated in SNc (*Foxa2*, *Gata3*,

Pax5, *Pitx3*, *En1* and *En2)*, but only *Pitx3* and *En1* are also downregulated in VTA of DAT:TH

KO^Low animals. Together, our data suggests that DA is involved in maintaining expression of the

DA-related genes required for DA biosynthesis and DA signaling by a novel regulatory loop

with different susceptibility in SNc and VTA.

One of the genes downregulated in both SNc and VTA is *Chrnb3*, that encodes the nicotinic

acetylcholine receptor. These results strengthen the role of the nicotinic acetylcholine receptors

in the development of PD. However, the observed downregulation in both regions could actually

represent the counter reaction of brain to decreased DA, since mice with α5 nicotinic receptor

subunit gene knockout have less severe phenotype after treatment with 6-OHDA (Leino et al.

2018).

Another gene downregulated in both presynaptic regions encodes the cytochrome c oxidase

assembly protein COX20. This finding could indicate that DA deficiency contributes to

mitochondrial dysfunction involved in pathogenesis of PD, since COX20 is required for

Cytochrome c oxidase assembly, required for normal mitochondrial function (Bourens et al.

2014; Elliott, Saracco, and Fox 2012).

Furthermore, we observe downregulation of five additional genes involved in assembly and

proper function of the mitochondrial respiratory chain: *Ndufaf4* and *Ndufab1* connected to

mitochondrial complex I, with *Ndufab1* downregulated in PD patients (Baertling et al. 2017;

Marcus et al. 2013; Saada et al. 2009; Simunovic et al. 2009; Triepels et al. 1999; Zurita Rendón

and Shoubridge 2012); *Apoo*, involved in complexes assuring stability of cristae (Ott et al. 2015);

Higd1a regulating cytochrome c oxidase (Hayashi et al. 2015); and *Tomm5*, a part of the

translocase complex of the mitochondrial outer membrane (Kato and Mihara 2008). However,

the decrease in expression of these particular genes does not necessarily implicate mitochondrial

dysfunction, but could also be an effect of a decrease in mitochondrial burden in the absence of

DA-related oxidative stress. A study utilizing the 6-OHDA PD mouse model showed that the DA

depletion does not elevate markers of oxidative stress in SNc, and moreover another study

indicates that decreased DA suppresses oxidative stress caused by ectopic DAT overexpression

(Chen et al. 2008; Smith and Cass 2007). This hypothesis is additionally supported by our data,

as we observed a decreased level of *Prdx3* in DAT:TH KO VTA, the gene encoding the only

mitochondrial member of peroxiredoxin family enzymes that plays a protective antioxidant role

by its peroxidase activity (Jiang et al. 2017; Liu et al. 2016).

The majority of genes upregulated in DAT:TH KO VTA are annotated with GO terms connected

to the regulation of trans-synaptic signaling and function of glutamatergic synapses.

Reciprocal signaling between DA and glutamate is well documented. Decreased levels of

midbrain DA resulting from unilateral 6-OHDA lesions, bilaterally increases levels of glutamate

in striatum and rostromedial tegmental nucleus, with parallel changes in expression of glutamate

receptors (Chang et al. 2019; Lindefors and Ungerstedt 1990; Rodríguez-Puertas et al. 1999).

There are additional levels of crosstalk between glutamate and dopamine-glutamate neurons and

their local synapses within the midbrain DA regions (Dobi et al. 2010; Morales and Root 2014;

Viereckel et al. 2016; Wang and Morales 2009; Yamaguchi et al. 2015). In our study we detect

increased levels of metabotropic glutamate receptor 1 and ionotropic glutamate receptor NMDA

3A. Increased glutamate signaling may play a role in a tuning of the VTA, since activation of

glutamate receptors modulates firing of dopaminergic neurons and regulates extracellular DA

levels (Ferrada et al. 2017; Floresco et al. 2003; Lodge and Grace 2006; Oakman et al. 1995).

Here we show data that suggests that DA is required for maintained expression of dopaminergic

genes in SNc, and to a lesser degree in VTA, suggesting the existence of a feedback loop

responsible for the maintenance of DA signaling. Furthermore, our study implicates an

unexpected diversity in responses to DA loss between individual animals. Future studies should

show whether the unusual class of animals expressing near-normal levels of the FLOXed Th

mRNA represent a compensatory response, partly compensating for the DA loss.

Acknowledgements:

The laboratory of J.H. was supported by NIH 5R01GM084128.

Materials and methods

Animals

All animal use procedures were approved by the Institutional Animal Care and Use Committee at

the University of Washington. Mice were housed under a 12-h, light–dark cycle in a

temperature-controlled room with food and water available ad libitum. Mice with inactivation of

the Th gene in DAT-expressing neurons (DAT:TH KO) were generated by crossing mice with

two floxed *Th* alleles ($Th^{lox/lox}$) with mice that have one deleted *Th* allele ($Th^{/+}$) and one allele

with Cre recombinase expressed under the control of the *Slc6a3* gene encoding DAT

($Slc6a3^{Cre/+}$). DAT:TH KO mice had the genotype $Th^{/lox}$-$Slc6a3^{Cre/+}$. C57Bl/6 were used as wild

type (WT) control animals.

RNA Extraction

Mice, 3-5-month-old, were sacrificed by decapitation after brief exposure to CO_2; the VS, DS,

SNc, VTA and PFC from each mouse were dissected from the brain. Tissues were immediately

frozen in liquid nitrogen and stored at −80 °C. Tissues were processed, and subsequently cDNA

libraries from isolated RNA were prepared in three independent batches; first one with samples

from 3 DAT:TH KO and 3 WT animals; second with samples from one mouse per group; third

with 2 DAT:TH KO and 2 WT animals. Each time tissues were homogenized using Bullet

Blender with 0.5 mm RNase Free Zirconium Oxide Beads (Next Advance) in RLT buffer from

RNeasy Plus Micro Kit (Qiagen). Total RNA was isolated using the RNeasy Plus Micro Kit

(Qiagen). The quality and quantity of RNA from each sample was determined using an Agilent

2100 Bioanalyzer with Agilent RNA 6000 Pico Chips. Messenger RNA was then obtained from

at least 100 ng of total RNA using NEBNext® Poly(A) mRNA Magnetic Isolation Module.

RNA-seq libraries were prepared using NEBNext® Ultra™ Directional RNA Library Prep Kit

for Illumina® following manufacturer's protocol. The library quality and insert length were

checked using the Agilent 2100 Bioanalyzer with DNA High Sensitivity DNA Kit. The first two

sets of 30 and 10 libraries were sequenced using Illumina HiSeq 2000 (BGI), third set of 20

libraries was sequenced using Illumina HiSeq 2500 (UCLA).

The quality of obtained fastq files was examined using FASTQC (S. 2010). The reads were

aligned to the mouse reference genome (GRCm38) using HISAT2 v2.1.0 (D. Kim, Langmead,

and Salzberg 2015) aligner with --rna-strandness RF parameter. Aligned reads were then

summarized with featureCounts from the SourceForge Subread package (Liao, Smyth, and Shi

2014). Summarizing, we obtained over 2.3 billion uniquely mapped 100 bp paired-end reads

(42206643± 4754672 uniquely mapped reads per library). A differential expression analysis was

performed with R/Bioconductor package DESeq2 including the batch variable into the model

design (Love, Huber, and Anders 2014). Multiple testing was corrected for using the Benjamini-

Hochberg procedure. Sequencing data have been deposited in the Sequence Read Archive under

accession code PRJNA550435.

References

Alessi, Amy, April D. Bragg, Justin M. Percival, Jean Yoo, Douglas E. Albrecht, Stanley C.

Froehner, and Marvin E. Adams. 2006. "γ-Syntrophin Scaffolding Is Spatially and

Functionally Distinct from That of the α/β Syntrophins." *Experimental Cell Research*

312(16):3084–95.

Baertling, Fabian, Laura Sánchez-Caballero, Mariël A. M. Van Den Brand, Liesbeth T. Wintjes,

Maaike Brink, Frans A. Van Den Brandt, Callum Wilson, Richard J. T. Rodenburg, and Leo

G. J. Nijtmans. 2017. "NDUFAF4 Variants Are Associated with Leigh Syndrome and

Cause a Specific Mitochondrial Complex i Assembly Defect." *European Journal of Human*

Genetics 25(11):1273–77.

Ballanger, Benedicte, Maude Beaudoin-Gobert, Sara Neumane, Justine Epinat, Elise Metereau,

Sandra Duperrier, Emmanuel Broussolle, Stephane Thobois, Frederic Bonnefoi, Christian

Tourvielle, Franck Lavenne, Nicolas Costes, Didier Lebars, Luc Zimmer, Véronique

Sgambato-Faure, and Léon Tremblay. 2016. "Imaging Dopamine and Serotonin Systems on

MPTP Monkeys: A Longitudinal PET Investigation of Compensatory Mechanisms." *The*

Journal of Neuroscience 36(5):1577–89.

Baunez, C. and T. W. Robbins. 1999. "Effects of Dopamine Depletion of the Dorsal Striatum

and Further Interaction with Subthalamic Nucleus Lesions in an Attentional Task in the

Rat." *Neuroscience* 92(4):1343–56.

Bezard, E. and C. E. Gross. 1998. "Compensatory Mechanisms in Experimental and Human

Parkinsonism: Towards a Dynamic Approach." *Progress in Neurobiology* 55(2):93–116.

Bezard, Erwan, Christian E. Gross, and Jonathan M. Brotchie. 2003. "Presymptomatic

Compensation in Parkinson's Disease Is Not Dopamine-Mediated." *Trends in*

Neurosciences 26(4):215–21.

Bezard, Erwan, Paula Ravenscroft, Christian E. Gross, Alan R. Crossman, and Jonathan M.

Brotchie. 2001. "Upregulation of Striatal Preproenkephalin Gene Expression Occurs before

the Appearance of Parkinsonian Signs in 1-Methyl-4-Phenyl- 1,2,3,6-Tetrahydropyridine

Monkeys." *Neurobiology of Disease* 8(2):343–50.

Björklund, Anders and Stephen B. Dunnett. 2007. "Dopamine Neuron Systems in the Brain: An

Update." *Trends in Neurosciences* 30(5):194–202.

Blesa, J., C. Pifl, M. A. Sánchez-González, C. Juri, M. A. García-Cabezas, R. Adánez, E.

Iglesias, M. Collantes, I. Peñuelas, J. J. Sánchez-Hernández, M. C. Rodríguez-Oroz, C.

Avendaño, O. Hornykiewicz, C. Cavada, and J. A. Obeso. 2012. "The Nigrostriatal System

in the Presymptomatic and Symptomatic Stages in the MPTP Monkey Model: A PET,

Histological and Biochemical Study." *Neurobiology of Disease* 48(1):79–91.

Bourens, Myriam, Aren Boulet, Scot C. Leary, and Antoni Barrientos. 2014. "Human COX20

Cooperates with SCO1 and SCO2 to Mature COX2 and Promote the Assembly of

Cytochrome c Oxidase." *Human Molecular Genetics* 23(11):2901–13.

Brichta, Lars and Paul Greengard. 2014. "Molecular Determinants of Selective Dopaminergic

Vulnerability in Parkinson's Disease: An Update." *Frontiers in Neuroanatomy*

8(December):1–16.

Carey, A. N., A. M. Lyons, C. F. Shay, O. Dunton, and J. P. McLaughlin. 2009.

"Endogenous   Opioid Activation Mediates Stress-Induced Deficits in Learning and

Memory." *Journal of Neuroscience* 29(13):4293–4300.

Castillo, Susan O., Judit S. Baffi, Miklós Palkovits, David S. Goldstein, Irwin J. Kopin, Jassir

Witta, Mark A. Magnuson, and Vera M. Nikodem. 1998. "Dopamine Biosynthesis Is

Selectively Abolished in Substantia Nigra/Ventral Tegmental Area but Not in Hypothalamic

Neurons in Mice with Targeted Disruption of the Nurr1 Gene." *Molecular and Cellular*

Neuroscience 11(1–2):36–46.

Chang, Yongli, Chengxue Du, Lingna Han, Shuxuan Lv, Jin Zhang, Guanyun Bian, Guoyi Tang,

Yiwei Liu, Tao Chen, and Jian Liu. 2019. "Enhanced AMPA Receptor-Mediated Excitatory

Transmission in the Rodent Rostromedial Tegmental Nucleus Following Lesion of the

Nigrostriatal Pathway." *Neurochemistry International* 122(September 2018):85–93.

Chen, L., Y. Ding, B. Cagniard, A. D. Van Laar, A. Mortimer, W. Chi, T. G. Hastings, U. J.

Kang, and X. Zhuang. 2008. "Unregulated Cytosolic Dopamine Causes Neurodegeneration

Associated with Oxidative Stress in Mice." *Journal of Neuroscience* 28(2):425–33.

Chin, Mark H., Wei-Jun Qian, Haixing Wang, Vladislav A. Petyuk, Joshua S. Bloom, Daniel M.

Sforza, Goran Laćan, Dahai Liu, Arshad H. Khan, Rita M. Cantor, Diana J. Bigelow,

William P. Melega, David G. Camp, Richard D. Smith, and Desmond J. Smith. 2008.

"Mitochondrial Dysfunction, Oxidative Stress, and Apoptosis Revealed by Proteomic and

Transcriptomic Analyses of the Striata in Two Mouse Models of Parkinson's Disease."

Journal of Proteome Research 7(2):666–77.

Chudasama, Y. and T. W. Robbins. 2006. "Functions of Frontostriatal Systems in Cognition:

Comparative Neuropsychopharmacological Studies in Rats, Monkeys and Humans."

Biological Psychology 73(1):19–38.

Cools, R., R. A. Barker, B. J. Sahakian, and T. W. Robbins. 2001. "Enhanced or Impaired

Cognitive Function in Parkinson's Disease as a Function of Dopaminergic Medication and

Task Demands." *Cerebral Cortex (New York, N.Y. : 1991)* 11(12):1136–43.

Cools, Roshan, Roger A. Barker, Barbara J. Sahakian, and Trevor W. Robbins. 2003. "L-Dopa

Medication Remediates Cognitive Inflexibility, but Increases Impulsivity in Patients with Parkinson's Disease." *Neuropsychologia* 41(11):1431–41.

Da Cunha, Claudio, Miriam Elizabeth Mendes Angelucci, Newton S. Canteras, Susan Wonnacott, and Reinaldo N. Takahashi. 2002. "The Lesion of the Rat Substantia Nigra Pars Compacta Dopaminergic Neurons as a Model for Parkinson's Disease Memory Disabilities." *Cellular and Molecular Neurobiology* 22(3):227–37.

Darvas, Martin, Charles W. Henschen, and Richard D. Palmiter. 2014. "Contributions of Signaling by Dopamine Neurons in Dorsal Striatum to Cognitive Behaviors Corresponding to Those Observed in Parkinson's Disease." *Neurobiology of Disease* 65(11):112–23.

Darvas, Martin and Richard D. Palmiter. 2009. "Restriction of Dopamine Signaling to the Dorsolateral Striatum Is Sufficient for Many Cognitive Behaviors." *Proceedings of the National Academy of Sciences of the United States of America* 106(34):14664–69.

Darvas, Martin and Richard D. Palmiter. 2011. "Contributions of Striatal Dopamine Signaling to the Modulation of Cognitive Flexibility." *Biological Psychiatry* 69(7):704–7.

Darvas, Martin, Amanda M. Wunsch, Jeffrey T. Gibbs, and Richard D. Palmiter. 2014. "Dopamine Dependency for Acquisition and Performance of Pavlovian Conditioned Response." *Proceedings of the National Academy of Sciences of the United States of America* 111(7):2764–69.

Dawson, Ted M. and Valina L. Dawson. 2003. "Molecular Pathways of Neurodegeneration in Parkinson's Disease." *Science* 302(5646):819–22.

Dobi, A., E. B. Margolis, H. L. Wang, B. K. Harvey, and Marisela Morales. 2010. "Glutamatergic and Nonglutamatergic Neurons of the Ventral Tegmental Area Establish Local Synaptic Contacts with Dopaminergic and Nondopaminergic Neurons." *Journal of*

Neuroscience 30(1):218–29.

Domanskyi, Andrii, Heike Alter, Miriam A. Vogt, Peter Gass, and Ilya A. Vinnikov. 2014.

"Transcription Factors Foxa1 and Foxa2 Are Required for Adult Dopamine Neurons

Maintenance." *Frontiers in Cellular Neuroscience* 8(September):1–11.

Elliott, Leah E., Scott A. Saracco, and Thomas D. Fox. 2012. "Multiple Roles of the Cox20

Chaperone in Assembly of Saccharomyces Cerevisiae Cytochrome C Oxidase." *Genetics*

190(2):559–67.

Elmore, Susan. 2007. "Apoptosis: A Review of Programmed Cell Death." *Toxicologic Pathology*

35(4):495–516.

Ferrada, Carla, Ramón Sotomayor-Zárate, Jorge Abarca, and Katia Gysling. 2017. "The

Activation of Metabotropic Glutamate 5 Receptors in the Rat Ventral Tegmental Area

Increases Dopamine Extracellular Levels." *NeuroReport* 28(1):28–34.

Floresco, Stan B., Anthony R. West, Brian Ash, Holly Moorel, and Anthony A. Grace. 2003.

"Afferent Modulation of Dopamine Neuron Firing Differentially Regulates Tonic and

Phasic Dopamine Transmission." *Nature Neuroscience* 6(9):968–73.

Gao, X., W. K. Scott, G. Wang, G. Mayhew, Y. J. Li, J. M. Vance, and Eden R. Martin. 2008.

"Gene-Gene Interaction between FGF20 and MAOB in Parkinson Disease." *Annals of*

Human Genetics 72(2):157–62.

Garrett Morgan, R., Jeffrey T. Gibbs, Erica J. Melief, Nadia O. Postupna, Emily E. Sherfield,

Angela Wilson, C. Dirk Keene, Thomas J. Montine, Richard D. Palmiter, and Martin

Darvas. 2015. "Relative Contributions of Severe Dopaminergic Neuron Ablation and

Dopamine Depletion to Cognitive Impairment." *Experimental Neurology* 271(11):205–14.

Gerfen, Charles R. and Heinz Steiner. 1998. "Role of Dynorphin and Enkephalin in the

Regulation of Striatal Output Pathways and Behavior." *Experimental Brain Research*

123(1):60–76.

Golden, Judith P., Joseph A. DeMaro, Amanda Knoten, Masato Hoshi, Elizabeth Pehek, Eugene

M. Johnson, Robert W. Gereau, and Sanjay Jain. 2013. "Dopamine-Dependent

Compensation Maintains Motor Behavior in Mice with Developmental Ablation of

Dopaminergic Neurons." *The Journal of Neuroscience* 33(43):17095–107.

Grimes, David A., Fabin Han, Michel Panisset, Lemuel Racacho, Fengxia Xiao, Ruobing Zou,

Kelly Westaff, and Dennis E. Bulman. 2006. "Translated Mutation in the Nurr1 Gene as a

Cause for Parkinson's Disease." *Movement Disorders* 21(7):906–9.

Gubellini, P. and P. Kachidian. 2015. "Animal Models of Parkinson's Disease: An Updated

Overview." *Revue Neurologique* 171(11):750–61.

Gubellini, Paolo, Barbara Picconi, Massimiliano Di Filippo, and Paolo Calabresi. 2010.

"Downstream Mechanisms Triggered by Mitochondrial Dysfunction in the Basal Ganglia:

From Experimental Models to Neurodegenerative Diseases." *Biochimica et Biophysica Acta*

- Molecular Basis of Disease 1802(1):151–61.

Halliday, Glenda, Mariese Hely, Wayne Reid, and John Morris. 2008. "The Progression of

Pathology in Longitudinally Followed Patients with Parkinson's Disease." *Acta*

Neuropathologica 115(4):409–15.

Haugas, Maarja, Laura Tikker, Kaia Achim, Marjo Salminen, and Juha Partanen. 2016. "Gata2

and Gata3 Regulate the Differentiation of Serotonergic and Glutamatergic Neuron Subtypes

of the Dorsal Raphe." *Development* 143(23):4495–4508.

Hayashi, Takaharu, Yoshihiro Asano, Yasunori Shintani, Hiroshi Aoyama, Hidetaka Kioka,

Osamu Tsukamoto, Masahide Hikita, Kyoko Shinzawa-Itoh, Kazuaki Takafuji, Shuichiro

Higo, Hisakazu Kato, Satoru Yamazaki, Ken Matsuoka, Atsushi Nakano, Hiroshi Asanuma,

Masanori Asakura, Tetsuo Minamino, Yu-ichi Goto, Takashi Ogura, Masafumi Kitakaze,

Issei Komuro, Yasushi Sakata, Tomitake Tsukihara, Shinya Yoshikawa, and Seiji

Takashima. 2015. "Higd1a Is a Positive Regulator of Cytochrome c Oxidase ." *Proceedings*

of the National Academy of Sciences 112(5):1553–58.

Henschen, Charles W., Richard D. Palmiter, and Martin Darvas. 2013. "Restoration of

Dopamine Signaling to the Dorsal Striatum Is Sufficient for Aspects of Active Maternal

Behavior in Female Mice." *Endocrinology* 154(11):4316–27.

Hornykiewicz, O. 1975. "Brain Monoamines and Parkinsonism." *National Institute on Drug*

Abuse Research Monograph Series (3):13–21.

Jackson-Lewis, Vernice, Michael Jakowec, Robert E. Burke, and Serge Przedborski. 1995.

"Time Course and Morphology of Dopaminergic Neuronal Death Caused by the Neurotoxin

1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine." *Neurodegeneration* 4(3):257–69.

Jackson, Chad R., Guo-Xiang Ruan, Fazila Aseem, Jane Abey, Karen Gamble, Greg Stanwood,

Richard D. Palmiter, P. Michael Iuvone, and Douglas G. McMahon. 2012. "Retinal

Dopamine Mediates Multiple Dimensions of Light-Adapted Vision." *The Journal of*

Neuroscience : The Official Journal of the Society for Neuroscience 32(27):9359–68.

Jacobs, F. M. J., J. V. Veenvliet, W. H. Almirza, E. J. Hoekstra, L. von Oerthel, A. J. A. van der

Linden, R. Neijts, M. G. Koerkamp, D. van Leenen, F. C. P. Holstege, J. P. H. Burbach, and

M. P. Smidt. 2011. "Retinoic Acid-Dependent and -Independent Gene-Regulatory Pathways

of Pitx3 in Meso-Diencephalic Dopaminergic Neurons." *Development* 138(23):5213–22.

Jiang, Min-Yao, Zhao-Dong Han, Wenjiao Li, Fei Yue, Jianheng Ye, Bowei Li, Zhiduan Cai,

Jian-Ming Lu, Weimin Dong, Xianhan Jiang, Weide Zhong, Huichan He, and Leyuan Liu.

2017. "Mitochondrion-Associated Protein Peroxiredoxin 3 Promotes Benign Prostatic

Hyperplasia through Autophagy Suppression and Pyroptosis Activation." *Oncotarget*

8(46):80295–302.

Kadkhodaei, B., A. Alvarsson, N. Schintu, D. Ramskold, N. Volakakis, E. Joodmardi, T.

Yoshitake, J. Kehr, M. Decressac, A. Bjorklund, R. Sandberg, P. Svenningsson, and T.

Perlmann. 2013. "Transcription Factor Nurr1 Maintains Fiber Integrity and Nuclear-

Encoded Mitochondrial Gene Expression in Dopamine Neurons." *Proceedings of the*

National Academy of Sciences 110(6):2360–65.

Kadkhodaei, B., T. Ito, E. Joodmardi, B. Mattsson, C. Rouillard, M. Carta, S. I. Muramatsu, C.

Sumi-Ichinose, T. Nomura, D. Metzger, P. Chambon, E. Lindqvist, N. G. Larsson, L.

Olson, A. Bjorklund, H. Ichinose, and T. Perlmann. 2009. "Nurr1 Is Required for

Maintenance of Maturing and Adult Midbrain Dopamine Neurons." *Journal of*

Neuroscience 29(50):15923–32.

Kalia, Lorraine V. and Anthony E. Lang. 2015. "Parkinson's Disease." *The Lancet*

386(9996):896–912.

Kato, Hiroki and Katsuyoshi Mihara. 2008. "Identification of Tom5 and Tom6 in the Preprotein

Translocase Complex of Human Mitochondrial Outer Membrane." *Biochemical and*

Biophysical Research Communications 369(3):958–63.

Kawazoe, Tomoya, Hideaki Tsuge, Takahito Imagawa, Kenji Aki, Seiki Kuramitsu, and Kiyoshi

Fukui. 2007. "Structural Basis of D-DOPA Oxidation by d-Amino Acid Oxidase:

Alternative Pathway for Dopamine Biosynthesis." *Biochemical and Biophysical Research*

Communications 355(2):385–91.

Kim, Daehwan, Ben Langmead, and Steven L. Salzberg. 2015. "HISAT: A Fast Spliced Aligner

with Low Memory Requirements." *Nature Methods* 12(4):357–60.

Kim, J. I., S. Ganesan, S. X. Luo, Y. W. Wu, E. Park, E. J. Huang, Lu Chen, and Jun B. Ding.

2015. "Aldehyde Dehydrogenase 1a1 Mediates a GABA Synthesis Pathway in Midbrain

Dopaminergic Neurons." *Science* 350(6256):102–6.

Koranda, Jessica L., Jackson J. Cone, Daniel S. McGehee, Mitchell F. Roitman, Jeff A. Beeler,

and Xiaoxi Zhuang. 2014. "Nicotinic Receptors Regulate the Dynamic Range of Dopamine

Release in Vivo." *Journal of Neurophysiology* 111:103–11.

Larhammar, Martin, Kalicharan Patra, Martina Blunder, Lina Emilsson, Christiane Peuckert,

Emma Arvidsson, Daniel Rönnlund, Julia Preobraschenski, Carolina Birgner, Christoph

Limbach, Jerker Widengren, Hans Blom, Reinhard Jahn, Åsa Wallén-Mackenzie, and Klas

Kullander. 2015. "SLC10A4 Is a Vesicular Amine-Associated Transporter Modulating

Dopamine Homeostasis." *Biological Psychiatry* 77(6):526–36.

Le, Wei dong, Pingyi Xu, Joseph Jankovic, Hong Jiang, Stanley H. Appel, Roy G. Smith, and

Demetrios K. Vassilatis. 2003. "Mutations in NR4A2 Associated with Familial Parkinson

Disease." *Nature Genetics* 33(1):85–89.

Leino, Sakari, Sini K. Koski, Raisa Hänninen, Tuukka Tapanainen, Saara Rannanpää, and Outi

Salminen. 2018. "Attenuated Dopaminergic Neurodegeneration and Motor Dysfunction in

Hemiparkinsonian Mice Lacking the A5 Nicotinic Acetylcholine Receptor Subunit."

Neuropharmacology 138:371–80.

De Leonibus, Elvira, Tiziana Pascucci, Sebastien Lopez, Alberto Oliverio, Marianne Amalric,

and Andrea Mele. 2007. "Spatial Deficits in a Mouse Model of Parkinson Disease."

Psychopharmacology 194(4):517–25.

Liao, Yang, Gordon K. Smyth, and Wei Shi. 2014. "FeatureCounts: An Efficient General

Purpose Program for Assigning Sequence Reads to Genomic Features." *Bioinformatics*

30(7):923–30.

Lindefors, Nils and Urban Ungerstedt. 1990. "Bilateral Regulation of Glutamate Tissue and

Extracellular Levels in Caudate-Putamen by Midbrain Dopamine Neurons." *Neuroscience*

Letters 115(2–3):248–52.

Liu, Zhilei, Yadong Hu, Yiyi Gong, Wenhao Zhang, Chongdong Liu, Qingtao Wang, and

Haiteng Deng. 2016. "Hydrogen Peroxide Mediated Mitochondrial UNG1-PRDX3

Interaction and UNG1 Degradation." *Free Radical Biology and Medicine* 99:54–62.

Lodge, D. J. and A. A. Grace. 2006. "The Laterodorsal Tegmentum Is Essential for Burst Firing

of Ventral Tegmental Area Dopamine Neurons." *Proceedings of the National Academy of*

Sciences 103(13):5167–72.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold

Change and Dispersion for RNA-Seq Data with DESeq2." 1–21.

Luk, Kelvin C., Victoria Kehm, Jenna Carroll, Bin Zhang, Patrick O'Brien, John Q.

Trojanowski, and Virginia M. Y. Lee. 2012. "Pathological α-Synuclein Transmission

Initiates Parkinson-like Neurodegeneration in Nontransgenic Mice." *Science (New York,*

N.Y.) 338(6109):949–53.

Lundblad, Martin, Mickael Decressac, Bengt Mattsson, and Anders Björklund. 2012. "Impaired

Neurotransmission Caused by Overexpression of α-Synuclein in Nigral Dopamine

Neurons." *Proceedings of the National Academy of Sciences of the United States of America*

109(9):3213–19.

Marcus, Dana, Michal Lichtenstein, Ann Saada, and Haya Lorberboum-Galski. 2013.

"Replacement of the C6ORF66 Assembly Factor (NDUFAF4) Restores Complex I Activity

in Patient Cells." *Molecular Medicine* 19(1):124–34.

Melief, E. J., J. T. Gibbs, X. Li, R. G. Morgan, C. D. Keene, T. J. Montine, R. D. Palmiter, and

M. Darvas. 2016. "Characterization of Cognitive Impairments and Neurotransmitter

Changes in a Novel Transgenic Mouse Lacking Slc10a4." *Neuroscience* 324(5):399–406.

Melief, Erica J., Jonathan W. McKinley, Jonathan Y. Lam, Nicole M. Whiteley, Alec W.

Gibson, John F. Neumaier, Charles W. Henschen, Richard D. Palmiter, Nigel S. Bamford,

and Martin Darvas. 2018. "Loss of Glutamate Signaling from the Thalamus to Dorsal

Striatum Impairs Motor Function and Slows the Execution of Learned Behaviors." *Npj*

Parkinson's Disease 4(1).

Morales, M. and D. H. Root. 2014. "Glutamate Neurons within the Midbrain Dopamine

Regions." *Neuroscience* 282:60–68.

Moriguchi, T. 2006. "Gata3 Participates in a Complex Transcriptional Feedback Network to

Regulate Sympathoadrenal Differentiation." *Development* 133(19):3871–81.

Mura, Anna and Joram Feldon. 2003. "Spatial Learning in Rats Is Impaired after Degeneration

of the Nigrostriatal Dopaminergic System." *Movement Disorders : Official Journal of the*

Movement Disorder Society 18(8):860–71.

Nakashima, Akira, Nobuhiro Hayashi, Yoko S. Kaneko, Keiji Mori, Hiromi Egusa, Toshiharu

Nagatsu, and Akira Ota. 2005. "Deletion of N-Terminus of Human Tyrosine Hydroxylase

Type 1 Enhances Stability of the Enzyme in AtT-20 Cells." 120:110–20.

Oakman, SA, PL Faris, PE Kerr, C. Cozzari, and BK Hartman. 1995. "Distribution of

Pontomesencephalic Cholinergic Neurons Projecting to Substantia Nigra Differs

Significantly from Those Projecting to Ventral Tegmental Area." *The Journal of*

Neuroscience 15(9):5859–69.

Ott, Christine, Eva Dorsch, Martin Fraunholz, Sebastian Straub, and Vera Kozjak-Pavlovic.
2015. "Detailed Analysis of the Human Mitochondrial Contact Site Complex Indicate a
Hierarchy of Subunits." *PLoS ONE* 10(3):1–15.

Patel, Suman, Kavita Singh, Seema Singh, and Mahendra Pratap Singh. 2008. "Gene Expression
Profiles of Mouse Striatum in Control and Maneb + Paraquat-Induced Parkinson's Disease
Phenotype: Validation of Differentially Expressed Energy Metabolizing Transcripts."
Molecular Biotechnology 40(1):59–68.

Pattarini, R., Y. Rong, C. Qu, and J. I. Morgan. 2008. "Distinct Mechanisms of 1-Methyl-4-
Phenyl-1,2,3,6-Tetrahydropyrimidine Resistance Revealed by Transcriptome Mapping in
Mouse Striatum." *Neuroscience* 155(4):1174–94.

Perez, Xiomara A., Neeraja Parameswaran, Luping Z. Huang, Kathryn T. O'Leary, and Maryka
Quik. 2008. "Pre-Synaptic Dopaminergic Compensation after Moderate Nigrostriatal
Damage in Non-Human Primates." *Journal of Neurochemistry* 105(5):1861–72.

Perlmann, Thomas and Asa Wallén-Mackenzie. 2004. "Nurr1, an Orphan Nuclear Receptor with
Essential Functions in Developing Dopamine Cells." *Cell and Tissue Research* 318(1):45–
52.

Piluso, Giulio, Massimiliano Mirabella, Enzo Ricci, Angela Belsito, Ciro Abbondanza, Serenella
Servidei, Annibale Alessandro Puca, Pietro Tonali, Giovanni Alfredo Puca, and Vincenzo
Nigro. 2000. "Γ1- and Γ2-Syntrophins, Two Novel Dystrophin-Binding Proteins Localized
in Neuronal Cells." *Journal of Biological Chemistry* 275(21):15851–60.

Postuma, Ronald B., Dag Aarsland, Paolo Barone, David J. Burn, Christopher H. Hawkes,
Wolfgang Oertel, and Tjalf Ziemssen. 2012. "Identifying Prodromal Parkinson's Disease:
Pre-Motor Disorders in Parkinson's Disease." *Movement Disorders* 27(5):617–26.

Quik, Maryka and Susan Wonnacott. 2011. "6 2* and 4 2* Nicotinic Acetylcholine Receptors As

Drug Targets for Parkinson's Disease." *Pharmacological Reviews* 63(4):938–66.

Reid, M. S., M. Herrera-Marschitz, T. Hökfelt, N. Lindefors, H. Persson, and U. Ungerstedt.

1990. "Striatonigral GABA, Dynorphin, Substance P and Neurokinin A Modulation of

Nigrostriatal Dopamine Release: Evidence for Direct Regulatory Mechanisms."

Experimental Brain Research 82(2):293–303.

Rodríguez-Puertas, R., M. Herrera-Marschitz, J. Koistinaho, and T. Hökfelt. 1999. "Dopamine

D1 Receptor Modulation of Glutamate Receptor Messenger RNA Levels in the Neocortex

and Neostriatum of Unilaterally 6-Hydroxydopamine- Lesioned Rats." *Neuroscience*

89(3):781–97.

S., Andrews. 2010. "FastQC: A Quality Control Tool for High Throughput Sequence Data."

Retrieved (http://www.bioinformatics.babraham.ac.uk/projects/fastqc).

Saada, Ann, Rutger O. Vogel, Saskia J. Hoefs, Mariël A. van den Brand, Hans J. Wessels, Peter

H. Willems, Hanka Venselaar, Avraham Shaag, Flora Barghuti, Orit Reish, Mordechai

Shohat, Martijn A. Huynen, Jan A. M. Smeitink, Lambert P. van den Heuvel, and Leo G.

Nijtmans. 2009. "Mutations in NDUFAF3 (C3ORF60), Encoding an NDUFAF4

(C6ORF66)-Interacting Complex I Assembly Protein, Cause Fatal Neonatal Mitochondrial

Disease." *American Journal of Human Genetics* 84(6):718–27.

Sawamoto, Nobukatsu, Paola Piccini, Gary Hotton, Nicola Pavese, Kris Thielemans, and David

J. Brooks. 2008. "Cognitive Deficits and Striato-Frontal Dopamine Release in Parkinson's

Disease." *Brain* 131(5):1294–1302.

Simon, Horst H., Lavinia Bhatt, Daniel Gherbassi, Paola Sgadó, and Lavinia Alberí. 2003.

"Midbrain Dopaminergic Neurons: Determination of Their Developmental Fate by

Transcription Factors." *Annals of the New York Academy of Sciences* 991:36–47.

Simunovic, Filip, Ming Yi, Yulei Wang, Laurel Macey, Lauren T. Brown, Anna M. Krichevsky,

Susan L. Andersen, Robert M. Stephens, Francine M. Benes, and Kai C. Sonntag. 2009.

"Gene Expression Profiling of Substantia Nigra Dopamine Neurons: Further Insights into

Parkinson's Disease Pathology." *Brain* 132(7):1795–1809.

Singh, Kavita, Seema Singh, Naveen Kumar Singhal, Amit Sharma, Devendra Parmar, and

Mahendra Pratap Singh. 2010. "Nicotine- and Caffeine-Mediated Changes in Gene

Expression Patterns of MPTP-Lesioned Mouse Striatum: Implications in Neuroprotection

Mechanism." *Chemico-Biological Interactions* 185(2):81–93.

Sivam, S. P. 1996. "Dopaminergic Regulation of Postnatal Development of Dynorphin Neurons

in Rat Striatum." *Neuropeptides* 30(1):103–7.

Sleeman, Isobel J., Eugene L. Boshoff, and Susan Duty. 2012. "Fibroblast Growth Factor-20

Protects against Dopamine Neuron Loss in Vitro and Provides Functional Protection in the

6-Hydroxydopamine-Lesioned Rat Model of Parkinson's Disease." *Neuropharmacology*

63(7):1268–77.

Smith, M. P. and W. A. Cass. 2007. "Oxidative Stress and Dopamine Depletion in an Intrastriatal

6-Hydroxydopamine Model of Parkinson's Disease." *Neuroscience* 144(3):1057–66.

Sulzer, David, Stephanie J. Cragg, and Margaret E. Rice. 2016. "Striatal Dopamine

Neurotransmission: Regulation of Release and Uptake." *Basal Ganglia* 6(3):123–48.

Tekin, Izel, Robert Roskoski, Nurgul Carkaci-Salli, and Kent E. Vrana. 2014. *Complex*

Molecular Regulation of Tyrosine Hydroxylase. Vol. 121.

Triepels, R., J. Smeitink, J. Loeffen, R. Smeets, C. Buskens, F. Trijbels, and L. Van Den Heuvel.

1999. "The Human Nuclear-Encoded Acyl Carrier Subunit (NDUFAB1) of the

Mitochondrial Complex I in Human Pathology." *Journal of Inherited Metabolic Disease*

22(2):163–73.

Ungerstedt, Urban. 1968. "6-Ohda Induced Degeneration of Central Monoamine Neurons

(Ungerstedt 1968).Pdf." 5:107–10.

Ungerstedt, Urban and Gordon W. Arbuthnott. 1970. "Quantitative Recording of Rotational

Behavior in Rats after 6-Hydroxy-Dopamine Lesions of the Nigrostriatal Dopamine

System." *Brain Research* 24(3):485–93.

Viereckel, Thomas, Sylvie Dumas, Casey J. A. Smith-Anttila, Bianca Vlcek, Zisis Bimpisidis,

Malin C. Lagerström, Asa Konradsson-Geuken, and Asa Wallén-Mackenzie. 2016.

"Midbrain Gene Screening Identifies a New Mesoaccumbal Glutamatergic Pathway and a

Marker for Dopamine Cells Neuroprotected in Parkinson's Disease." *Scientific Reports*

6(September):1–16.

Van Der Walt, Joelle M., Maher A. Noureddine, Raja Kittappa, Michael A. Hauser, William K.

Scott, Ron Mckay, Fengyu Zhang, Jeffrey M. Stajich, Kenichiro Fujiwara, Burton L. Scott,

Margaret A. Pericak-Vance, Jeffery M. Vance, and Eden R. Martin. 2004. "Fibroblast

Growth Factor 20 Polymorphisms and Haplotypes Strongly Influence Risk of Parkinson

Disease." *Am. J. Hum. Genet* 74:1121–27.

Wang, Hui-Ling and Marisela Morales. 2009. "Pedunculopontine and Laterodorsal Tegmental

Nuclei Contain Distinct Populations of Cholinergic, Glutamatergic and GABAergic

Neurons in the Rat." *European Journal of Neuroscience* 29(2):340–58.

Wise, Roy A. 2004. "Dopamine, Learning and Motivation." *Nature Reviews Neuroscience*

5(6):483–94.

Xu, Ming, Rosario Moratalla, Lisa H. Gold, Noboru Hiroi, George F. Koob, Ann M. Graybiel,

and Susumu Tonegawa. 1994. "Dopamine D1 Receptor Mutant Mice Are Deficient in

Striatal Expression of Dynorphin and in Dopamine-Mediated Behavioral Responses." *Cell*

79(4):729–42.

Xu, P. Y., R. Liang, J. Jankovic, C. Hunter, Y. X. Zeng, T. Ashizawa, D. Lai, and W. D. Le.

2002. "Association of Homozygous 7048G7049 Variant in the Intron Six of Nurr1 Gene

with Parkinson's Disease." *Neurology* 58(6):881–84.

Yamaguchi, Tsuyoshi, Jia Qi, Hui Ling Wang, Shiliang Zhang, and Marisela Morales. 2015.

"Glutamatergic and Dopaminergic Neurons in the Mouse Ventral Tegmental Area."

European Journal of Neuroscience 41(6):760–72.

You, Zhi-bing, Mario Herrera-marschitz, and Lars Terenius. 1999. "Modulation of

Neurotransmitter Release in the Basal Ganglia of the Rat Brain by Dynorphin Peptides 1."

Journal of Pharmacology and Experimental Therapeutics 290(3):1307–15.

Zetterström, R. H., Ludmila Solomin, Lottie Jansson, Barry J. Hoffer, Lars Olson, and Thomas

Perlmann. 1997. "Dopamine Neuron Agenesis in Nurr1-Deficient Mice." *Science (New*

York, N.Y.) 276(5310):248–50.

Zheng, Kangni, Bobak Heydari, and David K. Simon. 2003. "A Common NURR1

Polymorphism Associated with Parkinson Disease and Diffuse Lewy Body Disease."

Archives of Neurology 60(5):722–25.

Zhou, Qun Yong and Richard D. Palmiter. 1995. "Dopamine-Deficient Mice Are Severely

Hypoactive, Adipsic, and Aphagic." *Cell* 83(7):1197–1209.

Zhuang, Xiaoxi, Justine Masson, Jay A. Gingrich, Stephen Rayport, and René Hen. 2005.

"Targeted Gene Expression in Dopamine and Serotonin Neurons of the Mouse Brain."

Journal of Neuroscience Methods 143(1):27–32.

Zurita Rendón, Olga and Eric A. Shoubridge. 2012. "Early Complex I Assembly Defects Result

in Rapid Turnover of the ND1 Subunit." *Human Molecular Genetics* 21(17):3815–24.

Figures legends

Figure 1. Th expression level in SNc and VTA samples.

DAT:TH KO animals were split into two groups: DAT:TH KO[High] or DAT:TH KO[Low]. DAT:TH

KO animals were assigned to DAT:TH KO[High] group if they had *Th* expression level higher than

any WT animal in a given brain region. In all following analyses all samples from the same

animal are grouped into the same DAT:TH KO[Low or High] group. DESeq2 normalized counts are

based on featureCounts output. A) Normalized expression level of Th in both presynaptic

regions. The normalized counts are shown on log10 scale. B) Normalized expression level of Th

exon 1 is significantly lower in all DAT:TH KO animals in comparison to WT. Numbers over

bars represent p-value from Kruskal-Wallis test. Datapoints from "WT-" mouse are marked with

"-". C) Representative per base coverage plot of Th. The bottom part of the plots shows known

Th exons in orange and yellow rhombuses mark loxP sites flanking exon 1. Yellow square marks

the first exon, that should be removed by DAT:Cre recombinase in DAT:TH KO animals. Only

few Th exon 1 reads are detectable in all DAT:TH KO samples, however the remaining exons

have a high level of coverage in the DAT:TH KO[High] SNc and VTA samples.

Figure 2. Clustering analyses of SNc samples.

A) SNc PCA based on genes represented by at least 30 normalized counts across all samples.

The datapoint from "WT-" mouse is marked with "-". The dashed ellipse outlines a cluster of

samples with low level of *Th*. B) Hierarchical cluster analysis of SNc samples. Dendrogram and

heatmap were created based on the maximum distance between samples. Blue color indicates

large distance, with red diagonal line representing identity from comparison of the same sample.

Each row and column represent an individual library, with row/column labeling by color coded

boxes, following the same pattern as PCA plot. Leaves of dendrogram are color coded by

sequencing batch. Blue squares with "-" mark the sample form "WT-" mouse.

Figure 3. Cluster analyses.

Input data for all analyses are DESeq2 normalized count values based on featureCounts output.

A) Principal Component Analysis of all sequenced samples from three independent batches of

RNA-sequencing, coded by the shape of datapoints. Each data point represents an individual

library, color coded based on the brain region and genotype. The hollow square marks one

outlier PFC library. B) Hierarchical cluster analysis of all sequenced samples from three

independent batches of RNA-sequencing. Dendrogram and heatmap were created based on the

Euclidean distance between samples. On heatmap blue color indicate large distance, with red

diagonal line representing identity from comparison of the same sample. Each row and column

represent an individual library, with row/column labeled by color coded boxes based on the brain

region, following the same pattern as PCA plot. Leaves of dendrogram are color coded by

sequencing batch: 1^{st} – red, 2^{nd} – yellow, 3^{rd} – light blue.

Figure 4. Expression level of DE genes in postsynaptic brain regions.

All plotted genes have statistically significant DE in DAT:TH KO^{Low} at FDR <0.05. All figures

follow color coding from figure A. Datapoints from WT mouse with low level of *Th* mRNA are

marked by "-" sign. The lower number of WT samples in DS is due to a loss of 2 libraries during

preparation. A) Genes downregulated in DS. B) Genes upregulated in DS. C) DE genes in VS.

Figure 5. DE genes in SNc.

A) Volcano plots of DE analysis between WT and DAT:TH KO^{Low} samples. Data points are

color coded based on their adjusted p-value at FDR <0.05 and fold change. DE DA-related genes

are labeled. B) Heatmap representing hierarchical clustering of the SNc libraries based on

detected DE genes. Each column represents an expression patter of one DE gene. Color scale

represent Z-score, with red marking high expression level. Leaves of dendrogram and row labels

are by color coded based on the genotype and expression level of Th. WT sample with low level

of *Th* mRNA is marked by "-" sign.

Figure 6. Representative expression level of genes DE in SNc.

All plotted genes have statistically significant decreased expression level in DAT:TH KO^{Low} vs

WT samples at FDR <0.05. All figures follow color coding from figure B. Datapoints from WT

mouse with low level of *Th* mRNA are marked by "-" sign. A) log$_{10}$(Normalized Counts) of DA-

related downregulated transcription factors. B) log$_{10}$(Normalized Counts) of representative

downregulated dopaminergic markers.

Figure 7. Genes DE in VTA.

A) Volcano plots of DE analysis between WT and DAT:TH KO^{Low} VTA samples. Data points

are color coded based on their adjusted p-value at FDR <0.05 and fold change threshold. DE

DA-related genes are labeled. B) Heatmap representing hierarchical clustering of the VTA

samples based on detected DE genes. Each column represents an expression patter of one DE

gene. Color scale represent Z-score, with red marking high expression level. Leaves of

dendrogram and row labels are by color coded based on the genotype and expression level of *Th*.

WT sample with low level of *Th* mRNA is marked by "-" sign.

Figure 8. Expression levels of genes downregulated in both DAT:TH KO presynaptic

regions across VTA samples.

All plotted genes have statistically significant decreased expression level in DAT:TH KO$^{\text{Low}}$ SNc

and VTA at FDR <5%. Datapoints from WT mouse with low level of *Th* mRNA are marked by

"-" sign.

Supplementary Figure 1. RNA-Seq Samples profiles

Input data for all analyses are DESeq2 normalized count values based on featureCounts output.

A) Histograms representing the distributions of uniquely mapped reads per gene in examined

brain regions. B) Venn diagram showing numbers of genes detected uniquely in the individual

brain regions or in the intersections of two or more regions. Genes represented by fewer than 30

reads total were filtered out.

Table 1. Results of the GO enrichment analysis based on the DE genes in SNc and VTA at

FDR < 0.05.

P-adjusted values were corrected using Benjamini–Hochberg procedure. This table shows only

up to the top ten terms enriched based on the DE gene in SNc and VTA, from the total number of

33 and 85 significantly enriched categories.

Supplemental Information

SUPPLEMENTAL FILE 1 SF1.xlsx

Spreadsheet with the detailed results of DE analysis.

SUPPLEMENTAL FILE 2 SF2.xlsx

Spreadsheet with the results of GO analysis.

Figure 1

Figure 2

Figure 3

A

Genes downregulated in DS

B

Genes upregulated in DS

C

DE genes in VS

Figure 4

Figure 5

Figure 6

Figure 7

A

Figure 8

Supplementary Figure 1