Extraordinary Fast-Twitch Fiber Abundance Elite Weightlifters

Nathan Serrano[1], Lauren M. Colenso-Semple[1], Kara K. Lazauskus[1], Jeremy W. Siu[1], James R. Bagley[2], Robert G. Lockie[1], Pablo B. Costa[1], Andrew J. Galpin[1]

[1]Biochemistry and Molecular Exercise Physiology Laboratory, Center for Sport Performance, California State University, Fullerton, CA.

[1]Muscle Physiology Laboratory, San Francisco State University, San Francisco, CA.

Address for correspondence: Andrew J. Galpin, Ph.D., Center for Sport Performance, Department of Kinesiology, California State University, Fullerton, Fullerton, CA 92834; E-mail: agalpin@fullerton.edu.

Keywords: Fast-twitch, Olympic lifting, Strength, Single Fiber, Power, myosin heavy chain

ABSTRACT

Human skeletal muscle fibers exist across a continuum of slow → fast-twitch. The amount of each

fiber *type* (FT) influences muscle performance but remains largely unexplored in elite athletes,

particularly from strength/power sports. To address this nescience, v*astus lateralis* (*VL*) biopsies

were performed on World/Olympic (female, n=6, "WCF") and National-caliber (female, n=9,

"NCF"; and male, n=6, "NCM") American weightlifters. Participant accolades included 3

Olympic Games, 19 World Championships, 25 National records, and >170 National/International

medals. Samples were analyzed for myosin heavy chain (MHC) content via SDS-PAGE using two

distinct techniques: single fiber (SF) distribution (%) and homogenate (HG) composition. These

athletes displayed the highest MHC IIa concentrations ever reported in healthy *VL* (23±9% I, 5±3%

I/IIa, 67±13% IIa, and 6±10% IIa/IIx), with WCF expressing a notable 71±17% (NCF=67±8%,

NCM=63±16%). The heavyweights accounted for 91% of the MHC IIa/IIx fibers. When compared

to SF, HG overestimated MHC I (23±9 vs. 31±9%) and IIx (0±0 vs. 3±6%) by misclassifying I/IIa

fibers as I and IIa/IIx fibers as IIx. These findings suggest athlete caliber (World vs. National),

training experience, and body mass determine FT% more than sex and refutes the common

pronouncement that women possess more slow and fewer fast-twitch fibers than men. Our results

also show the abundance of pure MHC IIa and rarity of IIx in elite strength/power-trained athletes,

indicate a potential link between MHC IIa/IIx frequency and body mass, and question the fidelity

of HG as a measure of FT% distribution. The extreme fast-twitch abundance partially explains

how elite weightlifters generate high forces in rapid time-frames. These data highlight the need for

more cellular and molecular muscle research on elite anaerobic athletes.

INTRODUCTION

Legendary Italian physician Stefano Lorenzini made the first distinction of "red" and

"white" muscle fibers (myofibers) in 1678, and almost 200 years later (1873) French histologist

Louis-Antoine Ranvier confirmed the existence of two distinct myofiber *types* in vertebrate

skeletal muscle. Reintroduction of the skeletal muscle biopsy procedure in 1962 (1) allowed

scientists to begin exploring the topic in athletes and resulted in the discovery that each fiber type

(FT) is comprised of a unique myosin heavy chain (MHC) isoform signature. Human skeletal

muscle therefore contains three *pure* (MHC I, IIa, and IIx) and several *hybrid* (single myofibers

that co-expresses multiple MHC isoforms) FT (2). The pure and hybrid FT combine to form a

robust slow $\rightarrow$ fast continuum (MHC I $\rightarrow$ I/IIa $\rightarrow$ IIa $\rightarrow$ IIa/IIx $\rightarrow$ IIx) with each displaying

specific morphological, metabolic, and contractile properties (3-6). FT distribution (FT%), or the

relative quantity of each FT in a given muscle, influences whole muscle function (7) and is often

highly correlated with athletic performance (3, 7-13).

Extensive evidence indicates endurance athletes possess a slow-twitch myofiber majority

(9, 10, 12, 14, 15), yet relatively few investigations have explored FT in speed, power, or strength

athletes. Initial research in the 1970-80's found resistance-trained men expressed high quantities

(~60-65%) of fast-twitch fibers (11, 12, 15, 16), which was substantiated by later studies on elite

powerlifters (17) and national-caliber (*Olympic*) weightlifters (8). This trailblazing work provided

an important foundation, but used sub-elite participants (18) and/or laboratory methods that failed

to accurately resolve the highly prevalent hybrids (19-21) - which compromises measurement

fidelity and produces erroneous FT% conclusions (19, 22-25). More precise techniques were

developed in the early 1990's that allowed proper quantification of FT% by analyzing each single

myofiber (SF).

Since this time only 13 studies (Table 1) implemented SF in young speed, power, or

strength-trained individuals (5, 13, 19, 20, 22, 23, 25-30), and only 3 included females (n = 13,

total). Only 5/13 included athletes: unknown-caliber male sprinters (n = 6) (25), male soccer

players (n = 8) (24), elite female track and field runners (n = 6) (20), National-caliber male

bodybuilders (n = 8) (19), and a former World-champion male sprinter (n = 1) (13). Accurately

accounting for the full FT spectrum resulted in all five studies finding far lower MHC IIa

concentrations than expected (52%, 30%, 16%, 39%, and 34%, respectively). The extremely low

16% found by Parcell et al. (2003) (20) is possibly explained by sex as females are often purported

to possess more slow-twitch fibers than men (31, 32). Such sex-specific phenotypes are often the

case in murine models (31), but the topic remains unexplored in athletes. Moreover, these data are

difficult to interpret as the athletes sampled were from a combination of several dissimilar events

(i.e., pole vault, heptathlon, 400 m hurdles, etc.).

Table 1: Summary of literature reporting single muscle fiber myosin heavy chain (MHC)
fiber-type from the *vastus lateralis* in young speed, power, or strength-trained individuals.

Reference	Subjects	Condition	MHC Distribution (%)					
			I	I/IIa	IIa	IIa/IIx	IIx	I/IIa/IIx
Andersen (1994)	Sprinters; 6M (23y)	Post 12-week RE & Interval Training	41	1	52	5	0	0
Andersen (1994)	Soccer; 8M (23y)	National Players Post 12-week RE Training	59	3	30	9	0	<1
Williamson (2001)	Non Ath; 6M (25y) 6F (21 y)	Post 12-week RE Training	30 35	5 3	59 52	5 12	0 0	0 0
Parcell (2003)	Track & Field; 6F (23y)	Division I / Interntional - Caliber	57	9	16	14	1	1

Raue (2005)	Non Ath; 6M (24y) 6M (24y)	Post-Con RE Post-Ecc RE	38 25	1 7	34 39	27 25	0 2	<1 <1
Parcell (2005)	Non Ath; 10M (22y)	Post 8-week Sprint Cycle Training	34	8	44	12	0	2
Malisoux (2006)	Non Ath; 8M (23)	Post 8-week Plyometric Training	28	5	42	26	2	0
Kesidis (2008)	Bodybuilding; 8M (26 y)	National-Caliber	35	19	39	7	0	0
Trappe (2015)	Sprinter; 1M (? y)	Previously World Champion	24	5	34	9	24	0
Murach (2016)	Non Ath; 9M (25y)	Resistance Trained	17	10	60	11	<1	<1
Bagley (2017)	Non Ath; 15M (25y)	Resistance Trained	20	10	58	11	1	1
Arevalo (2017)	Non Ath; 13M (24y)	Resistance Trained	28	9	60	3	<1	<1
Tobias (2017)	Non Ath; 1F (32y)	Concurrently Trained	45	13	31	9	0	2

MHC = Myosin heavy chain; M = Male; F = Female; y = Year; RE = Resistance exercise;
Non Ath = Not a competitive athlete; Con = Concentric, Ecc = Eccentric

Numerous other knowledge gaps persist because in over 50 years of human muscle FT

research only two studies have utilized SF with elite (i.e., world or international) athletes (one

male sprinter and six female track and field) and no research has done so with any strength or

power athlete. Thus, the purpose of this study was to examine the FT% of elite weightlifters to

provide novel insight into the phenotype of competitive female and male strength and power

athletes.

METHODS

<u>Experimental Approach to the Problem</u>

Twenty-one elite ('*Olympic'*) Weightlifters (15 female, 6 male) underwent resting *vastus*

lateralis biopsies between 2-96 hours after competing in either the International Weightlifting

Federation World Championships or the USA Weightlifting American Open Finals (2017). All

procedures and risks were explained to the athletes prior to obtaining written consent and

completing medical and exercise history questionnaires. Performance records in the Snatch and

Clean and Jerk (1RM), competition medals, and other accolades were gathered from personal

interviews and publically available records from these or other sanctioned meets. Each muscle

sample was analyzed for MHC content using two distinct FT techniques: single fiber (SF) and

homogenate (HG). The University Institutional Review Board approved all experimental

procedures prior to any testing.

<u>Participants</u>

Participants were subdivided into three categories:

1. Olympic or World-caliber (n = 6 female): WCF

2. National-caliber female (n = 9): NCF

3. National-caliber male (n = 6): NCM

Athletes were considered "World-caliber" if they were on the most recent Olympic or

World team and competed at the most recent National event. Athletes were considered "National-

caliber" if they were top 5 placers at the 2017 American Open Finals meet but had never been on

a World or Olympic team. Athletes spanned multiple weight categories, had a minimum of two

years of National competition experience, had competed exclusively for the United States of

America, and were otherwise eligible for all American National meets (Table 2). Athlete accolades

at the time of data collection included participation in 3 Olympic Games, 19 World

Championships, 11 Pan American Championships, 49 National Championships, 32 American

Opens, 8 University National Championships, and 25 Junior World/Pan American/National

Championships. Participants also held 25 National records and >170 National/International

medals. One athlete had tested positive for substances prohibited by the World Anti-Doping

Agency and was suspended from the sport for two years prior to participating in the study.

Table 2. Descriptive information of elite female and male American weightlifters.

	Age (y)	Body Mass (kg)	Height (cm)	Years Competing (y)	Snatch Relative 1RM	Clean & Jerk Relative 1RM
WCF	28.2 ± 3.6*	81.2 ± 36.0	164.0 ± 11.1	7.7 ± 4.7*‡	1.32 ± 0.31*	1.69 ± 0.40*
NCF	23.6 ± 3.9	66.6 ± 11.0	164.8 ± 7.0	3.8 ± 0.8	1.29 ± 0.18‡	1.68 ± 0.19‡
NCM	26.0 ± 2.4	85.3 ± 26.9	169.0 ± 9.0	3.3 ± .08	1.64 ± 0.25	2.04 ± 0.29
Average	25.6 ± 3.8	76.1 ± 25.0	165.8 ± 8.7	4.8 ± 3.1	1.40 ± 0.28	1.79 ± .032

Data are described as mean ± standard deviation. WCF = Olympic or World-caliber female (n = 6), NCF = National-caliber female (n

= 9), NCM = National-caliber male (n = 6). Relative 1RM = competition record one repetition maximum divided by body mass. Years

competing = number of years competing in USA Weightlifting sanctioned meets. * = significantly different than NCF. ‡ =

significantly different than NCM. Significant = $p < 0.05$.

Procedures

Vastus Lateralis Muscle Biopsies

Following 30 minutes of supine rest, athletes underwent a mid-muscle belly

(approximately halfway between the greater trochanter and patella) biopsy of the *vastus lateralis*.

A detailed description of the biopsy procedure has been previously described by our lab (9, 22, 23,

33). Briefly, a small area of the thigh was numbed by injection of a local anesthetic

(Xylocaine/Lidocaine without epinephrine). An approximately ¼ inch incision was made in the

superficial cutaneous tissues. Muscle samples were obtained using the Bergström technique with

suction (1), immediately cleansed of excess blood and connective tissue, divided into

approximately 10-15 mg strips, placed into cold skinning solution (125 mM K propionate, 2.0 mM

EGTA, 4.0 mM ATP, 1.0 mM $MgCl_2$, 20.0 mM imidazole [pH 7.0], and 50% [vol ml/vol ml]

glycerol), and stored at -20° C for at least one week. Each sample was split such that a portion (~5

mg) could be used for single fiber isolation or homogenization. The incision site was cleaned,

pulled closed with a sterile Band-Aid, and covered with sterile gauze and cohesive bandage tape.

Myosin Heavy Chain Fiber Type Identification

All biopsy samples were analyzed for MHC via SDS-PAGE using two distinct techniques:

single fiber *distribution* (SF) and homogenate *composition* (HG). For SF, individual fibers (N =

2,147; 102 ± 3 fibers per athlete) were mechanically isolated with fine tweezers under a light

microscope and placed in 80 μl of sodium dodecyl sulfate (SDS) buffer (1% SDS, 23 mM EDTA,

0.008% bromophenol blue, 15% glycerol, and 715 mM b-mercaptoethanol [pH 6.8]). HG samples

(~5 mg) were hand homogenized and then diluted between 1:10 to 1:50 based on sample amount

and protein quantity. As described in detail elsewhere (5, 9, 22, 23, 27), 1-2 μl aliquots of both SF

or HG (run separately) were then loaded into individual wells in a 3.5% loading and 5% separating

gel (SDS-PAGE), run at 5°C for 15.5 hours (SE 600 Series; Hoefer, San Francisco, CA, USA),

and silver stained for MHC identification. The SF approach used known molecular weights and

standards to identify the MHC isoform (MHC I, I/IIa, IIa, IIa/IIx, and IIx) of each individual

myofiber. This enabled the most accurate calculation of the distribution/percent frequency (FT%)

of each FT contained within the muscle sample (21). For example, if 100 fibers were analyzed and

30 were identified as MHC I, 60 as MHC IIa, and 10 as MHC IIa/IIx, the FT% would be calculated

as 30% MHC I, 60% MHC IIa, and 10% MHC IIa/IIx. HG utilized densitometry (ImageJ, National

Institutes of Health, Bethesda, MD) to quantify the relative MHC protein composition (i.e., percent

area occupied by each pure isoform; MHC I, IIa, and IIx) of each sample, which is correlated

highly with FT area (34). Thus, SF indicates how frequently each isoform exists but cannot address

how much area each FT occupies within the muscle. HG addresses the latter, but cannot delineate

hybrids, therefore inaccurately quantifying FT% (9, 21-25, 27).

Statistical Analysis

Potential differences between groups in descriptive information were examined via

ANOVA. For SF, potential differences in FT% between groups were assessed via a 3 (group:

WCF, NCF, NCM) x 4 (fiber type: MHC I, I/IIa, IIa, IIa/IIx) ANOVA. For HG, potential

differences in FT composition between groups were examined via a 3 (group: WCF, NCF, NCM)

x 3 (fiber type: MHC I, IIa, IIx) ANOVA. Comparison of SF vs. HG was accomplished by a 2

(group: SF, HG) x 3 (fiber type: MHC I, IIa, IIx) ANOVA. Effect size was calculated with Cohen's

D (0.2 = small difference, 0.5 = medium difference, and 0.8 = large difference) to identify the

magnitude of difference between two groups. Pearson Product Moment Correlations (*r*) were

assessed for WCF, NCF, and NCM between 1RM, body mass, and SF FT%. All individual FT

data are reported in Table 3. Data are reported as mean ± standard deviation (SD), unless otherwise

noted. Significance was established *a priori* at an alpha level of *p < 0.05*. All analyses were

performed with SPSS (SPSS Statistics Version 24, IBM).

RESULTS

Descriptive

WCF were significantly older than NCF, but not NCM (Table 2). WCF also had

significantly more years of sport competition experience than NCF and NCM. Yet, NCM exceeded

both WCF and NCF in relative strength in both the Snatch 1RM and Clean and Jerk 1RM.

Table 3. Individual single muscle fiber type distribution (SF) and homogenate composition (HG)

of elite female and male American weightlifters**.** Data are reported as a percentage.

	MHC I		MHC I/IIa	MHC IIa		MHC IIa/IIx	MHC IIx	
Athlete	SF%	HG%	SF%	SF%	HG%	SF%	SF%	HG%
World-Caliber Female Weightlifters								
1	13	31	7	74	69	7	0	0
2	39	34	13	48	66	0	0	0
3	18	21	2	79	79	1	0	0
4	9	19	2	89	81	0	0	0
5	12	22	4	85	78	0	0	0
6*	10	17	9	52	70	28	0	13

National-Caliber Female Weightlifters

7	23	38	1	76	62	0	0	0
8*	14	18	1	63	68	22	0	15
9	32	43	6	62	57	0	0	0
10	29	39	8	58	61	4	0	0
11	20	26	2	78	74	0	0	0
12	29	46	5	66	54	0	0	0
13	25	30	2	73	70	0	0	0
14	19	25	6	74	75	0	0	0
15	34	35	6	56	65	4	0	0

National-Caliber Male Weightlifters

16	29	40	8	63	60	0	0	0
17	26	37	1	73	63	0	0	0
18	32	44	3	65	56	0	0	0
19	7	18	3	84	82	6	0	0
20*	26	36	3	54	57	17	0	7
21*	29	33	2	37	49	32	0	18

MHC = myosin heavy chain. * Denotes athlete in the heavyweight (or super) (>90 kg for

women and >105 kg for men) category.

Single Fiber Analysis (SF)

FT% for all lifters combined was 23 ± 9% I, 5 ± 3% I/IIa, 67 ± 13% IIa, and 6 ± 10%

IIa/IIx. No MHC IIx or I/IIa/IIx fibers were identified. No significant differences existed between

groups, despite WCF possessing 8% (absolute, not percent difference) less MHC I than NCF (d =

0.88) and NCM (d = 0.78) (Figure 1). The difference in MHC IIa between WCF and NCM (also

8%) was also not statistically significant, but had a moderate effect size (d = 0.50). The vast

majority of the MHC IIa/IIx fibers (91%) belonged to just five lifters, all of whom competed in

the heavyweight or super heavyweight categories (≥ 90 kg for women and ≥105 kg for men). Thus,

significant correlations were present between body mass and MHC IIa/IIx frequency for WCF (*r*

= 0.919, p = 0.010) and NCF (*r* = 0.826, p = 0.006) while a trend existed for NCM (*r* = 0.757, p =

0.080).

Figure 1. Myosin heavy chain (MHC) fiber type distribution of elite American weightlifters.

WCF = Olympic or World-caliber female (n = 6), NCF = National-caliber female (n = 9), NCM =

National-caliber male (n = 6). Data are reported as a percentage ± standard deviation.

Homogenate Analysis (HG)

FT composition for all lifters combined was 31 ± 9% I, 67 ± 9% IIa, and 3 ± 6% IIx. MHC

I tended (p = 0.08) to be lower in WCF (24 ± 7%) than NCF (33 ± 9%, p = 0.125, d = 1.33) and

NCM (35 ± 9%, p = 0.106, d = 1.14), yet MHC IIa was significantly higher (p = 0.046) in WCF

(74 ± 6%) than NCF (65 ± 7%, p = 0.145, d = 1.28) and NCM (61 ± 11%, p = 0.043, d = 1.39). FT

was significantly different (p < 0.001) between SF and HG for MHC I (p = 0.005) and MHC IIx

(p = 0.046), but not MHC IIa. SF MHC IIa/IIx and HG MHC IIx were highly correlated ($r = 0.96$,

p < 0. 001). No correlations existed for SF or HG between FT% and Snatch or Clean and Jerk

relative 1RM.

DISCUSSION

The current study resulted in the most detailed investigation of muscle phenotype in

Olympic and World-caliber anaerobic athletes published to date. Additionally, the data enabled

the first comparison and differentiation of World vs. National-caliber athletes at the single fiber

level. The current study was also the most precise description of FT% in strength or power sport

competitors, and the first ever in females. The MHC IIa abundance was the highest in healthy

muscle (VL) ever reported, especially for females. These data suggest athlete caliber and/or

training history influences FT% more than sex *per se* and also questions the pronouncement that

male athletes possess more fast-twitch myofibers than females. Our utilization of two different

typing methods confirmed the limitations of HG for FT% (inappropriately categorizes MHC I/IIa

as MHC I and MHC IIa/IIx as MHC IIx) and also allowed identification of a previously

undocumented relationship between body mass and MHC IIa/IIx concentrations. The unique

morphology and phenotypes in our participants highlight the need to further study elite anaerobic

athletes, particularly females.

WCF contained the highest concentration of MHC IIa (71%) ever reported in the literature

to our knowledge. NCF (67%) and NCM (63%) also possessed more MHC IIa than previous

research in competitive bodybuilders (40%) (16, 19) as well as power/weightlifters (8, 11, 12, 15,

16, 18), elite track and field athletes (20, 35), and resistance-trained men (18, 19, 22, 23, 27, 29,

36), which all ranged from 50-60%. Only six previous studies using SF have found pure MHC IIa

concentrations of >50%, with just two reporting 60% (Table 1). The resulting minimal MHC I

(~17-25%) in our athletes was strikingly lower than elite female track and field athletes (57%) (20)

and National-caliber bodybuilders (35%) (19). These pronounced differences are likely explained

by the substantial dissimilarities in training styles (e.g., external loading strategies, contraction

type and velocity, training frequency, etc.) between the various sports. More research is therefore

needed to continue delineating the subtle but significant differences in FT% between top-

performing athletes in various anaerobic sports and the specific role each training approach might

play on altering MHC I and IIa distribution. Although it did not reach statistical significance, large

ES were evident and MHC IIa frequencies of 74%-89% occurred in 66% of WCF but only in 44%

and 33% of NCF and NCM, respectively. Thus, scientists should continue to examine what

separates World from National-level athletes.

WCF differed from NCF and NCM in both sex and years competing in the sport (~8 vs. 3

y). Sex comparisons in athletes remains tenuous (31, 37) because nearly all investigations utilize

non-gold standard FT% methods (21) and sedentary (38, 39) or "recreationally active" individuals

(32). Not only do our findings contradict the claim that women possess more slow-twitch

myofibers than men (40), they illustrate the opposite when accounting for talent level (WCF <

NCF = NCM). The current cross-sectional study-design precludes direct analysis, but extensive

research affords strong support for training history as a critical determinate of FT% (2, 9, 26, 28,

30, 41-44). Chronic exercise generally decreases hybrids (30, 42) and induces style-specific shifts

in FT% such as increases in MHC I with endurance (9, 43) or MHC IIa with sprint (28), plyometric

(26), or strength training (36, 43-45). For example, MHC I concentrations in an individual with

extensive endurance exercise history were nearly double that of his non-exercising monozygous

twin (9). Another study reported an increase in MHC IIa from 46% to 60% following 19 weeks of

resistance training (36). MHC IIa/IIx fibers appear particularly responsible for exercise-induced

increases in MHC IIa and are thus uncommon in exercise-trained individuals (9, 20, 22-25, 29,

43). A reduction of MHC IIx in favor of IIa following chronic resistance exercise is also purported

extensively in the literature (28, 34), yet the overwhelming majority of this evidence comes from

experiments with methodologies directly shown here and elsewhere (9, 24, 25) to produce

erroneous FT% conclusions.

Most research from the 1970's – 2000's utilized either ATPase histochemistry or HG SDS-

PAGE to determine FT% (8, 14-17, 24, 25, 34, 36). Similar to SF, histochemistry allows

assessment of individual fibers for calculation of percent distribution, yet it does not enable

simultaneous delineate of hybrids (36). HG suffers the same drawback and actually indicates FT

area/composition (34) more so than distribution making it greatly influenced by the size of each

fiber; which is not uniform across all FT (particularly in resistance trained individuals) (46). All

three approaches hold strong merit and are often correlated to each other (34, 47) and performance

(8), but are clearly not interchangeable for maximally precise FT% assessment. In the current

study, HG accurately quantified MHC IIa (within 0-4%), but not I or IIx. MHC I was overestimated

by 8% percent (23 vs. 31%), which is largely explained by the non-differentiated MHC I/IIa fibers

(5%). HG also greatly exaggerated MHC IIx, particularly in individuals with >4% MHC IIa/IIx.

The inability of HG to account for MHC IIa/IIx explains why MHC IIx appear common in some

studies (48) even though their actual abundance in healthy human skeletal muscle is extraordinarily

rare; typically <0.1% (9, 22-25, 27) and 0 of the >2,100 isolated fibers from the current sample.

Thus, the seeming conversion of MHC IIx to IIa with exercise is more precisely IIa/IIx changing

to IIa.

288  For these reasons MHC IIa/IIx are typically inversely associated with muscle health and

289 physical activity (9, 43). Yet, the heavyweights (male and female) expressed irregularly high

290 concentrations (24%) and accounted for 91% of all MHC IIa/IIx myofibers. Terzis and colleagues

291 (2010) noted a similar abnormal abundance of MHC IIx (typed via HG, so likely IIa/IIx) in six

292 large (116 kg, body fat composition >22%), but presumably well strength-trained throwers (35).

293 Body composition was not assessed in the current study and little research exists on well-trained,

294 but obese individuals. Thus, additional studies across a broader spectrum of physical size are

295 required to truly interpret the correlations between body mass and MHC IIa/IIx prevalence.

296  Another juxtaposition was that of FT% and performance. Previous work in 94 kg male

297 competitive weightlifters found strong correlations between FT composition (HG) and percent FT

298 area to both snatch 1RM and vertical jump height (8), but not clean and jerk 1RM. We failed to

299 identify any such correlations, but also utilized multiple sexes and weight classes. Thus, while

300 FT% differed between our groups, that factor alone did not predict performance among our lifters.

301 Several possible explanations exist for this discrepancy. First, FT area may determine whole

302 muscle strength more than FT%. Second, neither studies found correlations to the clean and jerk,

303 which is heavier and slower than the snatch or vertical jump. This compliments previous isokinetic

304 research (23) and indicates FT% does not predict performance on strength tasks among strength-

305 trained individuals. FT% probably determines movement speed more than force production (7).

306 Further speculation on this point is unwarranted as limitations prohibited the ability to assess FT-

307 specific size or contractile properties, which likely differed significantly across our groups (49)

308 and are known to changes with training (3, 46).

CONCLUSION

This study provides novel insight into the muscle phenotype of elite competitive strength

and power athletes. Our data indicate athlete caliber, training history, and body mass dictate FT%

more than sex *per se*, but more work is needed to draw firm conclusions. The extreme fast-twitch

abundance partially explains how elite weightlifters are able to generate high forces in short time-

frames. Most athletes contained few hybrids and no MHC IIx or I/IIa/IIx, except the heavyweights

who possessed large quantities of IIa/IIx. Future research should use high fidelity techniques to

explore FT-specific distribution, size, and contractile properties in female and male athletes of

various caliber, sports, and body size; ideally across several years of competition. The resulting

knowledge could have practical significance if it enabled experimentation of differing training

volumes or recovery protocols based on athlete-specific FT properties (50). More detailed fiber

type profiles of elite strength and power athletes may eventually enable strength and conditioning

professionals to create more individualized and effective programs.

ACKNOWLEDGEMENTS

Funding for this project was provided by Renaissance Periodization. The authors would like to

thank Irene S. Tobias and Cameron Yen for their help with this project. Contact the corresponding

author at agalpin@fullerton.edu.

REFERENCES

1.   Bergstrom J. Muscle electrolytes in man determined by neutron activation analysis on needle biopsy specimens. The Scandinavian Journal of Clinical and Laboratory Investigations. 1962;14 (Suppl. 68).

2.   Pette D, Staron RS. Myosin isoforms, muscle fiber types, and transitions. Microsc Res Tech. 2000;50(6):500-9.

3.   Methenitis S, Karandreas N, Spengos K, Zaras N, Stasinaki AN, Terzis G. Muscle Fiber Conduction Velocity, Muscle Fiber Composition, and Power Performance. Med Sci Sports Exerc. 2016;48(9):1761-71.

4.   Galpin AJ, Raue U, Jemiolo B, Trappe TA, Harber MP, Minchev K, et al. Human skeletal muscle fiber type specific protein content. Anal Biochem. 2012;425(2):175-82.

5.   Tobias IS, Lazauskas KK, Arevalo JA, Bagley JR, Brown LE, Galpin AJ. Fiber type-specific analysis of AMPK isoforms in human skeletal muscle: advancement in methods via capillary nano-immunoassay. J Appl Physiol (1985). 2017.

6.   Bagley JR. Fibre type-specific hypertrophy mechanisms in human skeletal muscle: potential role of myonuclear addition. J Physiol. 2014;592(23):5147-8.

7.   Bottinelli R, Pellegrino MA, Canepari M, Rossi R, Reggiani C. Specific contributions of various muscle fibre types to human muscle performance: an in vitro study. J Electromyogr Kinesiol. 1999;9(2):87-95.

8.   Fry AC, Schilling BK, Staron RS, Hagerman FC, Hikida RS, Thrush JT. Muscle fiber characteristics and performance correlates of male Olympic-style weightlifters. J Strength Cond Res. 2003;17(4):746-54.

9.   Bathgate KE, Bagley JR, Jo E, Talmadge RJ, Tobias IS, Brown LE, et al. Muscle health and performance in monozygotic twins with 30 years of discordant exercise habits. Eur J Appl Physiol. 2018.

10.   Costill DL, Daniels J, Evans W, Fink W, Krahenbuhl G, Saltin B. Skeletal muscle enzymes and fiber composition in male and female track athletes. J Appl Physiol. 1976;40(2):149-54.

11.   Tesch PA, Karlsson J. Muscle fiber types and size in trained and untrained muscles of elite athletes. J Appl Physiol (1985). 1985;59(6):1716-20.

12.   Prince FP, Hikida RS, Hagerman FC. Human muscle fiber types in power lifters, distance runners and untrained subjects. Pflugers Arch. 1976;363(1):19-26.

13.   Trappe S, Luden N, Minchev K, Raue U, Jemiolo B, Trappe TA. Skeletal muscle signature of a champion sprint runner. J Appl Physiol (1985). 2015;118(12):1460-6.

14.   Gollnick PD, Armstrong RB, Saubert CWt, Piehl K, Saltin B. Enzyme activity and fiber composition in skeletal muscle of untrained and trained men. J Appl Physiol. 1972;33(3):312-9.

15.   Tesch PA, Thorsson A, Kaiser P. Muscle capillary supply and fiber type characteristics in weight and power lifters. J Appl Physiol Respir Environ Exerc Physiol. 1984;56(1):35-8.

16.   Tesch PA, Larsson L. Muscle hypertrophy in bodybuilders. Eur J Appl Physiol Occup Physiol. 1982;49(3):301-6.

17.   Fry AC, Webber JM, Weiss LW, Harber MP, Vaczi M, Pattison NA. Muscle fiber characteristics of competitive power lifters. J Strength Cond Res. 2003;17(2):402-10.

18.   Galpin AJ, Fry AC, Chiu LZ, Thomason DB, Schilling BK. High-power resistance exercise induces MAPK phosphorylation in weightlifting trained men. Appl Physiol Nutr Metab. 2012;37(1):80-7.

19.   Kesidis N, Metaxas TI, Vrabas IS, Stefanidis P, Vamvakoudis E, Christoulas K, et al.
Myosin heavy chain isoform distribution in single fibres of bodybuilders. Eur J Appl Physiol.
2008;103(5):579-83.
20.   Parcell AC, Sawyer RD, Craig Poole R. Single muscle fiber myosin heavy chain
distribution in elite female track athletes. Med Sci Sports Exerc. 2003;35(3):434-8.
21.   Pandorf CE, Caiozzo VJ, Haddad F, Baldwin KM. A rationale for SDS-PAGE of MHC
isoforms as a gold standard for determining contractile phenotype. J Appl Physiol (1985).
2010;108(1):222-2; author reply 6.
22.   Arevalo JA, Lynn SK, Bagley JR, Brown LE, Costa PB, Galpin AJ. Lower-Limb
Dominance, Performance, and Fiber Type in Resistance-trained Men. Med Sci Sports Exerc.
2017.
23.   Bagley JR, McLeland KA, Arevalo JA, Brown LE, Coburn JW, Galpin AJ. Skeletal
Muscle Fatigability and Myosin Heavy Chain Fiber Type in Resistance Trained Men. J Strength
Cond Res. 2017;31(3):602-7.
24.   Andersen JL, Klitgaard H, Bangsbo J, Saltin B. Myosin heavy chain isoforms in single
fibres from m. vastus lateralis of soccer players: effects of strength-training. Acta Physiol Scand.
1994;150(1):21-6.
25.   Andersen JL, Klitgaard H, Saltin B. Myosin heavy chain isoforms in single fibres from
m. vastus lateralis of sprinters: influence of training. Acta Physiol Scand. 1994;151(2):135-42.
26.   Malisoux L, Francaux M, Nielens H, Renard P, Lebacq J, Theisen D. Calcium sensitivity
of human single muscle fibers following plyometric training. Med Sci Sports Exerc.
2006;38(11):1901-8.
27.   Murach KA, Bagley JR, McLeland KA, Arevalo JA, Ciccone AB, Malyszek KK, et al.
Improving human skeletal muscle myosin heavy chain fiber typing efficiency. J Muscle Res Cell
Motil. 2016;37(1-2):1-5.
28.   Parcell AC, Sawyer RD, Drummond MJ, O'Neil B, Miller N, Woolstenhulme MT.
Single-fiber MHC polymorphic expression is unaffected by sprint cycle training. Med Sci Sports
Exerc. 2005;37(7):1133-7.
29.   Raue U, Terpstra B, Williamson DL, Gallagher PM, Trappe SW. Effects of short-term
concentric vs. eccentric resistance training on single muscle fiber MHC distribution in humans.
Int J Sports Med. 2005;26(5):339-43.
30.   Williamson DL, Gallagher PM, Carroll CC, Raue U, Trappe SW. Reduction in hybrid
single muscle fiber proportions with resistance training in humans. J Appl Physiol (1985).
2001;91(5):1955-61.
31.   Haizlip KM, Harrison BC, Leinwand LA. Sex-based differences in skeletal muscle
kinetics and fiber-type composition. Physiology (Bethesda). 2015;30(1):30-9.
32.   Norman B, Esbjornsson M, Rundqvist H, Osterlund T, von Walden F, Tesch PA.
Strength, power, fiber types, and mRNA expression in trained men and women with different
ACTN3 R577X genotypes. J Appl Physiol (1985). 2009;106(3):959-65.
33.   Bagley JR, Galpin AJ. Three-dimensional printing of human skeletal muscle cells: An
interdisciplinary approach for studying biological systems. Biochem Mol Biol Educ.
2015;43(6):403-7.
34.   Fry AC, Allemeier CA, Staron RS. Correlation between percentage fiber type area and
myosin heavy chain content in human skeletal muscle. Eur J Appl Physiol Occup Physiol.
1994;68(3):246-51.

35.   Terzis G, Spengos K, Kavouras S, Manta P, Georgiadis G. Muscle fibre type composition
and body composition in hammer throwers. J Sports Sci Med. 2010;9(1):104-9.
36.   Adams GR, Hather BM, Baldwin KM, Dudley GA. Skeletal muscle myosin heavy chain
composition and resistance training. J Appl Physiol (1985). 1993;74(2):911-5.
37.   Prince FP, Hikida RS, Hagerman FC. Muscle fiber types in women athletes and non-
athletes. Pflugers Arch. 1977;371(1-2):161-5.
38.   Miller MS, Callahan DM, Tourville TW, Slauterbeck JR, Kaplan A, Fiske BR, et al.
Moderate-intensity resistance exercise alters skeletal muscle molecular and cellular structure and
function in inactive older adults with knee osteoarthritis. J Appl Physiol (1985).
2017;122(4):775-87.
39.   Simoneau JA, Bouchard C. Human variation in skeletal muscle fiber-type proportion and
enzyme activities. Am J Physiol. 1989;257(4 Pt 1):E567-72.
40.   Welle S, Tawil R, Thornton CA. Sex-related differences in gene expression in human
skeletal muscle. PLoS One. 2008;3(1):e1385.
41.   Pette D, Staron RS. Mammalian skeletal muscle fiber type transitions. Int Rev Cytol.
1997;170:143-223.
42.   Pette D. The adaptive potential of skeletal muscle fibers. Can J Appl Physiol.
2002;27(4):423-48.
43.   Bagley JR, Murach KA, Trappe S. Microgravity-Induced Fiber Type Shift in Human
Skeletal Muscle Gravitational and Space Biology. 2012;26(1):34-40.
44.   Liu Y, Schlumberger A, Wirth K, Schmidtbleicher D, Steinacker JM. Different effects on
human skeletal myosin heavy chain isoform expression: strength vs. combination training. J
Appl Physiol (1985). 2003;94(6):2282-8.
45.   Kryger AI, Andersen JL. Resistance training in the oldest old: consequences for muscle
strength, fiber types, fiber size, and MHC isoforms. Scand J Med Sci Sports. 2007;17(4):422-30.
46.   Grgic J, Schoenfeld BJ. Are the Hypertrophic Adaptations to High and Low-Load
Resistance Training Muscle Fiber Type Specific? Front Physiol. 2018;9:402.
47.   Staron RS. Correlation between myofibrillar ATPase activity and myosin heavy chain
composition in single human muscle fibers. Histochemistry. 1991;96(1):21-4.
48.   Aagaard P, Magnusson PS, Larsson B, Kjaer M, Krustrup P. Mechanical muscle
function, morphology, and fiber type in lifelong trained elderly. Med Sci Sports Exerc.
2007;39(11):1989-96.
49.   Staron RS, Hagerman FC, Hikida RS, Murray TF, Hostler DP, Crill MT, et al. Fiber type
composition of the vastus lateralis muscle of young men and women. J Histochem Cytochem.
2000;48(5):623-9.
50.   Jones N, Kiely J, Suraci B, Collins DJ, de Lorenzo D, Pickering C, et al. A genetic-based
algorithm for personalized resistance training. Biol Sport. 2016;33(2):117-26.
461

Figure 1