Activation of rod input in a in a model of retinal degeneration reverses retinal remodeling and induces formation of normal synapses, circuitry and visual signaling in the adult retina

Running title: Restoration of visual signaling in the remodeled retina

Tian Wang[1], Johan Pahlberg[2§], Jon Cafaro[3§],
Alapakkam P. Sampath[2*], Greg D. Field[3*], and Jeannie Chen[1*]

Zilkha Neurogenetic Institute, Department of Physiology and Neuroscience, Keck School of Medicine, University of Southern California, Los Angeles, CA 90089
Department of Ophthalmology, Stein Eye Institute, University of California, Los Angeles, CA 90095
Department of Neurobiology, Duke University School of Medicine, Durham, NC, USA

[§]These authors contributed equally to this work

*Correspondence should be addressed to either A.P. Sampath, G. Field or J. Chen.
Email: asampath@jsei.ucla.edu, field@neuro.duke.edu, jeannie@usc.edu

Acknowledgements: This work was supported by National Institute of Health grants EY027193 (APS, GDF, and JC); EY12155 and EY027387 (JC); an unrestricted grant from Research to Prevent Blindness to the Department of Ophthalmology, UCLA; and Jules Stein Eye Institute Core Grant EY00331 (APS). We thank Dr. M. Scalabrino for comments on the manuscript, Dr. K.Martemyanov for providing the mGluR6 antibody, Dr. C. Craft for providing the cone arrestin (ARR3) antibody and Dr. S. Ruffins at the USC microscopy core for his help with confocal imaging.

Conflict of Interest: The authors declare no competing financial interests.

Key words: Neural plasticity, neural transmission, retinal circuitry, retinal degeneration, gene therapy, cGMP-gated channel, photoreceptor cell death.

Abstract

A major cause of human blindness is the death of rod photoreceptors. As rods degenerate, synaptic structures between rod and rod bipolar cells dissolve and the rod bipolar cells extend their dendrites and occasionally make aberrant contacts. Such changes are broadly observed in blinding disorders caused by photoreceptor cell death and is thought to occur in response to deafferentation. How the remodeled retinal circuit affect visual processing following rod rescue is not known. To address this question, we generated transgenic mice wherein a disrupted cGMP-gated channel (CNG) gene can be repaired at the endogenous locus and at different stages of degeneration by tamoxifen-inducible cre-mediated recombination. In normal rods, light-induced closure of CNG channels leads to hyperpolarization of the cell, reducing neurotransmitter release at the synapse. Similarly, rods lacking CNG channel exhibit a resting membrane potential that was ~10mV hyperpolarized compared to WT rods, indicating diminished glutamate release. Retinas from these mice undergo stereotypic retinal remodeling as a consequence of rod malfunction and degeneration. Upon tamoxifen-induced expression of CNG channels, rods recovered their structure and exhibited normal light responses. Moreover, we show that the adult mouse retina displays a surprising degree of plasticity upon activation of rod input. Wayward bipolar cell dendrites establish contact with rods to support normal synaptic transmission, which is propagated to the retinal ganglion cells. These findings demonstrate remarkable plasticity extending beyond the developmental period and support efforts to repair or replace defective rods in patients blinded by rod degeneration.

Significance Statement

Current strategies for treatment of neurodegenerative disorders are focused on the repair of the primary affected cell type. However, the defective neuron functions within a complex neural circuitry, which also becomes degraded during disease. It is not known whether a rescued neuron and the remodeled circuit will establish communication to regain normal function. We show that the adult mammalian neural retina exhibits a surprising degree of plasticity following rescue of rod photoreceptors. The wayward rod bipolar cell dendrites re-establish contact with rods to support normal synaptic transmission, which is propagated to the retinal ganglion cells. These findings support efforts to repair or replace defective rods in patients blinded by rod cell loss.

Introduction

Diseases that afflict sensory systems typically result from deficiencies within the sensory receptor cells themselves, either within sensory transduction or synaptic transmission (Bermingham-McDonogh and Reh, 2011). Deficits in visual processing are no exception, with the majority of blinding diseases resulting from the dysfunction or death of the primary input cells, the retinal rod and cone photoreceptors (Quartilho et al., 2016). Synaptic remodeling of retinal circuits, in particular between photoreceptor cells and their downstream neurons, occur early in retinal degeneration (Soto and Kerschensteiner, 2015). Remodeling of bipolar and horizontal cell dendrites is thought to occur in response to deafferentation (Marc and Jones, 2003). Changes that occur include homeostatic down-regulation of synaptic structures, exuberant extension of dendritic processes which sometimes contact off-target sites (Marc and Jones, 2003; Puthussery and Taylor, 2010), and even switching of post-synaptic receptor types from mGluR to iGluR expression (Chua et al., 2009). In genetically inherited forms of retinal degeneration, synaptic changes may already occur during a critical period of retinal development. It is not known how these changes in retinal circuitry may ultimately limit recovery of normal vision, although several approaches are being implemented to rescue dying photoreceptors using gene therapy, or replace them with stem cells (Scholl et al., 2016; Garg et al., 2017; Yao et al., 2018). To address this gap in knowledge, this study focuses on cellular plasticity in retinal circuits of young adult mice with rod degeneration, and how the synaptic structures and circuits that receive rod input respond to rod rescue.

We genetically engineered a mouse line in which rod function can be uniformly rescued via tamoxifen-induced cre-mediated recombination. The line was generated to lack expression of the cyclic nucleotide gated (CNG) channel beta-1 subunit (CNGB1) due to an insertion of a neomycin cassette at the endogenous gene to disrupt expression (Wang and Chen, 2014; Wang et al., 2017a). This mouse model recapitulates the effects of mutations in human *CNGB1* and *CNGA1* genes that cause autosomal recessive retinitis pigmentosa (Biel and Michalakis, 2007). Without the CNGB1 subunit, the CNG channels in rod outer segments fail to form normally functioning channels, which leads to a slow form of rod death that occurs over 4-6 months (Zhang et al., 2009; Wang et al., 2017a), or longer (Hüttl et al., 2005). Importantly, the neomycin cassette is flanked by loxP sites, which allows for cre-mediated excision and the expression of CNGB1 from the endogenous locus. Thus, this mouse line provides an opportunity to introduce precisely a 'cure' for the underlying genetic defect at different time points during degeneration.

We use this novel mouse line to determine the extent to which activating rod input in the
degenerating retina allows recovery of the structure and function of well-defined rod-driven
retinal circuits in young adult mice. The lack of CNG channels caused stereotypic degenerative
changes in the retina that included rhodopsin mislocalization, activation of Müller glia, and a
reduction of pre- and post-synaptic proteins between rods and rod bipolar cells by as early as 4
weeks of age. Signal transmission from rods to rod bipolar cells was abrogated and sensitivity of
retinal ganglion cells (RGCs) was reduced ~100-fold. Tamoxifen-induced restoration of CNG
channel expression initiated at 4 weeks of age led to an expected recovery of rod photoreceptor
function. Importantly, we show that initiation of rod input in the deafferented adult retina also
induced a high degree of structural plasticity between rods and their primary postsynaptic
partner, rod (ON) bipolar cells. Specifically, rod bipolar cell dendrites sprouted fine tips and
mGluR6 clusters formed on these tips which made new synapses with rods. This structural
transformation resulted in near-normal light responses in both bipolar cells and retinal ganglion
cells, the output neurons of the retina. Our findings indicate substantial plasticity in the adult
mammalian retina, suggesting favorable outcomes for interventions targeting the rescue of
dysfunctional rods from death.

Materials and Methods
Generation of transgenic mice. The use of mice in these experiments was in accordance with
the National Institutes of Health guidelines and the Institutional Animal Care and Use Committee
of our respective universities. Targeting of the neoloxP to the *Cngb1* locus in mouse embryonic
stem cells and generation of transgenic mice from verified stem cell clones were described
previously (Chen et al., 2010). The CAGGCre-ER^{TM} transgenic line, Tg(CAG-cre/Esr1*)5Amc/J,
was obtained from the Jackson Laboratory and crossed with *Cngb1$^{neo/neo}$* mice.

Tamoxifen treatment. One-hundred mg tamoxifen was dissolved in 500 µl 95% ethanol and
diluted with 4.5 ml corn oil to give final concentration of 20 mg/ml. Four-week-old cre-positive
were given a dose of 3 mg/25 g body weight by oral gavage for 4 or 7 consecutive days. In
control experiments shown in Fig. 1, some cre-negative mice did not receive tamoxifen. For all
other experiments, cre-negative littermate mice were also treated with tamoxifen to control for
the possible effect of tamoxifen on photoreceptor cell survival (Wang et al., 2017b).

PCR genotyping. Genomic DNA was isolated from the neural retina. Three PCR primers were
used to detect the presence or absence of the neoloxP cassette. Primer 1 sequence

(GTTTTATGTAGCAGAGCAGGGAC) is located on intron 19, primer 2 sequence
(GAGGAGTAGAAGGTGGCGC) is on neoloxP, and primer 3 sequence
(CCACTCCTTAGTACATACCTAAGC) is located on exon 20. Product size of 620 bp from
primer pairs (2+3) indicates the presence of neoloxP, and a 802 bp PCR band from primer pairs
(1+3) indicates the absence of the neoloxP insert.

Retinal morphology. Mice were rendered unconscious by isofluorane inhalation and
immediately followed by cervical dislocation. Retinal sections were prepared as previously
described (Concepcion and Chen, 2010; Wang and Chen, 2014). Briefly, before enucleation,
eyes were marked for orientation by cauterization on the superior aspect of the cornea. Eyes
were placed in ½ Karnovsky buffer (2.5% glutaraldehyde, 2% formaldehyde in 0.1 M cacodylate
buffer, pH 7.2). The cornea and lens were removed, and the remaining eyecup was further fixed
overnight. Fixed eyes were rinsed in 0.1 M cacodylate buffer, fixed for 1 h in 1% OsO_4,
dehydrated in graded EtOH and embedded in epoxy resin. Eyecups were hemi-sected along the
superior-inferior axis, and one μm sections along the central meridian were obtained for light
micrographs.

Immunocytochemistry. Eyecups were prepared as described above, except the tissues were
dissected in cold 4% formaldehyde in PBS and further fixed for 15 min on ice. For frozen
sections, eyecups were rinsed in cold PBS, placed in 30% sucrose for 1 h, embedded in Tissue-
Tek® O.C.T. Compound (Sakura® Finetek) and flash frozen in liquid N_2. Ten μm frozen
sections were obtained. For retinal flat mounts, four relaxing cuts (0^o, 90^o, 180^o, 270^o) were
made on the edge of the neural retina and the flattened tissue was immobilized on a piece of
nitrocellulose membrane (Whatman [®], GE Healthcare Life Sciences), photoreceptor side down,
as described (Anastassov et al., 2017). The tissues were incubated with the following
antibodies: rhodopsin 1D4 (generously provided by R. Molday), GFAP (AB5804, Millipore),
CtBP2 (612044, BD Biosciences), PKC (ab32376, Abcam), mGluR6 (generously provided by K.
Martemyanov), ARR3 (generously provided by C. Craft). Images were acquired on a Zeiss
LSM800 confocal microscope. For quantifications of mGluR6 puncta, images were imported into
Fiji (ImageJ2), adjusted to similar threshold and the number and areas of puncta were quantified
using the analyze particles function.

Western blots. Each isolated retina was homogenized in 150 µl buffer (150mM NaCl, 50mM Tris pH 8.0, 0.1% NP-40, 0.5% deoxycholic acid, 0.1 mM PMSF and complete mini protease inhibitor (Roche Applied Sciences), incubated with DNase I (30U, Roche Applied Sciences) at room temperature for 30 min. An equal amount of retinal homogenate from each sample was electrophoresed on 4-12% Bis-Tris SDS-PAGE Gel (Invitrogen). Protein was transferred onto nitrocellulose membrane (Whatman ®, GE Healthcare Life Sciences) and incubated overnight with the following primary antibodies: rabbit anti-PDE polyclonal antibody (PAB-06800, Cytosignal), rabbit anti-ROS-GC1 polyclonal antibody (sc50512, Santa Cruz), mouse Anti-$G_t\alpha$ antibody (371740, EMD4Biosciences), rabbit polyclonal anti-GCAP1 and GCAP2 antibodies (Hoyo et al., 2014; Wang and Chen, 2014), mouse anti-CNGB1 4B1 antibody (Poetsch et al., 2001), mouse anti-CNGα antibody PMc 1D1 (Cook et al., 1989), mouse NCKX1 8H6 antibody (Vinberg et al., 2015) and mouse anti-Actin antibody (MAB1501, Millipore). The membranes were then incubated with fluorescently labeled secondary antibodies (1:10,000, LI-COR biosciences, 926-31081) at room temperature for 1 hour and detected by Odyssey infrared imaging system.

Whole retina and single-cell recordings from rods and bipolar cells. Mice were maintained on a normal 12-hour day-night cycle and were dark-adapted overnight (>12-h) prior to experiments. All further manipulations were performed in total darkness under infrared illumination visualized with infrared image converters (BE Meyers, WA). Following euthanasia, eyes were enucleated, the lens and cornea were removed, and eyecups were stored in darkness at 32°C in Ames' media buffered with sodium bicarbonate (Sigma, Cat# A1420) equilibrated with 5% CO_2/ 95% O_2.

Trans-retinal electroretinograms (ERGs) were recorded from isolated retinas as described previously (Pahlberg et al., 2017). Retinas were mounted photoreceptor side-up over a machined hole in a recording chamber. The tissue was superfused in darkness with 35-37°C Ames' media buffered with sodium bicarbonate and equilibrated with 5% CO_2/ 95% O_2 (pH ~ 7.4). An additional 10 mM of BaCl was added to the solution facing the inner retina to mitigate Müller cell activity. The *trans*-retinal potential change to flashes of light, delivered from a standard light bench, was measured using Ag/AgCl half-cells connected to a differential amplifier (Model DP-311; Warner Instruments). Recordings were sampled at 1 kHz and low-pass filtered at 30 Hz.

Recordings of the photovoltage from individual rods and rod bipolar cells was made by whole-cell patch clamp from dark-adapted retinal slices as described previously (Pahlberg et al.,

2017). Briefly, a small piece of dark-adapted retina was embedded in low-gelling temperature

agar, slices were cut on a vibrating microtome, transferred into a recording chamber, and

superfused with Ames' media equilibrated with $5\%CO_2/95\%O_2$ while maintained at $35\text{-}37^{\circ}C$.

The pipette internal solution consisted of (in mM): 125 K-Aspartate, 10 KCl, 10 HEPES, 5 N-

methyl glucamine-HEDTA, 0.5 $CaCl_2$, 1 ATP-Mg, 0.2 GTP-Mg; pH was adjusted to 7.2 with N-

methyl glucamine hydroxide. Light-evoked responses were recorded following the delivery of 10

ms flashes from a blue LED ($\lambda_{max} \sim 470$ nm, full width half maximum ~ 30 nm) whose strength

varied from producing a just-measurable response, and increased by factors of 2. Recordings

were sampled at 1 kHz and low-pass filtered at 300 Hz.

Retinal ganglion cell recording, stimulation and analysis

Retinal ganglion cells were recorded from dorsal retina using a large scale, dense

hexagonal multi-electrode array covering $\sim 0.34mm^2$ of the retina (MEA, (Field et al., 2010) 519

electrodes with 30 µm spacing). The pigmented epithelium remained attached to the retina for

these recordings. The retina was perfused with Ames' solution ($30\text{-}31^{\circ}C$) bubbled with 95/5%

O_2/CO_2. Spikes were identified and assigned to specific RGCs on the MEA as previously

described (Yu et al., 2017). Dim flashes were delivered at 3 s intervals using a 490 nm LED.

Light intensity was controlled using pulse duration, 2-8 ms, and neutral density filters. Dim flash

responses were measured by counting spikes on each trial within a 100 ms window that was

centered on the peak of the peristimulus time histogram.

Experimental design and statistical analyses

Because our initial studies did not show gender-specific differences, the genders were

pooled. RGC response thresholds were measured from three *Cngb1ΔCaM* retinas (102-232

cells), five *Cngb1$^{neo/neo}$* retinas (100-336 cells) and three *Cngb1$^{neo/neo}$* rescue retinas (53-186

cells). Cumulitive threshold histograms were calculated in each tissue and averaged across all

retinas within a condition. A two tailed Kolmogorov-Smirnov goodness-of-fit hypothesis test was

used to assess the statistical difference between average cumulative histograms. The fraction

of cells for which no response surpassed threshold was also measured in each recorded retina.

A two sample t-test was used to evaluate significance between conditions.

Results

Generation of a novel animal model of genetically reversible rod degeneration

One challenge to identifying how plasticity among inner retinal neurons impacts functional recovery is the lack of an experimental system that is non-invasive and allows for stringent regulation of the timing and uniformity of rescue. For example, viral-mediated (gene therapy) approaches for treating rod dysfunction and death (1) take weeks for expression to occur; (2) they do not infect all targeted cells; (3) they may not drive proper protein expression levels; (4) and the subretinal injections used for viral delivery can damage the retina. A systematic investigation into the consequences of rod degeneration and subsequent rescue of the retinal circuitry requires an experimental system wherein both events occur uniformly in the retina. Towards this goal, a neoloxP cassette was inserted into intron 19 of the *Cngb1* gene by homologous recombination in mouse embryonic stem cells (Fig. 1A). Mice harboring this insertion were subsequently derived (*Cngb1$^{neo/neo}$*). The presence of the cassette disrupted a splice site and prevented CNGB1 expression (Fig. 1B). Expression of CNGA1 was also substantially attenuated (Fig. 1B), a phenomenon attributed to mis-trafficking (Hüttl et al., 2005) and structural stability conferred by association of both subunits. The expression levels of other major phototransduction proteins were minimally perturbed in retinas of 1-month old (1 MO) mice (Fig. 1B, NCKX1, GC1, PDE6A, and GNAT1). Consistent with previous reports on conventional *Cngb1* knockout mice (Hüttl et al., 2005; Zhang et al., 2009), the lack of CNG channel expression led to a progressive thinning of the outer nuclear layer over the course of 6 months (Fig. 1C). At two-weeks, the outer nuclear layer (ONL) containing primarily rod photoreceptor cell nuclei reached its maximum thickness. This thickness was reduced by ~20% in 1 MO mice and to ~50% in 2 MO mice. By 6 MO, the ONL was absent. Thus, these mice exhibit slow rod degeneration relative to other commonly used models of rod degenerative diseases, such as *rd1* (Farber and Lolley, 1974) and *rd10* mice (Chang et al., 2007).

As expected, the absence of the CNGB1 recapitulated the stereotypic sequence of events associated with rod degeneration (Marc and Jones, 2003; Puthussery and Taylor, 2010; Soto and Kerschensteiner, 2015). For example, rhodopsin mislocalization and activation of Müller glia were observed in 4-week old *Cngb1$^{neo/neo}$* mice (Fig. 1, compare F, G with control retina, D and E).

We also observed in *Cngb1$^{neo/neo}$* mice that synaptic contacts between rods and rod bipolar cells were structurally abnormal. Immunohistochemistry using a marker for the presynaptic ribbon protein CtBP2 (ribeye) and the post-synaptic glutamate receptor, mGluR6,

revealed clear differences between *Cngb1^{neo/neo}* and control retinas. In control retinas, these
structures were closely apposed, and both were contained within a well-defined outer plexiform
layer (Fig. 1H, OPL). However, in *Cngb1^{neo/neo}* retinas these structures were more dispersed,
with synaptic ribbons retracted from rod spherules and some were situated deep into the
photoreceptor nuclear layer (ONL, Fig. 1I, arrows). The ultrastructure of rod synapses was
further evaluated by transmission electron microscopy (TEM, Fig. 1J and K). A normal rod
spherule (blue) contains a single large mitochondria (yellow) and encompasses a synaptic triad
consisting of a single ribbon (red) along with horizontal (orange) and rod bipolar cell (green)
processes. The majority of imaged rod spherules from control retinas exhibited this structure.
Dyads were also frequently observed when one or another component was out of the plane of
the TEM section. Of the 27 fields taken from 4 control C57 retinas, 122 synapses were counted
(4.5 synapses per field) wherein 64% were triads and 36% were dyads. However, in 1 MO
Cngb1^{neo/neo}   retinas, the frequency of observing synapses and triadic structures were both
reduced (Fig. 1K). Of 31 TEM fields taken from 4 *Cngb1^{neo/neo}* retinas, 64 synapses were
counted (2.1 synapses per field) wherein only 22% were triads and 78% of were dyads. Note, at
1 MO, only 20% of rods had died, yet there was >50% reduction in contacts between rods and
rod bipolar cells. Furthermore the contacts that persisted were largely abnormal in structure.
These results demonstrate that *Cngb1^{neo/neo}* mice exhibit stereotypic slow rod degeneration and
that synaptic structures between rods and rod bipolar cells were disrupted.
292
Lack of CNGB1 expression attenuated rod photoresponses and eliminated rod bipolar
cell light responses.
Previous work has indicated that lack of CNGB1 expression compromises rod vision
(Biel and Michalakis, 2007). To verify compromised rod function in *Cngb1^{neo/neo}* mice, we
performed *ex vivo* whole-retina electroretinograms (ERG) under scotopic conditions. The ERG
reflects the averaged activity across all retinal neurons (Granit, 1933). ERGs from C57 retinas
exhibited a well characterized biphasic response (Fig 2A; see also (Saszik et al., 2002)) with the
initial negative-voltage deflection (a-wave) indicative of the rod hyperpolarization to the flash
stimulus, and the subsequent positive-voltage rebound indicative of predominantly the rod
bipolar cell depolarization. Recordings were also performed on control *Cngb1ΔCaM* mice in
which the calmodulin binding site was removed (Chen et al., 2010). No differences were
observed between *Cngb1ΔCaM* and C57 retinas (data not shown).
ERGs from 1 MO *Cngb1^{neo/neo}* retinas exhibit a diminished scotopic a-wave with reduced
sensitivity (Fig. 2B), indicating minimal rod signaling without CNGB1 expression. The light

response is not fully eliminated in these rods; this is likely due to residual activation of

homomeric channels composed of CNGA1 (see Discussion). Furthermore, ERGs from

Cngb1^{neo/neo} retinas did not exhibit a b-wave under scotopic conditions, indicating a lack of

measurable rod-to-rod bipolar signal transmission at this gross level. This observation

complements the abnormal synaptic structures observed between rods and rod bipolar cells via

light and electron microscopy. Together, these results indicate that synaptic transmission

between rods and rod bipolar cells is severely dysfunctional in *Cngb1^{neo/neo}* mice. Thus, we

sought to determine the extent to which normal synaptic structures and transmission between

rods and rod bipolar cells could be recovered by the rescue of CNGB1 expression in mature

retinas.

Cre-mediated excision of the NeoLoxP cassette leads to normal CNGB1 expression.

To activate CNGB1 expression in the the *Cngb1^{neo/neo}* retina, we utilized the CAGGCre-

ER^{TM} transgene (Hayashi and McMahon, 2002) to enable tamoxifen-dependent, cre-mediated

excision of the NeoloxP cassette. We previously demonstrated that mice derived from germline

excision of this cassette exhibit normal retinal morphology with a uniform and normal expression

level of CNGB1 (Chen et al., 2010). The homologous recombination strategy that introduced

the NeoloxP cassette also removed a stretch of 14 amino acids that encompassed the

calmodulin binding domain on CNGB1 (Grunwald et al., 1998). Importantly, rods that expressed

CNGB1ΔCaM exhibited normal light responses as mentioned above (see also (Chen et al.,

2010)). We hypothesize that utilizing the CAGGCre-ER^{TM} transgene would provide temporal

control over expression of the functional CNG channel. To determine the efficacy of cre-

mediated excision of the neoloxP cassette, four-week-old cre-positive and cre-negative

Cngb1^{neo/neo} littermate mice were divided into two groups. One group was given tamoxifen for

four-consecutive days by oral gavage, and the other group did not receive drug treatment.

A PCR strategy was designed to detect the extent of neoloxP excision in genomic DNA

extracted from isolated retinas: the primer pair (2+3) detects the presence of the neoloxP insert,

whereas primer pair (1+3) gives rise to a diagnostic band when the large neoloxP insert is

excised (Fig. 3A). After four-consecutive days of tamoxifen treatment, both sets of primers

produced positive bands. This result indicates a mixed population of cells at this stage, some of

which have undergone excision while others have not. However, when tamoxifen treatment was

given for seven-consecutive days a positive signal was detected only by primers (1+3). This

result indicates that following a 7-day tamoxifen treatment, most, if not all cells have undergone

neoloxP excision (Fig. 3A, bottom panels). Thus a 7-day treatment was used for further
structural and functional studies.

To assess the level of protein expression at 6- or 8-weeks (corresponding to 1 or 3
weeks after drug treatment), Western blots were prepared from whole retinal homogenates from
both cohorts (Fig. 3B). Expression of CNGB1 protein was observed only in tamoxifen-treated,
cre-positive mice. No expression was observed in cre-positive mice without drug treatment,
indicating a lack of basal recombinase activity. We next examined how excision of the neoloxP
insert affected the expression of CNGB1 and other major phototransduction proteins. We found
that following neoloxP excision, there was a striking increase in CNGB1 expression (Fig. 3C).
There was also an increase in the detected levels of other phototransduction proteins GC1,
PDE6A, GNAT1, and GCAP2 (Fig 3C). To determine if this is due to rod rescue, cre-negative
and cre-positive *Cngb1*$^{neo/neo}$ littermate mice were administered tamoxifen for 7 consecutive
days beginning at 4-weeks, and retinal sections were prepared from 3 MO mice (Fig. 3D and
3E). The ONL thickness was greater in cre-positive mice when compare to that from the cre-
negative sibling mice, and the rod outer segment structure was organized and of normal length.
Expression of CNGB1 exhibited a long term rescuing effect on rod survival (Fig. 3F and 3G,
tamoxifen was administered for 4 consecutive days starting at P28), consistent with a previous
report on AAV-mediated *Cngb1* gene replacement therapy (Koch et al., 2012). In sum, these
data show that the *Cngb1*$^{neo/neo}$ mice allowed us to regulate the expression of CNGB1 from the
endogenous locus in a temporally-controlled manner. Further, this excision is nearly complete
with a 7-day tamoxifen treatment and that upon expression of CNGB1, the rods exhibit normal
morphology and are stably rescued from cell death.

We measured the responsiveness of rod photoreceptors following drug treatment in
patch-clamp recordings from individual rods in retinal slices. In voltage-clamp (V_m = -40 mV),
rods from 1 MO *Cngb1*$^{neo/neo}$ mice displayed diminished response amplitudes (~6-fold) and a
~10-fold reduction in light sensitivity (Fig. 4A), a result consistent with the diminished a-wave in
ex vivo ERG recordings (Fig. 2B). In current-clamp (i = 0), *Cngb1*$^{neo/neo}$ rods exhibit a resting
membrane potential that was ~10 mV hyperpolarized compared to WT rods (-47 ± 1.3 mV (5)
vs. -37 ± 2.3 mV (6), mean ± SEM). These results are consistent with reduced CNG channel
expression (see Discussion) and indicate reduced glutamate release in darkness. However,
rods from tamoxifen treated *Cngb1*$^{neo/neo}$ mice displayed responses with characteristics very
similar to C57 mice (Fig. 4B), consistent with near-normal function and rescue of the
photoreceptor layer (Fig. 3).

Expression of CNGB1 induces normal synaptic structures between rods and rod bipolar
cells.

Given that tamoxifen administration in *Cngb1^{neo/neo}* mice rescued rods from death (Fig.
3E) and rescued normal rod light responses (Fig. 4), we next examined the synaptic contacts
between rods and rod bipolar cells to determine how rod rescue impacts these structures.
Tamoxifen treatment was initiated at 4-weeks for 7 consecutive days, and retinal structure was
examined at 3M. Comparisons were made between 1 MO C57 and *Cngb1^{neo/neo}* mice and 3 MO
tamoxifen-treated mice to examine the effect of rod rescue that was initiated at 1M. Synaptic
structures were labeled in retinal flat mounts stained for the presynaptic ribbon synapse protein
(CtBP2, blue) and postsynaptic mGluR6 (orange, Fig. 5A, 5D and 5G). To distinguish between
rod and cone synapses, cone pedicles were further labeled with the cone arrestin antibody
(ARR3, green). Rod bipolar cell morphology, visualized by PKCα staining, and the mGluR6
puncta that decorate their dendritic tips are shown in retinal cross sections (Fig. 5B, 5E and 5H).

Control retinas (C57), exhibited a close juxtaposition between the rod's single ribbon and
the mGluR6 puncta on the dendritic tips of rod bipolar cells (Fig. 5A and 5B; Fig. 1G). However,
in *Cngb1^{neo/neo}* retinas from 1 MO mice, both the number of synaptic ribbons and mGluR6
puncta were reduced (Fig. 5D). Rod bipolar cell dendrites were also unevenly distributed in
Cngb1^{neo/neo} retinas, and the size of the mGluR6 puncta appeared smaller and less uniform in
shape (Fig. 5E and Fig. 1H). In contrast, retinas from the tamoxifen-treated, cre-positive
littermates exhibited robust staining of synaptic ribbons along with their associated mGluR6
puncta (Fig. 5G). Furthermore, rod bipolar cell dendrites were evenly extended and the
mGluR6 puncta were larger in size and appeared more uniform in shape (Fig. 5H). To quantify
these changes, the number and size of mGluR6 puncta were measured (Fig. 5C, 5F and 5I).
For the C57 retina, mGluR6 puncta size of 20- to 30-unit area were the most numerous,
whereas in *Cngb1^{neo/neo}* retinas smaller size puncta were more frequent (compare Fig. 5C and
5F). The distribution shifted back to larger puncta sizes in the tamoxifen-treated mice (Fig. 5I).
These results indicate that inducing expression of CNGB1 in mature retina causes a recovery of
synaptic structures between rods and rod bipolar cells.

Rescue of CNGB1 expression in mature retina recovers rod bipolar light responses.
The results above indicate a structural recovery of synapses following expression of
CNGB1. In addition to this structural recovery, *ex vivo* whole-retina ERGs revealed a recovery
of the rod bipolar cell-driven b-wave with amplitudes similar to control retinas (Fig. 6A). Thus,
structural and functional measures broadly indicate recovery of synaptic function between rods

and rod bipolar cells. To examine further synaptic function before and following rod rescue, we
performed patch clamp recordings from rod bipolar cells in retinal slices. In untreated
Cngb1^{neo/neo} mice there was a complete absence of functional transmission of between rods and
rod bipolar cells (Fig. 6C); 1 MO *Cngb1^{neo/neo}* rod bipolar cells never yielded light-evoked
responses (n=15 from 5 retinas), a result consistent with a lack of b-wave in *ex vivo* ERG
recordings which represent mass-action of largely rod bipolar cells (Fig. 2B). However, in
Cngb1^{neo/neo} mice administered tamoxifen for 7 days at 4 weeks of age and recorded at 3 MO,
rod bipolar cells exhibited robust light-evoked responses similar to control animals (Fig. 6D).
The extent of functional recovery in rod bipolar cells was characterized in plots of the response
amplitude versus the flash strength. These intensity-response relationships were fit with a Hill
curve and compared quantitatively to control responses. The half-maximal flash strength ($I_{1/2}$)
increased by ~2-fold in rescued animals (Fig. 6E), consistent with some rod loss (see
Discussion). However, other features of the rod bipolar light response that are critical for
function near visual threshold had recovered to near control values. For example, the Hill
exponent for the fit of the response-intensity relationship matched that in control, indicating a
similar nonlinear relationship between the flash strength and the response amplitude. The
extent of nonlinearity reflects the rate of glutamate release from rod synapses (Sampath and
Rieke, 2004). In addition, the time course of rod bipolar cell responses was similar in rescued
animals (Fig. 6A, 6C, 6E – dashed line), further indicating the anatomical and physiological
recovery of synaptic transmission in tamoxifen-treated animals. These results indicate that
rescuing rod function in the mature mouse retina produces a cascade of structural and
functional recovery in synaptic transmission between rods and rod bipolar cells, and thus the
primary rod pathway (Dacheux and Raviola, 1986).
431
Rescuing rods recovers absolute sensitivity of retinal output
Retinal ganglion cells (RGCs) provide the sole output from the retina and can integrate
input from thousands of rods, making them the most light-sensitive cells in the retina
(Chichilnisky and Rieke, 2005; Field and Sampath, 2017). RGC sensitivity relies on functioning
photoreceptors and highly tuned synaptic connections via the primary rod pathway (Field and
Rieke, 2002; Sampath and Rieke, 2004). To understand how rescuing rod function in the 1 MO
Cngb1^{neo/neo} retina impacts RGC sensitivity, we used a large-scale multi-electrode array (MEA)
to record spikes from hundreds of RGCs. We tested the sensitivity of the RGCs by stimulating
the retina with brief, dim flashes (0.001-10 Rh*/rod) and compared RGC responses in 3 MO
control *Cngb1ΔCaM*, *Cngb1^{neo/neo}*, and 4-5 MO *Cngb1* tamoxifen rescued mice.   Flashes

producing less than 1 isomerization per rod faithfully produced spike rate modulations in many
RGCs from control mice (Fig 7A shows an example cell). The same flash intensities did not
reliably modulate the spike output of most RGCs in retinas from 3 MO old $Cngb1^{neo/neo}$ mice (Fig
7B shows an example cell). Indeed, most RGCs from $Cngb1^{neo/neo}$ mice did not show reliable
responses until flash intensities exceeded 1 Rh*/rod (Fig $7B_3$). However, similar to the control
retinas, RGC responses were often evident at low flash intensities in 4-5 MO $Cngb1$ rescued
mice (Fig 7C shows an example cell). These example cells suggest that rod and circuit
functionality are broadly and stably rescued in some RGCs for $Cngb1$ rescued mice.

To measure the extent that sensitivity across the RGC population recovered in $Cngb1$
rescued mice, we quantified the response-threshold for all RGCs (N=1954) in MEA recordings
from 11 mice (3 control $Cngb1\Delta CaM$ mice, 3 $Cngb1^{neo/neo}$ mice, 5 tamoxifen-treated $Cngb1$
mice). RGC response thresholds were quantified as the lowest flash intensity needed to drive
the average spike rate change two standard deviations above baseline (eg. Fig $7A_3$, $7B_3$, $7C_3$;
see Methods). Average RGC response threshold distributions were similar between control and
$Cngb1$ rescued mice, but were significantly higher in untreated $Cngb1^{neo/neo}$ mice (Fig 7D; KS
test, $p<0.05$). Additionally, the fraction of RGCs for which no-threshold response could be
measured was similar between control and $Cngb1$ rescued mice but significantly higher in
untreated $Cngb1^{neo/neo}$ mice (Fig 7E; t-test, $p<0.05$). These results indicate a broad and lasting
recovery of rod and circuit functions in $Cngb1$ rescued mice.

Discussion

In contrast to other neurons, rods and cones are depolarized in darkness and tonically
release glutamate through ribbon synapses (Molday and Moritz, 2015), leading to saturation of
rod-to-rod bipolar cell synapses (Sampath and Rieke, 2004). Light exposure causes graded
hyperpolarization of the photoreceptor cell and suppression of glutamate release. Reductions in
glutamate release from photoreceptors that occur during the early process of retinal
degeneration lead to homeostatic changes in the downstream neurons and degrade the retinal
circuit. This is seen in the dissolution of synaptic structures, dendritic sprouting, formation of
ectopic contacts, and gliosis (Marc et al., 2003; Puthussery and Taylor, 2010). Although
strategies to rescue and restore function in defective photoreceptors have shown success for
regaining some visual function, gene and stem cell therapies for visual restoration are often
implemented in the adult; how well these rescued neurons reinstate their detailed circuitries in
the remodeled retina is not known. Here we examined functional restoration at the level of inner

retinal cells and defined rod-driven circuits in the young adult mouse retina. We show that

repairing a primary genetic defect in rods not only restored rod function, but also recovered

normal synaptic connectivity with remodeled second order rod bipolar cells.

Rod-to-rod bipolar cell synaptic contacts are reduced in *Cngb1neo/neo* retina and do not

support synaptic transmission

The relatively slow photoreceptor degeneration we observe in the *Cngb1neo/neo* mouse

model, and observed in human patients (Bareil et al., 2001; Biel and Michalakis, 2007), may be

due to the fact that it is not a functional null. A small but measurable light response persisted in

Cngb1neo/neo rods from 1 MO mice (Fig. 4B). The residual light response is probably due to the

presence of homomeric channels composed of CNGA1 subunits, which are capable of

mediating a diminished and desensitized cGMP-dependent current (Kaupp et al., 1989). The

small current would reduce Ca^{2+} influx to the outer segment, causing increased levels of cGMP

through stimulation of guanylyl cyclases by Ca^{2+}-free guanylyl cyclase activating proteins 1 and

2 (Mendez et al., 2001; Dizhoor et al., 2010). Elevated cGMP has been shown to be a driver of

rod degeneration through activation of protein kinase G (Ma et al., 2015; Wang et al., 2017a).

Despite the diminished rod light responses in the *Cngb1neo/neo* mice, the utter lack of

evidence for light-evoked transmission between rods and rod bipolar cells was unexpected,

especially given our ultrastructural evidence showing that the number of triads in the

Cngb1neo/neo rod spherules was reduced, but not fully eliminated (Fig. 1). Therefore, it is

surprising that ERGs from *Cngb1neo/neo* mice did not exhibit a scotopic b-wave, which is largely

contributed by rod bipolar cells (Fig. 2B; (Saszik et al., 2002). This result was further

corroborated by patch-clamp measurements from rod bipolar cells that revealed no light

response, even for bright flashes delivering ~2000 Rh*/rod (Fig. 6C). We speculate that this

defect in synaptic transmission is due to diminished glutamate release at the ribbon synapse

given that the lack of CNG channels should act as a source of "equivalent light", similar to light-

induced closure of CNG channels (see also Sampath and Rieke, 2004; Dunn et al., 2006).

Supporting this idea, the resting membrane potential of *Cngb1neo/neo* rods are ~10 mV

hyperpolarized due to their smaller dark current (Fig. 4A). At the rod's normal resting potential in

darkness (~40 mV), calcium enters through the voltage gated channel (Ca_v1.4) and supports

tonic glutamate release at the ribbon synapse (Waldner et al., 2018). Thus the hyperpolarizing

shift in resting potential of *Cngb1neo/neo* rods predicts attenuated glutamate release from the rod

spherule.

Interestingly, suppression of glutamate release at the rod synapse is strongly correlated
with synaptic remodeling. Examples include blockade of glutamate release by tetanus toxin
(Cao et al., 2015), in knockout mice that lack the presynaptic $Ca_V1.4$ Ca^{2+} channel (Mansergh et
al., 2005), and in human patients diagnosed with congenital stationary night blindness (CSNB2)
that harbor null mutations in the gene encoding $Ca_V1.4$ (Bech-Hansen et al., 1998; Boycott et
al., 2000). Calcium entry through $Ca_V1.4$ channel is required for neurotransmitter release at the
ribbon synapse of both rods and cones. The absence of CaBP4 (Haeseleer et al., 2004) or $\alpha2\delta4$
(Wang et al., 2017c) that bind and regulate the activity of $Ca_V1.4$, also manifest in retinal
remodeling in knockout mice. These plastic changes occurred with minimal photoreceptor cell
loss, suggesting that synaptic remodeling is likely driven by suppression of neural transmission,
or deafferentation, rather than photoreceptor cell death *per se*. Modest synaptic changes were
also observed in RIBEYE knockout retinas. RIBEYE is an essential component of the synaptic
ribbon, and its absence abolished all presynaptic ribbons in the retina and severely impaired
fast and sustained neurotransmitter release (Maxeiner et al., 2016). Spontaneous miniature
release continues to occur without the synaptic ribbon, which may explain the milder retinal
remodeling phenotype observed in the RIBEYE knockout retina (Maxeiner et al., 2016).

Previous studies on two independent lines of *Cngb1^-/-* mice have similarly reported a
substantial reduction of light responses from *Cngb1^-/-* rods, but in both mouse lines the presence
of a rod bipolar cell driven b-wave was observed (Hüttl et al., 2005; Zhang et al., 2009). The
reason behind the discrepancy between those and our results is not clear, but may be due to
differences in: 1) the degree of rod hyperpolarization caused by the number of functioning
homomeric CNGA1 channels and hence the amount of glutamate released by rods in the
different mouse lines, 2) the *in vivo* vs. *ex vivo* ERG measurements, or 3) mouse genetic
backgrounds.

Adult rod bipolar cells demonstrate plastic changes to establish functional contacts with
rescued rods

The developmental time window for the formation of the rod to rod bipolar cell synapse
in mice appears to be from eye opening to postnatal day 30 (P30), during which synaptic
proteins are expressed, pre- and post-synaptic molecular complexes form, and the rod bipolar
cells develop the appropriate number of dendritic tips that make synaptic contacts with rods
(Anastassov et al., 2017). Some of the molecules that guide neurite growth during development
are absent at maturity (D'Orazi et al., 2014), and if functional connectivity of the neural retina
can only occur during a critical period in development, then one would expect that the adult

retina may lack the ability to make such connections when rod activity is switched on after this
time window. Such developmental processes would have been disrupted in *Cngb1^{neo/neo}* retinas,
wherein pronounced retinal remodeling is evident by P30 (Fig. 1). We show that tamoxifen-
induced CNGB1 expression between P28-P34 led to establishment of the rod's circulating
current in darkness and normal light responses (Fig. 4). Concomittantly, structural changes
were observed at the synapse: rod bipolar cells elaborated fine dendritic tips, mGluR6 receptors
clustered on these tips which came in close contact with presynaptic ribbons (Fig. 5G-5I). These
newly formed synaptic structures supported normal neural transmission, as shown by ERG
recordings and patch recordings from rod bipolar cells (Fig. 6), and light sensitivity in increased
in RGCs (Fig. 7). We hypothesize that these changes may be initiated by glutamate release at
the rod's synapse, similar to that which occurs at the cortex, where focal uncaging of glutamate
in mouse cortical layer 2/3 pyramidal neurons triggered spinogenesis from the dendrite shaft in
a location-specific manner (Kwon and Sabatini, 2011).

Plasticity at the photoreceptor/bipolar cell synapse has also been observed in a model of
photocoagulation of rabbit retina, where the laser ablation acutely removes a patch of
photoreceptors while leaving the inner retina intact (Beier et al., 2017). After some days, nearby
photoreceptors slowly migrate toward and fill in the lesioned area (Sher et al., 2013). As they do
so, they form functional contacts with the deafferented bipolar cells (Sher et al., 2013; Beier et
al., 2017). Another example of plasticity at the photoreceptor/bipolar cell synapse is the AAV-
mediated gene therapy to replace retinoschisin (RS1) in adult mice (Ou et al., 2015). Retinal
development of the RS1 knockout mice appears to proceed normally. However, the absence of
RS1, a cell adhesion protein, eventually causes splitting of the retina and a failure of synaptic
maintenance that manifests in reduction of the ERG b-wave amplitude (Sikkink et al., 2007).
This defect was reversed upon RS1 gene replacement (Ou et al., 2015). Here we demonstrate
for the first time that activation of rod input in young adults reversed synaptic changes that
occurred during development and established functional contacts with their downstream
neurons in the retinal circuitry of the adult retina. These results support the therapeutic potential
of repairing or replacing defective rods in the degenerating retina. However, a critical time
window for rescue likely exists: recent clinical trials for Leber Congenital Amaurosis to replace
RPE65 in human patients for treating a type of Leber's congenital amaurosis (LCA) caused by
RPE65 mutations show limited success in visual improvement, and the retina continued to
degenerate in some patients (Cideciyan et al., 2013). Future experimentation will address
whether a critical time window of rescue exists for these approaches.

References

Anastassov IA, Wang W, Dunn FA (2017) Synaptogenesis and synaptic protein localization in the postnatal development of rod bipolar cell dendrites in mouse retina. J Comp Neurol.

Bareil C, Hamel CP, Delague V, Arnaud B, Demaille J, Claustres M (2001) Segregation of a mutation in CNGB1 encoding the beta-subunit of the rod cGMP-gated channel in a family with autosomal recessive retinitis pigmentosa. Hum Genet 108:328-334.

Bech-Hansen NT, Naylor MJ, Maybaum TA, Pearce WG, Koop B, Fishman GA, Mets M, Musarella MA, Boycott KM (1998) Loss-of-function mutations in a calcium-channel alpha1-subunit gene in Xp11.23 cause incomplete X-linked congenital stationary night blindness. Nat Genet 19:264-267.

Beier C, Hovhannisyan A, Weiser S, Kung J, Lee S, Lee DY, Huie P, Dalal R, Palanker D, Sher A (2017) Deafferented Adult Rod Bipolar Cells Create New Synapses with Photoreceptors to Restore Vision. J Neurosci 37:4635-4644.

Bermingham-McDonogh O, Reh TA (2011) Regulated reprogramming in the regeneration of sensory receptor cells. Neuron 71:389-405.

Biel M, Michalakis S (2007) Function and dysfunction of CNG channels: insights from channelopathies and mouse models. Molecular neurobiology 35:266-277.

Boycott KM, Pearce WG, Bech-Hansen NT (2000) Clinical variability among patients with incomplete X-linked congenital stationary night blindness and a founder mutation in CACNA1F. Can J Ophthalmol 35:204-213.

Cao Y, Sarria I, Fehlhaber KE, Kamasawa N, Orlandi C, James KN, Hazen JL, Gardner MR, Farzan M, Lee A, Baker S, Baldwin K, Sampath AP, Martemyanov KA (2015) Mechanism for Selective Synaptic Wiring of Rod Photoreceptors into the Retinal Circuitry and Its Role in Vision. Neuron 87:1248-1260.

Chang B, Hawes NL, Pardue MT, German AM, Hurd RE, Davisson MT, Nusinowitz S, Rengarajan K, Boyd AP, Sidney SS, Phillips MJ, Stewart RE, Chaudhury R, Nickerson JM, Heckenlively JR, Boatright JH (2007) Two mouse retinal degenerations caused by missense mutations in the beta-subunit of rod cGMP phosphodiesterase gene. Vision Res 47:624-633.

Chen J, Woodruff ML, Wang T, Concepcion FA, Tranchina D, Fain GL (2010) Channel modulation and the mechanism of light adaptation in mouse rods. The Journal of neuroscience : the official journal of the Society for Neuroscience 30:16232-16240.

Chichilnisky EJ, Rieke F (2005) Detection sensitivity and temporal resolution of visual signals near absolute threshold in the salamander retina. The Journal of neuroscience : the official journal of the Society for Neuroscience 25:318-330.

Chua J, Fletcher EL, Kalloniatis M (2009) Functional remodeling of glutamate receptors by inner retinal neurons occurs from an early stage of retinal degeneration. J Comp Neurol 514:473-491.

Cideciyan AV, Jacobson SG, Beltran WA, Sumaroka A, Swider M, Iwabe S, Roman AJ, Olivares MB, Schwartz SB, Komaromy AM, Hauswirth WW, Aguirre GD (2013) Human retinal gene therapy for Leber congenital amaurosis shows advancing retinal degeneration despite enduring visual improvement. Proceedings of the National Academy of Sciences of the United States of America 110:E517-525.

Concepcion F, Chen J (2010) Q344ter mutation causes mislocalization of rhodopsin molecules that are catalytically active: a mouse model of Q344ter-induced retinal degeneration. PLoS One 5:e10904.

Cook NJ, Molday LL, Reid D, Kaupp UB, Molday RS (1989) The cGMP-gated channel of bovine rod photoreceptors is localized exclusively in the plasma membrane. J Biol Chem 264:6996-6999.

D'Orazi FD, Suzuki SC, Wong RO (2014) Neuronal remodeling in retinal circuit assembly,
disassembly, and reassembly. Trends Neurosci 37:594-603.
Dacheux RF, Raviola E (1986) The rod pathway in the rabbit retina: a depolarizing bipolar and
amacrine cell. The Journal of neuroscience : the official journal of the Society for
Neuroscience 6:331-345.
Dizhoor AM, Olshevskaya EV, Peshenko IV (2010) Mg2+/Ca2+ cation binding cycle of guanylyl
cyclase activating proteins (GCAPs): role in regulation of photoreceptor guanylyl
cyclase. Mol Cell Biochem 334:117-124.
Farber DB, Lolley RN (1974) Cyclic guanosine monophosphate: elevation in degenerating
photoreceptor cells of the C3H mouse retina. Science 186:449-451.
Field GD, Rieke F (2002) Nonlinear signal transfer from mouse rods to bipolar cells and
implications for visual sensitivity. Neuron 34:773-785.
Field GD, Sampath AP (2017) Behavioural and physiological limits to vision in mammals. Philos
Trans R Soc Lond B Biol Sci 372.
Field GD, Gauthier JL, Sher A, Greschner M, Machado TA, Jepson LH, Shlens J, Gunning DE,
Mathieson K, Dabrowski W, Paninski L, Litke AM, Chichilnisky EJ (2010) Functional
connectivity in the retina at the resolution of photoreceptors. Nature 467:673-677.
Garg A, Yang J, Lee W, Tsang SH (2017) Stem Cell Therapies in Retinal Disorders. Cells 6.
Granit R (1933) The components of the retinal action potential in mammals and their relation to
the discharge in the optic nerve. J Physiol 77:207-239.
Grunwald ME, Yu WP, Yu HH, Yau KW (1998) Identification of a domain on the beta-subunit of
the rod cGMP-gated cation channel that mediates inhibition by calcium-calmodulin. J
Biol Chem 273:9148-9157.
Haeseleer F, Imanishi Y, Maeda T, Possin DE, Maeda A, Lee A, Rieke F, Palczewski K (2004)
Essential role of Ca2+-binding protein 4, a Cav1.4 channel regulator, in photoreceptor
synaptic function. Nat Neurosci 7:1079-1087.
Hayashi S, McMahon AP (2002) Efficient recombination in diverse tissues by a tamoxifen-
inducible form of Cre: a tool for temporally regulated gene activation/inactivation in the
mouse. Dev Biol 244:305-318.
Hoyo NL, Lopez-Begines S, Rosa JL, Chen J, Mendez A (2014) Functional EF-hands in
neuronal calcium sensor GCAP2 determine its phosphorylation state and subcellular
distribution in vivo, and are essential for photoreceptor cell integrity. PLoS Genet
10:e1004480.
Hüttl S, Michalakis S, Seeliger M, Luo DG, Acar N, Geiger H, Hudl K, Mader R, Haverkamp S,
Moser M, Pfeifer A, Gerstner A, Yau KW, Biel M (2005) Impaired channel targeting and
retinal degeneration in mice lacking the cyclic nucleotide-gated channel subunit CNGB1.
J Neurosci 25:130-138.
Kaupp UB, Niidome T, Tanabe T, Terada S, Bonigk W, Stuhmer W, Cook NJ, Kangawa K,
Matsuo H, Hirose T, et al. (1989) Primary structure and functional expression from
complementary DNA of the rod photoreceptor cyclic GMP-gated channel. Nature
342:762-766.
Koch S, Sothilingam V, Garcia Garrido M, Tanimoto N, Becirovic E, Koch F, Seide C, Beck SC,
Seeliger MW, Biel M, Muhlfriedel R, Michalakis S (2012) Gene therapy restores vision
and delays degeneration in the CNGB1(-/-) mouse model of retinitis pigmentosa. Hum
Mol Genet 21:4486-4496.
Kwon HB, Sabatini BL (2011) Glutamate induces de novo growth of functional spines in
developing cortex. Nature 474:100-104.
Ma H, Butler MR, Thapa A, Belcher J, Yang F, Baehr W, Biel M, Michalakis S, Ding XQ (2015)
cGMP/Protein Kinase G Signaling Suppresses Inositol 1,4,5-Trisphosphate Receptor
Phosphorylation and Promotes Endoplasmic Reticulum Stress in Photoreceptors of
Cyclic Nucleotide-gated Channel-deficient Mice. J Biol Chem 290:20880-20892.

Mansergh F, Orton NC, Vessey JP, Lalonde MR, Stell WK, Tremblay F, Barnes S, Rancourt
DE, Bech-Hansen NT (2005) Mutation of the calcium channel gene Cacna1f disrupts
calcium signaling, synaptic transmission and cellular organization in mouse retina. Hum
Mol Genet 14:3035-3046.
Marc RE, Jones BW (2003) Retinal remodeling in inherited photoreceptor degenerations. Mol
Neurobiol 28:139-147.
Marc RE, Jones BW, Watt CB, Strettoi E (2003) Neural remodeling in retinal degeneration. Prog
Retin Eye Res 22:607-655.
Maxeiner S, Luo F, Tan A, Schmitz F, Sudhof TC (2016) How to make a synaptic ribbon:
RIBEYE deletion abolishes ribbons in retinal synapses and disrupts neurotransmitter
release. Embo J 35:1098-1114.
Mendez A, Burns ME, Sokal I, Dizhoor AM, Baehr W, Palczewski K, Baylor DA, Chen J (2001)
Role of guanylate cyclase-activating proteins (GCAPs) in setting the flash sensitivity of
rod photoreceptors. Proc Natl Acad Sci U S A 98:9948-9953.
Molday RS, Moritz OL (2015) Photoreceptors at a glance. J Cell Sci 128:4039-4045.
Ou J, Vijayasarathy C, Ziccardi L, Chen S, Zeng Y, Marangoni D, Pope JG, Bush RA, Wu Z, Li
W, Sieving PA (2015) Synaptic pathology and therapeutic repair in adult retinoschisis
mouse by AAV-RS1 transfer. J Clin Invest 125:2891-2903.
Pahlberg J, Frederiksen R, Pollock GE, Miyagishima KJ, Sampath AP, Cornwall MC (2017)
Voltage-sensitive conductances increase the sensitivity of rod photoresponses following
pigment bleaching. J Physiol 595:3459-3469.
Poetsch A, Molday LL, Molday RS (2001) The cGMP-gated channel and related glutamic acid-
rich proteins interact with peripherin-2 at the rim region of rod photoreceptor disc
membranes. J Biol Chem 276:48009-48016.
Puthussery T, Taylor WR (2010) Functional changes in inner retinal neurons in animal models
of photoreceptor degeneration. Adv Exp Med Biol 664:525-532.
Quartilho A, Simkiss P, Zekite A, Xing W, Wormald R, Bunce C (2016) Leading causes of
certifiable visual loss in England and Wales during the year ending 31 March 2013. Eye
(Lond) 30:602-607.
Sampath AP, Rieke F (2004) Selective transmission of single photon responses by saturation at
the rod-to-rod bipolar synapse. Neuron 41:431-443.
Saszik SM, Robson JG, Frishman LJ (2002) The scotopic threshold response of the dark-
adapted electroretinogram of the mouse. J Physiol 543:899-916.
Scholl HP, Strauss RW, Singh MS, Dalkara D, Roska B, Picaud S, Sahel JA (2016) Emerging
therapies for inherited retinal degeneration. Sci Transl Med 8:368rv366.
Sher A, Jones BW, Huie P, Paulus YM, Lavinsky D, Leung LS, Nomoto H, Beier C, Marc RE,
Palanker D (2013) Restoration of retinal structure and function after selective
photocoagulation. J Neurosci 33:6800-6808.
Sikkink SK, Biswas S, Parry NR, Stanga PE, Trump D (2007) X-linked retinoschisis: an update.
J Med Genet 44:225-232.
Soto F, Kerschensteiner D (2015) Synaptic remodeling of neuronal circuits in early retinal
degeneration. Front Cell Neurosci 9:395.
Vinberg F, Wang T, Molday RS, Chen J, Kefalov VJ (2015) A new mouse model for stationary
night blindness with mutant Slc24a1 explains the pathophysiology of the associated
human disease. Human molecular genetics 24:5915-5929.
Waldner DM, Bech-Hansen NT, Stell WK (2018) Channeling Vision: CaV1.4-A Critical Link in
Retinal Signal Transmission. Biomed Res Int 2018:7272630.
Wang T, Chen J (2014) Induction of the unfolded protein response by constitutive G-protein
signaling in rod photoreceptor cells. J Biol Chem 289:29310-29321.
Wang T, Tsang SH, Chen J (2017a) Two pathways of rod photoreceptor cell death induced by
elevated cGMP. Hum Mol Genet 26:2299-2306.

Wang X, Zhao L, Zhang Y, Ma W, Gonzalez SR, Fan J, Kretschmer F, Badea TC, Qian HH,
Wong WT (2017b) Tamoxifen Provides Structural and Functional Rescue in Murine
Models of Photoreceptor Degeneration. J Neurosci 37:3294-3310.
Wang Y, Fehlhaber KE, Sarria I, Cao Y, Ingram NT, Guerrero-Given D, Throesch B, Baldwin K,
Kamasawa N, Ohtsuka T, Sampath AP, Martemyanov KA (2017c) The Auxiliary Calcium
Channel Subunit alpha2delta4 Is Required for Axonal Elaboration, Synaptic
Transmission, and Wiring of Rod Photoreceptors. Neuron 93:1359-1374 e1356.
Yao K, Qiu S, Wang YV, Park SJH, Mohns EJ, Mehta B, Liu X, Chang B, Zenisek D, Crair MC,
Demb JB, Chen B (2018) Restoration of vision after de novo genesis of rod
photoreceptors in mammalian retinas. Nature 560:484-488.
Yu WQ, Grzywacz NM, Lee EJ, Field GD (2017) Cell type-specific changes in retinal ganglion
cell function induced by rod death and cone reorganization in rats. J Neurophysiol
118:434-454.
Zhang Y, Molday LL, Molday RS, Sarfare SS, Woodruff ML, Fain GL, Kraft TW, Pittler SJ (2009)
Knockout of GARPs and the beta-subunit of the rod cGMP-gated channel disrupts disk
morphogenesis and rod outer segment structural integrity. J Cell Sci 122:1192-1200.
744

Figure 1. Retinas from *Cngb1^neo/neo* mice exhibit stereotypic degenerative changes. (A) The 1.8 kb neomycin cassette, flanked by loxP sites, was inserted into intron 19 of the *Cngb1* gene. (B). Western blots of retinal homogenates from control and *Cngb1^neo/neo* mice show that the neomycin insertion blocked expression of CNGB1, and down regulated expression of CNGA1 channel proteins. C. Light micrograph of representative retinal sections prepared from *Cngb1^neo/neo* mice at the indicated ages. Scale bar = 20 µm. (D-I) Cryosections from 1 MO C57 (left panels) and 1 MO *Cngb1^neo/neo* mice (right panels). Rhodopsin is localized to the outer segment (D) and GFAP to the inner limiting membrane proximal to the ganglion cell layer (E) in C57 retina. In contrast, the outer segment of mutant retina is shortened, and rhodopsin is mislocalized to the outer nuclear layer (F), and GFAP immunoreactivity extends to the entire retina (G). Nuclei are stained with DAPI (blue). (H) C57 and (I) *Cngb1^neo/neo* retinal sections stained for synaptic ribbons (CtBP2, blue) of photoreceptors and mGluR6 puncta (red) of bipolar cell dendrites. Transmission electron microscopy of C57 (J) and *Cngb1^neo/neo* (K) retinal sections. Color coding is as follows: rod spherule (blue), mitochondria (yellow), synaptic ribbon (red), horizontal cell (orange) and rod bipolar cell (green). OS, outer segment; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; GCL, ganglion cell layer.

762
763
764 **Figure 2.** Characterization of rod function by *ex-vivo* electroretinography (ERGs). (A) ERG
765 recordings from a 3 MO C57 mouse show normal a- and b-waves (rod and RBC responses,
766 respectively). Flashes generated 2, 9, 35, 140, 560, and 2200 Rh*/rod (B) ERGs from 1 MO
767 *Cngb1^{neo/neo}* mice show total absence of the b-wave, consistent with disrupted rod-to-RBC
768 signaling. Flashes generated 550, 2200, 8800, 18,000, and 35,000 Rh*/rod.

A

NeoloxP

primers 1+3 ——————— 802 bp
primers 2+3 ——— 620 bp

Cre   -   +   +
TM   -   4d 7d   C

Deletion +
(primers 1+3)

Deletion -
(primers 2+3)

B

        6W   8W
TM   -   -   +   -   +

cre

CNGB1

C

cre   -   +   +
TM   -   4d 7d   C

CNGB1

GC1

PDE6A

GNAT1

GCAP2

Actin

D

3M cre(-)

E

OS

IS

ONL

3M cre(+)

20 um

F

5M cre(-)

G

5M cre(+)

Figure 3. Excision of the floxed neomycin cassette restores CNG channel expression and rescues rod cell death. (A) PCR primers 1, 2, and 3 were designed to detect the presence or absence of the neoloxP cassette. Littermate mice were treated with tamoxifen (TM) for the indicated number of days starting at 4 weeks and retinal DNA was extracted from mice at 8 weeks. Control retinal DNA "c" is from a germline-floxed mouse wherein the neoloxP cassette has been removed in all tissues (*Cngb1ΔCaM*). (B) Western blots of retinal homogenates from cre-negative and cre-positive littermate mice of the indicated ages (6 weeks, 8 weeks) that were treated with TM or vehicle (corn oil) beginning at 4 weeks old for four consecutive days. (C) Western blot of retinal homogenate prepared from the contralateral eye from mice used in (A). Representative retinal morphology (N = 3) of 3 MO cre-negative (D) or cre-positive littermates (E) mice treated with tamoxifen for 7 consecutive days beginning at 4 weeks. TM-treatment of cre-positive mice showed improved outer segment (OS) length and thicker outer nuclear layer (ONL), indicating a halt on cell death. (F) cre-negative and (G) cre-positive littermates treated with tamoxifen for 4 consecutive days starting at P28, and retinal morphology was examined at 5 months of age (representative image from N ≥ 3). Scale bar = 20 µm.

786
787

Figure 4. Light sensitivity is improved following expression of CNGB1. (A) Single cell recordings
show small, desensitized response families in *Cngb1^{neo/neo}* mice likely reflecting residual CNG
channels composed of CNGA1 monomers. Flashes generated 79, 160, 310, 270, 1300, and
2500 Rh*/rod. (B) Following tamoxifen treatment, rod responses showed amplitudes and
sensitivity resembling those of C57; flash strengths were 9, 17, 34, 68, 140, 270, and 540
Rh*/rod.   (C) Response-intensity relationships from single-cell recordings display ~10-fold
reduction in sensitivity between WT (black dots) and *Cngb1^{neo/neo}* (red dots) ($I_{1/2}$ values were 27 ±
4 (n=5) and 360 ± 8 (n=9), respectively). This sensitivity shift is nearly restored following
reintroduction of the CNGB1 (blue dots; $I_{1/2}$ = 43 ± 3 (n=10)).

798
799

Figure 5. Expression of CNGB1 reverses pre- and postsynaptic retinal remodeling in
Cngb1^neo/neo retina. Shown are representative images from N>3 independent experiments. A, D
and G are retinal flat mounts from 1 MO C57, 1 MO *Cngb1^neo/neo* and 3 MO rescued *Cngb1* mice
treated with tamoxifen for 7 consecutive days beginning at 4 weeks of age, respectively. The flat
mounts were stained with the pre-synaptic ribbon marker, CtBP2 (blue), and post-synaptic
marker mGluR6 (orange). Cone pedicles were visualized using cone arrestin, ARR3 (green). B,
E and H are retinal cross sections from mice of the same genotype as the flat mounts. The
retinal sections were stained with antibodies to mGluR6 (red) and the rod bipolar cell marker,
PKCα (teal). C, F and I are frequency vs. mGluR6 puncta size distributions for 1 MO C57, 1 MO
Cngb1^neo/neo and 3 MO rescued *Cngb1* mice treated with tamoxifen.

811
812
Figure 6. Physiological responses from rod bipolar cells in retinal slices. A. *Ex-vivo* ERG
responses from 3 to 6 MO mice after tamoxifen treatments show near normal a- and b-waves.
Flashes generated 2, 9, 35, 140, 550, and 2200 Rh*/rod. Please compare against Fig. 2A.
Voltage-clamp rod bipolar cell recordings (V_m = -60 mV) from the following mice: B. C57 rod
bipolar cells (2-3 MO); C. *Cngb1^{neo/neo}* rod bipolar cells (1 MO); D. *Cngb1* tamoxifen-treated (3
MO). Flashes generated 2, 4, 8, 16, 31, 62 and 130 Rh*/rod for C57 rod bipolar cells, and 280,
560, 1100 and 2200 Rh*/rod for *Cngb1^{neo/neo}* rod bipolar cells. Light-evoked responses were
never observed in *Cngb1^{neo/neo}* rod bipolar cells (15 cells across 5 retinas – flashes generated
2200 Rh*/rod). E.Response-intensity relationships from mean data show that this relationship is
shifted to higher flash strengths in rescued mice, reflecting some rod loss. The Hill exponent of
rod bipolar cells were similar to normal following rod recovery (Hill exponent = 1.6 ± 0.05
(n=12)), in support of a restoration of the normal rod-to-rod bipolar cell synaptic structure and
the dark rate of glutamate release.

Figure 7. Dim flash responses from retinal ganglion cells (RGCs) in whole mount retina show
recovery from early rod rescue.   The top panels (A-C$_1$) show spike times of 100 trials of 3
example cells to a single dim flash (0.75 Rh*/rod).   The middle panels (A-C$_2$) shows PSTHs for
three increasingly bright dim flashes (0.002, 0.02, 0.75, Rh*/rod).   The bottom panels (A-C$_3$)
show the mean spike rate ± SD, measured on each trial in a 100 ms window around the peak of
the PSTH. Dim flash thresholds were estimated from these curves for 1954 cells. The average
cumulative distribution function show higher thresholds responses in *Cngb1$^{neo/neo}$* mice RGCs
than *Cngb1ΔCaM* and tamoxifen-treated *Cngb1$^{neo/neo}$* mice RGCs (D).   The shaded regions
illustrating SEMs across (3-5 experimental preparations).   Additionally, flash thresholds could
not be identified in a larger portion of RGCs from *Cngb1$^{neo/neo}$* mice (E).