1   **A Scube2-Shh feedback loop links morphogen release to morphogen**

2   **signaling to enable scale invariant patterning of the ventral neural tube**

3   **Authors:** Zachary M. Collins[1], Kana Ishimatsu[1], Tony Y.C. Tsai[1], and Sean G. Megason[1,*]
4

[1] Department of Systems Biology, Harvard Medical School, Boston, MA 02115, USA.

*Author for correspondence (megason@hms.harvard.edu)

Running Title: Scaling of Shh patterning by Scube2

Key words: Scaling, morphogen, patterning, sonic hedgehog, robustness, size

Summary Statement: The Shh morphogen gradient can scale to different size tissues by feedback

between Scube2 mediated release of Shh and Shh based inhibition of Scube2 expression

Author contributions

Z.M.C. conducted experiments and data analysis. Z.M.C and S.G.M. conceived the study,

designed the experiments, and wrote the paper. K.I and Z.M.C. developed the size reduction technique.

T.Y.C.T helped develop the image analysis technique and generated the *tg(shha:memCherry)* reporter

line. S.G.M. supervised the overall study.

Abstract

To enable robust patterning, morphogen systems should be resistant to variations in gene expression and tissue size. Here we explore how a Shh morphogen gradient in the ventral neural tube enables proportional patterning in embryos of varying sizes. Using a surgical technique to reduce the size of zebrafish embryos and quantitative confocal microscopy, we find that patterning of neural progenitors remains proportional after size reduction. Intriguingly, a protein necessary for Shh release, Scube2, is expressed far from the source of sonic hedgehog production. *scube2* expression levels control Shh signaling extent during ventral neural patterning and conversely Shh signaling represses the expression of *scube2*, thereby restricting its own signaling. *scube2* is disproportionately downregulated in size-reduced embryos, providing a potential mechanism for size-dependent regulation of Shh. This regulatory feedback is necessary for pattern scaling, as demonstrated by a loss of scaling in *scube2* overexpressing embryos. In a manner akin to the expander-repressor model of morphogen scaling, we conclude that feedback between Shh signaling and *scube2* expression enables proportional patterning in the ventral neural tube by encoding a tissue size dependent morphogen signaling gradient.

Introduction

When Lewis Wolpert first posed the "French Flag Problem", he was seeking the answer to this fundamental question: What systems enable proportional patterning in embryos independent of embryo size? By the time Wolpert formalized this problem, developmental biologists had long known that embryos scale their patterning programs in response to changes in embryo size (Wolpert, 1969). For example, sea urchin larva pattern normally from a single blastomere up to the four-cell stage and amphibian embryos can survive bisection and pattern proportionally at a reduced size (Driesch, 1892; Morgan, 1895; Spemann, 1938; Cooke, 1981). Significant scaling of pattern formation to tissue availability seems to be a near universal property of developing organisms. Yet, as we pass the 50-year

anniversary of Wolpert's work, how morphogen gradients scale to pattern domains of varied sizes

remains unclear in many systems.

Recent theoretical studies have proposed mechanisms that could account for scaling of

morphogen-mediated patterning (Ben-Zvi and Barkai, 2010; Umulis and Othmer, 2013). Amongst the

most prominent of these is a model termed expander-repressor integral feedback control (Ben-Zvi and

Barkai, 2010). In this model, a morphogen represses the expression of another gene, known as the

expander, that affects the range of the morphogen itself cell-non-autonomously. In such models,

morphogen signaling will expand until it has reached an encoded equilibrium. This equilibrium is

controlled by the morphogen's repression of the expander, thus enabling "measurement" of the size of

the domain in need of patterning. The first biological example of this mechanism was proposed in

Xenopus axial patterning. In this original model, ADMP expands BMP signaling by binding Chordin and

inhibiting shuttling of BMP towards the ventral side (Francois et al., 2009). However, more recent

experimental work implicated another factor, Sizzled, which may play a central role in scaling in a

mathematically equivalent manner (Ben-Zvi et al., 2014; Inomata et al., 2013). Expander-like

relationships have also been proposed to regulate scaling of Dpp gradients during wing disc growth and

even scaling of synthetic patterns in bacterial colonies (Ben-Zvi et al., 2011; Cao et al., 2016;

Hamaratoglu et al., 2011).

Though scaling of early axis patterning following size reduction has been extensively studied,

the molecular mechanisms through which tissues and organs subsequently scale their patterning has

received less attention (Ben-Zvi et al., 2008; Inomata et al., 2013). Previously, scaling of patterning

during organ growth has been considered in the fly wing disc, which grows remarkably in size while

maintaining proportion (Averbukh et al., 2014; Ben-Zvi et al., 2011; Hamaratoglu et al., 2011). In

vertebrates, the developing neural tube has been a powerful model to study morphogen-mediated

patterning (Briscoe and Small, 2015). While neural tube patterning does not expand isometrically over

time with growth, embryos of different species maintain consistent embryonic proportions in the face of

significant variation in organ size during initial patterning (Kicheva et al., 2014; Uygur et al., 2016).

The vertebrate ventral neural tube is patterned by the morphogen Sonic Hedgehog (Shh; Marti

et al., 1995; Roelink et al., 1995). Shh is produced by the notochord and floorplate and induces ventral

cell fates over a long range in a dose-dependent manner (Briscoe et al., 2001; Zeng et al., 2001). Shh

ligands themselves are dually lipid-modified and are highly lipophilic (Pepinsky et al., 1998; Porter et al.,

1996a; Porter et al., 1996b). While mechanisms of Shh transport have long been disputed, biochemical

evidence supports soluble Shh as a primary component of long-range signaling, and release of Shh

ligands from cell membranes is critical for gradient formation (Burke et al., 1999; Chen et al., 2004;

Zeng et al., 2001). Shh release was largely thought to be achieved by the protein Dispatched, but

recent work has identified Scube2 as a more potent factor in promoting Shh release (Burke et al., 1999;

Creanga et al., 2012; Kawakami et al., 2002; Tukachinsky et al., 2012).

Scube2 is a Signal sequence containing protein with a CUB domain and EGF-like repeats. The

role of Scube2 in Shh signaling was first identified from work using the zebrafish *you* mutant which

corresponds to *scube2* (Hollway et al., 2006; Kawakami et al., 2005; van Eeden et al., 1996; Woods

and Talbot, 2005; Yang et al., 2002). Interestingly, while *scube2* mutants have defects in ventral

patterning, *scube2* is predominantly expressed in the dorsal and intermediate neural tube in both mice

and zebrafish (Grimmond et al., 2001; Kawakami et al., 2005; Woods and Talbot, 2005). Additionally,

epistasis experiments indicated that Scube2 acts upstream of Patched to stimulate Shh signaling

(Woods and Talbot, 2005). This effect was also found to be cell-non-autonomous, as mosaic injection

of *scube2* mRNA was capable of rescuing Shh-signaling defects over a long range (Hollway et al.,

2006; Woods and Talbot, 2005). Studies in cell culture then demonstrated that Scube2 releases Shh

from secreting cells cell-non-autonomously (Creanga et al., 2012; Tukachinsky et al., 2012). Recent

work concluded that Scube2 may be responsible for catalyzing the shedding of lipids from the Shh

ligands, but this model is disputed by previous findings that released Shh remains dually lipid-modified

(Creanga et al., 2012; Jakobs et al., 2014; Jakobs et al., 2016; Tukachinsky et al., 2012). Scube2's cell

non-autonomous role in Shh release and unexpected expression pattern led us to wonder whether

Scube2 may regulate pattern scaling by acting as an expander, as has recently been hypothesized

elsewhere (Shilo and Barkai, 2017). In this work, we use quantitative imaging of cell fate specification in

zebrafish to investigate the scaling of ventral neural patterning and the regulatory role of Scube2.

Results

Ventral neural patterning scales with embryo size

Studying mechanisms of scaling during growth or between species of different sizes is difficult

because many properties of the patterning system depend on stage or species-specific variables. To

study scaling of pattern formation in embryos with comparable genetic backgrounds at matched time

points, we developed a technique to reduce the size of zebrafish embryos inspired by classical work in

amphibians, as we recently described (Ishimatsu et al., 2018; Morgan, 1895; Spemann, 1938). Two

lateral cuts are made across the blastula stage embryo: one to remove cells near the animal pole, to

avoid damaging signaling centers crucial to early D-V patterning, and a second to remove yolk near the

vegetal pole. (Fig. 1A). With this technique, a significant fraction of embryos pattern normally and

develop at a reduced size (Fig. 1B-C).

We measured scaling of neural patterning in size-reduced embryos using quantitative imaging

(Megason SG, 2009; Xiong et al., 2013). High-resolution image-stacks of 18-24 hours postfertilization

(hpf) stage matched zebrafish embryos were collected under identical settings, during the same

imaging session, from matched Anterior-Posterior positions in control and experimentally perturbed

embryos (Fig. 1D-E). Imaging volumes were analyzed with custom built software to manually

demarcate the dorsoventral axis and width of the neural tube along the length of the dataset (Fig. 1F,

Fig. S1). Image intensity values were extracted in a set number of bins along the D-V axis for the left

and right halves of the neural tube to normalize for variability in neural tube height. This system allowed

for the quantitative and unbiased comparison of 3-4 somite lengths of neural imaging data from multiple

embryos.

Using this imaging platform, we compared the expression of *patched2*—a direct transcriptional

target of Shh—with the *tg(ptch2:kaede)* reporter in wild type and size-reduced embryos (Fig. 1G-J)

(Huang et al., 2012) . When quantified relative to their respective neural tube dorsal-ventral heights,

tg(ptch2:kaede) response gradients maintained nearly identical intensity distributions despite neural

tube height being 15.0% (+/- 2.8%) smaller in size-reduced embryos in this dataset (N=5), indicating

that Shh responses scale following size reduction (Fig. 1I-J). When viewed on an absolute scale,

control and size-reduced embryos show clear shifts in the response gradient as measured by the

position at which 50% of mean control maximum intensity is reached (p=0.0076) (Fig. 1J). To quantify

this effect at the level of cell fate specification, we utilized a triple transgenic imaging strategy based on

reporter lines marking *nkx2.2a* (p3 progenitors), *olig2* (pMN and some p3 progenitors), and *dbx1b* (p0,

d6 progenitors) (Fig. 1K-L) (Gribble et al., 2007; Jessen et al., 1998; Kinkhabwala et al., 2011; Kucenas

et al., 2008). Anterior-posterior averaged intensity profiles were then segmented to form cell fate

profiles (see methods and Fig. S2). Using this method, we generated cell fate profiles which can be

compared between embryos (Fig. S2). After normalizing for their altered D-V height (which was

reduced in this population by 12.2% +/- 2.4% compared to controls), the average of these cell fate

profiles of size-reduced embryos were virtually indistinguishable from those of full sized embryos (Fig.

1M). Furthermore, differences between progenitor domain boundary positions were visible when size

normalization was removed (Fig. 1N). Statistically significant shifts in the positions of the p2 and pMN

upper boundaries were observed only when compared in their absolute coordinates (Fig. 1N). This

further demonstrates that ventral neural patterning adjusts to changes in total D-V height.

Figure 1- Neural tube patterning scales following embryonic size reduction.

(A) Surgical size reduction of 128-256 cell stage embryos in which cells and yolk are removed to produce smaller embryos (images adapted from Ishimatsu et al. 2018). (B) Example of a size-reduced embryo at 24 hpf (lower) with a normal-sized sibling (upper). (C) Example of a size-reduced larva at six days post fertilization (lower) with a normal-sized sibling (upper). (D) Schematic of an embryo mounted

for imaging; anterior-posterior length of the imaging window is shown with blue lines. Red lines indicate

the 3-D extent of the imaging window. (E) 3-D rendering of a confocal z-stack on an example

tg(ptch2:kaede) mem-mCherry mRNA injected image volume. Scale bar represents 100 µm. (F)

Flattened "fillet" profile of segmented imaging data from a z-stack imaged as in E (See methods).

(G,H,K,L) Scale bar represents 20 µm. (G-H) Transverse view of 20 hpf *tg(ptch2:kaede)*, *mem-mCherry*

mRNA injected control (G) and size-reduced embryos (H). Size reduction led to a decrease in DV

height of 15.0% +/- 2.8% in this dataset. (I-J) *ptch2:kaede* intensity profiles from segmented imaging

data of embryos from G-H plotted either normalizing for D-V neural tube height (I) or relative to their

absolute neural tube heights (J). Error bars indicate standard deviation. No significant shift is observed

in the position of 50% mean maximum control intensity on a relative scale between control and size-

reduced embryos(I) (unpaired t-test p=0.5981) while a very statistically significant shift in the position is

observed when comparing absolute positional values (J) (unpaired t-test p=0.0076; Control N=5, Size

Reduced N=5). (K-L) Transverse views of *mem-mtagBFP2* mRNA injected 24 hpf *tg(nkx2:mGFP;*

olig2:dsred; dbx1b:GFP) control (K) and size-reduced (L) embryos. Size reduced embryos had an

average 12.2% +/- 2.4% reduction in neural tube DV height relative to control embryos in this dataset.

(M-N) Results of progenitor domain segmentation analysis (See methods) of embryos treated as in K-L

plotted either normalizing for differences in neural tube height (relative scale) (M) or with respect to their

endogenous heights (absolute scale) (N). Only when compared on an absolute scale are statistically

significant shifts seen in the dorsal boundaries of the p1-p2 and pMN domains (absolute heights

unpaired t-test p^{p2}=1.0172e-4 and p^{pMN} =0.0016, Control N=5, Size Reduced N=4). Thresholds for P

value significance asterisks: Extremely significant: (< 0.0001) = ****, (0.0001 to 0.001) = ***; Very

significant: (0.001 to 0.01) = **; Significant: (0.01 to 0.05) = *.

Scube2 levels control Shh signaling

Based on its role in the cell-non-autonomous regulation of Shh release and its dorsal expression pattern, we hypothesized a potential role for Scube2 in enabling scaling of Shh gradients. This hypothesis depends on *scube2* expression levels having a dose dependent effect on Shh signaling. However, previous work concluded that Scube2 is only required for Shh signaling as a permissive factor (Kawakami et al., 2005; Woods and Talbot, 2005). To examine the role of Scube2 in ventral neural patterning, we performed a morpholino knockdown of *scube2* in *tg(ptch2:kaede)* reporter embryos using a previously validated translation inhibiting morpholino (Fig. 2A-C) (Woods and Talbot, 2005). We observed a decrease in Shh signaling following morpholino injection, as demonstrated by a statistically significant suppression of maximum *ptch2:kaede* intensity (Fig. 2C)(Woods and Talbot, 2005)(Woods and Talbot, 2005)(Woods and Talbot, 2005)(Woods and Talbot, 2005)(Woods and Talbot, 2005). Additionally, quantification of *nkx2.2a, olig2*, and *dbx1b* domain sizes in embryos injected with *scube2* morpholino showed a contraction of ventral progenitor domains (Fig. 2D-F). Ventral shifts in the upper boundaries of the pMN and p3 domains were statistically significant, due in part to near complete elimination of the *nkx2.2a+* p3 domain (Fig. 2F). Unexpectedly, expansion of the p1-2 domain dorsally was also observed, potentially implying long range Shh signaling is required for complete p1-2 induction.

Previous work concluded that Scube2 was a permissive factor based on a lack of ectopic expression of downstream Shh signaling markers as observed by whole mount *in situ* hybridization experiments (Woods and Talbot, 2005). However, our high-resolution quantitative imaging reveals that injection of *scube2* mRNA leads to the expansion of Shh signaling, as shown by broader distributions of *tg(ptch2:kaede)* fluorescence (Fig. 2G-I). Embryos injected with *scube2* mRNA showed significant dorsal shifts in the position of half control maximum *tg(ptch2:kaede)* intensities (p=0.0047), a measure of absolute Shh signaling range (Fig. 2H). Maximum *tg(ptch2:kaede)* fluorescence in *scube2* overexpressing embryos was somewhat higher on average, but not statistically significant (p=0.0512),

which may suggest the stronger effect of *scube2* overexpression is to extend the range of Shh

signaling. In addition, *scube2* overexpression affected cell type patterning in the ventral neural tube, as

measured in triple transgenic *nkx2.2a, olig2,* and *dbx1b* reporter embryos (Fig. 2J-L). Quantification of

these cell fate profiles revealed large increases in p3 and pMN domain sizes, a decrease in the size of

the p2-p1 domains, and unchanged patterning of the p0-d6 domains and more dorsal cell types.

Ventralization was measured by comparing dorsal boundaries of the p3 and pMN, which were

statistically significantly shifted (Fig. 2L). These data indicate that not only is Scube2 required for long

range Shh signaling, but that *scube2* overexpression amplifies endogenous Shh signaling. Additionally,

this suggests that Scube2-stimulated Shh release is a limiting factor in normal patterning.

Figure 2- Scube2 expression levels regulate Shh signaling in the ventral neural tube.

(A-B,D-E, G-H, J-K) Transverse view of a confocal z-stack, scale bar represents 20 μm (A-B)

Representative image of 22 hpf *tg(ptch2:kaede*) reporter line embryos injected with (A) *mem-mCherry*

mRNA with a control morpholino or (B) injected with *mem-mCherry* mRNA and *scube2* morpholino. (C)

Quantification of mean intensity distributions in segmented neural tissue from z-stacks of embryos as

treated in A-B. Maximum intensities of morpholino treated embryos were very statistically significantly

reduced compared to controls (p= 0.0040). (D-E) Representative image of 20 hpf *tg(dbx1b:GFP,*

olig2:dsred, nkx2.2a:memGFP) reporter line embryos injected with (D) mem-*mTagBFP2* mRNA alone

or (E) co-injected with *scube2* morpholino. (F) Mean result of automated segmentation of progenitor

domain sizes (see methods for details) for embryos treated as in D-E. Statistical comparisons of

progenitor domain boundaries are shown with connected lines and significance is marked by asterisks.

p1-2 domain upper boundaries were shifted dorsally following morpholino treatment (p^{p1-2}= 0.0019),

while the upper boundary of pMN and p3 domains were both significantly contracted in morpholino

injected embryos (p^{pMN}= 0.0158 and p^{p3}= 9.87e-9). (G-H) Representative image of 20 hpf

*tg(*ptch2:kaede) reporter line embryos injected with (G) *mem-mcardinal* mRNA alone or (H) co-injected

with *scube2* mRNA. (I) Quantification of mean intensity distributions of embryos as treated in G-H.

Dorsoventral position at which half of average control maximum intensity was reached was very

significantly shifted in Scube2 overexpressing embryos (unpaired t-test p=0.00470). (J-K)

Representative image of 20 hpf *tg(dbx1b:GFP, olig2:dsred, nkx2.2:mGFP*) reporter line embryos

injected with (J) *mem-mTagBFP2* mRNA alone or (K) co-injected with *scube2* mRNA. (L) Mean results

of automated progenitor domain segmentation of J-K. Statistical comparisons of progenitor domain

boundaries are shown with connected lines and significance marked by number of asterisks. pMN and

p3 domains were drastically shifted dorsally in *scube2* mRNA injected embryos (p^{MN}=8.6748e-04, and

p^{p3}=0.0034 respectively).

Shh negatively regulates Scube2 expression over a long-range

To study Scube2's expression, we developed the *tg(scube2:moxNG)* reporter line containing

7.6KB of the endogenous regulatory sequences driving the extremely bright moxNeonGreen

fluorescent protein (Fig. 3A) (Costantini et al., 2015). The expression of *tg*(*scube2:moxNG*) we

observed is consistent with previously reported *in situ* hybridizations (Grimmond et al., 2001; Kawakami

et al., 2005; Woods and Talbot, 2005). *Tg(scube2:moxNG)* embryos showed very low expression close

to the sources of Shh in the floor plate and notochord—as visualized with a transgenic *shh:memCherry*

reporter line—and high levels of expression in the dorsal-intermediate neural tube (Fig. 3A-C). Time

lapse imaging of *tg(scube2:moxNG)* embryos revealed weak mesodermal expression in the early

embryo, which faded during the onset of neurulation and was replaced by high levels of expression in

the dorsal and intermediate neural tube (Fig. 3D-G, Movie S1-2).

To test whether Scube2 is downregulated by Shh signaling, we injected mRNA encoding a

potent activator of Shh signaling, dnPKA, at the single cell stage and observed the resulting embryos

(Hammerschmidt et al., 1996). Embryos injected with *dnpka* mRNA showed near complete ablation of

neural *tg(scube2:moxNG)* expression (Fig. 4 A-C). To test whether Shha ligands themselves were

capable of suppressing Scube2 expression at a distance, we mosaically overexpressed *shha* in

tg(scube2:moxNG) embryos by injecting a single blastomere at the 16-cell stage with either *mem-*

mTagBFP2 alone or with *shha* mRNA (Fig. 4 D-F). We expected local inhibition of Scube2 reporter

activity near secreting cells within a few cell diameters. Surprisingly, *tg(scube2:moxNG)* expression was

nearly completely eliminated in these embryos indicating potent cell-non-autonomous repression of

Scube2 by Shh. When quantified, these embryos demonstrate a highly significant reduction of peak

tg(scube2:moxNG) intensities (Fig. 4F). To test whether Shh's inhibition of Scube2 is required for its

endogenous low ventral expression, we treated embryos with sonidegib, a potent Smoothened

antagonist starting at the dome stage. Resulting embryos showed expanded *scube2* expression

towards the floor plate and notochord (Fig. 4G-I). Shifts in ventral boundaries were quantified by

measuring the D-V position at which 50% of the maximum intensity of the control population was

reached. These measurements were statistically significantly shifted in sonidegib-treated embryos

relative to controls, indicating that endogenous Shh signaling is responsible for a lack of ventral *scube2*

expression (Fig. 4I). A ventral expansion *tg(scube2:moxNG)* was also observed following Cyclopamine

treatment, which is consistent with the previous findings of a genomewide screen for genes regulated

by Shh signaling (Fig. S3) (Xu et al., 2006). To further probe the transcriptional regulation of Scube2's

expression we performed a small scale CRISPR mutagenesis screen and found that Pax6a/b are

necessary for driving Scube2 expression. Coinjection of pax6a and pax6b sgRNAs with Cas9 caused

significant downregulation of *tg(scube2:moxNG)* relative to control embryos injected with a sgRNA

targeting *tyrosinase,* an unrelated pigment gene (Fig. S4).

Figure 3- Scube2 is expressed distantly from

Shh secreting cells.

(A) Schematic of the *scube2:moxNG* transgenic

expression reporter construct used to generate the

tg(scube2:moxNG) line. (B) Wide-field fluorescence

image of *tg(scube2:moxNG)* embryo at 20 hpf. (C)

Transverse view of *mem-mTagBFP2*-injected

tg(scube2:moxNG; shh:mem-mCherry) embryo at 20

hpf. Scale bar represents 20 µm. (D-G) Transverse

view from a time-lapse imaging dataset of

tg(scube2:moxNG; shh:memCherry) embryo which

was injected at the single cell stage with *mem-*

mTagBFP2 mRNA. Time in hours post fertilization is

displayed in the bottom right corner. Scale bar

represents 50 µm. (D) At early neurulation stages

there is weak mesodermal expression of

scube2:moxNG in the notochord. In addition,

expression of *scube2:moxNG* is visible in neural

progenitors as the neural plate converges. (E) By

12.5 hpf a pronounced gap in expression of neural

progenitors between *shh:mem-mCherry* and

scube2:moxNG cells is visible. (F) Expansion of the

scube2+ domain dorsally is visible as cells continue

to converge. (G) *scube2* expression is constricted to

the dorsal-intermediate neural tube.

Figure- 4 Shh signaling represses Scube2 expression.

(A-B, D-E, G-H) Scale bar represents 20 μm (A-B) Transverse view of 18 hpf *tg(scube2:moxNG)*

reporter line embryos injected with (A) *mem-mTagBFP2* mRNA alone or (B) co-injected with *dnpka*

mRNA. (C) Quantification of mean reporter intensity of embryos as treated in A-B. Maximum

scube2:moxNG intensity values were very significantly reduced in *dnpka* mRNA-injected embryos (p=

0.0014). (D-E) Transverse view 20 hpf *tg(scube2:moxNG)* reporter line embryos injected at the single

cell stage with *mem-mCherry* mRNA and then injected in one blastomere at the 8-16 cell stage with

either (D) *mem-mTagBFP2* mRNA alone or (E) co-injected with *shha* mRNA. (F) Quantification of mean

reporter intensity of embryos as treated in D-E. Maximum *scube2:moxNG* reporter intensity is extremely

significantly reduced in *shha* injected embryos (p= 9.14e-06). (G-H) Transverse view 22 hpf

tg(scube2:moxNG; *mem:mCherry)* embryos treated with a DMSO control (G) or treated with 50 μm

Sonidegib (H). (I) Quantification of mean reporter intensity of embryos as treated in G-H. The black

bracket marks the position of 50% of control maximum intensity used for statistical testing. These

values were very significantly shifted ventrally in drug treated embryos relative to control (unpaired t-

test p=0.0014.).

Scube2 diffuses over long distances during patterning

While Scube2 is known to act cell non-autonomously from transplantation experiments and

Scube2-conditioned media has a potent Shh release stimulating effect *in vitro*, the localization of

Scube2 protein during development is unknown (Woods and Talbot, 2005; Creanga et al., 2012). *In*

vitro, Scube2 is thought to associate with Heparin Sulfate Proteoglycans, and Scube2 had been

hypothesized to diffuse from secreting cells in the dorsal neural tube, the need for which was later

disputed (Jakobs et al., 2016; Kawakami et al., 2005; Hollway et al., 2006). To examine Scube2's

localization, we developed Scube2 fluorescent fusion proteins by tagging the C-terminus as previously

validated in cell culture with other tags (Fig. 5A) (Creanga et al., 2012). The resulting Scube2-mCitrine

fusion proteins were functional and rescued Scube2 CRISPR mutants at comparable rates to wildtype

Scube2 (Fig. S5). Mosaic injection of *scube2-mCitrine* mRNA at the 32-64 cell stage revealed that

Scube2-mCitrine diffuses distantly from producing cells (Fig. 5B). Following single cell mRNA injection,

Scube2-mCitrine fusions were secreted and did not remain associated with cell membranes, as

demonstrated by their presence in the extracellular space between cells marked with mem-mCherry

(Fig. 4C).

To assay Scube2's rate of diffusion we performed Fluorescence Recovery After Photobleaching

(FRAP) at the dome stage, during which cell movement is minimal. FRAP was performed in a 100µm x

100µm region and recovery was observed at 10 second intervals over 5 minutes (Fig. 5 D-E, Movie S3-

4). Image data from the bleached region was then normalized to its initial intensity and fitted to recovery

curves using standard methods (Munjal et al., 2015). We find that the addition of Scube2 to mCitrine

causes only minor changes to its diffusion (Fig. 5F-G). No significant differences were observed

between the calculated mobile fractions of Scube2-mCitrine (0.283 +/- 0.018) and Sec-mCitrine

(0.338+/- 0.019; unpaired t-test p=0.0698). When the time to 50% recovery of the mobile fraction is

calculated, Scube2-mCitrine has modestly slower diffusion time than Sec-mCitrine alone (Fig. 5 F-G;

p=0.0405). These data further support Scube2's diffusion in the extracellular space, which likely

mediates its long range of effect.

To observe distributions of Scube2 during development, we generated a transgenic line

expressing the full length Scube2 protein fused to moxNeonGreen under control of Scube2 regulatory

sequences (Fig. 5H-K). We validated the functionality of this *Tg(scube2:scube2-moxNG)* line using a

morpholino which bound only endogenous Scube2 at the splice junction of exon6, and not

Tg(scube2:scube2-moxNG) derived RNA which lacks this splice junction (Figure S6).

Tg(scube2:scube2-moxNG) embryos were markedly resistant to treatment with this morpholino,

validating the in vivo functionality of this construct (Figure S6). *Tg(scube2:scube2-moxNG)* embryos

showed broad distributions of Scube2 during patterning (Fig. 5H-K). Throughout early patterning

Scube2-moxNeonGreen is visible near ventral cells marked by *tg(shha:mem-mCherry)*, although

Scube2 is expressed largely in the dorsal neurectoderm at this timepoint (Fig. 5H-I, Movie S5-6). By 24

hpf *tg(scube2:scube2-moxNG)* fluorescence is found distributed throughout the embryo, although

expression from *tg(scube2:moxNG)* is localized to the dorsal-intermediate neurectoderm. These data

further suggest that Scube2's long range of effect can be explained by diffusion from secreting cells in

the intermediate and dorsal neural tube to the source of Shh in the floor plate and notochord.

Figure 5- Scube2 diffuses from secreting cells and is broadly distributed during

patterning.

(A) Schematic of Scube2-mCitrine fluorescent fusion protein design. (B-C) Scale bar represents 100

µm. (B) Scube2-mCitrine fluorescence at the sphere stage from embryos injected in one blastomere at

the 64-cell stage with *scube2-mCitrine* mRNA. Cells from the injected clone are marked by intracellular

(secretory system) fluorescence. (C) Scube2-mCitrine fluorescence from embryos injected at the single

cell stage with *scube2-mCitrine* and *membrane mCherry* mRNA. (D-E) Fluorescence recovery after

photobleaching at the dome stage of Scube2-mCitrine (D) and Secreted-mCitrine (E). (F) Raw FRAP

recovery traces normalized to maximum intensity pre-bleach and minimum intensity following

bleaching. Red lines represent Secreted-mCitrine and blue lines represent Scube2-mCitrine. (G) A

boxplot comparing the time at which 50% recovery of mobile fraction is achieved from dome stage

FRAP data. Bracket with asterisk denotes unpaired t-test comparison between Scube2-mCitrine and

Sec-mCitrine (p =.0405) (H-K) Scale bar represents 20 µm. (H) Transverse view of an 11.5 hpf

tg(sc2:sc2-moxNG;shh:mem-mCherry) embryo. (I) Transverse view of a 14 hpf *tg(sc2:sc2-moxNG;*

shh:mem-mCherry) embryo. (G) Transverse view of a 24 hpf *tg(scube2:scube2-moxNG)* embryo. (K)

Horizontal view of the embryo from J.

Feedback regulation of Scube2 levels is necessary for pattern scaling

To examine the regulation of Scube2 in size-reduced embryos, we performed our size reduction

technique on *tg(scube2:moxNG; shha:mem-mCherry)* embryos and imaged them at 20 hours post

fertilization. Unlike other observed patterning genes, *scube2* expression levels did not scale in size-

reduced embryos but were instead severely reduced (Fig. 6A-C). This finding is consistent with an

expander-repressor-like model of Scube2-Shh. In this regime, inhibition of *scube2* expression in size-

reduced embryos would contract Shh signaling, enabling adjustment of Shh signaling for a decreased

tissue size (Fig. 6D-E).

Next, we examined whether feedback control of *scube2* expression levels by Shh signaling is

required for pattern scaling by saturating *scube2* levels by over-expression. If Scube2 is responsible for

adjusting Shh signaling during scaling, we would expect *scube2*-overexpressing size-reduced embryos

to have the same absolute Shh response profiles as controls, which would fail to scale following size

normalization (Fig. 6F-G). If scaling of ventral patterning is not dependent on Scube2, we would expect

maintenance of pattern scaling with size-proportionate increases in *ptch2:kaede* distributions in both

populations. We overexpressed *scube2* by mRNA injection in *ptch2:kaede* reporter embryos. When

normalized for differences in D-V heights, size-reduced *scube2*-overexpressing embryos showed a

disproportionate expansion of the Ptch signaling gradient compared to normal-sized *scube2*-

overexpressing embryos (Fig. 6H-K). Dorsal expansion of Shh signaling is quantified using the position

of 50% of average maximum control intensity, which is statistically significantly shifted dorsally in size-

reduced embryos (Fig. 6J). Importantly, when *tg(ptch2:kaede)* response profiles are plotted on an

absolute rather than relative scale, they nearly exactly overlap (Fig. 6K). This overlap without size

normalization suggests that *scube2* overexpression encodes a response profile which is independent of

embryo size and is not secondarily tuned by another scaling related factor. This strongly suggests that

control of *scube2* expression levels is required for scaling the Shh response gradient.

Figure 6- Scube2 expression is size-dependent and required for pattern scaling.

(A-B) Transverse view of *mem-mTagBFP2* mRNA-injected *tg(scube2:moxNG; shh:mem-mCherry)*

control (A) or size-reduced (height reduction of 17.1% +/- 7.7%) (B) embryos at 20 hpf. (C)

Quantification of mean *tg(scube2:moxNG)* intensity versus ventral-to-dorsal position of embryos from

A-B. Maximum intensity values are statistically significantly reduced in treated embryos (p=3.03e-4). (D-

E) Schematic of expander-repressor-like feedback control of Shh signaling by Scube2 and its ability to

enable pattern scaling. Repression of Scube2 by Shh encodes an equilibrium level of Shh signaling

across the tissue by linking morphogen spread to tissue size. (F-G) Schematic representation of the

experiment as performed in H-K, where Scube2 levels are at saturation due to overexpression, and

size-reduced embryos (G) are disproportionately affected. (H-I) Transverse view of 20 hpf

tg(ptch2:kaede) control (H) and size-reduced (I) embryos injected with *mem-mCherry* and *scube2*

mRNA. (J) Quantification of mean *tg(ptch2:kaede)* intensity versus normalized ventral-to-dorsal position

of embryos treated as in H-I. Very significant shifts are observed in the dorsal position of 50% of the

maximum intensity value (unpaired t-test p=0.0092). (K) Intensity profiles from J rescaled to reflect the

absolute scale of the measurements (DV height reduction of 15.2% +/- 2.1%). When measured in these

coordinates, no difference is observed in the position of half of maximum intensity (p= 0.9760).

Discussion

Our work uncovers that the morphogen Sonic Hedgehog can self-regulate to enable scale-

invariant patterning through linking morphogen signaling to inhibition of Scube2. We discovered that

patterning of the neural tube adjusts to tissue availability following surgical size reduction in zebrafish

embryos. Using overexpression experiments we demonstrate that Scube2's activity during patterning is

not just permissive—overexpression of *scube2* quantiatively enhances Shh signaling (Woods and

Talbot, 2005). Utilizing a transgenic reporter line which we developed, we characterized the expression

of Scube2 during neural patterning and found that Shh signaling is responsible for its repression in the

ventral neural tube. Using Scube2 fluorescent fusion proteins we found that Scube2 is broadly

distributed from secreting cells, explaining its previously reported cell non-autonomous activity

(Creanga et al., 2012; Woods and Talbot, 2005). Unlike other patterning genes, *scube2* responds to

changes in neural tube height by disproportionately decreasing its expression, and overexpression of

Scube2 inhibits scaling of the Shh signaling gradient by circumventing its feedback control. The

expression of *scube2* thus can be seen as comparable to the "size-dependent factor" Sizzled, which is

thought to enable scaling in early D-V patterning by tuning its expression levels to embryo size by

chordin-dependent feedback inhibition (Inomata et al., 2013).

The relationship between Scube2 and Shh has important similarities to proposed "expander-

repressor" models of morphogen scaling (Barkai and Ben-Zvi, 2009; Ben-Zvi and Barkai, 2010; Inomata

et al., 2013). As with expanders in these models, *scube2* enhances morphogen range, is repressed by

morphogen signaling, and acts cell non-autonomously at a distance from its source. However, Scube2's

reported role in morphogen release may be distinct from the proposed mechanism of expanders.

Expanders extend the range of morphogens by promoting their diffusion or inhibiting their degradation

(Ben-Zvi and Barkai, 2010). While release of Shh ligands from secreting cells would support their

transport, the potentially irreversible nature of this effect and local action at the morphogen source

would make distinct predictions for Scube2's effects on morphogen distributions. Nonetheless our study

marks the first observation of an expander-repressor-like relationship outside of the BMP/Dpp signaling

pathway in a developing organism. This finding raises the possibility that expander-repressor-like

relationships may be common motifs in the regulation of morphogen gradients.

We began this work in part due to interest in the discrepancy between the area of Scube2's

activity in the ventral neural tube and its expression in the dorsal neural tube. Our work with Scube2

fluorescent protein fusions revealed that Scube2 is diffusive and is distributed broadly from producing

cells. Scube2's diffusion from producing cells could easily account for the distance between its

expression domain and area of effect (Figure 5). Scube2's broad distribution and considerable

extracellular diffusivity bolsters the hypothesis that Scube2 may serve as a chaperone for Shh during its

transport as hypothesized previously (Tukachinsky et al., 2012). Cell culture experiments have

indicated that Scube2 cooperates with Dispatched in a cholesterol-dependent "hand off" by binding

different domains on Shh's cholesterol moiety (Tukachinsky et al., 2012). Continued binding of Scube2

to the hydrophobic sterol domain may facilitate the un-hindered diffusion of Shh through the

extracellular millieu. This model is consistent with the dose dependency we observe in our Scube2

overexpression experiments (Figure 2G-L) and may help solve the puzzle of the long-range transport of

dually lipid modified hedgehog in vertebrates. Further investigation of the strength and duration of

Scube2 and Shh's binding in vivo may shed light on this relationship. Unfortunately, direct imaging of

this phenomenon is hampered by the lack of fully functional Shh fluorescent protein fusions

(Chamberlain et al., 2008).

While some evidence suggests that Scube2 plays a role in lipid-shedding, these observations

conflict with previous HPLC analysis and the findings of independent groups which demonstrated that

Shh species released by Scube2 are dually lipid modified (Creanga et al., 2012; Tukachinsky et al.,

2012). If correct, a model of Scube2 in which it acts only transiently at the cell surface of producing

cells—either by enabling the formation of multimeric Shh complexes or lipid shedding—would have

interesting implications for its role as an expander. Expanders are often formalized as having a dose

dependent reversible effect on morphogen spread, while a transient role of Scube2 in Shh multimeric

complex formation or shedding would be localized and irreversible. Mathematical modeling may reveal

interesting implications of each proposed mechanism in Scube2-Shh feedback interactions during

pattern scaling.

Scube2 is one of several recently identified elements of the Shh signaling pathway that exerts

cell non-autonomous effects. Recent work has shown that Hhip—initially characterized as a membrane-

tethered hedgehog antagonist—acts over a long range that cannot be explained by ligand

sequestration (Kwong et al., 2014). Additionally, the Hedgehog receptor, Patched, may also have cell

non-autonomous inhibitory effects on Smoothened through regulating inhibitory sterols or sterol

availability (Bidet et al., 2011; Roberts et al., 2016). Together with known feedback relationships and

the diffusivity of Scube2 that we demonstrated here, these mechanisms interlink Shh signaling between

neighboring cells and may enable tissue level properties, such as the scaling of pattern formation we

observed.

However, scaling of neural patterning is unlikely to be achieved by regulation of Shh signaling

alone. BMP signaling in the dorsal neural tube is known to pattern dorsal progenitors. Scaling of BMP

signaling in dorsal neural patterning may be achieved via a similar or carry-over mechanism to early D-

V axis patterning and should be explored in further studies. In the early D-V patterning system, both

existing models propose expander-like relationships between elements of the BMP signaling pathway.

The first model proposed ADMP as a scaling related factor, while more recent research has

demonstrated that Sizzled has an indispensable role in scaling (Ben-Zvi et al., 2008; Ben-Zvi et al.,

2014; Inomata et al., 2013). During neural patterning, the BMP antagonists Noggin, Follistatin, and

Chordin are expressed in the notochord while BMP ligands are expressed in the roof plate. Intriguingly,

while Sizzled does not seem to be expressed during neural patterning, ADMP is expressed in the

notochord and thus may play a role in the scaling of BMP-mediated patterning of the dorsal neural tube

(Willot et al., 2002).

BMP signaling is known to increase the thresholds for Shh-dependent cell fate specification,

making signaling integration between these pathways a potential additional candidate regulator of

scaling (Liem et al., 2000; McHale et al., 2006). Inhibition of either Shh or BMP signaling causes

expansion of signaling by the alternative program. In normal patterning, cells do not measure ratios of

BMP and Shh. In fact, Dbx1 positive progenitors in the medial neural tube require little to no Shh or

BMP signaling present in order to be specified (Pierani et al., 1999). In addition, recent experiments

with precise control of Shh and BMP concentrations in an explant system have shown that cells choose

either ventral or dorsal fates in the presence of significant BMP and Shh signaling (Zagorski et al.,

2017). Regulation of *scube2* expression may be another way to enable crosstalk between signaling

pathways, as *scube2* is not expressed in the dorsal most cells of the spinal cord, suggesting repression

by dorsal factors. Specification of the dorsal boundary of *scube2* expression may encode yet more

information about the size of the tissue which would then affect Shh spread.

Methods

Generation of Transgenic Lines

The construct used to make *tg(scube2:moxNG)* was generated by isothermal assembly of PCR-

amplified *scube2* regulatory elements obtained from the CHORI-211 BAC library. Regulatory elements

were in part chosen based on annotations of H3K4me1 and H3K4me3 binding (Aday et al., 2011).

Selected regulatory sequences spanned 1677bp of upstream intergenic sequence and 5962bp of the

area spanning exons 1-5 *scube2*. Regulatory sequences were cloned into a pMT backbone by placing

a zebrafish codon-optimized moxNeonGreen fluorescent protein and sv40 poly-A tail just downstream

of the endogenous *scube2* Kozak sequence (Costantini et al., 2015). The construct used to make

tg(scube2:scube2-moxNG) was generated using the same regulatory sequences as

tg(scube2:moxNG), with the addition of cDNA corresponding to exons 6-23 of the Scube2 coding

sequence downstream of exon 5 and moxNeonGreen attached at the c-terminus with a 10 amino acid

long GA rich linker. The construct used to make *tg(shh:mem-mCherry)* was derived from the previously

reported *tg(shh:GFP*), by replacement of GFP with mem-mCherry (Megason, 2009; Shkumatava et al.,

2004).

Transgenic lines were generated by injecting plasmid DNA for each construct along with Tol2

mRNA into wild type (AB) embryos at the single cell stage, as described previously (Kawakami, 2004).

moxNeonGreen positive embryos were then selected for raising. Upon reaching sexual maturity, F0s

were outcrossed and screened for founders. Founders were isolated and raised as single alleles.

Monoallelic versions of each line are shown throughout the paper.

Zebrafish Strains

For wild type lines, AB fish were used. All fish were kept at 28°C on a 14-hour-light/10-hour-dark

cycle. Embryos were collected from natural crosses. All fish-related procedures were carried out with

the approval of Institutional Animal Care and Use Committee (IACUC) at Harvard University.

tgBAC(*ptch2:kaede*) (Huang et al., 2012; renamed from *ptch1* due to a change in zebrafish gene

nomenclature), *tg(nkx2.2a:mGFP)* (Jessen et al., 1998), *tg(olig2:GFP)* (Shin et al., 2003),

tg(olig2:dsRed) (Kucenas et al., 2008), and tgBAC(*dbx1b:GFP)* (Kinkhabwala et al., 2011) have been

described previously.

Size Reduction Technique.

Size reduction was performed as described in our previous report (Ishimatsu et al., 2018). Embryo

sizes were reduced by sequentially removing ~1/3 of the cells from the animal cap, then wounding the

yolk. These surgeries are performed in 1/3 ringers solution, and embryos are immobilized in a thin layer

of 2% methyl cellulose. Surgeries can be performed either with glass needles – as previously described

– or using a loop of thin stainless-steel wire that is inserted through a glass capillary tube and mounted

on a halved chopstick as done here. Healthy uninjected embryos show a maximum success rate of

~60% while embryos which have undergone injection or were spawned by older females have

significantly lower success rates. In each size reduction experiment, embryos are screened for health

and the largest size reductions; those with insufficient size reduction or morphological defects are

discarded.

Construct Generation and Injections of mRNAs and Morpholinos

Scube2-mCitrine was generated from cDNA obtained from the Talbot lab (Woods and Talbot, 2005).

Fluorescent protein fusions were made by attaching mCitrine or moxNeonGreen with a 10 amino acid

GA rich linker to the c-terminus of Scube2. Membrane-mTagBFP2 constructs were generated using

membrane localization tags reported previously (Megason, 2009; Subach et al., 2011). These

constructs were each sub-cloned into a pMTB backbone. mRNA for all experiments was synthesized

from pCS or pMTB backbones via *in vitro* transcription using the mMESSAGE mMACHINE system

(Ambion). Embryos were injected at the single cell stage using a Nanoject system set to 2.3nl of

injection volume containing 70-90pg of RNA for each mRNA injected. Injected embryos were then

screened for brightness, and damaged embryos were removed. Scube2 morpholino injections were

performed with 7ng of Scube2 MO2 and 3.5ng of p53 MO to control for phenotypic variability, while

control morpholino injections were performed using 10.5ng of p53 MO only (Gerety and Wilkinson,

2011; Woods and Talbot, 2005).

Sonidegib and Cyclopamine Treatment

Stock solution of 1 mM Sonidegib suspended in DMSO was used for treatment as generously given by

the lab of Rosalind Segal. Embryos were placed in egg water containing a concentration of 50µM for

the treatment condition, and equal parts DMSO were added to the sham control. Cyclopamine was

dissolved in 100% ethanol to make 50mM stock solution and was diluted for treatment in egg water to

100 µM. Treatment began at 7 hpf and continued until imaging at 22 hpf.

Confocal Imaging

For quantitative imaging, embryos were staged and mounted in our previously described dorsal mount

(Kimmel et al., 1995; Megason, 2009; Xiong et al., 2013) in egg water with 0.01% tricaine (Western

Chemical, Inc.). Embryos were manipulated for proper positioning with hair loops, before gently

lowering the coverslip. Embryos were not depressed by the coverslip or impinged by the mold, enabling

imaging of their normal proportions. Imaging was performed on embryos staged at 18-24 hpf, unless

otherwise noted in corresponding figure legends. Live imaging was performed using a Zeiss 710

confocal/2-photon microscope, Zen image acquisition software, and C-Apochromat 40X 1.2 NA

objective. For fluorescent protein excitation, 405 nm (BFP), 488 nm (GFP/moxNeonGreen), 514 nm

(mCitrine), 561 nm (mCherry/dsRed) and 594 nm (mCardinal) lasers were used. The imaging field was

centered in each embryo on the somite 6/7 boundary for consistent positioning between images. For

quantitative analysis, imaging datasets are only compared between sibling embryos imaged on the

same day with the same settings. This approach aims to avoid clutch effects or variability in detector

sensitivity and laser power that occur over time. Typical imaging settings with the 40x objective were as

follows: image size of 1024x1024 pixels with .21µm per pixel and an interval of 1µm in the Z direction.

For display purposes, images are rendered in cross sectional views (X-Z axis) which are then rotated

for display, with image intensities for co-injection markers adjusted evenly within datasets for

displayable brightness. FRAP, early stage embryo imaging and time-lapses were performed using a 1.0

NA 20x objective. Brightfield and widefield fluorescence images of whole embryos were obtained using

an Olympus MVX10 and a Leica MZ12.5 dissecting microscope.

FRAP Experiments and Analysis

Imaging for FRAP was performed using a 1.0 NA 20x objective at dome stage. Bleaching was

performed for two minutes in a 100μmx100μm area in the center of the frame with a 488nm Argon

laser. Imaging was performed with a low laser power to reduce bleaching, and images were obtained at

10 second intervals over five minutes to quantify recovery. FRAP data in the bleached region was then

normalized to the minimum and maximum intensity for each respective time trace. Normalized recovery

intensities were then fitted to the following exponential to determine the mobile fraction (*A*) in MATLAB:

$y = A(1 - e^{-\tau t})$ (Munjal et al., 2015). These fitted traces were also used to determine the point at which

50% maximum recovery was reached in Figure 5G.

Image Analysis

Images were analyzed using a custom MATLAB-based image analysis software that enables rapid

segmentation of neural tube imaging data. Neural imaging data is segmented by the user sequentially

from anterior to posterior. Over a set step size (usually 50 pixels), the user selects points at the base of

the floor plate cell and top of the roof plate cell that divide the neural tube into its two halves (Fig. S1A).

The user then selects the widest point of the neural tube in each image. Imaging data from mature

neurons, found laterally, and within the lumen of the neural tube, found medially, are disregarded using

a set percentage of neural width (Fig. S1B). Once these positions are recorded, imaging data is then

recovered as average pixel intensity in 25 bins from ventral to dorsal across 3-4 somites of A-P length.

This binning and averaging strategy enables comparison of data between embryos that accounts for

variations in neural tube D-V height. During the segmentation process, the researcher is blinded to the

title of the dataset which contains information about its treatment condition. For distribution plots,

binned intensities are reported for each embryo as the average intensities for each bin along the entire

AP axis of the imaging volume. Each embryo's average intensity profile is then treated as an

independent sample and averaged for displayed distribution profiles and standard deviations. To avoid

artifacts caused by rounding in the calculations of half maximum control intensity positional values are

extracted from the spline-interpolated intensity profiles for each individual dataset.

Progenitor domain segmentation is performed on average intensity profiles from each embryo in

a dataset in the following manner: first, all intensity profiles in the data set undergo background

subtraction and cross channel fluorescence caused by the extreme brightness of the *dbx1b:GFP* line is

removed from the *olig2:dsred* channel. Intensity profiles are then fed to a peak finding algorithm to

identify local maxima. Both *dbx1b+* and *nkx2.2a+* progenitor domains are found in the green channel,

so a maximum of two peaks is allowed. In the red channel, only one peak is specified to identify

olig2:dsred signal. Average peak intensity values for each domain are then calculated for the entire

control dataset, and 50% of this value in the case of the *nkx2.2a* and *dbx1b* domains is used as the

threshold for calculating domain width. Given its greater spread along the D-V axis, a threshold of 25%

of peak height is used in calculating width of the *olig2+* domain. Domain widths are then extracted from

spline-interpolated intensity profiles to avoid errors introduced by rounding to the next bin. Segmented

widths and positions of *nkx2.2a*, *olig2*, and *dbx1b* expression are then averaged for plotting purposes.

Domain plots are generated by assigning all *nkx2.2a+* progenitors to the p3 fate, *olig2+* progenitors

lacking *nkx2.2a* expression overlap to the pMN fate, and *dbx1b+* progenitors to the p0-d6 fate. These

domain sizes and positions are then used to reconstruct domains in-between or flanking them, which

include the p2-p1 domain between pMN and p0-d6, the floorplate below p3, and the d5-roofplate above

p0-d6. These heights and positions are then used to generate the stacked bar plots shown.

Statistical Analysis

Statistical comparisons of maximum average intensity and position of 50% maximum intensity

are performed by an unpaired T-test. Although each dataset contains hundreds of measurements of

each binned intensity value over the A-P axis of a z-stack, only the average of these measurements for

each embryo is treated as a data-point for calculation of the standard deviation and statistical

significance tests. This is done to avoid oversampling that would exaggerate statistical significance. In

all measurements, statistical significance is markedly increased if analysis is performed by treating all

underlying intensity measurements as samples. Thresholds for calculating the position of half maximum

are determined from the average maximum of the corresponding control dataset for each experiment.

Position is then determined from the fitted trend-line to avoid inaccuracies due to rounding. To calculate

the significance of shifts in boundary positions, upper domain boundaries for each embryo were

compared in an un-paired t-test between embryos from each population. When the progenitor domain

segmentation algorithm finds there is no domain present, the boundary is set to 0.

CRISPR Screen for Scube2 Regulators

Cas9 protein was generated and purified in lab as described (Gagnon et al., 2014). Three guide RNA

sequences targeting the first one-to-three exons of each gene were selected based on their quality

using the web-tool CHOP-CHOP and synthesized using standard methods (Gagnon et al., 2014).

Equivalent guide RNA and Cas9 protein concentrations were used in all samples for mosaic knockout.

Phenotypes were assessed at 18-20 hpf by confocal microscopy.

Acknowledgments

We thank Lisa Goodrich, Rosalind Segal, and Wolfram Goessling for their comments and helpful

discussion. The Scube2 construct was a gift from the Talbot lab.

Competing interest

The authors declare no competing or financial interests.

Funding

Z.M.C was supported in part by the program in Biological and Biomedical Sciences at Harvard

University. S.G.M. and Z.M.C. were supported by R01-GM107733 and R01-DC015478. T.Y.-C.T. was

supported by the Damon Runyon Cancer Foundation Fellowship.

References

References

Aday, A. W., Zhu, L. J., Lakshmanan, A., Wang, J. and Lawson, N. D. (2011). Identification
of cis regulatory features in the embryonic zebrafish genome through large-scale profiling of
H3K4me1 and H3K4me3 binding sites. *Dev. Biol.* **357**, 450–462.

Averbukh I, Ben-Zvi D, Mishra S, Barkai N, Averbukh, I., Ben-Zvi, D., Mishra, S. and
Barkai, N. (2014). Scaling morphogen gradients during tissue growth by a cell division rule.
Development **141**, 2150–6.

Barkai, N. and Ben-Zvi, D. (2009). "Big frog, small frog"--maintaining proportions in embryonic
development: delivered on 2 July 2008 at the 33rd FEBS Congress in Athens, Greece. *FEBS J.*
276, 1196–207.

Ben-Zvi, D. and Barkai, N. (2010). Scaling of morphogen gradients by an expansion-repression
integral feedback control. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 6924–9.

Ben-Zvi, D., Shilo, B.-Z., Fainsod, A. and Barkai, N. (2008). Scaling of the BMP activation
gradient in Xenopus embryos. *Nature* **453**, 1205–11.

Ben-Zvi, D., Pyrowolakis, G., Barkai, N. and Shilo, B.-Z. (2011). Expansion-repression
mechanism for scaling the Dpp activation gradient in Drosophila wing imaginal discs. *Curr. Biol.*
21, 1391–6.

Ben-Zvi, D., Fainsod, A., Shilo, B.-Z. and Barkai, N. (2014). Scaling of dorsal-ventral
patterning in the Xenopus laevis embryo. *Bioessays* **36**, 151–6.

Bidet, M., Joubert, O., Lacombe, B., Ciantar, M., Nehmé, R., Mollat, P., Brétillon, L.,
Faure, H., Bittman, R., Ruat, M., et al. (2011). The Hedgehog Receptor Patched Is
Involved in Cholesterol Transport. *PLoS One* **6**, e23834.

Briscoe, J. and Small, S. (2015). Morphogen rules: design principles of gradient-mediated embryo
patterning. *Development* **142**, 3996–4009.

Briscoe, J., Chen, Y., Jessell, T. M. and Struhl, G. (2001). A hedgehog-insensitive form of
patched provides evidence for direct long-range morphogen activity of sonic hedgehog in the
neural tube. *Mol. Cell* **7**, 1279–91.

Burke, R., Nellen, D., Bellotto, M., Hafen, E., Senti, K.-A. A., Dickson, B. J. and
Basler, K. (1999). Dispatched, a Novel Sterol-Sensing Domain Protein Dedicated to the Release
of Cholesterol-Modified Hedgehog from Signaling Cells. *Cell* **99**, 803–815.

Cao, Y., Ryser, M. D., Payne, S., Li, B., Rao, C. V. and You, L. (2016). Collective Space-
Sensing Coordinates Pattern Scaling in Engineered Bacteria. *Cell* **165**, 620–630.

Chamberlain, C. E., Jeong, J., Guo, C., Allen, B. L. and McMahon, A. P. (2008).
Notochord-derived Shh concentrates in close association with the apically positioned basal body in
neural target cells and forms a dynamic gradient during neural patterning. *Development* **135**,
1097–106.

Chen, M.-H., Li, Y.-J., Kawakami, T., Xu, S.-M. and Chuang, P.-T. (2004). Palmitoylation is
required for the production of a soluble multimeric Hedgehog protein complex and long-range
signaling in vertebrates. *Genes Dev.* **18**, 641–59.

Cooke, J. (1981). Scale of body pattern adjusts to available cell number in amphibian embryos.
Nature **290**, 775–778.

Costantini, L. M., Baloban, M., Markwardt, M. L., Rizzo, M., Guo, F., Verkhusha, V. V.
and Snapp, E. L. (2015). A palette of fluorescent proteins optimized for diverse cellular
environments. *Nat. Commun.* **6**, 7670.

Creanga, A., Glenn, T. D., Mann, R. K., Saunders, A. M., Talbot, W. S. and Beachy, P.
A. (2012). Scube/You activity mediates release of dually lipid-modified Hedgehog signal in soluble
form. *Genes Dev.* **26**, 1312–25.

Driesch, H. (1892). Entwicklungsmechanische Studien: I. Der Werthe der beiden ersten
Furchungszellen in der Echinogdermenentwicklung. Experimentelle Erzeugung von Theil- und
Doppelbildungen. *Zeitschrift für wissenschaftliche Zool.*

Francois, P., Vonica, A., Brivanlou, A. H. and Siggia, E. D. (2009). Scaling of BMP gradients
in Xenopus embryos. *Nature* **461**, E1–E1.

Gagnon, J. A., Valen, E., Thyme, S. B., Huang, P., Ahkmetova, L., Pauli, A., Montague,
T. G., Zimmerman, S., Richter, C., Schier, A. F., et al. (2014). Efficient Mutagenesis by
Cas9 Protein-Mediated Oligonucleotide Insertion and Large-Scale Assessment of Single-Guide
RNAs. *PLoS One* **9**, e98186.

Gerety, S. S. and Wilkinson, D. G. (2011). Morpholino artifacts provide pitfalls and reveal a novel
role for pro-apoptotic genes in hindbrain boundary development. *Dev. Biol.* **350**, 279–89.

Gribble, S. L., Nikolaus, O. B. and Dorsky, R. I. (2007). Regulation and function of Dbx genes
in the zebrafish spinal cord. *Dev. Dyn.* **236**, 3472–83.

Grimmond, S., Larder, R., Van Hateren, N., Siggers, P., Morse, S., Hacker, T., Arkell,
R. and Greenfield, A. (2001). Expression of a novel mammalian epidermal growth factor-
related gene during mouse neural development. *Mech. Dev.* **102**, 209–211.

Hamaratoglu, F., de Lachapelle, A. M., Pyrowolakis, G., Bergmann, S. and Affolter, M.
(2011). Dpp signaling activity requires Pentagone to scale with tissue size in the growing
Drosophila wing imaginal disc. *PLoS Biol.* **9**, e1001182.

Hammerschmidt, M., Bitgood, M. J. and McMahon, A. P. (1996). Protein kinase A is a
common negative regulator of Hedgehog signaling in the vertebrate embryo. *Genes Dev.* **10**, 647–
58.

Hollway, G. E., Maule, J., Gautier, P., Evans, T. M., Keenan, D. G., Lohs, C., Fischer,
D., Wicking, C. and Currie, P. D. (2006). Scube2 mediates Hedgehog signalling in the
zebrafish embryo. *Dev. Biol.* **294**, 104–18.

Inomata, H., Shibata, T., Haraguchi, T. and Sasai, Y. (2013). Scaling of dorsal-ventral
patterning by embryo size-dependent degradation of Spemann's organizer signals. *Cell* **153**,
1296–311.

Ishimatsu, K., Hiscock, T. W., Collins, Z. M., Sari, D. W. K., Lischer, K., Richmond, D.
L., Bessho, Y., Matsui, T. and Megason, S. G. (2018). Size-reduced embryos reveal a
gradient scaling-based mechanism for zebrafish somite formation. *Development* **145**, dev161257.

Jakobs, P., Exner, S., Schürmann, S., Pickhinke, U., Bandari, S., Ortmann, C., Kupich,
S., Schulz, P., Hansen, U., Seidler, D. G., et al. (2014). Scube2 enhances proteolytic Shh
processing from the surface of Shh-producing cells. *J. Cell Sci.* **127**, 1726–37.

Jakobs, P., Schulz, P., Ortmann, C., Schürmann, S., Exner, S., Rebollido-Rios, R.,
Dreier, R., Seidler, D. G. and Grobe, K. (2016). Bridging the gap: heparan sulfate and
Scube2 assemble Sonic hedgehog release complexes at the surface of producing cells. *Sci. Rep.*
6, 26435.

Jessen, J. R., Meng, A., McFarlane, R. J., Paw, B. H., Zon, L. I., Smith, G. R. and Lin,
S. (1998). Modification of bacterial artificial chromosomes through Chi-stimulated homologous
recombination and its application in zebrafish transgenesis. *Proc. Natl. Acad. Sci.* **95**, 5121–5126.

Kawakami, K. (2004). Transgenesis and Gene Trap Methods in Zebrafish by Using the Tol2
Transposable Element.pp. 201–222.

Kawakami, T., Kawcak, T., Li, Y.-J., Zhang, W., Hu, Y. and Chuang, P.-T. (2002). Mouse
dispatched mutants fail to distribute hedgehog proteins and are defective in hedgehog signaling.
Development **129**, 5753–65.

Kawakami, A., Nojima, Y., Toyoda, A., Takahoko, M., Satoh, M., Tanaka, H., Wada, H.,
Masai, I., Terasaki, H., Sakaki, Y., et al. (2005). The zebrafish-secreted matrix protein
you/scube2 is implicated in long-range regulation of hedgehog signaling. *Curr. Biol.* **15**, 480–8.

Kicheva, A., Bollenbach, T., Ribeiro, A., Valle, H. P., Lovell-Badge, R., Episkopou, V.
and Briscoe, J. (2014). Coordination of progenitor specification and growth in mouse and chick
spinal cord. *Science (80-.).* **345**, 1254927–1254927.

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. F. (1995).

Stages of embryonic development of the zebrafish. *Dev. Dyn.* **203**, 253–310.

Kinkhabwala, A., Riley, M., Koyama, M., Monen, J., Satou, C., Kimura, Y., Higashijima,
S.-I. and Fetcho, J. (2011). A structural and functional ground plan for neurons in the hindbrain
of zebrafish. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 1164–9.

Kucenas, S., Takada, N., Park, H.-C., Woodruff, E., Broadie, K. and Appel, B. (2008).
CNS-derived glia ensheath peripheral nerves and mediate motor root development. *Nat. Neurosci.*
11, 143–151.

Kwong, L., Bijlsma, M. F. and Roelink, H. (2014). Shh-mediated degradation of Hhip allows cell
autonomous and non-cell autonomous Shh signalling. *Nat. Commun.* **5**, 4849.

Liem, K. F., Jessell, T. M. and Briscoe, J. (2000). Regulation of the neural patterning activity of
sonic hedgehog by secreted BMP inhibitors expressed by notochord and somites. *Development*
127, 4855–4866.

Liu, T.-L., Upadhyayula, S., Milkie, D. E., Singh, V., Wang, K., Swinburne, I. A.,
Mosaliganti, K. R., Collins, Z. M., Hiscock, T. W., Shea, J., et al. (2018). Observing
the cell in its native state: Imaging subcellular dynamics in multicellular organisms. *Science* **360**,
eaaq1392.

McHale, P., Rappel, W.-J. and Levine, H. (2006). Embryonic pattern scaling achieved by
oppositely directed morphogen gradients. *Phys. Biol.* **3**, 107–120.

Megason, S. G. (2009). In toto imaging of embryogenesis with confocal time-lapse microscopy.
Methods Mol. Biol. **546**, 317–32.

Morgan, T. H. (1895). Half embryos and whole embryos from one of the first two blastomeres. *Anat.*
Anz. **10**, 623–638.

Munjal, A., Philippe, J.-M., Munro, E. and Lecuit, T. (2015). A self-organized biomechanical
network drives shape changes during tissue morphogenesis. *Nature* **524**, 351–355.

Pepinsky, R. B., Zeng, C., Wen, D., Rayhorn, P., Baker, D. P., Williams, K. P., Bixler,
S. A., Ambrose, C. M., Garber, E. A., Miatkowski, K., et al. (1998). Identification of a
Palmitic Acid-modified Form of Human Sonic hedgehog. *J. Biol. Chem.* **273**, 14037–14045.

Pierani, A., Brenner-Morton, S., Chiang, C. and Jessell, T. M. (1999). A sonic hedgehog-
independent, retinoid-activated pathway of neurogenesis in the ventral spinal cord. *Cell* **97**, 903–
15.

Porter, J. A., Ekker, S. C., Park, W.-J., von Kessler, D. P., Young, K. E., Chen, C.-H.,
Ma, Y., Woods, A. S., Cotter, R. J., Koonin, E. V, et al. (1996a). Hedgehog Patterning

Activity: Role of a Lipophilic Modification Mediated by the Carboxy-Terminal Autoprocessing
Domain. *Cell* **86**, 21–34.

Porter, J. A., Young, K. E. and Beachy, P. A. (1996b). Cholesterol Modification of Hedgehog
Signaling Proteins in Animal Development. *Science (80-.).* **274**, 255–259.

Roberts, B., Casillas, C., Alfaro, A. C., Jägers, C. and Roelink, H. (2016). Patched1 and
Patched2 inhibit Smoothened non-cell autonomously. *Elife* **5**,.

Shilo, B.-Z. and Barkai, N. (2017). Buffering Global Variability of Morphogen Gradients. *Dev. Cell*
40, 429–438.

Shin, J., Park, H.-C., Topczewska, J. M., Mawdsley, D. J. and Appel, B. (2003). Neural
cell fate analysis in zebrafish using olig2 BAC transgenics. *Methods Cell Sci.* **25**, 7–14.

Shkumatava, A., Fischer, S., Müller, F., Strahle, U. and Neumann, C. J. (2004). Sonic
hedgehog, secreted by amacrine cells, acts as a short-range signal to direct differentiation and
lamination in the zebrafish retina. *Development* **131**, 3849–58.

Spemann, H. (1938). *Embryonic Development and Induction*. Yale Univ.; New Haven:

Subach, O. M., Cranfill, P. J., Davidson, M. W. and Verkhusha, V. V. (2011). An Enhanced
Monomeric Blue Fluorescent Protein with the High Chemical Stability of the Chromophore. *PLoS*
One **6**, e28674.

Tukachinsky, H., Kuzmickas, R. P. P., Jao, C. Y. Y., Liu, J. and Salic, A. (2012).
Dispatched and Scube Mediate the Efficient Secretion of the Cholesterol-Modified Hedgehog
Ligand. *Cell Rep.* **2**, 308–320.

Umulis, D. M. and Othmer, H. G. (2013). Mechanisms of scaling in pattern formation.
Development **140**, 4830–43.

Uygur, A., Young, J., Huycke, T. R., Koska, M., Briscoe, J. and Tabin, C. J. (2016).
Scaling Pattern to Variations in Size during Development of the Vertebrate Neural Tube. *Dev. Cell*
37, 127–135.

van Eeden, F. J. J., Granato, M., Schach, U., Brand, M., Furutani-Seiki, M., Haffter, P.,
Hammerschmidt, M., Heisenberg, C. P. P., Jiang, Y. J. J., Kane, D. A. A., et al.
(1996). Mutations affecting somite formation and patterning in the zebrafish, Danio rerio.
Development **123**, 153–164.

Willot, V., Mathieu, J., Lu, Y., Schmid, B., Sidi, S., Yan, Y.-L., Postlethwait, J. H.,
Mullins, M., Rosa, F. and Peyriéras, N. (2002). Cooperative Action of ADMP- and BMP-
Mediated Pathways in Regulating Cell Fates in the Zebrafish Gastrula. *Dev. Biol.* **241**, 59–78.

Wolpert, L. (1969). Positional information and the spatial pattern of cellular differentiation. *J. Theor.*
Biol. **25**, 1–47.

Woods, I. G. and Talbot, W. S. (2005). The you gene encodes an EGF-CUB protein essential for
Hedgehog signaling in zebrafish. *PLoS Biol.* **3**, e66.

Xiong, F., Tentner, A. R., Huang, P., Gelas, A., Mosaliganti, K. R., Souhait, L.,
Rannou, N., Swinburne, I. a, Obholzer, N. D., Cowgill, P. D., et al. (2013). Specified
neural progenitors sort to form sharp domains after noisy Shh signaling. *Cell* **153**, 550–61.

Xu, J., Srinivas, B. P., Tay, S. Y., Mak, A., Yu, X., Lee, S. G. P., Yang, H.,
Govindarajan, K. R., Leong, B., Bourque, G., et al. (2006). Genomewide expression
profiling in the zebrafish embryo identifies target genes regulated by Hedgehog signaling during
vertebrate development. *Genetics* **174**, 735–52.

Yang, R.-B., Ng, C. K. D., Wasserman, S. M., Colman, S. D., Shenoy, S., Mehraban,
F., Komuves, L. G., Tomlinson, J. E. and Topper, J. N. (2002). Identification of a novel
family of cell-surface proteins expressed in human vascular endothelium. *J. Biol. Chem.* **277**,
46364–73.

Zagorski, M., Tabata, Y., Brandenberg, N., Lutolf, M. P., Tkačik, G., Bollenbach, T.,
Briscoe, J. and Kicheva, A. (2017). Decoding of position in the developing neural tube from
antiparallel morphogen gradients. *Science* **356**, 1379–1383.

Zeng, X., Goetz, J. A., Suber, L. M., Scott, W. J., Schreiner, C. M. and Robbins, D. J.
(2001). A freely diffusible form of Sonic hedgehog mediates long-range signalling. *Nature* **411**,
716–20.

837
838
839

Supplemental Figures:

Figure S1- Segmentation of neural imaging data.

(A) Image of a 20 hpf *tg(ptch2:kaede)* reporter embryo undergoing selection of the "axis of reflection"

which serves to mark a measurement of Dorsoventral height and separate the left and right halves of

the neural tube. These positions are picked by the user, first by picking the bottom of the floor plate cell,

then inputting the top coordinate of the roof plate cell. (B) Image of a 20 hpf *tg(ptch2:kaede)* reporter

embryo after a user has selected the width of the spinal cord. The algorithm then calculates how much

imaging data to collect based on a ratio which avoids mature neurons and the lumen of the spinal cord.

(C) After collection of average intensities in each bin, data is stored as shown. Average profiles for

generating distribution plots and segmenting domains are gathered by averaging these data along the

A-P axis for both halves of the neural tube.

Figure S2- Progenitor domain width determination and bar plot generation.

(A) Averaged image intensity profiles from the green channel of both sides of a segmented neural tube

from a *tg(dbx1b:GFP;olig2:dsred;nkx2.2a:memGFP)* embryo. Black diamonds represent peaks found

by a peak finding algorithm, while open circles and lines show the calculated domain boundaries and

width for this embryo based on the universally applied threshold in this dataset. Thresholds are

determined by 50% of average peak intensity of the control population for each dataset. (B) Example

domain determination of *nkx2.2a* and dbx1b+ cells. Red lines mark the predicted *nkx2.2a* domain,

which correlates with the boundary of their fluorescence. Green lines mark the predicted width of the

dbx1b domain which correlates well with visible fluorescence of this domain. Some anterior-posterior

variability in domain size is observed. (C) Formatted as in part A, this plots the averages *olig2:dsred+*

intensity, peak, and determined width. (D) Example domain determination of *olig2+* cells. Blue lines

mark the predicted *olig2:dsred* domain, which correlates with the boundary of their fluorescence.

Thresholds are determined by 25% of average peak intensity of the control embryos in each dataset.

(E) Example stacked bar plot generated only from this embryo using calculated domain positions to

determine domain sizes.

Figure S3- Cyclopamine treatment of *tg(scube2:moxNG)* embryos.

(A-C) Transverse view 22 hpf *tg(scube2:moxNG; mem:mCherry)* embryos treated with a DMSO control

(A) or 100µM Cyclopamine (B). (C) Quantification of mean reporter intensity of embryos as treated in A-

B. The black bar marks the position of 50% of control maximum intensity which was used for statistical

testing. These values were statistically significantly shifted ventrally in drug treated embryos relative to

control (unpaired t-test p= P=.0001145.).

Figure S4- Mosaic pax6a/b CRISPR mutants have lowered *scube2* expression.

(A-B) *tg(scube2:moxNG)* embryos imaged at 18 hpf that were injected at the single cell stage with

mem-mTagBFP2 mRNA and injected at the 8-16 cell stage with *mem-mCherry* mRNA, Cas9 protein,

and sgRNAs targeting either the tyrosinase pigment gene as a control (A) or *pax6a* and *b* (B).

888

Figure S5- Rescue of scube2 CRISPR mutants with *scube2* or *scube2-mCitrine* mRNA.

(A) Results of a *scube2* mutant in-cross. The allele was generated by mutagenesis with CRISPR using

three guides targeting *scube2* coding sequence (B) Embryos rescued by the injection of *scube2* mRNA

co-injected with *mem-mCardinal* which were screened for being mem-mCardinal positive (Liu et al.,

2018). (C) Embryos rescued by the injection of *scube2-mCitrine* mRNA co-injected with *mem-*

mCardinal which were also screened for *mem-mCardinal* fluorescence.

| AB ix injected with Exon 6 MO | Tg(*scube2:sc2-ng*) +/- ix injected with Exon6 MO |

Mutant phenotype: 85% (29/34) Mutant Phenotype: 28% (4/14)

Figure S6- Rescue of morpholino knockdown by *tg(scube2:scube2-moxNeonGreen)*

expression.

(A) Schematic of Scube2 Exon 6 targeting morpholino and expected resistance in the genome of

tg(scube2:scube2-moxNeonGreen) embryos. Splice junction targeted by the Exon 6 morpholino is

largely absent from the transgenic full length Scube2-mCitrine construct, allowing a test of rescue

capacity. (B) *tg(ptch2:kaede)* response profiles of embryos injected wither with p53 morpholino only,

the previously published Scube2 MO2 (Woods and Talbot, 2005), or Scube2 Exon 6 morpholino. (C)

Overview photo of wildtype embryos injected with Scube2 Exon6 morpholino showing the tail curling

indicative of Shh signaling phenotypes. (D) Overview photo of *tg(scube2:scube2-moxNeonGreen)*

embryos injected with Scube2 Exon6 morpholino.

Supplemental Movies:

Movie S1- *tg(scube2:moxNG; shh:mem-mCherry)* **timelapse transverse view**

Movie S2- *tg(scube2:moxNG; shh:mem-mCherry)* **timelapse maximum intensity projection**

dorsal view

Movie S3- Scube2-mCitrine FRAP imaging in a dome stage embryo

Movie S4- Sec-mCitrine FRAP imaging in a dome stage embryo

Movie S5- *tg(scube2:scube2-moxNG; shh:mem-mCherry)* **timelapse transverse view**

Movie S6- *tg(scube2:scube2-moxNG; shh:mem-mCherry)* **timelapse maximum intensity**

projection dorsal view