1

A transcriptional cycle suited to daytime N$_2$ fixation in the unicellular cyanobacterium

Candidatus *Atelocyanobacterium thalassa* (UCYN-A)

María del Carmen Muñoz-Marin[a, b], Irina N. Shilova[a#], Tuo Shi[a, c], Hanna Farnelid[a, d], Ana Maria

Cabello[a] & Jonathan P. Zehr[a*]

[a] Ocean Sciences Department, University of California, Santa Cruz, CA, 95064, USA.

[b] Departamento de Bioquímica y Biología Molecular, Universidad de Córdoba, Córdoba, 14071,

Spain.

[c] State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen, 361102,

China.

[d] Centre for Ecology and Evolution in Microbial Model Systems (EEMiS), Linnaeus University,

SE-391 82, Kalmar, Sweden.

Running Head: Diel whole genome transcription in UCYN-A

*Address correspondence to Prof. Jonathan Zehr, zehrj@ucsc.edu.

Present address: Second Genome, Inc, South San Francisco, CA, 94080, USA

27   **Abstract**

The symbiosis between a marine alga and a N_2-fixing cyanobacterium (UCYN-A) is

geographically widespread in the oceans and is important in the marine N cycle. UCYN-A is

uncultivated, and is an unusual unicellular cyanobacterium because it lacks many metabolic

functions, including oxygenic photosynthesis and carbon fixation, which are typical in

cyanobacteria. It is now presumed to be an obligate symbiont of haptophytes closely related to

Braarudosphaera bigelowii. N_2-fixing cyanobacteria use different strategies to avoid inhibition of

N_2 fixation by the oxygen evolved in photosynthesis. Most unicellular cyanobacteria temporally

separate the two incompatible activities by fixing N_2 only at night, but surprisingly UCYN-A

appears to fix N_2 during the day. The goal of this study was to determine how the unicellular

UCYN-A coordinates N_2 fixation and general metabolism compared to other marine

cyanobacteria. We found that UCYN-A has distinct daily cycles of many genes despite the fact

that it lacks two of the three circadian clock genes found in most cyanobacteria. We also found

that transcription patterns in UCYN-A are most similar to marine cyanobacteria that are capable

of aerobic N_2 fixation in the light such as *Trichodesmium* and heterocyst-forming cyanobacteria,

rather than *Crocosphaera* or *Cyanothece* species, which are more closely related to unicellular

marine cyanobacteria evolutionarily. Our findings suggest that the symbiotic interaction has

resulted in a shift of transcriptional regulation to coordinate UCYN-A metabolism with the

phototrophic eukaryotic host, thus allowing efficient coupling of N_2 fixation (by the

cyanobacterium) to the energy obtained from photosynthesis (by the eukaryotic unicellular alga)

in the light.

52   **Importance**

The symbiotic N_2-fixing cyanobacterium UCYN-A and its eukaryotic algal host, which is

closely related to *Braarudosphaera bigelowii*, have been shown to be globally distributed and

important in open ocean N_2 fixation. These unique cyanobacteria have reduced metabolic

capabilities, even lacking genes for oxygenic photosynthesis and carbon fixation. Cyanobacteria

generally use energy from photosynthesis for nitrogen fixation, but require mechanisms for

avoiding inactivation of the oxygen-sensitive nitrogenase enzyme by ambient oxygen (O_2) or the

O_2 evolved through photosynthesis. This study shows that the symbiosis between the N_2-fixing

cyanobacterium UCYN-A and its eukaryotic algal host has led to adaptation of its daily gene

expression pattern in order to enable daytime aerobic N_2 fixation, which is likely more

energetically efficient than fixing N_2 at night, as in other unicellular marine cyanobacteria.

Introduction

Nitrogen (N_2)-fixing microorganisms (diazotrophs), which reduce atmospheric N_2 to biologically available ammonium, are critical components of aquatic and terrestrial ecosystems because they supply fixed inorganic N (1). Cyanobacteria are particularly important in N_2 fixation because they can fuel the energy intensive N_2 reduction reaction using energy supplied by oxygenic photosynthesis. In the oceans, the filamentous, non-heterocyst-forming cyanobacterium *Trichodesmium* and the heterocyst-forming symbiont of diatoms (*Richelia* and related cyanobacteria) were believed to be the major N_2-fixing microorganisms until the discovery of the unicellular cyanobacteria *Crocosphaera*, *Cyanothece* and Candidatus *Atelocyanobacterium thalassa* (UCYN-A) in the open ocean. *Crocosphaera* and *Cyanothece* are free-living marine cyanobacteria, but UCYN-A is unusual in that it lacks oxygenic photosynthesis and is a symbiont of a haptophyte alga (related to *Braarudosphaera bigelowii*). The UCYN-A symbiosis is geographically widespread and is important in oceanic N_2 fixation (2-5). The UCYN-A genome has been greatly reduced, with massive metabolic streamlining including the loss of the oxygen-evolving Photosystem II (PSII), the carbon-fixing enzyme RuBisCO, and the entire tricarboxylic acid (TCA) cycle (6). UCYN-A has been shown to supply fixed N to the haptophyte in exchange for fixed carbon (4, 7), but it is not known how these two single-celled organisms coordinate metabolism and cell growth over the daily division cycle.

N_2 fixation requires energy and reductant, but the nitrogenase enzyme is inactivated by oxygen (O_2). Cyanobacteria generally have access to sufficient energy from photosynthesis but require mechanisms for avoiding inactivation of nitrogenase and N_2 fixation by ambient oxygen (O_2) or the O_2 evolved through photosynthesis. *Trichodesmium* and heterocystous cyanobacteria such as *Richelia* and *Nostoc* fix N_2 during the day, whereas the free-living unicellular *Crocosphaera* and *Cyanothece* fix N_2 at night. Interestingly, the symbiotic UCYN-A appears to fix

N_2 during the day (8-10), in contrast to most other unicellular marine N_2-fixing cyanobacteria,

such as *Crocosphaera* and *Cyanothece*.

The processes of N_2-fixation and photosynthesis in cyanobacteria are regulated daily to

increase cellular fitness and ecological competitiveness (11-13). Most cyanobacteria have

circadian rhythms (11, 14, 15) that are involved in controlling daily cycles of gene transcription

and protein synthesis by signal transduction pathways involving the circadian clock *kai* genes.

UCYN-A lacks two of the three *kai* genes (*kaiA* and *kaiB*) known in most other cyanobacteria,

whereas the non-N_2-fixing cyanobacterium *Prochlorococcus* only lacks *kaiA*. Thus, the daily

whole genome expression pattern in UCYN-A is of interest to determine if there are daily patterns

as in all other cyanobacteria compared to evolutionarily-related unicellular cyanobacteria.

We used a whole genome transcription array that targets two genetically distinct uncultivated

sub-lineages of UCYN-A (UCYN-A1 and UCYN-A2), which have similar, but genetically distinct

hosts. We compared the UCYN-A whole genome diel transcription patterns to those of

Cyanothece sp. ATCC 51142 (16) and *Crocosphaera watsonii* WH 8501 (17) (both unicellular

night-time N_2-fixers) and of *Trichodesmium erythraeum* IMS101 (a filamentous non-heterocystous

day-time N_2-fixer). We also compared expression to whole genome expression of

Prochlorococcus sp. MED4 (18) (a marine non-N_2-fixer) in order to determine how UCYN-A

gene expression compares to general cyanobacterial gene expression in a sympatric open ocean

species. We found that many genes in UCYN-A have distinct diel expression patterns and that

UCYN-A has unusual gene expression patterns in comparison to unicellular N_2-fixing

cyanobacteria that fix N_2 in the dark; however, it shares some general patterns with daytime N_2-

fixing cyanobacteria, with heterocysts of heterocyst-forming cyanobacteria and with unicellular

non-N_2-fixing cyanobacteria. Results suggest that optimal metabolism for open ocean

cyanobacteria is aligned to the light period, and that symbiosis has enabled the unicellular UCYN-

A to shift N_2 fixation to the daylight period.

Results and discussion

UCYN-A has a daily rhythm of gene transcription

UCYN-A has clear diel patterns of gene transcription, with a large fraction of genes that had

periodicity of transcript levels over the dark and light periods (27%).

About a third of the UCYN-A genome (31% genes) targeted by the array were transcribed

at detectable levels (365 of 1194 total genes in UCYN-A1 and 394 of 1244 total genes in UCYN-

A2, respectively) (Table S1). Approximately 85% of these genes have differences in transcript

levels between dark and light periods, accounting for 27% of the total genes in each strain (Table

S1 and S2). *C. watsonii*, *Cyanothece* sp. and *Trichodesmium* cultures also had a large fraction of

genes with changes in the transcript levels between dark and light periods (39% in *C. watsonii*,

20% in *Cyanothece* sp. and 34% in *Trichodesmium*) (Tables S1 and S2).

The UCYN-A transcription values ($\log_2$-transformed) ranged from 2 to 13.5 with the median

of 6.0. In both sub-lineages, genes coding for nitrogenase (*nif*), F_0F_1-ATP synthase (*atpA, atpB*),

cytochrome b_6f complex (*petB*, *petC*, *petF, petL*) and the photosynthetic gene *psaC* were the most

highly transcribed in comparison to all detected genes (Table S5). Transcript levels of the same

genes were also high for both sub-lineages in metatranscriptomes collected during the TARA

expedition in the South Atlantic Ocean (19).

The two UCYN-A sub-lineages had similar periodicity of transcript levels to each other,

despite divergence in gene sequences at the amino-acid level (average 14% genome-wide), cell

morphology (19) and genome size (Figure 1). There were four gene clusters based on the time of

day exhibiting the highest relative transcript level (Figure 1). Cluster I had the highest relative

transcript level during the day (with a maximum 10 h into the light period) and included genes

involved in cell division (e.g. *ftsZ, murG, minE, murB*), DNA replication (e.g. *topA, rpoE,*

DPO3B), ABC transporters (e.g. *nikA, nikB, pstC, cbiO*), carbohydrate and lipid metabolism (e.g.

pdhA, pgi, fabG, fabH) and a few photosynthesis genes (*petL, psaD and ccsB*). The transcripts for

the *petL* gene, encoding subunit 6 of the cytochrome b_6f complex and the only nitrogen fixation-

related gene in this cluster (*nifK*) had a substantial change at this time (more than 3-fold).

140   The transcript abundance of genes from clusters II and III had similar patterns, with an

increase before sunrise and a decrease during the dark period. The highest relative transcript levels

for clusters II and III were 4h and 1h after sunrise, respectively, and included genes involved in

nitrogen fixation (*nifHDK* operon) that increased 4-fold during the light period. However, these

clusters also included genes involved in oxidative phosphorylation (e. g., NADH dehydrogenases

subunits and ATP synthase related genes), carbohydrate catabolism such as those involved in

glycolysis (e.g. *gap1, fbaA, pgi, eno*), the pentose phosphate pathway (*opcA* and *zwf*) and

photosynthesis (e.g. cytochrome b_6f complex subunit genes). In most cyanobacteria, genes

encoding proteins involved in carbohydrate catabolism are highly transcribed during the night and

are essential for survival under dark conditions.

150   The gene with the most dramatic difference in transcript levels between the light and dark

151 periods encoded the membrane protein COP23 (23 kDa circadian oscillating protein), which had

152 more than a 5-fold change in transcript abundance in both UCYN-A strains (Figure 1). COP23, a

153 protein which may have a critical role in membrane function, has only been detected in nitrogen-

154 fixing cyanobacteria (20).

155   Cluster IV had genes with the highest transcript level during the night and lowest during the

day and included genes encoding photosystem I (PSI) subunits, a carbohydrate porin (*oprB*) and

also genes encoding ribosomal proteins with 2- and 4-fold changes during the night period. Cluster

IV had the lowest number of genes compared with the other clusters. Surprisingly, the PSI genes

(*psaA* and *psaB)* were expressed during the night as in many anoxygenic phototrophic bacteria

(21), whereas in most oxygenic cyanobacteria (including mats) these genes are expressed during

the day (22).

The results show that UCYN-A has a daily rhythm of gene expression with strong

periodicities of transcript levels over the diel cycle. Daily patterns of gene transcription in

cyanobacteria are typically regulated by a circadian rhythm mediated by *kai* gene products (11).

Rhythmic daily transcription patterns are still possible without the full suite of *kai* genes, for

example, the marine cyanobacterium *Prochlorococcus* sp. MED4 lacks one of the circadian genes,

kaiA, yet it maintains strong diel gene transcription patterns (18). However, *Prochlorococcus* sp.

PCC 9511 loses the typical periodicities of the circadian clock under continuous light (23). In the

case of UCYN-A, it lacks two of the three *kai* genes (24), which is unique among cyanobacteria,

and furthermore, the *kaiC* gene was not transcribed at detectable levels. It is unclear what controls

the UCYN-A diel gene expression pattern, but it could be that 1) there are unidentified

components of a clock and signal transduction pathway, or that 2) the pattern could be driven by

the physiological differences between light and dark conditions, which might be primarily driven

by energy supplied by the eukaryotic partner. It is possible that the diel transcription patterns in

UCYN-A are primarily regulated by the daily host metabolism, which itself is likely to be

circadian. However, it is not yet known whether the UCYN-A diel cycle is maintained under

constant conditions in UCYN-A, or whether the diel pattern is maintained in the absence of the

partner alga.

UCYN-A transcription patterns are similar to aerobic marine daytime N_2-fixers and

non- N_2-fixers

UCYN-A had diel whole genome expression patterns that were different from those of

phylogenetically closely related unicellular cyanobacteria (17). Only a few genes (such as those

encoding ATP synthase) had the same daily pattern among all cyanobacteria, presumably differing

because of physiology (e.g. N_2-fixing or not). The unicellular cyanobacteria *C. watsonii* WH 8501

and *Cyanothece* sp. ATCC 51142, which fix N_2 during the night, expressed many genes in an

opposite pattern compared to the day-time N_2-fixing *T. erythraeum* and UCYN-A (Figure 2 and

Tables S4 and S5). Interestingly, the diel transcription patterns of N_2 fixation and PSI genes in

UCYN-A were opposite to those in *Cyanothece* sp. ATCC 51142 and *C. watsonii* WH 8501 and

more similar to those of *T. erythraeum* (Figure 2 and Tables S4 and S5).

As observed for the activity of nitrogenase, it has been demonstrated that levels

of *nif* transcripts and the biosynthesis of different components of the nitrogenase complex are very

sensitive to O_2 (22, 25-27), most likely to avoid energy losses associated with the degradation of

this enzyme by O_2. Thus, the different patterns observed in the genes involved in N_2-fixation in the

cyanobacteria studied here presumably are due to the different mechanisms used to protect the

nitrogenase complex from the O_2 produced by photosynthesis. *T. erythraeum* and UCYN-A had

the maximum transcript levels of the nitrogenase and PSI genes just prior to dawn, but maintained

high levels of transcripts for both sets of genes during the day. The peak of transcript levels just

before dawn is likely due to the advantage of synthesizing nitrogenase in preparation for N_2

fixation in the early hours of the day (28).

The diel expression patterns of genes that are unrelated to N_2 fixation in the aerobic day-time

N_2-fixers (*T. erythraeum* and UCYN-A) were also more similar to those of non-N_2-fixing

sympatric cyanobacteria of the genus *Prochlorococcus* and to heterocysts of heterocyst-forming

cyanobacteria than to the nighttime N_2-fixing cyanobacteria (*C. watsonii* and *Cyanothece* sp.). The

transcript levels of genes encoding ribosomal proteins in both UCYN-A and *T. erythraeum* were

higher during the night, probably because the reduced nitrogen required for the synthesis of new

proteins was obtained during the day (Figure 2 and Tables S4 and S5). Similar patterns were

observed in *Prochlorococcus* with higher transcript levels during the night (Figure 2 and Tables S4

and S5) while genes encoding ribosomal proteins in *C. watsonii* WH 8501 and *Cyanothece* sp.

ATCC 41142 had maximum transcript levels during the day (Figure 2 and Tables S4 and S5).

Intriguingly, these results imply that both UCYN-A and *T. erythraeum* have adopted day-time

gene transcription patterns for the main metabolic pathways minimizing cellular processes in the

dark. The night-time patterns of the transcript levels of the ribosomal proteins (genes) would make

it possible to have proteins synthesized in order to make the most efficient use of the light period,

as in *Prochlorococcus*. Because UCYN-A and *Trichodesmium* are likely to be the two most

abundant N_2-fixing cyanobacteria in the open ocean, it appears that direct coupling of N_2 fixation

to photosynthesis is important in the oligotrophic environment (as long as low oxygen

concentrations are maintained in the cell).

Phosphorus is a vital element for cellular energetics and growth and is acquired by oceanic

bacterioplankton primarily as phosphate (29-31). The UCYN-A phosphate ABC transporter had

the same diel pattern as in *Trichodesmium* for genes involved in DNA replication, with higher

transcript levels during the day (Table S5), but maximum transcript abundances during the late

afternoon in *Crocosphaera* and *Cyanothece* (17, 32). High levels of phosphate transporters during

the day could meet the increased demand for inorganic phosphate (33, 34) during DNA

replication, which occurs during the day in UCYN-A and *Trichosdesmium*. Similar patterns were

observed in the heterocyst-forming *Richelia* with peak expression of P acquisition genes at

approximately 15:00, suggesting the apparent rhythmicity of P acquisition could be a common

feature of daytime N_2-fixers (35).

The initiation factor of DNA replication, DnaA, is a protein highly conserved in prokaryotes

although it is absent in red algae, the cyanobacterial symbiont *Nostoc azollae* (36) and also the

spheroid bodies of diatoms (37). The genome of UCYN-A lacks the *dnaA* gene as well. Recent

studies suggested that DnaA is not essential for DNA replication and the lack of *dnaA* could

suggest a preadaptation of the genome to enable the symbiosis (38). In UCYN-A and *T.*

erythraeum, genes for DNA replication (*dnaE and* RNaseHI), DNA topoisomerases, DNA gyrases

and cell division (*ftsZ*, *mre*, *min*) had maximum transcript levels during the day (i.e., after

midday), and minimum levels at night (Figure 3A and Figure S1). In contrast, the nighttime N_2-

fixing *Cyanothece* sp. ATCC 51142 and *C. watsonii* WH 8501 confine cell division to the period

of transition from dark to light at sunrise. The temporal delay in cell division in *Cyanothece* and

Crocosphaera has been suggested to reflect the need to recover energy reserves with light-derived

energy after night-time metabolic activity (39). The similarity of the pattern in UCYN-A to

Trichodesmium is consistent with UCYN-A shifting metabolism to the daytime.

Microscopy counts of the *B. bigelowii* -UCYN-A2 symbiosis were performed eight times

during two diel cycles in order to observe the timing of cell division (Figures 3B and C and Table

S6). In both diel cycles, single host cells with two associated UCYN-A2 cells (or groups of cells),

corresponding to approximately 60% of total cell counts, were present at night between 21:00 and

03:00. The delay observed between the higher transcription levels after midday and actual cell

division at 21:00 may be explained by the need of the cell to coordinate the assembly of the cell

division machinery prior to cell division.

Unique UCYN-A transcription patterns

Although many gene transcription patterns in UCYN-A are more similar to

Trichodesmium than to other unicellular N_2-fixing cyanobacteria, some of the patterns were

unique to UCYN-A. Such unique gene transcription patterns in the UCYN-A symbiosis may

provide clues to possible roles of specific genes involved in adaptation to N_2-fixing symbiosis

revealing metabolic interdependence between host and symbiont. In order to compare the

transcriptomic patterns of these specific genes with the rest of the N_2-fixers, we performed

network analysis of these genes using Pearson correlation. Whereas most of the key genes of

the major pathways in UCYN-A had higher transcript levels during the day, the other

unicellular N_2-fixing cyanobacteria had maximum transcript levels at night (Figure 4). For

example, glycolysis genes in UCYN-A had the highest levels of transcripts at sunrise and

midday (maximum light conditions) in contrast to the other cyanobacteria (Figures 4 and 5).

The metabolic pathway that generates reductant for biosynthesis activities (NADPH), the

pentose phosphate pathway (PPP), had similar patterns. The allosteric effector *opcA*, which

redirects carbon flow to the first enzyme of the PPP (glucose-6-P dehydrogenase (*zwf*)) (18,

40), had a different periodic transcript level pattern in UCYN-A (Figures 4 and 6) compared to

other cyanobacteria (41, 42).

N_2 fixation in UCYN-A depends on the light period for the supply of photosynthate from

the host during the day, as well as possibly producing ATP by cyclic photophosphorylation with

PSI. Because UCYN-A cannot fix carbon dioxide, it has to obtain reduced carbon compounds

in the same way. Based on genome and transcriptomic profiles, we propose a pathway of

carbon metabolism for the regeneration of reductant and ATP in UCYN-A, which is needed for

N_2 fixation (Figure 7). Carbohydrate porins or ABC transporters could transport the

carbohydrates from the host to the cyanobacteria during the day and the carbon compounds

metabolized through the oxidative pentose phosphate (OPP) or glycolysis pathways. Pyruvate is

required for generation of reductant for nitrogenase and also to generate acetyl-CoA for

synthesis of fatty acids.

Because UCYN-A lacks photosystem II, which normally supplies electrons to

photosystem I by splitting water, UCYN-A needs alternative electron donors if it uses PSI to

make the reductant NADPH. The NADH generated by the OPP pathway or by glycolysis could

reduce the plastoquinone pool via the NDH-1 complex and transfer electrons to ferredoxin

though the PQ pool, cytochrome b_6f plastocyanin and the action of PSI. Ferredoxin could

deliver electrons to the ferredoxin:NADPH oxidoreductase (FNR), which might provide

reductant and ATP directly to the dinitrogenase reductase. To increase the ATP/e$^-$ ratio,

UCYN-A can redirect electrons from PSI to NDH-1 in cyclic phosphorylation. This mechanism

to supply nitrogenase with electrons was proposed years ago for heterocysts (43).

Together, the results are consistent with the assumption that UCYN-A uses host-supplied

carbohydrates during the day while other unicellular cyanobacteria synthesize their own

carbohydrates during the day and use them during the evening or at night. The unique distribution

of these metabolic processes suggests that UCYN-A has developed the ability for light-driven,

day-time N_2 fixation under oxic conditions as a result of symbiosis.

Apart from fixed carbon, several other compounds may be made available to UCYN-A,

which may be endosymbiotic, and relies on the host for all of its essential nutrients.

Interestingly, UCYN-A has the whole pathway for the synthesis of the cyanobacterial type of

vitamin B_{12}, pseudocobalamin, that can be required for the activity of several vital enzymes in

central metabolism (44) (Table S8). Transcription of genes involved in B_{12} synthesis were

detected in all cyanobacteria, and some of them had diel patterns (Table S2 and S8). It is

unknown if UCYN-A has enzymes that require pseudocobalamin or whether it can be used by

the host. However, in order for the host to use pseudocobalamin, it would have to be remodeled

in order to be accessible to the haptophyte (45). The role of pseudo-B_{12} biosynthesis in UCYN-

A is unclear, but the fact that UCYN-A retains this entire pathway, in such a reduced genome,

indicates that it is likely to have an important role, perhaps in symbiosis.

It is still unclear how N_2 fixation in UCYN-A avoids the oxygen evolved by the

photosynthetic host alga. There are only two possible pathways for consuming O_2 in UCYN-A,

including aerobic (cytochrome-dependent) respiration and the photocatalyzed reduction of O_2 to

H_2O in PSI which occurs in the heterocysts of cyanobacteria like *Nostoc* sp. PCC 7120 (46-48).

The latter, called the Mehler reaction, results in the production of the superoxide radical O_2^-,

which is subsequently reduced to water (49, 50).

In UCYN-A, the cytochrome *c* oxidase *coxA* gene was transcribed during the night (cluster

IV) but also rarely during the day, along with a few N_2 fixation genes (cluster I) (Figure 1).

Moreover, we also found higher transcript levels during the day for the antioxidant enzyme

superoxide dismutase (*sod1*) and two peroxiredoxins (*prxR*), which have the ability to detoxify

peroxide (Figures 1 and 7). Both antioxidants would protect the nitrogenase against the reactive

oxygen species produced by UCYN-A or the haptophyte host (Figure 7).

It is not currently possible to directly determine the oxygen protection mechanisms in this

uncultured microorganism because 1) transcription cannot necessarily be related to function and 2)

it is not possible to do physiological experiments with this low-abundance microorganism that has

yet to be obtained in an axenic culture. Consequently, the question of protection from O_2 cannot be

directly addressed experimentally, but our results suggest that some of the proteins in UCYN-A

could help to protect nitrogenase from the O_2 generated by host photosynthesis.

Because UCYN-A has genome reduction normally associated with endosymbiosis (e.g. in

Paulinella chromatophora; (51)), the unique gene transcription patterns of UCYN-A may provide

insights into the evolution of endosymbiosis and organellar evolution. Future studies are needed to

determine if the rhythm of these patterns is maintained under constant conditions as in a circadian

rhythm, whether the host has a circadian rhythm and/or the daily cycle in UCYN-A simply

responds to metabolite availability from the host. It will also be interesting to determine how PSI

is involved in supporting the energy or reductant requirements of N_2 fixation. Such experiments

will have to await the establishment of a pure culture.

Materials and Methods

Diel sampling of UCYN-A

Surface seawater samples for UCYN-A transcription and catalyzed reported deposition-

fluorescence *in situ* hybridization (CARD-FISH) analyses were collected using a bucket from the

end of the Scripps Institution of Oceanography (SIO) Ellen Browning Scripps Memorial Pier in La

Jolla, CA, USA. Two replicates were collected from the bucket at each time point within 48 hours

between 28[th] July and 1[st] August 2014 for transcriptomic analysis and between 3[rd] - 8[th] May 2016

for CARD-FISH. A total of 16 samples were collected every 3-6 hours (two replicates taken at

each of eight time points): 12:00-L6, 15:00-L9, 21:00-D3, 00:00-D6, 06:00-2D12, 09:00-2L3,

15:00-2L9 and 18:00-2L12. L and D stand for light and dark period, respectively, 2L and 2D the

second light-dark cycle, and the number the corresponding hours entering light or dark period.

For the CARD-FISH assay, from each seawater replicate, 190 mL of seawater was fixed with

10 mL 37% formaldehyde (1.87% v/v final concentration) at 4ºC in the dark for 1 hour. After

fixation, 100 mL was filtered at a maximum vacuum pressure of 100 mm Hg onto a 0.6 μm pore-

size, 25 mm diameter polycarbonate membrane filter (Millipore Isopore^{TM}, EMD Millipore,

Billerica, MA, USA) with a support filter of 0.8 μm pore-size, 25 mm diameter polycarbonate

cellulose acetate membrane filter (Sterlitech Corporation, Kent, WA, USA). The filters were kept

at -80ºC until processed.

Samples for RNA extraction were collected by filtering a total of 500 mL from each seawater

replicate through 0.22 μm pore-size, 47 mm diameter Supor filters (Pall Corporation, Port

Washington, NY, USA) using a peristaltic pump. Filters were placed in sterile 2 mL bead-beating

tubes with sterile glass beads, flash-frozen in liquid nitrogen and stored at -80ºC until extraction.

Double CARD-FISH assay

The double CARD-FISH assay was carried out following the protocol designed by Cabello

et al. 2016 and Cornejo-Castillo et al. 2016. All of the probes, competitors and helpers used in this

work are compiled in Table S7. More details are described in Supplementary Information.

Microscopic evaluation and counting was performed with the Carl Zeiss Axioplan-2 Imaging

Fluorescent Microscope (Zeiss, Berlin, Germany) in 3 transects (8.0 x 0.1 mm^2 each) across the

filter piece. Cell dimensions were estimated using AxioVision 4.8 and Image J software (52).

Diel sampling of Trichodesmium erythraeum IMS101 cultures

Biological triplicate cultures of *T. erythraeum* were grown in rectangular canted neck

polycarbonate cell culture flasks with a 0.2 μm pore-size vent cap and 225 cm^2 surface area

(Corning Inc., Corning, NY, USA). The cultures were maintained at 26ºC on a 12h:12h light:dark

cycle at 50 μmol quanta m^{-2} s^{-1} in YBCII media (53) supplemented with 2.8 μmol L^{-1} ferric

ammonium citrate. The light was set on at 7:00 and off at 19:00 hours. The cultures were 10-fold

diluted from the inoculum and were verified to be axenic by staining with DAPI and visualizing

cells under an epifluorescence microscope (Carl Zeiss, Thornwood, NY, USA). Growth and cell

density were monitored until the cultures reached exponential phase (~10-14 days after

inoculation), during which the cells were harvested for the diel transcription assay. Samples were

taken at 3 hours intervals starting at the onset of the light period until the end of the dark period for

a total of 24 hours. A total of 27 samples were collected from these nine time points: 7:00-D12,

10:00-L3, 13:00-L6, 16:00-L9, 19:00-L12, 22:00-D3, 1:00-D6, 4:00-D9 and 7:00-2D12, where L

and D stand for light and dark period, respectively, 2D the second light-dark cycle, and the number

the corresponding hours entering light or dark period. At each time point, 200 mL each of

triplicate cultures (replicates from different flasks) was filtered onto a 5 μm pore-size, 47 mm

diameter polycarbonate membrane filter (Osmonics, Minnetonka, MN, USA). The filters were

immediately frozen in liquid nitrogen and stored at -80º C until processing.

RNA extraction and processing for hybridization to the microarray

Environmental RNA containing transcripts from UCYN-A cells was extracted using the

Ambion RiboPure Bacteria kit (Ambion[®], ThermoFisher), with modifications that included

mechanical lysis using glass beads (Biospec, Bartlesville, OK). The extracted RNA was treated

with Turbo-DNA-free[TM] DNase Kit (Ambion[®], ThermoFisher) to remove genomic DNA.

Sufficient environmental RNA was obtained for two replicates at 4 sampling times (L6, L9, D3

and 2L12): L6-1, L6-2, L9-1, L9-2, D3-1, D3-2, 2L12-1 and 2L12-2. L and D stand for light and

dark period, respectively, 2L and 2D the second light-dark cycle, and the number the

corresponding hours entering light or dark period.

Total RNA for *T. erythraeum* was extracted using the Ambion RiboPure Bacteria kit

(Ambion®, ThermoFisher), followed by in solution DNase digestion with the RNase-free DNase

kit and on-column cleanup with the RNeasy MiniElute kit (Qiagen, Valencia, CA, USA).

RNA purity, concentration and quality were determined using a NanoDrop 1000 (Thermo

Scientific, Waltham, MA, USA) and a 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA,

USA) using the RNA 6000 Nano kit (Agilent Technologies). Only samples with RNA Integrity

Number >7.0 and ratios of A260/A230 and A260/A280 ≥1.8 were processed further.

From environmental RNA samples that contained UCYN-A, double-stranded (ds) cDNA

was synthesized and amplified following the procedure described in Shilova et al..(54). Briefly,

400 ng RNA from each sample was used, and 1 μL of 1:100 dilution (corresponding to 4.7

attomoles of ERCC-0016) of the (External RNA Control Consortium, (55)) RNA spike-in mix 1

(Ambion®) was added before amplification to monitor the technical performance of the assay (55).

Double-stranded cDNA was synthesized and amplified using the TransPlex Whole

Transcriptome Amplification kit (WTA-2, Sigma-Aldrich, St Louis, MO, USA) and antibody-

inactivated hot-start Taq DNA Polymerase (Sigma-Aldrich). The amplified cDNA was purified

with the GenElute PCR cleanup kit (Sigma-Aldrich), and the quality and quantity of ds-cDNA was

determined with NanoDrop 1000 and a 2100 Bioanalyzer using the Agilent DNA 7500 kit

(Agilent Technologies). Four hundred ng of total RNA yielded on average 12 μg of ds-cDNA. The

labeling and hybridization of cDNA samples (1.0 μg of ds-cDNA) to the microarray was done at

Roy J. Carver Center for Genomics (CCG) Facility (University of Iowa, Iowa city, Iowa, USA)

according to the Agilent Technology for arrays protocol.

For *T. erythraeum,* at least 30 μg of unamplified total RNA with a concentration of 1.0 μg

μL^{-1} per sample was provided for 27 samples. A control sample was generated by mixing equal

amount of total RNA, based on NanoDrop measured concentration, from each of the 27 samples

resulting in 28 samples in total. Reverse transcription of the total RNA, labeling of cDNA, and

hybridization to the array were performed at the Roche NimbleGen facility according to the

manufacturer's protocol (Roche NimbleGen, Inc., Madison, WI, USA).

Design of the UCYN-A array

The oligonucleotide expression array of UCYN-A was designed using UCYN-A1 and

UCYN-A2 genes using eArray web-based tool (Agilent Technology Inc.;

https://earray.chem.agilent.com/earray/) similar to the array design described in Shilova et al.(54).

The gene sequences were obtained from the National Center of Biotechnology Information (NCBI,

http://www.ncbi.nlm.nih.gov). Briefly, six probes of 60 nucleotides (nt) length were designed for

each gene, and a total of 6618 probes (1199 genes) and 6862 probes (1246 genes) were designed

for UCYN-A1 and UCYN-A2, respectively. These probes were replicated (4 times in the 8x60K

array slides and 13 times in the 4x180K array slide) which allowed internal evaluation of signals.

The sequences of all oligonucleotide probes were tested *in silico* for possible cross-hybridization

as described below. The probe sequences were used as queries in the BLASTN against the

available nt databases in June 2014: Marine microbes, Microbial Eukaryote Transcription and

Non-redundant Nucleotides in the Community Cyberinfrastructure for Advanced Microbial

Ecology Research and Analysis (CAMERA, http://camera.calit2.net/,(56)). Agilent technology

allows 5% nt mismatch in the whole probe region, thus sequences with a range of 95–100% nt

identity to the target probe are detected. Therefore, all probes with BLASTN hits with ≥95% over

100% nt length were deleted. Next, probe sequences that passed the cross-hybridization filter,

were clustered using CD-HIT-EST(57, 58) at 95% nt similarity to select unique probes for UCYN-

A1 and unique probes for UCYN-A2. Finally, to select probes specific for each strain, the probes

with ≥95% nt identity to the genes in the other strain were deleted. However, a few probes that

showed cross-hybridization between both strains for highly conserved genes (such as the

nitrogenase gene, *nifH*) were retained. In summary, 6120 probes for 1194 genes of UCYN-A1 and

6324 probes for 1244 genes of UCYN-A2 were chosen.

In addition, standard control probes as part of the Agilent Technology Array (IS-62976-8-

V2_60Kby8_GX_EQC_201000210 with ERCC control probes added) were included randomly to

feature locations on the microarray slide. The final design of the microarray was synthesized on

two platforms: ca. 62976 experimental and 1319 control probes on the 8x60K array slide and ca.

180880 experimental and 4854 control probes on the 4x180K array slide. The probe sequences are

available at NCBI Gene Expression Omnibus (GEO) under accession number GSE100124.

Design of the T. erythraeum IMS101 array

A custom oligonucleotide array for *T. erythraeum* was designed using the Roche NimbleGen

platform: (NimbleGen design ID: 080610_Trich_erth_UCSC_TS_expr) according to the complete

genome assembly of *T. erythraeum* IMS101 (NC_008312). The genome sequence is publically

available via gateways including GenBank (https://www.ncbi.nlm.nih.gov/nuccore/NC_0083120),

IMG

(http://img.jgi.doe.gov:80/cgibin/pub/main.cgi?section=TaxonDetail&page=taxonDetail&taxon_oi

d=637000329),   and   UCSC   genome   browser   (http://microbes.ucsc.edu/cgi-

bin/hgGateway?db=tricEryt_IMS101). Up to six 60-nt long tiling probes were designed to

target each of the 4788 genes in the genome, resulting in a total of 28235 probes. The probes were

duplicated on the array to allow internal evaluation of hybridization signals. Moreover, tiling 60 nt

oligonucleotide probes were also designed to target the intergenic regions >60 bp in length at a

150 bp interval, leading to a total of 11175 probes targeting 3877 intergenic regions (average 2.9

probes per intergenic region), however hybridization data for intergenic probes are not presented

here. All the probes were rank ordered and selected based on the following criteria: 1) they must

have a minimum annealing temperature of 68ºC; 2) there is no cross contamination among the

probes for different genes and for different intergenic regions. In addition to the experimental

probes, standard control probes were also included on the microarray for quality assessment of the

sample preparation, the hybridization process and the intensity measurements. The final

microarray slides were printed in 4-plex (4x72K) format with 67645 experimental probe features

and 7454 control probe features on one array. The full microarray platform descriptions and data

for *T. erythraeum* are available at NCBI GEO under accession number GSE99896. Microarray

hybridization signals were quantified using a GenePix 4000B Scanner (Molecular Devices,

Sunnyvale, CA, USA) at the Roche NimbleGen facility.

Microarray data analysis

All data analyses were performed with R (www. R-project.org) and the Bioconductor

Project(59), specifically using the Biobase(60), Linear Models for Microarray LIMMA (61),

arrayQualityMetrics(62), affyPLM(63, 64), and genefilter packages.

1) UCYN-A microarray

Transcription values for each gene were obtained using median polish summarization, and

values were normalized using quantile normalization (63, 64) (Figure S2). The transcription values

for UCYN-A at L6, L9, D3 and 2L12 are the mean transcription of the two replicates (L6-1, L6-2,

L9-1, L9-2, D3-1, D3-2, 2L12-1 and 2L12-2). Raw and normalized microarray data for UCYN-A

were submitted to NCBI GEO under accession number GSE100124. To determine if transcription

of a gene was detected, the signal-to-noise ratio (SNR) of each chip was calculated as: $SNR = (S_i-$

$BG)/BG$; where S_i is the hybridization signal for the gene and BG is the chip background signal

determined as average of the lowest 5% of all signals. Transcription was considered detected if

SNR of a transcript was ≥ 5 (as in (Shilova et al. 2014) Transcription values were centered and

scaled across genes and samples, and a distance matrix was calculated using Pearson's correlation

coefficient. The distance matrix was then used in hierarchical clustering by a complete

agglomeration method to identify clusters of genes with similar patterns of transcription during the

diel transcription.

2) T. erythraeum microarray

The raw microarray data for *T. erythraeum* were subjected to robust multichip average

(RMA) analysis (65) and quantile normalization (63, 64) (Figure 3S). Transcription values for

each gene were obtained using median polish summarization (54). Final transcription value for

each sample was a mean of up to twelve technical replicates (Blocks 1 and 2 with up to six

replicate probes in each block in the *T. erythraeum* microarray design). A gene was selected for

further analysis if it had $\log_2$ transcription above 64 in at least 25% of samples and an interquartile

range across all samples on the $\log_2$ scale of at least 0.5. This filtering resulted in 4128 genes,

which were used in further analysis.

3) Comparison of diel transcription patterns for all cyanobacteria

Transcription data for *Prochlorococcus* sp. MED4, *Cyanothece* sp. ATCC 51142 and

Crocosphaera watsonii WH 8501 was collected from previous published data (16-18).

Cyanothece sp. ATCC 51142 and *C. watsonii* WH 8501 microarray data were downloaded

from ArrayExpress (http://www.ebi.ac.uk/aerep/) using accession no. E-TABM-386 and E-

TABM-737, respectively. The genes with periodic transcriptional patterns for all studied

cyanobacteria (*Prochlorococcus* sp. MED4, *Cyanothece* sp. ATCC 51142, *C. watsonii* WH

8501, *T. erythraeum* and UCYN-A) were identified using the R package "cycle" based on

Fourier analysis, and the genes with FDR<0.25 were selected for further comparison (66)

(Table S2). To compare the diel transcription patterns among the cyanobacteria, gene

transcription values for each cyanobacterium were selected for over 36 hours. Eight points

were selected for UCYN-A (L6, L9, D3, D6, 2D12, 2L3, 2L9, 2L12), 9 points for *T.*

erythraeum (D12, L3, L6, L9, L12, D3, D6, D9, 2D12), 6 points for *Cyanothece* sp. ATCC

51142 (L2, L6, L10, D2, D6, D10), 8 points for *C. watsonii* WH 8501 (D11, L1, L6, L11, D1,

D6, 2D11, 2L1) and 19 points for *Prochlorococcus* sp. MED4 (D12 - 2L12 every 2 hours). L

and D stand for light and dark period, respectively, 2L and 2D the second light-dark cycle, and

the number the corresponding hours entering light or dark period. Because the studies had a

few dissimilar sampling times, the missing values were interpolated using the Stineman

algorithm implemented in the *imputeTS* package (67). A network was constructed based on the

Pearson correlation and using 'make_network' function in phyloseq (68). The maximum

distance between connecting nodes was selected as 0.5 unless otherwise noted in figure

legends.

Acknowledgments

We thank J.C. Meeks (University of California, Davis) for discussions, F. Azam (Scripps

Institution of Oceanography, UC San Diego) for access to Scripps facilities, and K. Turk-Kubo

and M. Hogan for lab and field support. J.M. García-Fernández and J. Díez-Dapena (University of

Córdoba, Spain) and Marine Landa (University of Santa Cruz, CA, USA) for helping us to

improve the manuscript. Microarray data have been deposited at NCBI Gene Expression Omnibus

(GEO) under accession numbers GSE100124 and GSE99896. The following secure token has

been created to allow review of record GSE100124 while it remains in private status:

ufchawekhzmlpaj.

Author contributions

M.M.M. designed the UCYN-A array, designed and performed the research and analyzed the data.

I.N.S. analyzed the *T. erythraeum* array data, aided with the design of the UCYN-A array and

comparison of transcription among cyanobacteria. T.S. designed the *T. erythraeum* array and

performed the diel sampling of *T. erythraeum* cultures. H.F. aided sampling diel UCYN-A

samples and performed the phylogenetic tree. A.M.C carried out and counted the CARD-FISH diel

samples. J.P.Z. conceptualized the study, and M.M.M., I.N.S., T.S., H.F. and J.P.Z. drafted and

edited the manuscript and figures. All authors read and approved the final manuscript.

Competing financial interests

References

1. Postgate JR. 1982. The fundamentals of nitrogen fixation. Cambridge University Press,
London.

2. Montoya JP, Holl CM, Zehr JP, Hansen A, Villareal TA, Capone DG. 2004. High rates
of N_2 fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean. Nature
430:1027-1031.

3. Cabello AM, Cornejo-Castillo FM, Raho N, Blasco D, Vidal M, Audic S, de Vargas C,
Latasa M, Acinas SG, Massana R. 2015. Global distribution and vertical patterns of a
prymnesiophyte-cyanobacteria obligate symbiosis. ISME J 10:693-706.

4. Martínez-Pérez C, Mohr W, Löscher C, Dekaezemacker J, Littmann S, Yilmaz P,
Lehnen N, Fuchs BM, Lavik G, Schmitz R, LaRoche J, Kuypers MM. 2016. Small
unicellular diazotrophic symbiont is a key player in the marine nitrogen cycle. Nature
Microbiology doi:DOI: 10.1038/NMICROBIOL.2016.163.

5. Farnelid H, Turk-Kubo K, Muñoz-Marin M, Zehr J. 2016. New insights into the
ecology of the globally significant uncultured nitrogen-fixing symbiont UCYN-A.
Aquatic Microbial Ecology 77:125-138.

6. Tripp HJ, Bench SR, Turk KA, Foster R, Desany BA, Niazi F, Affourtit JP, Zehr JP.
2010. Metabolic streamlining in an open-ocean nitrogen-fixing cyanobacterium. Nature
464.

7. Thompson AW, Foster RA, Krupke A, Carter BJ, Musat N, Vaulot D, Kuypers MMM,
Zehr JP. 2012. Unicellular cyanobacterium symbiotic with a single-celled eukaryotic
alga. Science 337:1546-1550.

8. Church M, Short C, Jenkins B, Karl D, Zehr J. 2005. Temporal patterns of nitrogenase
gene (*nifH*) expression in the oligotrophic North Pacific Ocean. Applied and
Environmental Microbiology.

9. Krupke A, Musat N, LaRoche J, Mohr W, Fuchs BM, Amann RI, Kuypers MMM,
Foster RA. 2013. In situ identification and N_2 and C fixation rates of uncultivated
cyanobacteria populations. Systematic and Applied Microbiology 36:259-271.

10. Thompson A, Carter BJ, Turk-Kubo K, Malfatti F, Azam F, Zehr JP. 2014. Genetic
diversity of the unicellular nitrogen-fixing cyanobacteria UCYN-A and its
prymnesiophyte host. Environ Microbiol 16:3238-49.

11. Cohen SE, Golden S. 2015. Circadian rhythms in cyanobacteria. Microb Mol Biol Rev
79:373-385.

12. Bell-Pedersen D, Cassone VM, Earnest DJ, Golden SS, Hardin PE, Thomas TL, Zoran
MJ. 2005. Circadian rhythms from multiple oscillators: Lessons from diverse
organisms. Nature Reviews Genetics 6:544-556.

13. Reimers AM, Knoop H, Bockmayr A, Steuer R. 2017. Cellular trade-offs and optimal
resource allocation during cyanobacterial diurnal growth. Proc Natl Acad Sci USA
doi:https://doi.org/10.1073/pnas.1617508114.

14. Golden SS, Ishiura M, Johnson CH, Kondo T. 1997. Cyanobacterial circadian rhythms.
Annual Review of Plant Physiology and Plant Molecular Biology 48:327-354.

15. Kondo T, Ishiura M. 2000. The circadian clock of cyanobacteria. BioEssays 22:10-15.

16. Toepel J, Welsh E, Summerfield TC, Pakrasi HB, Sherman LA. 2008. Differential
transcriptional analysis of the cyanobacterium *Cyanothece* sp. strain ATCC 51142
during light-dark and continuous-light growth J. Bacteriol 190:3904-3913.

17. Shi T, Ilikchyan IN, Rabouille S, Zehr JP. 2010. Genome-wide analysis of diel gene
expression in the unicellular N_2-fixing cyanobacterium *Crocosphaera watsonii* WH
8501. ISME J 4:621-632.

18. Zinser ER, Lindell D, Johnson ZI, Futschik ME, Steglich C, Coleman ML, Wright MA,
Rector T, Steen R, McNulty NP, Thompson LR, Chisholm SW. 2009. Choreography of
the transcriptome, photophysiology, and cell cycle of a minimal photoautotroph,
Prochlorococcus. PLoS ONE 4:e5135.

19. Cornejo-Castillo FM, Cabello AM, Salazar G, Sanchez-Baracaldo P, Lima-Mendez G,
Hingamp P, Alberti A, Sunagawa S, Bork P, de Vargas C, Raes J, Bowler C, Wincker
P, Zehr JP, Gasol JM, Massana R, Acinas SG. 2016. Cyanobacterial symbionts
diverged in the late Cretaceous towards lineage-specific nitrogen fixation factories in
single-celled phytoplankton. Nat Commun 7:11071.

20. Rong-Fong L, Kun-Da T, Tan-Chi H. 2003. Factors affecting the circadian degradation
of COP23 in *Synechococcus* RF-1. Botanical Bulletin-Academia Sinica Taipei 44:151-
158.

21. Klatt CG, Liu Z, Ludwig M, Kühl M, Jensen SI, Bryant DA, Ward DM. 2013.
Temporal metatranscriptomic patterning in phototrophic *Chloroflexi* inhabiting a
microbial mat in a geothermal spring. ISME Journal 7:1775-1789.

22. Steunou AS, Jensen SI, Brecht E, Becraft ED, Bateson MM, Kilian O, Bhaya D, Ward
DM, Peters JW, Grossman AR, Kühl M. 2008. Regulation of *nif* gene expression and
the energetics of N_2 fixation over the diel cycle in a hot spring microbial mat. The
ISME Journal 2:364-378.

23. Holtzendorff J, Partensky F, Mella D, Lennon J-F, Garczarek L. 2008. Genome
streamlining results in loss of robustness of the circadian clock in the marine
cyanobacterium *Prochlorococcus marinus* PCC 9511. J. Biol Rhythms 23:187-199.

24. Axmann I, Hertel S, Wiegard A, Dörrich A, Wilde A. 2014. Diversity of KaiC-based
timing systems in marine Cyanobacteria. Marine Genomics 14:3-16.

25. Fay P. 1992. Oxygen relations of nitrogen fixation in cyanobacteria. Microbiology and
Molecular Biology Reviews 56:340-373.

26. Staal M, Rabouille S, Stal LJ. 2007. On the role of oxygen for nitrogen fixation in the
marine cyanobacterium *Trichodesmium* sp. Envionmental Microbiology 9:727-736.

27. Bürgmann H, Widmer F, Sigler WV, Zeyer J. 2003. mRNA extraction and reverse
transcription-PCR protocol for detection of *nif*H gene expression by *Azotobacter*
vinelandii in soil. Appl. Environ. Microbiol. 69:1928-1935.

28. Chen YB, Dominic B, Mellon MT, Zehr JP. 1998. Circadian rhythm of nitrogenase
gene expression in the diazotrophic filamentous nonheterocystous cyanobacterium
Trichodesmium sp. strain IMS 101. J Bacteriol 180:3598-3605.

29. Zubkov MV, Martin AP, Hartmann M, Grob C, Scanlan D. 2015. Dominant oceanic
bacteria secure phosphate using a large extracellular buffer. Nature Commun 6.

30. Zubkov MV, Mary I, Woodward EMS, Warwick PE, Fuchs BM, Scanlan DJ, Burkill
PH. 2007. Microbial control of phosphate in the nutrient-depleted North Atlantic
subtropical gyre. Environ. Microbiol 9:2079-2089.

31. Michelou VK, Lomas MW, Kirchman DL. 2011. Phosphate and adenosine-5´-
triphosphate uptake by cyanobacteria and heterotrophic bacteria in the Sargasso Sea.
Limnol Oceanogr 56:323-332.

32. Stockel J, Welsh EA, Liberton M, Kunnvakkam R, Aurora R, Pakrasi HB. 2008. Global
transcriptomic analysis of *Cyanothece* 51142 shows a robust diurnal oscillation of
central metabolic processes. *Proc Natl Acad Sci* USA 105:6156-6161.

33. Davey M, Tarran G, Mills M, Ridame C, Geider R, Roche JL. 2008. Nutrient limitation
of picophytoplankton photosynthesis and growth in the tropical North Atlantic. Limnol
Oceanogr 53:1722-1733.

34. Klausmeier C, Litchman E, Levin S. 2004. Phytoplankton growth and stoichiometry
under multiple nutrient limitation. Limnol Oceanogr 49:1463-1470.

35. Harke MJ, Frischkorn KR, Haley ST, Aylward FO, Zehr J, Dyhrman ST. 2018.
Periodic and coordinated gene expression between a diazotroph and its diatom host.
ISME Journal doi:https://doi.org/10.1038/s41396-018-0262-2.

36. Ran L, Larsson J, Vigil-Stenman T, Nylander J, Ininbergs K, Zheng W, Lapidus A,
Lowry S, Haselkorn R, Bergman B. 2010. Genome erosion in a nitrogen-fixing
vertically transmitted endosymbiotic multicellular cyanobacterium. PLoS ONE
5:e11486.

37. Nakayama T, Kamikawa R, Tanifuji G, Kashiyama Y, Ohkouchi N, Archibald JM,
Inagaki Y. 2014. Complete genome of a nonphotosynthetic cyanobacterium in a diatom
reveals recent adaptations to an intracellular lifestyle. *Proc Natl Acad Sci*
USA:201405222.

38. Ohbayashi R, Watanabe H, Ehira S, Kanesaki Y, Chibazakura T, Yoshikawa H. 2016.
Diversification of DnaA dependency for DNA replication in cyanobacterial evolution.
The ISME J 10:1113-1121.

39. Wilson ST, Aylward FO, Ribalet F, Barone B, Casey JR, Connell PE, Eppley JM,
Ferrón S, Fitzsimons JN, Hayes CT, Romano AE, Turk Kubo K, Vislova A, Armbrust
EV, Caron DA, Church M, Zehr JP, Karl D, DeLong E. 2017. Coordinated regulation
of growth, activity and transcription in natural populations of the unicellular nitrogen-
fixing cyanobacterium *Crocosphaera*. Nature Microbiol 2.

40. Hagen KD, Meeks JC. 2001. The unique cyanobacterial protein OpcA is an allosteric
effector of glucose-6-phosphate dehydrogenase in *Nostoc punctiforme* ATCC 29133.
Journal of Biological Chemistry 276:11477-11486.

41. Min HT, Golden SS. 2000. A new circadian class 2 gene, *opcA*, whose product is
important for reductant production at night in *Synechococcus elongatus* PCC 7942. J
Bacteriol 182:6214-6221.

42. Lee S, Ryu J, SY, Jeon J, Song J, Cho H, Choi S, Choi D, Tandeau de Marsac N, Park
Y. 2007. Transcriptional regulation of the respiratory genes in the cyanobacterium
Synechocystis sp. PCC 6803 during the early response to glucose feeding. Plant Physiol
145:1018-1030.

43. Maldener I, Muro-Pastor AM. 2010. Cyanobacterial heterocysts. *In* Sons JW (ed),
Encyclopedia of Life Sciences (ELS) doi:DOI:
10.1002/9780470015902.a0000306.pub2, Chichester, UK.

44. Martens J, Barg H, Warren M, Jahn D. 2002. Microbial production of vitamin B12.
Appl Microbiol Biotechnol 58:275-285.

45. Helliwell KE, Lawrence AD, Holzer A, Kudahl UJ, Sasso S, Kräutler B, Scanlan DJ,
Warren MJ, Smith AG. 2016. Cyanobacteria and eukaryotic algae use different
chemical variants of vitamin B12. Curr Biol 26:999-1008.

46. Milligan AJ, Berman-Frank I, Gerchman Y, Dismukes GC, Falkowski PG. 2007. Light-
dependent oxygen consumption in nitrogen-fixing cyanobacteria plays a key role in
nitrogenase protection. J Phycol 43:845-852.

47. Valladares A, Herrero A, Pils D, Schmetterer G, Flores E. 2003. Cytochrome c oxidase
genes required for nitrogenase activity and diazotrophic growth in *Anabaena* sp. PCC
7120. Mol Microbiol 47:1239-1249.

48. Jones KM, Haselkorn R. 2002. Newly identified cytochrome c oxidase operon in the
nitrogen-fixing cyanobacterium *Anabaena* sp. strain PCC 7120 specially induced in
heterocyts. J Bacteriol 184:2491-2499.

49. Lafiti A, Ruiz M, Cheng-Cai Z. 2009. Oxidative stress in cyanobacteria. FEMS
Microbiol Rev 33:258-278.

50. Milligan AJ, Berman-Frank I, Gerchman Y, Dismukes GC, Falkowski PG. 2007. Light-
dependent oxygen consumption in nitrogen-fixing cyanobacteria plays a key role in
nitrogenase protection. J Phycol 43:845-852.

51. Nakayama T, Archibald JM. 2012. Evolving a photosynthetic organelle. BMC Mol
Biol 10.

52. Collins TJ. 2007. Introduction to ImageJ for light microscopy. Microsc Microanal
13:1674-1675.

53. Chen Y, Zehr JP, Mellon MT. 1996. Growth and nitrogen fixation of the diazotrophic
filamentous nonheterocystous cyanobacterium *Trichodesmium* sp. IMS 101 in defined
media: evidence for a circadian rhythm J Phycol 32:916-923.

54. Shilova IN, Robidart JC, Tripp HJ, Turk-Kubo K, Wawrik B, Post AF, Thompson AW,
Ward B, Hollibaugh JT, Millard A. 2014. A microarray for assessing transcription from
pelagic marine microbial taxa. The ISME J 8:1476-1491.

55. Lemire A, Lea K, Batten D, Gu SJ, Whitley P, Bramlett K. 2011. Development of
ERCC RNA spike-in control mixes. J Biomol Tech 22:S46.

56. Sun S, Chen J, Li W, Altinatas I, Lin A, Peltier S, Stocks K, Allen EE, Ellisman M,
Grethe J, Wooley J. 2011. Community cyberinfrastructure for Advanced Microbial
Ecology Research and Analysis: the CAMERA resource Nucl Acids Res 39:D546-
D551.

57. Huang Y, Niu B, Gao Y, Fu L, Li W. 2010. CD-HIT Suite: a web server for clustering
and comparing biological sequences. Bioinformatics 26:680-682.

58. Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets
of protein or nucleotide sequences. Bioinformatics 22:1658 - 1659.

59. Gentleman R, Carey V, Bates D, Bolstad B, Dettling M, Dudoit S, Ellis B, Gautier L,
Ge Y, Gentry J, Hornik K, Hothorn T, Huber W, Iacus S, Irizarry R, Leisch F, Li C,
Maechler M, Rossini A, Sawitzki G, Smith C, Smyth G, Tierney L, Yang J, Zhang J.
2004. Bioconductor: open software development for computational biology and
bioinformatics. Genome Biology 5:R80.

60. Huber W, Carey JV, Gentleman R, Anders S, Carlson M, Carvalho SB, Bravo CH,
Davis S, Gatto L, Girke T, Gottardo R, Hahne F, Hansen DK, Izarry AR, Lawrence M,
Love IM, Waldron L, Morgan. 2015. Orchestrating high-throughput genomic analysis
with Bioconductor. Nat Meth 12:115121.

61. Smyth GK. 2005. Limma: Linear models for microarray data., p 397-420. *In* Gentleman
R, Carey V, Dudoit S, Irizarry R, Huber W (ed), Bioinformatics and Computational
Biology Solutions using R and Bioconductor. Springer, New York.

62. Kauffmann A, Gentleman R, Huber W. 2009. arrayQualityMetrics-a bioconductor
package for quality assessment of microarray data. Bioinformatics 25:415-416.

63.   Bolstad BM, Collin F, Brettschneider J, Simpson K, Cope L, Irizarry RA, Speed TP.
726   2005. Quality Assessment of Affymetrix GeneChip Data, p 33-47. *In* Springer NY (ed),
727   Bioinformatics and Computational Biology Solutions using R and Bioconductor.

728   64.   Bolstad B. 2004. Low level analysis of high-density oligonucleotide Array Data:
729   Background, Normalization and SummarizationUniversity of California, Berkeley.

730   65.   Irizarry R, Hobbs B, Collin F, Beazer-Barclay Y, Antonellis K, Scherf U, Speed T.
731   2003. Exploration, normalization, and summaries of high density oligonucleotide array
732   probe level data. Biostatistics 4:249.

733   66.   Futschik ME, Herzel H. 2008. Are we overestimating the number of cell-cycling genes?
734   The impact of background models on time series analysis. Bioinformatics 24:1063-
735   1069.

736   67.   Moritz S, Sardá A, Bartz-Beielstein T, Zaefferer M, Stork J. 2015. Comparison of
737   different methods for univariate time series imputation in R. arXiv:151003924

738   68.   McMurdie P, Holmes S. 2013. Phyloseq: An R package for reproducible interactive
739   analysis and graphics of microbiome census data. PLoS ONE 8:e61217.
740

 Titles and legends to main figures

Figure 1. Four different clades based on Pearson correlation of the transcription profile of

UCYN-A1 and UCYN-A2 genes over light–dark cycles. The transcription value of each gene at

each time point was normalized to the mean at all time points and divided by standard deviation

(SD) (*Y* axis, log 2 scale). The *X* axis represents time points where D and L stand for dark and

light, respectively, followed by the corresponding hour into the light or dark periods. The second

light-dark cycle is shown as 2D followed by the number of the corresponding hours entering light

or dark period. The shaded area represents the dark period. In each cluster, most representative

genes are listed in the table attached to the plot. UCYN-A1 genes are coded in green and UCYN-

A2 genes are coded in orange.

753
Figure. 2. A. Transcriptional network based on Pearson correlation of gene transcription over the

diel cycle in all studied cyanobacteria. The genes are connected if correlation coefficient for their

transcription patterns is higher than 0.5. The genes shown are diel genes with variable transcription

patterns among the studied cyanobacteria. The arrows point to genes for glycolysis, PPP and N_2

fixation in the studied diazotrophs. The purple circles demarcate genes for ribosomal proteins

included in the analysis. Abbreviations: *Prochlorococcus* sp. MED4 (Pro), *Cyanothece* sp. ATCC

51142 (Cya), *C. watsonii* WH 8501 (Cro), *T. erythraeum* (Tricho), Pentose Phosphate Pathway

(PPP), Photosystem I (PSI). **B.** Four time course plots are attached for the N_2-fixing cyanobacteria

showing the diel transcription patterns of photosystem I, N_2 fixation and genes for ribosomal

proteins.

Figure 3. Transcription of genes for replication and cell division in UCYN-A. **Upper Panel: A)**

Diel transcription patterns for cell division and replication genes in UCYN-A1 and UCYN-A2

over the light-dark cycle. Hierarchical clustering of genes was based on Pearson correlation

between their transcription profiles. The transcription values of genes at each time point were

standardized, and the blue-red scale shows how many standard deviations a transcription value

was lower or higher, respectively, from the mean transcription values over the diel cycle (Z score).

Gene ID and gene product corresponding to each gene for UCYN-A1 and UCYN-A2 are shown.

Time is shown on X-axis as light (L) and dark (D), respectively, followed by the corresponding

hour after the sunrise and sunset periods started. The second light-dark cycle is shown as 2D

followed by the number of the corresponding hours entering light or dark period.

Lower Panel: Epifluorescence micrographs of dividing UCYN-A2 detected with CARD-FISH

(19). **B)** Two big clusters of UCYN-A2 cells and the haptophyte host attached. Left Panel: the

nucleus of the host and the UCYN-A2 cells were visualized with-DAPI stain (blue). Right Panel:

The UCYN-A2 (red) and its haptophyte host (green). **C)** Two different associations of UCYN-A2

with its haptophyte dividing in samples from Scripps Pier.

Figure 4. Network showing the Pearson correlation for gene transcriptions in the unicellular N_2-

fixing cyanobacteria *Cyanothece* sp. ATCC 51142 (*Cyanothece*), *C. watsonii* WH 8501

(*Crocosphaera*) and UCYN-A. Shown here are key genes in major metabolic pathways with

distinct diel transcription patterns. The genes are connected if their correlation coefficient for

transcription patterns is higher than 0.2. PPP, pentose phosphate pathway; PSI, photosystem I.

Figure 5. Transcriptional profiles of the genes for glycolysis over light–dark cycles in the

cyanobacteria studied here. The transcription value of each gene at each time point was

normalized to the mean at all time points and divided by standard deviation (SD) (*Y* axis, log

scale). The *X* axis represents time points where D and L stand for dark and light, respectively,

followed by the corresponding hour into the light or dark periods. The second light-dark cycle is

shown as 2D followed by the number of the corresponding hours entering light or dark period. The

shaded area represents the dark period.

Figure 6. Transcriptional profiles of *opcA* (allosteric effector) and *zwf* (glucose-6-P

dehydrogenase) over light–dark cycles in the cyanobacteria studied here. The transcription value

of each gene at each time point was normalized to the mean at all time points and divided by

standard deviation (SD) (*Y* axis, log scale). The *X* axis represents time points where D and L stand

for dark and light, respectively, followed by the corresponding hour into the light or dark periods.

The second light-dark cycle is shown as 2D followed by the number of the corresponding hours

entering light or dark period. The shaded area represents the dark period.

Figure 7. Schematic model of UCYN-A showing the possible main cellular functions, metabolic

pathways and transporters.

Organism
- UCYN-A1
- UCYN-A2

Panel I

UCYN-A1		UCYN-A2	
fabD	ftsZ	-	-
-	tpi	fabH	-
lpxC	-	lpxC	rpe
sqd1	pdha	-	pdha
hypD	-	-	thiL
cbio	acp	-	acp
pstA	glnA	-	glnA
pstC	g6pd	pstcC	-
minD	-	-	accD
-	accA	minE	-
-	-	mreC	topA
g6pi	-	-	rpoE
murG	dpo3	-	-
murE	-	-	sufB
-	fepB	murJ	-
-	fepGD	murB	-

Panel III

UCYN-A1	UCYN-A2
atpF	atpF
atpB	atpB
-	atpA
pgd	-
rpiA	pstA
cop23	cop23
iscA	iscA
-	eno
-	ndhK
-	ndhC
-	psaJ
-	psaL
-	tktA
-	pgd

Panel II

UCYN-A1		UCYN-A2	
ndhJ	petC	ndhJ	petC
ndhD	petF	ndhD	petF
hupS	nikB	-	-
-	sufB	hypE	-
-	-	hypD	petH
g3pd	-	-	lpxA
-	-	fba	lpxB
atpA	fabH	atpA	-
atpI	fabF	atpI	-
-	prpS	atpF	-
atpC	accC	-	-
psaC	-	psaC	gpi
minE	nifU	-	-
-	nifH	ftsZ	nifH
-	nifD	murD	nifD
prxR	sod1	prxR	-
petA	-	petA	nifK

Panel IV

UCYN-A1		UCYN-A2	
psaA	-	psaA	rps13
psaB	-	-	rpl13
oprB	-	oprB	rpl18
rpl11	-	rpl11	rpl5
nusG	-	-	rps9
secE	-	-	rpl24
rps19	-	rps19	rpl2
rpl22	-	rpl22	rpl3
rpl16	-	-	rpl23
rpl24	-	-	rpl29
rps8	-	-	rpl1
rpl6	-	-	guaB
rpl15	-	rpl15	sqdB
dnaE	-	-	acyl
truA	-	-	rpoA
-	-	coxB	-

A.

Ribosomal proteins (Pro, Tricho and UCYN-A)

N₂ fixation (Cya and Cro)

Glycolysis and PPP in UCYN-A

Glycolysis and PPP (Tricho, Pro, Cya and Cro)

N₂ fixation (Tricho and UCYN-A)

Ribosomal proteins (Cya and Cro)

Organism
- ● *C. watsonii* WH 8501
- ▲ *Cyanothece* sp. ATCC 51142
- ■ *Prochlorococcus* sp. MED4
- ✛ *T. erythraeum*
- ⊠ UCYN-A

Pathway
- Cell division
- Glycolysis
- N₂ fixation
- Nucleotide
- PPP
- PSI
- Replication
- Respiration
- Ribosomal proteins
- Transport Fe
- Transport PO₄

B.

UCYN-A

Trichodesmium erythraeum IMS101

Crocosphaera watsonii WH 8501

Cyanothece ATCC 51142

A.

Gene_ID	Strain	Description
646530139	UCYN-A1	septum site-determining protein *minD*
646530123	UCYN-A1	DNA gyrase subunit A
2528848158	UCYN-A2	rod shape-determining protein *mreC*
2528847831	UCYN-A2	DNA polymerase III- delta prime subunit
2528847919	UCYN-A2	pyridoxal phosphate-dependent enzyme
2528847263	UCYN-A2	DNA gyrase subunit A
2528847442	UCYN-A2	cell division protein *ftsZ*
2528848139	UCYN-A2	Superfamily II DNA and RNA helicases
646530138	UCYN-A1	cell division topological specificity factor *minE*
2528847429	UCYN-A2	DNA topoisomerase I
646530091	UCYN-A1	DNA topoisomerase IV subunit A
646530302	UCYN-A1	cell division protein *ftsZ*
2528847278	UCYN-A2	cell division topological specificity factor *minE*
2528848219	UCYN-A2	Glycosyltransferases involved in cell wall biogenesis
646530433	UCYN-A1	DNA/RNA helicase, superfamily II

L6   L9   D3   D6   2D12   2L3   2L9   2L12

B.

C.

Unicellular N$_2$-fixing cyanobacteria

Cyanothece

Crocosphaera

UCYN-A

Pathway

- Cell division
- Glycolysis
- N$_2$ fixation
- Nucleotide
- PPP
- PSI
- Replication
- Respiration
- Ribosome
- Transport Fe
- Transport PO$_4$

Glycolysis

Pentose phosphate

Gene_ID	Description	Pathway
Tery_3301	MutS2 family protein	Replication and Repair
Tery_2841	glucose inhibited division protein A	Cell Growth and Death
Tery_4190	single-stranded-DNA-specific exonuclease *RecJ*	Replication and Repair
Tery_0168	Holliday junction DNA helicase *RuvA*	Replication and Repair
Tery_3645	DNA mismatch repair protein *MutS*	Replication and Repair
Tery_0165	DNA recombination protein, *RuvA*	Replication and Repair
Tery_4942	cell cycle protein	Cell Growth and Death
Tery_0212	DNA polymerase III, delta subunit	Replication and Repair
Tery_2316	DNA ligase, NAD-dependent	Replication and Repair
Tery_1165	protein of unknown function DUF59	Cell Growth and Death
Tery_0163	maf protein	Cell Growth and Death
Tery_0413	phage SPO1 DNA polymerase-related protein	Replication and Repair
Tery_0001	chromosomal replication initiator protein *DnaA*	Replication and Repair
Tery_4604	DNA repair protein *RecO*	Replication and Repair
Tery_4232	ATP-dependent DNA helicase *PcrA*	Replication and Repair
Tery_0348	transposase, IS605 *OrfB* family	Replication and Repair
Tery_3302	cell cycle protein	Cell Growth and Death
Tery_3384	cell division topological specificity factor *MinE*	Cell Growth and Death
Tery_3269	endonuclease III	Replication and Repair

Number of genes	*T. erythraeum* IMS101	*C. watsonii* WH8501	*Cyanothece* ATCC 51142	UCYN-A	*Prochlorococcus* MED4
Targeted on the microarray	4,788	4,407	5,040	1,194	1,698
With detected transcription	4,305	4,140	4,312	365	1.511
Have diel pattern of transcription	1,620	1,708	998	316	1,370
Percent					
With detected transcription	89.9	93.9	85.6	30.6	89.0
Have diel pattern of transcription	33.8	38.8	19.8	26.5	80.7

	Sampling Time	Host associated with 1 UCYN-A2 cluster (cells ml^{-1})	Host associated with 2 UCYN-A2 clusters (cells ml^{-1})	Host associated with 2 UCYN-A2 clusters (%)
Diel cycle 1	12 pm	36.5	0.0	0.0
	3 pm	38.0	0.0	0.0
	6 pm	9.1	0.0	0.0
	9 pm	12.2	15.2	55.6
	12 am	6.1	9.1	60.0
	3 am	41.0	10.6	20.6
	6 am	7.6	0.0	0.0
	9 am	38.0	0.0	0.0
Diel cycle 2	12 pm	44.1	0.0	0.0
	3 pm	53.2	0.0	0.0
	6 pm	36.5	0.0	0.0
	9 pm	16.7	28.9	63.3
	12 am	34.9	51.7	59.6
	3 am	51.7	4.6	8.1
	6 am	66.8	0.0	0.0
	9 am	27.3	0.0	0.0

Probe Name	Target	Sequence (5' to 3')	Reference
UPRYM69	Host-A1	CACATA**G**GAACATCCTCC	Cornejo-Castillo et al.(6)
UPRYM69 competitor	Host-A2 used as Host-A1 competitor	CACATT**T**GGAACATCCTCC	Cornejo-Castillo et al.(6)
UBRADO69	Host-A2	CACATT**T**GGAACATCCTCC	Cornejo-Castillo et al.(6)
UBRADO69 competitor	Host-A1 used as Host-A2 competitor	CACATA**G**GAACATCCTCC	Cornejo-Castillo et al.(6)
Helper A-PRYM	*Haptophyta*	GAAAGGTGCTGAAGGAGT	Cornejo-Castillo et al.(6)
Helper B-PRYM	*Haptophyta*	AATCCCTAGTCGGCATGG	Cornejo-Castillo et al.(6)
UCYN-A1 732	UCYN-A1	GTTA**C**GGTCCAGTAGCAC	Krupke et al.(5)
UCYN-A1 competitor	UCYN-A2 used as UCYN-A1 competitor	GTTG**G**CGGTCCAGTAGCAC	Cornejo-Castillo et al.(6)
UCYN-A2 732	UCYN-A2	GTTG**G**CGGTCCAGTAGCAC	Cornejo-Castillo et al.(6)
UCYN-A2 competitor	UCYN-A1 used as UCYN-A2 competitor	GTTA**C**GGTCCAGTAGCAC	Krupke et al.(5)
Helper A-732	UCYN-A	GCCTTCGCCACCGATGTTCTT	Krupke et al.(5)
Helper B-732	UCYN-A	AGCTTTCGTCCCTGAGTGTCA	Krupke et al.(5)