Precisely control mitochondria with light to manipulate cell fate decision

Patrick Ernst[1], Ningning Xu[1], Yun Lu[1,2], Jing Qu[2], Herbert Chen[3], Matthew S. Goldberg[4], Jianyi Jay Zhang[1], Victor Darley-Usmar[5], Brian O'Rourke[6], Xiaoguang Liu[1], and Lufang Zhou[1,2*]

Departments of Biomedical Engineering[1], Medicine [2], Surgery[3], Neurology[4], and Pathology[5], University of Alabama at Birmingham, Birmingham, Alabama, USA, 35294
[6]Division of Cardiology, Department of Medicine, The Johns Hopkins University, Baltimore, Maryland, USA, 21205

Running title: Optogenetic control of mitochondria

* Correspondence:
Lufang Zhou, PhD
Division of Cardiovascular Disease
Department of Medicine
University of Alabama at Birmingham
703 19th Street South, ZRB 306
Birmingham, AL 35294, USA
Email: lfzhou@uab.edu
Phone: 205-975-3764

Keyword: optogenetics, mitochondria, parkin, apoptosis, autophagy

Abstract

Mitochondrial dysfunction has been implicated in many pathological conditions or diseases. The normal functioning of mitochondria relies on maintaining the inner mitochondrial membrane (IMM) potential (*a.k.a.* $\Delta\Psi_m$) that is essential for ATP synthesis, Ca^{2+} homeostasis, redox balance and regulation of other key signaling pathways such as mitophagy and apoptosis. However, the detailed mechanisms by which $\Delta\Psi_m$ regulates cellular function remain incompletely understood, partially due to difficulty of manipulating $\Delta\Psi_m$ with spatiotemporal resolution, reversibility, or cell type specificity. To address this need, we have developed a new generation optogenetic-based technique for controllable mitochondrial depolarization with light. We demonstrate successful targeting of the heterologous Channelrhodopsin-2 (ChR2) fusion protein to the IMM and formation of functional cationic channels capable of light-induced selective $\Delta\Psi_m$ depolarization and mitochondrial autophagy. Importantly, we for the first time show that optogenetic-mediated mitochondrial depolarization can be well-controlled to differentially influence the fate of cells expressing mitochondrial ChR2: while sustained moderate light illumination induces substantial apoptotic cell death, transient mild light illumination elicits cytoprotection *via* mitochondrial preconditioning. Finally, we show that Parkin overexpression exacerbates, instead of ameliorating, mitochondrial depolarization-mediated cell death in HeLa cells. In summary, we provide evidence that the described mitochondrial-targeted optogenetics may have a broad application for studying the role of mitochondria in regulating cell function and fate decision.

Introduction

Mitochondria lie at the crossroad of cellular metabolic and signaling pathways and thus play a pivotal role in regulating many intracellular processes such as energy production, reactive oxygen species (ROS) production, Ca^{2+} handling, and cell fate decision (1). Not surprisingly, defects of mitochondrial function have been implicated in a variety of human maladies such as aging, cardiovascular disease, cancer, and neurodegeneration (2). The normal functioning of mitochondria largely relies on maintaining the inner membrane potential ($\Delta\Psi_m$) to preserve the protonmotive force necessary to drive oxidative phosphorylation and redox balance. Interestingly, while profound depolarization of $\Delta\Psi_m$ is detrimental and induces cell injury (3), partial mitochondrial dissipation can be cytoprotective (4, 5). The detailed mechanism underlying the differential regulatory role of $\Delta\Psi_m$ on cell function and fate remains incompletely understood, partially due to the difficulty of controlling $\Delta\Psi_m$ or mitochondrial membrane permeability (MMP) with spatiotemporal resolution, reversibility, and with cell type specificity in an experimental setting. Currently, manipulation of $\Delta\Psi_m$ is mainly *via* pharmacological intervention, i.e. using chemicals to uncouple the mitochondria (4, 5) or to induce permeability transition pore (mPTP) opening (6). However, pharmacological approaches often cause unknown side effects and lack the ability to probe spatiotemporal domains. Several research groups, including us, use laser flashes to trigger local or cell wide mitochondrial depolarization *via* the photooxidation mechanism (7-10). A limitation of this approach is that it requires high-energy laser illumination, which may induce uncontrollable oxidative stress and unpredictable outcomes. A new approach capable of precisely controlling MMP or $\Delta\Psi_m$ is highly desired for advancing mitochondrial research.

Optogenetics has recently emerged as a technique that utilizes genetically-encoded light-sensitive ion channels, such as channelrhodopsin 2 (ChR2) (11-14) or halorhodopsin (NpHR) (15, 16), to precisely and remotely manipulate the activity of cells or animals (11, 13). ChR2 is a seven-transmembrane domain protein that contains the light-isomerizable chromophore all-trans-retinal (17). When excited by blue light, the *all-trans* retinal undergoes *trans-cis* isomerization (18), activating non-selective cationic channels and inducing depolarizing inward currents. Due to its high spatiotemporal resolution, ChR2-based optogenetic techniques have evolved as a powerful tool in basic and translational research. For example, ChR2 has been expressed in neurons (19) to monitor and control their membrane potential, intracellular acidity, Ca^{2+} influx, and other cellular processes (see ref (14) for review). In addition, ChR2 has been

used to manipulate the membrane excitability of cardiomyocytes (20-22), skeletal muscle cells (23) and cell lines expressing voltage-gated ion channels (24, 25).

In addition to remote control of excitable cells, optogenetic techniques have been used to induce and study cell death. For instance, Yang *et al.* showed that an optogenetic approach can induce apoptotic cell death in human glioma cells *via* depolarization of cell membrane potential and influx of Ca^{2+} (26). In another study, Hill *et al.* reported a two-photon chemical apoptotic targeted ablation technique (27). By combining this technique with organelle-targeted fluorescent proteins and biosensors, the authors successfully achieved precise ablation of specific populations of neurons in the mouse brain. Of note, none of these published optogenetic approaches directly targets the inner mitochondrial membrane (IMM) permeability to induce cell death. We hypothesize that functional expression of ChR2 in cellular IMM forms cationic channels capable of light-induced controlled $\Delta\Psi_m$ depolarization and manipulation of cell function and fate. We test this hypothesis in a variety of cell types and our results show that the mitochondrial leading sequence (MLS) of ABCB10, an ATP binding cassette transporter, effectively targeted ChR2 into IMM. We demonstrate that this new generation optogenetic approach can induce controlled $\Delta\Psi_m$ depolarization that influences cell functions such as apoptosis and mitophagy at high spatial resolution.

Results

Targeting ChR2 to mitochondria. The MLS of cytochrome *c* oxidase subunit VIII (*a.k.a.* mito) has been widely used to import proteins such as GFP (28) or pericam (29) to mitochondria. Therefore, we first examined its capability to target ChR2 to mitochondria. A mutant of ChR2, ChR2(H134R) (30), was fused with eYFP and inserted into vector pCMV-myc-mito to generate plasmid mito-ChR2(H134R)-eYFP. The construct and the control vectors (mito-eYFP) were transfected to H9C2 cells, respectively. The expressions of ChR2-eYFP and eYFP in mitochondria were examined by staining mitochondria with Mitoracker® Deep Red and analyzing the colocalization of the green and red channels using confocal microscopy. Results show that eYFP was perfectly colocalized with mitochondria (Fig. S1A) but ChR2-eYFP was not (Fig. S1B). In addition, ChR2-eYFP appeared to be improperly processed by the endoplasmic reticulum (ER), implying that the size of the ChR2(H134R)-eYFP precursor might be impairing import. Thus, we next doubled the copy number of mito sequence and constructed plasmid (mito)$_2$-ChR2(H134R)-eYFP. However, this modification did not improve the import of

ChR2(H134R)-eYFP to mitochondria (data not shown). Thereafter, a series of mito-based ChR2-eYFP plasmids such as mito-ChR2 (H134R)-eYFP-ER (adding ER extraction sequence), mito-ChR2(H134R)(T2-23aa)-eYFP (removing the cell membrane targeting sequence) and (mito)$_3$-ChR2(H134R)-eYFP (to triplicate mito sequence) were constructed. Unfortunately, none of them could efficiently target ChR2-eYFP precursor to the mitochondria (data not shown). We also studied the MLS of ROMK, a candidate of mitochondrial K_{ATP} channel located on IMM (31). Similarly, ROMK MLS successfully imported eYFP into mitochondria (data not shown) but not ChR2-eYFP (Fig. S1C).

ATP binding cassette (ABC) transporters comprise a large family of membrane translocases (32) and are found in all species from microbe to human (33). Of them, ABCB10 is an IMM erythroid transporter with an unusually long MLS (140 amino acids) (34). Studies showed that ABCB10 MLS could lead various membrane proteins to mitochondria (35), while its depletion targeted ABCB10 to the ER (34). This MLS was fused with eYFP or ChR2(H134R)-eYFP, to generate ABCB-eYFP and ABCB-ChR2(H134R)-eYFP vectors, respectively, which were then transfected into H9C2 cells. To our great satisfaction, ABCB-eYFP was exclusively localized in mitochondria (Fig. 1A). Excitingly, ABCB10 MLS successfully translocated ChR2-eYFP precursor into H9C2 mitochondria, as demonstrated by the colocalization of green (eYFP) and red (MitoTracker) channels (Fig. 1B). Importantly, this method of targeting ChR2 to the IMM did not significantly affect mitochondrial membrane potential and reactive oxygen species (ROS) level (Fig. S2). The disposition of ChR2-YFP fusion protein in H9C2 mitochondria was further examined by immunostaining. As shown in figure 1C, the ChR2 protein expression, detected by the ChR2 antibody, colocalized well with MitoTracker. The capability of ABCB10 MLS to translocate ChR2-eYFP to mitochondria was also confirmed in other cell types such as HeLa cells (Fig. 1D) and human induced pluripotent stem cell-derived cardiomyocytes (hiPSC-CMs) (Fig. 1E). Altogether, our data demonstrate successful mitochondrial ChR2 import and expression.

Light illumination-elicited targeted mitochondrial depolarization. After demonstrating mitochondrial ChR2 expression, we examined if light illumination could cause targeted mitochondrial depolarization in ABCB-ChR2-expressing cells. In the example shown in figure 2A, H9C2 cells expressing ABCB-ChR2 (fused with eYFP) were split into two zones with zone 1 receiving blue (475 nm) LED pulse illuminations and zone 2 as control (i.e. no LED illumination). The change of $\Delta\Psi_m$ was measured with TMRM using confocal microscopy (Fig. 2B), which

showed that TMRM fluorescence faded significantly in the illuminated cells (i.e. zone 1) but did not change in the unilluminated cells (i.e. zone 2) (Fig. 2C and Video S1). The averaged TMRM intensities before and after light illumination in the unilluminated and illuminated cells were summarized in figure 2D, showing that $\Delta\Psi_m$ depolarized about 80% in the illuminated cells but remained polarized in the non-illuminated cells. The data suggest that mitochondrial optogenetics can induce mitochondrial depolarization with spatial resolution. In addition to impaired mitochondrial membrane potential, light illumination also caused increased ROS production in ABCB-ChR2 expressing cells, as compared with mock-transfected or ABCB-ChR2-expressing cells without blue light illumination (Fig S3). To verify that mitochondrial depolarization in the illuminated cells was caused by light-induced activation of IMM ChR2 channels instead of imaging laser-induced photooxidation and photobleaching, we performed two control experiments. We first examined the effect of imaging laser alone on $\Delta\Psi_m$ in ABCB-ChR2-expressing cells. As shown in figure S4A (and Video S2A), confocal scanning laser itself had no effect on TMRM. However, turn on blue LED illumination led to significant $\Delta\Psi_m$ depolarization (Fig S4B and Video S2B) similar to that shown in figure 2. In another experiment, we applied the same imaging and LED illumination as that used in the experiment shown in figure 2 to ABCB-eYFP-expressing cells. We found that neither eYFP nor TMRM fluorescent intensity changed upon LED illumination (Fig. S5 and Video S3). Taken together, these data demonstrate functional expression of ChR2 in the IMM that allows light-induced spatially-selective mitochondrial depolarization.

Mitochondrial optogenetic-mediated cell death and the underlying mechanisms. Next, we examined the effect of sustained light illumination on ABCB-ChR2-expressing cells. Twenty-four hours moderate (0.5 mW/mm^2) LED illumination had negligible effects on the viability of non-transfected and mock-transfected HeLa cells (Fig S6), as well as ABCB-eYFP-expressing HeLa cells, but caused significant cell death in ABCB-ChR2-expressing HeLa cells (Fig. 3A), as measured by Trypan Blue staining. The LED illumination-induced cytotoxicity in ABCB-ChR2-expressing cells was confirmed by the lactate dehydrogenase (LDH) assay and AlamarBlue assay. As shown in figure 3B, cell viability dropped 80% in the light-illuminated ABCB-ChR2 cell cultures but only slightly in the ABCB-eYFP-expressing HeLa cells. Consistently, substantial cytotoxicity was detected in the light-illuminated ABCB-ChR2-expressing cells but not in the ABCB-eYFP-expressing cells (Fig 3C). Importantly, we found that pretreating cells with cyclosporine A, a selective mitochondrial permeability transition pore (mPTP) inhibitor, did not

prevent the light-induced cell death (Fig 3D) and cytotoxicity (Fig 3E), indicating that the mitochondrial-targeted optogenetic-mediated cell death is independent of mPTP opening. This finding also denotes that ChR2 proteins formed mPTP-like ion channels in IMM. Of note, antioxidants such as MitoTEMPO and MitoQ failed to rescue cells from light-induced cell death (data not shown), suggesting that ROS production is not a crucial factor in mitochondrial optogenetic-mediated cytotoxicity.

We also examined the time and light irradiance dependence of optogenetic-mediated cell death. As shown in figure 4A, cell viability dropped nearly linearly within 12 hours of light (0.5 mW/mm^2) illumination and tapered off slightly thereafter. Regarding the effect of light intensity, we found that six hours of moderate (e.g., 1 mW/mm^2 and 3 mW/mm^2) light illumination caused minimal cell death in ABCB-ChR2-expressing cell cultures, as measured by Hoechst and Propidium Iodide (PI) staining (Fig S7). Increasing light intensity to 5 mW/mm^2 or 7 mW/mm^2 caused a substantial (62% and 85%, respectively) decrease in cell viability (Fig 4B and 4C). Importantly, light illumination at the similar intensities (e.g., 3 mW/mm^2 and 5 mW/mm^2) did not induce detectable cell death in mock-transfected cultures (Fig 4B, 4C and Fig S7). When light intensity was further increased (e.g. >8 mW/mm^2), illumination (24 hours) begun to cause cell death in the mock-transfected cells (data not shown).

To explore the mechanistic pathway underlying the mitochondrial optogenetic-mediated cell death, we treated cells with an apoptosis inhibitor and necroptosis inhibitor, respectively. Cell death analysis showed that Z-VAD-FMK (20µM), a caspase inhibitor, substantially reduced light illumination-induced cell death, but 7-Cl-O-Nec-1 (100µM), a necroptosis inhibitor, had little effect on cell viability (Fig 5A), suggesting that the optogenetic-mediated light-induced cell death is likely through the caspase-dependent apoptotic pathway. We also noticed that mitochondrial membrane potentials were less depolarized in Z-VAD-FMK-treated cell cultures upon light illumination (Fig 5B). The light illumination-elicited activation of the intrinsic apoptosis pathway in ABCB-ChR2-expressing cells was confirmed by cytochrome C release experiments. As shown in figure 5C, immunocytochemistry revealed that cytochrome C was colocalized with TOM20, a mitochondrial outer membrane protein, in both the mock-transfected cells and ABCB-ChR2-expressing cells not illuminated by LED light. However, light-illuminated ABCB-ChR2-expressing cells showed a punctate/diffuse cytochrome C staining, which was disparate from TOM20 immunostaining, indicating the release of cytochrome C into the cytosol (Fig 5C).

Mitochondrial optogenetic-mediated mitochondrial autophagy. We next examined whether optogenetics-mediated mitochondrial depolarization and damage elicits mitochondrial autophagy (*a.k.a.* mitophagy). As HeLa cells do not express endogenous Parkin that is required for the canonical PINK1/Parkin mediated mitophagy pathway, we overexpressed mCherry-Parkin in ABCB-ChR2-eYFP-expressing HeLa cells, which were then subjected to LED illumination (24 hours, 0.5 mW/mm^2). The cellular localization of Parkin was examined by confocal microscopy at different time points during light illumination. As shown in figure 6A, at the baseline (i.e. 0 hour) mitochondria (represented by eYFP fluorescence) are tubular like and mCherry-Parkin was clearly cytosolic and diffuse, as expected. After 4 hours of LED illumination, mCherry-Parkin began to translocate from the cytosol to some mitochondria, as indicated by colocalization of Parkin and eYFP. After 8 hours of LED illumination, mitochondria became fragmented and mCherry-Parkin translocated to most of the mitochondria. After 24 hours, most of the mCherry-Parkin was found to surround mitochondria, as shown in the enlarged view of the boxed region, demonstrating that Parkin was recruited to mitochondrial outer membranes. On the contrary, no evident mitochondrial translocation of Parkin was observed in the light-illuminated ABCB-eYFP-expressing cells (Fig S8A) and the un-illuminated ABCB-ChR2-eYFP-expressing cells (Fig S8B) in 24 hours. Furthermore, light-illuminated ABCB-ChR2-eYFP-expressing cells displayed less mitochondrial mass compared with other groups, especially after 24 hours, suggesting induction of Parkin-dependent mitophagy. The optogenetic-induced mitochondrial autophagy was confirmed by the aggregation of the autophagosome marker LC3 (Fig 6B, red) and lysosome marker LysoTracker (Fig. 6C, red) and their colocalization with mitochondria (represented by YFP). Neither maker aggregated and exhibited colocalization with YFP in the un-illuminated ABCB-ChR2-eYFP cells (Fig. S8C and S8E) or the mock-transfected cells (Fig. S8D). Together, these data demonstrate mitochondrial optogenetic-induced mitophagy.

As activation of mitophagy is thought to be a cell survival mechanism by eliminating damaged mitochondria, we asked whether Parkin overexpression and activation of Parkin-mediated mitophagy pathway reduces optogenetic-mediated cell death. ABCB-ChR2-expressing HeLa cells, with or without Parkin expression, were illuminated with light (24 hours, 0.5 mW/mm^2) and then subjected to $\Delta\Psi_m$ and cell death analysis. As expected, expression of Parkin did not affect mitochondrial membrane potential (Fig. 7A) and cell viability (data not shown) in the absence of light illumination. However, expression of Parkin surprisingly exacerbated light-induced mitochondrial depolarization (Fig 7A). In addition, although Parkin expression did not cause evident difference in cell viability in ABCB-ChR2-expressing cells

immediately after 24 hours light illumination, significantly more severe cell death was detected in the Parkin-expressing HeLa cells 24 hours after stopping LED illumination (Fig 7B). The pro-apoptotic effect of Parkin activation has been reported in recent studies (36, 37). Furthermore, in our hands Parkin did not abrogate or promote cell death in FCCP-treated (10 µM, 24 hours) HeLa cells (data not shown).

Mitochondrial optogenetic-mediated preconditioning and cytoprotection. As previous studies indicated that mitochondrial preconditioning protects cells from stress, we examined if mitochondrial optogenetic-mediated partial mitochondrial depolarization reduces apoptotic cell death in ABCB-ChR2-expressing cells. As shown in figure 8A, 6 hours of moderate light (4 mW/mm^2) illumination caused nearly 60% decrease in cell viability, which was consistent with data reported in figure 5. Intriguingly, cell viability in those cultures pretreated with 2 hours of mild (0.2 mW/mm^2) light illumination was significantly higher compared with the non-preconditioned group (i.e. without pretreatment) (Fig. 8B), as summarized in figure 8C. In addition, confocal imaging showed that light illumination caused 78% reduction in the mitochondrial membrane potential and preconditioning significantly alleviated mitochondrial depolarization (Fig. 8D-8G).

Discussion

Although optogenetics has been widely used to control the activity of excitable cells *in vitro* and *in vivo* (25, 38, 39), application of this innovative technique to manipulate intracellular organelles such as mitochondria has been limited until recently (40, 41). One challenge to developing a mitochondrial-based optogenetic approach is targeted expression of light-gated channel proteins in the highly folded IMM. In the present study, we have examined the capability of several commonly used MLS, including *mito*, to target the heterologous ChR2 proteins to mitochondria. While Taktch *et al.* have reported that tandem repeat *mito* sequence can target ChR2 to mitochondria (40), triplicating *mito* sequence leads to little ChR2 mitochondrial localization in our hands. Instead, we provide evidence that the ABCB10 MLS can effectively translocate ChR2 to the IMM in a variety of cells types including H9C2, HeLa and cardiomyocyte. In addition, with the ABCB MLS the plasmid membrane targeting sequence of ChR2 does not need be removed for mitochondrial localization. We further demonstrate that the expressed ChR2 proteins form functional cationic ion channels in the IMM, enabling targeted induction of mitochondrial depolarization by blue light. We confirm the optogenetic-mediated

$\Delta\Psi_m$ dissipation by measuring $\Delta\Psi_m$ using confocal live cell imaging and flow cytometry assay, as well as by the translocation of Parkin from the cytosol to mitochondria, which is a well-established marker of mitochondrial depolarization.

The advantages of the optogenetic-based approach over the conventional pharmacological methods to induce mitochondrial depolarization include high spatial resolution *via* optics and specific targeting of ChR2 using cell type-specific promoters or vectors. The light stimulus can be controlled over the high spatial extent, particularly at subcellular dimensions, due to its optical nature. Thus, optogenetic-based light illumination would allow controllable manipulation of individual mitochondria or a group of mitochondria within a single cell. The capability of inducing mitochondrial membrane permeability in specific mitochondria with light can enable novel research, such as investigating how $\Delta\Psi_m$ impacts mitochondrial dynamics and network properties, Ca^{2+} handling, motility, mitophagy, or ROS-dependent signal propagation within or across cells. Spatial resolution is especially important for studies of neurons, which have much of their mitochondrial mass located in distal axons or dendrites. There has been considerable debate regarding how neurons handle damaged or dysfunctional mitochondria with some studies concluding that Parkin-mediated mitophagy only occurs near mature lysosomes in the cell body and that retrograde transport is required for degradation of distal mitochondria, while others concluding that Parkin-mediated mitophagy occurs in distal axons without retrograde transport to the soma (42, 43). Moreover, there may be distinct mechanisms by which neurons handle proximal vs. distal dysfunctional mitochondria and it is important to better define these mechanisms given that mitochondrial dysfunction is implicated in many neurodegenerative diseases and that degeneration of distal axon terminals may be one of the earliest stages and a better therapeutic target (44). The methods described here are ideally suited for investigating this because illumination can selectively depolarize only distal or only proximal mitochondria *in vitro* or *in vivo* using fiber optics.

In addition to high spatial resolution, the optogenetics approach provides specific targeting of tissues or cell types. Chemical uncoupling agents, such as the ionophores FCCP and CCCP, are widely used for the induction of mitochondrial depolarization. However, interpretation of the results can be confounded, especially in intact cells, because of their known off-target effects. For instance, It has been reported that both FCCP and CCCP inhibit the movement of mitochondria in neurites of chick sensory neurons in culture (45). Studies also showed that FCCP mediates conductance of several ions (e.g., H^+, Na^+ and Cl^-) in the plasma membrane (46), affecting membrane electrical characteristics (such as causing cell depolarization) (47)'(48).

On the contrary, the mitochondrial optogenetics approach shown here allows direct light-elicited $\Delta\Psi_m$ depolarization by targeted expression of ChR2 proteins in the IMM. Moreover, compared with the conventional photoactivation method that requires high energy laser flash, optogenetic-mediated $\Delta\Psi$m depolarization requires very low energy (mW/mm^2 level), as optogenetics is not simply photoexcitation of targeted cells; rather, it delivers gain of function of precise events. The low intensity light illumination would minimize side effects (such as photobleaching) on the illuminated cells or the proximal un-illuminated cells.

Apoptosis is a process of programmed cell death that occurs in multicellular organisms. While it is well established that mitochondria play key roles in activating programmed cell death, the underlying detailed mechanisms remain incompletely understood. For instance, the role of inner membrane permeabilization in the execution of pro-apoptotic protein (e.g. cytochrome C) release from the intermembrane space into cytosol is a subject of ongoing controversy. Some studies show that mitochondrial depolarization and IMM permeabilization are required for cytochrome C release and apoptotic cell death (49). However, others suggest that cytochrome C release can occur in the absence of $\Delta\Psi_m$ depolarization (50), mitochondrial depolarization may not be involved in the activation of apoptosis (51), or PTP opening is a consequence of apoptosis (49). The disparity may be attributed to the methods used in different studies, such as pharmacological intervention versus genetic manipulation. In the present study, using the mitochondrial optogenetics approach we demonstrate that increased IMM permeability alone is sufficient to trigger apoptotic cell death, which is time and (light) dosage dependent. The activation of intrinsic apoptosis is confirmed by induction of cytochrome C release and prevention of cell death by a caspase inhibitor (Fig. 5). Importantly, blocking mPTP did not prevent the light-induced cell death, which provides additional evidence that the expressed ChR2 formed functional cationic pores in IMM. Importantly, we also show that mild short-term light illumination ameliorates apoptotic cell death in ABCB-ChR2-expressing cells, supporting the notion of mitochondrial preconditioning (Fig. 8). Thus, this innovative mitochondrial optogenetics technique can be used to well control IMM depolarization and differentially modulate cell fate, a capability not possessed by the existing pharmacological approaches.

Loss-of-function mutations in Parkin have been causally linked to early-onset Parkinson's disease (52). In addition, studies have shown that Parkin overexpression can prevent cell death in response to a variety of stimuli (53-55). Thus, considerable enthusiasm has emerged recently regarding the potential of Parkin as a therapeutic target. However, here we surprisingly observe that activation of Parkin exacerbates light-induced mitochondrial depolarization and cell death.

While our findings appear contradictory to the conventional notion that Parkin is cytoprotective, the pro-apoptotic effect of Parkin has been reported in two recent studies (36, 37). The apparent paradoxical role of Parkin appears attributed to the nature of the stressor to induce cell death, i.e., whether the stimulus induces mitochondrial depolarization and activation of Parkin. For instance, Carroll *et al.* (36) have shown that Parkin overexpression enhances HeLa cell apoptosis induced by mitochondrial depolarizing agents such as CCCP and valinomycin but has no effects on cell death induced by other stimuli that fail to activate Parkin. Similarly, Zhang *et al.* reported that Parkin activation significantly exacerbates valinomycin-induced apoptosis but does not promote CCCP-induced cell death in HeLa cells (37). They further demonstrated that the level of Parkin recruitment-induced Mcl-1 degradation and polyubiquitination is a key factor determining whether Parkin-dependent mitophagy versus apoptotic cell death is activated. While the reason underlying the discriminate effect of Parkin on the fate of CCCP-treated cells in those two studies requires further investigation, the severity of mitochondrial damage might be a critical factor. If mitochondrial damage is too severe to be repaired by mitophagy, the PINK1/Parkin pathway may promote apoptosis. The effect of Parkin on cell death decisions may also depend the level of its activation: with transient activation inducing mitophagy to repair mitochondrial networks and sustained activation triggering apoptosis to eliminate damaged cells. Our findings warrant consideration when designing Parkin-based therapeutic strategies.

One limitation of the developed mitochondrial optogenetics technique is that although it enables targeted mitochondrial depolarization at high spatial (cellular or subcellular) resolution, the time resolution of light-induced mitochondrial depolarization is much less. Similar to that shown in a recent study (40), optogenetic-induced mitochondrial depolarization takes several minutes to accomplish. Using other rhodopsin protein variants with higher conductance or greater light sensitivity may help to achieve better temporal control of mitochondrial depolarization (56). The lack of mitochondrial repolarization may be partially due to the slow deactivation of ChR2 channels. Tkatch *et al.* (40) showed that optogenetic-induced mitochondrial depolarization can recover slowly in cells expressing the ChR(SSFO) mutant. Alternatively, co-expressing Channelrhodopsin and Bacteriorhodopsin or Archaerhodopsins in the IMM and stimulating cells with alternating blue and yellow light might achieve reversible control of $\Delta\Psi_m$. The energetic states or types of cell may also affect the feasibility of inducing faster or more reversible mitochondrial depolarization. For instance, mitochondrial oscillations can be readily triggered in isolated guinea pig cardiomyocytes (7, 9), but it is much more difficult in isolated mouse cardiomyocytes.

In summary, we have developed and characterized a mitochondrial optogenetics approach to induce targeted mitochondrial depolarization, which provides an alternative to the traditional optical or pharmacological means of modulating MMP and $\Delta\Psi_m$ to understand their roles in determining cell fate. We expect the research tools shown here and the knowledge acquired will facilitate the design of novel optogenetic-based therapeutic strategies for the treatment of diseases involving mitochondrial dysfunction.

Materials and methods

Cell culture. HeLa and H9C2 cells were obtained from ATCC and cultured in Dulbecco's Modified Eagle Medium (Gibco) supplemented with 10% (v/v) Fetal Bovine Serum (Gibco) and 2 mM L-Glutamine (Gibco). Cells were maintained in 5% CO_2 at 37 °C with confluence between 10% and 80%. hiPSC-CMs were reprogrammed from human cardiac fibroblasts as described previously (57) and maintained on Matrigel Membrane Matrix (Thermo Fisher Scientific) in mTeSR™ medium (Stem Cell Technologies) until 75% confluency. CM differentiation was performed by using a small molecule–based protocol as described previously (58, 59). Beating hiPSC-CM began to appear 9-12 days after differentiation was initiated.

Plasmid construction. The ROMK and ABCB10(1-140AA) (named ABCB for simplification) MLS were synthesized by GeneArt (Life Technologies). pCMV-myc-mito vector was purchased from Life Technologies (catalog number V82220), and pLenti-EF1a-hChR2(H134R)-EYFP-WPRE was a gift from Karl Deisseroth (Addgene plasmid # 20942). To construct the CMV-myc-mito based plasmids (e.g. pCMV-mito-ChR2-eYFP), pCMV-myc-mito-GFP vector was digested with SalI and XbaI and ligated with ChR2-eYFP fragment obtained from pLenti-EF1a-hChR2(H134R)-EYFP-WPRE vector *via* PCR. The ROMK based plasmid was constructed by replacing the mito sequence with the ROMMK sequence. The pCAG-ABCB-ChR2-eYFP and pCAG-ABCB-eYFP vectors were constructed using the Gibson Ligation Kit following the manufacturer's instructions. The sequences of constructed vectors were confirmed by DNA sequencing at the University of Alabama at Birmingham Helfin Center Genetics Core.

Adenovirus production. The gene of interest (i.e. CMV-ABCB140-ChR2-YFP-ER) was PCR amplified from vector pCAG-CMV-ABCB-ChR2-eYFP using blunt-end primers. This gene fragment was cloned into the pENTR-D-TOPO vector purchased from Life Technologies (catalog number K2400) to generate the adenovirus entry vector. The entry vector sequence

was confirmed using the M13 Forward and M13 Reverse primers. Adenoviral expression vector was constructed by LR recombination using the ViraPower™ Adenoviral Gateway™ Expression Kit (Life technologies catalog number K4930). To produce adenovirus, the expression vector was digested with PacI, purified, and then transfected into HEK 293A cells pre-seeded in T-flasks using Lipofectamine 2000. Cells were cultured until visible regions of cytopathic effect were observed in approximately 80% of the culture. Cells were then squirted from the flask bottom and lysed via three successive freeze/thaw cycles to release the crude adenovirus. To amplify the adenovirus, 293A cells were infected with the crude adenovirus at a MOI of approximately 5. High titer adenovirus was harvested as described above when the 293A cells were floating or lightly attached to the culture dish, 2-3 days after infection.

Confocal imaging and colocalization analysis. Plasmids containing ChR2-eFYP or eYFP with various MLS (e.g., mito, ROMK and ABCB) were transfected to H9C2 or HeLa cells cultured on a 15mm glass-bottom dish using the Lipofectoamine 3000 transfection reagent. hiPSC-CMs were transduced with the adenovirus containing ABCB-ChR2-eYFP at MOI of 50. 48 hours after transfection/transduction, cells were loaded with MitoTracker Deep Red (250 nM) for 30 minutes. The 515nm argon laser and 635nm LD laser lines of an Olympus FV1000 confocal microscope were used to image eYFP and MitoTracker, respectively. Images were processed offline using ImageJ software (Wayne Rasband, NIH) for colocalization analysis.

Light illumination-induced targeted mitochondrial depolarization. The ABCB-ChR2-eYFP or ABCB-eYFP-expressing H9C2 cells were loaded with mitochondrial membrane potential fluorescent dye TMRM (20 nM) and illuminated with 475nm LED pulses (5 mW/mm^2). The blue light was delivered by a 10X objective lens coupled to a LED driver, which was controlled by a Myopacer stimulator (IonOptix, Westwood, MA). The fluorescence of TMRM during light illumination was recorded using an Olympus FV1000 confocal microscope and analyzed using ImageJ software.

Cell death and cytotoxicity assay. Cells were illuminated by LED light of different duration and irradiance (as indicated in the corresponding figure legends). Right after light treatment cells were detached with 0.25% Trypsin-EDTA, stained with Trypan Blue, and counted using a hemocytometer as described previously (25). The cytotoxicity and cell viability were also measured using the LDH assay and AlamarBlue (AB) assay, respectively. For the LDH assay,

at the end of light illumination, 200 µl cell culture medium was removed and added to a 96 well plate, 100 µl of the reaction mixture was added to each well and the plate was incubated for 30 minutes at room temperature, protected from light. The absorbance was measured at 490nm and 680nm using a microplate reader. LDH activity was calculated following the manufacture's instruction. For the AB assay, 10 µl AlamarBlue reagent was directly added to cells in 100 µl culture medium. Cell cultures were incubated for 4 hours in an incubator (37 $^{\circ}$C, protected from light). The absorbance was measured at 570nm using a spectrometer.

Immunocytochemistry. Cells were fixed in 4% Formaldehyde in PBS and then treated with PBS containing 10% goat serum and 0.3% Triton® X-100 to block non-specific staining. Cells were then stained with the following primary antibodies: ChR2, TOMM20, LC3, or Cytochrome C at a ratio of 1:200. After overnight room temperature incubation, secondary antibodies AlexaFluor 555 and/or AlexaFluor 647 were diluted in 1% BSA, 1% goat serum, 0.3% Triton® X-100 in PBS at a ratio of 1:200, and then added to the cells. Coverslips were mounted on slides and imaged using an Olympus FV1000 confocal microscope.

Parkin translocation assay. HeLa cells were co-transfected with plasmid encoding mCherry-tagged Parkin (60) (Addgene plasmid #23956;) and encoding either ABCB-ChR2-YFP or ABCB-YFP using the Neon Transfection System (Invitrogen) and seeded in 35 mm glass-bottom dishes (MatTek, Ashland, MA). Two days after transfection cells were subjected to LED illumination (0.5 mW/mm^2, 24 hours). Confocal imaging of YFP and mCherry expression and localization was done immediately before treatment, and after 4, 8, and 24 hours of treatment.

Detection of the colocalization of mitochondria and lysosomes. HeLa cells were transduced with ABCB-ChR2-eYFP adenovirus and transfected with mCherry-Parkin plasmid. Forty-eight hours later LED stimulation (0.5 mW/mm^2) of the cells began. After 24 hours of stimulation, cells were stained with 50 nM Lysotracker Deep Red at 37 °C for 30 minutes, followed by three washes. Fluorescent images were observed with confocal microscopy. Green fluorescence (i.e. YFP) represents mitochondria and red (LysoTracker) represents lysosomes. The colocalization of mitochondria and lysosomes indirectly indicates mitophagy. All light stimulation was done at a frequency of 4 Hz with 90 ms pulses. Un-illuminated mock-transfected and ABCB-ChR2-eYFP expressing cell cultures were used as control.

Mitochondrial preconditioning. HeLa cells were transduced with ABCB-ChR2-eYFP adenovirus and transfected with mCherry-Parkin plasmid. Forty-eight hours later cultures were subjected to their respective treatments: no light stimulation for control group, 2 hours of no light stimulation followed by 6 hours of stimulation at 4 mW/mm^2 for stimulation-only group, and 2 hours of stimulation at 0.2 mW/mm^2 followed by 6 hours of stimulation at 4 mW/mm^2 for pre-conditioning group. Cell cultures were then stained with Mitoview 633 (25 nM) at 37 °C for 15 minutes, washed of dye, and then immediately followed by confocal imaging. All light stimulation was done at a frequency of 4 Hz with 90 ms pulses.

Flow Cytometry. Cells were trypsinized and washed twice with PBS before resuspension in staining buffer (10 mM HEPES/NaOH (pH 7.4), 140 mM NaCl, 2.5 mM $CaCl_2$) containing Mitoview 633 (25 nM) and stained following the manufacture's instruction. Cells were then washed twice with PBS + 1% BSA before analysis with a BD LSR II Flow Cytometer (BD Biosciences, San Jose, CA).

Materials. Mouse monoclonal antibodies for GFP, ATP5A, TOM20 and LC3 were purchased from Abcam (Cambridge, MA). Mouse monoclonal ChR2 antibody was obtained from Progen (Heidelberg, Germany). Oregon Green 488 anti-rabbit IgG, MitoSox, TMRM, and MitoTracker Deep Red FM were purchased from Life Technologies (Carlsbad, CA). MitoView 633 was purchased from Biotium (Fremont, CA). All other reagents were from Sigma (St Louis, MO).

Statistics. Analysis and presentation of the data were performed using OriginLab software (Northampton, MA). Comparisons between groups were performed using paired or unpaired 2-tailed Student's t test. Data were considered significantly different at $p < 0.05$ or below. Results are presented as mean $\pm$ SEM.

Materials & Correspondence. All relevant materials that support our experimental findings are available from the corresponding author upon reasonable request.

Acknowledgements

The authors thank Drs. Kah Yong Goh, Jiajia Song and Qince Li for their technical support.

Competing interests

The authors declare no competing interests.

Funding

This work is supported by National Institute of Health (NIH) R21 HL127599 and R01 HL121206 (to L. Zhou) and American Heart Association Predoctoral Fellowship 18PRE34060188 (to P. Ernst).

References

1. Honda, H. M., Korge, P., and Weiss, J. N. (2005) Mitochondria and ischemia/reperfusion injury. *Annals of the New York Academy of Sciences* **1047**, 248-258
2. Wallace, D. C., and Fan, W. (2009) The pathophysiology of mitochondrial disease as modeled in the mouse. *Genes & development* **23**, 1714-1736
3. Desagher, S., and Martinou, J. C. (2000) Mitochondria as the central control point of apoptosis. *Trends Cell Biol.* **10**, 369-377
4. Weisova, P., Anilkumar, U., Ryan, C., Concannon, C. G., Prehn, J. H., and Ward, M. W. (2012) 'Mild mitochondrial uncoupling' induced protection against neuronal excitotoxicity requires AMPK activity. *Biochimica et biophysica acta* **1817**, 744-753
5. Brennan, J. P., Southworth, R., Medina, R. A., Davidson, S. M., Duchen, M. R., and Shattock, M. J. (2006) Mitochondrial uncoupling, with low concentration FCCP, induces ROS-dependent cardioprotection independent of KATP channel activation. *Cardiovascular research* **72**, 313-321
6. Hausenloy, D. J., Maddock, H. L., Baxter, G. F., and Yellon, D. M. (2002) Inhibiting mitochondrial permeability transition pore opening: a new paradigm for myocardial preconditioning? *Cardiovascular research* **55**, 534-543
7. Aon, M. A., Cortassa, S., Marban, E., and O'Rourke, B. (2003) Synchronized whole cell oscillations in mitochondrial metabolism triggered by a local release of reactive oxygen species in cardiac myocytes. *J. Biol. Chem.* **278**, 44735-44744
8. Brady, N. R., Elmore, S. P., van Beek, J. J., Krab, K., Courtoy, P. J., Hue, L., and Westerhoff, H. V. (2004) Coordinated behavior of mitochondria in both space and time: a reactive oxygen species-activated wave of mitochondrial depolarization. *Biophys. J.* **87**, 2022-2034
9. Zhou, L., Aon, M. A., Liu, T., and O'Rourke, B. (2011) Dynamic modulation of Ca(2+) sparks by mitochondrial oscillations in isolated guinea pig cardiomyocytes under oxidative stress. *Journal of molecular and cellular cardiology* **51**, 632-639
10. Goh, K. Y., Qu, J., Hong, H., Liu, T., Dell'Italia, L. J., Wu, Y., O'Rourke, B., and Zhou, L. (2016) Impaired mitochondrial network excitability in failing guinea-pig cardiomyocytes. *Cardiovasc Res* **109**, 79-89
11. Boyden, E. S. (2011) A history of optogenetics: the development of tools for controlling brain circuits with light. *F1000 Biol. Rep.* **3**, 11
12. Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G., and Deisseroth, K. (2005) Millisecond-timescale, genetically targeted optical control of neural activity. *Nat. Neurosci.* **8**, 1263-1268
13. Deisseroth, K. (2011) Optogenetics. *Nat Methods* **8**, 26-29
14. Zhang, F., Wang, L. P., Boyden, E. S., and Deisseroth, K. (2006) Channelrhodopsin-2 and optical control of excitable cells. *Nat Methods* **3**, 785-792
15. Adams, D. S., Tseng, A. S., and Levin, M. (2013) Light-activation of the Archaerhodopsin H(+)-pump reverses age-dependent loss of vertebrate regeneration: sparking system-level controls in vivo. *Biology open* **2**, 306-313
16. Gradinaru, V., Thompson, K. R., and Deisseroth, K. (2008) eNpHR: a Natronomonas halorhodopsin enhanced for optogenetic applications. *Brain Cell Biol.* **36**, 129-139
17. Nagel, G., Szellas, T., Huhn, W., Kateriya, S., Adeishvili, N., Berthold, P., Ollig, D., Hegemann, P., and Bamberg, E. (2003) Channelrhodopsin-2, a directly light-gated cation-selective membrane channel. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 13940-13945
18. Hegemann, P., Gartner, W., and Uhl, R. (1991) All-trans retinal constitutes the functional chromophore in Chlamydomonas rhodopsin. *Biophysical journal* **60**, 1477-1489
19. Rein, M. L., and Deussing, J. M. (2012) The optogenetic (r)evolution. *Molecular genetics and genomics : MGG* **287**, 95-109

20. Arrenberg, A. B., Stainier, D. Y., Baier, H., and Huisken, J. (2010) Optogenetic control of cardiac function. *Science* **330**, 971-974

21. Bruegmann, T., Malan, D., Hesse, M., Beiert, T., Fuegemann, C. J., Fleischmann, B. K., and Sasse, P. (2010) Optogenetic control of heart muscle in vitro and in vivo. *Nat Methods* **7**, 897-900

22. Entcheva, E. (2013) Cardiac Optogenetics. *Am. J. Physiol. Heart Circ. Physiol.*

23. Asano, T., Ishizua, T., and Yawo, H. (2012) Optically controlled contraction of photosensitive skeletal muscle cells. *Biotechnol. Bioeng.* **109**, 199-204

24. Jia, Z., Valiunas, V., Lu, Z., Bien, H., Liu, H., Wang, H. Z., Rosati, B., Brink, P. R., Cohen, I. S., and Entcheva, E. (2011) Stimulating cardiac muscle by light: cardiac optogenetics by cell delivery. *Circulation. Arrhythmia and electrophysiology* **4**, 753-760

25. Li, Q., Ni, R. R., Hong, H., Goh, K. Y., Rossi, M., Fast, V. G., and Zhou, L. (2017) Electrophysiological Properties and Viability of Neonatal Rat Ventricular Myocyte Cultures with Inducible ChR2 Expression. *Sci. Rep.* **7**, 1531

26. Yang, F., Tu, J., Pan, J. Q., Luo, H. L., Liu, Y. H., Wan, J., Zhang, J., Wei, P. F., Jiang, T., Chen, Y. H., and Wang, L. P. (2013) Light-controlled inhibition of malignant glioma by opsin gene transfer. *Cell Death Dis.* **4**, e893

27. Hill, R. A., Damisah, E. C., Chen, F., Kwan, A. C., and Grutzendler, J. (2017) Targeted two-photon chemical apoptotic ablation of defined cell types in vivo. *Nat Commun* **8**, 15837

28. Karbowski, M., Arnoult, D., Chen, H., Chan, D. C., Smith, C. L., and Youle, R. J. (2004) Quantitation of mitochondrial dynamics by photolabeling of individual organelles shows that mitochondrial fusion is blocked during the Bax activation phase of apoptosis. *J. Cell Biol.* **164**, 493-499

29. Kettlewell, S., Cabrero, P., Nicklin, S. A., Dow, J. A., Davies, S., and Smith, G. L. (2009) Changes of intra-mitochondrial Ca2+ in adult ventricular cardiomyocytes examined using a novel fluorescent Ca2+ indicator targeted to mitochondria. *J. Mol. Cell. Cardiol.* **46**, 891-901

30. Nagel, G., Brauner, M., Liewald, J. F., Adeishvili, N., Bamberg, E., and Gottschalk, A. (2005) Light activation of channelrhodopsin-2 in excitable cells of Caenorhabditis elegans triggers rapid behavioral responses. *Curr. Biol.* **15**, 2279-2284

31. Foster, D. B., Ho, A. S., Rucker, J., Garlid, A. O., Chen, L., Sidor, A., Garlid, K. D., and O'Rourke, B. (2012) Mitochondrial ROMK channel is a molecular component of mitoK(ATP). *Circ. Res.* **111**, 446-454

32. Higgins, C. F. (1992) ABC transporters: from microorganisms to man. *Annu. Rev. Cell Biol.* **8**, 67-113

33. Higgins, C. F. (2001) ABC transporters: physiology, structure and mechanism--an overview. *Res. Microbiol.* **152**, 205-210

34. Graf, S. A., Haigh, S. E., Corson, E. D., and Shirihai, O. S. (2004) Targeting, import, and dimerization of a mammalian mitochondrial ATP binding cassette (ABC) transporter, ABCB10 (ABC-me). *J. Biol. Chem.* **279**, 42954-42963

35. Miyazaki, E., Kida, Y., Mihara, K., and Sakaguchi, M. (2005) Switching the sorting mode of membrane proteins from cotranslational endoplasmic reticulum targeting to posttranslational mitochondrial import. *Mol. Biol. Cell* **16**, 1788-1799

36. Carroll, R. G., Hollville, E., and Martin, S. J. (2014) Parkin sensitizes toward apoptosis induced by mitochondrial depolarization through promoting degradation of Mcl-1. *Cell Rep* **9**, 1538-1553

37. Zhang, C., Lee, S., Peng, Y., Bunker, E., Giaime, E., Shen, J., Zhou, Z., and Liu, X. (2014) PINK1 triggers autocatalytic activation of Parkin to specify cell fate decisions. *Current biology : CB* **24**, 1854-1865

38. Ichas, F., Jouaville, L. S., and Mazat, J. P. (1997) Mitochondria are excitable organelles capable of generating and conveying electrical and calcium signals. *Cell* **89**, 1145-1153

39. Zhou, L., and O'Rourke, B. (2012) Cardiac mitochondrial network excitability: insights from computational analysis. *Am J Physiol Heart Circ Physiol* **302**, H2178-2189

40. Tkatch, T., Greotti, E., Baranauskas, G., Pendin, D., Roy, S., Nita, L. I., Wettmarshausen, J., Prigge, M., Yizhar, O., Shirihai, O. S., Fishman, D., Hershfinkel, M., Fleidervish, I. A., Perocchi, F., Pozzan, T., and Sekler, I. (2017) Optogenetic control of mitochondrial metabolism and Ca2+ signaling by mitochondria-targeted opsins. *Proceedings of the National Academy of Sciences* **114**, E5167-E5176

41. Ernst, P. X., N; Goh, K; O;Rourke, B; Zhang, J.J; Liu, X.M; Zhou, L. (2017) Optogenetic-induced mitochondrial membrane potential depolarization and targeting cell death. In *AHA Scientific Sessions*, Anaheim, CA

42. Ashrafi, G., Schlehe, J. S., LaVoie, M. J., and Schwarz, T. L. (2014) Mitophagy of damaged mitochondria occurs locally in distal neuronal axons and requires PINK1 and Parkin. *The Journal of cell biology* **206**, 655-670

43. Cai, Q., Zakaria, H. M., Simone, A., and Sheng, Z. H. (2012) Spatial parkin translocation and degradation of damaged mitochondria via mitophagy in live cortical neurons. *Current biology : CB* **22**, 545-552

44. Burke, R. E., and O'Malley, K. (2013) Axon degeneration in Parkinson's disease. *Exp Neurol* **246**, 72-83

45. Hollenbeck, P. J., Bray, D., and Adams, R. J. (1985) Effects of the uncoupling agents FCCP and CCCP on the saltatory movements of cytoplasmic organelles. *Cell Biol Int Rep* **9**, 193-199

46. Berezhnov, A. V., Soutar, M. P., Fedotova, E. I., Frolova, M. S., Plun-Favreau, H., Zinchenko, V. P., and Abramov, A. Y. (2016) Intracellular pH Modulates Autophagy and Mitophagy. *The Journal of biological chemistry* **291**, 8701-8708

47. Doebler, J. A. (2000) Effects of protonophores on membrane electrical characteristics in NG108-15 cells. *Neurochem Res* **25**, 263-268

48. To, M. S., Aromataris, E. C., Castro, J., Roberts, M. L., Barritt, G. J., and Rychkov, G. Y. (2010) Mitochondrial uncoupler FCCP activates proton conductance but does not block store-operated Ca(2+) current in liver cells. *Arch Biochem Biophys* **495**, 152-158

49. Baines, C. P., Kaiser, R. A., Sheiko, T., Craigen, W. J., and Molkentin, J. D. (2007) Voltage-dependent anion channels are dispensable for mitochondrial-dependent cell death. *Nat Cell Biol* **9**, 550-555

50. Kim, C. N., Wang, X., Huang, Y., Ibrado, A. M., Liu, L., Fang, G., and Bhalla, K. (1997) Overexpression of Bcl-X(L) inhibits Ara-C-induced mitochondrial loss of cytochrome c and other perturbations that activate the molecular cascade of apoptosis. *Cancer Res* **57**, 3115-3120

51. Wang, C., and Youle, R. J. (2009) The role of mitochondria in apoptosis*. *Annu Rev Genet* **43**, 95-118

52. Kitada, T., Asakawa, S., Hattori, N., Matsumine, H., Yamamura, Y., Minoshima, S., Yokochi, M., Mizuno, Y., and Shimizu, N. (1998) Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* **392**, 605-608

53. Dai, H., Deng, Y., Zhang, J., Han, H., Zhao, M., Li, Y., Zhang, C., Tian, J., Bing, G., and Zhao, L. (2015) PINK1/Parkin-mediated mitophagy alleviates chlorpyrifos-induced apoptosis in SH-SY5Y cells. *Toxicology* **334**, 72-80

54. Zhao, C., Chen, Z., Xu, X., An, X., Duan, S., Huang, Z., Zhang, C., Wu, L., Zhang, B., Zhang, A., Xing, C., and Yuan, Y. (2017) Pink1/Parkin-mediated mitophagy play a protective role in cisplatin induced renal tubular epithelial cells injury. *Exp Cell Res* **350**, 390-397

55. Zhang, H. T., Mi, L., Wang, T., Yuan, L., Li, X. H., Dong, L. S., Zhao, P., Fu, J. L., Yao, B. Y., and Zhou, Z. C. (2016) PINK1/Parkin-mediated mitophagy play a protective role in manganese induced apoptosis in SH-SY5Y cells. *Toxicol In Vitro* **34**, 212-219

56. Yizhar, O., Fenno, L. E., Prigge, M., Schneider, F., Davidson, T. J., O'Shea, D. J., Sohal, V. S., Goshen, I., Finkelstein, J., Paz, J. T., Stehfest, K., Fudim, R., Ramakrishnan, C., Huguenard, J. R., Hegemann, P., and Deisseroth, K. (2011) Neocortical excitation/inhibition balance in information processing and social dysfunction. *Nature* **477**, 171-178

57. Zhang, L., Guo, J., Zhang, P., Xiong, Q., Wu, S. C., Xia, L., Roy, S. S., Tolar, J., O'Connell, T. D., Kyba, M., Liao, K., and Zhang, J. (2015) Derivation and high engraftment of patient-specific cardiomyocyte sheet using induced pluripotent stem cells generated from adult cardiac fibroblast. *Circ Heart Fail* **8**, 156-166

58. Zhu, W., Gao, L., and Zhang, J. (2017) Pluripotent Stem Cell Derived Cardiac Cells for Myocardial Repair. *J Vis Exp*

59. Gao, L., Kupfer, M. E., Jung, J. P., Yang, L., Zhang, P., Da Sie, Y., Tran, Q., Ajeti, V., Freeman, B. T., Fast, V. G., Campagnola, P. J., Ogle, B. M., and Zhang, J. (2017) Myocardial Tissue Engineering With Cells Derived From Human-Induced Pluripotent Stem Cells and a Native-Like, High-Resolution, 3-Dimensionally Printed Scaffold. *Circ Res* **120**, 1318-1325

60. Narendra, D., Tanaka, A., Suen, D. F., and Youle, R. J. (2008) Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. *The Journal of cell biology* **183**, 795-803

Figure Legends

Figure 1. **Representative confocal images showing intracellular localization of ChR2-eYFP or eYFP in different cell types.** A): H9C2 cells expressing ABCB-eYFP; B): H9C2 cells expressing ABCB-ChR2-eYFP; C): Immunoblotting of H9C2 cells expressing ABCB-ChR2-eFYP with ChR2 antibody; D): HeLa cells expressing ABCB-ChR2-eYFP; and E): hiPSC-CMs expressing ABCB-ChR2-eYFP. In all experiments cells were stained with MitoTracker Deep Red (250 nM). Yellow indicates colocalization of YFP (in A), ChR2-eYFP (in B, D, and E) or ChR2 (in C) and the mitochondrial marker.

Figure 2. **Mitochondrial-targeted optogenetics induces selective mitochondrial depolarization in H9C2 cells expressing ABCB-ChR2-eYFP.** A): Schematic figure showing that cells in zone 1 were illuminated by LED (5 mW/mm^2) and cells in zone 2 were not exposed to illumination; B) and C): Confocal images showing mitochondrial membrane potential ($\Delta\Psi_m$), measured by TMRM fluorescent dye (20 nM), of cells in zone 1 and zone 2 before (B) and after (C) light illumination; D): Normalized $\Delta\Psi_m$ in cells in the un-illuminated (i.e. zone 1) and illuminated (i.e. zone 2) zones before and after light illumination. 2-4 cells from each zone were analyzed in each cell culture and four cell cultures were examined.

Figure 3. **Mitochondrial-targeted optogenetics induces cell death in HeLa cells expressing ABCB-ChR2-eYFP.** A): Representative transmitted light images of ABCB-ChR2-eYFP or ABCB-eYFP-expressing cells with or without LED pulse illumination (0.5 mW/mm^2, 24 hours); B) and C): Comparison of cell viability measured by LDH assay (B) and cytotoxicity measured by alamarBlue assay (C) in ABCB-ChR2-eYFP or ABCB-eYFP cells with or without LED illumination; D) and E): Effect of a mPTP blocker, CsA, on cell viability (D) and cytotoxicity (E) in ABCB-ChR2-eYFP-expressing or mock-transfected cell cultures with or without LED illumination. n represents the number of cell cultures.

Figure 4. **Time and light irradiance dependence of mitochondrial optogenetic-mediated cell death.** A): Time dependence of moderate LED illumination (0.5 mW/mm^2)-induced cell death in ABCB-ChR2-eYFP-expressing HeLa cell cultures. Four cell cultures were repeated at each time point. Cell viability was determined using the trypan blue method; B): LED irradiance dependence of illumination-induced cell death. ABCB-ChR2-eYFP or mock-transfected HeLa cultures were illuminated by LED of various intensity for six hours. At least four cell cultures

were repeated for each group. Cell death were measured by Propidium iodide (PI) staining; C): Representative images of PI staining in cultures illuminated by LED of different irradiance summarized in (B).

Figure 5. Mitochondrial optogenetics induces cell death *via* apoptotic pathways. A): Z-VAD-FMK, a pan caspase inhibitor, substantially ameliorated light-elicited cell death in HeLa cell cultures expressing ABCB-ChR2. 7-Cl-O-Nec-1, a necroptosis inhibitor, had no significant effect on the light-induced cell death. n represents the numbers of cell cultures; B): Z-VAD-FMK alleviated light-induced mitochondrial depolarization in ABCB-ChR2-expressing cells; and C): Light illumination (0.5 mW/mm^2, 24 hours) induced cytochrome C release into cytosol in ABCB-ChR2-expressing cells.

Figure 6. **Light illumination induces mitochondrial autophagy in ABCB-ChR2-eYFP-expressing cells.** A): Representative confocal imagines showing time-dependent localization of Parkin-mCherry in light illuminated cells expressing ABCB-ChR2-eYFP and Parkin-mCherry. Images were taken at 0, 4, 8, and 24 hours during light illumination (0.5 mW/mm^2). Red represents Parkin-mCherry, green represents eYFP, and overlay represents Parkin accumulation in mitochondria; B): Representative confocal imagines showing LC3 (red) aggregation and colocalization with mitochondria (represented by YFP) in illuminated (0.5 mW/mm^2, 24 hours) ABCB-ChR2-eYFP expressing cells; and C): Representative confocal imagines showing lysosomes (red, represented by LysoTracker) aggregation and colocalization with mitochondria (represented by YFP) in illuminated (0.5 mW/mm^2, 24 hours) ABCB-ChR2-eYFP expressing cells.

Figure 7. Parkin activation exacerbates mitochondrial optogenetic-mediated cell death. A): Flow cytometry showing mitochondrial membrane potential (measured by MitoView) in the illuminated (0.5 mW/mm^2, 24 hours) or un-illuminated HeLa cells with or without Parkin overexpression; and B): Parkin overexpression and activation did not influence HeLa cell viability right after light illumination (0.5 mW/mm^2, 24 hours) but promoted cell death 24 hours later. 5 cell cultures were examined for each group.

Figure 8. Optogenetic-mediated mitochondrial preconditioning protects cells against apoptotic cell death. A): Representative images of Hoechst and Propidium Iodide (PI) staining in cultures illuminated by LED of moderate irradiance (4 mW/mm^2 for 6 hours); B)

Representative images of Hoechst and PI staining in cultures illuminated with mild (0.2 mW/mm^2 for 2 hours) LED followed by moderate (4 mW/mm^2 for 6 hours) LED illumination; C): Summary of cell viability in the unilluminated, illuminated only, and preconditioned groups; D-F): Representative confocal images showing MitoView fluorescence in the un-illuminated (D), illuminated only (E) and preconditioned and illuminated (F) cell cultures; and G): Normalized mitochondrial membrane potential ($\Delta\Psi_m$, measured by MitoView) in different groups. n=3 each group.