Cell-type-specific genomics reveals histone modification

dynamics in mammalian meiosis

Authors

Kwan-Wood Gabriel Lam, Kevin Brick, Gang Cheng, Florencia Pratto and R.

Daniel Camerini-Otero*

Affiliations

Genetics and Biochemistry Branch, National Institute of Diabetes, Digestive and

Kidney Diseases, National Institutes of Health, Bethesda, MD, USA

*Corresponding author

Abstract

Meiosis is the specialized cell division during which parental genomes recombine to create genotypically unique gametes. Despite its importance, mammalian meiosis cannot be studied *in vitro,* greatly limiting mechanistic studies*. In vivo,* meiocytes progress asynchronously through meiosis and therefore the study of specific stages of meiosis is a challenge. Here, we describe a method for isolating pure sub-populations of nuclei that allows for detailed study of meiotic sub-stages. Interrogating the H3K4me3 landscape revealed dynamic chromatin transitions between sub-stages of meiotic prophase I, both at sites of genetic recombination and at gene promoters. We also leveraged this method to perform the first comprehensive, genome-wide survey of histone marks in meiotic prophase, revealing a heretofore unappreciated complexity of the epigenetic landscape at meiotic recombination hotspots. Ultimately, this study presents a straightforward, scalable framework for interrogating the complexities of mammalian meiosis.

Main

Meiosis is the specialized cell division required to form gametes. In meiosis the DNA of a parent cell is replicated, then haploid gametes are produced through two successive cell divisions. Distinct mechanisms are required to assure that chromosomes segregate accurately in each division and these events take place through a staged cascade of nuclear events. Unique to meiosis is the formation of programmed DNA double-strand breaks (DSBs) that result in recombination between parental haplotypes and whose repair tethers homologous chromosomes at the first meiotic division. Meiotic DSB formation, repair and recombination occur in a continuum of sub-stages (leptonema, zygonema, pachynema, and diplonema) known collectively as meiotic prophase I (MPI), and each stage is characterized by specific nuclear events[1]. Despite its importance, technical challenges preclude the isolation and study of specific meiotic sub-stages and many of the molecular mechanisms of meiosis remain poorly studied in higher eukaryotes. This is largely because mammalian meiosis cannot be adequately recapitulated *in vitro*[2]. *In vivo*, meiosis either occurs in a difficult-to-isolate niche, such as the fetal ovary, or in an asynchronously dividing tissue such as in adult testis.

Extant methods to study sub-stages of meiotic prophase have been dominated by the use of crude enrichment strategies. One common approach has been to study spermatocytes in juvenile mice, where the first wave of meiosis progresses relatively synchronously[3,4]. Nonetheless, most cells in juvenile testis are not at the stage of interest because this synchrony is not absolute, and there are many non-meiotic cells. Furthermore, differences between the first and the subsequent rounds of spermatogenesis[5–7] may preclude generalization to adults. Chemically induced synchronization of meiosis has recently been developed[8], however the application is technically cumbersome and the consequences of chemical treatment have not been explored. An alternative to enrichment is the isolation of meiotic sub-stages. Fluorescence-activated cell sorting (FACS) or sedimentation based methods have been used to isolate particular sub-stages[9–11]. However,

since these methods rely on imprecise metrics such as cell size and chromatin

content, the isolation and distinction between many sub-stages is challenging.

Thus, isolation-based approaches have not been broadly adapted to the study of

the dynamic molecular events during meiosis.

We have devised a straightforward, yet highly specific strategy to isolate pure sub-

populations of meiotic nuclei. This approach uses a variant of FACS[12] in which

antibodies to intra-nuclear markers are used to select nuclei of interest. The

proteins present in different meiotic sub-stages are well studied[13–16] and we

leverage this knowledge to isolate nuclei in which specific combinations of proteins

are expressed. This allowed for the isolation of meiotic sub-populations ranging

from 74-96% purity. Nuclei are fixed before sorting, which preserves *in vivo*

interactions and prevents degradation during sample processing. Importantly, this

fixation also means our method is the first to allow for downstream interrogation of

genome-wide protein-DNA interactions in meiotic sub-stages.

Although comprehensively studied in somatic cells, little is known about histone

modification dynamics in meiotic cells. A case in point are the histone modifications

that precede meiotic DSB formation, an early event in meiotic recombination.

Programmed DSBs are targeted to sites defined by DNA binding and subsequent

H3K4/K36-trimethylation by the PRDM9 protein[17,18]. PRDM9 gene expression[19],

nuclear protein levels of PRDM9[20] and microscopy-based estimates of DSB

formation[21,22] all imply that PRDM9 marks sites for DSB formation at the onset of

MPI.

Using chromatin immunoprecipitation followed by high throughput sequencing

(ChIP-Seq) on sorted populations of nuclei, we determined that Histone H3 Lysine

4 trimethylation (H3K4me3) varies from stage-to-stage at meiotic DSB hotspots,

gene promoters and a vast number of heretofore unannotated sites. These are the

first genome-wide analyses of the dynamic chromatin landscape in meiosis. Also,

we found that the histone code at DSB hotspots is more complex than has been

previously appreciated[17,18], with multiple histone tail modifications exhibiting

enrichment. This includes histone acetylation marks unlikely to have been

deposited by the DSB-defining histone methyltransferase PRDM9.

Together, these data demonstrate the simplicity, utility and feasibility of our

approach. This method opens the door to a far better understanding of the

molecular mechanisms underlying meiosis and other complex developmental

processes.

Results

Isolation of stage-specific spermatocyte nuclei

We first devised a discriminative panel of intra-nuclear proteins to distinguish

between five, classically-defined MPI sub-stages; leptonema, zygonema, early

pachynema, late pachynema and diplonema (Fig. 1). Nuclei were extracted from

formaldehyde-fixed testes from adult male mice, then incubated with antibodies

against the desired proteins (see methods). Nuclei were also stained with DAPI to

allow measurement of DNA content. The combinatorial signals from these

fluorescent markers were subsequently used to isolate specific sub-populations of

spermatocyte nuclei using FACS (Fig. 2a and b; see methods).

We used the meiosis-specific protein SCP3 to distinguish post-replicative meiotic

prophase nuclei (4C) from fully replicated mitotic nuclei (also 4C). We then

combined SCP3 with other markers for FACS. Our combination of antibodies

required two separate sorts; one to isolate leptotene nuclei and a second to isolate

the other four sub-stages (Fig. 2a and b). In male mice, STRA8 regulates the entry

to meiotic S phase, and is expressed from primitive spermatogonia to leptotene

spermatocytes[16]. We therefore used STRA8 to isolate leptotene nuclei from later

stages. To isolate the four other meiotic stages, we used the presence or absence

of H1t and quantitative differences in SCP1 intensity (Fig. 1, Fig. 2a). H1t (testis-

specific histone　H1) is present from mid pachynema through MPI[23], whereas

SCP1 is a marker of progressive chromosome synapsis from zygotene to
pachytene stage and the SCP1 signal is diminished in diplonema when
chromosomes desynapse[14] (Fig. 1). Thus, we collected seven populations,
assessed the purity of each population and determined the gating thresholds for
the four meiotic sub-stages (Fig. 2b and c).

The purity of each sorted population was assessed using immunofluorescence
microscopy to stage the nuclei according to the known localization of markers (Fig.
1, Fig. 2a and c, Supplementary Fig. 1, see methods)[13–16]. The purity of the sorted
leptotene population was 94.6% - 97.9% (n = 2), a >50-fold increase in purity
compared to the percentage of leptotene cells in whole testis (1.6% - 1.8% (n = 2),
Fig. 2c). The purity of the other four populations ranged from 73.9 - 90.4%, each
more than 20-fold enriched relative to the starting population from whole testis (Fig.
2c). The least pure sample was that of zygotene (71.3% - 78.7% (n = 3)), likely
because we rely entirely on quantitative differences in SCP1 to distinguish
between adjacent stages (Fig. 1, Fig. 2b and c). Narrowing of the gate can increase
purity but at the cost of yield (data not shown). Quantitative thresholding issues
also explain the 9.0% - 16.9% (n = 3) of zygotene nuclei found in the sorted early
pachytene population (Fig. 1, Fig. 2b and c).

From a single adult mouse, we obtained between 115,000 and 400,000 nuclei for
each meiotic sub-stage (Fig. 2d). This range broadly reflects differences in the
proportion of each sub-stage in the adult testis, where leptotene and zygotene
nuclei are less abundant. Thus, the different sub-stages are isolated in sufficient
numbers for experiments using standard genomics protocols[24].

Stage-specific ChIP-Seq reveals histone modification dynamics in meiosis
We next used ChIP-Seq to examine the dynamics of H3K4me3 in MPI. H3K4me3
is of key biological interest in meiosis because it regulates multiple independent
dynamic processes during MPI; H3K4me3 is found at gene promoters and its
presence correlates with active transcription[25,26], whereas PRDM9-mediated

H3K4me3 marks the future sites of meiotic DSBs genome-wide[27,28]. We therefore

performed H3K4me3 ChIP-Seq in the five aforementioned sorted populations to

assess the temporal dynamics of this histone mark in MPI.

We performed ChIP-Seq for H3K4me3 using a highly-specific antibody

(EpiCypher, 13-0028) (Supplementary Fig. 2). To allow for quantitative cross-

comparison of H3K4me3 signals from samples of differing quality, we used a

spike-in control containing H3K4me3-modified nucleosomes[29] (see methods). We

observed a highly dynamic H3K4me3 signal at hotspots in MPI; the H3K4me3

signal first appears at leptonema, and is maximal at zygonema, and gone by early

pachynema (Fig. 3a and b). The maximal signal at zygonema was somewhat

surprising because DSBs are formed and nuclear PRDM9 protein levels are

maximal at leptonema[20]. Intriguingly, at strong hotspots, H3K4me3 is relatively

weak at zygonema compared to leptonema (Supplementary Fig. 3). This implies

that DSB formation or repair removes H3K4me3-modified nucleosomes, similar to

findings in *Saccharomyces cerevisae*[30].

Most H3K4me3 ChIP-Seq studies, including all studies of H3K4me3 in meiosis

have used antibodies that extensively cross-react with H3K4me2[31]

(Supplementary Fig. 2). To allow cross-comparison with published data we

performed ChIP-Seq with the most commonly used antibody (Abcam ab8580; [31])

and observed a broadly similar dynamic of histone modification at hotspots as

compared to the specific antibody (Supplementary Fig. 4); the hotspot signal is still

maximal at zygotene, but the non-specific antibody gives a higher leptotene signal

and a residual signal in early pachynema. We hypothesize that these differences

reflect dynamic changes in H3K4 dimethylation at hotspots.

The H3K4me3 level at gene promoters is correlated with gene expression in mitotic

cells[26]. We therefore investigated if the temporal dynamics of H3K4me3 at

transcription start sites (TSSs) is predictive of meiotic gene expression patterns.

Stage-specific gene expression in MPI has been deduced from classical cell

sorting of meiotic cells followed by RNA-Seq[11,32]. We found that the H3K4me3

profiles at TSSs were positively correlated with gene expression through MPI

(Spearman's R = 0.27; see methods). This correlation was highly significant, far

higher than the maximum correlation observed in randomized comparisons

(empirical P < 0.0001; N = 10,000 bootstraps, R_{max} = 0.03, see methods). To

explore the relationship between H3K4me3 and gene expression in more detail,

we clustered transcripts based on the temporal H3K4me3 pattern (Fig. 3c; optimal

clusters = 5; k-means clustering using gap statistic[33]). There was significant

correlation with gene expression in all clusters (P < 0.0001; N = 10,000 bootstraps,

range; 0.12 ≤ R ≤ 0.44) (Fig. 3c). Each cluster is composed of a large fraction of

transcripts where H3K4me3 and gene expression correlate very well

(Supplementary Fig. 5). Nonetheless, there are also many transcripts for which

H3K4me3 and expression are not correlated. For instance, in cluster 1, many

genes have higher gene expression late in MPI than we would predict from

H3K4me3 data (far right panels; Supplementary Fig. 5). This may be explained by

mRNA accumulation through MPI or by H3K4me3 in early MPI marking poised,

but not yet active promoters[34]. In contrast, genes in cluster 5 are less affected by

these confounding effects and the correlation is far higher for this cluster. This

seems to indicate a concerted wave of gene activation and expression as cells

approach the end of MPI. Together, these data demonstrate that ChIP-Seq of

H3K4me3 in stage-specific nuclei reveals a dynamic that parallels that of gene

expression. These data also reveal an intricate and poorly understood interplay

between H3K4me3 and gene expression during MPI.

HIstone modification dynamics at other genomic elements

Dynamic H3K4-trimethylation is observed at both gene promoters and DSB

hotspots in MPI. However, 16% of H3K4me3 peaks (4,664/28,528) are outside of

these sites (Supplementary Fig. 6a). These unannotated H3K4me3 peaks may

represent enhancers, cryptic promoters or other functional elements. To explore

these dynamics in greater detail, we used the H3K4me2/3 antibody (Abcam

ab8580), as this should capture H3K4me2 modifications that mark enhancers.

Indeed, using this antibody, 18,738/53,518 (35%) peaks occur at non-hotspot, non-

TSS sites (Supplementary Fig. 7a). Using peaks from this antibody, we derived a

MPI profile for each peak across the five MPI stages, then performed unbiased k-

means clustering. To validate this approach, we examined clustering of DSB

hotspots; as expected, hotspots primarily occur in a single large cluster with a

maximum signal at zygonema (cluster 3; Supplementary Fig. 7b-d; 89% of

hotspots occur in this cluster). 11% of unannotated peaks also occur in this cluster

(cluster 3; Supplementary Fig. 7d) and may represent sites of PRDM9 binding that

were not detected in hotspot mapping experiments. Alternatively, we cannot rule

out that a cluster of cryptic functional sites simply exhibit a similar dynamic. The

remaining unannotated peaks exhibit an MPI dynamic that broadly mirrors that at

TSS (Supplementary Fig. 7b, d). It seems likely therefore that functional elements

at these sites play a role in regulating meiotic progression. These dynamic, but

unannotated sites represent a completely unexplored aspect of the regulation of

mammalian meiosis.

Clustering also allowed us to study H3K4me3 dynamics at sites involved in other

important processes during MPI. In particular, we examined H3K4me3 sites that

are used for DSB targeting in mice that lack functional PRDM9[27,28]. H3K4me3

peaks used as "default" hotspot locations were predominantly those with maximal

signal in early MPI (Supplementary Fig. 6 and 7).

Identifying other histone marks at hotspots

At DSB hotspots defined by PRDM9, H3K4me3 is necessary for DSB formation[35].

Nonetheless, H3K4me3 alone is not sufficient to define DSB sites, as this histone

mark is also present at other functional sites, such as gene promoters. In mice,

H3K36me3 is the only other histone mark described genome-wide at DSB

hotspots[17], however other histone marks may also be involved in defining DSB

sites[36–38]. We therefore used our method to isolate target nuclei from adult mice

for a systematic survey of multiple histone modifications.

We performed ChIP-Seq using antibodies against 16 additional histone
methylation and acetylation marks in isolated nuclei (Fig. 4a and b). These marks
were either previously analyzed at individual hotspots in mice[36,38] or reported to be
enriched at DSB sites in other organisms[39,40]. In this exploratory phase, we did not
perform ChIP-Seq in all five sub-populations, but instead, in a combined
population that includes all stages with potential relevance to understanding
histone modifications at DSB sites (leptonema to early pachynema; SCP3+ and
H1t− population).

H4K8ac, H4K12ac, H4K20me3, H3K4ac, H3K79me1, H3K79me3, H3K27me1,
H3K9me2, H3K9me3 and H3K27me3 were not enriched at hotspots relative to
controls (Fig. 4a, b; Supplementary Fig. 8). With the exceptions of H3K27me1, and
H3K9me2, these experiments showed enrichment at the expected functional
genomic regions (Supplementary Fig. 9), lending confidence the lack of
enrichment at hotspots does not result from failed ChIP experiments. Nonetheless,
transient or weak signals at hotspots could still be missed.

H3K4me1, H3K27ac and H4ac5 showed marginal enrichment at hotspots (Fig. 4a,
b; Supplementary Fig. 8). The weak signals at hotspots are unlikely the result of
sub-optimal ChIP-quality, as strong signals are seen at gene promoters and
enhancers (Supplementary Fig. 9). Instead, these marks may be rapidly turned-
over or present only in a sub-population of early spermatocytes. Indeed, a weak
DSB-dependent H4ac5 signal was previously reported at two mouse hotspots[36].

PRDM9 can trimethylate both H3K4 and H3K36[35] and as expected, H3K4me3 and
H3K36me3 were both enriched at hotspots (Fig. 4a and b; Supplementary Fig. 8).
H3K4me2 and H3K9ac were also strongly enriched at DSB hotspots genome-
wide; this clearly resolves the ambiguity from previous studies where these marks
were enriched at one but not another hotspot[36]. H3K4me2 is a likely intermediate
of PRDM9 H3K4-trimethylation (see above), however since PRDM9 lacks a
histone acetyltransferase domain, it is unlikely that PRDM9 directly acetylates

H3K9. In stage-specific experiments, the dynamics of H3K9ac through MPI
resemble those of H3K4me3: a signal is first observed at hotspots in leptonema,
maximal at zygonema and absent by early pachynema (Fig. 4c). Thus, H3K9ac is
a *bona-fide* marker of meiotic DSB hotspots genome-wide, explicitly demonstrating
for the first time that proteins other than PRDM9 modify nucleosomes at the sites
of DSB hotspots.

H3K4me3 is positively correlated with DSB frequency[28,41], but most variation in
hotspot strength is not accounted for by changes in H3K4me3 (Spearman R^2 =
0.46) (Supplementary Fig. 10a). H3K36me3 was previously reported to slightly
improve the correlation between H3K4me3 and DSB strength[42], therefore we tried
to better predict hotspot strength using a combination of histone marks that are
enriched at DSB hotspots. Multiple linear regression with all histone marks only
slightly improved the correlation with hotspot strength (max R^2 = 0.52;
Supplementary Fig. 10b; see methods) suggesting that these extra histone marks
(including H3K36me3 and H3K9ac) do not offer an independent readout of DSB
usage as compared to H3K4me3.

A histone code distinguishes DSB hotspots from PRDM9-independent
H3K4me3 sites
In the absence of PRDM9, DSB hotspots occur at sites of non-PRDM9-mediated
H3K4me3[28]. Nonetheless, in *wild-type* mice, PRDM9-defined H3K4me3 sites are
used. In agreement with previous results[17,18], we found that H3K36me3 is a potent
discriminator between DSB hotspots and other H3K4me3-marked sites in the
genome (Fig. 4d). Among the other histones, H3K4me2 can best distinguish
hotspots (Fig. 4d; Supplementary Fig. 11). We excluded histone marks made by
PRDM9 (H3K4me3 and H3K36me3), then performed principal component
analysis to explore if any combination(s) of histone marks better define DSB
hotspots. The most discriminative PC (PC1) captured signal from H3K9ac and
other histone acetylation marks (Fig. 4e), but did not discriminate hotspots as well
as H3K4me2 or H3K36me3 alone. These data suggest that H3K36me3, H3K4me2

and to a lesser extent H3K9ac and other histone acetylation marks, distinguish the
chromatin at sites of PRDM9-marked DSB hotspots from that at other functional
sites where H3K4 is tri-methylated.

Discussion

In this study, we developed a method to isolate highly pure populations of meiotic
nuclei from whole testis. This method is rapid, requires very little starting material,
and resolves a major hurdle to studying meiosis in mammals. We have
demonstrated that we can efficiently sort five populations of MPI nuclei, however,
in principle, any number of specific stages can be purified if the requisite antibodies
are available. We obtained populations of up to 96% purity for a given meiotic sub-
stage, negating the need to artificially synchronize meiosis using chemicals or the
need to use juvenile mice to obtain enriched populations of meiocytes. This opens
the door to many detailed studies of meiosis and also to experiments in
challenging-to-breed knockout mice and importantly in humans, where other
synchronization strategies are impractical.

Importantly, fixation prior to sorting allows for genome-wide interrogation of
transient protein-DNA interactions in sorted populations. To this end, we profiled
H3K4me3 using ChIP-Seq in five MPI populations, capturing extensive dynamics
of this histone mark. The H3K4me3 marks made at DSB hotspots by PRDM9
appear in leptonema, are maximal in zygonema, and are removed by early
pachynema. Intriguingly, the presence of robust H3K4me3 signal at zygonema
implies that PRDM9 trimethylates more nucleosomes than are used during DSB
formation. An alternate explanation would require a large fraction of DSBs to form
in zygonema. This is unlikely because homolog synapsis, which occurs in
zygonema, begins to shut down the DSB machinery in yeast[43]. An analogous
system appears to operate in mammals[44,45]. These are the first data to
demonstrate excess H3K4-trimethylation by PRDM9 in *wild-type* mice, however

recent work in mice co-expressing both endogenous and transgenic PRDM9

reached a similar conclusion[35].

H3K4me3 at gene promoters is also dynamic through MPI. By comparing

H3K4me3 at promoters with gene expression, we found that transcript levels are

broadly predicted by H3K4me3. The most cohesive group of transcripts exhibit a

coordinate increase in H3K4me3 and in expression late in MPI. This likely reflects

the resumption of extensive transcription during mid-pachytene and its continued

increase into diplotene[46]. A large subset of transcripts exhibit gene expression that

appears decoupled from H3K4me3 levels. Some such transcripts exhibit a strong

H3K4me3 signal at the TSS in early MPI, but expression is maximal late in MPI;

this may result from accumulation of long-lived mRNA and/or from H3K4me3

marks at the sites of poised RNA polymerase[34]. Indeed, these transcripts are

intriguing candidates for further understanding gene regulation in MPI. Because

we isolate only nuclei, cytoplasmic mRNA cannot be sequenced in our sorted

populations. We therefore used mRNA-Seq data from a previously published

study[32]. However, we anticipate that recent developments in transcriptomics using

RNA from formaldehyde-fixed nuclei[47] will enable us to adapt our method to study

transcription in sorted populations of meiotic nuclei. This may help to clarify the

relationship between histone marks and active transcription. Finally, a large

fraction of dynamic histone marks (H3K4me2/3) in meiosis do not coincide with

gene promoters or DSB hotspots. The functions of these sites are unknown

although some are likely to be enhancers. Together, they represent a large,

heretofore unstudied set of functional sites that may be modulating mouse meiosis.

To examine the chromatin landscape at DSB hotspots, we also performed the first

systematic genome-wide survey on a wide repertoire of histone marks. Our ability

to efficiently sort early MPI nuclei eliminated the need to perform these

experiments in juvenile mice; this negates the logistical challenge of collecting

sufficient juvenile animals and assures consistent and high levels of enrichment.

Aside from H3K4me3[28] and H3K36me3[17,18], H3K9ac, H3K4me2, H3K4me1,

H3K27ac and H4ac5 were enriched at DSB hotspots genome-wide. Incorporating
the extra information about these histone marks only marginally increases our
ability to predict hotspot strength, implying that even the expanded chromatin
landscape at hotspots remains a poor predictor of DSB formation. The genome is
replete with H3K4me3 marks that are used for DSB formation in the absence of
PRDM9, however, enigmatically, these sites are not used if PRDM9 is functional.
We demonstrated that both H3K36me3[17,40,42], and multiple histone marks
distinguish between these sites. Although this unique histone "code" may help the
DSB machinery to distinguish PRDM9-defined H3K4me3 from other H3K4me3
marks, it is equally possible that direct interactions mediated by PRDM9 itself are
important for this distinction. It seems unlikely that PRDM9 is responsible for all of
the histone tail modifications at hotspots, therefore other histone remodelers are
likely acting at the sites of PRDM9-marked H3K4/K36me3. H3K9ac and H3K4me3
are strength correlated and have a similar temporal dynamic implying that they
occur at a comparable frequency at hotspots. In mitotic cells, H3K9 acetylation is
actively promoted at H3K4 trimethylated nucleosomes[48]. It is therefore likely that
H3K9ac is a constitutive response to H3K4 trimethylation. Nonetheless, histone
acetylation may be functionally important *per se* in the context of DSB formation;
for example, in fission yeast, H3K9 to H3A9 mutation eliminates H3K9ac and
consequently reduces DSB formation[40].
392
393 Overall, we demonstrated that our straightforward and robust method can explore
394 previously hidden dynamics of mammalian meiotic recombination. Importantly, our
395 method is widely applicable to other organisms, tissues and cell types, paving the
396 way for further understanding of the temporal dynamics of other developmental
397 processes.
398

Acknowledgements

We thank members from Camerini-Otero's lab for discussions, and Galina Petukhova for insightful comments. We are grateful to Mary Ann Handel for sharing H1t antibodies. This study used the high performance computational capabilities of the Biowulf Linux cluster at the National Institutes of Health, Bethesda, MD (http://biowulf.nih.gov). We thank members from the NHLBI Flow Cytometry Core for assistance with nuclei sorting, and members from the NIDDK Genomics Core for assistance with high-throughput sequencing. This work was funded by the NIDDK Intramural Research Program (R.D.C.O).

Author contributions

K-W.G.L., K.B., G.C., F.P. and R.D.C-O. conceived the study and designed the experiments. K-W.G.L. performed the experiments. K-W.G.L., K.B. and F.P. analyzed the data. K-W.G.L., K.B., F.P. and R.D.C-O. wrote the manuscript.

Ethical compliance

The authors confirm that all experiments were performed in accordance with the NIH Animal Care and Use regulations.

418 **Table 1.** Summary of cell-type-specific H3K4me3 ChIP-Seq.

Sub-stage	Experiment	Antibody	Purity (%)	Starting chromatin DNA (ng)	SPoT (HS; %)	SPoT (All; %)	Peaks (#)
Leptonema	1	EpiCypher	96.9	2,398	2.4	4.5	9,528
	2	EpiCypher	95.3	805	2.2	7.6	13,095
	2	Abcam	95.3	805	3.3	20.3	30,943
Zygonema	1	EpiCypher	79.7	2,688	4.7	6.4	15,699
	2	EpiCypher	64.6	755	3.6	4.2	11,314
	2	Abcam	64.6	755	5.3	17.0	29,114
Early Pachynema	1	EpiCypher	81.6	1,793	1.6	4.2	6,647
	2	EpiCypher	89.5	4,585	1.7	2.3	4,883
	2	Abcam	85.2	960	2.0	14.0	17,561
Late Pachynema	1	EpiCypher	84.1	4,936	1.7	2.6	5,189
	2	EpiCypher	71.3	915	1.6	3.8	9,124
	2	Abcam	71.3	915	1.5	20.1	19,711
Diplonema	1	EpiCypher	88.7	6,502	1.8	3.1	6,645
	2	EpiCypher	82.3	1,115	1.7	5.4	12,071
	2	Abcam	82.3	1,115	1.6	21.0	18,191

Fig. 1 Immunofluorescence staining of spermatocyte nuclei.

Immunofluorescence images and signal quantification of stage-specific spermatocyte nuclei through meiosis prophase I. Details for signal quantification are described in Methods. Microscopic images are selected from two independent experiments in which two different combinations of primary antibodies are used; one using SCP3, H1t and SCP1, another one using SCP3 and STRA8. *Early and late pachytene nuclei cannot be unambiguously differentiated in the absence of H1t staining, and are therefore merged for counting and signal quantification. Source data are provided as a Source Data file.

Fig. 2 Experimental design for isolating stage-specific spermatocyte nuclei.
a Signal strength of intra-nuclear markers across meiotic sub-stages observed in immunofluorescence staining. Signal strength is classified as absent (−), very weak (−/+), weak (+), medium (++), or strong (+++). Combinatorial markers for isolating stage-specific nuclei are highlighted. **b** Flow cytometric strategies for isolating populations of stage-specific nuclei. Meiotic 4C nuclei are gated by DAPI and SCP3 signals (top left). Stage-specific nuclei are sorted into populations based on combinatorial signals of intra-nuclear markers in two separate sorts; one using antibodies against STRA8 for leptonema (top right), and the other one using antibodies against H1t and SCP1 for seven populations (P1 to P7) (bottom). **c** Distributions and purities of each specific type of nuclei in whole-testis and in

sorted populations. The five selected populations of leptonema, zygonema, early

pachynema, late pachynema, and diplonema are highlighted as LE, ZY, EP, LP,

and DI, respectively. Purities (mean with standard error) of these populations are

derived from two or three independent sorts. Source data are provided as a Source

Data file. **d** Numbers of nuclei in each sub-population collected in an adult mouse.

Data (mean with standard error) are derived from two independent sorts.

Fig. 3 H3K4me3 dynamics across meiotic sub-stages.

a H3K4me3 dynamics at DSB hotspots across five meiotic sub-stages. H3K4me3 signals at DSB hotspots are normalized by the spike-in H4K4me3 mononucleosomes. **b** H3K4me3 dynamics at different genomic elements in five sorted sub-populations of meiotic nuclei. **c** Stage-specific H3K4me3 at TSSs recapitulate meiotic gene expression patterns. Genes with only one isoform are selected for this analysis. H3K4me3 profiles at TSSs are grouped into five clusters (top), while the respective mRNA levels are shown in the bottom panel. L = Leptonema, Z = Zygonema, P = Early and Late Pachynema, D = Diplonema.

Fig. 4 Histone modifications at DSB hotspots.

a Distributions of histone marks along a genomic region. Locations of DSB hotspots are shaded in green, TSS in orange. **b** Histone marks are enriched at DSB hotspots. Red depicts enrichment relative to IgG, blue is depletion. **c** H3K9ac dynamics at DSB hotspots across five meiotic sub-stages. **d-e** Principal component analysis using histone marks can distinguish DSB hotspots from PRDM9-independent H3K4me3 sites. **d** Histone modifications can distinguish DSB hotspots from other H3K4me3 sites. An ROC curve for each histone mark (gray) or PC (blue) is shown. Selected histone modifications and PC1 are highlighted. **e** Contributions of histone marks to each PC. Only marks that contribute more than expected are shown. Deeper blue indicates a stronger contribution (numbers show percent contribution).

Methods

Mouse strains

C57BL/6J (B6) mice were either obtained from The Jackson Laboratory (Stock no.

000664) or bred in-house. All experiments were done on adult mice (≥ 8 weeks of

age).

Antibodies

The following antibodies were used for primary immunofluorescence staining: anti-

SCP3 (D-1) (Santa Cruz, sc-74569), anti-STRA8 (Abcam, ab49602), anti-H1t (a

gift from Mary Ann Handel and a custom-made antibody), and anti-SCP1 [Biotin]

(Novus Biologicals, NB300-229B).

The following antibodies were used for secondary immunofluorescence staining:

goat anti-mouse IgG conjugated with Cy3 (Jackson ImmunoResearch, 115-167-

003), goat anti-rabbit IgG conjugated with FITC (Jackson ImmunoResearch, 115-

097-003), goat anti-mouse IgG conjugated with DyLight488 (Abcam, ab98757),

goat anti-rabbit IgG conjugated with Cy3 (Jackson ImmunoResearch, 111-165-

003), goat anti-guinea pig IgG conjugated with Alexa Fluor 488 (Thermo Fisher

Scientific, A-11073), streptavidin conjugated with Alexa Fluor 647 (Thermo Fisher

Scientific, S21374).

The following antibodies were used for ChIP: anti-H3K4me3 (Abcam, ab8580),

anti-H3K4me3 (EpiCypher, 13-0028), anti-H3K9ac (Active Motif, 39918), anti-

H3K4me2 (Active Motif, 39914), anti-H3K36me3 (Active Motif, 61102), anti-H4ac5

(Millipore, 06-946), anti-H3K4me1 (Abcam, ab8895), anti-H3K27ac (Abcam,

ab177178), anti-H4K8ac (Abcam, ab15823), anti-H4K12ac (Active Motif, 39928),

anti-H3K27me3 (Millipore, 07-449), anti-H4K20me3 (Millipore, 07-463), anti-

H3K79me1 (Abcam, ab2886), anti-H3K4ac (Abcam, ab176799), anti-H3K79me3

(Abcam, ab2621), anti-H3K9me2 (Abcam, ab1220), anti-H3K9me3 (Active Motif,

39766), anti-H3K27me1 (Millipore, 07-448), anti-H3 (Abcam, ab1791), normal
rabbit IgG (Millipore, 12-370).

Nuclei preparation

Testes from adult mice were de-capsulated and fixed with 1% formaldehyde for 10

min followed by 5 min of quenching at RT. Fixed tissues were homogenized and

filtered through a 70- or 100-μm cell strainer. After washing with chilled 1 x PBS,

nuclei were extracted using nucleus extraction buffer (15 mM Tris-HCl pH 7.4, 0.34

M sucrose, 15 mM NaCl, 60 mM KCl, 0.2 mM EDTA, 0.2 mM EGTA) on ice for 5

min and were homogenized with 20 strokes with loose pestle followed by 10

strokes with tight pestle. Nuclei were filtered through a 40-μm cell strainer and

resuspended in chilled PBTB buffer (1 x PBS with 0.1% Triton X-100, 5% BSA and

protease inhibitor).

Immunofluorescence staining

Nuclei were incubated in 10% of normal serum at RT for 10 min. This blocking step

helps reduce non-specific binding of antibodies. Nuclei were then labelled with

different combinations of primary antibodies (1 μg of antibodies to 11 million

events) in 10% of normal serum at 20°C for 40 min. Nuclei were washed,

resuspended in PBTB, blocked with serum, and then labelled with secondary

antibodies (1:250 dilution) in 10% of normal serum at 20°C for 30 min. Controls for

each secondary antibody were also prepared for setting up the threshold of

background fluorescent signal for FACS. Nuclei were washed, resuspended in

PBTB, and stored at 4°C until sorting.

Counting stage-specific nuclei from whole-testis

Nuclei isolated from whole-testis were stained with antibodies against SCP3, H1t

and SCP1, and subsequently with secondary antibodies as described.

Immunofluorescence-labelled nuclei were spread on microslides, stained with

mounting medium with DAPI (Vetashield), sealed with cover slips, and viewed

using fluorescence microscopy. Each of the five types of MPI nuclei (SCP3+) were

counted; these were characterized by the patterning of the SCP3 and H1t proteins
(Fig. 1). A total of 678 nuclei were counted from two independent experiments. To
extrapolate counts to the percentage of cells of each stage in whole testis (top
panel in Fig. 2c), we use the following formula:
543
544   WT_{stage} (%) = N_{stage}/678 * (WT_{4C}/WT_{all}) * 100
545
546   N_{stage} = number of nuclei of each stage
547   WT_{4C} = number of total 4C (SCP3+) nuclei from sorting
548   WT_{all} = number of total singlet nuclei (1C to 4C) from sorting
549

Quantification of immunofluorescence signal intensity

Nuclei isolated from whole-testis were stained with antibodies against SCP3 and
STRA8 or with antibodies against SCP3, H1t and SCP1, subsequently with
secondary antibodies, and spread as previously described. Immunofluorescence
signals of each marker were quantified using Volocity 6.2.1 (PerkinElmer). 4C
nuclei were identified using signals from the DAPI channel by excluding objects
with size < 40 μm^2. Touching objects were separated using the parameter of 30−50
μm^2. Stage-specific nuclei in MPI were characterized by the presence of SCP3
signal and by the patterning of synaptonemal complex[14]. Signals from all channels
were recorded and exported for signal quantification.
560

Isolation of meiotic sub-populations with FACS

Nuclei were filtered through a 40-μm cell strainer, and stained with DAPI for 30 min
or longer at RT before sorting. All sorting experiments were performed on either a
BD FACSAria II or a BD FACSAria Fusion flow cytometer at a flow rate of ~20,000
events/sec. Singlets were gated using both forward scatter and side scatter. Nuclei
from primary spermatocytes (4C) were gated based on DNA content deduced from
the DAPI signal, and sorted into populations of interest based on fluorochrome
intensity into collection tubes containing PBTB. Sorted nuclei were collected by
centrifugation. Nuclei were examined under microscope and counted with a

hemocytometer. Finally, excess buffer was removed and nuclei pellets were stored

at −80°C.

Assessment of purity of sorted populations

An aliquot of nuclei suspension from each sorted population was spread on

microslides, further stained with mounting medium with DAPI (Vectashield), and

sealed with coverslips for purity examination. About 150 nuclei from each sorted

population were checked at 400X magnification using immunofluorescence

microscopy. The staging was primarily accessed by the patterning of SCP3[14] and

further confirmed by signals of STRA8 or of H1t and SCP1 accordingly.

Chromatin extraction

Chromatin extracted for ChIP-Seq was either sheared by sonication or Micrococcal

nuclease (MNase) digestion. Chromatin for all H3K4me3/H3K9ac-ChIP-Seq for

studying histone dynamics in stage-specific populations was sheared by

sonication, whereas ChIP-Seq for identifying histone marks at hotspots were

performed on MNase-digested chromatin. Experimental procedures are listed

below in detail.

Shearing chromatin by sonication:

Frozen nuclei pellets were thawed at RT for 10 min. Nuclei were lysed with Lysis

buffer 1 (0.25% Triton X-100, 10 mM EDTA, 0.5 mM EGTA, 10 mM Tris-HCl pH 8)

and incubated at RT for 10 min. Nuclei pellets were subsequently washed with

Lysis buffer 2 (200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 10 mM Tris-HCl pH 8)

and lysed with RIPA buffer (10 mM Tris-HCl pH 8, 1 mM EDTA, 0.5 mM EGTA,

1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS, 140 mM NaCl plus

protease inhibitor). Chromatin was sheared into ~50−300 bp fragments by

sonication using Bioruptor (diagenode). The amount of chromatin DNA was

measured using a Qubit dsDNA HS Assay Kit (Thermo Fisher).

MNase digestion:

Frozen nuclei pellets were thawed at RT for 10 min. Nuclei pellets were

resuspended with MNase buffer (50 mM Tris-HCl pH 8, 1 mM $CaCl_2$, 4 mM $MgCl_2$,

4% NP-40 plus protease inhibitor). MNase digestion was performed in a

concentration of 3U MNase (USB, Affymetrix) per one million nuclei at 37°C for 5

min. The reaction was stopped by adding a final concentration of 10 mM EDTA

and incubated at 4°C for 5 min. Soluble chromatin was collected and diluted with

RIPA buffer.

Chromatin Immunoprecipitation

For stage-specific H3K4me3-ChIP-Seq, modified mononucleosomes (SNAP-ChIP

K-MetStat Panel, EpiCypher) were added at a spike-in volume of 1 μl per ~1.5 μg

chromatin DNA. Chromatin was immunoprecipitated with 0.8–5 μg antibodies (or

5 μl unpurified serum) in 0.5–1 ml RIPA buffer at 4°C overnight (see

Supplementary Table 1 for details). The immuno-complexes were captured using

20–75 μl Dynabeads Protein G (30 mg/ml, Novex) at 4°C for 2 hr. The beads were

washed with buffers previously described[27]. ChIPed DNA was eluted using a IPure

kit v2 (diagenode).

Sequencing library construction

ChIP-seq libraries were constructed with a KAPA Hyper Prep Kit (Kapa

Biosystems) following steps for generating 1 μg of library DNA. DNA libraries were

cleaned up with an Agencourt AMPure XP system (Beckman Coulter). The DNA

concentration and fragment size of these libraries were measured with a Qubit

dsDNA HS Assay Kit (Thermo Fisher) and an Agilent High Sensitivity DNA Kit

(Agilent), respectively.

High throughput DNA sequencing

DNA sequencing was performed on the Illumina HiSeq 2500 or HiSeq X.

Sequencing tags were aligned to the mouse mm10 reference genome using BWA

mem 0.7.12[49]. For ChIP-Seq samples containing the K-MetStat spike-in panel, we

aligned to a custom mm10 genome to which we added the sequences of the DNA
bound by each K-MetStat nucleosome.

Inter-sample normalization and peak strength estimation using spike-in
nucleosomes

We retained only sequencing reads with an alignment q-score >= 20, that were
mapped in a "proper pair" (for paired-end data), that passed vendor QC and that
were mapped as a primary alignment. For paired-end data, we only retained reads
from fragments with an insert size between 100 bp and 200 bp. Normalization
using the K-MetStat spike-in was performed as described previously [29], with slight
modification. Briefly, each spike-in nucleosome is bound to a unique DNA
sequence. Thus, the number of reads recovered for each nucleosome is a proxy
for the frequency of that nucleosome in the library. We first converted paired-end
reads to fragments, then calculated the enrichment of fragments mapping to each
spike-in sequence after ChIP:

E_{si} = (barcode fragments in ChIP / barcode fragments in input)

Next, we calculated H3K4me3 ChIP-Seq fragment coverage in all 25 bp non-
overlapping windows in the genome. The per-locus enrichment was then
calculated as :

E_{locus} = (fragment coverage in ChIP) / (fragment coverage in Input)

The histone modification density was then calculated as :

HMD (%) = 100 * (E_{locus} / $E_{H3K4me3\_A}$)

$E_{H3K4me3\_A}$ is the E_{si} value for the H3K4me3_A spike_in. For our final estimate of
H3K4me3 signal, we normalized the HMD by the global average HMD. The ChIP-

Seq signal at peaks and other genomic intervals was calculated as the sum of the
signal in all overlapping 25 bp windows.

Inter-sample normalization and peak strength estimation using "stable"
promoters

The K-MetStat panel cannot be used to normalize data from H3K9ac ChIP-Seq.
Thus, we used an alternative strategy using stable promoters. NCIS[50] was used to
estimate the contribution of background to each sequencing library. We then
calculated the NCIS normalized ChIP-Seq fragment coverage in all 25 bp windows
in the genome (c = ChIP coverage - (Input coverage * NCIS factor)). The ChIP-
Seq signal at the central ± 1 Kbp region around GENCODE transcript 5' ends
(TSS) was quantified as the sum of the coverage in all 25 bp widows overlapping
the site. TSS coverage was estimated in each of the five meiotic populations. TSS
that overlapped a DSB hotspot, TSSs with a negative strength after correction and
TSS with the strongest (1%) and weakest (15%) H3K4me3 signal at each stage
were discarded. The $\log_2$ ratio of NCIS-corrected H3K4me3 was calculated at each
TSS between all pairs of stages. TSSs with an absolute $\log_2$ (fold-change) ≤ 1.8
between all stages were retained. This resulted in 362 TSSs for H3K4me3 and
187 TSSs for H3K9ac. The median signal across these intervals was calculated
for each ChIP and used as a normalization factor. The strength of each 25 bp
genome interval was calculated as the NCIS-corrected coverage divided by the
normalization factor. The ChIP-Seq signal at peaks and other genomic intervals
was calculated as the sum of the signal in all overlapping 25 bp windows. Note
that for H3K4me3 ChIP-Seq, this internal normalization gives comparable results
to those using the K-MetStat spike-in panel (Supplementary Fig. 4).

Correlating H3K4me3 profiles at TSSs with mRNA expression levels
We used RNA-Seq data from spermatocyte populations obtained by classical cell-
sorting to quantify gene expression[32]. Gene expression at GENCODE vM20
transcripts was quantified for each RNA-Seq dataset using kallisto 0.45.0 (kallisto
quant --single --single --single-overhang --fragment-length=250 --sd=100 --

seed=42). Only transcripts with maximal expression > 0.2 tags per million (TPM),

with at least a 1.25-fold change in expression and with non-zero expression and

H3K4me3 signals at all stages were considered for subsequent analysis. To

minimize the confounding effects of multiple isoforms, only TSS with a single

transcript in this list were retained.

The temporal expression profile for each transcript was described as a four unit

vector (Leptonema, Zygonema, Pachynema, Diplonema). Similarly, a temporal

profile of H3K4me3 at the TSS was generated. We averaged the Early Pachnema

and Late Pachynema H3K4me3 signals to allow direct comparison with the RNA-

Seq data.

The overall correlation between H3K4me3 and gene expression was calculated by

concatenating all gene expression vectors and comparing with a concatenated

vector of H3K4me3. To calculate the expected random correlation, we shuffled the

order of each 4-unit transcript H3K4me3 vector before concatenation. 10,000

iterations of this process were performed.

k-means clustering was used to group similar H3K4me3 temporal profiles and to

compare each set to gene expression. We used an implementation of the gap-

statistic to determine the optimal number of clusters (R; factoextra package).

Within each set of clusters the method described in the previous paragraph was

employed to determine the correlation between H3K4me3 and expression

temporal profiles and to derive an empirical p-value for this correlation. Clustering

by gene expression instead of by H3K4me3 signal gives similar correlations,

though the number of clusters can change (data not shown).

Peak calling

Peaks for H3K4me3 ChIP-Seq were called using MACS2 (version 2.1.2)[51] with

default parameters except (-q 0.1 --broad) and with a stage-matched input DNA

sample as a control. Peaks overlapping DSB hotspots or gene promoters were

ascertained using bedtools (version v2.27.1)[52] after removing blacklisted regions

described in[53]. DSB hotspots were defined and reported in previous studies[28].

Transcription start sites (TSSs) were defined as the ± 0.5 Kbp region around

GENCODE v20 transcripts[54].

Unbiased clustering

ChIP-Seq peaks from all five populations were merged and peak strength was

calculated (as described above) for each merged peak in each population. The

H3K4me3 profile for each peak was described as a five unit vector (LE, ZY, EP,

LP, DI) and scaled by subtracting the mean and dividing by the standard deviation.

k-means clustering was used to cluster. Five clusters was determined as optimal

using the gap-statistic.

Multiple linear regression

Hotspots overlapping a TSS or overlapping a site used for DSB formation in

$Prdm9^{-/-}$ mice were discarded. Only hotspots with read coverage > 0 for all histone

marks were used for regression analyses because we performed regression on

the log10 transformed coverage values. The leaps package in R was used to

perform an all-subsets regression using the seven histone marks enriched at DSB

hotspots.

Principal Component Analysis

All H3K4me3 peaks from LE, ZY, EP and LP were used for analysis. Each interval

was resized to ± 250 bp around the center. Sequencing reads for each histone

mark were counted at each feature. Input DNA reads were also counted and

subtracted from the count for each mark, following NCIS[50] correction. DSB

hotspots and GENCODE TSSs were expanded to ± 1,500 bp to determine

H3K4me3 peaks that overlapped hotspots and TSSs, respectively. Peaks that

overlapped both a hotspot and TSS were discarded as the potential compound

signal would confound these analyses. Only autosomal peaks were used.

H3K36me3 and H3K4me3 were excluded. The R prcomp command was used for

Principal Component Analysis. Variables were scaled to have unit variance and
zero centered.

ROC curves were built by ranking intervals by the dependent variable. Intervals
were ranked either from high to low or from low to high and the ROC with the higher
area under the curve was used.

Code availability

The custom pipeline used for these analyses is deposited at zenodo
(https://doi.org/10.5281/zenodo.2651204)[55].

Data availability

765 The authors declare that all data generated or analysed during this study are
766 included in this published article (and its supplementary information files). The
767 sequencing data reported in this paper are archived at the Gene Expression
768 Omnibus (www.ncbi.nlm.nih.gov/geo) as accession no. GSE121760. The source
769 data underlying Figs 1 and 2c are provided as a Source Data file.

References

1. Baudat, F., Imai, Y. & de Massy, B. Meiotic recombination in mammals: localization and regulation. *Nat. Rev. Genet.* **14**, 794–806 (2013).

2. Handel, M. A., Eppig, J. J. & Schimenti, J. C. Applying 'Gold Standards' to In-Vitro-Derived Germ Cells. *Cell* **157**, 1257–1261 (2014).

3. Bellve, A. R., Cavicchia, J. C. & Millette, C. F. Spermatogenic cells of the prepuberal mouse: isolation and morphological characterization. *J. Cell Biol.* (1977).

4. Goetz, P., Chandley, A. C. & Speed, R. M. Morphological and temporal sequence of meiotic prophase development at puberty in the male mouse. *J. Cell Sci.* **65**, 249–263 (1984).

5. Kluin, P. M., Kramer, M. F. & de Rooij, D. G. Spermatogenesis in the immature mouse proceeds faster than in the adult. *Int. J. Androl.* **5**, 282–294 (1982).

6. Zelazowski, M. J. *et al.* Age-Dependent Alterations in Meiotic Recombination Cause Chromosome Segregation Errors in Spermatocytes. *Cell* **171**, 601–614.e13 (2017).

7. Grive, K. J. *et al.* Dynamic transcriptome profiles within spermatogonial and spermatocyte populations during postnatal testis maturation revealed by single-cell sequencing. *PLoS Genet.* **15**, e1007810 (2019).

8. Hogarth, C. A. *et al.* Turning a spermatogenic wave into a tsunami: synchronizing murine spermatogenesis using WIN 18,446. *Biol. Reprod.* **88**, 40–41 (2013).

9. Bastos, H. *et al.* Flow cytometric characterization of viable meiotic and postmeiotic cells by Hoechst 33342 in mouse spermatogenesis. *Cytometry A* **65**, 40–49 (2005).

10. Barchi, M., Geremia, R., Magliozzi, R. & Bianchi, E. Isolation and analyses of enriched populations of male mouse germ cells by sedimentation velocity: the centrifugal elutriation. *Methods Mol. Biol.* **558**, 299–321 (2009).

11. Gaysinskaya, V., Soh, I. Y., Heijden, G. W. & Bortvin, A. Optimized flow cytometry

isolation of murine spermatocytes. *Cytometry A* **85**, 556–565 (2014).

12. Bonn, S. *et al.* Cell type–specific chromatin immunoprecipitation from multicellular

complex samples using BiTS-ChIP. *Nat. Protoc.* **7**, 978 (2012).

13. Meuwissen, R. L. *et al.* A coiled-coil related protein specific for synapsed regions of

meiotic prophase chromosomes. *EMBO J.* **11**, 5091–5100 (1992).

14. Dobson, M. J., Pearlman, R. E., Karaiskakis, A., Spyropoulos, B. & Moens, P. B.

Synaptonemal complex proteins: occurrence, epitope mapping and chromosome

disjunction. *J. Cell Sci.* **107 (Pt 10)**, 2749–2760 (1994).

15. Inselman, A., Eaker, S. & Handel, M. A. Temporal expression of cell cycle-related

proteins during spermatogenesis: establishing a timeline for onset of the meiotic

divisions. *Cytogenet. Genome Res.* **103**, 277–284 (2003).

16. Mark, M. *et al.* STRA8-deficient spermatocytes initiate, but fail to complete, meiosis

and undergo premature chromosome condensation. *J. Cell Sci.* **121**, 3233–3242

(2008).

17. Powers, N. R. *et al.* The Meiotic Recombination Activator PRDM9 Trimethylates

Both H3K36 and H3K4 at Recombination Hotspots In Vivo. *PLoS Genet.* **12**,

e1006146 (2016).

18. Grey, C. *et al.* In vivo binding of PRDM9 reveals interactions with noncanonical

genomic sites. *Genome Res.* **27**, 580–590 (2017).

19. Margolin, G., Khil, P. P., Kim, J., Bellani, M. A. & Camerini-Otero, R. D. Integrated

transcriptome analysis of mouse spermatogenesis. *BMC Genomics* **15**, 39 (2014).

20. Sun, F. *et al.* Nuclear localization of PRDM9 and its role in meiotic chromatin

modifications and homologous synapsis. *Chromosoma* **124**, 397–415 (2015).

21. Moens, P. B. *et al.* The time course and chromosomal localization of recombination-

related proteins at meiosis in the mouse are compatible with models that can

resolve the early DNA-DNA interactions without reciprocal recombination. *J. Cell*

Sci. **115**, 1611–1622 (2002).

22. Cole, F. *et al.* Homeostatic control of recombination is implemented progressively in

mouse meiosis. *Nat. Cell Biol.* **14**, 424–430 (2012).

23. Moens, P. B. Histones H1 and H4 of surface-spread meiotic chromosomes.

Chromosoma **104**, 169–174 (1995).

24. Adli, M. & Bernstein, B. E. Whole-genome chromatin profiling from limited numbers

of cells using nano-ChIP-seq. *Nat. Protoc.* **6**, 1656–1668 (2011).

25. Bernstein, B. E. *et al.* Genomic maps and comparative analysis of histone

modifications in human and mouse. *Cell* **120**, 169–181 (2005).

26. ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the

human genome. *Nature* **489**, 57–74 (2012).

27. Smagulova, F. *et al.* Genome-wide analysis reveals novel molecular features of

mouse recombination hotspots. *Nature* **472**, 375–378 (2011).

28. Brick, K., Smagulova, F., Khil, P., Camerini-Otero, R. D. & Petukhova, G. V. Genetic

recombination is directed away from functional genomic elements in mice. *Nature*

485, 642–645 (2012).

29. Grzybowski, A. T., Chen, Z. & Ruthenburg, A. J. Calibrating ChIP-Seq with

Nucleosomal Internal Standards to Measure Histone Modification Density Genome

Wide. *Mol. Cell* **58**, 886–899 (2015).

30. Mimitou, E. P., Yamada, S. & Keeney, S. A global view of meiotic double-strand

break end resection. *Science* **355**, 40–45 (2017).

31. Shah, R. N. *et al.* Examining the Roles of H3K4 Methylation States with

Systematically Characterized Antibodies. *Mol. Cell* **72**, 162–177.e7 (2018).

32. Gaysinskaya, V. *et al.* Transient reduction of DNA methylation at the onset of

meiosis in male mice. *Epigenetics Chromatin* **11**, 15 (2018).

33. Tibshirani, R., Walther, G. & Hastie, T. Estimating the number of clusters in a data

set via the gap statistic. *J. R. Stat. Soc. Series B Stat. Methodol.* **63**, 411–423

(2001).

34. Vastenhouw, N. L. *et al.* Chromatin signature of embryonic pluripotency is

established during genome activation. *Nature* **464**, 922–926 (2010).

35. Diagouraga, B. *et al.* PRDM9 Methyltransferase Activity Is Essential for Meiotic DNA

Double-Strand Break Formation at Its Binding Sites. *Mol. Cell* **69**, 853–865.e6

(2018).

36. Buard, J., Barthès, P., Grey, C. & de Massy, B. Distinct histone modifications define

initiation and repair of meiotic recombination in the mouse. *EMBO J.* **28**, 2616–2624

(2009).

37. Walker, M. *et al.* Affinity-seq detects genome-wide PRDM9 binding sites and

reveals the impact of prior chromatin modifications on mammalian recombination

hotspot usage. *Epigenetics Chromatin* **8**, 31 (2015).

38. Getun, I. V. *et al.* Functional Roles of Acetylated Histone Marks at Mouse Meiotic

Recombination Hot Spots. *Mol. Cell. Biol.* **37**, (2017).

39. Merker, J. D. *et al.* The histone methylase Set2p and the histone deacetylase

Rpd3p repress meiotic recombination at the HIS4 meiotic recombination hotspot in

Saccharomyces cerevisiae. *DNA Repair* **7**, 1298–1308 (2008).

40. Yamada, S., Ohta, K. & Yamada, T. Acetylated Histone H3K9 is associated with

meiotic recombination hotspots, and plays a role in recombination redundantly with

other factors including the H3K4 methylase Set1 in fission yeast. *Nucleic Acids Res.*

41, 3504–3517 (2013).

41. Lange, J. *et al.* The Landscape of Mouse Meiotic Double-Strand Break Formation,

Processing, and Repair. *Cell* **167**, 695–708.e16 (2016).

42. Yamada, S. *et al.* Genomic and chromatin features shaping meiotic double-strand

break formation and repair in mice. *Cell Cycle* **16**, 1870–1884 (2017).

43. Thacker, D., Mohibullah, N., Zhu, X. & Keeney, S. Homologue engagement controls

meiotic DNA break number and distribution. *Nature* **510**, 241–246 (2014).

44. Wojtasz, L. *et al.* Mouse HORMAD1 and HORMAD2, two conserved meiotic

chromosomal proteins, are depleted from synapsed chromosome axes with the help

of TRIP13 AAA-ATPase. *PLoS Genet.* **5**, e1000702 (2009).

45. Kauppi, L. *et al.* Numerical constraints and feedback control of double-strand breaks

in mouse meiosis. *Genes Dev.* **27**, 873–886 (2013).

46. Page, J. *et al.* Inactivation or non-reactivation: what accounts better for the silence

of sex chromosomes during mammalian male meiosis? *Chromosoma* **121**, 307–326

(2012).

47. Cao, J. *et al.* The single-cell transcriptional landscape of mammalian

organogenesis. *Nature* **566**, 496–502 (2019).

48. Gates, L. A., Foulds, C. E. & O'Malley, B. W. Histone Marks in the 'Driver's Seat':

Functional Roles in Steering the Transcription Cycle. *Trends Biochem. Sci.* **42**, 977–

989 (2017).

49. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. *arXiv [q-bio.GN]* (2013).

50. Liang, K. & Keleş, S. Normalization of ChIP-seq data with control. *BMC*

Bioinformatics **13**, 199 (2012).

51. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137

(2008).

52. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**, 841–842 (2010).

53. Smagulova, F., Brick, K., Pu, Y., Camerini-Otero, R. D. & Petukhova, G. V. Erratum:

The evolutionary turnover of recombination hot spots contributes to speciation in

mice. *Genes Dev.* **30**, 871 (2016).

54. Mudge, J. M. & Harrow, J. Creating reference gene annotation for the mouse

C57BL6/J genome assembly. *Mamm. Genome* **26**, 366–378 (2015).

55. kevbrick. *kevbrick/lametal2019: Analytic pipeline for Lam et al. (Nature*

Communications revision). (2019). doi:10.5281/zenodo.2656746

56. Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible

platform for exploring deep-sequencing data. *Nucleic Acids Res.* **42**, W187–91

(2014).

Supplementary Figures

STRA8

a

SCP3

913
914
915
916

917 **Supplementary Fig. 1** Microscopic examination of sorted nuclei after FACS.
918 Whole-field immunofluorescence images of sorted nuclei in the populations of **a**
919 leptonema and **b** zygonema. Lep=leptonema; Zyg=zygonema; ePac=early
920 pachynema.
921
922

Supplementary Fig. 2 H3K4me3 antibodies have varying levels of substrate specificity.

We performed ChIP-Seq across five MPI stages using two H3K4me3 antibodies; EpiCypher 13-0028 (H3K4me3; two replicates) and Abcam ab8580 (abK4me). The EpiCypher K-MetStat panel is a collection of nucleosomes with different histone modifications that was spiked into each H3K4me3 ChIP-Seq experiment to validate the specificity of the antibody and to allow cross sample normalization

for ChIP-efficiency (see methods). The relative enrichment of sequences bound
to each modified nucleosome are shown here, compared to the relative
frequency in the sample before ChIP was performed. The Epicypher antibody is
highly specific to H3K4me3, however, as previously demonstrated[31], the Abcam
antibody shows substantial cross-reactivity with H3K4me2. Note that although
the enrichment for H3K4me2 is relatively low compared to H3K4me3, the
relative abundance of H3K4me2 in the genome compared to H3K4me3[31] makes
it difficult to assess the quantitative impact of this cross-reactivity in ChIP-Seq
experiments.

940
941
Supplementary Fig. 3 H3K4me3 is relatively elevated at stronger DSB hotspots
in leptonema. **a** Snapshot of H3K4me3 at four DSB hotspots. The DSB hotspots
are progressively stronger from left to right. At the three weaker hotspots (left)
the H3K4me3 signal is maximal at zygonema. At the strongest hotspot (right) the
H3K4me3 signal is strongest at leptonema. **b-c** Systematic trend towards
stronger leptonema H3K4me3 at stronger DSB hotspots. **b** The H3K4me3 ratio
between leptonema and zygonema is shown. **c** A more pronounced trend is seen
using the Abcam ab8580 H3K4me3 antibody (abK4me).

Supplementary Fig. 4 H3K4me3 dynamics are similar using different
normalization approaches.

Coverage heatmaps at DSB hotspots through MPI. **a,c,e** Coverage is normalized
using NCIS and stable TSS signals (see methods). **b,d** Coverage is normalized

using the K-MetStat spike-in. The spike-in normalization cannot be used for the
H3K4me2/3 sample because of antibody cross-reactivity (Abcam ab8580
antibody). Both normalization methods yield similar results for H3K4me3 replicate
1 (R1; **a,b**) and replicate 2 (R2; **c,d**). A slight increase in the leptonema signal and
a marginal signal at early pachynema are seen with the H3K4me2/3 ChIP-Seq
(**e**).
961

MPI stage

All transcripts in cluster 5

Clustering by expression

Representative individual transcripts

Supplementary Fig. 5 H3K4me3 dynamics correlate with gene expression

profiles from male juvenile mice.

Each of the plots represents a single cluster of transcripts from Fig. 3 (Cluster 1-5

are from left to right on Fig. 3c, respectively). Throughout, red depicts H3K4me3

ChIP-Seq signal and blue depicts gene expression from RNA-Seq. Uppermost

panels show all transcripts in the cluster; lines represent individual transcripts.

Transcripts were next sub-clustered by gene expression profile (middle panels; k-

means clustering; optimal k chosen using gap statistic). Clusters in which profiles

correlate more or less than expected are highlighted (* = P < 0.01, ** = P < 0.001;

1,000 bootstraps of shuffling stages for each gene). Six representative transcripts

from each cluster are shown in the lower panel below each cluster. Transcripts of

genes known to play a role in MPI were preferentially selected; if < 6 such genes

were present in a cluster, the remainder were randomly selected.

Supplementary Fig. 6 Extensive dynamics revealed by unbiased clustering of H3K4me3 peaks. a 16% of all H3K4me3 peaks occur at sites not defined at DSB hotspots or transcription start sites (TSS). These are classified as "Other". 5.8% of H3K4me3 peaks coincide with both a DSB hotspot and a GENCODE TSS (HS & TSS). **b** Unbiased k-means clustering of MPI H3K4me3 profiles. Cluster number is indicated in parentheses in the bottom left corner. Each black line represents the MPI profile of an individual H3K4me3 peak. 10,000 randomly chosen profiles are shown. Purple lines depict the mean profile for each cluster. Profiles are normalized by the mean and standard deviation. The optimum number of clusters was identified using the gap statistic (optimum k = 5; range tested = 2 < k < 24). Cluster number is shown on each panel in parentheses. **c** Cluster composition shown as the percentage of each type of H3K4me3 peak within each cluster. Cluster 2 (N=11,336) is the only cluster composed primarily of peaks at DSB hotspots. **d** The percentage of peaks of each type across clusters is shown. Notably, 96% of DSB hotspots occur in cluster 2. 74% of HS & TSS peaks also occur in this cluster, suggesting that H3K4me3 at many of these sites is PRDM9-mediated. **e** In mice lacking PRDM9, PRDM9-independent H3K4me3 peaks are used for targeting the DSB machinery. These sites are termed "default hotspots" and are defined as H3K4me3 peaks that coincide with a DSB hotspot in *Prdm9*[-/-] mice[28]. Default hotspots form more frequently than expected at H3K4me3 peaks that are present in early MPI (cluster 1; N = 6,223). Peaks with a late dynamic (cluster 4; N=7,744) are used less than expected. The "Not used" peaks are all H3K4me3 peaks that do not coincide with either a PRDM9-defined or a default hotspot.

1005

**Supplementary Fig. 7 Extensive dynamics revealed by unbiased clustering
of H3K4me2/3 peaks. a** 35% of all H3K4me2/3 peaks (using Abcam ab8580
antibody) occur at sites not defined at DSB hotspots or transcription start sites
(TSS). These are classified as "Other". 3.7% of H3K4me3 peaks coincide with
both a DSB hotspot and a GENCODE TSS (HS & TSS). **b** Unbiased k-means
clustering of MPI H3K4me2/3 profiles. Cluster number is indicated in
parentheses in the bottom left corner. Each black line represents the MPI profile
of an individual H3K4me2/3 peak. 10,000 randomly chosen profiles are shown.
Purple lines depict the mean profile for each cluster. Profiles are normalized by
the mean and standard deviation. The optimum number of clusters was identified
using the gap statistic (optimum k = 6; range tested = 2 < k < 24). Cluster number
is shown on each panel in parentheses. **c** Cluster composition shown as the
percentage of each type of H3K4me2/3 peak within each cluster. Cluster 3 is the
only cluster composed primarily of peaks at DSB hotspots. **d** The percentage of
peaks of each type across clusters is shown. Notably, 80% of DSB hotspots
occur in cluster 3. 52% of HS & TSS peaks also occur in this cluster, suggesting
that H3K4me3 at many of these sites is PRDM9-mediated. **e** In mice lacking
PRDM9, PRDM9-independent H3K4me3 peaks are used for targeting the DSB
machinery. These sites are termed "default hotspots" and are defined as
H3K4me3 peaks that coincide with a DSB hotspot in *Prdm9*[-/-] mice[28]. Default
hotspots form more frequently than expected at H3K4me3 peaks that are present
in early MPI (cluster 2). Peaks with a late dynamic (cluster 5, 6) are used less
than expected. "Not used" peaks are all H3K4me3 peaks that do not coincide
with either a PRDM9-defined or a default hotspot.

1030

1031
1032

Supplementary Fig. 8 Heatmaps of histone modification ChIP-Seq at DSB hotspots. Coverage data were averaged in 150 bp windows and converted to reads per Kbp per million (RPKM) to facilitate cross-comparison. Heatmaps were generated using deeptools[56]. It is important to note that some enrichment at DSB hotspots is seen when using a non-specific antibody (IgG; last panel).

1039

1040

Supplementary Fig. 9 Enrichment of histone modifications at functional
genomic elements. Histone modification coverage was plotted around functional
sites in the genome. Enrichment is shown as mean signal - mean signal in IgG
ChIP-Seq (red=high; blue=low). L1 LINEs were obtained from the repeatmasker
database. Enhancers were obtained from the UCSC table browser RefSeq
functional elements table.

1047

1048

1049

1050
1051

Supplementary Fig. 10 Correlations between histone modifications and DSB hotspot strength. a Pearson correlation coefficients of log-transformed read counts and hotspot strength (strength). Only histone marks enriched at DSB hotspots were considered. Hotspots with above-zero coverage for all histone marks and hotspot strength were used (N = 11,253). **b** Multiple linear regression with histone marks only marginally improves correlation with DSB hotspots. The lower panel depicts the histone marks used (red; x) for models with each number of features. The order of addition was determined using all-subsets regression with the leaps package in R. Small increases in R^2 may simply arise because additional samples reduce stochastic noise in strength estimates.

Supplementary Fig. 11: Histone marks distinguish hotspots from other H3K4me3 sites.

a ROC curves for individual histone marks and principal components (PCs) for discriminating hotspots from other H3K4me3 sites in the genome. Dashed line indicates no predictive ability. **b** The area under the ROC curves (AUC). **c** Most of the variance in histone mark signals is captured by few PCs.