Excitability changes in the motor cortex in preparation for self-paced and cue-guided movements: a transcranial magnetic stimulation study

J. Ibáñez[1], R. Hannah[1,2], L. Rocchi[1], J.C. Rothwell[1]

1. Department of Movement Disorders, Institute of Neurology, UCL, UK
2. Department of Psychology, University of California San Diego, USA

CORRESPONDENCE: Jaime Ibáñez, UCL Institute of Neurology, London, WC1N 3BG, UK, email: j.ibanez@ucl.ac.uk

ABBREVIATED TITLE: Cortical excitability changes before cue-guided and self-paced movements

CONFLICT OF INTEREST: The authors declare no competing financial interests.

ACKNOWLEDGEMENTS: JI was supported in part by Grant No. #H2020-MSCA-IF-2015-700512 from the European Commission (JI). RH was supported by a Biotechnology and Biological Sciences Research Council (BBSRC) (Grant No. BB/N016793/1).

SIGNIFICANCE (120 words):

Brain responses to transcranial magnetic stimulation provide insight into how the brain

prepares planned movements. However, most experiments involve temporally-controlled

stimulus-driven actions. We provide evidence that cortical excitability evolves over the

same time course in self-paced and reaction time movements. This is paralleled by

previously unreported similarities in the way movement onset times are altered (speeded)

when stimuli are delivered some hundreds of milliseconds before movements are to be

started. This contrasts starkly with movements timed according to predictable

countdowns. Here we see equivalent temporal evolution of pre-movement excitability,

but the timing of these movements is unaffected by the external stimuli, thus suggesting

a higher control of the time of action initiation.

ABSTRACT (250 words):

A simple of model of movement preparation, involving subthreshold accumulation of

motor commands readily explains two features of a simple reaction time task. One is

preparatory inhibition of corticospinal excitability, which is viewed as necessary to

prevent premature release of movement. The other is intersensory facilitation in which

receipt of a second stimulus at the same time as the imperative signal shortens the time

taken to trigger a reduction in inhibition and increase in excitation. This allows

preparatory activity to reach threshold more quickly and release action.

Here we question that interpretation by asking what happens in movements that are self-

paced or timed to be coincident with an external event (predictive movements).

Surprisingly we find that both types of movement show preparatory inhibition at

approximately the same time prior to movement initiation as in a reaction time movement.

This suggests that rather than preventing premature release, premovement inhibition is an

integral part of preparation to move, and may reflect preparatory neural population

dynamics shared between actions initiated in different contexts. We also observed

intersensory facilitation in self-paced movements when a sensory event preceded

predicted movement onset by about 200-300ms. We suggest that this is consistent with

the idea that the preparatory state requires a separate trigger event before movement is

initiated. Intersensory facilitation was not seen in predictive movements, perhaps because

they filter out external signals that could interfere with internal timing events.

Introduction (650 words)

Simple reaction time (RT) movements are usually viewed as the stimulus-triggered

release of prepared, "subthreshold", movement commands (Tanji and Evarts, 1976; Prut

and Fetz, 1999). But what happens if movements are made without reference to an

external trigger? For example, self-paced (SP) movements, or movements that are made

to coincide with an external event rather than being triggered by it ("predictive" task: PT).

If a movement is self-paced or predictive, then are movement commands still stored in

advance and released by an internal trigger? Or does movement occur immediately

preparation is complete? Perhaps no preparation is needed at all.

The aim of the present experiments was to compare movement preparation in these three

types of movement (RT, PT and SP). In particular, we tested whether 2 key characteristics

of RT movements differed in the three tasks.

The first characteristic was preparatory inhibition. In RT tasks, corticospinal excitability

is reduced around the expected time of the imperative stimulus (Hasbroucq et al., 1997).

This has been interpreted as preventing the premature release of the prepared

"subthreshold" motor plan (Touge et al., 1998; Duque et al., 2010, 2017). It is compatible

with data from animal experiments showing that there are changes in the firing of

neurones in motor cortex at a time when there is no overt EMG activity in the periphery

(Omrani et al., 2017). If preparatory inhibition is necessary to prevent premature release

of movement commands (Duque et al., 2010), then it may be reduced in PT movements

where the time to move is known, and it should not be present at all in SP movements

when there is intentionally no constraint on the time of execution.

The second characteristic of RT movements investigated here was inter-sensory

facilitation, in which the presence of a second sensory stimulus at around the time of the

imperative stimulus reduces reaction times and speeds release of the prepared movement

(Nickerson, 1973; Pascual-Leone et al., 1992b). It has been postulated that the additional

input shortens the time taken to identify the imperative stimulus (Pascual-Leone et al.,

1992b). SP and PT movements may not require an external trigger since they could start

immediately preparation was complete. If so they would not display inter-sensory

facilitation.

Recent work on the relationship between movement preparation and execution leads to

rather different predictions. Recordings of neural population activity in cortical motor

areas of monkeys show that activity of neurons in the preparatory period may differ from

that in the same neurons during execution (Churchland et al., 2010; Kaufman et al.,

2014a). Preparatory activity is not a "subthreshold" version of that seen during execution

and therefore, preparatory suppression would seem to be unnecessary (Kaufman et al.,

2014b). Indeed, in human experiments, the degree of preparatory suppression was

positively correlated to the RT, suggesting that, if anything, preparatory suppression was

causally involved in preparation to move rather than in preventing its premature release

(Hannah et al., 2018). Thus, preparatory suppression might be seen in SP and PT

movements as well as in RT tasks.

Recordings of neural activity also show that all types of movement are preceded by the

same patterns of activity consisting of a preparation period during which discharge rates

change without any overt EMG activity in the periphery (Lara et al., 2018). This is

followed by a transition to an execution phase that appears to be caused by a separate and

non-specific timing signal (Haith et al., 2016; Kaufman et al., 2016a). If this timing signal

corresponds to an internal trigger for movement, then it may be susceptible to inter-

sensory facilitation in SP and PT movements.

Our results show that both SP and PT movements display preparatory suppression of

corticospinal excitability consistent with the idea that it is a necessary part of movement

preparation rather than preventing premature release. SP movements also display inter-

sensory facilitation-like effects, with onset times of speeded responses occurring earlier

than predicted, suggesting that execution depends on some form of trigger. In contrast,

PT movements do not display inter-sensory facilitation, which could indicate that there is

active filtering of irrelevant sensory events that might distract from the internal trigger

used in predictive timing.

Materials and Methods

Participants

In total, 28 right-handed, healthy participants (9 females, 28 ± 5 years old, age range 19

- 36 years) participated in this study. All of them reported no contraindications to TMS

(Rossi et al., 2011) and had normal or corrected to normal visual acuity. The study was

approved by the University College London Ethics Committee and warranted to be in

accordance with the Declaration of Helsinki. All participants signed a written informed

consent prior to the experimental session.

Recordings

Participants sat in a comfortable chair with both forearms resting on a pillow placed on

their lap and index fingers resting on a keyboard which registered movement times. A

screen was placed approximately one meter in front of them. Participants wore ear

defenders during the TMS experiments to reduce the influence of loud sounds generated

by the discharges of the TMS coils.

EMG signals were obtained from the right first dorsal interosseous (FDI) muscle for

experiment 1 and of both hands in experiment 2. EMG activity from the right abductor

digiti minimi (ADM) muscle was also recorded in the two experiments. MEPs recorded

from the right FDI were the main outcome of this study. The ADM was used as a control

muscle for the MEPs: it was activated by the TMS stimuli but, unlike the right FDI, it

was not directly involved in the movement, so it was only supposed to show a monotonic

reduction of excitability up until the time at which muscles showed voluntary activation

(Duque et al., 2010). Surface recording electrodes were placed on the muscle bellies, with

references on the closest metacarpophalangeal joint. The ground electrode was placed on

the right wrist. EMG signals were amplified, band-pass filtered between 20 Hz and 2000

Hz (Digitimer D360, 2015 Digitimer Ltd, United Kingdom) and acquired at 5000 Hz

sampling rate with a data acquisition board (CED-1401, Cambridge Electronic Design

Ltd 2016) connected to a PC and controlled with the Signal and Spike[2] software (also by

CED).

Once EMG sensors were set, the participants' TMS hotspot was located. This was done

by finding the point over M1 giving the largest MEPs in the contralateral FDI for a given

stimulus intensity. The TMS coil was always held at a 45^{o} angle to the sagittal plane with

the handle pointing backwards. Once the hotspot was found, the resting motor threshold

(RMT) and 1 mV intensity were determined. The RMT was estimated by adjusting the

TMS output until 5 out of 10 MEPs could be obtained over 50 µV. The 1 mV intensity

was defined as that able to elicit MEPs of around 1 mV amplitude. The experimental

paradigms were implemented using custom-made Matlab routines (Mathworks, MA,

USA). Synchronization of TMS pulses with EMG and movement events was realised

using Cogent 2000[1]'s utilities to control the parallel port of the PC running the

experimental paradigms. Data analysis was carried out using custom-made Matlab

functions and SPSS software (IBM, NY, USA).

Experiment 1 – TMS recordings preceding RT and PT movements

In this experiment, participants performed two types of movement paradigms: 1) a PT

task in which movements were timed with an external countdown signal (Fig. 1-A); and

2) a RT task in which movements were initiated following and imperative stimulus (Fig.

1-B). During task execution, TMS pulses were delivered at predetermined time points

relative to the average task- and subject-specific movement times to probe motor cortical

excitability. The PT task consisted of a resting phase of 2 s followed by a delay period of

1 s during which four circles moved from the extremes of a cross towards its centre with

a velocity inversely proportional to the remaining distance to the centre of the cross

(initial distance 4.5 cm). Participants were instructed to time their movements with the

merging of the circles so that button presses with the FDI were performed as soon as the

circles fully overlapped. After each movement, the trial ended by giving participants

feedback about the button press times. This feedback was displayed for a random period

of time between 1-3 s, and it consisted of the time at which the button press had been

---

[1] By Cogent 2000 team at the FIL and the ICN and Cogent Graphics developed by John
Romaya at the LON at the Wellcome Department of Imaging Neuroscience

detected, expressed in ms, and a font colour code indicative of the performance: green

text was used for button presses done between 50-100 ms relative to the time at which

circles overlapped, thus encouraging participants to aim at pressing the button within this

interval; yellow text was used for button presses in the intervals 0-50 ms and 100-150 ms;

finally, red text was used in any other case. Additionally, participants were given "too

early" and "too late" messages when button presses were performed during the delay

period or more than 200 ms after its end. The RT task had the same states with matched

durations as the PT task but, in this case, the movement of the circles during the delay

period followed a random and uninformative path along the four arms of the cross (Fig.

1). In this case participants were instructed to wait until the circles suddenly appeared in

the centre of the cross, which was considered the "GO" cue. Participants were specifically

asked to avoid any predictions of when the imperative cues were appearing. For that, they

were told to specifically use the sudden overlapping of the circles in the centre of the

screen to make a fast and reactive movement.

At the start of each task, participants practised the two paradigms until they showed

consistent movement times. Thirty additional movements were then performed with each

paradigm so that the subject- and task-specific average movement onset times (based on

the FDI EMG activity) could be estimated. Then, the actual recording took place,

consisting in two blocks per paradigm (interleaving the blocks of the two paradigms). In

each block, six conditions were tested for 10 trials each, using a randomized order of

conditions. These conditions differed from each other with regards to the timing of the

TMS pulse: 1) no TMS delivered (control condition); 2) TMS at the beginning of the

delay period (baseline condition); 3) TMS halfway through the delay period; 4) TMS 200

ms before the average movement onset time; 5) TMS 60 ms before the average movement

onset time; and 6) TMS 30 ms before the average movement onset time.

Experiment 2 – TMS during the resting phases between SP movements

The task involved participants sitting still and comfortably, with both index fingers

resting on buttons on a keyboard. They were instructed to make ballistic bilateral button

presses every 4-8 seconds, whilst avoiding pre-movement muscle activation and ensuring

movements were made in a similar way along the whole experiment (Fig. 1-C).

Movements of the left index finger (non-targeted by the TMS) allowed accurate estimates

of the onset of voluntary movements (Schneider, 2004) without being affected by the

TMS induced delays of voluntary actions in cases where the stimulus was given in close

proximity with the intended movement onset time (Ziemann et al., 1997). Participants

were instructed to perform their movements spontaneously and to avoid any form of

internal countdown to decide when to initiate the movements. It was stressed to

participants that they must not let the TMS pulses alter their decision to move. A resting

period of time followed by a button press was considered a trial, and 12 blocks were

performed by each participant with 65 trials making up a block. During blocks, EMG was

monitored to ensure the hand was in a relaxed condition between movements.

A custom-made Matlab program was used to determine the timing of a TMS stimulus on

a given trial based on the button press times registered in the previous 5 trials performed

by each participant without TMS (this number of trials was empirically chosen to allow

the code program to quickly adapt to changes in participants' behaviours). TMS pulse

timing were distributed so that in 4% of the trials, stimuli were delivered early after the

previous movement (3 s after the previous button press); 8% of the trials were non-TMS

trials, which were then used to monitor inter-movement intervals in the absence of

external stimuli along the experiment. Finally, in 82% of the trials, TMS pulse timings

were defined based on the probability density function of inter-movement intervals

considering the 5 most recent non-TMS trials. For that, a Gaussian fit was estimated and

the next TMS firing time was selected according to the left-hand side of this probability

density function. TMS firing times were thus programmed to be delivered at a time

interval relative to the previous movement such that it was always below the average

inter-movement interval estimated. It was important to ensure that time intervals left by

participants between button presses were not too long, as this would result in reduced

chances of delivering TMS pulses at points in time close to the button presses. In the cases

when participants waited for over 10 s between movements for 3-4 trials in a row,

participants were given an indication by the experimenter to reduce the inter-movement

time intervals.

Analysis of results and statistics

All the analyses were done using the onsets of the EMG as the reference points indicating

the times of movement initiations. In order to obtain EMG references in each trial, the

absolute value of the EMG recordings was estimated and then a moving average of 5 ms

and a low pass filter (fc < 5Hz) were applied to obtain a smoothed envelope of the EMG

signal. EMG recordings of all trials whilst the participant was at rest were analysed to

obtain a subject-specific baseline level. A threshold set at five times the baseline value

was used to determine EMG onset times. This level was also used to detect and remove

trials with pre-TMS or pre-movement activation of all the muscles registered. All trials

were then visually inspected and manually corrected to ensure that EMG-based

movement onsets were estimated properly and that no building-up of EMG activity was

apparent before the TMS pulses. Once trials were corrected, EMG onset times and peak

to peak amplitudes of MEPs were estimated. MEP amplitudes were estimated from the

acquired EMG signals without applying any additional filters. A logarithmic

transformation of MEP amplitudes was performed before the statistical tests to ensure

normality in the distributions of amplitudes.

In Experiment 1, MEP amplitudes and times of EMG onsets of all trials were labelled

according to the type of paradigm (PT v RT) and to the time at which TMS was delivered.

Movement onset times were referenced to the ones obtained with each participant in the

control trials without TMS (subtracting the average movement onset times). A two-way

repeated measures ANOVA (rmANOVA) with factors TIME and PARADIGM was

performed to test for changes in movement onset times. A three-way rmANOVA (TIME

x PARADIGM x MUSCLE) was performed to test for changes in MEPs. Post-hoc

comparisons were done between the baseline and 200 ms before movement TMS

conditions.

In Experiment 2, two types of trials were extracted per session: TMS trials in which a

TMS stimulus was delivered before the movement, and non-TMS trials in which no TMS

pulse was scheduled so they could be used as control trials to define the correct times of

subsequent TMS pulses. To check if TMS pulses were inadvertently used by the

participants as cues to start the movements in TMS trials, we compared inter-movement

intervals in TMS and non-TMS trials using a paired t-test. Additionally, to analyse the

effect of TMS pulses on the times at which participants decided to move, non-TMS trials

were used to simulate the length of the TMS-movement intervals had participants not

been biased by TMS stimuli. To do this, all non-TMS trials obtained from each session

were used by a simulation algorithm (5000 iterations) that: *i*) randomly selected 5 trials

in each iteration; *ii*) obtained a simulated TMS time for the "next" trial in the same way

as in the actual experiment; *iii*) randomly selected a new trial of the same participant; *iv*)

obtained the time interval between the movement onset time and the simulated TMS time

and kept it if it was positive (*i.e.*, if the TMS was delivered before the movement).

Regarding the TMS recordings in experiment 2, since the times of the MEPs relative to

the posterior movements could not be well controlled (due to the free nature of SP

movements), standard bootstrap statistics were applied to all participants' MEPs

recordings in order to identify, in an unbiased way, intervals of interest, *i.e.*, intervals that

contained significant increases or decreases in MEP amplitudes. To do this, the following

steps were repeated for 100 iterations: 1) 200 TMS-trials per participant were chosen at

random. Since the number of TMS trials obtained with each participant was different, this

step balanced the influence of each participant's data when combining data from all

participants. 2) MEPs selected from each participant were referenced to MEPs in the

interval [-1500 : -500] ms relative to the onset of movements, *i.e.* MEPs were subtracted

the mean and divided by the standard deviation of the MEPs in the defined interval. 3)

MEP values from all participants were merged. 4) A sliding window of 40 ms in steps of

20 ms was applied from -1 s to the movement onset. For each window, 40 MEPs were

picked at random with replacement (standard bootstrapping procedure) and used to

calculate a mean. This was repeated 1000 times, thus generating 1000 means for every

window. These values were arranged in ascending order and the 25^{th} and 975^{th} values

were taken as confidence intervals. This whole process was repeated 100 times (using

different sets of 200 trials per participant each time), and an average of all estimated

confidence intervals was taken to produce the definitive confidence intervals of MEP

changes across the time in preparation for the movements. Subsequently, a two-way

rmANOVA was performed in SPSS with factors TIME and MUSCLE to test for changes

in MEP amplitudes. Here, the TIME factor included the baseline period [-1500:-500] ms,

and the periods of time in which significant decreases or increases of the right FDI MEPs

were observed (*i.e.*, time intervals when both confident interval lines obtained from the

previous bootstrap analysis were either positive or negative).

Overall, all results are reported as group mean ± standard error of the mean (SEM).

Bonferroni *post-hoc* corrections are applied in the cases of multiple comparisons. *P*

values < 0.05 are considered to be significant. The Greenhouse–Geisser procedure was

applied where necessary to correct for violations of sphericity in rmANOVAs.

Results

Experiment 1

The average movement onset times (based on EMG) obtained in the baseline trials

without TMS were -35 ± 6 ms and 200 ± 9 ms for the PT and RT paradigms, respectively.

The average time intervals between the estimated EMG onset times and the button press

events were 93 ± 3 ms (PT) and 91 ± 3 ms (RT). Fig. 2 shows the observed average

movement times in the two paradigms for different TMS firing times. There appear to be

two main effects. First, if the TMS pulses are given 30 or 60ms prior to average movement

times, then reactions are delayed. Previous work as ascribed this to the effect to the silent

period following the TMS-evoked MEP. The second effect is that reaction times are

reduced in the RT task when TMS pulses are given 200ms prior to average movement

times at around the time of the imperative stimulus. Previous work refers to this as a form

of intersensory facilitation caused by the auditory click and scalp stimulation from the

TMS pulse.

These conclusions are borne out in the statistical analysis. rmANOVA showed a

significant main effect of TIME ($F_{[4,6]} = 18.154$; $P < 0.001$), reflecting delayed responses

when TMS pulses were delivered in proximity to the average movement times (*i.e.*, higher

movement times were found when TMS pulses where delivered 60 ms and 30 ms before

the average movement times). In addition, there was a significant interaction of

PARADIGM x TIME ($F_{[4,6]} = 11.943$; $P = 0.003$). Post-hoc comparison between

paradigms for trials in which TMS pulses were delivered 200 ms before the average

movement time revealed a significant reduction of the movement times in the RT

movements relative to the PT (mean difference of -29.8 ± 8.8 ms; *P = 0.008*). No

significant difference was found between paradigms for baseline TMS trials (mean

difference -3.8 ± 5.0 ms; $P = 0.473$).

Resting motor threshold and 1mV levels were 53 ± 3 % and 63 ± 3 % of the maximum

stimulator output, respectively. Fig. 3 shows average MEP results. rmANOVA showed a

main effect of factors MUSCLE ($F_{[1,9]} = 8.327$; $P = 0.018$; difference between FDI and

ADM: 0.7 ± 0.2 log(μV)) and TIME ($F_{[4,6]} = 5.731$; $P = 0.010$). The post-hoc comparison

between MEPs obtained from TMS pulses delivered either at the baseline point or 200

ms before the average movement times revealed a significant reduction of MEPs at the

latter point (mean difference: -0.5 ± 0.1 log(μV); $P = 0.009$). There was a significant

interaction of MUSCLE x TIME ($F_{[4,6]} = 11.943$; $P < 0.001$), due to the diverging

directions of MEP changes in the FDI and ADM at the time of movement initiation

(Duque et al., 2010). Post-hoc comparisons between baseline trials and trials with TMS

delivered 200 ms before the average movement time revealed significant reductions of

MEPs both in the FDI (mean difference -0.5 ± 0.2 log(μV); *P = 0.018*) and in the task-

irrelevant ADM (mean difference -0.4 ± 0.1 log(μV); *P = 0.040*). No significant

differences were found in the reductions (TMS 200 ms before movements) of the FDI

MEPs between the RT and PT paradigms ($P = 0.590$). MEPs obtained with TMS pulses

delivered 200 ms before average movement times relative to baseline MEPs showed an

average reduction of 31 % and 35 % for PT and RT tasks, respectively.

Experiment 2

The average inter-movement intervals in TMS and non-TMS trials was 4.94 ± 0.16 s and

4.99 ± 0.18 s. A paired t-test on the inter-movement intervals of TMS and non-TMS trials

revealed no significant difference between both conditions ($P = 0.415$). Fig. 4 shows a

comparison between real and simulated distributions (across all participants) of time

intervals between TMS firing times and consecutive movement onset times. The

experimental design was such that distribution of TMS times relative to the time of

movement should reflect the left-hand side of a normal distribution (before movement),

with the peak occurring close to the onset of movement. Instead, the obtained

distributions show a trough around 300 ms followed by a peak at ~100 ms before the

onset of the muscle activation. This suggests that, in trials in which TMS is delivered less

than 300 ms before participants were about to move, movements were speeded, thus

modifying the normal distribution expected to result from the design of the paradigm. The

paired sample t-test returned significant differences between the simulated and the real

distributions 300 ms before the onsets of the movements and in the interval [-150 ms : -

50 ms] also relative to the movement (p < 0.001 in both intervals). We refer to this

speeding up as a form of intersensory facilitation.

Resting motor threshold and 1 mV levels were 56 ± 3 % and 66 ± 4 % of the maximum

stimulator output, respectively. Fig. 5 shows the summary of the MEP results obtained

both using bootstrap statistics on the grouped (z-scored) data and from the posterior

comparison between the intervals of interest. The bootstrap analysis returned a significant

($P < 0.01$) interval of MEP reduction that applied both to the FDI and the ADM muscles.

For the FDI, this period with reduced MEPs peaked at -120 ms (relative to the estimated

muscle activation onsets) and it was followed by an (also significant) increase of

excitability towards the time of initiation of the movement. According to these results,

the period of reduced excitability was defined to be at the time interval [-160 ms : -100

ms] relative to the movement (*i.e.*, the two consecutive 40 ms windows showing the

strongest MEP reduction effects in the bootstrap analysis). Due to the lack of a dense

enough population of MEPs at the late phase before movements were initiated, we used

the MEPs located within the final 80 ms interval before starting the movements as

representative of the movement initiation phase (Pascual-Leone et al., 1992b; Chen and

Hallett, 1999; Chen et al., 1999). rmANOVA showed main effects of MUSCLE ($F_{[1,17]} =$

26.246; $P < 0.001$; mean difference between FDI and ADM: 1.0 ± 0.2 log(μV)) and TIME

($F_{[2,16]} = 9.046$; $P = 0.01$). The post-hoc comparison of MEPs within the baseline and

excitability reduction intervals revealed a significant reduction of MEPs at the latter time

point (mean difference: 0.3 ± 0.1 log(μV); $P = 0.006$). There was also a significant

interaction of MUSCLE x TIME ($F_{[2,16]} = 13.041$; $P < 0.001$). Post-hoc comparisons

between MEPs for baseline and reduced excitability conditions and for each muscle

separately revealed a significant effect in both cases ($P = 0.019$ for FDI; $P = 0.020$ for

ADM).

Discussion (1500 words)

In the present experiments we probed the temporal evolution of M1 excitability in

paradigms with different constraints on movement initiation times. The results showed

that reduced excitability can be observed prior to a voluntary movement whether it is self-

paced, predictive or reactive. In addition, we found that TMS had distinct biasing effects

on movement times across the different paradigms. While the initiation of a predictive

movement was unaffected by TMS, RT and SP movements were speeded in a way

resembling intersensory facilitation effects commonly reported in RT tasks (Nickerson,

1973). Overall, the results support the notion that preparatory cortical excitability changes

in M1 do not have a direct functional role in preventing premature movement releases

and they raise the question about the possible similarities of the mechanisms involved in

triggering planned voluntary movements in the presence or absence of specific external

cues.

Different movement preparation paradigms exhibit similar patterns of preparatory M1

excitability

A comparison of the results obtained in the three paradigms tested here suggests a

common temporal evolution of corticospinal excitability before movements, regardless

of the nature of the trigger that initiates them. It has previously been hypothesized that

M1 excitability reduction before movements reflects proactive control of motor cortical

outputs preventing premature responses (Bestmann and Duque, 2015; Duque et al., 2017).

However, this does not explain why we see the same premovement suppression prior to

SP movements, in which there is explicitly no temporal constraint on initiation time.

Strictly speaking, suppression is also unnecessary in the PT movements since if

preparation was initiated at the correct time it would automatically evolve to reach

threshold and initiate a movement to coincide with the external event. Only if suppression

was used to reduce the variability of onset times would it serve a function.

Instead, our results appear a better fit for the alternative hypothesis that smaller responses

reflect the dynamics of neural populations evolving towards more stable states from

which to initiate movement (Churchland et al., 2010; Shenoy et al., 2013; Kaufman et al.,

2014b, 2016b; Hannah et al., 2018). Recent studies in primates show that the time spent

in this preparatory state can be compressed or extended depending on task demands (Lara

et al., 2018). This observation might explain why we observed a peak in MEP suppression

in SP movements which, relative to RT and PT movements, was shifted towards time

points closer to the EMG onsets. It may also be relevant to the longer RTs in paradigms

with unpredictable imperative stimuli (Alink et al., 2010). These could be explained by

the need to make a transition through reduced excitability states after the imperative

stimulus. Indeed, previous work has shown that, in reaction time tasks without prior

warning cues, MEPs remain unchanged before the imperative signal but show a marked

suppression right after the onset of the imperative cue (Duque et al., 2014). Altogether,

our results support the notion that MEP changes in preparation for movements can largely

be explained as being a result of action- and state-specific M1 evolutions towards

movement rather than being a reflection of proactive control mechanisms over M1.

The size and spatio-temporal patterns of the observed excitability reduction in preparation

for movements, which was comparable across conditions, matched the results of previous

TMS studies based on RT paradigms. On average, MEP amplitudes decreased by 30%,

in line with changes reported before (Duque et al., 2014; Quoilin et al., 2016). The FDI

muscle (task-relevant) presented deeper decreases of MEP amplitudes than the control

ADM muscle across conditions, in agreement with previous research suggesting a

somatotopic gradient in excitability changes related to the location, certainty and

specificity of forthcoming actions (Greenhouse et al., 2015).

Intersensory facilitation depends on the predictability of the imperative stimulus

Different TMS biasing effects found in PT and RT movement times indicate that these

paradigms differ with respect to how motor commands are withheld and/or released.

Movement initiation in RT movements is contingent upon external cues and the link

between sensory processing and release of motor programs is presumed to be direct

(Pascual-Leone et al., 1992b). Speeded responses in RT movements induced by TMS

have been interpreted as intersensory facilitation (Nickerson, 1973; Terao et al., 1997),

and have been suggested to be due to a shortening of the time for identification of external

stimuli as the go-signal (Pascual-Leone et al., 1992b). In eye movement paradigms with

predictable onsets of imperative cues, previous studies have proposed that inner

mechanisms aimed at anticipating the timing of the environmental cues need to be

deployed and used to trigger motor commands (Janssen and Shadlen, 2005; Badler and

Heinen, 2006). Based on this, in the PT task tested here the lack of speeded responses can

be attributed to the usage of an alternative mechanism to trigger movements, which is

internally driven, once the timing of external stimuli is learned. In this case, external

signals during the delay period of PT movements may be either downregulated (Alink et

al., 2010) or simply disregarded (Rohenkohl et al., 2012).

Biasing effects of TMS in SP movements

Analysing brain responses to external stimuli before SP movements with a degree of

temporal precision is technically challenging and only few studies have attempted it.

Wasaka and colleagues used a SP paradigm to show that sensory suppression processes

in these actions partially resembled those seen in preparation for RT movements

(Shimazu et al., 1999; Wasaka et al., 2003). Castellote and colleagues showed that

StartReact responses during the resting periods before self-initiated movements are

closely similar to StartReac responses in RT paradigms (Valls-Sole et al., 1999; Castellote

et al., 2013). Interestingly, Castellote's experiment showed a biasing effect of startling

stimuli on movement times that tightly resembles that obtained here (Fig. 3), *i.e.*, if

delivered with a certain anticipation (~300 ms) before the forthcoming movements, the

startling stimuli speed-up the release of the required movements. In that work, the features

of the responses matched those obtained in StartReact paradigms using RT tasks, which

allowed authors to suggest that the mechanisms engaged in the preparation for SP and

cue-driven actions could share common elements (Castellote et al., 2013). In our case,

the intensity of the applied stimuli (1 mV TMS and participants using ear defenders ,

which lessen the likelihood of startle response) suggests that the observed effects are

closer to intersensory facilitation (Pascual-Leone et al., 1992b, 1992a). Precisely

quantifying how much movements are sped-up would help verify this idea (Valls-Sole et

al., 2008), but doing so is challenging because of the lack of a more precise knowledge

about when movements would be performed in the absence of external stimuli. Based on

the fact that stimuli used in the RT and SP paradigms were equal and responses to TMS

comparable, it is conceivable that effects observed in both cases reflect the use of a similar

neural strategy to trigger actions that is not shared by PT movements.

Technical considerations and future work

Studying corticospinal changes before SP movements with TMS has inherent limitations.

We were able to obtain movement times and MEPs in our SP paradigm despite the

apparent difficulties in accurately probing excitability at specific, well-defined times

relative to movement onset. However, unlike with cue-guided movements, the

estimations of TMS times (relative to the movements) in the SP paradigm were done

based on the times of the subsequent movements of the non-stimulated hand, which also

presents intersensory facilitation effects (Hannah et al., 2018), is not affected by cortical

silent period-related delaying effects (Ziemann et al., 1997), but may still have been

biased by the TMS pulse in a different way than the stimulated side. Therefore, precise

estimations of the excitability reduction peak time in this case are not definitive.

Previous studies testing the possible contribution of spinal mechanisms to the decrease

seen in MEP amplitudes in preparation for movements in reaction time tasks have led to

contradictory results (Duque et al., 2010; Lebon et al., 2016; Hannah et al., 2018). The

techniques used here to probe corticospinal excitability do not allow measuring whether

spinal inhibitory processes are taking place in preparation for movements performed in

the three paradigms tested. Knowing if and how spinal inhibition applies to cue-driven

and SP movements could help further proving the hypothesis that observed preparatory

changes are not driven by proactive control mechanisms. Such analysis should be

addressed in future work.

Conclusions

Taken together, results from the three compared paradigms suggest that TMS recordings

of excitability changes in preparation for planned movements share a common temporal

profile, likely reflecting aspects of neural population-level changes to generate the desired

actions. Results also suggest possible shared neural mechanisms involved in triggering

reaction time and self-paced movements that are different from those in actions timed

with fully predictable cues.

References

Alink A, Schwiedrzik CM, Kohler A, Singer W, Muckli L (2010) Stimulus Predictability Reduces
Responses in Primary Visual Cortex. J Neurosci 30:2960–2966.

Badler JB, Heinen SJ (2006) Anticipatory Movement Timing Using Prediction and External Cues. J
Neurosci 26:4519–4525.

Bestmann S, Duque J (2015) Transcranial Magnetic Stimulation: Decomposing the Processes Underlying
Action Preparation. Neurosci:1–14.

Castellote JM, Van Den Berg MEL, Valls-Sole J (2013) The startreact effect on self-initiated movements.
Biomed Res Int 2013.

Chen R, Corwell B, Hallett M (1999) Modulation of motor cortex excitability by median nerve and digit
stimulation. Exp Brain Res 129:77–86.

Chen R, Hallett M (1999) The time course of changes in motor cortex excitability associated with
voluntary movement. Can J Neurol Sci 26:163–169.

Churchland MM, Cunningham JP, Kaufman MT, Ryu SI, Shenoy K V. (2010) Cortical Preparatory
Activity: Representation of Movement or First Cog in a Dynamical Machine? Neuron 68:387–400
Available at: http://dx.doi.org/10.1016/j.neuron.2010.09.015.

Duque J, Greenhouse I, Labruna L, Ivry RB (2017) Physiological Markers of Motor Inhibition during
Human Behavior. Trends Neurosci xx:1–18.

Duque J, Labruna L, Cazares C, Ivry RB (2014) Dissociating the influence of response selection and task
anticipation on corticospinal suppression during response preparation. Neuropsychologia 65:287–
296.

Duque J, Lew D, Mazzocchio R, Olivier E, Ivry RB, Louvain D, Brussels B-, Clinica N (2010) Evidence
for Two Concurrent Inhibitory Mechanisms during Response Preparation. J Neurosci 30:3793–
3802.

Greenhouse I, Sias A, Labruna L, Ivry RB (2015) Nonspecific Inhibition of the Motor System during
Response Preparation. J Neurosci 35:10675–10684.

Haith AM, Pakpoor J, Krakauer JW (2016) Independence of Movement Preparation and Movement
Initiation. J Neurosci 36:3007–3015 Available at:
http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.3245-15.2016 [Accessed August 28, 2018].

Hannah R, Cavanagh XSE, Tremblay S, Simeoni S, Rothwell XJC (2018) Selective Suppression of Local
Interneuron Circuits in Human Motor Cortex Contributes to Movement Preparation. 38:1264–1276.

Hasbroucq T, Kaneko H, Akamatsu M, Possamaï C-A (1997) Preparatory inhibition of cortico-spinal
excitability: a transcranial magnetic stimulation study in man. Cogn Brain Res 5:185–192.

Janssen P, Shadlen MN (2005) A representation of the hazard rate of elapsed time in macaque area LIP.
Nat Neurosci 8:234–241.

Kaufman MT, Churchland MM, Ryu SI, Shenoy K V. (2014a) Cortical activity in the null space:
permitting preparation without movement. Nat Neurosci 17:440–448 Available at:

http://www.nature.com/articles/nn.3643 [Accessed August 28, 2018].

Kaufman MT, Churchland MM, Ryu SI, Shenoy K V. (2014b) Cortical activity in the null space:
Permitting preparation without movement. Nat Neurosci 17:440–448 Available at:
http://dx.doi.org/10.1038/nn.3643.

Kaufman MT, Seely JS, Sussillo D, Ryu SI, Shenoy K V., Churchland MM (2016a) The Largest
Response Component in the Motor Cortex Reflects Movement Timing but Not Movement Type.
eNeuro 3 Available at: http://eneuro.org/cgi/doi/10.1523/ENEURO.0085-16.2016 [Accessed
August 28, 2018].

Kaufman MT, Seely JS, Sussillo D, Ryu SI, Shenoy K V., Churchland MM (2016b) The Largest
Response Component in the Motor Cortex Reflects Movement Timing but Not Movement Type.
eNeuro 3 Available at: http://eneuro.sfn.org/cgi/doi/10.1523/ENEURO.0085-16.2016.

Lara AH, Elsayed GF, Zimnik AJ, Cunningham J, Churchland MM (2018) Conservation of preparatory
neural events regardless of how movement is initiated. Elife:7:e31826.

Lebon F, Greenhouse I, Labruna L, Vanderschelden B, Papaxanthis C, Ivry RB (2016) Influence of Delay
Period Duration on Inhibitory Processes for Response Preparation. Cereb Cortex 26:2461–2470.

Nickerson RS (1973) Intersensory facilitation of reaction time: Energy summation or preparation
enhancement? Psychol Rev 80:489–509.

Omrani M, Kaufman MT, Hatsopoulos NG, Cheney PD (2017) Perspectives on classical controversies
about the motor cortex. J Neurophysiol:jn.00795.2016 Available at:
http://jn.physiology.org/lookup/doi/10.1152/jn.00795.2016.

Pascual-Leone A, Brasil-neto J, Valls-Sole J, Cohen LG, Hallett M (1992a) Simple reaction time to focal
transcranial magnetic stimulation. Brain, 115, 109-122 :109–122.

Pascual-Leone A, Valls-Sole J, Wassermann EM, Brasil-neto J, Cohen LG, Hallett M (1992b) Effects of
Focal Transcranial Magnetic Stimula Tion on Simple Reaction Time To Acoustic , Visual and
Somatosensory. Significance:1045–1059.

Prut Y, Fetz EE (1999) Primate spinal interneurons show pre-movement. 401:590–594.

Quoilin C, Lambert J, Jacob B, Klein PA, Duque J (2016) Comparison of motor inhibition in variants of
the instructed-delay choice reaction time task. PLoS One 11:1–16.

Rohenkohl G, Cravo AM, Wyart V, Nobre AC (2012) Temporal Expectation Improves the Quality of
Sensory Information. J Neurosci 32:8424–8428 Available at:
http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.0804-12.2012.

Rossi S, Hallett M, Rossini P, Pascual-Leone A (2011) Screening questionnaire before TMS: An update.
Clin Neurophysiol 122:1686 Available at: http://dx.doi.org/10.1016/j.clinph.2010.12.037.

Schneider C (2004) Timing of cortical excitability changes during the reaction time of movements
superimposed on tonic motor activity. J Appl Physiol 97:2220–2227 Available at:
http://jap.physiology.org/cgi/doi/10.1152/japplphysiol.00542.2004.

Shenoy K V., Sahani M, Churchland MM (2013) Cortical Control of Arm Movements: A Dynamical
Systems Perspective. Annu Rev Neurosci 36:337–359 Available at:
http://www.annualreviews.org/doi/10.1146/annurev-neuro-062111-150509.

Shimazu H, Kaji R, Murase N, Kohara N, Ikeda A, Shibasaki H, Kimura J, Rothwell JC (1999) Pre-
movement gating of short-latency somatosensory evoked potentials. Neuroreport 10:2457–2460
Available at: http://www.ncbi.nlm.nih.gov/pubmed/10574351 [Accessed August 31, 2018].

Tanji J, Evarts E V (1976) Anticipatory activity of motor cortex neurons in relation to direction of an
intended movement. J Neurophysiol 39:1062–1068 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/824409 [Accessed July 23, 2018].

Terao Y, Ugawa Y, Suzuki M, Sakai K, Hanajima R, Gemba-Shimizu K, Kanazawa I (1997) Shortening
of simple reaction time by peripheral electrical and submotor-threshold magnetic cortical
stimulation. Exp Brain Res 115:541–545.

Touge T, Taylor JL, Rothwell JC (1998) Reduced excitability of the cortico-spinal system during the
warning period of a reaction time task. Electroencephalogr Clin Neurophysiol 109:489–495.

Valls-Sole J, Kumru H, Kofler M (2008) Interaction between startle and voluntary reactions in humans.
Exp Brain Res 187:497–507.

Valls-Sole J, Rothwell JC, Goulart F, Cossu G, Muñoz E (1999) Patterned ballistic movements triggered
by a startle in healthy humans. J Physiol 516:931–938.

Wasaka T, Hoshiyama M, Nakata H, Nishihira Y, Kakigi R (2003) Gating of somatosensory evoked
magnetic fields during the preparatory period of self-initiated finger movement. Neuroimage
20:1830–1838.

Ziemann U, Tergau F, Netz J, Hömberg V (1997) Delay in simple reaction time after focal transcranial
magnetic stimulation of the human brain occurs at the final motor output stage. Brain Res 744:32–
40.

Figure 1. Movement initiation tasks and TMS recordings. (A) In PT movements - experiment 1- participants had to press a button with their index finger after the end of a 1-s countdown (the time it took the 4 white circles to reach the centre of the cross). (B) In RT movements -experiment 1- participants had to perform the same movement as in the PT task, but here they were specifically instructed to perform movements in response to seeing the 4 white circles suddenly appearing on the centre of the cross. During the 1-s delay period in this case, circles moved randomly along the arms of the cross. Both in PT and RT movements, single-pulse TMS was delivered when the four circles appeared

on the screen at the beginning of the delay period (BASELINE), half way through the delay period and 200/60/30 ms before the average movement times in each subject and for each task. (C) In experiment 2, participants performed SP movements consisting in simultaneously pressing two keyboard buttons with the index fingers of their two hands. An algorithm was run in parallel to characterize the times at which movements were performed in the non-TMS trials. This information was in turn used to distribute TMS pulses in subsequent TMS trials with different time intervals between the stimuli and the movements.

Figure 2. Means and SEs of the movement times (relative to non-TMS trials) for the RT (green) and PT (red) tested in Experiment 1. (** indicates $P < 0.01$)

Figure 3. Normalized means and SEs of the FDI (left panel) and ADM (right panel) MEP amplitudes in RT (green) and PT (red) movements tested in Experiment 1. The found significant difference in MEPs between baseline and -200 ms TMS time conditions is indicated as well. (** indicates *P* < 0.01)

Figure 4. Upper and lower confidence limits (black lines) of the observed intervals between the TMS trigger times and the subsequent movement onset times in the SP paradigm (green). The simulated frequencies based on the non-TMS trials are represented in grey. The horizontal red bars represent time points in which a significant ($P < 0.01$) difference between the two distributions is observed.

Figure 5. Changes in the FDI (A) and ADM (C) MEPs across time before movements in the SP paradigm. Grey dots show normalized MEP amplitudes (all subjects and trials). Red traces represent the upper and lower confident limits obtained using a bootstrap analysis with all data points ($p < 0.001$). Bars at the bottom identify three intervals of interest used to extract MEPs for a subsequent group level rmANOVA. These are aimed to contain MEPs observed at baseline (grey), and during the periods of reduced excitability –black- and movement initiation phase (white). Normalized group means and SEs of FDI (B) and ADM (D) MEP amplitudes in the three selected intervals of interest.