Title

Structural insights into substrate recognition by the SOCS2 E3 ubiquitin ligase

Authors

Wei-Wei Kung*, Sarath Ramachandran*, Nikolai Makukhin*, Elvira Bruno, Alessio Ciulli

*These authors contributed equally to this work

Address

Division of Biological Chemistry and Drug Discovery, School of Life Sciences, University of Dundee, James Black Centre, Dow Street, Dundee, DD1 5EH, United Kingdom

Abstract

The suppressor of cytokine signaling 2 (SOCS2) acts as substrate recognition subunit of a Cullin5 E3 ubiquitin ligase complex. SOCS2 binds to phosphotyrosine-modified epitopes as degrons for ubiquitination and proteasomal degradation, yet the molecular basis of substrate recognition has remained elusive. We solved co-crystal structures of SOCS2-ElonginB-ElonginC in complex with phosphorylated peptides from substrates growth hormone receptor (GHR-pY595) and erythropoietin receptor (EpoR-pY426) at 1.98 Å and 2.69 Å, respectively. Both peptides bind in an extended conformation recapitulating the canonical SH2 domain-pY pose, yet capture different conformations of the EF loop via specific hydrophobic interactions. The flexible BG loop, for the first time fully defined in the electron density, does not contact the substrate degrons directly. Cancer-associated SNPs located around the pY pocket weaken substrate-binding affinity in biophysical assays. Our findings reveal insights into substrate recognition and specificity by SOCS2, and provide a blueprint for small molecule ligand design.

Correspondence and requests for materials should be addressed to A.C. (a.ciulli@dundee.ac.uk)

ORCID ID: 0000-0002-8654-1670

Introduction

Cytokines are small glycoproteins that play important roles in the differentiation, development and function of lymphoid and myeloid cells [1]. The Janus kinase (JAK) – signal transducer and activator of transcription (STAT) signaling pathway plays a critical role enabling cells to respond to specific cytokines by regulating gene expression. Suppressor of cytokine signalling (SOCS) proteins, which comprise of cytokine inducible SH2–containing protein (CIS) and SOCS1 – SOCS7, negatively regulate cytokine receptors and inhibit the JAK-STAT signaling pathway [2].

SOCS proteins share a conserved domain architecture comprising of an N-terminal extended SH2 subdomain (ESS) that functions as a bridge at the interface between the SH2 domain and the SOCS box enabling ubiquitination of captured substrate[3–5], followed by a central Src-homology 2 (SH2) domain that recognizes a phosphotyrosine (pY) containing sequence [6], and a C-terminal SOCS box that interacts with the adaptor ElonginB-ElonginC complex (EloBC) [7–9]. All SOCS proteins bind to EloBC and recruit Cullin5 with high specificity, forming different SOCS-EloBC-Cullin5-Rbx2 (CRL5^{SOCS}) E3 ligases that catalyse ubiquitin transfer and subsequent proteasomal degradation of specific substrates, as a mechanism to regulate diverse biological processes [10–13]. SOCS proteins serve as substrate recognition modules that impart substrate specificity to each CRL5^{SOCS} E3 complex.

Expression of SOCS proteins is induced by cytokine stimulation. Upon cytokine binding, the oligomerized receptors activate the JAK family kinases that phosphorylate specific tyrosine residues on the receptor, including the docking sites for the STAT proteins. The docked STAT proteins are sequentially phosphorylated, they dimerize and translocate into the nucleus, initiating gene transcription of several downstream proteins including the SOCS proteins. SOCS proteins suppress the JAK-STAT pathway via three distinct but often concomitant mechanisms: 1) KIR mediated direct JAK inhibition [3,14]; 2) Blocking STAT activation by competing for receptor pY sites [15]; 3) Targeting the receptor for proteasomal degradation via SOCS E3 ligase activity [16,17]. Some of the SOCS-substrate interactions have been structurally characterized, including SOCS1-JAK [18], SOCS3-gp130 [4,19], SOCS3-gp130-JAK2 [20] and SOCS6-cKit [21].

SOCS2, one of the members of the SOCS family, is implicated in the disorders in immune system, central nervous system and cancer, and is thus emerging as a promising target for

cancer therapies [22–25]. SOCS2 has been shown as the primary suppressor of growth
hormone (GH) pathway where a gigantism phenotype was observed in a *SOCS2*[-/-] mice [26].
Paradoxically, the SOCS2 overexpressed transgenic mice also led to the same phenotype
[27]. Attenuation of GHR signalling relies on two phosphorylation sites at GHR that are
recognised by SOCS2 [16,28]. The pY487 site of GHR interacts with CRL5[SOCS2] E3 ligase
that targets the GHR for ubiquitination and proteasomal degradation [16]. A downstream
pY595 site interacts with SOCS2, STAT5b and SHP2 (SH2 domain-containing
phosphatase 2), enabling SOCS2 to inhibit the signaling by blocking this receptor site from
STAT5b [15,27,29,30]. Nonetheless, deletion of both sites is required to remove the inhibitory
effect of SOCS2 on the GH signaling [16,28]. Analysis of the binding affinity of SOCS2 for
these two phosphorylation sites of GHR reveals that the pY595 region exhibits a higher
affinity towards SOCS2 (K_D = 1.6 µM) compared to pY487 region (K_D = 11.3 µM) [5,31,32]. An
11-mer phosphorylated peptide spanning the pY595 region of GHR was sufficient to pull
down the whole CRL5[SOCS2] complex from human cell lysates[31] as well as CIS, the closest
homologue to SOCS2 from the same family, which plays a role in anti-tumor immunity
controlling the differentiation of CD4 T helper cell, and the IL-2 and IL-4 response [31,33]. In
addition to GHR and CIS, other substrates have been identified to interact with SOCS2,
including the erythropoietin receptor (EpoR) at pY426 [34], the leptin receptor at pY1077 [35],
the epidermal growth factor receptor [36] and the insulin-like growth factor-I receptor [37]. The
first crystal structure of SOCS2-ElonginB-ElonginC (SBC) was reported in 2006 [32],
however the structural basis for substrate recognition to SOCS2 has yet remained elusive.
87
Here, we determine the first co-crystal structures of SBC in complex with phosphorylated
epitope peptides from its physiological targets GHR and EpoR. Our structures reveal the
peptides are accommodated in an extended conformation to capture specific interactions
with SOCS2. A key flexible region of SOCS2, known as the BG loop, is defined in the
electron density for the first time and shown not to contact the bound substrates. Structural
analyses supported by biophysical and mutagenesis investigations allowed identification of
hotspot residues on the substrate degrons and functional elucidation of disease-relevant
single nucleotide polymorphisms (SNPs) of SOCS2. Our findings provide fresh insights
into the molecular recognition and selectivity between SOCS2 and target substrates, and
provide an important template for future structure-guided ligand design.
98
Results

To elucidate the molecular basis of substrate recognition by SOCS2, we subjected the

SOCS2-ElonginB-ElonginC (SBC) complex to extensive co-crystallization trials with 11-

residue phosphopeptides of either EpoR or GHR that span the regions surrounding Tyr426

and Tyr 595 region, respectively. The affinity of SBC for EpoR_pY426 (K_D of 6.9 μM) and

GHR_pY595 (K_D of 1.1 μM) was measured by isothermal titration calorimetry (ITC), and

found to be consistent with the literature [31,32] (Figure S1). Attempts to co-crystallize wild-

type SBC protein constructs [31,32] with either GHR or EpoR peptide were unsuccessful as

resulting crystals only diffracted poorly. To improve crystal quality, we engineered a cluster

of three mutations K115A/K117A/Q118A on SOCS2 that was predicted to significantly

reduce surface conformational entropy and thermodynamically favor crystal packing [38].

Crystallization attempts with this new S^{KKQ}BC triple-mutant construct

(K115A/K117A/Q118A on SOCS2) eventually yielded high-resolution datasets.

SBC-EpoR co-crystal structure

The structure of SBC in complex with EpoR_pY426 peptide (SBC-EpoR) was solved and

refined at 2.69 Å with 19.64 % R_{work} and 23.51 % R_{free} (Table 1). The overall subunit and

domain arrangements of the SBC-EpoR structure is consistent with those of the apo SBC

structures [32,39,40] (Figure 1a). Electron density for nine out of eleven non-terminal

EpoR_pY426 residues are well defined in the structure (Figure 1b). A classic SH2 domain-

pY peptide interaction is observed where the pY residue is anchored at the pY pocket and

the flanking residues are extending across the SH2 domain. The pY residue is tightly

locked by an intricate hydrogen-bonding network formed by residues Arg73, Ser75, Ser76,

Thr83 and Arg96 of SOCS2 (Figure 1c). Additional hydrogen bonds are formed along the

backbone of EpoR_pY426 peptide from Glu(-1) to Leu(+3) with SOCS2 residues Thr93,

Asn94, Asp107 and one structural water (Figure 1d). Multiple hydrophobic interactions

also support the binding of EpoR_pY426 C-terminal residues, Ile(+2), Leu(+3) and Pro(+5)

that are well accommodated within a hydrophobic patch created by Leu95, Leu106,

Ser108, Ile109, Val112, Leu116 and Leu150 of SOCS2 (Figure 1e).

Crystal structure of SBC-GHR

Encouraged by the success in solving an SBC-EpoR structure, to deepen understanding

of the SOCS2 binding epitopes, we co-crystallized SBC with an 11-mer GHR_pY595

peptide (SBC-GHR) and solved the structure at 1.98 Å resolution with 19.00 % R_{work} and

22.66% R_{free} (Table 1). In contrast to SBC-EpoR, which contains one protomer in the

asymmetric unit, the SBC-GHR contains two copies of protomer. Alignment of these two

protomers via the backbone atoms of the EloB subunit reveals a hinge motion between the

SH2 domain and the SOCS box (Figure S2). Such motion is a common feature in SOCS

box and F-box containing proteins and it has been shown to be important as it facilitates

accurate orientation and positioning of a target substrate protein relative to the

multisubunit CRL complex [39,41]. Contrary to the SBC-EpoR complex, which contains a

single copy of peptide per SH2 domain, two copies of GHR_pY595 peptides were found

binding per SH2 domain of SOCS2 (giving a total of four copies within the asymmetric

unit). The two peptides run in an anti-parallel direction relative to each other across the

SH2 domain, with well-defined electron density surrounding them both (Figure 2a). One of

the peptides (referred to as peptide A hereafter) binds to SH2 domain in a canonical

manner, where the pY is recognized by the positively charged pY pocket between the

central β strands and αA (Figure 2b). In contrast, the second peptide, peptide B, has its pY

residue exposed to solvent and interacting only with His149 of SOCS2 (Figure 2b).

Specific interaction of the GHR_pY595 phosphopeptide

The unusual simultaneous binding observed for the GHR substrate peptide to SOCS2

SH2 domain is imparted mainly by the region comprising Ser(+2) to Val(+6) from each

peptide, which pair such that they form an anti-parallel beta sheet (Figure 2c). Extensive

hydrogen bonds are formed between the backbone of the two peptides and backbone

residues of SOCS2 and structural waters (Figure 2c). Further hydrophobic interactions

appear to reinforce the binding, which impart specificity for GHR. The Ile(+3) and Ile(+5) of

peptide A and peptide B settle in a hydrophobic patch of the SH2 domain formed by

Leu95, Leu106, Ser108, Leu116 and Leu150 (Figure 2d). Another hydrophobic interaction

that is distinct in SBC-GHR compared to SBC-EpoR is that formed by the side chain of

Val(-3) of peptide A, that nicely fits into a hydrophobic pocket comprising of Thr88, Ala90,

Thr93, Leu95 and Val148 from SOCS2 (Figure 2e). In the SBC-GHR structure, a cobalt

ion is modeled at a positive peak that disappeared only at 21 σ level in the unbiased Fo-Fc

electron density map. This cobalt ion satisfies the formation of an octahedral coordination

geometry with the side chains of His(+4) of peptide B, His149 of SOCS2 and with three

surrounding water molecules (Figure 2f).

The BG loop of SOCS2 is observed for the first time in an open conformation

SOCS2 is known to recognize two GHR binding sites at regions around pY487 and pY595,

respectively. We therefore hypothesized that the two copies of the GHR peptides bound in

the crystal structure might mimic a physiological folded conformation of GHR, presenting

each of the phosphorylated sites bound simultaneously to SOCS2. To test this hypothesis,

we utilized an 11-residue GHR_pY487 phosphopeptide (NIDFpYAQVSDI, K_D of 2.3 μM by

ITC, Figure S1), mixed with the GHR_pY595 peptide and SBC in equimolar 1:1:1 ratio for

co-crystallization. In this crystal structure (hereafter referred to as SBC-GHR$_2$), still two

copies of the GHR_pY595 peptide, but no GHR_pY487, are observed bound, yielding a

structure very similar to the previous SBC-GHR structure (Figure S3). However, an

important observation in SBC-GHR$_2$ compared to our other co-crystal structures was that

the regions of SOCS2 corresponding to residues 134-162 is now fully visible in the

electron density. This is unprecedented, and the first time this region, also called BG loop

is structurally defined in full. The BG loop connects the βB and βG strands of an SH2

domain (Figure 3a). The first part of the BG loop (residues 134-148 in SOCS2) differs in

length and sequence among SOCS proteins (Figure 3b). We refer herein to this more

variable region as the "specificity BG loop", because its conformation, together with the

configuration of the adjacent EF loop, governs accessibility of the pY binding pocket and

contributes to substrate specificity in SH2 domains [42]. A particular region in the middle of

the specificity BG loop (residues 136-145) is found to be disordered in all previously

determined SOCS2 structures (PBD code, 2C9W, 4JGH and 5BO4) as well as our other

co-crystal structures SBC-GHR and SBC-EpoR. In this SBC-GHR$_2$ structure, the BG loop

is in an open conformation stabilised by crystal contacts, Pro140 (BG loop) to Arg186

(SOCS box) and Pro140 (BG loop) to Ile90 (EloB), as clearly defined by the unbiased omit

map at this region (Figure 3c, Table 1).

Conformational changes of the EF and BG loop

The configuration of the EF and BG loops play an important role in governing the

accessibility of the binding pocket and specificity toward ligand binding [42]. A comparison of

the SOCS2 structures in the presence and absence of peptides bound highlight

conformational changes in EF (residue 107-116) and BG loop. In the absence of substrate

peptide, the EF loop curls up placing the Ile110 and Cys111 at the hydrophobic SH2

domain (Figure 4a). Upon binding of a substrate peptide, the EF loop opens up forming

backbone interactions with GHR_pY595 (Figure 4b), or rearranges itself to allow a specific

interaction with EpoR_pY426 (Figure 4c). This specific interaction between EF loop and

EpoR_pY426 involves hydrophobic interactions between Ile109 and Val112 of SOCS2,

Val112 of a SOCS2 symmetry mate, and Pro(+5) of EpoR_pY426, resulting in a

differential binding mode from GHR_pY595 to SOCS2. The BG loop of SOCS2 is

observed to be in an open conformation in SBC-GHR$_2$ and SBC-EpoR (disordered)

structures. A superimposition of the two substrate complex structures suggest that the BG

loop opens up further in SBC-GHR structure to accommodate two GHR peptides (Figure

4d).

Biophysical characterization of specificity between GHR_pY595 and SOCS2

To evaluate the specificity of the protein-peptide interaction in solution, we designed

single-point mutations on the peptide first, and compared their binding to wild-type peptide

by two orthogonal biophysical methods: a direct binding assay using SPR (SBC

immobilized on the chip) and ^{19}F ligand-observed displacement NMR assays. In the ^{19}F

NMR displacement assay, the fluorine signal of a purposely-designed reporter ligand (also

referred to as spy molecule) was monitored as a mean to quantify the extent of the

competition between the tested peptides and the spy molecule. A Carr-Purcell-Meibom-Gill

(CPMG) pulse sequence was applied to estimate the spin-spin relaxation time (T_2) of the

spy molecule in the absence and presence of protein [43–45]. By adding competitor to disrupt

the protein-spy interaction, the binding affinity of a competitor can be calculated based on

the degree of displacement of the spy[46,47]. The spy molecule used in our assay is

phosphate **3**, a fluorinated pY analogue that specifically binds to the pY pocket with a K_D of

50 μM (Scheme 1, and Figure S4). The two assays were found to be robust and reliable

where the measured K_D and K_i values at R^2 of 0.74 correlation (Figure S5).

First, we focused on the unique interaction formed by Val(-3) of GHR_pY595, which

inserts into a small hydrophobic cavity of SOCS2 (Figure 2e). This interaction was

investigated by mutating Val(-3) in the GHR peptide to Tyr and Arg, as representative

bulky and charged residues, respectively. We hypothesized this structural change would

disrupt the fit at this small hydrophobic pocket. Mutant V(-3)R exhibited between a 6- and

a 14-fold loss of binding affinity to SBC, depending on the assay, suggesting the charged

group strongly disrupts the interaction (Table 2). By contrast, the V(-3)Y was less

disruptive, with only a two-fold loss in affinity.

Next, to map the relative importance and contribution of each individual amino acids to the

binding affinity with SBC, alanine scan of the substrate peptides was invoked. The peptide

sequences were designed such that individual amino acids were separately mutated into

alanine except pY, which is known to abolish binding if mutated even to unphosphorylated

Y [31]. The resulting library comprised of the original wild-type sequences, ten derivatives

from GHR_pY595 and nine from EpoR_pY426, and was characterized in parallel using

SPR and [19]F NMR competition assay (Table 3). Alanine substitution at pY(-3), pY(-1),

pY(+3) and pY(+4) of the GHR_pY595 resulted in at least two-fold weakened binding

(increase in K_D) compared to the wild type. In contrast, a similar 2-fold weakening in

binding affinity was observed in the EpoR_pY426 peptide upon alanine substitution at pY(-

1), pY(+2) and pY(+3). These results are consistent with observations from our crystal

structures that peptide-SOCS2 binding is mediated by hydrophobic interaction including

pY(-3), pY(+3) and pY(+5) on the GHR_pY595 and pY(+2) and pY(+3) on the

EpoR_pY426. The binding affinity for each peptide also dropped by at least two-fold with

Ala substitution at pY(-1) position, indicating the importance of the residue just upstream of

pY.

SNPs study

Several single nucleotide polymorphisms (SNPs) on SOCS2 are reported in the Catalogue

of Somatic Mutation in Cancer database (COSMIC) as potentially linked to cancers such

as tumours of the lung, breast, and pancreas [48]. We thus next decided to characterize the

interaction of selected SNP SOCS2 mutants with substrate peptides GHR_pY595,

EpoR_pY426 and GHR_pY487 by SPR. Inspection of our SBC co-crystal structures

guided us to select five known SNPs: N94D, R96L, R96Q that are located in the pY-pocket

and involved in direct recognition of pY; L106V that is highly conserved at the hydrophobic

SH2 domain involving in substrate interaction; and C133Y that participates in the SH2

hydrophobic core (Figure 5a). All mutant proteins expressed and purified similarly to wild-

type, and the mutations did not appear to affect the structural integrity and solubility of the

constructs, as observed by [1]H NMR (Figure S6).

The L106V and C133Y mutations did not affect binding affinities compared to wild type

(Figure 5b). In contrast, the N94D and R96L mutations drastically impaired substrate

binding, leading to almost undetectable binding response by SPR. Because of the low

signal-to-noise, reliable K_D values could not be measured with these protein mutants. For

the R96Q mutation no signal response was detected, suggesting that binding was

completely abolished and highlighting the most disruptive of the mutation studies.

Mutations at R96 site abrogate SOCS2 binding was also reported by Rupp *et al.* with

R96C [49]. In addition to SPR, the SNP mutants were characterized using [19]F NMR by

monitoring the transverse relaxation rate (R_2). L106V and C133Y exhibited similar rate, 10

s^{-1} and 14 s^{-1}, comparable to wild type at 11 s^{-1}, suggesting these mutations did not affect

pY recognition. Mutations on N94 and R96 showed a substantial disruption to pY

recognition, resulting in a significant disruption of binding as the no protein control at

around 3 to 4 s^{-1}, consistent with the SPR data (Figure 5c).

Discussion

SOCS2 is a substrate binding protein of the CRL5 E3 complex that negatively regulates the JAK-STAT signaling by targeting substrate receptors for degradation and blocking STAT5b activation by competing with receptor pY sites. The details of these interactions have remained elusive and to date structural information remained limited to apo SOCS2. Herein, we have disclosed two novel structures of the SBC in complex with substrate peptides EpoR_pY426 and GHR_pY595. Both peptides recapitulate a canonical substrate-binding mode to SH2 domain of SOCS2. Residues at pY(-1), pY(+1) and pY (+3) positions of the GHR_pY595 and EpoR_pY426 peptide contains similar properties; whereas residues at pY(-3), pY(+2) and pY(+5) position are different in properties and sizes (Figure 6a). In particular, the Val(-3) of GHR_pY595 and Pro(+5) of EpoR_pY426 peptide catch different hydrophobic interactions resulting in exclusive binding modes in SOCS2 comparing with the substrate peptides of SOCS3 and SOCS6 (Figure 6b, 6c) [4,21].

The BG loop of SOCS2 had not been fully revealed in previous published structures. Here, we report an open conformation BG loop, which is contradictory to other SOCS structures with bound peptides, for example SOCS3:gp130 and SOCS6:c-kit [19,21]. In the SOCS3 and SOCS6 peptide-bound structures, the BG loop folds up as a hairpin interacting with the substrate peptide, forming a triple-stranded β sheet structure (Figure 7a, 7b). The corresponding BG loop region in SOCS2 is either fully disordered or in an open conformation (SBC-EpoR and SBC-GHR$_2$), suggesting that this region does not participate in substrate recognition. Nevertheless, interestingly, a similar triple-stranded β sheet structure is observed in the SBC-GHR$_2$ structure, where the peptide B replaces the first β-sheet of the BG loop and makes backbone interactions with BG loop and peptide A (Figure 7c). Additionally, a tyrosine phosphatase SHP-2, which contains SH2 domains, features a comparable structure [50]. Its BG loop folds up as a hairpin and forms a triple-stranded β sheet interaction with two bound peptides (Figure 7d). Although the canonical phosphotyrosine binding site is conserved for both SOCS2 and SHP2, the positioning of the non-canonical phosphotyrosine varies significantly. The distance between Cα atoms of phosphotyrosines within the bound peptides are 23 Å and 13 Å in SBC-GHR and SHP2-substrate co-structures, respectively (Figure 7e). Longer distance between the canonical and non-canonical pY enable antiparallel beta strand interactions between four amino acids of GHR peptides unlike antiparallel interactions between two amino acids in SHP-2 substrate peptides. Furthermore, a longer EF loop in SOCS2 pushes the two GHR peptide, away from itself towards the open BG loop, enabling β sheet formation.

The BG loop along with the EF loop forms a hydrophobic channel in SOCS3 and SOCS6.

This channel imparts specificity and restricts the binding of substrates. In contrast the open

conformation of BG loop in SOCS2 could be critical in enabling SOCS2 to accommodate a

wider range of substrates including GHR, EpoR, SOCS1 and SOCS3 amongst others. A

comparison of buried surface area of the substrate peptides among SOCS proteins, reveal

that EpoR and GHR binds with SOCS2 with only 595 Å^2 and 641 Å^2, respectively, in

contrast to areas of 1714 Å^2 for SOCS6/c-KIT and 1761 Å^2 for SOCS3/gp130 complexes.

Unlike SOCS3 and SOCS6 complex structures, the pY flanking residues from EpoR and

GHR do not participate in extensive side-chain hydrogen bonding interactions. Together,

these observations are consistent with greater binding affinities of SOCS3 and SOCS6

substrates compared to SOCS2 substrates. The lower potency of affinity for SOCS2

substrates could contribute to its relatively greater promiscuity to multiple substrates.

The observation of the dual-peptide binding mode to SOCS2 was unexpected, however is

not unprecedented with SH2 domains, as reported previously with the tyrosine

phosphatase SHP-2 [50]. In the co-crystal structure of SHP-2:pY peptide solved by Zhang et

al., one pY of the peptide is recognized at the pY pocket and the other one is solvent

exposed as in our structures described herein. Besides, two peptides run antiparallel to

each other and form an antiparallel four-stranded β sheet with BG loop. Zhang *et al*.

suggested that the dimerization of peptide binding in SHP-2 requires at least one pY

containing peptide and demonstrates enhanced binding affinity to protein. For our SBC-

GHR structure, despite preparing several protein-peptide samples for co-crystallization at

1:1 molar ratio, all dataset collected from crystals were consistent with a 1:2 (protein-

peptide) binding mode. In both cases, the binding of the non-canonical binding peptide

relies on hydrophobic interaction with the SH2 domain hydrophobic core, and backbone to

backbone interaction with the canonical binding peptide and protein (Figure 6d). The 1:1

binding mode for EpoR could be justified by the presence of Pro(+5) in EpoR, that acts as

a strand breaker and prevents backbone to backbone interaction with the second peptide.

Indeed, from the alanine scan (Table 3), we observe that when Pro(+5) is replaced by

alanine, the affinity improves - implying the possibility of secondary peptide binding.

We put forth two distinct models that might explain the dual peptide recognition mode and

its role in specific tuning of GHR signaling response. First, a "cis" recognition mode, where

the GHR tail folds back as a hairpin structure presenting two binding epitopes around

distinct phosphorylation sites (e.g. pY487 and pY595) for recognition (Figure 8a).
However, the crystallography data from our follow-up experiment as described in the SBC-
GHR$_2$ structure is not consistent with this hypothesis, as two instances of the pY595
peptide were found bound despite a molar ratio of 1:1 for pY487 and pY595 peptides
being present in the co-crystallization buffer. However, we cannot exclude that
simultaneous recognition of the two distinct epitopes would require a loop of the same tail
twisting back onto itself to enhance the binding affinity of second epitope. Alternatively, we
envisage a "trans" recognition mode, where SOCS2 recognize two separate receptor tails
of the activated dimerized GHR receptors at the cell membrane (Figure 8b). SOCS2 might
additionally play a role as scaffold bringing two substrates in close proximity, for example
by recruiting one instance of phosphorylated substrate to assist the binding of un-
phosphorylated one for post-translational modification. This mechanism evokes potential
similarities with some phosphodegrons which require two sites to be phosphorylated,
utilizing a first kinase to "prime" phosphorylation events, followed by a second kinase for
follow-on phosphorylation [51]. An example of such a mechanism is the β-catenin
degradation mediated by the β-TrCP [52]. Further biophysical investigation is warranted to
address the extent to which these potential mechanisms might be invoked for SOCS2
function.
367
SOCS2 is an attractive therapeutic target due to its links to cancer, diabetes, neurological
and inflammatory diseases [23,24,53–57]. Breast, lung, liver and ovarian cancer have been
correlated with down-regulation in SOCS2 [58–63]. In addition to the JAK-STAT pathway, a
recent study has identified the involvement of SOCS2 in the NF-κB (nuclear factor kappa-
light-chain-enhancer of activated B cells) pathway that regulates the immune and
inflammatory responses [64,65]. NF-κB is found to be constitutively activated in many types of
cancer and influences a diverse array of pro-tumorigenic functions, therefore NK-κB plays
an pivotal role in cancer initiation and progression [66]. SOCS2 negatively regulates TNFα
induced NF-κB activation by targeting NDR1, a serine-threonine kinase, for proteasomal
degradation. Hence SOCS2 deficiency may lead to an increased level of NDR1, which
was is reported to result in aggressive behaviour of PC3 prostate cancer cells [64]. These
evidences highlight the potential in targeting SOCS2 for drug discovery for inflammation
and cancer biology. We have revealed structural insights into the SOCS2-peptide
interactions by X-ray crystallography and identified hotspot using alanine scanning,
mutation study and SNPs study. This information provides a template to guide the
structure-based rational design of SOCS2 ligands that are instrumental in the development

of novel chemical tools to address biological question of SOCS2, and in the quest for
novel small molecule ligands binding to SOCS2 as potential therapeutics. SOCS2 binders
at the pY binding pocket can be used as inhibitors of the CRL5^{SOCS2}, which would be
expected to prevent degradation of target substrate receptors, thus prolonging the activity
of cytokine signaling pathway and upregulating expression of endogenous STAT5b-
responsive gene expression. In a distinct application, a SOCS2 binder could be used as
E3 ligase ligand handle for designing new chemical degraders to hijack SOCS2 CRL
activity and trigger the degradation of unwanted proteins inside cell [67–69]. This approach,
also known as proteolysis targeting chimeras (PROTACs), offers the advantage of
inducing rapid and selective intracellular depletion of the target protein, as opposed to
mere blockade of a single interaction or activity, which pairs more closely to genetic target
validation and often results in greater maximal efficacy of intervention in a signaling
pathway. PROTAC-mediated protein degradation has been shown to occur at very low
compound concentration (pM to nM range) also potentially allows targeting of intractable
protein targets that are beyond the reach of conventional small-molecule approaches that
require full occupancy of a target binding site e.g. receptor antagonists and enzyme
inhibitors. A limited set of E3 ligases have been targeted so far for PROTACs, notably VHL
[70,71] and cereblon [72,73], so extending the approach to other ligandable E3 ligases would be
an important advance to the field. The usage of SOCS2 binders for PROTACs may be
limited to SOCS2 expressing cells, however, this provides an additional layer of tissue
specific degradation of target proteins. PROTACs designed using SOCS2 binders as
recruiters would be ideal for degrading targets specifically in cells affected with leukemia
and gastrointestinal sarcoma which have been recorded to have upregulated SOCS2
expression [87]. Low levels of SOCS2 in other normal tissues would help minimize toxicity.
In addition, a pan-SOCS recruiting PROTACs would help diversifying the target tissue
range. Our peptide-bound co-crystal structures suggest that SOCS2 might be ligandable
and provide a blueprint for the rational structure-guided design of novel SOCS2 inhibitors
and SOCS2 ligand handles for PROTACs.
412
413
414

Materials and Methods

Cloning and protein expression

The human SOCS2 (amino acids 32 – 198) and the ElonginB (amino acids 1 – 104) and ElonginC (amino acids 17 – 112) plasmids were used for protein expression as previously reported [39], and as templates for mutagenesis. SOCS2 mutants with N94D, R96L, R96Q, L106V or C133Y mutation were introduced using a PCR-based method site-directed mutagenesis. SOCS2 wild type and mutants were co-expressed and purified as previously described [31,39].

Crystallization and structure determination of SBC-GHR

To improve crystallization, surface entropy reducing mutations were introduced into SOCS2 construct (amino acids 32 – 198). Three mutation clusters (K63A/E64A/E67A; K113A and K115A/K117A/Q118A) were identified with the SER server [38]. SER-assisted crystallization attempts yielded crystals with the K115A/K117A/Q118A SOCS2-EloBC (S^{KKQ}BC). Five times molar excess of GHR_pY595 (PVPDpYTSIHIV-amide) was incubated with S^{KKQ}BC, followed by removing unbound peptide using a protein concentrator. Sample was concentrated to 22 mg/ml with an additional 0.1 M of sodium cacodylate pH7.2 added to the sample. Diffraction-quality crystals were obtained with 0.005 M Cobalt (II) chloride, 0.1 M MES pH 6.5, 1.0 M ammonium sulphate at 4 °C using hanging drop vapour diffusion method at 2:1 protein:precipitant ratio. Crystals were cryo-protected using 20% MPD prior to vitrification in liquid nitrogen.

Diffraction data were collected at 100 K at Diamond Light Source beamline i04 using Pilatus 6M-F detector at 0.98 Å wavelength. Indexing and integration was processed by XDS [74] and scaling and merging with AIMLESS within the CCP4 program suite [75,76]. The experimental phases was obtained by identifying positions of arsenic atom using MR-SAD Phases in the PHENIX software suite [77,78] with a model provided. The provided template was a lower resolution SBC-GHR co-crystal obtained previously (unpublished work). The structure was reconstructed by AutoBuild [79,80] and manually built in Coot [81]. The resulting structure was refined iteratively with REFMAC5 [82].

Crystallization and structure determination of SBC-EpoR

Five times molar excess of EpoR_pY426 (ASFEpYTILDPS-amide) was incubated with S^{KKQ}BC (5mg/ml). Unbound peptide was removed by a protein concentrator (sartorius Vivaspin) while the mixture was concentrated to 20 mg/ml concentration. Sodium

cacodylate pH7.2 was added to a final concentration of 0.1 M prior to crystallisation.

Crystallisation drops were set up in a ratio of 1:1 protein:precipitant in 18% ethanol, 0.1M

HEPES pH7.5, 0.1M $MgCl_2$ using hanging drop at 4°C. Crystals were cryo-protected using

20 % PEG400 prior to flash-cooled.

Diffraction data were collected at 100 K on beamline i24 at Diamond Light Source. Data

were recorded to Pilatus3 6M-F detector at 0.97 Å wavelength. Data were indexed,

integrated, and reduced using XDS [74] and AIMLESS [75,76]. The phase was obtained by

molecular replacement (MR) using Phaser [78] with the coordinates of SOCS2-EloB-EloC

(PBD ID: 2C9W) as a search model. The presence of the EpoR_pY426 was observed in

the initial electron density map. Model building was conducted manually with Coot [81] and

refined with cycles of retrained refinement with REFMAC5 [82].

Crystallization and structure determination of SBC-GHR$_2$

GHR_pY595 (PVPDpYTSIHIV-amide) and GHR_pY487 (NIDFpYAQVSDI-amide) were

mixed with S^{KKQ}BC at 1:1:1 stoichiometric ratio with a final concentration of 20 mg/ml and

additional 0.1 M sodium cacodylate pH7.2. Drops of the complex were mixed 2:1 with

0.005 M cobalt chloride, 0.1 M MES pH6.5 and 1.0 M ammonium sulphate in the sitting-

drop vapor diffusion format at 4°C. 20 % MPD was applied to crystal before flash-cooling.

Data collection of the SBC-GHR$_2$ co-crystal was at 100 K on beamline i24 at Diamond

Light Source. Images were indexed, intergraded and reduced using XDS [74] and AIMLESS

[75,76]. A molecular replacement solution was obtained by Phaser [78] using SBC-GHR as

search model. Refinement was performed using REFMAC5 [82] and model building was

performed in COOT [81].

Synthetic details

All chemicals, unless otherwise stated were commercially available and used without

further purification. Solvents were anhydrous and reactions preformed under positive

pressure of nitrogen. Flash column chromatography was performed using a Teledyne Isco

Combiflash Rf or Rf200i. As prepacked columns RediSep Rf Normal Phase Disposable

Columns were used. NMR spectra were recorded on a Bruker 500 Ultrashield. ^{13}C spectra

were ^1H decoupled. Chemical shifts (δ) are reported in ppm relative to solvent (CD$_3$OD: δ_H

= 3.31ppm, δ_C = 49.0 ppm) as internal standard. Low resolution MS and analytical HPLC

traces were recorded on an Agilent Technologies 1200 series HPLC connected to an

Agilent Technologies 6130 quadrupole LC/MS, connected to an Agilent diode array detector. The column used was a Waters XBridge column (50 mm × 2.1 mm, 3.5 μm particle size) and the compounds were eluted with a gradient of 5−95% acetonitrile/water + 0.1% formic acid over 3 min. Preparative HPLC was performed on a Gilson Preparative HPLC System with a Waters X-Bridge C18 column (100 mm x 19 mm; 5 μm particle size) and a gradient of 5 % to 95 % acetonitrile in water over 10 min, flow 25 mL/min, with 0.1 % formic acid in the aqueous phase.

Cbz-O-bis(dimethylamino)phosphono)-L-tyrosine (**1**)

O-bis(dimethylamino)phosphono)-*L*-tyrosine [83] (485 mg, 1.54 mmol) and NaHCO$_3$ (260 mg, 3.1 mmol) were dissolved in the mixture THF/H$_2$O = 1:1 (10 mL) and *N*-(benzyloxycarbonyloxy)succinimide (383 mg, 1.54 mmol) was added. The reaction mixture was stirred at room temperature overnight. After the addition of 5% NaHSO$_4$ the product was extracted with ethyl acetate, washed with brine, dried over MgSO$_4$, concentrated by rotary evaporation under reduced pressure. After drying, Cbz-*O*-bis(dimethylamino)phosphono)-*L*-tyrosine **1** (620 mg, 89%) was obtained as pale yellow. ^{1}H NMR (CD$_3$OD): 2.69 (d, *J* = 10.1 Hz, 12H), 2.92 (dd, *J* = 14.0, 9.3 Hz, 1H), 3.18 (dd, *J* = 14.0, 4.9 Hz, 1H), 4.40 (dd, *J* = 4.9, 9.3 Hz, 1H), 5.03 (s, 2H), 7.06 (d, *J* = 8.5 Hz, 2H), 7.21 (d, *J* = 8.5 Hz, 2H), 7.25-7.36 (m, 5H). ^{31}P NMR (CD$_3$OD): 18.2.

(S)-2-amino-3-(4-((bis(dimethylamino)phosphoryl)oxy)phenyl)-N-methylpropanamide (**2**)

To a mixture of the compound **1** (160 mg, 0.35 mmol), HATU (135 mg, 0.35 mmol), HOAt (48 mg, 0.35 mmol) and DIPEA (150 μL, 1 mmol) in DMF (1 mL), 2M methylamine solution in THF (0.5 mL) was added under stirring at room temperature. After two hours, LC-MS analysis showed complete conversion of the starting material and formation of the desired product. The mixture was diluted with ethyl acetate, washed with 5% NaHSO$_4$, brine, dried over MgSO$_4$, concentrated by rotary evaporation under reduced pressure. The crude product was dissolved in the mixture ethanol/ethyl acetate= 1:1 (8 mL). Hydrogenation was carried out using H-Cube at 80 ˚C, Pd/C, 1 atm, at 1mL/min. The solvent was evaporated under vacuum to afford **2** (108 mg, 92%) which was directly used in the next step without any further purification. ^{1}H NMR (CD$_3$OD): 2.66 (s, 3H), 2.71 (d, *J* = 10.1 Hz, 12H), 2.81 (m, 1H), 2.96 (m, 1H), 3.51 (m, 1H), 7.09 (dd, *J* = 8.5, 1.0 Hz, 2H), 7.20 (d, *J* = 8.5 Hz, 2H). ^{31}P NMR (CD$_3$OD): 18.3.

(S)-4-(3-(methylamino)-3-oxo-2-(2,2,2-trifluoroacetamido)propyl)phenyl dihydrogen phosphate (**3**)

A solution of the compound **2** (80 mg, 0.24 mmol) and DIPEA (85 µL, 0.48 mmol) in DCM (2 mL) was cooled to -78 °C, and trifluoroacetic anhydride (34 µL, 0.24 mmol) was added. The reaction mixture was stirred 1h at -78 °C. After solvent evaporation the residue was dissolved in acetonitrile (0.5 mL) and 2M HCl was added (2 mL). The mixture was stirred at room temperature overnight until no presence of the starting materials was detected by LC-MS. The solvents were evaporated and residue was purified by HPLC to afford phosphate **3** (30 mg, 34%) as a white solid. ^{1}H NMR (CD$_3$OD): 2.69 (s, 3H), 2.96 (dd, J = 13.8, 6.4 Hz, 1H), 3.15 (dd, J = 13.8, 6.4 Hz, 1H), 4.57 (dd, J = 8.8, 6.4 Hz, 1H), 7.13 (dd, J = 8.5, 1.1 Hz, 2H), 7.22 (d, J = 8.5 Hz, 2H). ^{13}C NMR (CD$_3$OD): 26.3, 37.8, 56.5, 117.3 (q, J = 286.7 Hz), 121.3 (d, J = 4.5 Hz), 131.3, 134.1, 151.9 (d, J = 6.8 Hz), 158.7 (q, J = 37.5 Hz), 172.4. ^{31}P NMR (CD$_3$OD): 3.7. ^{19}F NMR (CD$_3$OD): -75.6.

Peptide synthesis

All peptides were prepared via solid-phase peptide synthesis on 10 mmol scale using standard Fmoc chemistry on Rink amide resin (0.68 mmol/g) on an INTAVIS ResPepSL automated peptide synthesizer. *O*-(dibenzylphosphono)-*N*-Fmoc-*L*-tyrosine was synthesised as described below. The peptides were cleaved with 2.5% triisopropylsilane and 2.5% water in TFA. The crude peptides were isolated from the cleavage mixture by precipitation with cold ether, dissolved in the mixture water/DMF=1/1 and purified by preparative HPLC under the following conditions: Waters X-Bridge C18 column (100 mm x 19 mm; 5 µm particle size), gradient of 5-95 % acetonitrile in water over 10 min, flow 25 mL/min, with 0.1 % formic acid in the aqueous phase, UV detection at λobs = 190 and 210 nm). The poor soluble peptides were purified according to the literature procedure [84]: the impurities were extracted by DCM from the solution of peptides in 20% acetic acid. The purity and identity of the peptides were determined by the analytical LCMS on an Agilent Technologies 1200 series HPLC connected to an Agilent Technologies 6130 quadrupole LC/MS linked to an Agilent diode array detector.

O-(dibenzylphosphono)-N-Fmoc-L-tyrosine

To a solution of Fmoc-tyrosine (2 g, 5 mmol) in anhydrous THF (12 mL) *N*-methylmorpholine (540 µL, 5 mmol) and *tert*-butyldimethylsilyl chloride (740 mg, 4.9 mmol) were added. After 15 min 4,5-dicyanoimidazole (1.8 g, 15 mmol) and diisopropylphosphoramidite (3.4 mL, 10 mmol) were added and the reaction mixture was

stirred at room temperature for 4 h. After cooling to 0°C 70% *tert*-butyl hydroperoxide (2

ml, 15 mmol) was introduced. After stirring for 2 h at 0°C, 10% $Na_2S_2O_5$ (20 ml) was added

and stirring continued for one more hour. The product was extracted with ethyl acetate,

washed with a 5% solution of $KHSO_4$, brine, dried over $MgSO_4$, concentrated by rotary

evaporation under reduced pressure, and further purified by column chromatography on

silica gel using a gradient elution of 0% to 10% of MeOH in DCM to afford *O*-

(dibenzylphosphono)-*N*-Fmoc-*L*-tyrosine (3g, 90%) as a pale yellow solid. NMR spectra

were in agreement with the published data [85].

Isothermal titration calorimetry

Experiments were performed with ITC200 instrument (Malvern) in 100mM HEPES pH7.5,

50mM NaCl, 0.5mM TCEP at 298K stirring the sample at 750 rpm. The ITC titration

consisted of 0.4 μL initial injection (discarded during data analysis) followed by 19 of 2 μL

injections at 120 seconds interval between injections. The GHR_pY595 peptide

(PVPDpYTSIHIV-amide, 750 μM), EpoR_pY426 (ASFEpYTILDPS-amide, 750/1500 μM)

and GHR_pY487 (NIDFpYAQVSDI-amide, 1500 μM) were directly titrated into SBC (50

μM). Binding data was subtracted from a control titration where peptide was titrated into

buffer, and fitted using a one-set-of-site binding model to obtain dissociation constants,

binding enthalpy (ΔH), and stoichiometry (N) using MicroCal ITC-ORIGIN Analysis

Software 7.0 (Malven).

Surface plasmon resonance

Experiments were performed using Biacore T200 instrument (GH Healthcare) in 20 mM

HEPES pH7.5, 150 mM NaCl, 1mM TCEP, 0.005 % Tween20 buffer at 10 °C. Biotinylated

wild type SBC and mutants were immobilized onto a chip surface and injected a series of

seven concentrations (0.08, 0.25, 0.7, 2.2, 6.7, 20 and 60 μM) of peptide across the

sensor surface for 60 sec contact time and 120 sec disassociation time at 30 μl/min flow

rate. Data analysis was carried out using Biacore Evaluation Software (GE Healthcare). All

data were double-referenced for reference surface and blank injection. The processed

sensograms were fit to a steady-state affinity using a 1:1 binding model for K_D estimation.

^{19}F *Spy NMR displacement*

All NMR experiments were conducted using AV-500 NHz Bruker spectrometer equipped

with a 5 mm CPQCI 1H/19F/13C/15N/D Z-GRD cryoprobe) at 298 K. Spectra were

recorded using a CPMG pulse sequence that attenuates broad resonances with an

interpulse delay at 0.133 seconds for 80 scans. The transmitter frequency was placed

close to the resonance of O_1 = -35451 Hz (-75.3 ppm). Protein was used at 5 μM and spy

molecule (phosphate **3**) was used at 100 μM in buffer containing 20 mM HEPES pH8, 50

mM NaCl, 1mM DTT, 20 % D_2O. The GHR related peptides including wild type, V(-3)R, V(-

3)Y and alanine scanning were used at10 μM, whereas the EpoR related peptide including

wild type and alanine scanning were used at 50 μM. All NMR data were processed and

analyzed using TopSpin (Bruker).

Signal contrast between bound and free form of the spy is calculated as below

$$Contrast\ (\%) = \frac{Spy_{Free} - Spy_{Bound}}{Spy_{Free}}$$

600   where Spy_{free} is the peak integral or T_2 of spy in the absence of protein, while Spy_{bound} is

601   the signal of spy in complex with protein.

603   For competition studies, the percentage displacement is calculated as below

$$Displacement\ (\%) = \frac{Competitor - Spy_{Bound}}{Spy_{Free} - Spy_{Bound}}$$

606   where competitor is the peak integral of spy in the presence of competitor, protein and spy.

608   To calculate the dissociation constant of competitor (Ki), assuming all other experimental

609   conditions are identical in the absence and presence of competitor, the signal intensity is

610   proportional to the concentration of protein-spy complex, and can be expressed as below

611   [47]

$$\frac{Signal(I)}{Signal(0)} = \frac{[Eo] + [Lo] + K_D - [EI] - \sqrt{([Eo] + [Lo] + K_D - [EI])^2 - 4([Eo] - [EI])[Lo]}}{[Eo] + [Lo] + KD - \sqrt{([Eo] + [Lo] + K_D)^2 - 4[Eo][Lo]}}$$

614   where Signal(I) and Signal(0) are the fluorine intensities in the presence and absence of

615   competitor, respectively. [E_0] is the total concentration of protein. [L_0] is the total

616   concentration of ligand. [EL] is concentration of protein-ligand complex. From the observed

617   ratio of signal intensity, the Ki can be calculated.

T$_2$ relaxation measurement

620   Spectra were acquired for each sample at 298K, and 80 scans using a CPMG pulse

621   sequence with a relaxation delay of 0.05, 0.1, 0.2, 0.4 and 0.8 s. Protein was used at 5 μM

622   and spy was used at 100 μM in buffer containing 20 mM HEPES pH8, 50 mM NaCl, 1mM

623   DTT, 20 % D$_2$O. The T$_2$ relaxation time was calculated using Prism by fitting magnetisation

624   to a mono exponential function

$$M_{xy}(t) = M_{xy}(0)e^{-t/T_2}$$

627   where T$_2$ is the time constant of decay of component magnetisation in the xy-plane (M$_{xy}$)

628   that is perpendicular to the external field B$_0$.

Accession Code

The coordinates and structure factors for SBC in complex with EpoR_pY426 peptide, GHR_pY595 peptide, and GHR2_pY595 peptide have been deposited to the Protein Data Bank (PDB) with accession codes 6I4X, 6I5N, and 6I5J, respectively.

Supplemental information

Supplementary Figures S1–S14 are included.

Acknowledgements

This project has received funding from the European Research Council (ERC) under the European Union's Seventh Framework Programme (FP7/2007-2013) as a Starting Grant to A.C. (grant agreement No. ERC-2012-StG-311460 DrugE3CRLs). Biophysics and drug discovery activities at Dundee are supported by Wellcome Trust strategic awards (100476/Z/12/Z and 094090/Z/10/Z, respectively). W.K. was supported by a PhD Scholarship by the School of Life Sciences at the University of Dundee. We thank P. Fyfe for support with in-house X-ray facilities, the Diamond Light Source for beamtime (BAG proposals MX14980-13 and MX14980-20) and P. Romano and P. Aller for support at beamlines. We thank de Castro for support with NMR and helpful discussions.

Competing interests: The A.C. laboratory receives sponsored research support from Boehringer Ingelheim and Nurix, Inc. A.C. is a scientific founder, director and shareholder of Amphista Therapeutics, a company that is developing targeted protein degradation therapeutic platforms.

References

1. Sato, N. & Miyajima, A. Multimeric cytokine receptors: common versus specific
functions. *Curr. Opin. Cell Biol.* **6**, 174–179 (1994).

2. Starr, R. *et al.* A family of cytokine-inducible inhibitors of signalling. *Nature* **387**, 917–
921 (1997).

3. Yasukawa, H. The JAK-binding protein JAB inhibits Janus tyrosine kinase activity
through binding in the activation loop. *EMBO J.* **18**, 1309–1320 (1999).

4. Babon, J. J. *et al.* The Structure of SOCS3 Reveals the Basis of the Extended SH2
Domain Function and Identifies an Unstructured Insertion That Regulates Stability.
Mol. Cell **22**, 205–216 (2006).

5. Bullock, A. N., Rodriguez, M. C., Debreczeni, J. É., Songyang, Z. & Knapp, S.
Structure of the SOCS4-ElonginB/C Complex Reveals a Distinct SOCS Box
Interface and the Molecular Basis for SOCS-Dependent EGFR Degradation.
Structure **15**, 1493–1504 (2007).

6. Yoshimura, a *et al.* A novel cytokine-inducible gene CIS encodes an SH2-containing
protein that binds to tyrosine-phosphorylated interleukin 3 and erythropoietin
receptors. *EMBO J.* **14**, 2816–2826 (1995).

7. Kamura, T. *et al.* The Elongin BC complex interacts with the conserved SOCS-box
motif present in members of the SOCS, ras, WD-40 repeat, and ankyrin repeat
families. *Genes Dev.* **12**, 3872–3881 (1998).

8. Zhang, J. G. *et al.* The conserved SOCS box motif in suppressors of cytokine
signaling binds to elongins B and C and may couple bound proteins to proteasomal
degradation. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 2071–2076 (1999).

9. Kamura, T. *et al.* VHL-box and SOCS-box domains determine binding specificity for
Cul2-Rbx1 and Cul5-Rbx2 modules of ubiquitin ligases. *Genes Dev.* **18**, 3055–3065
(2004).

10. Piessevaux, J., Lavens, D., Peelman, F. & Tavernier, J. The many faces of the
SOCS box. *Cytokine and Growth Factor Reviews* (2008).
doi:10.1016/j.cytogfr.2008.08.006

11. Babon, J. J., Sabo, J. K., Zhang, J. G., Nicola, N. A. & Norton, R. S. The SOCS Box
Encodes a Hierarchy of Affinities for Cullin5: Implications for Ubiquitin Ligase
Formation and Cytokine Signalling Suppression. *J. Mol. Biol.* **387**, 162–174 (2009).

12. Linossi, E. M. & Nicholson, S. E. The SOCS box-adapting proteins for ubiquitination
and proteasomal degradation. *IUBMB Life* **64**, 316–23 (2012).

13. Bulatov, E. & Ciulli, A. Targeting Cullin–RING E3 ubiquitin ligases for drug discovery:
structure, assembly and small-molecule modulation. *Biochem. J.* **467**, 365–386
(2015).

14. Sasaki, A. *et al.* Cytokine-inducible SH2 protein-3 (CIS3/SOCS3) inhibits Janus
tyrosine kinase by binding through the N-terminal kinase inhibitory region as well as
SH2 domain. *Genes to Cells* **4**, 339–351 (1999).

15.   Ram, P. A. & Waxman, D. J. SOCS/CIS protein inhibition of growth hormone-
694   stimulated STAT5 signaling by multiple mechanisms. *J. Biol. Chem.* **274**, 35553–
695   35561 (1999).

696   16.   Vesterlund, M. *et al.* The SOCS2 ubiquitin ligase complex regulates growth hormone
697   receptor levels. *PLoS One* **6**, (2011).

698   17.   Piessevaux, J., De Ceuninck, L., Catteeuw, D., Peelman, F. & Tavernier, J. Elongin
699   B/C recruitment regulates substrate binding by CIS. *J. Biol. Chem.* (2008).
700   doi:10.1074/jbc.M803742200

701   18.   Liau, N. P. D. *et al.* The molecular basis of JAK/STAT inhibition by SOCS1. *Nat.*
702   *Commun.* (2018). doi:10.1038/s41467-018-04013-1

703   19.   Bergamin, E., Wu, J. & Hubbard, S. R. Structural Basis for Phosphotyrosine
704   Recognition by Suppressor of Cytokine Signaling-3. *Structure* **14**, 1285–1292
705   (2006).

706   20.   Kershaw, N. J. *et al.* SOCS3 binds specific receptor-JAK complexes to control
707   cytokine signaling by direct kinase inhibition. *Nat. Struct. Mol. Biol.* **20**, 469–476
708   (2013).

709   21.   Zadjali, F. *et al.* Structural basis for c-KIT inhibition by the suppressor of cytokine
710   signaling 6 (SOCS6) ubiquitin ligase. *J. Biol. Chem.* **286**, 480–490 (2011).

711   22.   Haan, S. SOCS2 physiological and pathological functions. *Front. Biosci.* **8**, 760
712   (2016).

713   23.   Keating, N. & Nicholson, S. E. SOCS-mediated immunomodulation of natural killer
714   cells. *Cytokine* (2018). doi:10.1016/j.cyto.2018.03.033

715   24.   Turnley, A. M. Role of SOCS2 in growth hormone actions. *Trends in Endocrinology*
716   *and Metabolism* (2005). doi:10.1016/j.tem.2005.01.006

717   25.   Rico-Bautista, E., Flores-Morales, A. & Fernández-Pérez, L. Suppressor of cytokine
718   signaling (SOCS) 2, a protein with multiple functions. *Cytokine Growth Factor Rev.*
719   (2006). doi:10.1016/j.cytogfr.2006.09.008

720   26.   Metcalf, D. *et al.* Gigantism in mice lacking suppressor of cytokine signalling-2.
721   *Nature* **405**, 1069–1073 (2000).

722   27.   Greenhalgh, C. J. *et al.* Growth Enhancement in Suppressor of Cytokine Signaling 2
723   (SOCS-2)-Deficient Mice Is Dependent on Signal Transducer and Activator of
724   Transcription 5b (STAT5b). *Mol. Endocrinol.* **16**, 1394–1406 (2002).

725   28.   Greenhalgh, C. J. *et al.* SOCS2 negatively regulates growth hormone action in vitro
726   and in vivo. *J. Clin. Invest.* **115**, 397–406 (2005).

727   29.   Hansen, J. a, Lindberg, K., Hilton, D. J., Nielsen, J. H. & Billestrup, N. Mechanism of
728   inhibition of growth hormone receptor signaling by suppressor of cytokine signaling
729   proteins. *Mol. Endocrinol.* **13**, 1832–1843 (1999).

730   30.   Stofega, M. R., Herrington, J., Billestrup, N. & Carter-Su, C. Mutation of the SHP-2
731   binding site in growth hormone (GH) receptor prolongs GH-promoted tyrosyl

phosphorylation of GH receptor, JAK2, and STAT5B. *Mol. Endocrinol.* **14**, 1338–
1350 (2000).

31. Bulatov, E. *et al.* Biophysical Studies on Interactions and Assembly of Full-size E3
Ubiquitin Ligase. *J. Biol. Chem.* **290**, 4178–4191 (2015).

32. Bullock, A. N., Debreczeni, J. E., Edwards, A. M., Sundstrom, M. & Knapp, S.
Crystal structure of the SOCS2-elongin C-elongin B complex defines a prototypical
SOCS box ubiquitin ligase. *Proc. Natl. Acad. Sci.* **103**, 7637–7642 (2006).

33. Yang, X. O. *et al.* The signaling suppressor CIS controls proallergic T cell
development and allergic airway inflammation. *Nat. Immunol.* **14**, 732–740 (2013).

34. Eyckerman, S. *et al.* Design and application of a cytokine-receptor-based interaction
trap. *Nat. Cell Biol.* **3**, 1114–1119 (2001).

35. Lavens, D. A complex interaction pattern of CIS and SOCS2 with the leptin receptor.
J. Cell Sci. **119**, 2214–2224 (2006).

36. Goldshmit, Y., Walters, C. E., Scott, H. J., Greenhalgh, C. J. & Turnley, A. M.
SOCS2 induces neurite outgrowth by regulation of epidermal growth factor receptor
activation. *J. Biol. Chem.* (2004). doi:10.1074/jbc.M312873200

37. Dey, B. R., Spence, S. L., Nissley, P. & Furlanetto, R. W. Interaction of Human
Suppressor of Cytokine Signaling (SOCS)-2 with the Insulin-like Growth Factor-I
Receptor. *J. Biol. Chem.* **273**, 24095–24101 (1998).

38. Goldschmidt, L., Cooper, D. R., Derewenda, Z. S. & Eisenberg, D. Toward rational
protein crystallization: A Web server for the design of crystallizable protein variants.
Protein Sci. **16**, 1569–1576 (2007).

39. Gadd, M. S., Bulatov, E. & Ciulli, A. Serendipitous SAD solution for DMSO-Soaked
SOCS2-ElonginC-ElonginB crystals using covalently incorporated dimethylarsenic:
Insights into substrate receptor conformational flexibility in cullin RING ligases. *PLoS
One* **10**, (2015).

40. Kim, Y. K. *et al.* Structural basis of intersubunit recognition in elongin BC-cullin 5-
SOCS box ubiquitin-protein ligase complexes. *Acta Crystallogr. Sect. D Biol.
Crystallogr.* **69**, 1587–1597 (2013).

41. Liu, J. & Nussinov, R. The mechanism of ubiquitination in the cullin-RING E3 ligase
machinery: Conformational control of substrate orientation. *PLoS Comput. Biol.* **5**,
(2009).

42. Kaneko, T. *et al.* Loops govern SH2 domain specificity by controlling access to
binding pockets. *Sci. Signal.* **3**, (2010).

43. Carr, H. & Purcell, E. Effects of Diffusion on Free Precession in Nuclear Magnetic
Resonance Experiments. *Phys. Rev.* **94**, 630–638 (1954).

44. Meiboom, S., Gill, D. & Gillt, D. Modified Spin-Echo Method for Measuring Nuclear
Relaxation Times Modified Spin-Echo Method for Measuring Nuclear Relaxation
Times*. *Cit. Rev. Sci. Instruments J. Chem. Phys. J. Appl. Phys. J. Chem. Phys. J.
Appl. Phys. Spin-Echo Method Meas. Relax. Times Two-Line NMR Spectra Rev.*

Sci. Instrum **29**, 214504–2174 (1958).

45.   Dalvit, C., Fagerness, P. E., Hadden, D. T. A., Sarver, R. W. & Stockman, B. J.
774   Fluorine-NMR experiments for high-throughput screening: Theoretical aspects,
775   practical considerations, and range of applicability. *J. Am. Chem. Soc.* **125**, 7696–
776   7703 (2003).

777   46.   de Castro, G. V. & Ciulli, A. Spy vs. spy: selecting the best reporter for 19F NMR
778   competition experiments. *Chem. Commun.* **55**, 1482–1485 (2019).

779   47.   Wang, Y. Sen, Liu, D. & Wyss, D. F. Competition STD NMR for the detection of
780   high-affinity ligands and NMR-based screening. *Magn. Reson. Chem.* **42**, 485–489
781   (2004).

782   48.   Forbes, S. A. *et al.* COSMIC: Somatic cancer genetics at high-resolution. *Nucleic*
783   *Acids Res.* **45**, D777–D783 (2017).

784   49.   Rupp, R. *et al.* A Point Mutation in Suppressor of Cytokine Signalling 2 (Socs2)
785   Increases the Susceptibility to Inflammation of the Mammary Gland while Associated
786   with Higher Body Weight and Size and Higher Milk Production in a Sheep Model.
787   *PLoS Genet.* (2015). doi:10.1371/journal.pgen.1005629

788   50.   Zhang, Y. *et al.* Simultaneous binding of two peptidyl ligands by a Src homology 2
789   domain. *Biochemistry* **50**, 7637–7646 (2011).

790   51.   Mészáros, B., Kumar, M., Gibson, T. J., Uyar, B. & Dosztányi, Z. Degrons in cancer.
791   *Science Signaling* **10**, (2017).

792   52.   Liu, C. *et al.* Control of β-catenin phosphorylation/degradation by a dual-kinase
793   mechanism. *Cell* (2002). doi:10.1016/S0092-8674(02)00685-2

794   53.   Alexander, W. S. & Hilton, D. J. The Role of Suppressors of Cytokine Signaling
795   (SOCS) Proteins in Regulation of the Immune Response. *Annu. Rev. Immunol.* **22**,
796   503–529 (2004).

797   54.   Yoshimura, A., Naka, T. & Kubo, M. SOCS proteins, cytokine signalling and immune
798   regulation. *Nature Reviews Immunology* **7**, 454–465 (2007).

799   55.   Linossi, E. M., Babon, J. J., Hilton, D. J. & Nicholson, S. E. Suppression of cytokine
800   signaling: The SOCS perspective. *Cytokine and Growth Factor Reviews* **24**, 241–
801   248 (2013).

802   56.   Galic, S., Sachithanandan, N., Kay, T. W. & Steinberg, G. R. Suppressor of cytokine
803   signalling (SOCS) proteins as guardians of inflammatory responses critical for
804   regulating insulin sensitivity. *Biochem. J.* **461**, 177–188 (2014).

805   57.   Inagaki-Ohara, K., Kondo, T., Ito, M. & Yoshimura, A. SOCS, inflammation, and
806   cancer. *JAK-STAT* **2**, e24053 (2013).

807   58.   Wikman, H. *et al.* Identification of differentially expressed genes in pulmonary
808   adenocarcinoma by using cDNA array. *Oncogene* **21**, 5804–5813 (2002).

809   59.   Sutherland, K. D. *et al.* Differential hypermethylation of SOCS genes in ovarian and
810   breast carcinomas. *Oncogene* **23**, 7726–7733 (2004).

811   60.   Farabegoli, F., Ceccarelli, C., Santini, D. & Taffurelli, M. Suppressor of cytokine
812   signalling 2 (SOCS-2) expression in breast carcinoma. *J. Clin. Pathol.* **58**, 1046–
813   1050 (2005).

814   61.   Qiu, X. *et al.* Reduced expression of SOCS2 and SOCS6 in hepatocellular
815   carcinoma correlates with aggressive tumor progression and poor prognosis. *Mol.*
816   *Cell. Biochem.* **378**, 99–106 (2013).

817   62.   Zhu, J. G. *et al.* Expression of SOCSs in human prostate cancer and their
818   association in prognosis. *Mol. Cell. Biochem.* **381**, 51–59 (2013).

819   63.   Haffner, M. C. *et al.* Favorable prognostic value of SOCS2 and IGF-I in breast
820   cancer. *BMC Cancer* **7**, (2007).

821   64.   Paul, I. *et al.* The ubiquitin ligase Cullin5 SOCS2 regulates NDR1/STK38 stability
822   and NF-κB transactivation. *Sci. Rep.* **7**, (2017).

823   65.   Oeckinghaus, A. & Ghosh, S. The NF-kappaB family of transcription factors and its
824   regulation. *Cold Spring Harbor perspectives in biology* **1**, (2009).

825   66.   Hoesel, B. & Schmid, J. A. The complexity of NF-κB signaling in inflammation and
826   cancer. *Molecular Cancer* **12**, (2013).

827   67.   Hughes, S. J. & Ciulli, A. Molecular recognition of ternary complexes: a new
828   dimension in the structure-guided design of chemical degraders. *Essays Biochem.*
829   **61**, 505–516 (2017).

830   68.   Lucas, X. & Ciulli, A. Recognition of substrate degrons by E3 ubiquitin ligases and
831   modulation by small-molecule mimicry strategies. *Current Opinion in Structural*
832   *Biology* **44**, 101–110 (2017).

833   69.   Ottis, P. & Crews, C. M. Proteolysis-Targeting Chimeras: Induced Protein
834   Degradation as a Therapeutic Strategy. *ACS Chem. Biol.* **12**, 892–898 (2017).

835   70.   Zengerle, M., Chan, K. H. & Ciulli, A. Selective Small Molecule Induced Degradation
836   of the BET Bromodomain Protein BRD4. *ACS Chem. Biol.* **10**, 1770–1777 (2015).

837   71.   Maniaci, C. *et al.* Homo-PROTACs: bivalent small-molecule dimerizers of the VHL
838   E3 ubiquitin ligase to induce self-degradation. *Nat. Commun.* **8**, 830 (2017).

839   72.   Winter, G. E. *et al.* Phthalimide conjugation as a strategy for in vivo target protein
840   degradation. *Science (80-.).* **348**, 1376 LP-1381 (2015).

841   73.   Lu, J. *et al.* Hijacking the E3 Ubiquitin Ligase Cereblon to Efficiently Target BRD4.
842   *Chem. Biol.* **22**, 755–763 (2015).

843   74.   Kabsch, W. XDS. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 125–132 (2010).

844   75.   Evans, P. R. & Murshudov, G. N. How good are my data and what is the resolution?
845   *Acta Crystallogr. Sect. D Biol. Crystallogr.* **69**, 1204–1214 (2013).

846   76.   Winn, M. D. *et al.* Overview of the CCP4 suite and current developments. *Acta*
847   *Crystallographica Section D: Biological Crystallography* **67**, 235–242 (2011).

77.   McCoy, A. J., Storoni, L. C. & Read, R. J. Simple algorithm for a maximum-likelihood

SAD function. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **60**, 1220–1228 (2004).

78.   McCoy, A. J. *et al.* Phaser crystallographic software. *J. Appl. Crystallogr.* **40**, 658–
674 (2007).

79.   Adams, P. D. *et al.* PHENIX: A comprehensive Python-based system for
macromolecular structure solution. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**,
213–221 (2010).

80.   Terwilliger, T. C. *et al.* Iterative model building, structure refinement and density
modification with the PHENIX AutoBuild wizard. in *Acta Crystallographica Section D:*
Biological Crystallography **64**, 61–69 (2007).

81.   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of
Coot. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 486–501 (2010).

82.   Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal
structures. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **67**, 355–367 (2011).

83.   Chao, H. G. *et al.* Synthesis and Application of Fmoc-O-
[bis(dimethylamino)phosphono]-tyrosine, a Versatile Protected Phosphotyrosine
Equivalent. *J. Org. Chem.* (1995). doi:10.1021/jo00129a001

84.   Biosystems, A. Cleavage, Deprotection, and Isolation of Peptides after Fmoc
Synthesis. *Tech. Bull.* 1–12 (1998). doi:10.1038/nrd.2016.29

85.   Perich, J. W. & Reynolds, E. C. The Facile One-Pot Synthesis of Nα-(9-
Fluorenylmethoxycarbonyl)-O-(O′, O″-dialkylphosphoro)-L-tyrosines Using Dialkyl
N,N-Diethylphosphoramidites. *Synlett* **1991**, 577–578 (1991).

86.   Armougom, F. *et al.* Expresso: Automatic incorporation of structural information in
multiple sequence alignments using 3D-Coffee. *Nucleic Acids Res.* (2006).
doi:10.1093/nar/gkl092

87.   Hruz T., Laule O., Szabo G., Wessendorp F., Bleuler S., Oertle L., Widmayer P.,
Gruissem W. and Zimmermann P. Genevestigator V3: a reference expression
database for the meta-analysis of transcriptomes. *Advances in Bioinformatics*
(2008). 420747

Figures and Schemes

2
Figure 1. Structural and interaction detail of the SBC-EpoR co-crystal
(a) Domain and protein arrangement of the SBC-EpoR co-crystal. Protein chains are
shown in cartoon, with EloB (cyan), EloC (magenta) and SOCS2, comprising of SOCS box
(yellow) and SH2 domain (green). The EpoR_pY426 peptide is shown in orange stick. (b)
The Fo-Fc ligand omit map of the EpoR_pY426 peptide (green mesh) contoured at 2.0 σ
level to highlight densities for the EpoR_pY426 peptide (orange stick) (c) Hydrogen bond
interactions (dash) between the pY of EpoR_pY426 peptide (orange stick) and SOCS2
(green). (d) Hydrogen bond interaction (dash) between the EpoR_pY426 peptide (orange
stick), SOCS2 (green stick) and water (red sphere). (e) Hydrophobic interaction between
EpoR_pY426 peptide (orange stick) and SOCS2 (surface). Hydrophobic residues on
SOCS2 are colored in pink.

Figure 2. Structural and interaction detail of the SBC-GHR co-crystal

Two copies of GHR_pY595 were observed in the co-crystal. One copy is shown as peptide
A (yellow stick) and the other one as peptide B (cyan stick). (a) The Fo-Fc ligand omit map
of the peptides (green mesh) contoured at 2.5 σ level to highlight densities for the peptide
A and peptide B (yellow and cyan stick). (b) Cartoon diagram of the SH2 domain of
SOCS2 (green) with peptides A and peptide B. (c) Hydrogen bond interactions (dash)
among peptide A, peptide B, SOCS2 (green stick) and water (red sphere). (d) (d)(e)
Hydrophobic interaction of the peptides with C-terminal half and N-terminal half of the SH2
domain (surface), respectively. SOCS2 residues involved in hydrophobic interactions are
coloured in pink. (f) The coordination of cobalt ion (pink sphere) with His149 of SOCS2
(green), His(+4) of peptide B and water molecules (red sphere).

(a) Secondary structure elements in SOCS2 are shown above the sequence alignment.
SOCS proteins with structure available were aligned using T-Coffee expresso mode for
sequence alignment with structural information [86]. (b) Locations of the EF loop (orange)
and the BG loop (green) on SOCS2 (white). (c) The Fo-Fc ligand omit map of the
previously disordered BG loop (green mesh) contoured at 1.5 σ level.

27 **Figure 3. The BG loop of SOCS2**

Figure 4. Conformation changes of the EF and BG loops

(a) The EF loop curls up in the apo SOCS2 (white, PDB code: 2C9W), placing Ile110 and Cys111 (stick) at the SH2 hydrophobic pocket. (b) The EF loop makes backbone interactions with GHR peptide (pink) in the SBC-GHR$_2$ structure. (c) The Ile109 and Val 112 of SOCS2 (cyan) and Val112 of SOCS2 symmetry mate (white) make unique hydrophobic interaction with Pro(+5) of EpoR peptide (blue). (d) Superposition of the SOCS2 from SBC-GHR$_2$ (green) and SBC-EpoR (cyan) displays conformational changes of the BG loop.

Scheme 1. Synthesis of spy molecule 3

(a)

(b)

Protein	SPR K_D (µM)		
	GHR_pY595	GHR_pY487	EpoR_pY426
Wild type	1.4 ± 0.12	5.1 ± 0.77	12.5 ± 0.63
L106V	2.4 ± 0.11	8.9 ± 0.70	14.1 ± 1.23
C133Y	2.2 ± 0.10	7.7 ± 0.71	13.4 ± 0.82
N94D	n.d.	weak	n.d.
R96L	weak	weak	n.d.
R96Q	n.d.	n.d.	n.d.

(c)

48
Figure 5. Characterization of SNP containing SOCS2 by SPR and ^{19}F-NMR

(a) Sequence conservation mapped onto the SH2 domain of SOCS2 surface.
Conservation surface representation based upon the ClustalW multiple sequence
alignment of CIS and SOCS1 – SOCS7 sequences where highly conserved residues are
shown in red/orange colour and variable residue positions coloured blue. (b) The SBC
proteins binding affinity for phosphorylated substrate peptides were determined by SPR.
Weak denotes signal detected, however, saturation was not achieved. Hence, binding too
weak to be detected. N.d. denotes signal not detected. Values reported are the means ±
s.e.m. from four independent experiments (c) Transverse relaxation rate (R_2) of SBC
proteins. Error bars are R_2 errors generated from five-point data fitting.

59
60

62
63
Figure 6. Comparison of the GHR and EpoR binding
(a) Overlay of the GHR_pY595 (yellow) and EpoR (orange) peptides. Side chains circled
in blue have similar properties and red have different properties. (b) Different hydrophobic
interactions (pink) caught by GHR_pY595 (yellow) and EpoR (orange) peptide. (c) Overlay
of the substrate peptide of SOCS proteins. The GHR_pY595 (yellow) and EpoR (orange)
peptides catch distinct hydrophobic cavities (pink) on SOCS2. These interactions
differentiate SOCS2 substrate binding mode from other substrates of SOCS proteins. The
SOCS3 substrate peptide gp130 is in cyan (PDB ID: 2HMH) and SOCS6 substrate peptide
c-Kit is in green (PDB ID: 2VIF). (d) The binding of the non-canonical GHR (cyan) relies on
backbone to backbone interaction and hydrophobic interaction from pY(+3) and pY(+5) of
the GHR_pY595 to the SH2 domain hydrophobic core (pink). The binding mode of EpoR
(orange) makes it unfavourable to establish these interactions.

(a)

(b)

(c)

(d)

(e)

Figure 7. The triple-stranded β-sheet interaction between substrate peptide and protein

(a) The SOCS3 (blue) in complex with gp130 (cyan) (PDB code: 2HMH) (B) The SOCS6 (magenta) in complex with c-Kit (pink) (PBD code: 2VIF) (C) The SOCS2 (green) in complex with GHR_pY595 peptides. Peptide A is shown in yellow and peptide B in cyan. (d) The SHP2 (purple) in complex with peptides (white). (e) overlap of SH2 domains from SOCS2-GHR (green)   and SHP2-pY peptide (cyan) highlights the difference in positioning of non-canonical pY from GHR and SHP2-substrate peptide

87
88
Figure 8. Illustration of GHR peptide binding modes

(a) Illustration of the "cis" recognition binding mode where the two binding sites on GHR,
pY487 and pY595, folded into a hairpin structure for SOCS2 binding. SOCS2 is shown in
white surface, the hairpin structure is illustrated by connecting the two peptides (yellow
cartoon) with red dash lines. (b) Illustration of the "trans" binding mode where SOCS2
(white surface) could grab two receptor tails (cyan and yellow dash line) at the same time
after dimerization of receptors at the membrane.

Tables

Table 1. Data collection and refinement statistics

	SBC-EpoR	SBC-GHR	SBC-GHR$_2$
PDB code	6I4X	6I5N	6I5J
Data collection			
Wavelength (Å)	0.9686	0.9795	0.9686
Space group	I 1 2 1	P 2$_1$ 2$_1$ 2	P 2 2$_1$ 2$_1$
Cell dimensions			
a, b, c (Å)	41.29, 56.33, 203.39	113.18, 156.76, 57.57	57.83, 113.71, 156.94
α, β, γ (°)	90.00, 91.53, 90.00	90.00, 90.00, 90.00	90.00, 90.00, 90.00
Molecules/ASU	1	2	2
Resolution	29.36–2.69 (2.82–2.69)	113.55–1.98 (2.01–1.98)	92.08–2.80 (2.95–2.80)
R$_{merge}$ (%)	10.8 (51.4)	9.4 (103.5)	19 (72.1)
<I/σ (I)>	9.6 (2.4)	17.1 (2.2)	7.3 (2.7)
Completeness (%)	93.2 (63.3)	100 (100)	100 (100)
Redundancy	4.9 (4.2)	13.3 (13.2)	7.8 (7.8)
CC$_{1/2}$	0.99 (0.82)	1.0 (0.9)	0.98 (0.74)
Refinement			
Resolution (Å)	2.69	1.98	2.8
Unique reflections	12265 (791)	72170(7105)	26275(2606)
R$_{work}$/R$_{free}$ (%)	19.64/23.51	19.00/22.66	20.96/26.38
Wilson B factor (Å^2)	44.1	24.2	42.5
Average B factor (Å^2)	46.9	32.0	42.5
No. non-hydrogen atoms Protein/ligand/water	2688/83/11	6433/344/512	5775/327/33
R.M.S.D.			
Bond lengths (Å)	0.002	0.010	0.002
Bond angles (°)	0.451	1.064	0.412
Ramachandran analysis			
Preferred regions (%)	96.12	97.82	96.83
Allowed regions (%)	3.58	2.18	3.17
Outliers (%)	0.3	0.00	0.00

98 Values in parentheses are for the highest resolution shell

99 **Table 2. Binding affinities of the GHR peptide and Val(-3) analogues**

GHR peptide	Sequence	K_D by SPR (μM)	K_i by NMR (μM)
Wild type	PVPDpYTSIHIV	1.5 ± 0.09	1.5 ± 0.06
V(-3)R	P**R**PDpYTSIHIV	9.0 ± 1.05	20.6 ± 2.09
V(-3)Y	P**Y**PDpYTSIHIV	3.3 ± 0.13	3.3 ± 0.24

101 Residue mutated in bold
102 Values reported are the means ± s.e.m. from four independent experiments

Table 3. Affinity measurement for the GHR_pY595 and EpoR_pY426 derivatives by SPR and ^{19}F NMR

Sequence	Ala. position	SPR K_D ± SE (μM)	NMR K_i (μM)
PVPDpYTSIHIV	GHR wild type	1.5 ± 0.09	1.5 ± 0.06
AVPDpYTSIHIV	pY(-4)	1.7 ± 0.14	0.9 ± 0.17
P**A**PDpYTSIHIV	pY(-3)	2.7 ± 0.19	3.4 ± 0.25
PV**A**DpYTSIHIV	pY(-2)	1.4 ± 0.12	4.1 ± 0.75
PVP**A**pYTSIHIV	pY(-1)	6.2 ± 0.37	21.7 ± 3.34
PVPDpY**A**SIHIV	pY(+1)	1.0 ± 0.09	0.5 ± 0.07
PVPDpYT**A**IHIV	pY(+2)	1.4 ± 0.07	1.2 ± 0.06
PVPDpYTS**A**HIV	pY(+3)	3.7 ± 0.22	3.4 ± 0.66
PVPDpYTSI**A**V	pY(+4)	6.7 ± 0.78	18.5 ± 2.67
PVPDpYTSIH**A**V	pY(+5)	2.4 ± 0.13	1.4 ± 0.36
PVPDpYTSIHI**A**	pY(+6)	1.9 ± 0.09	1.6 ± 0.21
ASFEpYTILDPS	EpoR wild type	13.3 ± 0.59	7.1 ± 1.01
A**A**FEpYTILDPS	pY(-3)	14.2 ± 0.65	7.5 ± 1.04
AS**A**EpYTILDPS	pY(-2)	16.3 ± 1.02	7.2 ± 0.42
ASF**A**pYTILDPS	pY(-1)	21.7 ± 1.49	23.5 ± 2.99
ASFEpY**A**ILDPS	pY(+1)	17.1 ± 0.38	10.3 ± 1.28
ASFEpYT**A**LDPS	pY(+2)	30.8 ± 1.24	17.2 ± 0.89
ASFEpYTI**A**DPS	pY(+3)	32.3 ± 1.42	20.4 ± 0.88
ASFEpYTIL**A**PS	pY(+4)	18.0 ± 0.91	14.1 ± 1.74
ASFEpYTILD**A**S	pY(+5)	8.9 ± 1.25	6.9 ± 1.39
ASFEpYTILDP**A**	pY(+6)	16.4 ± 0.29	11.3 ± 1.29

Residue mutated in bold
Values reported are the means ± s.e.m. from four independent experiments