The landscape of DNA methylation associated with the transcriptomic network in laying hens and broilers generates insight into embryonic muscle development in chicken

Zihao Liu[£], Xiaoxu Shen[£], Shunshun Han[£], Yan Wang[£], Qing Zhu, Can Cui, Haorong He, Jing Zhao, Yuqi Chen, Yao Zhang, Lin Ye, Zhichao Zhang, Diyan Li, Xiaoling Zhao and Huadong Yin[#]

Farm Animal Genetic Resources Exploration and Innovation Key Laboratory of Sichuan Province, Sichuan Agricultural University, Chengdu, Sichuan 611130, PR China

[£] These authors contributed equally to this work.

[#] Corresponding author:

Huadong Yin, Farm Animal Genetic Resources Exploration and Innovation Key Laboratory of Sichuan Province, Sichuan Agricultural University, Chengdu, Sichuan 611130, PR China. E-mail: yinhuadong@sicau.edu.cn

Abstract

As DNA methylation is one of the key epigenetic mechanisms involved in embryonic development, elucidating its relationship with non-coding RNA and genes is essential for understanding early development. In this study, we performed single-base-resolution bisulfite sequencing together with RNA-seq to explore the genetic basis of embryonic muscle development in chicken. Comparison of methylome profiles between broilers and laying hens revealed that lower methylation in broilers might contribute to muscle development. Differential methylated region (DMR) analysis between two chicken lines showed that the majority of DMRs were hypo-DMRs for broilers. Differential methylated genes were significantly enriched in muscle development-related terms at E13 and E19. Furthermore, by constructing the network of the lncRNAs, we identified a lncRNA, which we named MYH1-AS, that potentially regulated muscle development. These findings reveal an integrative landscape of late period of embryonic myogenesis in chicken and give rise to a comprehensive understanding of epigenetic and transcriptional regulation, in skeletal muscle development. Our study provides a reliable data resource for further muscle studies.

Keywords: DNA methylation, lncRNA, chicken, muscle development

Introduction

Epigenetics mechanisms, including DNA methylation, histone modification, non-coding RNAs and chromatin remodeling, have been the subject of intense research over recent years because of their essential roles in various biological processes [1,2]. These epigenetic mechanisms have been reported to be involved in human diseases[3], oogenesis and spermatogenesis[4] as well as in adipose and muscle development[5-7]. DNA methylation is an epigenetic mechanism that exerts considerable influence on the regulation of gene expression without changing the DNA sequence[8]. A role for DNA methylation in muscle development has been illustrated in human[9], pig[5,6], rabbit[10], bovine[11] and chicken[12].

The embryonic stage is critical for muscle development in mammals, as the number of muscle fibers in the developing embryo remains stable after birth. Previous reports have demonstrated a function of DNA methylation in embryonic muscle development. For instance, Carrio et al.[13] built the methylome of myogenic stem cells and demonstrated the importance of DNA methylation-mediated regulation of the cell-identity Myf5 super-enhancer during muscle-stem cell differentiation. Long noncoding RNAs have also been proven to be important in the regulation of muscle development. For example, linc-MD1 interacts with miR-133 and miR-135 to regulate the expression of transcription factors MAML1 and MEF2C that activate muscle-specific gene expression[7]. Recently, the

regulatory relationship between DNA methylation and lncRNAs has

drawn extensive research attentions and a database of methylation and

lncRNA regulatory relationships has been built for human diseases

studies[14]. However, studies on the role for this regulatory relationship in

muscle development are limited. Zhang at el.[5] reported the function of the

lincRNA and DNA methylation regulatory relationship in muscle

development in pig. Yang at el.[6] revealed that DNA methylation

potentially affects gene expression in skeletal muscle to influence the

propensity for obesity and body size.

After long-term artificial breeding for different purposes, laying

hens and broilers show great differences in the development of skeletal

muscles. The skeletal muscle growth rate of broilers far exceeds that of

laying hens even under optimal feeding conditions, and broilers can

exhibit weights 5 times more than laying hens at 6 weeks of age. The

comparatively similar genetic backgrounds and genomes of these two

chicken lines allow for comparative studies of muscle development at the

epigenetic level.

Several genome-wide methylation studies have been reported in

chicken, and a relationship between DNA methylation level of promoters

and expression level of genes were identified[15-17]. Furthermore, the global

methylation landscape of muscle development was described in chicken

using juvenile and later laying-period hens[12]. However, a role for DNA

methylation in chicken embryonic muscle development has not been fully

clarified.

Here we used whole genome bisulfite sequencing to determine the

methylomes of 12 standardized broilers and 12 standardized laying hens.

We sequenced the whole transcriptome of these 24 samples by RNA-seq

simultaneously for the multi-Omics integrative analyses, to explore the

effect of DNA methylation and lncRNA relationship on muscle

development.

Results

Overview of DNA methylation

In the genomic methylation data among 24 samples (from 12

broilers and 12 laying hens), the average sequence depth is about 30.3X.

Approximately 3.4 billion reads were generated by the Illumina HiSeq in

total and an average of 71.99% clean reads were mapped to the *Gallus*

gallus genome (version 5.0) (Supplementary Table S1). The coverage

analysis revealed that approximately 82% of the *Gallus gallus* genome

were covered by reads at least one-fold, whereas nearly 78% of genome

was covered by more than five-fold and 75% of genome was covered

more than 10-fold (Supplementary Table S2). These results indicated a

reliable sequencing outcome.

The methylation level of each developmental stages is displayed in

Fig 1a, which indicates that the layers and broilers have a similar global

methylation profile. Similar proportions of CpGs in three sequence

contexts (mCG, mCHG, and mCHH) were observed among four

developmental stages (Fig. 1b). Next, the methylation level distributions

of CpGs were analyzed at four developmental stages. In general, CpGs

showed a high methylation level in the mCG context and a low

methylation level in mCHG and mCHH contexts (Fig. 1c and

Supplementary Fig. 1a). We then measured the methylation level of

different regions of genes and compared these levels at different stages

and populations. Interestingly, we found that broilers showed statistically

lower methylation levels at all stages in the mCG context than layers (Fig.

1d). We quantified the numbers of CpG islands (CGIs) in different

regions at different stages (Supplementary Fig. 1b). More CGIs were

located in gene promoter regions in broilers than layers, which indicates

that methylation in CGIs may be involved in faster muscle development

in broilers, as CGIs located at promoter regions are important for

controlling gene expression.[18].

We also examined the methylation level of lncRNAs assembled in

RNA-seq using a similar approach and compared levels with the analysis

of gene methylations. Generally, broilers still showed a lower methylation

level in various types of lncRNAs in mCG and mCHH contexts compared

with layers; similar methylation levels were observed among different

types of lncRNAs (Fig. 2b and Supplementary Fig. 2c–d). Genes and

lncRNAs had similar global methylation levels and both showed significant difference in broilers compared with layers (Fig. 2a and Supplementary Fig. 2a–b). These results suggest that faster muscle development of broilers may be due to the lower methylation level in late embryonic stage compared with those in layers.

We also analyzed the genomic distribution patterns of DNA methylation in genes and lncRNAs. We divided the upstream region (2 kb), first exon, first intron, internal exon, internal intron, last exon and downstream region (2 kb) of genes and lncRNAs across the genome as different features and their methylation levels were measured through 20 bins. In general, the 5′ upstream and 3′ downstream regions showed lower methylation levels than other gene regions. We also compared the methylation level of features of genes with features of lncRNA (Fig. 2c–d). LncRNAs have relatively higher methylation levels around the transcription start site (TSS) compared with genes ($P < 0.001$). In addition, methylation levels of different types of repeat regions were also analyzed across the genome. Beside the significant differences between broilers and layers, short interspersed nuclear elements (SINE) showed lower methylation levels across the four stages in the mCG context (Fig. 3 and Supplementary Fig. 3).

Identification of differential methylation regions (DMRs) and genes.

To explore the potential causes of the divergence in muscle

development between broilers and layers, the differential methylation loci were identified in DSS package. DMRs were identified in E10, E13, E16 and E19 based on differential methylation loci. The DMRs were subsequently annotated to the genome, and the distribution of the DMRs in the whole genome was analyzed (Fig. 4a and Supplementary Table S4–S7). In general, the majority of DMRs was located in intronic regions, and a small portion of DMRs was distributed in the promoters of genes (Fig. 4a). Proportion analysis revealed that broilers had more hypomethylated regions across the genome in the four developmental stages, indicating that low methylation in muscle development-related genes may account for the fast muscle development in broilers (Fig. 4b).

Differential methylation genes (DMGs) were defined as genes with at least one overlapping DMR in its exon/intron regions. Gene Ontology (GO) enrichment analyses were then performed to investigate potential biological functions of the DMGs. In general the DMGs in the four developmental stages were most significantly enriched in terms related to the nervous system. However, many muscle-related terms were also found, especially for DMGs at E13 and E19, such as muscle organ development (47 genes; Q-value < 0.001), myotube cell development (12 genes; Q-value < 0.005), positive regulation of muscle organ development (17 genes; Q-value < 0.001), and muscle cell differentiation (51 genes; Q-value < 0.003) (Fig. 4c, Supplementary Table S8–S11).

Because DMRs were not unanimous on the genomic position among different developmental stages, we merged the genomic position of DMRs from the 24 samples to generate common DMRs and re-calculated the methylation level for each common DMR. Clustering analysis was performed using the common DMRs and displayed using heatmap analysis. Different developmental stages were shown to cluster together, which is indicative of the high quality of sampling and DMR calling in this experiment (Fig. 5a). Moreover, the principle component analysis (PCA) result was consistent with the clustering analysis (Fig. 5b).

Integrative analyses of DNA methylation and transcriptome

To further explore whether methylation influences gene and lncRNA expression in chicken, RNA-seq was used to measure the expression of genes and identified lncRNAs. We identified 20656 lncRNAs in total. Most of the lncRNAs were lincRNAs (63.6%) (Fig. 6a, 6b). Heatmap of 24 samples and PCA suggested developmental stages accounted for most variances (Fig. 6c). We divided genes and lncRNAs into four groups on the basis of their expression level (highest, medium high, medium low and lowest) using quantile method. We then measured methylation levels in different groups of genes and lncRNAs. In general, broilers and layering hens had similar methylation levels. A negative correlation was observed between genes and methylation of promoters in both broilers and layers: the highest expression level group showed the lowest

methylation level around the TSS, whereas the lowest expression level group showed the highest methylation level (Fig. 6d, e). Interestingly, the trend of negative correlation between expression and methylation was observed in downstream regions of lncRNAs but not around the TSS (Fig. 6f, g). Moreover, the lncRNAs were usually methylated at higher levels around the TSS compared with genes (Fig. 6d–g).

Next, differential expression gene (DEG) and lncRNA (DEL) calling was performed, and the cis-targets and trans-targets of lncRNAs were predicted. The DMRs were assigned to lncRNAs generated from RNA-seq in this study (Supplementary Table S12–S15) and the differential methylation lncRNA (DM lncRNA) were defined as DEL that overlapped with DMR. The result showed that 55 DM lncRNAs were identified (13,16,11,15 in 4 stages, respectively) (Supplementary Table S16). We then searched for DM lncRNAs with potential in regulating muscle development. In particular, we found that the expression of one lncRNA (which we named as MYH1-AS; Fig. 7a) was highly correlated with the methylation level of the DMR assigned to it (Spearman, Cor=-0.7513, $P < 10^{-4}$; Fig. 7b). The expression of MYH1-AS was detected to dramatically increase in broilers compared to laying hens at E16 and E19 (Fig 8a). As the lncRNA was predicted by by lncTar to target several genes like MYH1A, MYH1G and MYH1E, the expression correlations between the lncRNA and its targets were calculated to search

for its most likely target. MYH1E showed the highest correlation with

MYH1-AS (Fig. 7d), indicating MYH1E as a potential target of

MYH1-AS. To further explore the role of MYH1-AS in muscle

development, the gene-lncRNA networks were constructed based on their

mRNA expression connectivity using WGCNA, and the subnetwork of

MYH1-AS was extracted from the whole network. MYH1-AS had a high

correlation with several muscle-related genes in this subnetwork (Fig. 7d).

The relationship between the connectivity and correlation is shown in

Figure 7f. Interestingly, genes that were highly negatively correlated with

MYH1-AS did not show high connectivity with MYH1-AS. All genes

showing high connectivity with MYH1-AS were also highly positively

correlated with the lncRNA (Fig 7e-f). A total of 168 genes with both

high connectivity and correlation with MYH1-AS, were selected to

perform GO enrichment analysis to confirm the role of MYH1-AS in

muscle (Fig. 7g and Supplementary S17). The results showed that the

majority of terms enriched by these genes were muscle-related.

The expressions of MYH1-AS produced by RNA-seq were verified

by qPCR and a similar trend was observed, indicating a reliable

sequencing outcome (Fig 8 a, b). Subsequently, a siRNA was designed to

perform MYH1-AS silencing assay. As shown in fig 8c, expression of

MYH1-AS was significantly reduced after transfecting, indicative of

efficiency of siRNA used in this experiment (Fig 8c). Then the mRNA

expression of muscle related genes (MyoD1, MyoG and MyH3) were measured at 48h after MYH1-AS silencing. It resulted in a reduced mRNA expression in silencing groups compared to control groups (Fig 8d-f). Besides, the microscope was used to monitor the morphological change in myotubes after silencing. We found that MYH1-AS silencing resulted in a reduced number of myotube (Fig 8g-h). Further western blot assay revealed that the protein expression of MyhC and MyoG was repressed in silencing groups (Fig 8i). Those results suggest that lncRNA MYH1-AS may function in muscle differentiation.

Discussion

The chicken provides a unique model to perform embryology research because of the accessibility of egg. As chicken is an important food source for the human diet, the muscle development of chicken is an important topic worth of study. Here we used broilers and laying hens to explore the muscle development in chicken in the late embryonic period as they are artificially selected for different commercial use (depositing meat and laying eggs, respectively) thereby are divergent in muscle development. Because of the crucial role of methylation in embryogenesis, we performed whole genome bisulfite sequencing and RNA-seq to systematically explore the prenatal methylation landscape during chicken muscle development. Previous methylome studies have been performed using prenatal chicken or born chicken muscle[12,19,20],

however, these studies failed to generate a comprehensive methylation landscape of embryonic stages. We focused on more systematical study at embryonic stage range from E10 to E19 between two chicken lines and aimed to elucidate the detain of embryonic muscle development.

The methylation level and proportion of different methylations (mCG, mCHG, mCHH) of each developmental stage indicated that layers and broilers have a similar global methylation profile. We also measured the methylation level of different types of CpG (Fig. 1e–g), and results were consistent with previous studies in chicken muscle[15]. The distribution proportions of CpG in the genome were different from those in the study of Zhang et al[20], as the CpG proportions in repeat regions accounted for less genomic proportion in our study. One possibility for the discrepancy may be because the previous study used data from born chicken, whereas our analyses were performed in data from prenatal chicken. More studies are required to clarify these differences.

We next comprehensively compared the methylation level of genes and lncRNAs among different developmental stages and chicken lines (Fig. 2a). In general, layers showed a significantly higher methylation level than broilers in the mCG context in both genes and lncRNAs, which may be responsible for the differences in muscle development. Furthermore, we compared the methylation levels of different types of lncRNAs (sense, intronic, antisense and lincRNA) and there were no

significant differences, although layers and broilers still revealed
significant variances. Next, genomic methylation around genes and
lncRNAs were measured across the genome, and the TSSs were found to
be low methylated in genes (Fig. 2c). The broilers and layers showed
similar trends around the TSS, which is consistent with patterns reported
in previous studies in chicken[12,15], as well as in bovine muscle tissue[11] and
pig[21]. However, the TSSs of lncRNAs were usually methylated at higher
levels compared with genes, which may explain why mRNA expression
of lncRNAs are usually lower than genes ($P < 10^{-8}$) because methylation
events in the promoter region usually affect gene expression[22]. In addition,
the methylation levels of different types of transpose elements (TEs)
(SINE, LINE, LTR, DNA and satellites) were also measured and TEs
were methylated at higher levels in layers compared with broilers. TEs
are usually inactivated in animals but were reported to function in the
early development of human and other mammals to provide
cis-regulatory elements that coordinate the expression of groups of
genes[23]. As epigenetic regulation is important for the activity of TEs[24],
these differences in the two chicken lines may also account for the
divergence in development.

The clustering heatmap and PCA were performed using common
DMRs among four developmental stages. The expected classifications
were observed in both analyses, indicating the reliable outcomes of

sequencing and DMR calling. Moreover, we found that DMRs between two chicken lines mainly distributed in intron regions and intergenic regions. These results are consistent with previous studies in chicken[12], indicative of the important role of methylation in development regulation. However, as methylation in gene body region affects gene expression in several sophisticated ways[18], further studies on how methylation of the intron regions can influence gene expression are required to elucidate the complicated epigenetic mechanism underlying muscle development in chickens. We analyzed the proportion of hypermethylated and hypomethylated regions and the majority of DMRs were detected to be hypomethylated regions in broilers, indicating that low methylation may be responsible for fast muscle development. This result is consistent with the preceding results in this study. Genes with overlapped with DMR at different times were regarded as DMGs and used for GO enrichment analysis. We found that DMGs at E13 and E19 were significantly enriched in muscle-related terms, suggesting that methylation plays an important role in embryonic stage muscle development. Additionally, DMGs among four stages were significantly enriched in nerve development-related terms, which may relate to the impact of domestication and artificial breeding. Integrative analysis was conducted to study the association between methylation level and mRNA expression. We noticed that mRNA level and methylation level around TSSs were

negatively correlated in genes but not lncRNAs, indicating that DNA

methylation regulates lncRNA expression in a more complex way than

gene expression.

To explore which lncRNA may potentially influence muscle

development, the DM lncRNAs were identified and the correlation

between DM lncRNA and the assigned DMR were measured. In

particular, MYH1-AS showed a high correlation with its target MYH1E

and the DMR located in its intron region. Further WGCNA analysis

revealed that several muscle-related genes were highly correlated with

MYH1-AS in its subnetwork. For example, *MYLK2*, a muscle-specific

gene, expresses skMLCK specifically in skeletal muscles[25,26]. ABLIM1

was reported to be related to muscle weakness and atrophy[27]. Increased

PDK4 expression may be required for the stable modification of the

regulatory characteristics of PDK observed in slow-twitch muscle in

response to high-fat feeding[28], and other genes in the network, such as

MyoZ1, MYPN and ZBTB16 genes, were also revealed to be muscle- or

meat quality-related genes[29-32]. This indicates that MYH1-AS may

function in muscle development. Notably, as we noticed that high

correlation did not exactly indicate high connectivity (Fig. 7f), we also

performed GO enrichment analysis using 168 genes, which had top 50%

both high connectivity and correlation values with MYH1-AS in its

network as input. The majority of the resulting GO terms were

muscle-related terms (Fig. 7f–g), which is strongly indicative of MYH1-AS functioning in muscle development. Therefore, these results suggest that MYH1-AS is regulated by DNA methylation and participates in muscle development during embryonic stages. Subsequent silencing and western blot assay verified our analysis results, suggesting the reliability of our analysis and the role of MYH1-AS in muscle differentiation. However, how the lncRNA regulates muscle development requires more studies.

Our experiment revealed a comprehensive DNA methylome and transcriptome landscape during embryonic developmental stages. We identified one lncRNA, MYH1-AS, that may potentially play a part in muscle development in chicken, and our study provides evidence for this conclusion. Moreover, we provided a basis and a reliable resource for further investigating the genetic regulation of methylation and gene expression in embryonic chicken. However, more studies are needed to elucidate the detailed mechanism on how DNA methylation impacts lncRNA expression and how the lncRNA regulates myogenesis.

Acknowledgements

We thank Edanz Group (www.edanzediting.com/ac) for editing a draft of this manuscript.

Materials and Methods

Sample collection

The fertilized eggs of Rose and WhiteLoghorn were incubated in the same condition. The breast muscle and blood were collected at E10, E13, E16, E19. After sex determination, only samples identified as male were kept for next experiment. A total of 24 embryonic chicken were used in the study to form eight groups: E10, E13, E16, E19 for Rose and WhiteLoghorn, respectively. Each group included 3 individuals as biological replicates.

DNA and RNA extraction

Genomic DNA was extracted using an animal genomic DNA kit (Tiangen, China) following the manufacturer's instructions. The DNA integrity and concentration were measured by agarose gel electrophoresis and NanoDrop spectrophotometer, respectively. Total RNA was isolated using TRIzol (TAKARA, Dalian, China) 110 reagent according to the manufacturers' instruction. RNA was reverse 111 transcribed by TAKARA PrimeScriptTM RT reagent kit (TAKARA) 112 according to the manufacturers' instruction.

Library construction and sequencing

Bisulfite sequencing libraries were prepared using the TruSeq Nano DNA LT kit (Illumina, San Diego, CA, USA). The genomic DNAs were then fragmented into 100–300 bp by sonication (Covaris, USA) and purified using a MiniElute PCR Purification Kit (QIAGEN, Silicon Valley Redwood City, CA, USA). The fragmented DNAs were end

repaired and a single 'A' nucleotide was appended to the 3′ end of each

fragment. After ligating the DNAs to the sequencing adapters, the

genomic fragments were bisulfite converted via a Methylation-Gold kit

(ZYMO, Murphy Ave. Irvine, CA, USA). The converted DNA fragments

were PCR amplified and sequenced as paired-end reads using the

Illunima HiSeq xten platform by the Biomarker Technologies company

(Beijing, China).

Data alignment and process

The raw data in the FastQ format generated by the Illumina HiSeq

were pre-processed by removing reads containing adapters, N (unknown

bases) > 10%, and those which over 50% of the sequence exhibited low

quality value (Qphred score ≤ 10). During the process, we also calculated

the Q20, Q30, CG content for each sample data. The reads remained after

this procedure were clean reads and used for subsequent analysis. The

methylation data were aligned to reference genome Gallus gallus 5.0 by

Bismark software[33]. Meanwhile, the number of aligned clean reads in

unique position of reference genome were calculated as unique mapped

reads number. The proportion of the number of aligned reads in the total

number of reads was calculated as the mapping rate. Subsequently, the

methylation level of single base was then calculated by the ratio of the

number of methylated reads to the sum of total reads covered the locus.

Finally, we used a binominal distribution teat approach to determine

whether a locus was regarded as methylated locus with the criteria: coverage depth > 4 and FDR<0.05[33].

The transcriptional libraries were sequenced on an Illumina HiSeq xten platform at the Biomarker Technologies Company (Beijing, China). The obtained transcriptome data were filtered by removing sequences containing adaptors, low-quality reads (Q-value $<$ 20), and reads containing more than 10% of unknown nucleotides (N) and were aligned to reference genome Gallus gallus 5.0 by HISAT2[34] then the transcript assembly and FPKM calculation were performed using the StringTie[35]. Transcripts mapped to the coding genes of reference were used to subsequent differential expression gene calling.

LncRNA identification

In order to identify the potential lncRNA, the assembled transcripts generated from the StringTie were submitted to CPC[36], CNCI[37], CPAT[38] and pfam[39] software with defeat parameters to predict the potential lncRNAs. Only transcripts predicted as lncRNA shared among four tools were regarded as candidate lncRNA. Then the cis-target gene of lncRNA were defined as neighbor gene in 100 kb genomic distance from the lncRNA and were identified using in-house script. The trans-target prediction of lncRNAs was performed by LncTar software[40].

DMLs and DMRs calling

The differential methylation locus (DMLs) and differential

methylation regions (DMRs) between broilers and layers at each

comparison were detected separately using Dispersion Shrinkage for

Sequencing Data (DSS) package in R[41-44]. The differential methylation

regions (DMRs) were then calculated in with default parameters.

Subsequently, DMRs were annotated using ChIPseeker package in R[45].

Gene overlapped with at least one DMR is defined as differential

methylation gene (DMG). Common DMRs among 4 developmental

stages were identified by merging all positions of DMRs in 24 samples

and re-calculating the methylation level for each merged DMR position

with an average approach using mCpG data.

DEGs and DELs calling

449   The differential expression genes (DEGs) calling and the differential

450 expression lncRNA (DEL) calling between two populations at each time

451 point were performed separately using the DEseq[46]. The results were

452 filtering with the criteria: (1) fold change >2 (2) FDR<0.5. The transcripts

453 satisfied both standards were regarded as DEGs or DELs.

Functional enrichment analysis and WGCNA analysis

455   Gene ontology enrichment analyses were conducted for DMGs at

456 E10, E13, E16, E19 comparisons respectively to explore their potential

457 roles in muscle development. These analyses were performed by

458 clusterProfiler package implemented in R[47]. A hypergeometric test was

459 applied to map DMGs to terms in the GO database to search for

significantly enriched terms in DMGs compared to the genome background.

The WGCNA analysis was performed using WGCNA package implemented in R ENREF_48[48]. We used all the differential expression lncRNAs and all the genes as input. Then, variable coefficient was used to filter transcripts with low expression change. The variable coefficient was calculated as follow: $C_v = \sigma/\mu$. The σ is the standard deviation and μ represents the mean value of expression of input transcripts. Only transcripts with ranked top 30% high C_v value were used for WGCNA analysis. After the entire network was constructed, only genes with connectivity more than 0.15 were selected for subsequent subnetwork analysis.

Validation for RNA-seq by quantitative Real-time RCP(Q-PCR)

Total RNA was purified and reversely transcribed into cDNA using PrimerScriptR RT reagent Kit with gDNA Eraser (Takara Biotechnology (Dalian) Co., Ltd) following the specification. Quantities of mRNA were then measured with qRT-PCR using a CFX96TM real-time PCR detection system (Bio-Rad, USA). The qRT-PCR assays were then performed with a volume of 20 μL containing 10 μL SYBR Green Mixture, 7 μL deionized water, 1 μL template of cDNA, 1 μL of each primer and with following thermal conditions: 95 °C for 5 min, 45 cycles of 95 °C for 10 sec, 60 °C for 10 sec, 72 °C for 10 sec. Primer sequences

used for qRT-PCR assays are displayed in Supplementary Table 17. β-actin gene was used as internal control. Each qPCR assay was carried out in triplicate. The relative gene expression was calculated by using the $2-\Delta\Delta Ct$ method.

Cell cultures

Post-hatch chickens (7-day-old commercial generation Avian broiler chicks) were purchased from Wenjiang Charoen Pokphand Livestock & Poultry Co., Ltd. The pectoralis muscle was removed and used for preparation of primary myogenic cultures. About 5 g of muscle was finely minced and treated with 0.1% collagenase I (Sigma, MO, USA) followed by 0.25% trypsin (Hyclone, UT, USA) to release cells. Then, the cell suspension was subjected to Percoll density centrifugation to separate myoblasts from contaminating myofibril debris and nonmyogenic cells. Cells were plated in 25 cm3 cell culture bottles with complete medium [DMEM/F12 (Invitrogen, Carlsbad, CA) +15% FBS (Gibco, NY, USA) +10% horse serum (Hyclone, UT, USA) +1% penicillin-streptomycin (Solarbio, Beijing, China) +3% chicken embryo extraction]. The cells cultured at 37 °C and 5% CO_2 with saturating humidity, which were allowed to proliferate in growth medium for 2-4 d, and the medium was refresh every 24 h. To induce differentiation, satellite cells were grown to 80% confluence in growth medium, and the replaced with differentiation medium composed of DMEM, 2% horse serum and 1%

penicillin-streptomycin, and the medium was refreshed every 24 h.

LncRNA silencing

Chicken satellite cells were cultivated in 6-well plates and transfected with siRNAs: 5'-GGAAGGGAGUAGGUGGUAATT-3' and 5'-UUACCACCUACUCCCUUCCTT -3'; Sangon Biotech, Shanghai, China) when grown to a density of approximate 70% in plates. In contrast, control cells were transfected with negative siRNA with same other condition. The transfection reagent was Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA). The knockdown efficiency was assessed by quantitative RT-PCR of lncRNA MYH1-AS.

Microscopy

Cellular morphology was evaluated in differentiated myotubes by phase-contrast microscopy without preliminary fixation. Pictures were produced using the Olympus IX73 inverted microscope (OLYMPUS, Tokyo, Japan) and the Hamamatsu C11440 digital camera (HAMAMATSU, Shizuoka, Japan).

Western blot assay

The cells were collected from the cultures, placed in the RIPA lysis buffer on ice (BestBio, Shanghai, China). The whole proteins were subjected to 10% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred to polyvinylidene

fluoride membranes (PVDF; Millipore Corporation, Billerica, MA, USA). The PVDF membrane was incubated with 5% defatted milk powder at room temperature for 1 h, then incubation with the following specific primary antibodies at 4℃ overnight: anti-MyoG (Abcam), anti-MyHC (Abcam) and anti-β-Actin (Abcam). The secondary antibodies HRP-labeled rabbit IgG (Cell Signaling) were added at room temperature for 1h. Following each step, the membranes were washed five times with PBS-T for 3 min. The proteins were visualized by enhanced chemiluminescence (Amersham Pharmacia Biotech, Piscataway, NJ, USA) with a Kodak imager (Eastman Kodak, Rochester, NY, USA). Quantification of protein blots was performed using the Quantity One 1-D software (version 4.4.0) (Bio-Rad, Hercules, CA, USA) on images acquired from an EU-88 image scanner (GE Healthcare, King of Prussia, PA, USA).

Fig 1. Genome-wide profiles of DNA methylation among different

sample groups. **(a)** Genomic methylation level in either layers or broilers

at E10, E13, E16, E19, respectively. Methylation level were range from 0

to 1. **(b-d)** Proportion of mCpG in different genomic features at different

developmental stages in mCG, mCHG and mCHH contexts, respectively.

(e-g) Methylation level of CpGs was equally divided into 10 intervals and

the percentage of each interval were measured using E10 as example.

Fig 2. Comparatively measurement of methylation level of genes

and lncRNA. **(a)** Comparison of methylation level of genes or lncRNAs

between layers and broilers in three different contexts. **(b)** Measurement

of methylation level of different types of lncRNAs. * $P<0.05$, ** $P<0.01$

for comparison between two chicken lines. The red star means the

methylation level of layers is significantly higher than broilers whereas

the green star represents an opposite result. **(c-d)** Genomic methylation

around genes and lncRNAs were measured across the genome,

respectively. Transcripts were separated into seven regions (upstream,

first exon, first intron, inner exon, inner intron, last exon and downstream)

and each region was equally divided into 20 bins for visualization.

Fig 3. Methylation level of different types of TEs using E19 as an

example. **(a)** Comparatively measurement of methylation of SINE, LINE,

LTR, DNA, Satellite regions between two chicken lines in mCG context. **(b)** Methylation of different types of TEs for upstream, body and downstream regions in three different contexts using 20 bins across the whole genome.

Fig 4. Analyses of DMRs at 4 developmental stages. DMR calling were performed in mCG, mCHG and mCHH, respectively. **(a)** Numbers of DMRs in different genomic features (promoter, exon, intron, intergenic, and UTR regions). **(b)** Relative proportion of hyper DMRs to hypo DMRs in different CpG contexts. **(c)** The results of Gene Ontology (GO) analysis for genes with overlapped with DMR. Only part of the terms was selected for display. The red color means GO-BP terms, the blue color means GO-CC terms whereas green color represents GO-MF terms. The number in bracket means number of genes enriched in a specific term.

Fig 5. Heatmap clustering analysis and PCA analysis. **(a)** Heatmap clustering using merged common DMRs among 24 samples (see Materials and Methods). **(b)** The result of PCA analysis using common DMRs among 24 samples. Only the first component and the second component were visualized.

Fig 6. LncRNAs idenditication and correlation analysis between

methylome and transcriptome. **(a)** Number of different types of lncRNAs in all developmental stages. **(b)** Venn diagram of lncRNAs identified through different software. **(c)** Hierarchical cluster analysis of lncRNAs using their expression level. Replicates were merged together in the analysis. **(d-g)** The genes and lncRNAs were divided into five groups based on their expression levels, respectively. Then the methylation level around TSS and TES of each group were measured using 20 bins across the whole genome for layers and broilers.

Fig 7. Comprehensive analysis of lncRNA MYH1-AS. **(a)** visualization of the transcript of MYH1-AS and DMR overlapped it. **(b)** Correlation between methylation of DMR and expression of MYHA-AS using Spearman method. **(c)** Correlation between expression of MYH1-AS and expression of its potential target MYH1E. **(d)** The whole gene-lncRNA network and subnetwork including MYH1-AS extracted from the entire network. (**e**) Relationship between correlation and connectivity of gene and MYH1-AS. The red points represent genes with both high connectivity and correlation with MYH1-A and were selected for subsequent GO analysis. **(f)** Comparison of connectivity value between genes selected (red points) and all genes with in the subnetwork (background). * P <0.05, ** P <0.01 for comparison between selected genes and background. **(g)** Results of GO analysis for genes selected.

Fig 8. **(a)** Expression level of MYH1-AS in layers and broilers at different developmental stages. **(b)** Verification of lncRNA MYH1-AS expression at four developmental stages by qPCR. **(c)** lncRNA Silencing efficiency. ***** P <0.05, ****** P <0.01 for comparison between control and silenced group. **(d-f)** The mRNA expression of MyoD1, MyH3 and MyoG in control and MYH1-AS silenced groups, respectively. ***** P <0.05, ****** P <0.01 for comparison between control and silenced group. **(g-h)** The morphological changes in myotubes after silencing. **(8i)** The protein expression of MyHC and MyoG comparison between control and silenced group, respectively.

References

References

1.   Goldberg, A. D., Allis, C. D. and Bernstein, E. 2007, Epigenetics: a landscape takes shape.
Cell, **128**, 635-638.

2.   Guttman, M., Amit, I., Garber, M., et al. 2009, Chromatin signature reveals over a
thousand highly conserved large non-coding RNAs in mammals. *Nature*, **458**, 223.

3.   Feinberg, A. P. 2007, Phenotypic plasticity and the epigenetics of human disease. *Nature*,
447, 433-440.

4.   Sanford, J. P., Clark, H. J., Chapman, V. M. and Rossant, J. 1987, Differences in DNA
methylation during oogenesis and spermatogenesis and their persistence during early
embryogenesis in the mouse. *Genes & Development*, **1**, 1039-1046.

5.   Zhou, Z. Y., Li, A., Wang, L. G., et al. 2015, DNA methylation signatures of long intergenic
noncoding RNAs in porcine adipose and muscle tissues. *Scientific Reports*, **5**, 15435.

6.   Yang, Y., Liang, G., Niu, G., et al. 2017, Comparative analysis of DNA methylome and
transcriptome of skeletal muscle in lean-, obese-, and mini-type pigs. *Scientific Reports*,
7.

7.   Cesana, M., Cacchiarelli, D., Legnini, I., et al. 2011, A long noncoding RNA controls
muscle differentiation by functioning as a competing endogenous RNA. *Cell*, **147**,
358-369.

8.   Jaenisch, R. and Bird, A. 2003, Epigenetic regulation of gene expression: how the genome
integrates intrinsic and environmental signals.

9.   Miyata, K., Miyata, T., Nakabayashi, K., et al. 2015, DNA methylation analysis of human
myoblasts during in vitro myogenic differentiation: de novo methylation of promoters of
muscle-related genes and its involvement in transcriptional down-regulation. *Human*
Molecular Genetics, **24**, 410-423.

10.   Huszar, G. 1972, Developmental Changes of the Primary Structure and Histidine
Methylation in Rabbit Skeletal Muscle Myosin. *Nature New Biology*, **240**, 260-264.

11.   Huang, Y. Z., Sun, J. J., Zhang, L. Z., et al. 2014, Genome-wide DNA methylation profiles
and their relationships with mRNA and the microRNA transcriptome in bovine muscle
tissue (Bos taurine). *Scientific Reports*, **4**, 6546.

12.   Zhang, M., Yan, F. B., Li, F., et al. 2017, Genome-wide DNA methylation profiles reveal
novel candidate genes associated with meat quality at different age stages in hens.
Scientific Reports, **7**, 45564.

13.   Carriè´„ E., Dé"†Ezvillanueva, A., Lois, S., et al. 2015, Deconstruction of DNA methylation
patterns during myogenesis reveals specific epigenetic events in the establishment of the
skeletal muscle lineage. *Stem Cells*, **33**, 2025-2036.

14.   Zhi, H., Li, X., Wang, P., et al. 2017, Lnc2Meth: a manually curated database of regulatory
relationships between long non-coding RNAs and DNA methylation associated with
human disease. *Nucleic Acids Research*.

15.   Li, J., Li, R., Wang, Y., et al. 2015, Genome-wide DNA methylome variation in two
genetically distinct chicken lines using MethylC-seq. *Bmc Genomics*, **16**, 1-13.

16.   Li, Q., Wang, Y., Hu, X., Zhao, Y. and Li, N. 2015, Genome-wide Mapping Reveals
Conservation of Promoter DNA Methylation Following Chicken Domestication. *Sci Rep*, **5**,
8748.

17.   Li, Q., Li, N., Hu, X., et al. 2011, Genome-wide mapping of DNA methylation in chicken.
679   *Plos One*, **6**, e19428.

680   18.   Jones, P. A. 2012, Functions of DNA methylation: islands, start sites, gene bodies and
681   beyond. *Nature Reviews Genetics*, **13**, 484-492.

682   19.   Li, S., Zhu, Y., Zhi, L., et al. 2016, DNA Methylation Variation Trends during the Embryonic
683   Development of Chicken. *Plos One*, **11**, e0159230.

684   20.   Hu, Y., Xu, H., Li, Z., et al. 2013, Comparison of the genome-wide DNA methylation
685   profiles between fast-growing and slow-growing broilers. *Plos One*, **8**, e56411.

686   21.   Wang, H., Wang, J., Ning, C., et al. 2017, Genome-wide DNA methylation and
687   transcriptome analyses reveal genes involved in immune responses of pig peripheral
688   blood mononuclear cells to poly I:C. *Scientific Reports*, **7**, 9709.

689   22.   Lorincz, M. C., Dickerson, D. R., Schmitt, M. and Groudine, M. 2004, Intragenic DNA
690   methylation alters chromatin structure and elongation efficiency in mammalian cells.
691   *Nature Structural & Molecular Biology*, **11**, 1068-1075.

692   23.   Garcia-Perez, J. L., Widmann, T. J. and Adams, I. R. 2016, The impact of transposable
693   elements on mammalian development. *Development*, **143**, 4101-4114.

694   24.   Waterland, R. A. and Jirtle, R. L. 2003, Transposable elements: targets for early nutritional
695   effects on epigenetic gene regulation. *Molecular and cellular biology*, **23**, 5293-5300.

696   25.   Kamm, K. E. and Stull, J. T. 2001, Dedicated myosin light chain kinases with diverse
697   cellular functions. *Journal of Biological Chemistry*, **276**, 4527-4530.

698   26.   Zhi, G., Ryder, J. W., Huang, J., et al. 2005, Myosin light chain kinase and myosin
699   phosphorylation effect frequency-dependent potentiation of skeletal muscle contraction.
700   *Proc Natl Acad Sci U S A*, **102**, 17519-17524.

701   27.   Ohsawa, N., Koebis, M., Mitsuhashi, H., Nishino, I. and Ishiura, S. 2015, ABLIM1 splicing is
702   abnormal in skeletal muscle of patients with DM 1 and regulated by MBNL, CELF and
703   PTBP 1. *Genes to Cells*, **20**, 121-134.

704   28.   Holness, M. J., Kraus, A., Harris, R. A. and Sugden, M. C. 2000, Targeted upregulation of
705   pyruvate dehydrogenase kinase (PDK)-4 in slow-twitch skeletal muscle underlies the
706   stable modification of the regulatory characteristics of PDK induced by high-fat feeding.
707   *Diabetes*, **49**, 775-781.

708   29.   Ying, F., Gu, H., Xiong, Y. and Zuo, B. 2017, Analysis of differentially expressed genes in
709   gastrocnemius muscle between DGAT1 transgenic mice and wild-type mice. *BioMed
710   research international*, **2017**.

711   30.   Caremani, M., Yamamoto, D. L., Nigro, V., Lombardi, V., Bang, M. L. and Linari, M. 2014,
712   The Role of Myopalladin in Skeletal Muscle. *Biophysical Journal*, **106**, 767a.

713   31.   Shum, A. M., Mahendradatta, T., Taylor, R. J., et al. 2012, Disruption of MEF2C signaling
714   and loss of sarcomeric and mitochondrial integrity in cancer-induced skeletal muscle
715   wasting. *Aging (Albany NY)*, **4**, 133.

716   32.   Luo, B., Ye, M., Xu, H., et al. 2018, Expression analysis, single-nucleotide polymorphisms
717   of the Myoz1 gene and their association with carcase and meat quality traits in chickens.
718   *Italian Journal of Animal Science*, 1-9.

719   33.   Krueger, F. and Andrews, S. R. 2011, Bismark: a flexible aligner and methylation caller for
720   Bisulfite-Seq applications. *Bioinformatics*, **27**, 1571-1572.

721   34.   Kim, D., Langmead, B. and Salzberg, S. L. 2015, HISAT: a fast spliced aligner with low

722    memory requirements. *Nature Methods*, **12**, 357.

723 35. Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T. C., Mendell, J. T. and Salzberg, S. L.
724    2015, StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.
725    *Nature Biotechnology*, **33**, 290-295.

726 36. Kong, L., Zhang, Y., Ye, Z. Q., et al. 2007, CPC: assess the protein-coding potential of
727    transcripts using sequence features and support vector machine. *Nucleic Acids Research*,
728    **35**, W345.

729 37. Sun, L., Luo, H., Bu, D., et al. 2013, Utilizing sequence intrinsic composition to classify
730    protein-coding and long non-coding transcripts. *Nucleic Acids Research*, **41**, e166.

731 38. Wang, L., Park, H. J., Dasari, S., Wang, S., Kocher, J. P. and Wei, L. 2013, CPAT:
732    Coding-Potential Assessment Tool using an alignment-free logistic regression model.
733    *Nucleic Acids Research*, **41**, e74-e74.

734 39. Finn, R. D., Tate, J., Mistry, J., et al. 2011, A: The Pfam protein families database. *Nucleic*
735    *Acids Research*, **32**, D138.

736 40. Li, J., Ma, W., Zeng, P., et al. 2015, LncTar: a tool for predicting the RNA targets of long
737    noncoding RNAs. *Briefings in Bioinformatics*, **16**, 806.

738 41. Wu, H., Wang, C. and Wu, Z. 2012, A new shrinkage estimator for dispersion improves
739    differential expression detection in RNA-seq data. *Biostatistics*, **14**, 232-243.

740 42. Feng, H., Conneely, K. N. and Wu, H. 2014, A Bayesian hierarchical model to detect
741    differentially methylated loci from single nucleotide resolution sequencing data. *Nucleic*
742    *acids research*, **42**, e69-e69.

743 43. Wu, H., Xu, T., Feng, H., et al. 2015, Detection of differentially methylated regions from
744    whole-genome bisulfite sequencing data without replicates. *Nucleic acids research*, **43**,
745    e141-e141.

746 44. Park, Y. and Wu, H. 2016, Differential methylation analysis for BS-seq data under general
747    experimental design. *Bioinformatics*, **32**, 1446-1453.

748 45. Yu, G., Wang, L. G. and He, Q. Y. 2015, ChIPseeker: an R/Bioconductor package for ChIP
749    peak annotation, comparison and visualization. *Bioinformatics*, **31**, 2382-2383.

750 46. Anders, S., Mccarthy, D. J., Chen, Y., et al. 2013, Count-based differential expression
751    analysis of RNA sequencing data using R and Bioconductor. *Nature Protocols*, **8**, 1765.

752 47. Yu, G., Wang, L.-G., Han, Y. and He, Q.-Y. 2012, clusterProfiler: an R package for
753    comparing biological themes among gene clusters. *Omics: a journal of integrative*
754    *biology*, **16**, 284-287.

755 48. Langfelder, P. and Horvath, S. 2008, WGCNA: an R package for weighted correlation
756    network analysis. *BMC bioinformatics*, **9**, 559.

Fig 1a

Methylation Level of Different Developmental Stages

Fig 1b

Fig 1c

Fig 1d

Fig 1e

mCG

Fig 1f

mCHG

Fig 1g

mCHH

Fig 5a

Heatmap of 24 Samples Using Merged Common DMRs

Fig 5b

PCA Of 24 Samples Using Merged Common DMRs

Fig 8a

lncRNA Expression by RNA-seq

mRNA Expression level of MYH1-AS

Layer
Broiler

E 10   E 13   E 16   E 19

Fig 8b

qPCR Verification of lncRNA Expression

Relative mRNA Expression of lncRNA

Layer
Broiler

E10   E13   E16   E19

Fig 8c

Silencing Efficiency

Relative lncRNA abundance

Control   siRNA

**

Fig 8d

MyoD1

Relative Myod mRNA abundance

Control   siRNA

**

Fig 8e

MyH3

Relative MyH3 mRNA abundance

Control   siRNA

**

Fig 8f

MyoG

Relative Myog mRNA abundance

Control   siRNA

**

Fig 8g

Fig 8h

Fig 8i

MyHC

MyoG

β-actin