Myocardin ablation in a cardiac-renal rat model

Anupam Mittal[1&2],PhD; Santanu Rana[3],PhD; Rajni Sharma[4],PhD; Akhilesh

Kumar[4],PhD; Rishikesh Prasad,[4]MSc, Satish K Raut[4],PhD; Sagartirtha Sarkar[3],PhD;

Uma Nahar Saikia,[5]MD; Ajay Bahl[2],DM; Perundurai S Dhandapany[1&6], PhD; Madhu*

Khullar[4],PhD.*

[1] Centre for Cardiovascular Biology and Disease, Institute for Stem Cell Biology and

Regenerative Medicine (inStem), Bangalore, India.

[2] Department of Cardiology, Post Graduate Institute of Medical Education and

Research, Chandigarh, India.

[3]Department of Zoology, University of Calcutta, India.

[4]Department of Experimental Medicine and Biotechnology, Post Graduate Institute of

Medical Education and Research, Chandigarh, India.

[5]Department of Histopathology, Post Graduate Institute of Medical Education and

Research, Chandigarh, India.

[6]The Knight Cardiovascular Institute, Oregon Health and Science University,

Portland, OR, USA Department of Medicine, Oregon Health and Science University,

Portland, OR, USA Department of Molecular and Medical Genetics, Oregon Health

and Science University, Portland, USA, OR,

Short/Running Title: Myocardin ablation in a cardiac-renal rat model

*Corresponding Authors

Prof. Madhu Khullar

Department of Experimental Medicine and Biotechnology

Postgraduate Institute of Medical Education and Research, Chandigarh, India

Phone no: 0172-2755229, +91-9316131057

Email Address: madhu.khullar@gmail.com

Dr. Dhandapany Perundurai

Centre for Cardiovascular Biology and Disease

Institute for Stem Cell Biology and Regenerative Medicine (inStem), Bangalore, India

Phone no: +91-8023666621

Email Address: dhan@instem.res.in
31

Number of Figures and Tables: 5 Figures and 1 Table

Supplementary File: 1

Abstract

Cardiorenal syndrome is defined by primary heart failure conditions influencing or

leading to renal injury or dysfunction. Dilated cardiomyopathy (DCM) is a major co-

existing form of heart failure (HF) with renal diseases. Myocardin (MYOCD), a

cardiac-specific co-activator of serum response factor (SRF), is increased in DCM

mice and patient cardiac tissues and plays a crucial role in the pathophysiology of

DCM. Inhibiting the increased MYOCD has shown to be partially rescuing the DCM

mice phenotype. However, expression levels of MYOCD in the cardiac tissues of the

cardiorenal syndromic patients and the beneficial effect of inhibiting MYOCD in a

cardiorenal syndrome model remains to be explored.

Here, we analyzed the expression levels of MYOCD in the DCM patients with and

without renal diseases. We also explored, whether cardiac specific silencing of

MYOCD expression could ameliorate the cardiac remodeling and improve cardiac

function in a renal artery ligated rat model (RAL). We observed an increase in

MYOCD levels in the endomyocardial biopsies of DCM patients associated with renal

failure compared to DCM alone. Silencing of MYOCD in RAL rats by a cardiac

homing peptide conjugated MYOCD siRNA resulted in attenuation of cardiac

hypertrophy, fibrosis and restoration of the left ventricular functions.

Our data suggest hyper-activation of MYOCD in the pathogenesis of the cardiorenal

failure cases. Also, MYOCD silencing showed beneficial effects by rescuing cardiac

hypertrophy, fibrosis, size and function in a cardiorenal rat model.

Introduction

DCM is a major cause of HF,[1] accounting for nearly 1/3rd of total cases. Many of

these patients subsequently display kidney dysfunction or injury leading to

cardiorenal syndrome. More than half of the heart failure patients show renal

diseases. Co-existence of cardiac and renal dysfunction in the patients increases the

mortality significantly compared to cardiac or renal disease alone patients.

Various molecular pathways including Renin-angiotensin-aldosterone system (RAAS)

are shown to be influencing the cardiorenal syndrome. Notably, circulating Ang II (an

essential component of RAAS) affects cardiac function by, increasing systemic

arteriolar vasoconstriction, vascular resistance, and cardiac afterload through AT1

receptor-mediated endothelial dysfunction [2].

Ang II has been shown to induce MYOCD under hypoxic condition[3]. MYOCD is a

cardiac-specific transcriptional co-activator present in cardiomyocytes and smooth

muscle cells. MYOCD is involved in heart development and cardiomyocyte

differentiation[4,5]. Also, MYOCD is required for maintenance of structural integrity,

cardiomyocyte survival, and heart function[5–7] . Besides, cardiomyocytes and smooth

muscle cells MYOCD expression has been detected in fibroblasts[8]. MYOCD has

been shown to promote fibroblast to myofibroblast differentiation and to inhibit cell

proliferation [9]. Forced expression of MYOCD in fibroblasts induces cardio-myogenic

properties alone[8] and/or in combination with other factors[10]. Transforming growth

factor (TGF-β) was shown to induce MYOCD expression in fibroblasts and vice-

versa[9]. TGF-β induction of MYOCD expression in the infarcted heart may have a

potential function in fibroblast-to-myofibroblast transition, similar to Myocardin related

transcription factor (MRTF)-A and MRTF-B which have been shown to be key

regulator in fibroblast to myofibroblast differentiation induced by TGF-β1[11]. Further,

deletion of MYOCD gene in the adult murine heart resulted in dilated

cardiomyopathy, and rapid death due to heart failure[5]. Upregulation of MYOCD

expression has been shown in cardiac hypertrophy [3,12,13] and MYOCD

overexpression in mouse cardiomyocytes resulted in activation of genes associated

with cardiac hypertrophy[12]. Increased cardiac MYOCD expression has been reported

in various cardiac ailments including DCM patients with end-stage HF [14,15]. MYOCD

has been shown to be a pro-hypertrophic factor in cardiac remodeling induced in

multiple models[3,12,13]. However, there is no report so far, suggesting the role of

MYOCD in cardiorenal syndrome.

In the present study, we analyzed the cardiac-specific expression of MYOCD in DCM

patients with renal disease and DCM alone cases. The results showed the MYOCD

is overexpressed in the DCM patients with renal disease compared to DCM alone

cases. In addition, the effects of cardiac-specific silencing of MYOCD was explored in

a cardiac renal syndrome rat model. The cardiac-specific silencing of MYOCD in rats

decreased the expression of upregulated hypertrophic and fibrotic genes leading to

restoration of left ventricular function.

100   Material and Methods

101   Study Population

Thirty consecutive biopsies were taken from left ventricle region from idiopathic DCM

(IDCM) patients, attending Cardiology Clinic at the Department of Cardiology,

Postgraduate Institute of Medical Education and Research, Chandigarh, India

between Jan 2011-2014. Inclusion criteria for recruitment of DCM patients,

diagnosed after echocardiography, defined by left ventricular ejection fraction (LVEF)

≤ 40% and chronic mild to severe HF (NYHA functional class II to IV). All patients

underwent left cardiac catheterization and coronary angiography before their

inclusion in the study. Exclusion criteria were: the presence of significant coronary

artery disease defined as lumen stenosis in >50% of any coronary artery, severe

primary valve disease, uncontrolled systemic, hypertension, hypertrophic or

restrictive cardiomyopathy, chronic systemic disease like myocarditis, thyrotoxicosis,

HIV disease and drug abuse. All recruited IDCM subjects were on optimal

medication, angiotensin-converting enzyme inhibitors, and beta-blockers but had

persistently low LVEF despite drug regime at the time of biopsy. Endomyocardial

biopsy from left ventricle region (n=14) taken from subjects undergoing surgery for

ventricular septal defect (VSD), served as controls. The VSD patients recruited in the

study have normal LVEF with no right or left ventricular dysfunction. The study was

approved by the Institutional Ethics Committee (8443-PG-1TRg-10/4497),

Postgraduate Institute of Medical Education and Research, Chandigarh and written

informed consent was taken from all patients for participation in the study. We

confirm that all experiments were performed in accordance with relevant guidelines

and regulations of Institutional Ethics Committee, Postgraduate Institute of Medical

Education and Research, Chandigarh. Inclusion and Exclusion criteria is given in

Supplementary methods. Baseline data of patients and control subjects is available in our earlier publication.[16]

Renal artery ligation model of cardiac renal syndrome (RAL)

RAL was developed in adult Wistar rats as described earlier[17]. Briefly, right renal artery of 20-week-old male rats (250–300 g; n =8) was ligated under anaesthesia, rats that underwent similar procedure without ligation served as sham-operated control group (n = 8). Animals were sacrificed 15 days after surgery and hearts were removed and either fixed in Karnovsky's fixative for histological study or immediately stored in liquid nitrogen for RNA and protein isolation. Cardiac hypertrophy was measured at the gross by heart weight (HW in mg) to body weight (BW in g) ratio. Expression of cardiac fibrosis (Col-1a, Col-4a CTGF, and FGF) and hypertrophy (ANP and β-MHC) genes at mRNA and protein levels was studied by qRT-PCR and western blotting respectively. Histopathological analysis (H&E staining, MT staining and EVG staining) was done in formalin fixed cardiac tissues to asses cardiac remodeling (hypertrophy and fibrosis). This study was approved by Institutional Animal Ethics Committee (IAEC), Postgraduate Institute of Medical Education and Research, Chandigarh and all experiments were done as per the guidelines and regulations of the committee.

Modulation of Myocardin Gene

Myocardin (MYOCD) was transiently over-expressed in H9c2 cells or CFs by transfection with 2μg of pcDNA3-MYOCD-HA_v1, using lipofectamine® 2000 transfection reagent (Invitrogen) according to the manufacturer's instructions. For MYOCD cellular knockdown, H9c2 cells or CFs were transfected with 1μg/ml of MYOCD siRNA (Table S1 in supplementary data) which targets both the cardiac

149   isoforms (Silencer® Select pre-designed MYOCD siRNA; ID s140959; Ambion,

150   USA). Cells transfected using scrambled sequence anti-miR (Ambion, USA) served

151   as a negative control. Transfection efficiency was measured using Cy3 labelled

152   negative siRNA by FACS Canto™ II Cell Analyser and by Olympus IX 73 Fluorescent

153   microscopy.

154   Cardiac specific silencing of MYOCD Gene

Cardiac MYOCD gene expression was inhibited in RAL rats using MYOCD specific

siRNA, conjugated with a cardiac specific 20mer homing peptide sequence

WLSEAGPVVTVRALRGTGSW encapsulated in stearic acid modified Carboxy

methyl chitosan as described earlier[18], as per protocol shown in Supplementary

Figure 1. Animals were sacrificed using sodium pantothenate as anaesthesia and

performing cervical dislocation. Hearts were removed, examined grossly, and

histopathological analysis (H&E staining, MT staining, EVG staining and hypertrophy

and fibrotic gene specific staining) was performed. RNA and protein was isolated

from left ventricular tissue. Expression of hypertrophic (ANP and β-MHC) and pre-

fibrotic genes (Col-1a, Col-3a, Col-4a CTGF, TGF β and FGF) was measured at

mRNA and protein levels.

166   Histology and Immunohistochemistry

Tissues were fixed in 4% paraformaldehyde, embedded in paraffin, and sectioned at

4-µm thickness. Hematoxylin/eosin for cell size and morphology and Masson's

trichrome for collagen staining were performed using standard procedures. β-MHC

and CTGF staining was performed on paraffin-embedded sections using anti-β-MHC

(1:100) and anti-CTGF (1:200). Nuclei were visualized using haematoxylin stain.

Images were obtained by Olympus IX 73 microscope.

MYOCD antibody specificity experiment

MYOCD antibody specificity was confirmed by immunostaining in cell lines which have high expression and sparse or null expression of MYOCD to ascertain that it is not binding to other family members of Myocardin like proteins and specific for MYOCD (Supplementary Figure 2). Imaging was done on Olympus FV1000 confocal microscopy.

Angiotensin II (Ang II) induced cardiomyocyte hypertrophy

H9c2 rat cardiomyocytes were cultured in Dulbecco's Modified Eagle's Medium (DMEM), supplemented with 10% foetal bovine serum under 5% CO_2 at 37°C. After 12 hrs of serum starvation, H9c2 cardiomyocytes were stimulated with 1µM angiotensin II (Ang-II) (Sigma Aldrich) or 7.5 ng/ml transforming growth factor-β (TGF-β) (Peprotech) for 24hrs, 48hrs and 72hrs. Expression of hypertrophic genes atrial natriuretic protein (ANP) was measured at mRNA and protein levels to confirm the development of hypertrophy.

Ang II induced fibrosis in cardiac fibroblasts

Primary adult rat cardiac fibroblasts (CFs) were isolated from ventricular tissues of 6 to 8 week old male Wistar rats, (150-200gm) as previously described[19]. After 12hrs of serum starvation, CFs were stimulated with 1µM Ang II for 24hrs (these were optimal conditions for time and dose of Ang II as standardized in our laboratory for induction of fibrosis). Expression of various fibrotic genes (Col-1a, Col-3a, Col-4a CTGF, and FGF-b) was measured at mRNA levels. Institutional Animal Ethics Committee, Postgraduate Institute of Medical Education and Research, Chandigarh approved all

animal experiments were performed in accordance with regulations specified by

CPCSEA.

Western Blotting

Cells and tissue protein lysate was prepared using RIPA lysis solution and 1X

Protease inhibitor (Cat no.87785, Thermo Scientific), followed by sonication at 30%

amplitude repeated for 3-4 times, 20secs each with 10secs interval. Protein

concentration was estimated by quick start^{TM} Bradford dye reagent, 1X (Cat no. 500-

0205, Bio-Rad). Laemmli Buffer was used for preparation of proteins. Protein lysate

(50µg/well) was then separated on the basis of their molecular weight on SDS-PAGE

Bio-Rad mini SDS-PAGE gel instrument. 50µg of protein sample along with 5µl of

PageRuler^{TM} pre-stained protein ladder (Cat no. 26617, Thermo Scientific), was

loaded and gel was run initially at 50 volts until the tracking dye crosses the stacking

gel, then at 100 volt till the dye come at the bottom of the gel. After SDS-PAGE,

PVDF membrane was activated in 100% methanol and equilibrated in 1X transfer

buffer along with SDS-PAGE gel and blotting sheets, at 4°C for 15 mins. Protein wet

transfer was carried out at 30mA for 10hr in Mini-Protean Tetra System (Bio-rad).

Ponceau staining was performed in order to check the protein transfer on PVDF

membrane. PVDF membrane was then washed with 1X TBST buffer and kept in

blocking solution (5% BSA in TBST buffer) for 2hrs on shaker at RT. After blocking,

PVDF membrane was washed 3 times in TBST buffer (1X) for 10mins each and was

then incubated in primary antibody (List of the antibodies used in the study with the

working dilution, source and catalog number are provided as Table S2 in

Supplementary data) for 4°C overnight. PVDF membrane was then washed 3 times

with 1X TBST buffer for 10mins each and then incubated with HRP labelled

secondary antibody for 40 mins at RT. PVDF membrane was then washed 3times

with 1X TBST buffer for 10mins each and target protein reactive with primary

antibodies was detected. After incubation with the appropriate secondary antibodies,

blots were visualized using the EZ-ECL detection system (Cat no.20-500-120A,

Biological Industries) and the FluorChem M luminescent image analyzer (Protein

Simple, Biotechne brand, USA). Densitometry analysis of band intensity was

performed using Image J sofware (NIH, USA).

Statistical Analysis

The statistical analysis was carried out using Statistical Package for Social Sciences

(SPSS Inc., Chicago, IL, and version 21.0 for Windows)/ Graph Pad Prism 5.0. All

quantitative variables were estimated using mean and measures of dispersion

(standard deviation and standard error). Normality of data was checked by measures

of skewness and Kolmogorov Smirnov tests of normality. Expression of genes were

compared using Mann Whitney or Student's t-test depending on the normality of

data. One way/Two way ANOVA was performed when multiple comparisons were

done within various experimental groups. All statistical tests were two-sided and

performed at a significance level of $p \leq 0.05^*$, $p \leq 0.01^{**}$ and $^{***}p \leq 0.001$.

Results

Cardiac MYOCD expression level is significantly increased in cardiorenal

syndrome patients

We analyzed the mRNA and protein expression levels of MYOCD in endomyocardial

biopsies of DCM patients with and without renal failure. The mRNA and protein

expression levels of MYOCD was increased in endomyocardial biopsies obtained

from DCM patients compared to healthy controls. However, the increase is more

pronounced in DCM patients with renal failure compared to DCM alone patients.

(Figure 1a & b).

MYOCD activation induces hypertrophic and fibroblast associated genes

To know the functional relevance of increase of MYOCD in cardiorenal patients, we treated cardiomyocytes cells (H9c2) and fibroblasts with Ang II (a well-known modulator of cardiorenal disease). The treatment showed increased mRNA and protein expression of MYOCD in both cell types (Supplementary Fig. 3a, 3b & 3c), along with hypertrophy markers including Atrial Natriuretic factor (ANP), Beta-Myosin heavy chain (β-MHC) and fibrotic genes at mRNA levels such as Collagen (Col) 1a, Col 3a, Col 4a, Transforming growth factor-β (TGF-β), Connective tissue growth factor (CTGF) and Fibroblast growth factor (FGF)- β respectively (Fig. 2a, 2b & 2c). Pre-treatment with losartan (an Ang II type I receptor antagonist) reversed these effects by increasing the levels of MYOCD, ANP and β-MHC mRNA levels in cardiomyocytes, whereas, treatment with known Ang II analogue had no significant effect on MYOCD, ANP and β-MHC expression in these cells (Supplementary Fig. 3d).

In vitro **silencing of MYOCD inhibited Ang II induced cardiomyocyte hypertrophy and fibrosis**

To examine the effect of MYOCD silencing on cardiomyocyte hypertrophy, Ang II treated H9c2 cells were transfected with MYOCD siRNA. siRNA transfection significantly reduced the MYOCD levels (Supplementary Figure 4a & 4b), leading to the reduction of ANP and β-MHC levels (Figure 3a & 3b).

Also, treatment of MYOCD siRNA significantly decreased MYOCD mRNA expression levels in the cardiac fibroblasts (Fig. 3c), whereas scrambled siRNA also showed some effect on MYOCD expression in these cells but the changes found in myocd

siRNA treated cells was much significant. MYOCD siRNA treated cells resulted in a

significant decrease in mRNA expression of Col 1a, Col 3a, Col 4a,TGF-β and FGF-β

genes in Ang II treated cardiac fibroblasts (Fig. 3d). We found changes in CTGF

expression both with Ang II treatment and myocd siRNA but those changes were

non-significant.

Cardiac-specific inhibition of MYOCD gene in a cardiorenal rat model improves

left ventricular functions

To know whether MYOCD inhibition in a cardiorenal rat model can rescue the above

observed *in vitro* phenotype, we performed renal artery ligation (RAL) in rats. This

model has been shown to an excellent model for cardiac-renal heart failure[20]. RAL in

rats resulted in cardiac hypertrophy, fibrosis and systolic dysfunction (Fig. 4a & 4b;

Fig. 5a & 5b and Table 1).

Next, we injected the MYOCD siRNA conjugated with homing peptide through tail

veins of the RAL rats as described earlier to be cardiac-specific and having no

bystander effects[18]. To study the effect of heart-specific MYOCD silencing on cardiac

function, echocardiography and HW/BW ratio were measured in MYOCD siRNA

treated RAL rats 15 days post ligation. HW/BW ratio was significantly reduced in

MYOCD siRNA treated RAL rats (Table 1). A significant decrease in LVEDD and

increase in LVEF was also seen in these rats (Table 1) indicating improvement in left

ventricular function by MYOCD inhibition.

MYOCD inhibition in the RAL model attenuated cardiac hypertrophy and

fibrosis

Ablation of MYOCD gene with MYOCD siRNA conjugated with homing peptide in

RAL rats resulted in significant reduction in cardiac MYOCD expression (Fig. 4c & d),

whereas scrambled siRNA did not affect the MYOCD expression. Also, a profound

reduction in the expression of hypertrophy markers, ANP and β-MHC (Fig. 4c & d)

and a decrease in cardiomyocyte size in MYOCD siRNA treated RAL rats compared

to untreated were noticed. (Fig. 4e III & IV). There was no evidence of inflammation

or coronary occlusion in these rats, suggesting absence of ischemic lesions (Fig. 4e).

The immuno-histochemical analysis also revealed reduction in β-MHC

(Supplementary Fig. 5a-c) expression in RAL rats treated with MYOCD siRNA.

Furthermore, MYOCD siRNA treatment, resulted in a significant reduction in cardiac

fibrotic gene expressions including Col 1a, Col 3a, Col 4a, TGF β, CTGF and FGF β

compared to untreated RAL rats (Fig. 5 c & d). Histopathological examination

showed decreased collagen deposition in the treated RAL cardiac tissues (Fig. 5e),

and reduced expression levels of fibrosis-related genes including Col 1a, Col 3a, Col

4a, CTGF, FGF-β and TGF-β. There is reduction in CTGF as evident from immunno-

histochemical staining of CTGF (Supplementary Fig. 5d-f). Collectively, these data

suggested that the MYOCD inhibition can reverse the heart failure in RAL model.

Discussion

MYOCD role have been reported in various animal models of HF and in end-stage

heart failure patients[14,15]. However, its role in cardiorenal patients remains

unexplored. We observed a significant upregulation of cardiac MYOCD in

endomyocardial biopsies from cardiorenal patients. This is the first observation on

the MYOCD expression levels in the cardiorenal failure patients.

Cardiorenal syndrome patients display a significant amount of circulating renin with

Ang II production, leading to arteriolar constriction and high venous pressure. Ang II

along with aldosterone release result in systemic pressure through increased sodium

re-absorption in the distal nephrons. These ultimately worsen the patient cardiac

functions[21]. Thus, consistent activation of RAAS contributes to poor prognosis of the

cardiorenal patients. RAAS activation is shown to be associated with upregulation of

MYOCD related transcription factors leading to increased transforming growth factor-

β (TGF-β) and fibronectin[11,22]. In the present study, direct inhibition of MYOCD in Ang

II treated cardiomyocytes and fibroblasts resulted in decreased expression of

hypertrophic and pro-fibrotic markers including TGF-β. These data suggest RAAS

activation can be mitigated by MYOCD inhibition .

Torrado et al. showed that moderate inhibition of MYOCD using shRNA in

doxorubicin-induced HF (DHF) model could partly restore diastolic dysfunction and

decreased the DHF animal mortality[23]. However, such an effect remains to be

explored in a cardiorenal model. We explored whether heart-specific MYOCD

targeting could ameliorate cardiac remodeling and improve cardiac function in a rat

renal artery ligation (RAL) model. RAL is a well-established model for studying the

cardiac remodeling in renal dysfunction associated heart failure[20].

Notably, we explored if MYOCD silencing pre- or post-ligation by siRNA could have

effective in a RAL rat model showing increased expression of cardiac MYOCD with

extensive cardiac hypertrophy and fibrosis. A cardiac specific homing peptide

conjugated MYOCD siRNA to specifically inhibit cardiac MYOCD without any specific

bystander effects was employed[18]. We observed that this cardiac specific MYOCD

silencing resulted in attenuation of cardiomyocyte hypertrophy and cardiac fibrosis

when given at either both at pre-ligation or post-ligation stages.

Further, we compared whether siRNA inhibition was more effective at pre- or post-

ligation to ascertain the efficacy of MYOCD inhibition on cardiac hypertrophy and

fibrosis under these two conditions. In pre-ligation treatment, MYOCD siRNA was

introduced before the initiation of cardiac remodeling process, and in post-ligation

treatment, MYOCD siRNA was given after the initiation of cardiac remodeling. Both

the treatment groups showed significant reduction in the cardiac remodeling and

improvement of cardiac function, but post-ligation treatment showed a greater

reduction. These results suggest that MYOCD inhibition is more effective at post

ligation stage, probably as MYOCD upregulation is maximal at this stage.

Our data suggest cardiac specific MYOCD silencing in cardiorenal rats model (RAL)

resulted in the significant reduction in expression of hypertrophy markers, ANP and

β-MHC and decrease in cardiomyocyte size. Further, the histopathological and

immuno-histochemical analysis also revealed reduction in cardiomyocyte size and

expression of hypertrophy marker, β-MHC will be beneficial in ameliorating the

process of cardiac hypertrophy. A significant decrease was observed in the cardiac

expression of Col 1a, Col 3a, Col 4a, TGF-β, CTGF and FGF-β genes in MYOCD

siRNA treated RAL rats. Histopathological examination corroborated the beneficial

effect of MYOCD siRNA, as myocardial collagen deposition decreased in siRNA

treated rats, suggesting the potential in controlling the cardiac fibrosis. MYOCD

silencing was done in a controlled manner so there are less chances of bystander

effects on the normal physiology of the myocytes and fibroblasts. Cardiomyocytes

were found to have normal morphology and functional capacity after the MYOCD

siRNA treatment.

Our data document for the first time, the upregulation of cardiac MYOCD in cardio-

renal patients and cardiac-specific MYOCD silencing in a cardiorenal model has an

integrated beneficiary effect on cardiac hypertrophy, fibrosis and function. Our data

also strengthen the claims that cardiac targeted MYOCD inhibition might be a

potential therapeutic target in reversing cardiac remodeling in the cardiorenal
syndrome.

Conflict of interest: NONE

Acknowledgement and Funding Sources
We would like to acknowledge Prof. Shyam K. Singh, PGIMER, Chandigarh for
providing us with the control tissues. We are also thankful to Prof. Joseph A Miano,
Rochester Institute, USA for providing MYOCD cardiac isoform plasmids as a kind
gift. Anupam Mittal was recipient of DS Kothari Postdoctoral Fellowship, UGC and
National Postdoctoral Fellowship by DST SERB. PSD is funded by American Heart
Association-Scientist Development Grant and Welcome Trust-Intermediate
Fellowship. While this manuscript is in review, Lyu et al., reported a size variability for
MYOCD protein. Further studies using the antibodies reported by Lyu et al will further
confirm our findings[24].

Contribution Statement
Anupam Mittal, Santanu Rana, Akhilesh Kumar and Satish K Raut were involved in
acquisition of data. Anupam Mittal, Madhu Khullar and Perundurai Dhandapany
was involved in analysis and interpretation of data. Anupam Mittal, Madhu Khullar,
and Perundurai Dhandapany were involved in drafting of the manuscript. Ajay
Bahl, Uma Nahar Saikia, Rishikesh Prasd and Sagartirtha Sarkar were involved in
technical support for patient data. Rajni Sharma was involved in statistical analysis
and analysis and interpretation of data.

390
391
392
393

394
395
396
References
1. Hershberger, R. E., Hedges, D. J. & Morales, A. Dilated cardiomyopathy: the

complexity of a diverse genetic architecture. *Nat. Rev. Cardiol.* **10**, 531–547

(2013).

2. Crowley, S. D. *et al.* Angiotensin II causes hypertension and cardiac hypertrophy

through its receptors in the kidney. *Proc. Natl. Acad. Sci.* **103**, 17985–17990

(2006).

3. Chiu, C.-Z., Wang, B.-W., Chung, T.-H. & Shyu, K.-G. Angiotensin II and the ERK

pathway mediate the induction of myocardin by hypoxia in cultured rat neonatal

cardiomyocytes. *Clin. Sci. Lond. Engl. 1979* **119**, 273–282 (2010).

4. Wang, D. *et al.* Activation of cardiac gene expression by myocardin, a

transcriptional cofactor for serum response factor. *Cell* **105**, 851–862 (2001).

5. Huang, J. *et al.* Myocardin is required for cardiomyocyte survival and

maintenance of heart function. *Proc. Natl. Acad. Sci.* **106**, 18734–18739 (2009).

6. Zhang, X., Azhar, G., Zhong, Y. & Wei, J. Y. Identification of a novel serum

response factor cofactor in cardiac gene regulation. *J. Biol. Chem.* **279**, 55626–

55632 (2004).

7. Huang, J. *et al.* Myocardin regulates expression of contractile genes in smooth

muscle cells and is required for closure of the ductus arteriosus in mice. *J. Clin.*

Invest. **118**, 515–525 (2008).

8. van Tuyn, J. *et al.* Activation of cardiac and smooth muscle-specific genes in

primary human cells after forced expression of human myocardin. *Cardiovasc.*

Res. **67**, 245–255 (2005).

9. Milyavsky, M. *et al.* Inactivation of myocardin and p16 during malignant

transformation contributes to a differentiation defect. *Cancer Cell* **11**, 133–146

(2007).

10. Wang, J. *et al.* Myocardin sumoylation transactivates cardiogenic genes in

pluripotent 10T1/2 fibroblasts. *Mol. Cell. Biol.* **27**, 622–632 (2007).

11. Crider, B. J., Risinger, G. M., Haaksma, C. J., Howard, E. W. & Tomasek, J. J.

Myocardin-related transcription factors A and B are key regulators of TGF-β1-

induced fibroblast to myofibroblast differentiation. *J. Invest. Dermatol.* **131**, 2378–

2385 (2011).

12. Xing, W. *et al.* Myocardin induces cardiomyocyte hypertrophy. *Circ. Res.* **98**,

1089–1097 (2006).

13. Wang, K., Long, B., Zhou, J. & Li, P.-F. miR-9 and NFATc3 regulate myocardin in

cardiac hypertrophy. *J. Biol. Chem.* **285**, 11903–11912 (2010).

14. Torrado, M. *et al.* Myocardin mRNA is augmented in the failing myocardium:

expression profiling in the porcine model and human dilated cardiomyopathy. *J.*

Mol. Med. Berl. Ger. **81**, 566–577 (2003).

15. Kontaraki, J. E., Parthenakis, F. I., Nyktari, E. G., Patrianakos, A. P. & Vardas, P.

E. Myocardial gene expression alterations in peripheral blood mononuclear cells

of patients with idiopathic dilated cardiomyopathy. *Eur. J. Heart Fail.* **12**, 541–548

(2010).

16. Mittal, A., Sharma, R., Prasad, R., Bahl, A. & Khullar, M. Role of cardiac TBX20 in

dilated cardiomyopathy. *Mol. Cell. Biochem.* **414**, 129–136 (2016).

17. Mir, S. A. *et al.* Inhibition of signal transducer and activator of transcription 3

(STAT3) attenuates interleukin-6 (IL-6)-induced collagen synthesis and resultant

hypertrophy in rat heart. *J. Biol. Chem.* **287**, 2666–2677 (2012).

18. Rana, S. *et al.* A spatio-temporal cardiomyocyte targeted vector system for

efficient delivery of therapeutic payloads to regress cardiac hypertrophy abating

bystander effect. *J. Control. Release Off. J. Control. Release Soc.* **200**, 167–178

(2015).

19. Shivakumar, K., Nair, R. R. & Valiathan, M. S. Paradoxical effect of cerium on

collagen synthesis in cardiac fibroblasts. *J. Mol. Cell. Cardiol.* **24**, 775–780

(1992).

20. Grossman, R. C. Experimental models of renal disease and the cardiovascular

system. *Open Cardiovasc. Med. J.* **4**, 257–264 (2010).

21. Clemmer, J. S., Pruett, W. A., Coleman, T. G., Hall, J. E. & Hester, R. L.

Mechanisms of blood pressure salt sensitivity: new insights from mathematical

modeling. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* **312**, R451–R466 (2017).

22. TGF-β-induced epithelial to mesenchymal transition | Cell Research. Available at:

https://www.nature.com/articles/cr20095. (Accessed: 18th April 2018)

23. Torrado, M., Iglesias, R., Centeno, A., López, E. & Mikhailov, A. T. Targeted

Gene-Silencing Reveals the Functional Significance of Myocardin Signaling in the

Failing Heart. *PLoS ONE* **6**, e26392 (2011).

24. Lyu, Q. *et al.* CRISPR-Cas9-Mediated Epitope Tagging Provides Accurate and

Versatile Assessment of Myocardin-Brief Report. *Arterioscler. Thromb. Vasc. Biol.*

38, 2184–2190 (2018).

479   **Figure legends**
480
Fig 1. Myocardin (MYOCD) mRNA and protein expression in myocardial tissues of
DCM patients with and without renal failure and controls a) Quantitative mRNA
expression of MYOCD in DCM cardiac tissues (n=30) with and without renal failure and
control group (n=15) as determined by qRT-PCR using Taqman probe chemistry. **b)**
Representative western blot for MYOCD and GAPDH proteins in control (C1–C4) and DCM
patients (P1–P5) in LV myocardium samples. Semi-Quantitative protein expression of
MYOCD in DCM patients and controls as estimated by software Image J. Normalization of
total RNA and protein was done using GAPDH as an internal control. Data given is Mean ±
SEM; *p<0.05 Control vs DCM; ***p<0.001 Control vs DCM

Fig 2. Effect of MYOCD overexpression on Ang II induced expression of hypertrophy
and fibrosis genes in H9c2 cardiomyocytes and fibroblasts respectively a) Quantitative
mRNA expression of Atrial Natriuretic factor (ANP) in Ang-II (1μM) and TGF β (7.5 ng/ml)
treated H9c2 cells as determined by qRT-PCR. **b)** Representative western blot for ANP and
GAPDH proteins in Ang-II and TGF-β treated H9c2 cells. **c)** Quantitative mRNA expression
of Collagen (Col) 1a, Col3a, Col4a, CTGF, TGF-β and FGF-β (fibrotic marker gene) in Ang II
treated cardiac fibroblasts as determined by qRT-PCR. Normalization of total RNA and
protein was done by using GAPDH as an internal control. Data given is Mean ± SEM of
triplicates; *p<0.05; **p<0.01; ***p<0.001 Control vs Ang II and Control vs TGF β.

Fig 3. MYOCD siRNA treatment on expression of hypertrophic genes and fibrotic
genes in H9c2 and cardiac fibroblast respectively a) mRNA expression of ANP, Beta
myosin heavy chain (β-MHC) and MYOCD after treatment with MYOCD siRNA in Ang II
treated H9c2 cells **b)** ANP and MYOCD protein expression after treatment with MYOCD
siRNA in Ang II treated H9c2 cells **c)** mRNA expression of MYOCD expression after
treatment with MYOCD siRNA in cardiac fibroblast cells **d)** mRNA expression of fibrotic gene
(Col1a, Col3a, Col4a, CTGF, TGF β and FGF β) after treatment with MYOCD siRNA in Ang-
II treated fibroblast cells. Normalization of total RNA and protein was done by using GAPDH
as an internal control. Data given is Mean ± SEM of triplicates; *p<0.05; **p<0.01;
***p<0.001Control vs Ang II and Ang II vs Ang II + MYOCD siRNA

Fig 4. Effect of MYOCD siRNA treatment on expression of hypertrophic genes (ANP
and β-MHC) and on cardiomyocyte size in RAL rats a) MYOCD, ANP and β-MHC mRNA

expression in RAL rats (n=8) **b)** Representative western blot for MYOCD, ANP and GAPDH

proteins in RAL rats. **c)** mRNA expression of MYOCD, ANP and β-MHC in MYOCD siRNA

treated RAL rats **d)** Representative western blot for MYOCD, ANP and GAPDH proteins in

MYOCD siRNA treated RAL rats (before and after ligation). **e)** Haematoxylin and Eosin (H&

E) stained Photomicrographs (10X) of rat hearts: I) Control group, II) RAL group III) MYOCD

siRNA treated (before ligation) RAL Group IV) MYOCD siRNA treated (after ligation) RAL

group. Bar diagram shows mean of cardiomyocyte area (n=50) as measured by Image J

analyzer. Normalization of total RNA and protein was done using GAPDH as an internal

control. Data given is Mean ± SEM; **p<0.01; ***p<0.001 for Control vs RAL; [##]p<0.01 for

RAL vs RAL treated with MYOCD siRNA before ligation (Pre-ligation); [$]p<0.05, [$$]p<0.01

[$$$]p<0.001 for RAL vs RAL treated with MYOCD siRNA after ligation (Post-Ligation)

Fig 5. Effect of MYOCD siRNA treatment on expression of fibrotic genes (Col 1a, Col

3a, Col 4a, TGF β, CTGF and FGF β) in RAL rats (n=8): a) mRNA expression of Col 1a,

Col 3a, Col 4a, TGF β, CTGF and FGF β in RAL rats. **b)** Representative western blot for

CTGF, FGF and GAPDH proteins in RAL rats. **c)** mRNA expression of Col 1a, Col 3a, Col

4a, TGF β, CTGF and FGF β in MYOCD siRNA treated RAL rats (before and after ligation).

d) Representative western blot for CTGF, FGF and GAPDH proteins in MYOCD siRNA

treated RAL rats (before and after ligation). **e)** Photomicrographs (10X) of rat heart tissues

with Masson's Trichrome staining which is collagen specific and evaluates fibrosis: I) control,

II) RAL rats showed extensive collagen deposition III) MYOCD siRNA (before ligation)

treated RAL rats show decreased collagen deposition IV) MYOCD siRNA (after ligation)

treated RAL rats show decreased collagen deposition. Normalization of RNA and protein was

done using GAPDH as an internal control. Data given is Mean ± SEM; *p<0.05; **p<0.01 for

Control vs RAL; [#]p<0.05; [##]p<0.01 for RAL vs RAL treated with MYOCD siRNA before

ligation; [$]p<0.05; [$$]p<0.01 for RAL vs RAL treated with MYOCD siRNA after ligation.

Table 1: Echocardiographic parameters and heart weight (HW) to body weight (BW) ratio in RAL model treated with myocd siRNA				
Treatment Groups	LVDD (mm)	FS (%)	LVEF (%)	HW/BW ratio
Group I (Sham Controls)	4.34 ± 0.20	62.08 ± 1.18	78 ± 8.8	3.2 ± 0.5
Group II (Ligated)	5.83 ± 0.38*	34.6 ± 3.8*	55 ± 6.7*	4.6 ± 0.7*
Group III (Ligated + post-myocd siRNA)	4.95 ± 0.67[#]	56.8 ± 1.97[#]	72 ± 5.9[#]	3.95 ± 0.3[#]
Group IV (Ligated+ pre-myocd siRNA)	5.05 ± 0.45[$]	50.6 ± 2.85[$]	61 ± 7.2[$]	3.7 ± 0.4[$]
Group V (Ligated+ scrambled siRNA)	5.6 ± 0.4	38 ± 2.4	52 ± 4.4	4.7 ± 0.5

545 LVDD: Left Ventricular Diastolic Diameter; FS: Fraction Shortening; LVEF: Left Ventricular ejection
546 fraction. *$p < 0.05$ Group I vs Group II; [#]$p < 0.05$ Group II vs Group III; [$]$p < 0.05$ Group II vs Group IV
547
548

a

b