The lipid phosphatase-like protein PLPPR1 increases cell adhesion by modulating RhoA/Rac1 activity

Chinyere Agbaegbu Iweka[1,2][#], Sharada Tilve[1][#], Yasuhiro Katagiri[1], Herbert M Geller[1]*

[1]Laboratory of Developmental Neurobiology, National Heart Lung and Blood Institute, NIH, Bethesda, MD.

[2]Department of Neuroscience, Interdisciplinary Program in Neuroscience, Georgetown University, Washington, DC.

[#] These authors contributed equally to this work.

*Correspondence:

Herbert M. Geller (E-mail: gellerh@nhlbi.nih.gov)

Laboratory of Developmental Neurobiology

Cell Biology and Physiology Center

National Heart, Lung, and Blood Institute

National Institutes of Health

9000 Rockville Pike

Bethesda, MD 20892

Phone: 301-451-9440

FAX: 301-594-8133

Running Title: PLPPR1 increases cell adhesion

SUMMARY

PLPPR1 alters cytoskeletal dynamics to influence cell adhesion by modulating RhoA and Rac1

activation through its association with RhoGDI, enabling PLPPR1 to impede the inhibitory

activity of CSPGs and LPA.

ABSTRACT

Phospholipid Phosphatase-Related Protein Type 1 (PLPPR1) is a member of a family of lipid

phosphatase related proteins, integral membrane proteins characterized by six transmembrane

domains. This family of proteins is enriched in the brain and recent data indicate potential

pleiotropic functions in several different contexts. An inherent ability of this family of proteins is

to induce membrane protrusions, and we have previously reported that members of this family

may act in concert. However, the function of PLPPR1 is not yet understood. Here, we report that

exogenous expression of PLPPR1 increases cell adhesion to the ECM substrate by altering

cytoskeletal dynamics and modulating RhoA and Rac1 activity through association with

RhoGDI, enabling PLPPR1 to overcome the inhibitory activity of CSPGs and LPA. Together,

these results establish a novel signaling pathway for PLPPR1 protein.

Abbreviations: PLPPR (Phospholipid Phosphatase-Related Protein), PRG (Plasticity-Related

Gene), FA (Focal Adhesions), CSPG (Chondroitin Sulfate Proteoglycan), PLL (Poly-L-Lysine),

LPA (Lysosphosphatidic Acid), LPP (Lipid Phosphate Phosphatase), FAFBSA (Fatty Acid Free

Bovine Serum Albumin), MLC (myosin light chain II), MYPT1 (myosin light chain phosphatase

1), ERM (Ezrin, Radixin and Moesin), GSK (Glycogen Synthase Kinase), GEF (Guanine

Exchange Factor), GAP (GTPase Activating Protein), RhoGDI (Rho Guanine Nucleotide

Dissociation Inhibitor), IRM (Interference Reflection Microscopy), TIRF (Total Internal Reflection

Fluorescence), STED (Stimulated Emission Depletion).

INTRODUCTION

The Phospholipid Phosphatase Related (PLPPR) proteins (PLPPR1 – PLPPR5) are

highly enriched in the brain and their expression is regulated, with the highest expression levels

during development (Wang and Molnár, 2005). This family of proteins were originally named

"Plasticity-Related Genes", due to the upregulation of PLPPR4, the first protein of this family to

be discovered, in sprouting axons following hippocampal deafferentation (Brauer et al., 2003).

More recent studies have shown that expression of PLPPR1 mRNA correlates with sprouting

corticospinal axons after injury (Fink et al., 2017) and neuronal remodeling in the hippocampus

after kainaic acid treatment (Savaskan et al., 2004). Furthermore, decreased PLPPR1 mRNA

has been associated with dysregulated neuronal migration (Khalaf-Nazzal et al., 2017; Pfurr et

al., 2017), suggesting a role for these proteins in neuronal migration as well as neurogenesis

and axon growth after injury.

We and others have previously shown that overexpression of PLPPR1 induces actin-rich

membrane protrusions in many different cell types (Broggini et al., 2016; Sigal et al., 2007;

Velmans et al., 2013; Yu et al., 2015). Overexpression of PLPPR5, the closest relative of

PLPPR1 in this family, produces a similar phenotype (Broggini et al., 2010). The members of

this family interact with each other and may function as a heteromeric complex in causing these

phenotypic changes (Yu et al., 2015), suggesting that these proteins may alter cytoskeletal

dynamics to induce membrane protrusions. However, the mechanism by which the PLPPR1

protein mediates this morphological change is yet unknown.

Here, we report that PLPPR1 overexpression leads to increased cell adhesion due to

changes in cytoskeletal dynamics. This increased adhesion results in decreased cell migration

as well as the ability of cells overexpressing PLPPR1 to overcome the inhibitory activity of

lysophosphatidic acid (LPA) and chondroitin sulfate proteoglycans (CSPGs). These changes in

cell adhesion are due to alterations in the RhoA-ROCK signaling pathway mediated by PLPPR1

through association with RhoGDI. Together, these results establish a novel signaling pathway

for the PLPPR family of proteins.

RESULTS

PLPPR1 reduces cell migration and increases cell adhesion

Overexpression of PLPPR1 induces actin-rich membrane protrusions in several different

cell types (Broggini et al., 2016; Savaskan et al., 2004; Sigal et al., 2007; Velmans et al., 2013;

Yu et al., 2015) implying a change in cytoskeletal dynamics. Many cellular processes including

cell migration and cell adhesion require dynamic changes to the cytoskeleton. Therefore, using

live cell imaging, we traced the migratory trajectories of individual Neuro2a cells expressing

either EGFP or EGFP-PLPPR1 on a uniform fibronectin substrate at a concentration determined

to be permissive for cell migration (Fig. 1A). We found that the average speed (Fig. 1B) and

average distance travelled by migrating cells overexpressing PLPPR1 (Fig. 1C) was significantly

reduced compared to EGFP-transfected cells.

Cell adhesion is a major determinant for cell migration and tissue morphogenesis. To

determine if PLPPR1 had any effect on cell adhesion, we performed a time-dependent cell

adhesion assay on Neuro2a cells expressing either EGFP or EGFP-PLPPR1. Our results show

that overexpression of PLPPR1 increased cell adhesion on a fibronectin substrate (Fig. 2A, B,

Fig. S1).

The PLPPR family share high sequence homology with the exception of their C-terminal

domain. Additionally, since the C-terminus of PLPPR1 is unique and essential to its ability to

induce membrane protrusions (Broggini et al., 2016; Yu et al., 2015), we questioned if the C-

terminal domain is essential to its ability to resist detachment from the fibronectin substrate.

Therefore, we deleted the 43 amino acid residues constituting the C-terminus of PLPPR1 and

expressed EGFP-PLPPR1ΔC43 in Neuro2a cells. We found that EGFP-PLPPR1ΔC43 also

resisted detachment to the fibronectin substrate after extensive washing, indicating that the

ability of PLPPR1 to increase cell adhesion is independent of its C-terminal domain (Fig. 2A, B).

Next, to examine the degree of attachment in cells overexpressing PLPPR1, we used

Interference Reflection Microscopy (IRM) imaging to observe and quantify events at the

interface between the cell surface and the fibronectin substrate. With IRM, the very fine cell

contours and the degree of contact cells exert on the fibronectin substrate can be visualized and

measured. Cell membranes in close contact with the substrate produce more interference and

darker pixels while membranes further from the substrate have less interference and hence,

lighter pixels (Fig. 2C). IRM imaging showed that the average area of attachment to the

fibronectin substrate was significantly greater in cells overexpressing PLPPR1 suggesting a

stronger adherence compared to EGFP-transfected cells (Fig. 2D). This confirms our results

that show increased resistance to detachment by trypsin/EDTA in cells overexpressing PLPPR1

(Fig. S1).

Our previous studies have demonstrated that while cytochalasin treatment eliminated

most of the membrane protrusions induced by overexpression of PLPPR1, some still remained,

suggesting that not all the protrusions induced by PLPPR1 contained actin filaments (Yu et al.,

2015). Using live-cell IRM imaging, we observed that some of these protrusions induced by

PLPPR1 remained strongly adhered to the ECM substrate, leaving a trail of fibrous structures as

the cell migrates (Video 1, Fig. 2E, F). To confirm that these trailing fibers are devoid of actin

filaments, we co-expressed mApple-FTR-940 (F-tractin) with either EGFP or EGFP-PLPPR1 in

Neuro2a cells. Confocal images show that these "trailing fibers" indeed lack actin filaments (Fig.

2G). The presence of such "trailing fibers", devoid of cytoskeletal components, have been

attributed to active migration in cancer cells (DePasquale, 1998; Haemmerli and Strauli, 1981)

and a stronger adherence of the cells to their substrate. Together, these results show that

overexpression of PLPPR1 decreases cell migration due to increased adhesion to the

fibronectin substrate.

PPLRI promotes the destabilization of actin cytoskeleton

The dynamic organization of the actin cytoskeleton causes changes in cell shape and

adhesion strength. This generates sufficient tension to create the pushing and pulling forces that

enable cell migration (Gardel et al., 2010; van Helvert et al., 2018). Confocal imaging

demonstrated that cells overexpressing PLPPR1 lack the network of organized linear stress

fibers observed in the cell periphery of EGFP-transfected cells (Fig. 3A). Since actin

polymerization and depolymerization are required for cell migration (Mogilner and Oster, 1996;

Ridley et al., 2003), actin turnover was determined by FRAP measurements in cells co-

expressing F-tractin with either EGFP or EGFP-PLPPR1 (Fig. 3B). The leading edge of the cell

was photo-bleached and fluorescence recovery was recorded. Results show a decrease in the

rate of fluorescence recovery of F-tractin in cells overexpressing PLPPR1 (Fig. 3C) with a short

half-time compared to EGFP-transfected cells (Fig. 3D). The mobile fraction of monomeric actin

was also decreased in cells overexpressing PLPPR1 compared to EGFP-transfected cells (Fig.

3E). Together, these results show a decrease in the rate of actin polymerization, suggesting

destabilized actin in cells overexpressing PLPPR1.

PLPPR1 increases nascent focal adhesions

The increased cell adhesion observed in cells overexpressing PLPPR1 led us to

evaluate changes in focal adhesion (FA) dynamics. Paxillin is expressed at focal adhesion sites,

therefore, cells expressing either EGFP or EGFP-PLPPR1 were plated on a fibronectin

substrate and immunostained for paxillin and F-actin (Fig. 4A). TIRF imaging revealed that the

distinct ventral stress fibers in the well-defined lamellipodial structure observed in EGFP-

transfected cells were absent in cells overexpressing PLPPR1 (white asterisks, Fig. 4A).

Additionally, F-actin containing stress fibers were sparse in cells overexpressing PLPPR1

compared to EGFP-transfected cells (Fig. 4A).

Focal adhesions (FA) are mechanosensitive and require the recruitment of several

proteins in order to change their composition and mature (Pasapera et al., 2010). This tension-

mediated maturation involves the recruitment of paxillin to nascent focal adhesion complexes

that are initially small in size, but then mature into focal adhesions as the size of the focal

complexes increase and are stabilized (Pasapera et al., 2010; Zaidel-Bar et al., 2003).

Measurement of the size of individual FA revealed more nascent paxillin in cells overexpressing

PLPPR1, compared to EGFP-transfected cells, suggesting reduced FA turnover (Fig. 4B). Live

cell imaging confirmed that the focal adhesion complexes in cells co-expressing EGFP-PLPPR1

and mCherry-paxillin remained nascent compared to the mature FA observed in EGFP-

transfected cells (Video 2). Furthermore, the average fluorescence intensity of paxillin-positive

FA was significantly increased in cells overexpressing PLPPR1 compared to EGFP-transfected

cells (Fig. 4C).

Phosphorylation of paxillin on Y31 is crucial for regulation of paxillin turnover and focal

adhesion maturation (Zaidel-Bar et al., 2007). Immunoblotting of cell lysates from Neuro2a cells

expressing either EGFP or EGFP-PLPPR1 revealed an increase in phosphorylation of paxillin in

cells overexpressing PLPPR1 (Fig. 4D, E). Paxillin phosphorylation at Y31 is mediated by FAK

(Bellis et al., 1995; Schaller and Parsons, 1995), therefore, we evaluated phosphorylation levels

of FAK. Phosphorylation of FAK at Y576 in cells overexpressing PLPPR1 was increased

compared to EGFP-transfected cells (Fig. 4D, F). Altogether, these results demonstrate reduced

stability and size of focal adhesions, due to increased phosphorylation of paxillin and FAK in

cells overexpressing PLPPR1.

PLPPR1 overcomes CSPG inhibition of neurite outgrowth.

Overexpression of PLPPR1 protein can overcome the inhibitory activity of myelin as well

as LPA-induced axon collapse and neurite retraction (Broggini et al., 2016). Furthermore,

PLPPR1 has been demonstrated to promote axon regeneration after spinal cord injury in mice

(Fink et al., 2017). Chondroitin sulfate proteoglycans are universally upregulated in the glial scar

following injury and are potent negative regulators of axon growth (Davies et al., 1999; Hynds

and Snow, 1999; Schwab et al., 1993). Therefore, we asked if overexpression of PLPPR1 could

overcome the inhibitory activity of CSPGs. Hippocampal neurons were transfected with EGFP

or EGFP-PLPPR1 and seeded on glass coverslips coated with either poly-L-lysine (PLL) or PLL

and CSPGs. Neurite outgrowth was analyzed by measuring both total neurite length and longest

neurite (Fig. 5A). Total neurite length of EGFP and EGFP-PLPPR1 transfected neurons plated

on PLL alone were 86.1 ± 6.5µm and 90.4 ± 6.1µm respectively. However, in the presence of

CSPGs, total neurite length was markedly decreased in EGFP-transfected neurons, 53.8 ±

4.4µm, p<0.001, (Fig. 5B). In comparison, total neurite length of neurons overexpressing

PLPPR1 exposed to CSPGs was not significantly different from neurons plated on PLL alone,

94.8 ± 6.0µm. Similarly, we observed a decrease in the length of the longest neurite of EGFP-

transfected neurons, while this inhibition was not observed in neurons overexpressing PLPPR1

(Fig. 5C). These results demonstrate that overexpression of PLPPR1 impedes the inhibitory

activity of CSPGs on neurite outgrowth.

Modulation of axon growth by PLPPR1 has been reported to be dependent on its C-

terminal domain (Broggini et al., 2016). Therefore, we expressed EGFP-PLPPR1ΔC43 in

Neuro2a cells and assessed its ability to overcome CSPG inhibition of neurite outgrowth (Fig.

S2A). Our results show no difference in total neurite length or in the longest neurite between

neurons expressing EGFP or EGFP-PLPPR1 on PLL alone. However, neurons expressing

PLPPR1ΔC43 had shorter neurites (Fig. S2B). This difference suggests that overexpression of

PLPPR1ΔC43, by itself, may interfere with neurite outgrowth, probably due to improper protein

localization as most of the truncated PLPPR1 protein tends to be retained in the endoplasmic

reticulum (Yu et al., 2015). In contrast, while EGFP-transfected neurons exhibited significant

decrease in total neurite length and longest neurite when plated onto CSPGs, the total neurite

length and the longest neurite of neurons expressing full-length PLPPR1 and PLPPR1ΔC43

remained unchanged suggesting that the ability to overcome CSPG inhibition is not dependent

on the C-terminus of PLPPR1 (Fig. S2B, C).

PLPPR1 attenuates LPA-induced neurite retraction.

LPA induces neurite retraction and cell rounding in several different cell types (Jalink et

al., 1993; Kranenburg et al., 1999; Tigyi et al., 1996). To confirm this in Neuro2a cells, serum

starved cells were exposed to either fatty acid free bovine serum albumin (FAFBSA) or LPA.

Live cell imaging revealed significant neurite retraction and cell rounding in cells exposed to

LPA compared to FAFBSA (Fig. S3A, Video 3A, B). Next, the effects of PLPPR1

overexpression on neurite retraction induced by LPA was examined. Neuro2a cells transfected

with EGFP or EGFP-PLPPR1 were exposed to either FAFBSA or LPA. 59% of cells expressing

EGFP underwent cell rounding and neurite retraction in response to LPA compared to 42% of

cells expressing EGFP-PLPPR1, $p < 0.05$, (Fig. 5D, E), demonstrating the ability of PLPPR1 to

attenuate LPA activity. Additionally, expression of PLPPR1ΔC43 resulted in a similar attenuated

response to LPA as full-length PLPPR1 (Fig. S3B, C), suggesting that the ability to impede LPA-

induced neurite retraction is not dependent on the C-terminus of PLPPR1.

PLPPR1 modulates RhoA activation

The RhoA signaling pathway is common to many molecules inhibitory to axon growth,

including CSPGs, myelin and LPA (Alabed et al., 2006; Jain et al., 2004; Kranenburg et al.,

1999; Monnier et al., 2003). Additionally, RhoA activation by LPA was reduced in cells

overexpressing PLPPR5, the closest relative to PLPPR1 (Broggini et al., 2010). Therefore, we

asked if PLPPR1 could similarly affect RhoA activation by LPA. Serum starved Neuro2a cells

expressing EGFP or EGFP-PLPPR1 were exposed to either FAFBSA, serum or LPA. Active

RhoA bound to GTP was immunoprecipitated using rhotekin-RBD beads and then subjected to

immunoblot analysis. While cells expressing EGFP had increased RhoA activation in response

to either serum or LPA, there was no change in RhoA activation in cells overexpressing

PLPPR1 (Fig. 5F, G).

PLPPR1 decreases LPA-induced phosphorylation of ROCK targets.

Rho-associated kinase (Rho-kinase/ROCK/ROK) is a major RhoA effector and is

implicated in the regulation of actinomyosin dynamics and inhibition of neurite outgrowth

(Amano et al., 1997; Lingor et al., 2007; Monnier et al., 2003). ROCK mediates its effect by

phosphorylating several substrates including cytoskeletal proteins, MLC, MYPT1 and the ERM

proteins (Kaneko-Kawano et al., 2012; Matsui et al., 1998; Riddick et al., 2008; Sutherland et

al., 2016; Yoneda et al., 2005). We evaluated the modulatory effect of PLPPR1 on the RhoA-

ROCK pathway by assessing the phosphorylation levels of several ROCK substrates in

response to LPA treatment. First, we measured the phosphorylation levels of MLC. Serum

starved Neuro2a cells expressing EGFP or EGFP-PLPPR1, were exposed to either FAFBSA,

serum or LPA and immunostained for phosphorylated MLC. Confocal imaging showed basal

levels of phosphorylated MLC with a diffuse cytoplasmic distribution after FAFBSA treatment

(Fig. 6A – C). However, in response to LPA, phosphorylation of MLC was increased and

localized to the plasma membrane in EGFP-transfected cells while no change in

phosphorylation levels of MLC was observed and the diffuse cytosolic localization was retained

in cells overexpressing PLPPR1 (Fig. 6A – C). GFP immunoblot of cell lysates from Neuro2a

cells expressing EGFP or EGFP-PLPPR1 and treated with either FAFBSA, serum or LPA show

that expression levels for both EGFP and EGFP-PLPPR1 does not change between treatment

conditions, FAFBSA, serum or LPA (Fig. 6D). Immunoblot analysis confirms the increase in

phosphorylation of MLC at S19 in EGFP-transfected cells, with no change in the phosphorylated

levels of MLC in cells overexpressing PLPPR1 (Fig. 6E, F). Similarly, phosphorylation of MYPT1

at T853 was increased in response to LPA treatment in EGFP-transfected cells while MYPT1

phosphorylation levels remained unchanged in cells overexpressing PLPPR1 (Fig. 6G, H).

260   Next, we evaluated the effect of PLPPR1 on LPA stimulated phosphorylation of the ERM

261   proteins. EGFP-transfected cells exhibited an increase in ERM phosphorylation in response to

262   LPA while phosphorylation levels of the ERM proteins in cells overexpressing PLPPR1

263   remained unchanged (Fig. 6I, J).

264   Activation of the ERM proteins results in their translocation to the plasma membrane

(Kim et al., 2011; Shaw et al., 1998). Therefore, we immunostained for phosphorylated ERM

proteins in cells expressing either EGFP or EGFP-PLPPR1. After FAFBSA treatment, cells

displayed low levels of phosphorylated ERM with a diffuse cytoplasmic distribution (Fig. S4).

However, exposure to LPA induced the translocation of phosphorylated ERM to the plasma

membrane in cells expressing EGFP while ERM localization remained unchanged in cells

overexpressing PLPPR1 (Fig. S4). Together, these results demonstrate that overexpression of

PLPPR1 attenuates phosphorylation of MLC, MYPT1 and the ERM proteins after LPA

stimulation.

273   Since LPA is known to induce phosphorylation of Akt, GSK3ß and Tau through

activation of the PI3K-PKB/Akt and PLCγ-PKC pathways respectively (Baudhuin et al., 2002;

Fang et al., 2002; Sayas et al., 2006), we asked if the effects of PLPPR1 overexpression extend

to these pathways. Immunoblot analysis of phosphorylation levels of GSK3ß at Y216 and S9, of

Tau at S396 and of Akt at S473 in transfected Neuro2a cells after LPA treatment, showed no

significant change in cells overexpressing PLPPR1 compared to EGFP-transfected cells (Fig.

S5).

280   To determine if overexpression of PLPPR1 had any effect on the phosphorylation levels

of MLC, ERM, MYPT1, Akt, GSK3ß or Tau without LPA stimulation, we performed an

immunoblot analysis of cell lysates from Neuro2a cells transfected with either EGFP or EGFP-

PLPPR1. Results show no significant changes in phosphorylation levels of these ROCK target

proteins in cells overexpressing PLPPR1 compared to EGFP-transfected cells (Fig. S6).

Altogether, these results demonstrate that PLPPR1 specifically attenuates activation of the

RhoA-ROCK signaling pathway in response to stimulation.

PLPPR1 modulates RhoA and Rac1 activation via RhoGDI interaction

Previously, we identified RhoGDI as a possible binding partner for PLPPR1 (Yu et al.,

2013). By direct interaction with the Rho GTPases, RhoGDI prevents their translocation from

the cytosol to the plasma membrane thereby preventing the nucleotide exchange of GDP for

GTP (Boulter et al., 2010; Dovas and Couchman, 2005; Sabbatini and Williams, 2013). To

confirm the interaction between RhoGDI and PLPPR1, we immunoprecipitated EGFP-PLPPR1

using GFP magnetic/agarose beads and immunoblotted for RhoGDI. We also

immunoprecipitated EGFP-PLPPR1∆C43, to determine if this putative interaction is dependent

on the C-terminal domain. Our results show that both EGFP-PLPPR1 and EGFP-PLPPR1∆C43

co-immunoprecipitated with RhoGDI, confirming that PLPPR1 associates with RhoGDI and this

interaction is independent of the C-terminus (Fig. 7A).

Inactive RhoA (bound to GDP) is tightly sequestered by RhoGDI in the cytosol

(Ellerbroek et al., 2003; Garcia-Mata et al., 2011; Michaelson et al., 2001; Sabbatini and

Williams, 2013; Takahashi et al., 1997; Yamashita and Tohyama, 2003). LPA stimulation

results in the displacement of RhoA from RhoGDI. This allows RhoA to disengage GDP, bind

GTP and become activated (Kranenburg et al., 1997; Yoshioka et al., 1999). Therefore, to

determine how overexpression of PLPPR1 would affect the RhoGDI – RhoA interaction after

LPA treatment, Neuro2a cells transfected with EGFP or EGFP-PLPPR1 were exposed to either

FAFBSA, serum or LPA. RhoGDI was immunoprecipitated and immunoblot analysis of RhoA

was performed. We found that the interaction between RhoA and RhoGDI was disrupted in

EGFP-transfected cells after treatment with either serum or LPA, as expected. However, the

interaction of RhoGDI for RhoA was maintained in cells overexpressing PLPPR1 even after

serum or LPA treatment (Fig. 7B). Immunoblot analysis of GFP expression showed that protein

expression levels did not change between treatment conditions (Fig. 7B).

Like RhoA, Rac1 activation is regulated by RhoGDI (Boulter et al., 2010;

DerMardirossian et al., 2004). We therefore assessed the effects of PLPPR1 overexpression on

the affinity of RhoGDI for Rac1 after LPA treatment. Neuro2a cells transfected with EGFP or

EGFP-PLPPR1 were exposed to either FAFBSA, serum or LPA. RhoGDI was

immunoprecipitated and immunoblotting was performed to detect Rac1 levels. Our results show

that, like RhoA, the interaction between RhoGDI and Rac1 was disrupted by LPA treatment in

cells expressing EGFP, however, this interaction was maintained in cells overexpressing

PLPPR1 (Fig. 7B). Altogether, these results demonstrate that PLPPR1 modulates LPA induced

activation of RhoA and Rac1, via its association with RhoGDI.

Co-expression of PLPPR1 and CA Rac1 reverts PLPPR1 induced morphological change

Rac1 activation is essential to lamellipodial formation (Ehrlich et al., 2002; Kozma et al.,

1997; Somanath and Byzova, 2009). Typical membrane ruffling lamellipodia were found absent

in cells overexpressing PLPPR1 (Fig. 4A). Therefore, given the possibility that PLPPR1 may

regulate Rac1 activity, we examined the morphology of cells expressing dominant negative (DN)

Rac1 T17N, constitutively active (CA) Rac1 G12V and cells co-expressing PLPPR1 with CA

Rac1. We observed that expression of DN Rac1 caused a morphological change, similar to the

"trailing fibers" observed in cells overexpressing PLPPR1 (Fig. 8A). Interestingly, co-expression

of CA Rac1 with PLPPR1 was able to revert the morphological change induced by PLPPR1

(Fig. 8A).

To confirm the effect of PLPPR1 on Rac1 activation, DN Rac1 and CA Rac1 were

expressed in Neuro2a cells and active Rac1 bound to GTP was immunoprecipitated using PAK-

PBD beads. Basal levels of Rac1-GTP were near zero in Neuro2a cells transfected with EGFP,

as well as in cells overexpressing PLPPR1 and DN Rac1 (Fig. 8B, C). As expected, there were

high levels of active Rac1 in cells expressing CA Rac1, however, co-expression of CA Rac1

with PLPPR1 significantly reduced active Rac1 (Fig. 8B, C), confirming that PLPPR1 modulates

Rac1 activation.

Arp2/3 is an actin nucleating and crosslinking complex that acts downstream of Rac1

and is essential to actin assembly and lamellipodia formation (Lai et al., 2008; Machesky and

Hall, 1997; Machesky and Insall, 1998; Wu et al., 2012). Therefore, we examined Arp2/3 in cells

overexpressing PLPPR1 using STED microscopy. Results show that while Arp2/3 was localized

to the actin-rich lamellipodia and exhibited a more diffuse cytoplasmic distribution in cells

transfected with EGFP, cells overexpressing PLPPR1 displayed a compact aggregation of

Arp2/3 (Fig. 8D), suggesting a disruption in Arp2/3 localization. Together, these results

demonstrate a regulatory function for PLPPR1 in Rac1 activation.

DISCUSSION

The PLPPR family of proteins were initially identified as "Plasticity-Related Genes" due

to their upregulation in sprouting axons (Brauer et al., 2003). PLPPR1 expression levels

correlate with axon growth and remodeling after kainic acid treatment in the hippocampus

(Savaskan et al., 2004). Additionally, PLPPR1 was identified in a screen for molecules that

promote regeneration after spinal cord injury (Fink et al., 2017). More recently, the PLPPR

family of proteins have been implicated in the regulation of neurogenesis and neuronal

migration (Khalaf-Nazzal et al., 2017; Lyons et al., 2010; Pfurr et al., 2017). Exogenous

expression studies of PLPPR1 in cultured cells demonstrate that the major phenotype is the

induction of filopodia in many different cell types (Broggini et al., 2016; Velmans et al., 2013; Yu

et al., 2015). The mechanism of action by which PLPPR1 induces these morphological changes

has been a subject of intense speculation.

Here, we demonstrate that PLPPR1 modulates the RhoA-ROCK signaling pathway. We

show that through association with RhoGDI, PLPPR1 may prevent GTP loading by stabilizing

the interaction between RhoGDI and GDP bound RhoA and Rac1. This stabilization results in

reduced RhoA and Rac1 activity, leading to alterations in cytoskeletal dynamics. This explains

the effect of PLPPR1 on cell adhesion and consequently cell migration enabling PLPPR1 to

overcome the inhibitory activity of CSPGs and LPA on neurite outgrowth (Fig. 9).

The PLPPR family of proteins share sequence and structural homology with the lipid

phosphate phosphatase family (LPP), and are characterized by six transmembrane domains

with intracellular N-terminal and C-terminal domains (Brauer and Nitsch, 2008; Strauss and

Brauer, 2013). This homology to lipid phosphatases and the discovery that overexpression of

PLPPR4 could attenuate LPA-induced axon collapse (Brauer et al., 2003), provided support for

the hypothesis that PLPPR signaling was through extracellular lipid phosphatase activity.

However, the PLPPR proteins lack key amino acid residues critical to the phosphatase activity

of the LPP protein family and are not known to exhibit any phosphatase activity (Strauss and

Brauer, 2013), thus, other signaling pathways have been suggested.

In addition to attenuating LPA-induced axon collapse, overexpression of PLPPR1 can

reduce the inhibitory response to myelin (Broggini et al., 2016), and here, we show a similar

response to CSPGs. Each of these guidance cues are thought to signal through activation of the

RhoA GTPase (Alabed et al., 2006; Ohtake et al., 2016) and reducing RhoA activity with C3

exoenzyme attenuates their inhibitory actions, as does inhibition of the downstream effector,

Rho kinase, ROCK (Dergham et al., 2002; Fournier et al., 2003). Here, we demonstrate that

overexpression of PLPPR1 reduced RhoA activation in response to LPA treatment. This is

similar to the effect of its close homolog, PLPPR5, on RhoA activation by LPA (Broggini et al.,

2010). Furthermore, we confirmed the modulatory effect of PLPPR1 on RhoA activation of

ROCK targets, MLC, MYPT1 and the ERM proteins. A previous study also implicated PIP5K, a

protein kinase downstream of RhoA, as being altered by PLPPR1 expression (Broggini et al.,

2016).

RhoA activation is controlled by regulatory proteins such as the GEFs, that catalyze the

exchange of GDP for GTP, and the GDIs, that sequester RhoA in the cytosol, preventing its

translocation to the plasma membrane and thus, its activation (Bokoch et al., 1994; Dovas and

Couchman, 2005; Hodge and Ridley, 2016). We identified RhoGDI as a possible binding target

for PLPPR1 in a proteomic screen for interacting proteins (Yu et al., 2013), and we have

confirmed this interaction by co-immunoprecipitating PLPPR1 with RhoGDI. Without stimulation,

RhoA-GDP is tightly bound to RhoGDI (Boulter et al., 2010; Hodge and Ridley, 2016;

Tkachenko et al., 2011). Upon LPA treatment, this interaction is disrupted, permitting RhoA to

translocate to the plasma membrane and become activated. We have demonstrated here that

overexpression of PLPPR1 maintains the interaction of RhoA with RhoGDI after LPA treatment

suggesting that PLPPR1 may play a role in stabilizing the RhoA-RhoGDI complex, thereby,

preventing the RhoA GDP exchange for GTP and its subsequent activation.

RhoGDI is known to regulate the activity of the small GTPases, Rac1 and Cdc42.

Expression of DN Rac1 prevents GTP loading, resulting in a failure to form proper lamellipodia

(Ehrlich et al., 2002; McCarty et al., 2005). Interestingly, we show that expression of DN Rac1

exhibits a similar morphological change as overexpression of PLPPR1 in Neuro2a cells. Our

results also show that co-expression of PLPPR1 with CA Rac1 reverts this morphological

change in Neuro2a cells. This is consistent with our results that demonstrate a stable Rac1-

RhoGDI complex in cells overexpressing PLPPR1, suggesting a modulatory effect on Rac1

activity.

The formation of membrane protrusions by exogenous expression of PLPPR1 has been

shown to be independent of Cdc42, Arp2/3 and the EnA/VASP proteins, suggesting that the

PLPPR1 has no modulatory effect on Cdc42 (Sigal et al., 2007). Additionally, co-expression of

PLPPR1 with dominant negative Cdc42 or with WASP-CRIB, which recruits and activates

Arp2/3, did not reverse the morphological change induced by PLPPR1 overexpression (Sigal et

al., 2007). It is noteworthy that the nucleation of actin filaments by Arp2/3 complex is also driven

413 by WAVE, downstream of Rac1 activation (Chen et al., 2017; Lai et al., 2008). Therefore, while

414 Arp2/3 may not be required to induce membrane protrusions, it still remains that localization of

415 Arp2/3 is disrupted in cells overexpressing PLPPR1 and this may be a direct consequence of

416 altered Rac1 activity.

417  The PLPPR proteins are similar in their N-terminal domains, with significant divergence

418 in their intracellular C-terminal regions (Brauer and Nitsch, 2008). The C-terminal domain of

419 PLPPR1 is important for the induction of membrane protrusions (Broggini et al., 2016; Yu et al.,

420 2015), likely due to the inability of PLPPR1 lacking its C-terminus, to be transported to the cell

421 membrane (Yu et al., 2015). This domain is also the site of the change in phosphorylation by

422 CSPGs (Yu et al., 2013), and has been reported to modulate Ras activity by interacting with

423 RasGRF1 (Broggini et al., 2016). However, our data show that the PLPPR1 C-terminal domain

424 is dispensable for its ability to overcome the inhibitory activity of CSPGs and LPA and is not

425 necessary for its association with RhoGDI, suggesting that while the C-terminal domain of

426 PLPPR1 may be important for the induction of membrane protrusions, it is not essential to its

427 modulation of RhoA and Rac1 activity. Interestingly, the C-terminal of PLPPR5, the closest

428 family member to PLPPR1, has also been reported to be essential for formation of protrusions

429 (Broggini et al., 2010). Whether there is a similar interaction of PLPPR5 with RhoGDI remains

430 an open question.

431  PLPPR1 overexpression ultimately results in increased adhesion and decreased

432 migration of cells on a fibronectin substrate. Cell migration is an integrated process that involves

433 the spatio-temporal coordination of actin cytoskeletal dynamics with focal adhesion turnover

434 (Gardel et al., 2010; Le Clainche and Carlier, 2008; Webb et al., 2002). Cellular adhesion to

435 fibronectin is mediated through the engagement and activation of integrins, which then promote

436 the establishment of focal adhesion complexes that are also regulated by the activity of the

437 Rac1 and RhoA GTPases (Nobes and Hall, 1995; Rottner et al., 1999; Sinnett-Smith et al.,

438 2001). Mechanical tension promotes the growth and maturation of focal adhesions that is driven

by RhoA mediated myosin II activity (Pasapera et al., 2010) and this is required for the

coordinated propulsive force that enables cell migration (Gardel et al., 2010; Lauffenburger and

Horwitz, 1996). Several studies have demonstrated that inhibition of Rho-ROCK activity

prevents the maturation of focal adhesions (Burridge and Guilluy, 2016; Sawada et al., 2002;

Sinnett-Smith et al., 2001). Therefore, it is conceivable that the reduced RhoA activity and the

nascent focal adhesion complexes that persists in cells overexpressing PLPPR1 may account

for the decrease in cell migration. This is consistent with the increase in phosphorylation of

paxillin and FAK observed in these cells, given that FAK mediated phosphorylation of paxillin

plays a role in the assembly, turnover and stability of focal adhesions complexes (Lawson et al.,

2012; Pasapera et al., 2010; Turner, 2000; Webb et al., 2004; Zaidel-Bar et al., 2007). It is

essential to note that while increased phosphorylation of paxillin and FAK is observed in cells

overexpressing PLPPR1, this does not imply direct kinase activity on paxillin and FAK by

PLPPR1 but suggests an indirect effect on the phosphorylation of paxillin and FAK.

Another member of the PLPPR family, PLPPR4, mediates cell adhesion through

activation of ß1-integrins (Liu et al., 2016). Through the calmodulin-binding domain (CBD) in its

long cytoplasmic C-terminal tail consisting of ~400 a min o acid residues that is unique to this

PLPPR family member alone, PLPPR4 interacts with protein phosphatase 2A (PP2A) to

organize the assembly of an integrin adhesome consisting of talin, paxillin and src, thereby

stimulating activation of ß1-integrin (Liu et al., 2016). While PLPPR1 lacks the key motifs in its

C-terminal domain that is characteristic of PLPPR4, and we have demonstrated that the

increased cell adhesion observed in PLPPR1 is independent of its C-terminal domain, it still

remains that PLPPR1 may mediate activation of ß1-integrins through other means to increase

cell adhesion. Additionally, there is evidence that PLPPR1 may associate with ß1-integrin (Yu et

al., 2013), however, this putative interaction is yet to be confirmed.

Mesenchymal type cell migration is characterized by increased focal adhesion sites at

the leading edge of the cell and an adhesive "pulling" during migration, causing parts of the

membrane protrusions to remain adhered to the ECM as the cell migrates away (Friedl and

Wolf, 2010; Schmidt and Friedl, 2010). We found that cells transfected with PLPPR1 did not

disengage from the substrate during migration, leaving long plasma membrane extensions that

we have termed "trailing fibers". The presence of these "trailing fibers" suggest an inability of the

cells to detach from the substrate due to increased adhesion.

While overexpression of PLPPR1 increases cell adhesion, biological functions, such as

alteration of neurogenesis (Pfurr et al., 2017), neuronal migration (Khalaf-Nazzal et al., 2017)

and regulation of axonal growth (Fink et al., 2017), have been inferred from experiments using

genetic screening approaches. Determining the role of PLPPR1 would require the ability to

silence PLPPR1 mRNA and confirm the knock-down using protein-detection techniques.

Unfortunately, there are no reliable commercial antibodies to detect this protein, thus,

preventing these studies.

Taken together, our results provide evidence for a novel signaling mechanism of

PLPPR1 – the modulation of RhoA and Rac1 activity by interaction with RhoGDI. As its original

name, Plasticity-Related Gene protein 3, infers, PLPPR1 plays a role in mediating plasticity in

response to extracellular cues, through modulation of the RhoA – ROCK signaling pathway via

its association with RhoGDI. This mechanism serves as a potential explanation for the ability of

PLPPR1 to overcome CSPG and myelin-mediated inhibition of neurite outgrowth to promote

axon regeneration after injury.

MATERIALS AND METHODS

Plasmid construction

DNA constructs: The mApple-FTR-940 (F-tractin) was originally from Michael Davidson's

laboratory (plasmid # 54902, Addgene, Cambridge, MA). pEGFP-Rac1 G12V and pEGFP-Rac1

T17N were both obtained from Clontech (Mountain view, CA). mApple-paxillin was kindly

provided by Clare Waterman (National Institutes of Health, Bethesda, MD). PLPPR1 cDNA was

amplified from a mouse brain cDNA library. Full-length PLPPR1 (amino acids 1–325) and

PLPPR1 lacking the last 43 amino acid residues of the C-terminal region (PLPPR1ΔC43, amino

acids 1–282) were inserted into the pEGFP-C1 vector with the EGFP tag fused to the N-

terminus of PLPPR1 cDNA using the In-Fusion cloning system (Clontech, Mountain View, CA).

A construct expressing C-terminal HA-tagged PLPPR1 (PLPPR1–HA) was kindly provided by

Andrew J. Morris (University of Kentucky, Lexington, KY) (Sigal et al., 2007). The

oligonucleotide primer pairs used were as follows:

PLPPR1 SalIfusC1F: 5'GAATTCTGCAGTCGACGCCACCATGGCTGTAGAAAACAACACTCA3'

PLPPR1 SacIIfusC1R:

5'GTGGATCCCGGGCCCGCGGTCAGGTGACTTCGGTCATGGAGGCG3'

PLPPR1ΔC43aaSacIIfusC1R:

5'GTGGATCCCGGGCCCGCGGTCAGGTTCCTCTGAAGTTATGAA3'

Cell culture and transfection

All animal procedures were performed in accordance with protocols approved by the

Institutional Animal Care and Use Committee (IACUC) at the National Institutes of Health.

Timed pregnant C57BL/6J mice were housed in a pathogen free facility with free access to food

and water under a standard 12h light/dark cycle.

Primary hippocampal neurons were isolated from embryonic day 18 – 20 C57BL/6J

pregnant mice and transfected with 1µg of either EGFP, EGFP-PLPPR1 or EGFP-

PLPPR1ΔC43 plasmid DNA per well using the 96-well shuttle Amaxa Nucleofector™ kit,

program CA-138 (Lonza, Walkersville, MD). Neurons were grown in Neurobasal media

supplemented with antibiotics (penicillin/streptomycin), glutamax, B27 (ThermoFisher Scientific,

Waltham, MA) and 2mM potassium chloride (KCl) for 48h.

Neuro2a cells were maintained in DMEM supplemented with penicillin/streptomycin

antibiotics and 10% fetal bovine serum (FBS). Twenty-four hours after plating, cells were

transfected with either EGFP or EGFP-PLPPR1 or EGFP-PLPPR1ΔC43, using Avalanche

Omni® transfection reagent (EZ Biosystems, College Park, MD).

For CSPG treatment and neurite outgrowth analysis, primary hippocampal neurons were

plated onto glass coverslips coated with either poly-l-lysine (PLL) or a soluble CSPG mix

containing neurocan, phosphacan, versican, and aggrecan isolated from embryonic chicken

brain (CC117 Millipore, Billerica, MA). Neurons were subsequently immunostained for anti-βIII

tubulin (Sigma Aldrich, St. Louis, MO) and imaged using a Nikon Eclipse Microscope with a

60×/1.4 NA oil immersion objective lens. Neurite outgrowth analysis was performed using the

NeuronJ plugin of ImageJ.

All cell culture plates (Corning, Corning NY), 6-well (3506), 12-well (3512), 24-well

(3527), 48-well (3548), MatTek dishes (35mm petri dish with 14mm microwells, No. 1.5, MatTek

Corporation, Ashland, MA) and Lab-Tek™ culture II chamber slides, (155382) (ThermoFisher

Scientific, Waltam, MA) were coated with 10µg/ml fibronectin (Millipore, Billerica, MA), unless

otherwise indicated, and incubated overnight at 4˚C. Excess fibronectin was rinsed off by

washing 3 times with Phosphate buffered saline (PBS).

Immunocytochemistry

Transfected Neuro2a cells were fixed in 4% PFA, 48h after plating and 24h after

transfection. Cells were briefly rinsed in PBS and incubated in blocking buffer (0.3% Triton X-

100 in PBS, supplemented with 10% normal goat serum, NGS) for 1h and then with primary

antibody diluted in 0.3% Triton X-100 in PBS, supplemented with 2.5% NGS. Cells were

washed in PBS and then incubated with secondary antibody diluted in 0.3% Triton X-100 in

PBS, supplemented with 2.5% NGS. See table for dilutions and source of antibodies. Cells were

then washed in PBS and then mounted with DAPI diluted in Fluoromount (Sigma-Aldrich). All

phalloidin (ThermoFisher Scientific, Waltam, MA) staining was performed as previously

described (Yu et al., 2015).

For STED microscopy, cells were mounted in prolong diamond mounting media

(ThermoFisher Scientific, Waltam, MA) and imaged on a Leica SP8 STED 3X/Confocal

Microscope using 100×/1.4 oil immersion objective lens (Leica Microsystems, Buffalo Grove, IL).

All other imaging was performed on a Zeiss 780 LSM confocal microscope using 63×/1.4 oil

immersion objective lens.

Live cell imaging

All live cell imaging was conducted on microscopes equipped with a heated stage at

37˚C and 5% CO_2 on either a Nikon A1R microscope (Nikon Instruments Inc., Melville, NY), a

Zeiss 780 LSM confocal microscope (Carl Zeiss, Thornwood, NY) or an Delta Vision OMX

microscope (GE, Malborough, MA).

Cell migration assay

Neuro2a cells transfected with either EGFP or EGFP-PLPPR1 were plated in Lab-Tek

culture chamber slides coated with fibronectin. Time lapse images were acquired for 10h at the

rate of 1 frame/10 min on a Nikon A1R microscope using a 10×/0.45 NA objective lens. Cell

migration was analyzed by tracking each transfected cell, using the middle of each cell body as

the point of reference. The distance travelled was determined using NIS Elements software

(Nikon Instruments Inc., Melville, NY).

Cell adhesion assay

Neuro2a cells transfected with either EGFP, EGFP-PLPPR1 or EGFP-PLPPR1∆C43

were plated in T75 culture flasks and detached with 5mM EDTA. Cells were replated in a 48-

well plate coated with fibronectin at a density of 2.0×10^5 cells per well. Cells were allowed to

adhere for 1h after plating and then washed briefly with DMEM culture media every 15 min

thereafter. Fluorescence detection was performed on a SpectraMax M3 microplate reader

(Molecular Devices, San Jose, CA). The cell density of adherent transfected cells was

determined by fluorescence using the Softmax Pro7 software (Molecular Devices, San Jose,

CA). Cell adhesion was calculated as the percentage of transfected cells that persists after each

wash, from unwashed transfected cells. Cells were subsequently fixed in 4% PFA, rinsed twice

in PBS and stained with DAPI. Images were acquired on a Nikon A1R microscope using a

10×/0.45 NA oil objective lens.

For the detachment assay, Neuro2a cells were plated on a 12-well plate coated with

fibronectin and transfected with either EGFP or EGFP-PLPPR1, 24 h after plating. After another

24h, cells were briefly washed with DMEM culture media to remove serum and subsequently

treated with Trypsin-EDTA diluted in DMEM (1:10) with increasing time intervals. Cells were

then rinsed with PBS, fixed with 4% PFA and stained with DAPI. 10 × 10 tiled images were

acquired on a Zeiss 780 LSM confocal microscope using a 10×/0.45 NA dry objective lens.

Interference reflection microscopy

Neuro2a cells transfected with either EGFP or EGFP-PLPPR1 were fixed with 4% PFA

for 15 min , 24h post transfection. Images were acquired with a Zeiss 780 LSM confocal

microscope using a 63×/1.4 NA oil immersion objective lens. Single-frame TIFF images were

extracted from TIFF stacks and processed using ImageJ. The area of each cell was traced from

bright field images to obtain an ROI (region of interest). The image was then rescaled to an 8-bit

scale of 0–255 based on pixel intensity values that fell within histogram bins exceeding 5%

of the mean histogram bin height. A band pass filter was applied based on convolution with a

41 × 41pixel kernel. The histogram was then stretched again based on a 5% cut-off of the

mean values. A binary threshold was fixed by selecting pixel intensities above 90 on the 0 – 255

scale. The image was then inverted and the number of pixels above threshold was divided by

the area of the ROI to determine intensity. A color map 'Gem' was used to display higher

intensity pixels as blue and lower intensity as orange which represent stronger and weaker

adhesions respectively. For quantification, the area of each cell was traced from bright field

images to obtain the region of interest. A threshold was applied to the IRM image of the same

cell and the number of pixels above threshold was divided by the ROI to calculate the area of

cell attachment.

To measure the size of focal adhesion (FA) contacts, Neuro2a cells were plated on

MatTek dishes at a density of 2×10^4 cells and transfected with either EGFP or EGFP-PLPPR1.

Cells were fixed and stained with anti-paxillin antibody. Images were acquired using TIRF

microscopy on an inverted TIRF OMX Delta Vision microscope system using a 60×/1.49 NA

TIRF objective lens. Counting of FAs in single cells (n = 10 cells) was done after noise removal

by thresholding and applying a size constraint to FAs using ImageJ software.

Actin turnover analyzed by FRAP

Neuro2a cells, plated on glass-bottom MatTek dishes coated with fibronectin, were co-

transfected with F-tractin (mApple-FTR-940) and either EGFP or EGFP-PLPPR1. FRAP

experiments were conducted on a Nikon A1R microscope using a 60×/1.4 NA oil objective lens.

A ROI was selected from the leading edge of transfected cells and a 564 nm laser at 20%

intensity was used as a pre-bleaching signal for F-tractin. The ROI was subsequently

photobleached by three lasers at 458nm, 564nm and 633nm, all set at 100% intensity. The

post-bleaching acquisition was carried out at 60 frames at a rate of 1 frame/0.6s. The

fluorescence intensity was determined at every time point and automatically generated by NIS

Elements software. Images were corrected for photobleaching and the fluorescence intensity of

the ROI for each frame was normalized to the average initial intensity of pre-bleaching frames

and to the area of the bleached cell. The fluorescence intensity for the pre-bleaching frames

was calculated as the average fluorescence intensity in the first 5 frames, F_{pre}. The value, F_0,

was designated as the average fluorescence intensity of the 8th frame after the photobleaching

phase that occurred in the 6th and 7th frames. Individual plots for each experiment was fitted to a

model function. The fluorescence recovery of F-tractin was best fitted by a single-exponential

function (double exponential function did not significantly improve the quality of the fit) using

Matlab, revealing the presence of mobile actin fractions with different recovery kinetics. The

following model was applied:

$$F(t) = A[1 - exp(-Bt)]$$

Where: A is the percentage of mobile fraction of actin and B is the time constant of fluorescence

recovery. A and B were obtained from the fitted curve. The mean ± SEM was calculated for the

percentage of fluorescence recovery and the time constant for each condition. The final graphs

show the fluorescence recovery, mobile fraction of monomeric actin and $t_{1/2}$ which indicates the

time it took for F-actin polymerization.

LPA treatment

Serum starved Neuro2a cells transfected with either EGFP or EGFP-PLPPR1 were

treated with fatty-acid free bovine serum albumin, FAFBSA, (Sigma Aldrich, St. Louis, MO), or

Lysophosphatidic acid, LPA (Cayman Chemical, Ann Arbor, MI) diluted in FAFBSA. Time lapse

images were acquired on a Nikon A1R microscope with a 20×/0.75 dry objective lens with no

time delay between frames for 20 min . Cell response was calculated as percentage of the total

transfected cell count that retracted their neurites or underwent cell rounding.

Western blot assay

Neuro2a cells seeded at a density of 4×10^5 cells/well in a 6-well culture dish were

transfected with either EGFP or EGFP-PLPPR1. Cells were either serum-starved for 16 – 18h

or maintained in DMEM supplemented with 10% FBS. Serum starved cells were then treated

with either FAFBSA, 10% serum or increasing concentrations of LPA: 4, 8 and 16µM for 2 min .

Media was gently but completely removed from all cells and rinsed once with warmed PBS. Cell

lysates were prepared in 2× SDS cell lysis buffer and clarified by centrifugation. Protein

quantitation was performed using Ionic Detergent Compatibility Reagent for Pierce™ 660nm

Protein Assay Reagent (ThermoFisher Scientific, Waltam, MA) by BCA method. Equal

concentration of proteins (20µg/lane) were separated on SDS-PAGE and transferred onto a

polyvinylidene difluoride (PVDF) membrane (Bio-Rad Laboratories, Inc. Hercules, CA).

Membranes were blocked for 1h in 5% nonfat dried milk in either 0.1% Tween20 in PBS (PBST)

or 0.1% Tween20 in 10mM Tris–HCl pH 7.5 (TBST) or blocked in 5% BSA in TBST. Membranes

were incubated with primary antibodies overnight at 4°C, washed in either PBST or TBST and

subsequently incubated with secondary antibody for 30 min . Antigen–antibody complexes were

detected with KPL LumiGlo chemiluminescent substrate (Seracare, Milford, MA). See Table 1

for source and dilutions for all primary and secondary antibodies used.

Co-immunoprecipitation assay

Neuro2a cells were seeded at a density of 4×10^5 cells/well in a 6-well culture dish and

transfected with either EGFP, EGFP-PLPPR1 or EGFP-PLPPR1ΔC43. Cells were harvested,

washed twice in ice-cold PBS and then resuspended in ice-cold cell lysis buffer (10mM Tris pH

7.5, 150mM NaCl, 0.5mM EDTA, 1% NP-40 supplemented with protease inhibitor cocktail set V

EDTA free (Millipore, Billerica, MA) and phosphatase inhibitors (Thermo Fisher Scientific,

Waltham, MA). Cell lysates were cleared and then incubated with GFP-Trap MA beads as

instructed in ChromoTek GFP-Trap® MA immunoprecipitation kit (ChromoTek Inc., Hauppauge,

NY). Cell lysates were subsequently washed twice in dilution buffer (10mM Tris-HCl pH 7.5,

150mM NaCl, 0.5mM EDTA, 0.5% NP-40 supplemented with protease and phosphatase

inhibitors) and then once in wash buffer (10mM Tris-HCl pH 7.5, 500mM NaCl, 0.5mM EDTA,

0.5% NP-40 supplemented with protease and phosphatase inhibitors). Protein was eluted and

subjected to SDS-PAGE as described above.

To immunoprecipitate RhoGDI, transfected cells were rinsed in warmed PBS, harvested

in cell lysis buffer (20mM Tris pH 7.5, 150mM NaCl, 0.5mM EDTA, 1% NP-40 supplemented

with protease and phosphatase inhibitors) and precleared with rabbit IgG serum and Protein G

agarose beads (Seracare, Milford, MA). Samples were subsequently incubated with primary

antibody overnight at 4°C and then incubated with Protein G agarose beads for 1h at 4°C.

Samples were washed twice in dilution buffer (20mM Tris pH 7.5, 150mM NaCl, 0.5mM EDTA,

0.5% NP-40 supplemented with protease and phosphatase inhibitors) and once in wash buffer

(20mM Tris pH 7.5, 500mM NaCl, 0.5mM EDTA, 0.5% NP-40 supplemented with protease and

phosphatase inhibitors). Protein was eluted and subjected to SDS-PAGE as described above.

RhoA and Rac1 pull-down assay

Serum starved Neuro2a cells transfected with either EGFP or EGFP-PLPPR1 were

treated with either FAFBSA, 10% serum or 16μM LPA. Briefly, cells were rinsed with warmed

PBS and cell lysates prepared in cell lysis buffer as instructed and provided by the RhoA or

Rac1 pull-down instruction manual and kit (Cytoskeleton, Inc. Denver, CO). Protein quantitation

was performed by BCA method and 600μg of protein was incubated with 50μg Rhotekin-RBD

beads for RhoA pull-down or with 10μg of PAK-PBD beads for Rac1 pull-down at 4°C for 1h.

Complexes were washed in wash buffer provided in the pull-down kit and protein was eluted in

2× SDS sample buffer and SDS-PAGE performed as described above.

Statistical analysis

All analyses were performed using GraphPad Prism software version 7 (GraphPad

Software, La Jolla, CA). Data was analyzed by unpaired Student's *t* tests, one-way ANOVA or

two-way ANOVA with Welch's correction or Tukey's post-hoc multiple comparison test to

determine significance. Normality of the distribution of the data was tested with Kolmogorov–

Smirnov normality tests using the column statistics function of GraphPad Software. All tests

were two-tailed with significance indicated as follows: *p<0.05; **p<0.01; ***p<0.001;

****p<0.0001. Unless otherwise specified, values represent the means ± SEM. All experiments

were repeated independently at least three times.

Primary Antibodies	Dilution	Source
Mouse anti-βIII tubulin	1:1000	Sigma Aldrich (SAB4700544)
Chicken anti-GFP	1:2000	Abcam (ab13970)
Rabbit anti-phospho-myosin light chain II (Ser19)	1:1000	Cell Signaling Technology (3671)
Rabbit anti-phospho-myosin light chain phosphatase I (Thr853)	1:1000	Cell Signaling Technology (4563)
Rabbit anti-phospho-GSK3β (Tyr216)	1:1000	Abcam (ab75745)
Rabbit anti-pERM	1:1000	Cell Signaling Technology (3141)
Rabbit anti-phospho-AKT (Ser473)	1:1000	Cell Signaling Technology (9271)
Rabbit anti-phospho-Tau (Ser396)	1:1000	Abcam (ab109390)
Rabbit anti-myosin light chain II	1:1000	Cell Signaling Technology (3672)
Rabbit anti-myosin light chain phosphatase I	1:1000	Cell Signaling Technology (2634)
Rabbit anti-GSK3β	1:1000	Cell Signaling Technology (27C10)
Rabbit anti-ERM	1:1000	Cell Signaling Technology (3142)
Rabbit anti-AKT	1:1000	Cell Signaling Technology (4685)
Rabbit anti-RhoGDI	1:1000	ThermoFisher Scientific (51-1000Z)
Anti-p34-Arc/ARPC2	1:100	Sigma Aldrich (07-227)
Mouse anti-Rac1	1:500	Cytoskeleton, Inc. (ARC03)
Mouse anti-Tau	1:2000	ThermoFisher Scientific (MA5-12808)
Mouse anti-RhoA	1:500	Cytoskeleton, Inc. (ARH03)

Mouse anti-β-actin	1:4000	Sigma Aldrich (A1978)
Rabbit anti-phospho-paxillin (Tyr31)	1:1000	ThermoFisher Scientific (44-720G)
Mouse anti-paxillin	1:1000	BD Biosciences (612405)
Rabbit anti-phospho FAK (Tyr576)	1:500	Epitomics (1700-1)
Rabbit anti-FAK	1:500	Epitomics (2103-1)

Secondary Antibodies	Dilution	Source
AlexaFluor 568, Goat anti-Mouse IgG	1:1000	Molecular Probes (A-11019)
Alexa Fluor 488, Goat anti-Chicken	1:1000	ThermoFisher Scientific (A-11039)
Goat anti-Chicken HRP	1:2000	Abcam (ab6877)
Veriblot for IP Detection Reagent (HRP)	1:4000	Abcam (ab131366)
Donkey anti-Rabbit IgG HRP-linked F(ab')$_2$ fragment	1:2000	Amersham-GE (NA9310)
Sheep anti-Mouse IgG HRP-linked F(ab')$_2$ fragment	1:2000 or 1:4000	Amersham-GE (NA9340)
Texas Red®-X phalloidin	1:500	ThermoFisher Scientific (T7471)

Online supplementary material

Fig. S1 shows cells overexpressing PLPPR1 resist trypsin detachment from the fibronectin

substrate. Fig. S2 shows that the C-terminal domain of PLPPR1 is not essential to its ability to

overcome CSPG inhibition of neurite outgrowth in primary hippocampal neurons. Fig. S3 shows

that the C-terminal domain of PLPPR1 is not required to impede LPA induced neurite retraction

and cell rounding. Fig. S4 shows that treatment of Neuro2a cells with LPA increases pERM

levels and induces localization to the cell membrane in EGFP transfected cells, while pERM

levels remain unchanged and localized in the cytosol in cells overexpressing PLPPR1. Fig. S5

shows that overexpression of PLPPR1 has no effect on phosphorylated levels of GSK3ß, Tau or

AKT after LPA treatment. Fig. S6 shows that without LPA stimulation, PLPPR1 has no effect on

ROCK targets, MLC, MYPT1, pERM, including GSK3ß, Tau and AKT.

Acknowledgements

We wish to thank Daniela Malide and the NHLBI Light Microscopy Core facility for their help and

guidance and Clare Waterman for materials provided.

This work was funded by the NIH Intramural Research Program.

The authors declare no competing financial interests.

Author contributions: C.A. Iweka, S. Tilve and H.M. Geller conceived the project. C.A. Iweka and

S. Tilve performed all the experiments. C.A. Iweka, S. Tilve and H.M. Geller wrote the

manuscript. All authors planned the experiments, analyzed and discussed the results and

commented on the manuscript.

References

Alabed, Y.Z., E. Grados-Munro, G.B. Ferraro, S.H. Hsieh, and A.E. Fournier. 2006.
Neuronal responses to myelin are mediated by rho kinase. *J. Neurochem.* 96:1616-
1625. doi: 10.1111/j.1471-4159.2006.03670.x.

Amano, M., K. Chihara, K. Kimura, Y. Fukata, N. Nakamura, Y. Matsuura, and K.
Kaibuchi. 1997. Formation of actin stress fibers and focal adhesions enhanced by Rho-
kinase. *Science*. 275:1308-1311.

Baudhuin, L.M., K.L. Cristina, J. Lu, and Y. Xu. 2002. Akt activation induced by
lysophosphatidic acid and sphingosine-1-phosphate requires both mitogen-activated
protein kinase kinase and p38 mitogen-activated protein kinase and is cell-line specific.
Mol. Pharmacol. 62:660-671. doi: 10.1124/mol.62.3.660.

Bellis, S.L., J.T. Miller, and C.E. Turner. 1995. Characterization of tyrosine
phosphorylation of paxillin in vitro by focal adhesion kinase. *J. Biol. Chem.* 270:17437-
17441.

Bokoch, G.M., B.P. Bohl, and T.H. Chuang. 1994. Guanine nucleotide exchange
regulates membrane translocation of Rac/Rho GTP-binding proteins. *J. Biol. Chem.*
269:31674-31679.

Boulter, E., R. Garcia-Mata, C. Guilluy, A. Dubash, G. Rossi, P.J. Brennwald, and K.
Burridge. 2010. Regulation of Rho GTPase crosstalk, degradation and activity by
RhoGDI1. *Nat. Cell Biol.* 12:477-483. doi: 10.1038/ncb2049.

Brauer, A.U., and R. Nitsch. 2008. Plasticity-related genes (PRGs/LRPs): a brain-
specific class of lysophospholipid-modifying proteins. *Biochim. Biophys. Acta*. 1781:595-
600. doi: 10.1016/j.bbalip.2008.04.004.

Brauer, A.U., N.E. Savaskan, H. Kuhn, S. Prehn, O. Ninnemann, and R. Nitsch. 2003. A
new phospholipid phosphatase, PRG-1, is involved in axon growth and regenerative
sprouting. *Nat. Neurosci.* 6:572-578. doi: 10.1038/nn1052.

Broggini, T., R. Nitsch, and N.E. Savaskan. 2010. Plasticity-related gene 5 (PRG5)
induces filopodia and neurite growth and impedes lysophosphatidic acid- and nogo-A-
mediated axonal retraction. *Mol. Biol. Cell*. 21:521-537. doi: 10.1091/mbc.E09-06-0506.

Broggini, T., L. Schnell, A. Ghoochani, J.M. Mateos, M. Buchfelder, K. Wiendieck, M.K.
Schafer, I.Y. Eyupoglu, and N.E. Savaskan. 2016. Plasticity Related Gene 3 (PRG3)
overcomes myelin-associated growth inhibition and promotes functional recovery after
spinal cord injury. *Aging (Albany N. Y.)*. 8:2463-2487. doi: 10.18632/aging.101066.

Burridge, K., and C. Guilluy. 2016. Focal adhesions, stress fibers and mechanical
tension. *Exp. Cell Res.* 343:14-20. doi: 10.1016/j.yexcr.2015.10.029.

Chen, B., H.T. Chou, C.A. Brautigam, W. Xing, S. Yang, L. Henry, L.K. Doolittle, T.
Walz, and M.K. Rosen. 2017. Rac1 GTPase activates the WAVE regulatory complex
through two distinct binding sites. *Elife*. 6. doi: 10.7554/eLife.29795.

Davies, S.J., D.R. Goucher, C. Doller, and J. Silver. 1999. Robust regeneration of adult
sensory axons in degenerating white matter of the adult rat spinal cord. *J. Neurosci.*
19:5810-5822.

DePasquale, J.A. 1998. Cell matrix adhesions and localization of the vitronectin
receptor in MCF-7 human mammary carcinoma cells. *Histochem. Cell Biol.* 110:485-
494.

Dergham, P., B. Ellezam, C. Essagian, H. Avedissian, W.D. Lubell, and L. McKerracher.
2002. Rho signaling pathway targeted to promote spinal cord repair. *J. Neurosci.*
22:6570-6577. doi: 10.1523/JNEUROSCI.22-15-06570.2002.

DerMardirossian, C., A. Schnelzer, and G.M. Bokoch. 2004. Phosphorylation of RhoGDI
by Pak1 mediates dissociation of Rac GTPase. *Mol. Cell*. 15:117-127. doi:
10.1016/j.molcel.2004.05.019.

Dovas, A., and J.R. Couchman. 2005. RhoGDI: multiple functions in the regulation of
Rho family GTPase activities. *Biochem. J.* 390:1-9. doi: 10.1042/BJ20050104.

Ehrlich, J.S., M.D. Hansen, and W.J. Nelson. 2002. Spatio-temporal regulation of Rac1
localization and lamellipodia dynamics during epithelial cell-cell adhesion. *Dev. Cell*.
3:259-270. doi: 10.1016/S1534-5807(02)00216-2.

Ellerbroek, S.M., K. Wennerberg, and K. Burridge. 2003. Serine phosphorylation
negatively regulates RhoA in vivo. *J. Biol. Chem.* 278:19023-19031. doi:
10.1074/jbc.M213066200.

Fang, X., S. Yu, J.L. Tanyi, Y. Lu, J.R. Woodgett, and G.B. Mills. 2002. Convergence of
multiple signaling cascades at glycogen synthase kinase 3: Edg receptor-mediated
phosphorylation and inactivation by lysophosphatidic acid through a protein kinase C-
dependent intracellular pathway. *Mol. Cell. Biol.* 22:2099-2110. doi:
10.1128/mcb.22.7.2099-2110.2002.

Fink, K.L., F. Lopez-Giraldez, I.J. Kim, S.M. Strittmatter, and W.B.J. Cafferty. 2017.
Identification of intrinsic axon growth modulators for intact CNS neurons after injury. *Cell*
Rep. 18:2687-2701. doi: 10.1016/j.celrep.2017.02.058.

Fournier, A.E., B.T. Takizawa, and S.M. Strittmatter. 2003. Rho kinase inhibition
enhances axonal regeneration in the injured CNS. *J. Neurosci.* 23:1416-1423. doi:
10.1523/JNEUROSCI.23-04-01416.2003.

Friedl, P., and K. Wolf. 2010. Plasticity of cell migration: a multiscale tuning model. *J.*
Cell Biol. 188:11-19. doi: 10.1083/jcb.200909003.

Garcia-Mata, R., E. Boulter, and K. Burridge. 2011. The 'invisible hand': regulation of
RHO GTPases by RHOGDIs. *Nat. Rev. Mol. Cell Biol.* 12:493-504. doi:
10.1038/nrm3153.

Gardel, M.L., I.C. Schneider, Y. Aratyn-Schaus, and C.M. Waterman. 2010. Mechanical
integration of actin and adhesion dynamics in cell migration. *Annu. Rev. Cell Dev. Biol.*
26:315-333. doi: 10.1146/annurev.cellbio.011209.122036.

Haemmerli, G., and P. Strauli. 1981. In vitro motility of cells from human epidermoid
carcinomas. A study by phase-contrast and reflection-contrast cinematography. *Int. J.*
Cancer. 27:603-610.

Hodge, R.G., and A.J. Ridley. 2016. Regulating Rho GTPases and their regulators. *Nat.*
Rev. Mol. Cell Biol. 17:496-510. doi: 10.1038/nrm.2016.67.

Hynds, D.L., and D.M. Snow. 1999. Neurite outgrowth inhibition by chondroitin sulfate
proteoglycan: stalling/stopping exceeds turning in human neuroblastoma growth cones.
Exp. Neurol. 160:244-255. doi: 10.1006/exnr.1999.7212.

Jain, A., S.M. Brady-Kalnay, and R.V. Bellamkonda. 2004. Modulation of Rho GTPase
activity alleviates chondroitin sulfate proteoglycan-dependent inhibition of neurite
extension. *J. Neurosci. Res.* 77:299-307. doi: 10.1002/jnr.20161.

Jalink, K., T. Eichholtz, F.R. Postma, E.J. van Corven, and W.H. Moolenaar. 1993.
Lysophosphatidic acid induces neuronal shape changes via a novel, receptor-mediated
signaling pathway: similarity to thrombin action. *Cell Growth Differ.* 4:247-255.

Kaneko-Kawano, T., F. Takasu, H. Naoki, Y. Sakumura, S. Ishii, T. Ueba, A. Eiyama, A.
Okada, Y. Kawano, and K. Suzuki. 2012. Dynamic regulation of myosin light chain
phosphorylation by Rho-kinase. *PLoS One.* 7:e39269. doi:
10.1371/journal.pone.0039269.

Khalaf-Nazzal, R., M.A. Stouffer, R. Olaso, L. Muresan, A. Roumegous, V. Lavilla, W.
Carpentier, I. Moutkine, S. Dumont, B. Albaud, N. Cagnard, H. Roest Crollius, and F.
Francis. 2017. Early born neurons are abnormally positioned in the doublecortin
knockout hippocampus. *Hum. Mol. Genet.* 26:90-108. doi: 10.1093/hmg/ddw370.

Kim, E.K., J.M. Park, S. Lim, J.W. Choi, H.S. Kim, H. Seok, J.K. Seo, K. Oh, D.S. Lee,
K.T. Kim, S.H. Ryu, and P.G. Suh. 2011. Activation of AMP-activated protein kinase is

essential for lysophosphatidic acid-induced cell migration in ovarian cancer cells. *J. Biol.*
Chem. 286:24036-24045. doi: 10.1074/jbc.M110.209908.

Kozma, R., S. Sarner, S. Ahmed, and L. Lim. 1997. Rho family GTPases and neuronal
growth cone remodelling: relationship between increased complexity induced by
Cdc42Hs, Rac1, and acetylcholine and collapse induced by RhoA and lysophosphatidic
acid. *Mol Cell Biol*. 17:1201-1211.

Kranenburg, O., M. Poland, M. Gebbink, L. Oomen, and W.H. Moolenaar. 1997.
Dissociation of LPA-induced cytoskeletal contraction from stress fiber formation by
differential localization of RhoA. *J. Cell Sci.* 110 (Pt 19):2417-2427.

Kranenburg, O., M. Poland, F.P. van Horck, D. Drechsel, A. Hall, and W.H. Moolenaar.
1999. Activation of RhoA by lysophosphatidic acid and Galpha12/13 subunits in
neuronal cells: induction of neurite retraction. *Mol. Biol. Cell*. 10:1851-1857. doi:
10.1091/mbc.10.6.1851.

Lai, F.P., M. Szczodrak, J. Block, J. Faix, D. Breitsprecher, H.G. Mannherz, T.E.
Stradal, G.A. Dunn, J.V. Small, and K. Rottner. 2008. Arp2/3 complex interactions and
actin network turnover in lamellipodia. *EMBO J.* 27:982-992. doi:
10.1038/emboj.2008.34.

Lauffenburger, D.A., and A.F. Horwitz. 1996. Cell migration: a physically integrated
molecular process. *Cell*. 84:359-369.

Lawson, C., S.T. Lim, S. Uryu, X.L. Chen, D.A. Calderwood, and D.D. Schlaepfer. 2012.
FAK promotes recruitment of talin to nascent adhesions to control cell motility. *J. Cell*
Biol. 196:223-232. doi: 10.1083/jcb.201108078.

Le Clainche, C., and M.F. Carlier. 2008. Regulation of actin assembly associated with
protrusion and adhesion in cell migration. *Physiol. Rev.* 88:489-513. doi:
10.1152/physrev.00021.2007.

Lingor, P., N. Teusch, K. Schwarz, R. Mueller, H. Mack, M. Bahr, and B.K. Mueller.
2007. Inhibition of Rho kinase (ROCK) increases neurite outgrowth on chondroitin
sulphate proteoglycan in vitro and axonal regeneration in the adult optic nerve in vivo. *J.*
Neurochem. 103:181-189. doi: 10.1111/j.1471-4159.2007.04756.x.

Liu, X., J. Huai, H. Endle, L. Schluter, W. Fan, Y. Li, S. Richers, H. Yurugi, K.
Rajalingam, H. Ji, H. Cheng, B. Rister, G. Horta, J. Baumgart, H. Berger, G. Laube, U.

Schmitt, M.J. Schmeisser, T.M. Boeckers, S. Tenzer, A. Vlachos, T. Deller, R. Nitsch,
and J. Vogt. 2016. PRG-1 Regulates Synaptic Plasticity via Intracellular PP2A/beta1-
Integrin Signaling. *Dev. Cell*. 38:275-290. doi: 10.1016/j.devcel.2016.06.019.

Lyons, D.M., P.S. Buckmaster, A.G. Lee, C. Wu, R. Mitra, L.M. Duffey, C.L.
Buckmaster, S. Her, P.D. Patel, and A.F. Schatzberg. 2010. Stress coping stimulates
hippocampal neurogenesis in adult monkeys. *Proc. Natl. Acad. Sci. U. S. A.* 107:14823-
14827. doi: 10.1073/pnas.0914568107.

Machesky, L.M., and A. Hall. 1997. Role of actin polymerization and adhesion to
extracellular matrix in Rac- and Rho-induced cytoskeletal reorganization. *J. Cell Biol.*
138:913-926. doi: 10.1083/jcb.138.4.913.

Machesky, L.M., and R.H. Insall. 1998. Scar1 and the related Wiskott-Aldrich syndrome
protein, WASP, regulate the actin cytoskeleton through the Arp2/3 complex. *Curr. Biol.*
8:1347-1356.

Matsui, T., M. Maeda, Y. Doi, S. Yonemura, M. Amano, K. Kaibuchi, S. Tsukita, and S.
Tsukita. 1998. Rho-kinase phosphorylates COOH-terminal threonines of
ezrin/radixin/moesin (ERM) proteins and regulates their head-to-tail association. *J. Cell
Biol.* 140:647-657.

McCarty, O.J., M.K. Larson, J.M. Auger, N. Kalia, B.T. Atkinson, A.C. Pearce, S. Ruf,
R.B. Henderson, V.L. Tybulewicz, L.M. Machesky, and S.P. Watson. 2005. Rac1 is
essential for platelet lamellipodia formation and aggregate stability under flow. *J. Biol.
Chem.* 280:39474-39484. doi: 10.1074/jbc.M504672200.

Michaelson, D., J. Silletti, G. Murphy, P. D'Eustachio, M. Rush, and M.R. Philips. 2001.
Differential localization of Rho GTPases in live cells: regulation by hypervariable regions
and RhoGDI binding. *J. Cell Biol.* 152:111-126. doi: 10.1083/jcb.152.1.111.

Mogilner, A., and G. Oster. 1996. Cell motility driven by actin polymerization. *Biophys. J.*
71:3030-3045. doi: 10.1016/S0006-3495(96)79496-1.

Monnier, P.P., A. Sierra, J.M. Schwab, S. Henke-Fahle, and B.K. Mueller. 2003. The
Rho/ROCK pathway mediates neurite growth-inhibitory activity associated with the
chondroitin sulfate proteoglycans of the CNS glial scar. *Mol. Cell. Neurosci.* 22:319-330.
doi: 10.1016/S1044-7431(02)00035-0.

Nobes, C.D., and A. Hall. 1995. Rho, rac, and cdc42 GTPases regulate the assembly of
multimolecular focal complexes associated with actin stress fibers, lamellipodia, and
filopodia. *Cell*. 81:53-62.

Ohtake, Y., D. Wong, P.M. Abdul-Muneer, M.E. Selzer, and S. Li. 2016. Two PTP
receptors mediate CSPG inhibition by convergent and divergent signaling pathways in
neurons. *Sci. Rep.* 6:37152. doi: 10.1038/srep37152.

Pasapera, A.M., I.C. Schneider, E. Rericha, D.D. Schlaepfer, and C.M. Waterman.
2010. Myosin II activity regulates vinculin recruitment to focal adhesions through FAK-
mediated paxillin phosphorylation. *J. Cell Biol.* 188:877-890. doi:
10.1083/jcb.200906012.

Pfurr, S., Y.H. Chu, C. Bohrer, F. Greulich, R. Beattie, K. Mammadzada, M. Hils, S.J.
Arnold, V. Taylor, K. Schachtrup, N.H. Uhlenhaut, and C. Schachtrup. 2017. The E2A
splice variant E47 regulates the differentiation of projection neurons via p57(KIP2)
during cortical development. *Development*. 144:3917-3931. doi: 10.1242/dev.145698.

Riddick, N., K. Ohtani, and H.K. Surks. 2008. Targeting by myosin phosphatase-RhoA
interacting protein mediates RhoA/ROCK regulation of myosin phosphatase. *J. Cell.*
Biochem. 103:1158-1170. doi: 10.1002/jcb.21488.

Ridley, A.J., M.A. Schwartz, K. Burridge, R.A. Firtel, M.H. Ginsberg, G. Borisy, J.T.
Parsons, and A.R. Horwitz. 2003. Cell migration: integrating signals from front to back.
Science. 302:1704-1709. doi: 10.1126/science.1092053.

Rottner, K., A. Hall, and J.V. Small. 1999. Interplay between Rac and Rho in the control
of substrate contact dynamics. *Curr. Biol.* 9:640-648.

Sabbatini, M.E., and J.A. Williams. 2013. Cholecystokinin-mediated RhoGDI
phosphorylation via PKCalpha promotes both RhoA and Rac1 signaling. *PLoS One*.
8:e66029. doi: 10.1371/journal.pone.0066029.

Savaskan, N.E., A.U. Brauer, and R. Nitsch. 2004. Molecular cloning and expression
regulation of PRG-3, a new member of the plasticity-related gene family. *Eur. J.*
Neurosci. 19:212-220. doi: 10.1046/j.1460-9568.2003.03078.x.

Sawada, K., K. Morishige, M. Tahara, Y. Ikebuchi, R. Kawagishi, K. Tasaka, and Y.
Murata. 2002. Lysophosphatidic acid induces focal adhesion assembly through

Rho/Rho-associated kinase pathway in human ovarian cancer cells. *Gynecol. Oncol.*
87:252-259. doi: 10.1006/gyno.2002.6831.

Sayas, C.L., A. Ariaens, B. Ponsioen, and W.H. Moolenaar. 2006. GSK-3 is activated by
the tyrosine kinase Pyk2 during LPA1-mediated neurite retraction. *Mol. Biol. Cell.*
17:1834-1844. doi: 10.1091/mbc.e05-07-0688.

Schaller, M.D., and J.T. Parsons. 1995. pp125FAK-dependent tyrosine phosphorylation
of paxillin creates a high-affinity binding site for Crk. *Mol. Cell. Biol.* 15:2635-2645.

Schmidt, S., and P. Friedl. 2010. Interstitial cell migration: integrin-dependent and
alternative adhesion mechanisms. *Cell Tissue Res.* 339:83-92. doi: 10.1007/s00441-
009-0892-9.

Schwab, M.E., J.P. Kapfhammer, and C.E. Bandtlow. 1993. Inhibitors of neurite growth.
Annu Rev Neurosci. 16:565-595. doi: 10.1146/annurev.ne.16.030193.003025.

Shaw, R.J., M. Henry, F. Solomon, and T. Jacks. 1998. RhoA-dependent
phosphorylation and relocalization of ERM proteins into apical membrane/actin
protrusions in fibroblasts. *Mol. Biol. Cell.* 9:403-419.

Sigal, Y.J., O.A. Quintero, R.E. Cheney, and A.J. Morris. 2007. Cdc42 and ARP2/3-
independent regulation of filopodia by an integral membrane lipid-phosphatase-related
protein. *J. Cell Sci.* 120:340-352. doi: 10.1242/jcs.03335.

Sinnett-Smith, J., J.A. Lunn, D. Leopoldt, and E. Rozengurt. 2001. Y-27632, an inhibitor
of Rho-associated kinases, prevents tyrosine phosphorylation of focal adhesion kinase
and paxillin induced by bombesin: dissociation from tyrosine phosphorylation of
p130(CAS). *Exp. Cell Res.* 266:292-302. doi: 10.1006/excr.2001.5219.

Somanath, P.R., and T.V. Byzova. 2009. 14-3-3beta-Rac1-p21 activated kinase
signaling regulates Akt1-mediated cytoskeletal organization, lamellipodia formation and
fibronectin matrix assembly. *J. Cell. Physiol.* 218:394-404. doi: 10.1002/jcp.21612.

Strauss, U., and A.U. Brauer. 2013. Current views on regulation and function of
plasticity-related genes (PRGs/LPPRs) in the brain. *Biochim. Biophys. Acta.* 1831:133-
138. doi: 10.1016/j.bbalip.2012.08.010.

Sutherland, C., J.A. MacDonald, and M.P. Walsh. 2016. Analysis of phosphorylation of
the myosin-targeting subunit of myosin light chain phosphatase by Phos-tag SDS-
PAGE. *Am. J. Physiol. Cell Physiol.* 310:C681-691. doi: 10.1152/ajpcell.00327.2015.

Takahashi, K., T. Sasaki, A. Mammoto, K. Takaishi, T. Kameyama, S. Tsukita, and Y.
Takai. 1997. Direct interaction of the Rho GDP dissociation inhibitor with
ezrin/radixin/moesin initiates the activation of the Rho small G protein. *J. Biol. Chem.*
272:23371-23375.

Tigyi, G., D.J. Fischer, A. Sebok, C. Yang, D.L. Dyer, and R. Miledi. 1996.
Lysophosphatidic acid-induced neurite retraction in PC12 cells: control by
phosphoinositide-Ca^{2+} signaling and Rho. *J. Neurochem.* 66:537-548.

Tkachenko, E., M. Sabouri-Ghomi, O. Pertz, C. Kim, E. Gutierrez, M. Machacek, A.
Groisman, G. Danuser, and M.H. Ginsberg. 2011. Protein kinase A governs a RhoA-
RhoGDI protrusion-retraction pacemaker in migrating cells. *Nat. Cell Biol.* 13:660-667.
doi: 10.1038/ncb2231.

Turner, C.E. 2000. Paxillin and focal adhesion signalling. *Nat. Cell Biol.* 2:E231-236.
doi: 10.1038/35046659.

van Helvert, S., C. Storm, and P. Friedl. 2018. Mechanoreciprocity in cell migration. *Nat.*
Cell Biol. 20:8-20. doi: 10.1038/s41556-017-0012-0.

Velmans, T., A. Battefeld, B. Geist, A.S. Farres, U. Strauss, and A.U. Brauer. 2013.
Plasticity-related gene 3 promotes neurite shaft protrusion. *BMC Neurosci.* 14:36. doi:
10.1186/1471-2202-14-36.

Webb, D.J., K. Donais, L.A. Whitmore, S.M. Thomas, C.E. Turner, J.T. Parsons, and
A.F. Horwitz. 2004. FAK-Src signalling through paxillin, ERK and MLCK regulates
adhesion disassembly. *Nat. Cell Biol.* 6:154-161. doi: 10.1038/ncb1094.

Webb, D.J., J.T. Parsons, and A.F. Horwitz. 2002. Adhesion assembly, disassembly
and turnover in migrating cells -- over and over and over again. *Nat. Cell Biol.* 4:E97-
100. doi: 10.1038/ncb0402-e97.

Wu, C., S.B. Asokan, M.E. Berginski, E.M. Haynes, N.E. Sharpless, J.D. Griffith, S.M.
Gomez, and J.E. Bear. 2012. Arp2/3 is critical for lamellipodia and response to
extracellular matrix cues but is dispensable for chemotaxis. *Cell.* 148:973-987. doi:
10.1016/j.cell.2011.12.034.

Yamashita, T., and M. Tohyama. 2003. The p75 receptor acts as a displacement factor
that releases Rho from Rho-GDI. *Nat. Neurosci.* 6:461-467. doi: 10.1038/nn1045.

Yoneda, A., H.A. Multhaupt, and J.R. Couchman. 2005. The Rho kinases I and II
regulate different aspects of myosin II activity. *J. Cell Biol.* 170:443-453. doi:
10.1083/jcb.200412043.

Yoshioka, K., S. Nakamori, and K. Itoh. 1999. Overexpression of small GTP-binding
protein RhoA promotes invasion of tumor cells. *Cancer Res.* 59:2004-2010.

Yu, P., C. Agbaegbu, D.A. Malide, X. Wu, Y. Katagiri, J.A. Hammer, and H.M. Geller.
2015. Cooperative interactions of LPPR family members in membrane localization and
alteration of cellular morphology. *J. Cell Sci.* 128:3210-3222. doi: 10.1242/jcs.169789.

Yu, P., T. Pisitkun, G. Wang, R. Wang, Y. Katagiri, M. Gucek, M.A. Knepper, and H.M.
Geller. 2013. Global analysis of neuronal phosphoproteome regulation by chondroitin
sulfate proteoglycans. *PLoS One*. 8:e59285. doi: 10.1371/journal.pone.0059285.

Zaidel-Bar, R., C. Ballestrem, Z. Kam, and B. Geiger. 2003. Early molecular events in
the assembly of matrix adhesions at the leading edge of migrating cells. *J. Cell Sci.*
116:4605-4613. doi: 10.1242/jcs.00792.

Zaidel-Bar, R., R. Milo, Z. Kam, and B. Geiger. 2007. A paxillin tyrosine phosphorylation
switch regulates the assembly and form of cell-matrix adhesions. *J. Cell Sci.* 120:137-
148. doi: 10.1242/jcs.03314.
997

Figure Legends

Figure 1. Cell migration is decreased in cells overexpressing PLPPR1. (A)

Trajectories of Neuro2a cells transfected with EGFP (left) or EGFP-PLPPR1 (right), are

plotted using the middle of each cell body as a point of reference. **(B)** Velocity and **(C)**

total distance travelled of migrating cells were determined using NIS Elements software.

Data represents mean ± SEM. EGFP, n = 38 cells; EGFP-PLPPR1, n = 52 cells. p-

values were calculated using student's *t*-test with Welch's analysis*,* ****p<0.0001.

Experiment was conducted in triplicate.

Figure 2. PLPPR1 increases cell adhesion. (A, B) Neuro2a cells expressing EGFP,

EGFP-PLPPR1 or EGFP-PLPP1△C43 were detached with 5mM EDTA, replated on

fibronectin coated 48-well plates and washed at 15, 30, 45 and 60min. Scale bar,

100μm. **(A)** Representative images at 15min before (top) and after washing (bottom).

(B) Quantification of the percentage of adherent cells after washing at different time

points. Data represents mean ± SEM. Individual p-values were calculated using Two-

way ANOVA with Tukey's post-hoc analysis, ****p<0.0001. **(C)** Representative IRM

images showing attachment of Neuro2a cells transfected with either EGFP (top) or

EGFP-PLPPR1 (below) to the fibronectin substrate. Scale bar, 50μm. Inset is a higher

magnification image of the boxed area. **(D)** Relative attachment of the cell to the

fibronectin substrate was calculated by thresholding darker pixels divided by the traced

cell area. A color map 'Gem' is used to display higher intensity pixels as blue and lower

intensity as orange which represent stronger and weaker adhesions respectively.

EGFP, n = 14 cells; EGFP-PLPPR1, n = 13 cells. Data represents mean ± SEM. p-

values were calculated using students *t*-test with Welch's analysis. **p<0.01. **(E)**

Representative confocal images of cells overexpressing EGFP-PLPPR-1 cells (right)

demonstrate trailing fibers (white arrow) which are absent in EGFP-transfected cells

(left). Nuclei are stained with DAPI (blue). Scale bar, 50µm. **(F)** Representative IRM

images of cells transfected with either EGFP (left) or EGFP-PLPPR1 (right). Inset is a

higher magnification of boxed area showing trailing fibers in cells overexpressing

PLPPR1. Scale bar, 50µm. **(G)** Representative images of cells co-expressing F-tractin

(mApple-FTR-940) with either EGFP (left) or EGFP-PLPPR1 (right) showing some

membrane protrusions are devoid of actin filaments. Scale bar, 50µm. Inset is a higher

magnification image of the boxed area. All experiments were conducted in triplicate.

Figure 3.

PLPPR1 expression induces changes in actin polymerization. (A) Actin stress

fibers in Neuro2a cells expressing either EGFP (left) or EGFP-PLPPR1 (right). Inset is a

higher magnification of boxed area. Scale bar, 50µm. **(B)** FRAP analysis of actin

polymerization in Neuro2a cells co-transfected with F-tractin (mApple-FTR-940) and

either EGFP (top) or EGFP-PLPPR1 (bottom). Top images demonstrate cell

morphology; the boxed areas were subjected to FRAP and imaged at the indicated

times post bleaching. Scale bar, 50µm. **(C)** Graph shows fluorescence recovery over

time in EGFP-transfected cells (black solid line) and cells overexpressing PLPPR1 (red

dotted line). **(D, E)** $t_{1/2}$ and mobile fraction were determined using Matlab. Data

represent mean ± SEM. p-values were calculated using Student's *t*-test with Welch's

analysis. *p<0.05.

Figure 4. Focal adhesion contacts are increased in cells overexpressing PLPPR1.

(A) Representative TIRF images of Neuro2a cells transfected with either EGFP or EGFP-PLPPR1, plated on fibronectin coated MatTek dishes and immunostained for paxillin and phalloidin for F-actin. White asterisks show structural difference of stress fibers in the lamellipodia. White arrows indicate focal adhesion (FA) contacts. Scale bar, 10µm. **(B)** Mean size and **(C)** mean fluorescence intensity of paxillin were determined from FAs isolated with a segmentation algorithm in ImageJ. Data represents mean ± SEM. p-values were calculated using Student's *t*-test with Welch's analysis, *p<0.05, ****p<0.0001. **(D)** Western blot of transfected Neuro2a cells, probed for pY31, paxillin, pY576 and FAK. Membranes were stripped and reprobed for GFP and ß-actin for loading control. **(E, F)** Densitometric analysis of phosphorylated protein versus total protein for paxillin and FAK was performed using Image Studio Lite. Data represents mean ± SEM. p-values were calculated using Student's *t*-test with Welch's analysis, *p<0.05. All experiments were performed in triplicate.

Figure 5. PLPPR1 attenuates LPA and CSPG induced neurite retraction by reducing RhoA activation. (A) Representative images showing EGFP (left) or EGFP-PLPPR1 (right) transfected hippocampal neurons cultured on either PLL (top) or PLL and CSPGs (bottom). Scale bar, 50µm. **(B)** Neurite outgrowth analysis of total neurite length and **(C)** longest neurite were assessed at 2 DIV using NeuronJ. EGFP/PLL, n = 112; EGFP-PLPPR1/PLL, n = 115; EGFP/CSPGs, n = 113; EGFP-PLPPR1, n = 134 neurons. Data represent mean ± SEM. p-values were calculated using Two-way

ANOVA with Tukey's posthoc analysis, **p<0.01, ***p<0.001, ****p<0.0001. **(D)** Time

lapse imaging of Neuro2a cells transfected with either EGFP (top) or EGFP-PLPPR1

(bottom) and exposed to LPA. Black arrows indicate retracting neurites. White asterisks

indicate nonretracting neurites. **(E)** Cell response was calculated as the percentage of

transfected cells that retracted their neurites. Data represent mean ± SEM. p-values

were calculated using Two-way ANOVA with Tukey posthoc analysis. *p<0.05,

****p<0.0001. Scale bar, 50µm. **(F)** RhoA-GTP was pulled down using rhotekin-RBD

beads from cell lysates prepared from transfected Neuro2a cells treated with either

FAFBSA, 10% serum or 16µM LPA for 2min. I Immunoblots were probed for RhoA. ß-

actin was used as loading control for total RhoA. Membranes were stripped and

reprobed for GFP. **(G)** Densitometric analysis was performed using Image Studio Lite.

Data represent mean ± SEM. p-values were calculated using Two-way ANOVA with

Tukey posthoc analysis, *p<0.05, ***p<0.001. All experiments were conducted in

triplicate.

Figure 6. PLPPR1 decreases LPA induced translocation and phosphorylation of

MLC, MYPT1 and ERM proteins. (A) Representative images of serum starved

Neuro2a cells expressing either EGFP or EGFP-PLPPR1 and exposed to FAFBSA or

16µM LPA for 2min were immunostained for pMLC. White arrows indicate localization of

pMLC after LPA treatment in EGFP (top) and EGFP-PLPPR1 (bottom) transfected cells.

White asterisks indicate the cell used for the profile plot in **B**. Scale bar, 50µm. **(B)**

Profile plot showing distribution and localization of pMLC in transfected cells. **(C)**

Differences in levels of pMLC was determined by assessing fluorescence intensity after

FAFBSA or LPA treatment using ImageJ. Immunoblots of cells transfected with either

EGFP or EGFP-PLPPR1 and treated with FAFBSA, serum or LPA were probed for **(D)**

GFP **(E)** pMLC and MLC **(G)** pMYPT1 and MYPT1 **(I)** pERM and ERM. **(F, H, J)**

Densitometry analysis of phosphorylated protein versus total protein was performed

using Image Studio Lite. Data represents mean ± SEM. p-values were calculated using

Two-way ANOVA with Tukey posthoc analysis, *p<0.05, **p<0.01, ***p<0.001,

****p<0.0001 respectively. All experiments were performed in triplicate.

Figure 7. PLPPR1 maintains RhoA and Rac1 interaction with RhoGDI after LPA

treatment. (A) Neuro2a cells expressing either EGFP, EGFP-PLPPR1 or EGFP-

PLPPR1ΔC43 were subjected to immunoprecipitation with GFP-MA^{TM} beads and

immunoblotted with anti-RhoGDI antibody. Input was probed for GFP. **(B)** Serum

starved transfected Neuro2a cells were treated with either FAFBSA, serum or LPA and

subjected to immunoprecipitation with RhoGDI antibody. Immunoblots of input and

immunoprecipitates were probed for either RhoGDI, RhoA or Rac1. Inputs were

stripped and reprobed for GFP and ß-actin antibody for loading control. All experiments

were conducted in triplicate.

Figure 8. Co-expression of constitutively active Rac1 with PLPPR1 rescues

morphological change. (A) Representative confocal images of Neuro2a cells co-

transfected HA-tag with either EGFP, EGFP-PLPPR1, EGFP-Rac1 T17N or cells co-

expressing EGFP-PLPPR1 with EGFP-Rac1 G12V showing a rescue of the

morphological change induced by overexpression of PLPPR1. Scale bar, 50µm. **(B)**

Rac1-GTP was pulled down using PAK-PBD beads from cell lysates prepared from Neuro2a cells co-transfected HA-tag with either EGFP, EGFP-PLPPR1, EGFP-Rac1 T17N or cells co-expressing with EGFP-PLPPR1 and EGFP-Rac1 G12V. Membranes were immunoblotted with Rac1 antibody. Membranes were stripped and reprobed with anti-GFP antibody. **(C)** Densitometric analysis of phosphorylated protein versus total protein was performed using Image Studio Lite. Data represents mean ± SEM. p-values were calculated using One-way ANOVA with Tukey posthoc analysis, **p<0.01, ****p<0.0001. **(D)** Cells transfected with either EGFP (top) or EGFP-PLPPR1 (bottom) were immunostained for Arp2/3, and phalloidin to visualize F-actin. White arrows indicate localization of Arp2/3. Inset is a higher magnification of boxed area. Scale bar, 50μm. All experiments were conducted in triplicate.

Figure 9. PLPPR1 modulates RhoA and Rac1 activity. Schematic showing the modulatory effect of PLPPR1 on RhoA and Rac1 activation through association with RhoGDI in response to CSPGs and LPA, to influence cell adhesion, cell migration and neurite outgrowth.

Movie S1. "Trailing fibers" observed in migrating PLPPR1 cells with Interference Reflection Microscopy.

Time lapse images of Neuro2a cells transfected with either EGFP or EGFP-PLPPR1 were acquired in bright field using Interference Reflection Microscopy (IRM) on Zeiss 780 LSM confocal microscope system using a 10×/0.45 NA objective lens. Movie shows 3min/frame.

Movie S2. Focal adhesions remain nascent in cells overexpressing PLPPR1 compared to EGFP-transfected cells.

Time lapse images of Neuro2a cells co-expressing mCherry-paxillin with either EGFP (left) or EGFP-PLPPR1 (right) were acquired on Delta Vision OMX microscope system using a 60×/1.49 NA TIRF objective lens. Movie highlight the size and nature of focal adhesions, including formation of trailing fibers in cells co-expressing mCherry-paxillin with EGFP-PLPPR1. Movie shows 3min/frame.

Movie S3. LPA induces neurite retraction and cell rounding in Neuro2a cells.

Serum starved Neuro2a cells were exposed to either FAFBSA (left) or 16µM LPA (right). Time lapse images were acquired on a Nikon A1R microscope with a 20×/0.75 dry objective lens with no time delay between frames for 20min. Movie shows 1min/frame.

PLPPR1 **CSPGs** **LPA**

Rac1
Rac1GDP RhoGDI Rac1GTP

RhoA
RhoGDI RhoAGDP RhoAGTP

ROCK

| Cell Migration | Cell Adhesion | Neurite Inhibition |

→ Activation
→ Inhibition
⇢ Predicted inhibition
⇢ Predicted Activation