1 **The lipid phosphatase-like protein PLPPR1 increases cell adhesion by modulating RhoA/Rac1**

2 **activity**

4 Chinyere Agbaegbu Iweka[1,2#], Sharada Tilve[1#], Caitlin Mencio[1], Yasuhiro Katagiri[1], Herbert M Geller[1]*

5 [1]Laboratory of Developmental Neurobiology, National Heart Lung and Blood Institute, NIH, Bethesda,
6 MD.
7
8 [2]Interdisciplinary Program in Neuroscience, Georgetown University, Washington, DC.
9

[#] These authors contributed equally to this work.

*Correspondence

Herbert M. Geller (E-mail: gellerh@nhlbi.nih.gov)

Laboratory of Developmental Neurobiology

Cell Biology and Physiology Center

National Heart, Lung, and Blood Institute

National Institutes of Health

9000 Rockville Pike

Bethesda, MD 20892

Phone: 301-451-9440

FAX: 301-594-8133

Running Title: PLPPR1 increases cell adhesion

SUMMARY

PLPPR1 increases cell adhesion and decreases cell motility by modulating RhoA and Rac1 activation

through its association with RhoGDI.

ABSTRACT

Phospholipid Phosphatase-Related Protein Type 1 (PLPPR1) is a member of a family of lipid phosphatase

related proteins, integral membrane proteins characterized by six transmembrane domains. PLPPR1 is

enriched in the brain and recent data indicate potential pleiotropic functions in several different contexts.

An inherent ability of PLPPR1 is to induce membrane protrusions, and we have previously reported that

members within this family may act in concert. However, the mechanism by which PLPPR1 produces

these actions is not yet understood. Here, we report that exogenous expression of PLPPR1 reduces cell

motility and increases cell adhesion to the ECM substrate by altering cytoskeletal dynamics and

modulating RhoA and Rac1 activity through association with RhoGDI. This signaling also allows

overexpression of PLPPR1 to overcome the inhibitory activity of CSPGs and LPA on neurites. Together,

these results establish a novel signaling pathway for the PLPPR1 protein.

Abbreviations: PLPPR (Phospholipid Phosphatase-Related Protein), FA (Focal Adhesions), CSPG

(Chondroitin Sulfate Proteoglycan), PLL (Poly-L-Lysine), LPA (Lysophosphatidic acid), LPP (Lipid

Phosphate Phosphatase), FAFBSA (Fatty Acid Free Bovine Serum Albumin), MLC (myosin light chain

II), MYPT1 (myosin light chain phosphatase 1), ERM (Ezrin, Radixin and Moesin), GSK (Glycogen

Synthase Kinase), GEF (Guanine Exchange Factor), GAP (GTPase Activating Protein), RhoGDI (Rho

Guanine Nucleotide Dissociation Inhibitor), IRM (Interference Reflection Microscopy), TIRF (Total

Internal Reflection Fluorescence), STED (STimulated Emission Depletion), FRAP (Fluorescence

Recovery After Photobleaching), ROCK (Rho-associated protein kinase), CA (Constitutively Active), DN

(Dominant Negative), FAK (Focal adhesion kinase), GTP (Guanosine triphosphate), GDP (Guanosine

diphosphate), WAVE (WASP-family verprolin-homologous protein).

INTRODUCTION

The Phospholipid Phosphatase Related (PLPPR) proteins (PLPPR1 – PLPPR5, previously

referred to as Plasticity Related Genes, PRGs) are highly enriched in the brain and their expression is

regulated, with the highest expression levels during development (Wang and Molnár, 2005). The first

protein of this family to be discovered, PLPPR4, was found in sprouting axons following hippocampal

deafferentation, implying a role in neuronal plasticity (Bräuer et al., 2003). Recent studies have shown

that expression of PLPPR1 mRNA correlates with sprouting corticospinal axons after injury (Fink et al.,

2017) and neuronal remodeling in the hippocampus after kainic acid treatment (Savaskan et al., 2004).

Furthermore, decreased PLPPR1 mRNA has been associated with dysregulated neuronal migration

(Khalaf-Nazzal et al., 2017; Pfurr et al., 2017), suggesting a role for these proteins in neuronal migration

as well as neurogenesis and axon growth after injury.

We and others have previously shown that overexpression of PLPPR1 induces actin-rich

membrane protrusions in many different cell types (Broggini et al., 2016; Sigal et al., 2007; Velmans et

al., 2013; Yu et al., 2015). Overexpression of PLPPR5, the closest relative of PLPPR1 in this family,

produces a similar phenotype (Broggini et al., 2010). The members of this family interact with each other

and may function as a heteromeric complex to alter cytoskeletal dynamics (Yu et al., 2015). However, the

mechanism by which PLPPR1 produces its effects is yet unknown.

Here we report that PLPPR1 expression leads to increased cell adhesion and changes in

cytoskeletal dynamics that result in decreased cell migration. This enables cells overexpressing PLPPR1

to overcome the inhibitory activity of lysophosphatidic acid (LPA) and chondroitin sulfate proteoglycans

(CSPGs). We show that these effects of PLPPR1 are due to alterations in the RhoA-ROCK signaling

pathway mediated by PLPPR1 through association with RhoGDI. These results establish a novel

signaling pathway for the PLPPR family of proteins.

RESULTS

PLPPR1 reduces cell migration and increases cell adhesion

Overexpression of PLPPR1 induces membrane protrusions in several different cell types

(Broggini et al., 2016; Savaskan et al., 2004; Sigal et al., 2007; Velmans et al., 2013; Yu et al., 2015),

implying a change in the cytoskeleton. Many cellular processes, including migration and adhesion, are

governed by dynamic changes to the cytoskeleton. To assess the potential role of PLPPR1 on these

processes, we expressed either EGFP or EGFP-PLPPR1 in Neuro2a cells, which do not express

detectable levels of PLPPR1 mRNA (data from (Iida, 2018; Llorens et al., 2013) accessible at the NCBI

GEO database), plated on a uniform fibronectin substrate. Live cell imaging was used to trace the

migratory trajectories of individual cells (Fig. 1A). We found that the velocity (Fig. 1B) and total distance

travelled (Fig. 1C) by migrating cells expressing PLPPR1 were significantly reduced compared to cells

expressing EGFP.

Cell adhesion is a major determinant for cell migration. We therefore performed both a cell

adhesion and a cell detachment assay. Neuro2a cells expressing either EGFP or EGFP-PLPPR1 were

plated on a fibronectin substrate. For the adhesion assay, cells were plated for one hour and subjected to

media washes every 15 min, at which times the number of attached cells was measured. For the

detachment assay, cells were allowed to attach overnight and their resistance to detachment by

trypsin/EDTA was measured. Our results show that cells expressing PLPPR1 were both more adherent

after media washing (Fig. 2A, B) and resistant to detachment by Trypsin/EDTA (Fig. S1).

To examine the intricacy of adhesion and mobility in cells expressing PLPPR1, we used

Interference Reflection Microscopy (IRM) to visualize and quantify the interface between the ventral cell

surface and the fibronectin substrate coated on glass. Cell membranes in close contact with the substrate

produce more interference detected as darker pixels while membranes further from the substrate have less

interference detected as lighter pixels (Fig. 2C). IRM showed that the average area of attachment to the

fibronectin substrate was significantly greater in cells expressing PLPPR1, suggesting a stronger

attachment compared to cells expressing EGFP (Fig. 2D). These observations confirm our results that

show increased attachment in cells expressing PLPPR1.

Using live-cell IRM, we observed the formation of a trail of fibrous structures as PLPPR1-

expressing cells migrate (Movie S1, Fig. 2E, F). These retraction fiber trails were formed in response to

cell movement, as opposed to previously seen actin-rich protrusions, suggesting that these "trailing

fibers" may be the result of the increased cell attachment. PLPPR1 overexpression is known to cause

filopodia like actin-rich protrusions (Velmans et al., 2013; Yu et al., 2015). To determine the nature of

these "trailing fibers", we co-expressed mApple-FTR-940 (F-tractin, a cellular probe for filamentous

actin) with either EGFP or EGFP-PLPPR1 in Neuro2a cells. Confocal images showed no expression of F-

tractin in these "trailing fibers" (Fig. 2G). The presence of such "trailing fibers", devoid of cytoskeletal

components, have been attributed to active migration in cancer cells (DePasquale, 1998; Haemmerli and

Strauli, 1981). This lack of actin implies they are not protrusions but may instead be remnants of cell

membrane that, due to the PLPPR1-induced increase in cell adhesion, could not detach readily from the

fibronectin substrate during migration.

PLPPR1 increases nascent focal adhesion complexes.

The increased adhesion observed in cells expressing PLPPR1 led us to take a closer look at focal

adhesions (FA). Focal adhesions are mechanosensitive and require the recruitment of several proteins in

order to change their composition and mature (Pasapera et al., 2010). This tension-mediated maturation

involves the recruitment of paxillin to nascent focal adhesion complexes that are initially small in size,

but then mature into focal adhesions as the size of the focal complexes increase and are stabilized

(Pasapera et al., 2010; Zaidel-Bar et al., 2003). Therefore, cells expressing either EGFP or EGFP-PLPPR1

were plated on a fibronectin substrate, immunostained for paxillin and F-actin and imaging performed

using TIRF (Fig. 3A). Measurement of the size of individual FA revealed more nascent paxillin in cells

overexpressing PLPPR1 as compared to cells expressing EGFP, suggesting reduced FA turnover (Fig.

3B). Live cell imaging confirmed that focal adhesion complexes in cells co-expressing EGFP-PLPPR1

and mCherry-paxillin remained nascent compared to the mature FA observed in cells expressing EGFP

(Movie S2).

Phosphorylation of paxillin on Y31 is crucial for the regulation of paxillin turnover and FA

complex maturation. Studies have demonstrated a reduced phospho-paxillin to total paxillin ratio in the

transition from nascent to mature FA (Zaidel-Bar et al., 2007). Therefore, we assessed phosphorylation

levels of paxillin to account for the nascent FAs observed in the PLPPR1-expressing cells.

Immunoblotting of Neuro2A cell lysates revealed an increase in phosphorylation of paxillin in cells

expressing PLPPR1 (Fig. 3C, D). Paxillin phosphorylation at Y31 is mediated by FAK (Bellis et al.,

1995; Schaller and Parsons, 1995), therefore, we evaluated phosphorylation levels of FAK.

Phosphorylation of FAK at Y576 in cells expressing PLPPR1 was increased compared to EGFP

expressing cells (Fig. 3C, D). Altogether, these results suggest that the increased phosphorylation of

paxillin and FAK in cells expressing PLPPR1 likely account for the increase in nascent FAs, resulting in

increased cell adhesion (Beningo et al., 2001).

PLPPR1 decreases the rate of actin polymerization

The dynamic organization of the actin cytoskeleton causes changes in cell shape and adhesion

strength. This generates tension and creates the pushing and pulling forces that enable cell migration

(Gardel et al., 2010; van Helvert et al., 2018). Given our results showing increased adhesion and

decreased migration in cells expressing PLPPR1, we investigated the effect of PLPPR1 on the

organization of the actin cytoskeleton. Confocal imaging demonstrated that Neuro2a cells expressing

PLPPR1 lack the network of organized linear stress fibers observed in cells expressing EGFP (Fig. 4A).

Because actin polymerization and depolymerization are required for stress fiber formation, we evaluated

actin turnover by Fluorescence Recovery After Photobleaching (FRAP) measurements in cells co-

expressing F-tractin with either EGFP or EGFP-PLPPR1 (Fig. 4B). The leading edge of the cell was

photo-bleached and fluorescence recovery was measured. Results show a decrease in the rate of

fluorescence recovery of actin in cells expressing PLPPR1 (Fig. 4C). The mobile fraction of monomeric

actin was also decreased in PLPPR1-expressing cells (Fig. 4D).

PLPPR1 overcomes CSPG and LPA neurite inhibition

Overexpression of PLPPR1 protein can overcome the inhibitory activity of LPA-induced axon

collapse and neurite retraction (Broggini et al., 2016). Furthermore, PLPPR1 has been demonstrated to

promote axon regeneration after spinal cord injury in mice (Fink et al., 2017). Chondroitin sulfate

proteoglycans (CSPGs) are universally upregulated in the glial scar following injury and, similar to LPA,

are potent negative regulators of axon growth (McKeon et al., 1991; Wang et al., 2008). Consequently,

we asked if overexpression of PLPPR1 could overcome the inhibitory activity of CSPGs. Hippocampal

neurons were transfected with EGFP or EGFP-PLPPR1 and seeded on glass coverslips coated with either

poly-L-lysine (PLL) or PLL and CSPGs (Fig. 5A). Neurite outgrowth was analyzed by measuring total

neurite length and the length of the longest neurite. Total neurite length of EGFP-transfected neurons was

reduced by 37.5% in the presence of CSPGs. This reduction was not observed in PLPPR1-transfected

neurons (Fig. 5B). Similarly, we observed a decrease in the length of the longest neurite of EGFP-

transfected neurons, while this inhibition was not observed in neurons overexpressing PLPPR1 (Fig. 5C).

LPA induces process retraction and cell rounding in several different cell types (Jalink et al.,

1993; Kranenburg et al., 1999; Tigyi et al., 1996). To confirm this in Neuro2a cells, serum starved cells

were exposed to either fatty acid free bovine serum albumin (FAFBSA) or LPA. Live cell imaging

revealed significant neurite retraction and cell rounding in cells exposed to LPA compared to FAFBSA

(Movie S3). Next, the effects of PLPPR1 expression on neurite retraction in Neuro2A cells induced by

LPA was examined. After transfection with pEGFP or pEGFP-PLPPR1, cells were exposed to either

FAFBSA or LPA. PLPPR1 expression reduced the response to LPA: 59% of cells expressing EGFP

underwent cell rounding and neurite retraction in response to LPA compared to 42% of cells expressing

PLPPR1 (Fig. 5D, E; Movie S4). Together, these results show that PLPPR1 expression impedes the

inhibitory activity of CSPGs on neurite growth and attenuates LPA-induced neurite retraction and cell

rounding.

PLPPR1 modulates RhoA activation

The RhoA signaling pathway is common to many molecules inhibitory to axon growth, including

CSPGs and LPA (Alabed et al., 2006; Jain et al., 2004; Kranenburg et al., 1999; Monnier et al., 2003).

Additionally, RhoA activation by LPA is reduced in cells overexpressing PLPPR5, the closest relative to

PLPPR1 (Broggini et al., 2010). Therefore, we asked if PLPPR1 could similarly affect RhoA activation

by LPA. Serum starved Neuro2a cells expressing EGFP or EGFP-PLPPR1 were treated with either

FAFBSA, serum or LPA. GTP-bound RhoA was immunoprecipitated using rhotekin-RBD beads and then

subjected to immunoblot analysis. While cells expressing EGFP had increased RhoA activation in

response to either serum or LPA, RhoA activation remained unchanged after serum or LPA treatment in

cells expressing PLPPR1 (Fig. 6A, B).

Rho-associated kinase (ROCK) is a major RhoA effector and is implicated in the regulation of

actomyosin dynamics and inhibition of neurite outgrowth (Amano et al., 1997; Lingor et al., 2007;

Monnier et al., 2003). ROCK mediates its effect by phosphorylating several substrates including

cytoskeletal proteins (Kaneko-Kawano et al., 2012; Matsui et al., 1998; Riddick et al., 2008; Sutherland et

al., 2016; Yoneda et al., 2005). We investigated the effect of PLPPR1 on the RhoA-ROCK pathway by

assessing the phosphorylation levels of the ROCK substrates MLC, MYPT1, and ERM in response to

LPA treatment. Cell lysates were collected from serum starved Neuro2a cells expressing EGFP or EGFP-

PLPPR1 that were treated with either FAFBSA, serum or LPA and immunoblotted for levels of pMLC,

pMYPT1, and pERM. Densitometry analysis demonstrated an increase in phosphorylation of S19 MLC,

T853 MYPT1 and T567, T564, T558 ERM proteins in EGFP-expressing cells in response to increasing

concentrations of LPA, with no change in the phosphorylated levels in cells expressing PLPPR1 (Fig

S2A). GFP immunoblots showed that expression levels for both EGFP and EGFP-PLPPR1 do not change

between treatment conditions (Fig. S2B).

LPA is also known to induce changes in phosphorylation of Akt, GSK3ß and Tau through

activation of the PI3K-PKB/Akt and PLCγ-PKC pathways respectively (Baudhuin et al., 2002; Fang et

al., 2002; Sayas et al., 2006). Therefore, we determined if the effects of PLPPR1 expression extend to

these pathways. Immunoblot analysis showed no significant change in phosphorylation levels of GSK3ß

(Y216 and S9), Tau (S396) and Akt (S473) in cells expressing PLPPR1 compared to cells expressing

EGFP after treatment with LPA (Fig. S3A).

To determine if PLPPR1 had any effect on the phosphorylation levels of ROCK target proteins

without LPA stimulation, we performed an immunoblot analysis of MLC, ERM and MYPT1, including

Akt, GSK3β or Tau, using cell lysates from Neuro2a cells expressing EGFP or EGFP-PLPPR1. Results

show no significant changes in phosphorylation levels of these target proteins in cells expressing PLPPR1

compared to cells expressing EGFP (Fig. S3B). Altogether, these results demonstrate that PLPPR1

specifically attenuates activation of the RhoA-ROCK signaling pathway in response to stimulation.

PLPPR1 modulates RhoA and Rac1 activation via RhoGDI interaction

Previously, we identified RhoGDI as a possible binding partner for PLPPR1 (Yu et al., 2013). By

direct interaction with the Rho GTPases, RhoGDI regulates their translocation from the cytosol to the

plasma membrane and impedes the nucleotide exchange of GDP for GTP (Boulter et al., 2010; Dovas and

Couchman, 2005; Sabbatini and Williams, 2013). To confirm the interaction between RhoGDI and

PLPPR1, we immunoprecipitated EGFP using GFP magnetic/agarose beads from cells expressing either

EGFP or EGFP-PLPPR1 and immunoblotted for RhoGDI. The PLPPR family share high sequence

homology except for the C-terminus, which is unique to each family member. Therefore, we also

expressed EGFP-PLPPR1ΔC43 to determine if this putative interaction is dependent on the C-terminal

domain. Our results show that both EGFP-PLPPR1 and EGFP-PLPPR1ΔC43 co-immunoprecipitated

with RhoGDI, confirming that PLPPR1 associates with RhoGDI and this association is independent of the

C-terminus (Fig. 6C).

Inactive RhoA (GDP bound form) is tightly sequestered by RhoGDI in the cytosol (Garcia-Mata

et al., 2011; Michaelson et al., 2001). LPA stimulation results in the displacement of RhoA from

RhoGDI, which allows RhoA to disengage from GDP, bind GTP and become active (Kranenburg et al.,

1997). To determine if expression of PLPPR1 would affect the RhoGDI – RhoA interaction after LPA

treatment, Neuro2a cells transfected with EGFP or EGFP-PLPPR1 were treated with either FAFBSA,

serum or LPA. RhoGDI was immunoprecipitated and immunoblot analysis of RhoA was performed. As

expected, after treatment with either serum or LPA, the interaction between RhoA and RhoGDI was

disrupted in cells expressing EGFP. However, the affinity of RhoGDI for RhoA was maintained in cells

expressing PLPPR1 even after serum or LPA treatment (Fig. 6D, E). Immunoblot analysis of GFP

expression showed that protein expression levels did not change between treatment conditions (Fig. 6D).

Like RhoA, Rac1 activation is regulated by RhoGDI (Boulter et al., 2010; DerMardirossian et al.,

2004). We assessed the effects of PLPPR1 expression on the affinity of RhoGDI for Rac1 after LPA

treatment. Neuro2a cells expressing EGFP or EGFP-PLPPR1 were exposed to either FAFBSA, serum or

LPA. RhoGDI was immunoprecipitated and immunoblotting was performed to detect Rac1 levels. Our

results show that, like RhoA, the interaction between RhoGDI and Rac1 was disrupted by LPA treatment

in cells expressing EGFP, however, this interaction was not disrupted in cells expressing PLPPR1 (Fig.

6D, F). Together, these results demonstrate that PLPPR1 modulates LPA induced activation of RhoA and

Rac1, likely via its association with RhoGDI.

Constitutively active (CA) Rac1 overexpression rescues PLPPR1 induced morphological change

Rac1 activation is essential to lamellipodial formation (Ehrlich et al., 2002; Kozma et al., 1997;

Somanath and Byzova, 2009). Given the possibility that PLPPR1 may regulate Rac1 activity, we

examined the morphology of cells expressing dominant negative (DN) Rac1 T17N, constitutively active

(CA) Rac1 G12V and cells co-expressing PLPPR1 with CA Rac1. We observed that expression of DN

Rac1 alone led to the formation of "trailing fibers" similar to those observed in cells expressing PLPPR1

(Fig. 7A). Interestingly, cells co-expressing CA Rac1 with PLPPR1 had fewer trailing fibers (Fig. 7A).

We quantified the percentage of cells that displayed trailing fibers and saw a significant reduction in this

phenotype when co-expressing CA Rac1 with PLPPR1 as compared to PLPPR1 alone (Fig 7B).

To further explore the effect of PLPPR1 on Rac1 activation, DN Rac1 and CA Rac1 were

expressed in Neuro2a cells and active Rac1 bound to GTP was pulled down using PAK-PBD beads. Basal

levels of Rac1-GTP were near zero in Neuro2a cells expressing EGFP and DN Rac1, as well as in cells

expressing PLPPR1 (Fig. 7C, D). High levels of active Rac1 were detected in cells expressing CA Rac1,

as expected. Interestingly, co-expression of CA Rac1 with PLPPR1 significantly reduced active Rac1

(Fig. 7C, D).

Arp2/3 is an actin nucleating complex that acts downstream of Rac1 and is essential to actin

assembly and lamellipodia formation (Lai et al., 2008; Machesky and Hall, 1997; Machesky and Insall,

1998; Wu et al., 2012). We examined Arp2/3 in cells expressing PLPPR1 using STED microscopy.

Results show that Arp2/3 was localized to the actin-rich lamellipodia and exhibited a diffuse cytoplasmic

distribution in cells transfected with pEGFP. Cells expressing PLPPR1 displayed a compact aggregation

of Arp2/3 (Fig. 7E), suggesting a disruption in Arp2/3 localization. Together, these results suggest a

modulatory effect of PLPPR1 on Rac1 activity, possibly through increased binding and extraction of

active Rac1 by RhoGDI.

DISCUSSION

The PLPPRs family has five members, of which PLPPR1 and PLPPR5 are most closely related.

Overexpression of either PLPPR1 or PLPPR5 in cultured cells leads to a morphological change,

including a significant increase in cellular protrusions in many cell types (Broggini et al., 2016; Velmans

et al., 2013; Yu et al., 2015). Recent studies using genetic screening approaches have identified changes

in PLPPR1 mRNA in models that show alterations of neurogenesis (Pfurr et al., 2017), neuronal

migration (Khalaf-Nazzal et al., 2017), regulation of axonal growth (Fink et al., 2017), as well as in

disease models such as schizophrenia (Windrem et al., 2017).   However, the mechanism of action by

which PLPPR1 induces morphological changes in cells and alters neural development has been a subject

of speculation. We now show that PLPPR1 expression, through modulation of GTPase signaling, alters

cell adhesion and motility (Fig. 8).

We first demonstrated that expression of PLPPR1 in Neuro2a cells results in decreased migration

due to increased cell adhesion. In addition, migrating cells expressing PLPPR1 left long plasma

membrane extensions that we have termed "trailing fibers". Migrating cells often leave behind membrane

fragments at the trailing edge of the cell (Regen and Horwitz, 1992), and their presence has been

suggested to contribute to decreased cell advance (Chen, 1981). The persistence of these "trailing fibers"

in cells expressing PLPPR1 suggest an inability of the cells to detach from the substrate during

movement, likely due to the increase in adhesion to the fibronectin substrate.

Cell adhesion is mediated through the engagement and activation of integrins, which then

promote the establishment of focal adhesions (Nobes and Hall, 1995; Rottner et al., 1999; Sinnett-Smith

et al., 2001). Mechanical tension promotes the growth and maturation of focal adhesion complexes that is

driven by RhoA-mediated myosin II activity (Pasapera et al., 2010). This is required for the coordinated

propulsive force that enables cell migration (Gardel et al., 2010; Lauffenburger and Horwitz, 1996).

PLPPR4 mediates cell adhesion through activation of ß1-integrins through the unique calmodulin-binding

domain in its C-terminal tail (Liu et al., 2016). While PLPPR1 lacks the key motifs in its C-terminal tail,

there is evidence that it may associate with β1-integrins (Yu et al., 2013). The nature of this interaction

and its role in the observed increase in cell adhesion has yet to be explored.

Several studies have demonstrated that inhibition of Rho-ROCK activity prevents the maturation

of focal adhesion complexes (Burridge and Guilluy, 2016; Sawada et al., 2002; Sinnett-Smith et al.,

2001). Therefore, it is conceivable that the reduced RhoA activity and the nascent focal adhesion

complexes that persist in cells expressing PLPPR1 may account for the increase in cell adhesion and

decrease in cell migration. This is consistent with the increase in phosphorylation of paxillin and FAK

observed in these cells, given that FAK mediated phosphorylation of paxillin plays a role in the assembly,

turnover and stability of focal adhesions (Lawson et al., 2012; Pasapera et al., 2010; Turner, 2000; Webb

et al., 2004; Zaidel-Bar et al., 2007). While increased phosphorylation of paxillin and FAK is observed in

Neuro2a cells expressing PLPPR1, it remains unclear if this effect is due to direct interactions between

PLPPR1 and focal adhesion complexes.

The PLPPR family of proteins share sequence and structural homology with the lipid phosphate

phosphatase family (LPP), which regulate LPA signaling through phosphatase activity (Bräuer and

Nitsch, 2008; Strauss and Bräuer, 2013). However, the PLPPR proteins lack key amino acid residues

essential to the phosphatase activity of the LPP protein family and are not known to exhibit any

phosphatase activity (Strauss and Bräuer, 2013). We show here that overexpression of PLPPR1 in

hippocampal neurons overcomes neurite outgrowth inhibition induced by CSPGs. Both CSPGs and LPA

are known to signal through activation of the RhoA GTPase (Alabed et al., 2006; Ohtake et al., 2016).

Expression of PLPPR1 in Neuro2a cells reduced RhoA activation in response to LPA treatment.

Furthermore, we confirmed the modulatory effect of PLPPR1 on RhoA activation of ROCK targets,

MLC, MYPT1 and the ERM proteins. RhoA activation is controlled by regulatory proteins such as the

GEFs and GDIs. GEFs catalyze the exchange of GDP for GTP, and GDIs sequester RhoA in the cytosol,

regulating its translocation to the plasma membrane (Bokoch et al., 1994; Dovas and Couchman, 2005;

Hodge and Ridley, 2016). RhoGDI was identified as a possible binding target for PLPPR1 in a proteomic

screen for interacting proteins (Yu et al., 2013). Our results confirmed this interaction by co-

immunoprecipitating PLPPR1 with RhoGDI. This association with RhoGDI was maintained in cells

expressing the PLPPR1 C-terminal deletion mutant, PLPPR1ΔC43, suggesting an alternative site of

association.

Without LPA stimulation, RhoA-GDP is tightly bound to RhoGDI (Boulter et al., 2010; Hodge

and Ridley, 2016; Tkachenko et al., 2011). Upon LPA treatment, this interaction is disrupted, enabling the

translocation of RhoA to the plasma membrane and its subsequent activation (Li et al., 2012). We have

demonstrated here that expression of PLPPR1 maintains the interaction of RhoA with RhoGDI after LPA

treatment suggesting that PLPPR1 may play a role in stabilizing the RhoA-RhoGDI complex, thereby,

modulating the RhoA GDP exchange for GTP and its subsequent activation.

We show that expression of DN Rac1 exhibits a similar morphological change as expression of

PLPPR1 in Neuro2a cells. Expression of DN Rac1 inhibits GTP loading by sequestering endogenous

GEFs and forming non-functional RhoGTPase – RhoGEF complexes. This leads to decreased membrane

ruffling and a failure to form proper lamellipodia (Debreceni et al., 2004; Ehrlich et al., 2002; McCarty et

al., 2005). The co-expression of PLPPR1 with CA Rac1 rescues the trailing fiber morphology. CA Rac1

blocks the GAP binding domain and inhibits GTPase activity (Zhang et al., 2006). With an excess of

mutated Rac1 fused to GTP, there may be an increased pool of GTP available to other RhoGTPases

leading to increased RhoA activity in cells expressing CA Rac1 rescuing the morphology. Nucleation of

actin filaments by Arp2/3 complex is driven by WAVE (WASP-family verprolin-homologous protein),

downstream of Rac1 activation (Chen et al., 2017; Lai et al., 2008). Therefore, while Arp2/3 may not be

required to induce membrane protrusions, localization of Arp2/3 is disrupted in cells overexpressing

PLPPR1 and this may be a direct consequence of altered Rac1 activity.

Taken together, our results provide evidence for a novel signaling mechanism of PLPPR1– the

modulation of RhoA and Rac1 activity by interaction with RhoGDI . This mechanism serves as a

potential explanation for the ability of PLPPR1 to overcome CSPG, lipid and myelin-mediated inhibition

of neurite outgrowth to promote axon regeneration after injury.

MATERIALS AND METHODS

Plasmid construction

pEGFP-PLPPR1 was prepared as described previously (Yu et al., 2015). Constitutively active Rac1

G12V (CA Rac1) and dominant negative Rac1 T17N (DN Rac1) were subcloned into XhoI/BamHI sites

of pEGFP-C1 (Clontech, Mountain view, CA). mApple-FTR-940 (F-tractin) was originally from Michael

Davidson's laboratory (plasmid # 54902, Addgene, Cambridge, MA). mApple-paxillin was kindly

provided by Dr. Clare Waterman (National Institutes of Health, Bethesda, MD). A construct expressing

C-terminal HA-tagged PLPPR1 (PLPPR1–HA) was a generous gift from Dr. Andrew Morris (University

of Kentucky, Lexington, KY) (Sigal et al., 2007).

Cell culture and transfection

All animal procedures were performed in accordance with protocols approved by the Institutional

Animal Care and Use Committee (IACUC) at the National Institutes of Health. Timed pregnant

C57BL/6J mice were housed in a pathogen free facility with free access to food and water under a

standard 12 h light/dark cycle.

Primary hippocampal neurons were isolated from embryonic day 18 – 20 C57BL/6J pregnant

mice and transfected with 1 µg of either pEGFP or pEGFP-PLPPR1 per well using the 96-well shuttle

Amaxa Nucleofector™ kit, program CA-138 (Lonza, Walkersville, MD). Neurons were grown in

Neurobasal media supplemented with antibiotics (penicillin/streptomycin), glutamax, B27 (ThermoFisher

Scientific, Waltham, MA) and 2 mM potassium chloride (KCl) for 48 h.

Neuro2a cells were maintained in DMEM supplemented with penicillin/streptomycin antibiotics

and 10% fetal bovine serum (FBS). Twenty-four hours after plating, cells were transfected with either

pEGFP or pEGFP-PLPPR1, using Avalanche Omni® transfection reagent (EZ Biosystems, College Park,

MD).

For CSPG treatment and neurite outgrowth analysis, transfected primary hippocampal neurons

were plated onto glass coverslips coated with either poly-l-lysine (PLL) or a soluble CSPG mix

containing neurocan, phosphacan, versican, and aggrecan isolated from embryonic chicken brain (CC117

Millipore, Billerica, MA). Neurons were subsequently immunostained for anti-βIII tubulin (Sigma

Aldrich, St. Louis, MO) and imaged using a Nikon Eclipse Microscope with a 60×/1.4 NA oil immersion

objective lens. Neurite outgrowth analysis was performed using the NeuronJ plugin of ImageJ.

All cell culture plates (Corning, Corning NY), 6-well (3506), 12-well (3512), 24-well (3527), 48-

well (3548), MatTek dishes (35 mm petri dish with 14mm microwells, No. 1.5, MatTek Corporation,

Ashland, MA) and Lab-Tek™ culture II chamber slides, (155382) (ThermoFisher Scientific, Waltam,

MA) were coated with 10 µg/ml fibronectin (Millipore, Billerica, MA), unless otherwise indicated, and

incubated overnight at 4˚C. Excess fibronectin was rinsed off by washing 3 times with phosphate buffered

saline (PBS).

Immunocytochemistry

Transfected Neuro2a cells were fixed in 4% paraformaldehyde (PFA), 48 h after plating and 24 h

after transfection. Cells were briefly rinsed in PBS and incubated in blocking buffer (0.3% Triton X-100

in PBS, supplemented with 10% normal goat serum, NGS) for 1 h and then with primary antibody diluted

in 0.3% Triton X-100 in PBS, supplemented with 2.5% NGS. Cells were washed in PBS and then

incubated with secondary antibody diluted in 0.3% Triton X-100 in PBS, supplemented with 2.5% NGS.

See Table 1 for dilutions and source of antibodies. Cells were then washed in PBS and then mounted with

DAPI diluted in Fluoromount (Sigma-Aldrich). All phalloidin (ThermoFisher Scientific, Waltam, MA)

staining was performed as previously described (Yu et al., 2015).

For STED microscopy, cells were mounted in prolong diamond mounting media (ThermoFisher

Scientific, Waltam, MA) and imaged on a Leica SP8 STED 3X/Confocal Microscope using 100×/1.4 oil

immersion objective lens (Leica Microsystems, Buffalo Grove, IL). All other imaging was performed on

a Zeiss 780 LSM confocal microscope using 63×/1.4 oil immersion objective lens.

Live cell imaging

All live cell imaging was conducted on microscopes equipped with a heated stage at 37˚C and 5%

CO_2 on either a Nikon A1R microscope (Nikon Instruments Inc., Melville, NY), a Zeiss 780 LSM

confocal microscope (Carl Zeiss, Thornwood, NY) or an Delta Vision OMX microscope (GE,

Marlborough, MA).

Cell migration assay

Neuro2a cells transfected with either pEGFP or pEGFP-PLPPR1 were plated in Lab-Tek culture

chamber slides coated with fibronectin. Time lapse images were acquired for 10 h at the rate of

1 frame/10 min on a Nikon A1R microscope using a 10×/0.45 NA objective lens. Cell migration was

analyzed by tracking each transfected cell, using the middle of each cell body as the point of reference.

The distance travelled was determined using NIS Elements software (Nikon Instruments Inc., Melville,

NY).

Cell adhesion assay

Neuro2a cells transfected with either pEGFP or pEGFP-PLPPR1 were plated in T75 culture

flasks and detached with 5mM EDTA in PBS. Cells were re-plated in a 48-well plate coated with

fibronectin at a density of 2.0×10^5 cells per well. Cells were allowed to adhere for 1 h after plating and

then washed briefly with DMEM culture media every 15 min thereafter. Fluorescence detection was

performed on a SpectraMax M3 microplate reader (Molecular Devices, San Jose, CA). The cell density of

adherent transfected cells was determined by fluorescence using the Softmax Pro7 software (Molecular

Devices, San Jose, CA). Cell adhesion was calculated as the percentage of transfected cells that persists

after each wash, from unwashed transfected cells. Cells were subsequently fixed in 4% PFA, rinsed twice

in PBS and stained with DAPI. Images were acquired on a Nikon A1R microscope using a 10×/0.45 NA

oil objective lens.

For the detachment assay, Neuro2a cells were plated on a 12-well plate coated with fibronectin

and transfected 24 h after plating with either pEGFP or pEGFP-PLPPR1. After another 24 h, cells were

briefly washed with DMEM culture media to remove serum and subsequently treated with Trypsin-EDTA

diluted in DMEM (1:10) with increasing time intervals. Cells were then rinsed with PBS, fixed with 4%

PFA and stained with DAPI. 10×10 tiled images were acquired on a Zeiss 780 LSM confocal

microscope using a 10×/0.45 NA dry objective lens.

Interference reflection microscopy

Neuro2a cells transfected with either pEGFP or pEGFP-PLPPR1 and 24 h post- transfection were

fixed with 4% PFA for 15 min. Images were acquired with a Zeiss 780 LSM confocal microscope using a

63×/1.4 NA oil immersion objective lens. Single-frame TIFF images were extracted from TIFF stacks and

processed using ImageJ. The area of each cell was traced from bright field images to obtain an ROI

(region of interest). The image was then rescaled to an 8-bit scale of 0–255 based on pixel intensity values

that fell within histogram bins exceeding 5% of the mean histogram bin height. A band pass filter was

applied based on convolution with a 41×41 pixel kernel. The histogram was then stretched again based

on a 5% cut-off of the mean values. A binary threshold was fixed by selecting pixel intensities above 90

on the $0 - 255$ scale. The image was then inverted and the number of pixels above threshold was divided

by the area of the ROI to determine intensity. A color map 'Gem' was used to display higher intensity

pixels as blue and lower intensity as orange which represent stronger and weaker adhesions respectively.

For quantification, the area of each cell was traced from bright field images to obtain the region of

interest. A threshold was applied to the IRM image of the same cell and the number of pixels above

threshold was divided by the ROI to calculate the area of cell attachment.

To measure the size of focal adhesion (FA) contacts, Neuro2a cells were plated on Fibronectin

coated MatTek dishes at a density of 2×10^4 cells and transfected with either pEGFP or pEGFP-PLPPR1.

Cells were fixed and stained with anti-paxillin antibody. Images were acquired using TIRF microscopy on

an inverted TIRF OMX Delta Vision microscope system using a 60×/1.49 NA TIRF objective lens.

Counting of FAs in single cells (n = 10 cells) was done after noise removal by thresholding and applying

a size constraint to FAs using ImageJ software.

Actin turnover analyzed by FRAP

Neuro2a cells, plated on glass-bottom MatTek dishes coated with fibronectin, were co-transfected

with F-tractin (mApple-FTR-940) and either pEGFP or pEGFP-PLPPR1. FRAP experiments were

conducted on a Nikon A1R microscope using a 60×/1.4 NA oil objective lens. A ROI was selected from

the leading edge of transfected cells and a 564 nm laser at 20% intensity was used as a pre-bleaching

signal for F-tractin. The ROI was subsequently photobleached by three lasers at 458 nm, 564 nm and 633

nm, all set at 100% intensity. The post-bleaching acquisition was carried out at 60 frames at a rate of

1 frame/0.6 s. The fluorescence intensity was determined at every time point and automatically generated

by NIS Elements software. Images were corrected for photobleaching and the fluorescence intensity of

the ROI for each frame was normalized to the average initial intensity of pre-bleaching frames and to the

area of the bleached cell. The fluorescence intensity for the pre-bleaching frames was calculated as the

average fluorescence intensity in the first 5 frames, F_{pre}. The value, F_0, was designated as the average

fluorescence intensity of the 8th frame after the photobleaching phase that occurred in the 6th and

7th frames. Individual plots for each experiment was fitted to a model function. The fluorescence recovery

of F-tractin was best fitted by a single-exponential function (double exponential function did not

significantly improve the quality of the fit) using Matlab, revealing the presence of mobile actin fractions

with different recovery kinetics. The following model was applied:

$$F(t) = A[1 - exp(-Bt)]$$

Where: A is the percentage of mobile fraction of actin and B is the time constant of fluorescence

recovery. A and B were obtained from the fitted curve. The mean ± SEM was calculated for the

percentage of fluorescence recovery and the time constant for each condition. The final graphs show the

fluorescence recovery, mobile fraction of monomeric actin and $t_{1/2}$ which indicates the time it took for F-

actin polymerization.

LPA treatment

Serum starved Neuro2a cells transfected with either pEGFP or pEGFP-PLPPR1 were treated with

fatty-acid free bovine serum albumin, FAFBSA, (Sigma Aldrich, St. Louis, MO), or Lysophosphatidic

acid, LPA (Cayman Chemical, Ann Arbor, MI) diluted in FAFBSA. Time lapse images were acquired on

a Nikon A1R microscope with a 20×/0.75 dry objective lens with no time delay between frames for 20

min. Cell response was calculated as percentage of the total transfected cell count that retracted their

neurites or underwent cell rounding.

Immunoblot assay

Neuro2a cells seeded at a density of 4×10^5 cells/well in a 6-well culture dish were transfected

with either pEGFP or pEGFP-PLPPR1. Cells were either serum-starved for 16 – 18 h or maintained in

DMEM supplemented with 10% FBS. Serum starved cells were then treated with either FAFBSA, 10%

serum or increasing concentrations of LPA: 4, 8 and 16 µM for 2 min. Media was gently but completely

removed from all cells and rinsed once with warmed PBS. Cell lysates were prepared in 2× SDS cell lysis

buffer and clarified by centrifugation. Protein quantitation was performed using Ionic Detergent

Compatibility Reagent for Pierce™ 660nm Protein Assay Reagent (ThermoFisher Scientific, Waltam,

MA) or BCA method. Equal amount of proteins (20 µg/lane) was separated by SDS-PAGE and

transferred onto a polyvinylidene difluoride (PVDF) membrane (Millipore, Billerica, MA). Membranes

were blocked for 1 h in 5% nonfat dried milk in either 0.1% Tween20 in PBS (PBST) or 0.1% Tween20

in 10 mM Tris–HCl pH 7.5 (TBST) or blocked in 5% BSA in TBST. Membranes were incubated with

primary antibodies overnight at 4˚C, washed in either PBST or TBST and subsequently incubated with

secondary antibody for 30 min. Antigen–antibody complexes were detected with KPL LumiGlo

chemiluminescent substrate (Seracare, Milford, MA). See Table 1 for source and dilutions for all primary

and secondary antibodies used.

Co-immunoprecipitation assay

Neuro2a cells were seeded at a density of 4×10^5 cells/well in a 6-well culture dish and

transfected with either pEGFP or pEGFP-PLPPR1. Cells were harvested, washed twice in ice-cold PBS

and then resuspended in ice-cold cell lysis buffer (10 mM Tris pH 7.5, 150 mM NaCl, 0.5 mM EDTA,

1% NP-40 supplemented with protease inhibitor cocktail set V EDTA free (Millipore, Billerica, MA) and

phosphatase inhibitors (Thermo Fisher Scientific, Waltham, MA). Cell lysates were cleared and then

incubated with GFP-Trap MA beads as instructed in ChromoTek GFP-Trap® MA immunoprecipitation

kit (ChromoTek Inc., Hauppauge, NY). Cell lysates were subsequently washed twice in dilution buffer

(10 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5% NP-40 supplemented with protease and

phosphatase inhibitors) and then once in wash buffer (10 mM Tris-HCl pH 7.5, 500 mM NaCl, 0.5 mM

EDTA, 0.5% NP-40 supplemented with protease and phosphatase inhibitors). Proteins were eluted and

subjected to SDS-PAGE as described above.

To immunoprecipitate RhoGDI, transfected cells were rinsed in warmed PBS, harvested in cell

lysis buffer (20 mM Tris pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 1% NP-40 supplemented with protease

and phosphatase inhibitors) and precleared with rabbit IgG serum and Protein G agarose beads (Seracare,

Milford, MA). Samples were subsequently incubated with anti-RhoGDI overnight at 4°C and then

incubated with Protein G agarose beads for 1 h at 4°C. Samples were washed twice in dilution buffer (20

mM Tris pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5% NP-40 supplemented with protease and

phosphatase inhibitors) and once in wash buffer (20 mM Tris pH 7.5, 500 mM NaCl, 0.5 mM EDTA,

0.5% NP-40 supplemented with protease and phosphatase inhibitors). Proteins were eluted and subjected

to SDS-PAGE as described above.

RhoA and Rac1 pull-down assay

Serum starved Neuro2a cells transfected with either pEGFP or pEGFP-PLPPR1 were treated with

either FAFBSA, 10% serum or 16 µM LPA. Briefly, cells were rinsed with warmed PBS and cell lysates

prepared in cell lysis buffer as instructed and provided by the RhoA or Rac1 pull-down instruction

manual and kit (Cytoskeleton, Inc. Denver, CO). Protein quantitation was performed by BCA method and

600 µg of protein was incubated with 50 µg Rhotekin-RBD beads for RhoA pull-down or with 10 µg of

PAK-PBD beads for Rac1 pull-down at 4°C for 1 h. Complexes were washed in wash buffer provided in

the pull-down kit and protein was eluted in 2× SDS sample buffer and SDS-PAGE performed as

described above.

Statistical analysis

All analyses were performed using GraphPad Prism software version 7 (GraphPad Software, La

Jolla, CA). Data was analyzed by unpaired Student's t tests, one-way ANOVA or two-way ANOVA with

Welch's correction or Tukey's post-hoc multiple comparison test to determine significance. Normality of

the distribution of the data was tested with Kolmogorov–Smirnov normality tests using the column

statistics function of GraphPad Software. All tests were two-tailed with significance indicated as follows:

*$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$. Unless otherwise specified, values represent the means

± SEM. All experiments were repeated independently at least three times.

Online supplementary material

Fig. S1 shows cells overexpressing PLPPR1 resist trypsin detachment from the fibronectin substrate. Fig.

S2 shows that expression of PLPPR1 has no effect on phosphorylated levels of GSK3ß, Tau or AKT after

LPA treatment. Fig. S3 shows that without LPA stimulation, PLPPR1 has no effect on ROCK targets,

MLC, MYPT1, pERM, including GSK3ß, Tau and AKT.

Acknowledgements

We wish to thank Daniela Malide and Xufeng Wu of the NHLBI Light Microscopy Core facility for their

help and guidance and Clare Waterman for materials provided.

This work was funded by the NIH Intramural Research Program.

The authors declare no competing financial interests.

Author contributions: C.A. Iweka, S. Tilve and H.M. Geller conceived the project. C.A. Iweka and S.

Tilve performed the experiments. C.A. Iweka, S. Tilve, C. Mencio and H.M. Geller wrote the manuscript.

All authors planned the experiments, analyzed and discussed the results and commented on the

manuscript.

Uncategorized References

Alabed, Y. Z., Grados-Munro, E., Ferraro, G. B., Hsieh, S. H. and Fournier, A. E. (2006).

Neuronal responses to myelin are mediated by Rho kinase. *J. Neurochem.* **96**, 1616-1625.

Amano, M., Chihara, K., Kimura, K., Fukata, Y., Nakamura, N., Matsuura, Y. and

Kaibuchi, K. (1997). Formation of actin stress fibers and focal adhesions enhanced by Rho-kinase.

Science **275**, 1308-11.

Baudhuin, L. M., Cristina, K. L., Lu, J. and Xu, Y. (2002). Akt activation induced by

lysophosphatidic acid and sphingosine-1-phosphate requires both mitogen-activated protein kinase kinase

and p38 mitogen-activated protein kinase and is cell-line specific. *Mol. Pharmacol.* **62**, 660-71.

Bellis, S. L., Miller, J. T. and Turner, C. E. (1995). Characterization of tyrosine

phosphorylation of paxillin in vitro by focal adhesion kinase. *J. Biol. Chem.* **270**, 17437-41.

Beningo, K. A., Dembo, M., Kaverina, I., Small, J. V. and Wang, Y. L. (2001). Nascent focal

adhesions are responsible for the generation of strong propulsive forces in migrating fibroblasts. *The*

Journal of cell biology **153**, 881-8.

Bokoch, G. M., Bohl, B. P. and Chuang, T. H. (1994). Guanine nucleotide exchange regulates

membrane translocation of Rac/Rho GTP-binding proteins. *J. Biol. Chem.* **269**, 31674-9.

Boulter, E., Garcia-Mata, R., Guilluy, C., Dubash, A., Rossi, G., Brennwald, P. J. and

Burridge, K. (2010). Regulation of Rho GTPase crosstalk, degradation and activity by RhoGDI1. *Nat.*

Cell Biol. **12**, 477-83.

Bräuer, A. U. and Nitsch, R. (2008). Plasticity-related genes (PRGs/LRPs): a brain-specific

class of lysophospholipid-modifying proteins. *Biochim. Biophys. Acta* **1781**, 595-600.

Bräuer, A. U., Savaskan, N. E., Kuhn, H., Prehn, S., Ninnemann, O. and Nitsch, R. (2003).

A new phospholipid phosphatase, PRG-1, is involved in axon growth and regenerative sprouting. *Nat.*

Neurosci. **6**, 572-8.

Broggini, T., Nitsch, R. and Savaskan, N. E. (2010). Plasticity-related gene 5 (PRG5) induces

filopodia and neurite growth and impedes lysophosphatidic acid- and nogo-A-mediated axonal retraction.

Mol. Biol. Cell **21**, 521-37.

Broggini, T., Schnell, L., Ghoochani, A., Mateos, J. M., Buchfelder, M., Wiendieck, K.,

Schafer, M. K., Eyupoglu, I. Y. and Savaskan, N. E. (2016). Plasticity Related Gene 3 (PRG3)

overcomes myelin-associated growth inhibition and promotes functional recovery after spinal cord injury.

Aging (Albany N. Y.) **8**, 2463-2487.

Burridge, K. and Guilluy, C. (2016). Focal adhesions, stress fibers and mechanical tension. *Exp.*

Cell Res. **343**, 14-20.

598 **Chen, B., Chou, H. T., Brautigam, C. A., Xing, W., Yang, S., Henry, L., Doolittle, L. K.,**

599 **Walz, T. and Rosen, M. K.** (2017). Rac1 GTPase activates the WAVE regulatory complex through two

600 distinct binding sites. *Elife* **6**.

601 **Chen, W. T.** (1981). Mechanism of retraction of the trailing edge during fibroblast movement. *J.*

602 *Cell Biol.* **90**, 187-200.

603 **Debreceni, B., Gao, Y., Guo, F., Zhu, K., Jia, B. and Zheng, Y.** (2004). Mechanisms of

604 guanine nucleotide exchange and Rac-mediated signaling revealed by a dominant negative trio mutant. *J.*

605 *Biol. Chem.* **279**, 3777-86.

606 **DePasquale, J. A.** (1998). Cell matrix adhesions and localization of the vitronectin receptor in

607 MCF-7 human mammary carcinoma cells. *Histochem. Cell Biol.* **110**, 485-94.

608 **DerMardirossian, C., Schnelzer, A. and Bokoch, G. M.** (2004). Phosphorylation of RhoGDI

609 by Pak1 mediates dissociation of Rac GTPase. *Mol. Cell* **15**, 117-27.

610 **Dovas, A. and Couchman, J. R.** (2005). RhoGDI: multiple functions in the regulation of Rho

611 family GTPase activities. *Biochem. J.* **390**, 1-9.

612 **Ehrlich, J. S., Hansen, M. D. and Nelson, W. J.** (2002). Spatio-temporal regulation of Rac1

613 localization and lamellipodia dynamics during epithelial cell-cell adhesion. *Dev. Cell* **3**, 259-70.

614 **Fang, X., Yu, S., Tanyi, J. L., Lu, Y., Woodgett, J. R. and Mills, G. B.** (2002). Convergence of

615 multiple signaling cascades at glycogen synthase kinase 3: Edg receptor-mediated phosphorylation and

616 inactivation by lysophosphatidic acid through a protein kinase C-dependent intracellular pathway. *Mol.*

617 *Cell. Biol.* **22**, 2099-2110.

618 **Fink, K. L., Lopez-Giraldez, F., Kim, I. J., Strittmatter, S. M. and Cafferty, W. B. J.** (2017).

619 Identification of intrinsic axon growth modulators for intact CNS neurons after injury. *Cell Rep.* **18**,

620 2687-2701.

621 **Garcia-Mata, R., Boulter, E. and Burridge, K.** (2011). The 'invisible hand': regulation of RHO

622 GTPases by RHOGDIs. *Nat. Rev. Mol. Cell Biol.* **12**, 493-504.

Gardel, M. L., Schneider, I. C., Aratyn-Schaus, Y. and Waterman, C. M. (2010). Mechanical

integration of actin and adhesion dynamics in cell migration. *Annu. Rev. Cell Dev. Biol.* **26**, 315-33.

Haemmerli, G. and Strauli, P. (1981). In vitro motility of cells from human epidermoid

carcinomas. A study by phase-contrast and reflection-contrast cinematography. *Int. J. Cancer* **27**, 603-10.

Hodge, R. G. and Ridley, A. J. (2016). Regulating Rho GTPases and their regulators. *Nat. Rev.*

Mol. Cell Biol. **17**, 496-510.

Iida, K. (2018). The RNA-binding protein Sfpq regulates long neuronal genes in transcriptional

elongation. *geo* **V1**, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE60246

Jain, A., Brady-Kalnay, S. M. and Bellamkonda, R. V. (2004). Modulation of Rho GTPase

activity alleviates chondroitin sulfate proteoglycan-dependent inhibition of neurite extension. *J. Neurosci.*

Res. **77**, 299-307.

Jalink, K., Eichholtz, T., Postma, F. R., van Corven, E. J. and Moolenaar, W. H. (1993).

Lysophosphatidic acid induces neuronal shape changes via a novel, receptor-mediated signaling pathway:

similarity to thrombin action. *Cell Growth Differ.* **4**, 247-55.

Kaneko-Kawano, T., Takasu, F., Naoki, H., Sakumura, Y., Ishii, S., Ueba, T., Eiyama, A.,

Okada, A., Kawano, Y. and Suzuki, K. (2012). Dynamic regulation of myosin light chain

phosphorylation by Rho-kinase. *PLoS One* **7**, e39269.

Khalaf-Nazzal, R., Stouffer, M. A., Olaso, R., Muresan, L., Roumegous, A., Lavilla, V.,

Carpentier, W., Moutkine, I., Dumont, S., Albaud, B. et al. (2017). Early born neurons are abnormally

positioned in the doublecortin knockout hippocampus. *Hum. Mol. Genet.* **26**, 90-108.

Kozma, R., Sarner, S., Ahmed, S. and Lim, L. (1997). Rho family GTPases and neuronal

growth cone remodelling: relationship between increased complexity induced by Cdc42Hs, Rac1, and

acetylcholine and collapse induced by RhoA and lysophosphatidic acid. *Mol. Cell Biol* **17**, 1201-1211.

Kranenburg, O., Poland, M., Gebbink, M., Oomen, L. and Moolenaar, W. H. (1997).

Dissociation of LPA-induced cytoskeletal contraction from stress fiber formation by differential

localization of RhoA. *J. Cell Sci.* **110 (Pt 19)**, 2417-27.

Kranenburg, O., Poland, M., van Horck, F. P., Drechsel, D., Hall, A. and Moolenaar, W. H.

(1999). Activation of RhoA by lysophosphatidic acid and Galpha12/13 subunits in neuronal cells:

induction of neurite retraction. *Mol. Biol. Cell* **10**, 1851-7.

Lai, F. P., Szczodrak, M., Block, J., Faix, J., Breitsprecher, D., Mannherz, H. G., Stradal, T.

E., Dunn, G. A., Small, J. V. and Rottner, K. (2008). Arp2/3 complex interactions and actin network

turnover in lamellipodia. *EMBO J.* **27**, 982-92.

Lawson, C., Lim, S. T., Uryu, S., Chen, X. L., Calderwood, D. A. and Schlaepfer, D. D.

(2012). FAK promotes recruitment of talin to nascent adhesions to control cell motility. *J. Cell Biol.* **196**,

223-32.

Li, Y., Kim, J. G., Kim, H. J., Moon, M. Y., Lee, J. Y., Kim, J., Kim, S. C., Song, D. K., Kim,

Y. S. and Park, J. B. (2012). Small GTPases Rap1 and RhoA regulate superoxide formation by Rac1

GTPases activation during the phagocytosis of IgG-opsonized zymosans in macrophages. *Free Radic.*

Biol. Med. **52**, 1796-805.

Lingor, P., Teusch, N., Schwarz, K., Mueller, R., Mack, H., Bahr, M. and Mueller, B. K.

(2007). Inhibition of Rho kinase (ROCK) increases neurite outgrowth on chondroitin sulphate

proteoglycan in vitro and axonal regeneration in the adult optic nerve in vivo. *J. Neurochem.* **103**, 181-9.

Liu, X., Huai, J., Endle, H., Schluter, L., Fan, W., Li, Y., Richers, S., Yurugi, H.,

Rajalingam, K., Ji, H. et al. (2016). PRG-1 regulates synaptic plasticity via intracellular PP2a/β1-

integrin signalin. *Dev. Cell* **38**, 275-90.

Llorens, F., Carulla, P., Villa, A., Torres, J. M., Fortes, P., Ferrer, I. and del Rio, J. A.

(2013). PrP(C) regulates epidermal growth factor receptor function and cell shape dynamics in Neuro2a

cells. *J. Neurochem.* **127**, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE38887

Machesky, L. M. and Hall, A. (1997). Role of actin polymerization and adhesion to extracellular

matrix in Rac- and Rho-induced cytoskeletal reorganization. *J. Cell Biol.* **138**, 913-26.

Machesky, L. M. and Insall, R. H. (1998). Scar1 and the related Wiskott-Aldrich syndrome

protein, WASP, regulate the actin cytoskeleton through the Arp2/3 complex. *Curr. Biol.* **8**, 1347-56.

Matsui, T., Maeda, M., Doi, Y., Yonemura, S., Amano, M., Kaibuchi, K., Tsukita, S. and

Tsukita, S. (1998). Rho-kinase phosphorylates COOH-terminal threonines of ezrin/radixin/moesin

(ERM) proteins and regulates their head-to-tail association. *J. Cell Biol.* **140**, 647-57.

McCarty, O. J., Larson, M. K., Auger, J. M., Kalia, N., Atkinson, B. T., Pearce, A. C., Ruf,

S., Henderson, R. B., Tybulewicz, V. L., Machesky, L. M. et al. (2005). Rac1 is essential for platelet

lamellipodia formation and aggregate stability under flow. *J. Biol. Chem.* **280**, 39474-84.

McKeon, R. J., Schreiber, R. C., Rudge, J. S. and Silver, J. (1991). Reduction of neurite

outgrowth in a model of glial scarring following CNS injury is correlated with the expression of

inhibitory molecules on reactive astrocytes. *J. Neurosci.* **11**, 3398-3411.

Michaelson, D., Silletti, J., Murphy, G., D'Eustachio, P., Rush, M. and Philips, M. R. (2001).

Differential localization of Rho GTPases in live cells: regulation by hypervariable regions and RhoGDI

binding. *J. Cell Biol.* **152**, 111-26.

Monnier, P. P., Sierra, A., Schwab, J. M., Henke-Fahle, S. and Mueller, B. K. (2003). The

Rho/ROCK pathway mediates neurite growth-inhibitory activity associated with the chondroitin sulfate

proteoglycans of the CNS glial scar. *Mol. Cell. Neurosci.* **22**, 319-30.

Nobes, C. D. and Hall, A. (1995). Rho, rac, and cdc42 GTPases regulate the assembly of

multimolecular focal complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell* **81**,

53-62.

Ohtake, Y., Wong, D., Abdul-Muneer, P. M., Selzer, M. E. and Li, S. (2016). Two PTP

receptors mediate CSPG inhibition by convergent and divergent signaling pathways in neurons. *Sci. Rep.*

6, 37152.

Pasapera, A. M., Schneider, I. C., Rericha, E., Schlaepfer, D. D. and Waterman, C. M.

(2010). Myosin II activity regulates vinculin recruitment to focal adhesions through FAK-mediated

paxillin phosphorylation. *J. Cell Biol.* **188**, 877-90.

Pfurr, S., Chu, Y. H., Bohrer, C., Greulich, F., Beattie, R., Mammadzada, K., Hils, M.,

Arnold, S. J., Taylor, V., Schachtrup, K. et al. (2017). The E2A splice variant E47 regulates the

differentiation of projection neurons via p57(KIP2) during cortical development. *Development* **144**, 3917-

3931.

Regen, C. M. and Horwitz, A. F. (1992). Dynamics of β1 integrin-mediated adhesive contacts

in motile fibroblasts. *J. Cell Biol.* **119**, 1347-59.

Riddick, N., Ohtani, K. and Surks, H. K. (2008). Targeting by myosin phosphatase-RhoA

interacting protein mediates RhoA/ROCK regulation of myosin phosphatase. *J. Cell. Biochem.* **103**, 1158-

70.

Rottner, K., Hall, A. and Small, J. V. (1999). Interplay between Rac and Rho in the control of

substrate contact dynamics. *Curr. Biol.* **9**, 640-8.

Sabbatini, M. E. and Williams, J. A. (2013). Cholecystokinin-mediated RhoGDI

phosphorylation via PKCalpha promotes both RhoA and Rac1 signaling. *PLoS One* **8**, e66029.

Savaskan, N. E., Bräuer, A. U. and Nitsch, R. (2004). Molecular cloning and expression

regulation of PRG-3, a new member of the plasticity-related gene family. *Eur. J. Neurosci.* **19**, 212-20.

Sawada, K., Morishige, K., Tahara, M., Ikebuchi, Y., Kawagishi, R., Tasaka, K. and

Murata, Y. (2002). Lysophosphatidic acid induces focal adhesion assembly through Rho/Rho-associated

kinase pathway in human ovarian cancer cells. *Gynecol. Oncol.* **87**, 252-9.

Sayas, C. L., Ariaens, A., Ponsioen, B. and Moolenaar, W. H. (2006). GSK-3 is activated by

the tyrosine kinase Pyk2 during LPA1-mediated neurite retraction. *Mol. Biol. Cell* **17**, 1834-44.

Schaller, M. D. and Parsons, J. T. (1995). pp125FAK-dependent tyrosine phosphorylation of

paxillin creates a high-affinity binding site for Crk. *Mol. Cell. Biol.* **15**, 2635-45.

Sigal, Y. J., Quintero, O. A., Cheney, R. E. and Morris, A. J. (2007). Cdc42 and ARP2/3-

independent regulation of filopodia by an integral membrane lipid-phosphatase-related protein. *J. Cell*

Sci. **120**, 340-52.

Sinnett-Smith, J., Lunn, J. A., Leopoldt, D. and Rozengurt, E. (2001). Y-27632, an inhibitor

of Rho-associated kinases, prevents tyrosine phosphorylation of focal adhesion kinase and paxillin

induced by bombesin: dissociation from tyrosine phosphorylation of p130(CAS). *Exp. Cell Res.* **266**, 292-

302.

Somanath, P. R. and Byzova, T. V. (2009). 14-3-3beta-Rac1-p21 activated kinase signaling

regulates Akt1-mediated cytoskeletal organization, lamellipodia formation and fibronectin matrix

assembly. *J. Cell. Physiol.* **218**, 394-404.

Strauss, U. and Bräuer, A. U. (2013). Current views on regulation and function of plasticity-

related genes (PRGs/LPPRs) in the brain. *Biochim. Biophys. Acta* **1831**, 133-8.

Sutherland, C., MacDonald, J. A. and Walsh, M. P. (2016). Analysis of phosphorylation of the

myosin-targeting subunit of myosin light chain phosphatase by Phos-tag SDS-PAGE. *Am. J. Physiol. Cell*

Physiol. **310**, C681-91.

Tigyi, G., Fischer, D. J., Sebok, A., Yang, C., Dyer, D. L. and Miledi, R. (1996).

Lysophosphatidic acid-induced neurite retraction in PC12 cells: control by phosphoinositide-Ca^{2+}

signaling and Rho. *J. Neurochem.* **66**, 537-48.

Tkachenko, E., Sabouri-Ghomi, M., Pertz, O., Kim, C., Gutierrez, E., Machacek, M.,

Groisman, A., Danuser, G. and Ginsberg, M. H. (2011). Protein kinase A governs a RhoA-RhoGDI

protrusion-retraction pacemaker in migrating cells. *Nat. Cell Biol.* **13**, 660-7.

Turner, C. E. (2000). Paxillin and focal adhesion signalling. *Nat. Cell Biol.* **2**, E231-6.

van Helvert, S., Storm, C. and Friedl, P. (2018). Mechanoreciprocity in cell migration. *Nat.*

Cell Biol. **20**, 8-20.

Velmans, T., Battefeld, A., Geist, B., Farres, A. S., Strauss, U. and Bräuer, A. U. (2013).

Plasticity-related gene 3 promotes neurite shaft protrusion. *BMC Neurosci.* **14**, 36.

Wang, H., Katagiri, Y., McCann, T. E., Unsworth, E., Goldsmith, P., Yu, Z. X., Tan, F.,

Santiago, L., Mills, E. M., Wang, Y. et al. (2008). Chondroitin-4-sulfation negatively regulates axonal

guidance and growth. *J. Cell Sci* **121**, 3083-3091.

Wang, W. Z. and Molnár, Z. (2005). Dynamic pattern of mRNA expression of plasticity-related

gene-3 (PRG-3) in the mouse cerebral cortex during development. *Brain Res. Bull.* **66**, 454-60.

Webb, D. J., Donais, K., Whitmore, L. A., Thomas, S. M., Turner, C. E., Parsons, J. T. and

Horwitz, A. F. (2004). FAK-Src signalling through paxillin, ERK and MLCK regulates adhesion

disassembly. *Nat. Cell Biol.* **6**, 154-61.

Windrem, M. S., Osipovitch, M., Liu, Z., Bates, J., Chandler-Militello, D., Zou, L., Munir,

J., Schanz, S., McCoy, K., Miller, R. H. et al. (2017). Human iPSC Glial Mouse Chimeras Reveal Glial

Contributions to Schizophrenia. *Cell Stem Cell* **21**, 195-208.e6.

Wu, C., Asokan, S. B., Berginski, M. E., Haynes, E. M., Sharpless, N. E., Griffith, J. D.,

Gomez, S. M. and Bear, J. E. (2012). Arp2/3 is critical for lamellipodia and response to extracellular

matrix cues but is dispensable for chemotaxis. *Cell* **148**, 973-87.

Yoneda, A., Multhaupt, H. A. and Couchman, J. R. (2005). The Rho kinases I and II regulate

different aspects of myosin II activity. *J. Cell Biol.* **170**, 443-53.

Yu, P., Agbaegbu, C., Malide, D. A., Wu, X., Katagiri, Y., Hammer, J. A. and Geller, H. M.

(2015). Cooperative interactions of LPPR family members in membrane localization and alteration of

cellular morphology. *J. Cell Sci.* **128**, 3210-22.

Yu, P., Pisitkun, T., Wang, G., Wang, R., Katagiri, Y., Gucek, M., Knepper, M. A. and

Geller, H. M. (2013). Global analysis of neuronal phosphoproteome regulation by chondroitin sulfate

proteoglycans. *PLoS One* **8**, e59285.

Zaidel-Bar, R., Ballestrem, C., Kam, Z. and Geiger, B. (2003). Early molecular events in the

assembly of matrix adhesions at the leading edge of migrating cells. *J. Cell Sci.* **116**, 4605-13.

Zaidel-Bar, R., Milo, R., Kam, Z. and Geiger, B. (2007). A paxillin tyrosine phosphorylation

switch regulates the assembly and form of cell-matrix adhesions. *J. Cell Sci.* **120**, 137-48.

Zhang, Z. G., Lambert, C. A., Servotte, S., Chometon, G., Eckes, B., Krieg, T., Lapiere, C.

M., Nusgens, B. V. and Aumailley, M. (2006). Effects of constitutively active GTPases on fibroblast

behavior. *Cell. Mol. Life Sci.* **63**, 82-91.

Figure Legends

Figure 1. Cell migration is decreased in cells overexpressing PLPPR1. (A) Trajectories of Neuro2a

cells transfected with pEGFP (left) or pEGFP-PLPPR1 (right), are plotted using the middle of each cell

body as a point of reference. **(B)** Average velocity and **(C)** total distance travelled of migrating cells were

determined using NIS Elements software. Data is represented as boxplot. EGFP (n = 38 cells; (B) Median

1.96, Range 1.19 to 2.75; (C) Median 117.7, Range 71.37 to 164.9); EGFP-PLPPR1 (n = 52 cells (B)

Median 1.33 Range 0 to 2.02; (C) Median 81.32 Range 39.95 to 121.1). p-values were calculated using

student's *t*-test with Welch's analysis, ****p<0.0001. Experiment was conducted in triplicate.

Figure 2. PLPPR1 increases cell adhesion. (A, B) Neuro2a cells expressing EGFP or EGFP-PLPPR1

were detached with 5 mM EDTA in PBS, re-plated on fibronectin coated 48-well plates and washed at 15,

30, 45 and 60 min. Scale bar, 100 µm. **(A)** Representative images at 15 min before (top) and after

washing (bottom). **(B)** Quantification of the percentage of adherent cells after washing at different time

points. Data represents mean ± SEM. Individual p-values were calculated using Two-way ANOVA with

Tukey's post-hoc analysis, ****p<0.0001. **(C)** Representative IRM images showing attachment of

Neuro2a cells transfected with either pEGFP (top) or pEGFP-PLPPR1 (below) to the fibronectin

substrate. Scale bar, 10 µm. Insets are higher magnification images of the boxed area. **(D)** Relative

attachment of the cell to the fibronectin substrate was calculated by thresholding darker pixels divided by

the traced cell area. An ImageJ color map 'Gem' is used to display higher intensity pixels as blue and

lower intensity as orange which represent stronger and weaker adhesions respectively. Data is represented

as boxplot (EGFP n = 14 cells; Median 19.57, Range 11.27 to 37.45) and (EGFP-PLPPR1, n = 13 cells;

Median 10.31, Range 6.60 to 31.92) calculated using Student's *t*-test with Welch's analysis. **p<0.01.

(E) Representative confocal images of cells overexpressing EGFP-PLPPR-1 cells (right) demonstrate

trailing fibers (white arrow) which are absent in cells expressing EGFP (left). Nuclei are stained with

DAPI (blue). Scale bar, 10 µm. **(F)** Representative IRM images of cells transfected with either pEGFP

(left) or pEGFP-PLPPR1 (right). Inset is a higher magnification of boxed area showing trailing fibers in

cells overexpressing PLPPR1. Scale bar, 10 µm. **(G)** Representative images of cells co-expressing F-

tractin (mApple-FTR-940) with either EGFP (left) or EGFP-PLPPR1 (right) showing membrane

protrusions are devoid of actin filaments. Scale bar, 10 µm. Inset is a higher magnification image of the

boxed area. All experiments were conducted in triplicate.

Figure 3. Focal adhesion contacts are increased in cells overexpressing PLPPR1. (A) Representative

TIRF images of Neuro2a cells transfected with either pEGFP or pEGFP-PLPPR1, plated on fibronectin

coated MatTek dishes and immunostained for paxillin. White dashed box area is presented at higher

magnification of paxillin staining. Scale bar, 10µm and 1µm. **(B)** Mean size of paxillin were determined

from FAs isolated with a segmentation algorithm in ImageJ. Data represents mean ± SEM. p-values were

calculated using Student's *t*-test with Welch's analysis, *p<0.05, ****p<0.0001. **(C)** Immunoblot of

transfected Neuro2a cells, probed for pY31, paxillin, pY576 and FAK. Membranes were stripped and

reprobed for GFP and β-actin for loading control. **(D)** Densitometric analysis of phosphorylated protein

versus total protein for paxillin and FAK was performed using Image Studio Lite. Data represents mean ±

SEM. p-values were calculated using Student's *t*-test with Welch's analysis, *p<0.05. All experiments

were performed in triplicate.

Figure 4. PLPPR1 expression induces changes in actin polymerization. (A) Actin stress fibers in

Neuro2a cells expressing either EGFP (left) or EGFP-PLPPR1 (right) Scale bar, 10 µm. Inset is a higher

magnification of boxed area. **(B)** FRAP analysis of actin polymerization in Neuro2a cells co-transfected

with F-tractin (mApple-FTR-940) and either pEGFP (top) or pEGFP-PLPPR1 (bottom). Top images

demonstrate cell morphology; the boxed areas were subjected to FRAP and imaged at the indicated times

post bleaching. Scale bar, 10µm. **(C)** Graph shows fluorescence recovery over time in cells expressing

EGFP (black solid line) and cells overexpressing PLPPR1 (red dotted line). **(D)** Mobile fraction was

determined using Matlab. Data is represented as boxplot (EGFP Median 0.80, Range 0.42 to 1) and

(PLPPR Median 0.62 Range 0.26 to 1). p-values were calculated using Student's *t*-test with Welch's

analysis. *p<0.05.

Figure 5. PLPPR1 attenuates LPA and CSPG mediated neurite inhibition and alters ROCK target

phosphorylation. (A) Representative images showing pEGFP (left) or pEGFP-PLPPR1 (right)

expressing hippocampal neurons cultured on either PLL (top) or PLL and CSPGs (bottom). Scale bar, 50

μm. **(B)** Analysis of total neurite length and **(C)** longest neurite were assessed at 2 DIV using NeuronJ.

EGFP/PLL, n = 112; EGFP-PLPPR1/PLL, n = 115; EGFP/CSPGs, n = 113; EGFP-PLPPR1, n = 134

neurons. Data represent mean ± SEM. p-values were calculated using Two-way ANOVA with Tukey's

posthoc analysis, **p<0.01, ***p<0.001, ****p<0.0001. **(D)** Time lapse imaging of Neuro2a cells

expressing either EGFP (top) or EGFP-PLPPR1 (bottom) and exposed to LPA. Black arrows indicate

retracting neurites. White asterisks indicate nonretracting neurites. **(E)** Cell response was calculated as the

percentage of transfected cells that retracted their neurites. Data represent mean ± SEM. p-values were

calculated using Two-way ANOVA with Tukey's posthoc analysis. *p<0.05, ****p<0.0001. Scale bar,

50μm. **(F)** Immunoblots of cells transfected with either pEGFP or pEGFP-PLPPR1 and treated with

FAFBSA (F), serum (S), or LPA were probed for pMLC and MLC, pMYPT1 and MYPT1, pERM and

ERM and **(G)** GFP. Densitometry analysis of phosphorylated MLC **(H)**, MYPT1 **(I)**, and ERM **(J)** versus

total protein was performed using Image Studio Lite. Data represents mean ± SEM. p-values were

calculated using Two-way ANOVA with Tukey's posthoc analysis, *p<0.05, **p<0.01, ***p<0.001,

****p<0.0001 respectively. All experiments were conducted in triplicate.

Figure 6. PLPPR1 reduces RhoA activation and maintains RhoA and Rac1 interaction with

RhoGDI after LPA treatment. (A) RhoA-GTP was pulled down using rhotekin-RBD beads from cell

lysates prepared from transfected Neuro2a cells treated with either FAFBSA (F), 10% serum (S) or 16

μM LPA for 2 min. Immunoblots were probed for RhoA. β-actin was used as loading control for total

RhoA. Membranes were stripped and reprobed for GFP. **(B)** Densitometric analysis was performed using

Image Studio Lite. Data represent mean ± SEM. p-values were calculated using Two-way ANOVA with

Tukey's posthoc analysis, **p<0.01, ***p<0.001. All experiments were performed in triplicate. **(C)**

Neuro2a cells expressing either EGFP, EGFP-PLPPR1 or EGFP-PLPPR1ΔC43 were subjected to

immunoprecipitation with GFP-MA™ beads and immunoblotted with anti-RhoGDI antibody. Input was

probed for GFP. **(D)** Serum starved transfected Neuro2a cells were treated with either FAFBSA (F),

serum (S) or LPA and subjected to immunoprecipitation with RhoGDI antibody. Immunoblots of input

and immunoprecipitates were probed for either RhoGDI, RhoA or Rac1. Inputs were stripped and

reprobed for GFP and β-actin antibody for loading control. Densitometric analysis of RhoA **(E)** and Rac1

(F) was performed using Image Studio Lite. Data represents mean ± SEM. p-values were calculated using

Two-way ANOVA with Tukey's posthoc analysis, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

respectively. All experiments were performed in triplicate.

Figure 7. Co-expression of constitutively active Rac1 with PLPPR1 rescues morphological change.

(A) Representative confocal images of Neuro2a cells expressing EGFP+HA, EGFP+PLPPR1-HA, EGFP-

DN Rac1+HA, EGFP-CA Rac1+HA and EGFP-CA Rac1+PLPPR1-HA. Cells expressing EGFP-CA

Rac1+PLPPR1-HA show a rescue of the morphological change induced by expression of PLPPR1. Scale

bar, 10 µm. **(B)** Percentage of cells expressing the PLPPR1 trailing fibers phenotype of Neuro2a cells

expressing EGFP+HA, EGFP-PLPPR1+HA, EGFP-DN Rac1+HA, EGFP-CA Rac1+HA and EGFP-CA

Rac1+PLPPR1-HA**. (C)** Rac1-GTP was pulled down using PAK-PBD beads from cell lysates prepared

from Neuro2a cells expressing EGFP+HA, EGFP+PLPPR1-HA, EGFP-DN Rac1+HA, EGFP-CA

Rac1+HA and EGFP-CA Rac1+PLPPR1-HA. Membranes were immunoblotted with Rac1 antibody.

Membranes were stripped and reprobed with anti-GFP antibody. **(D)** Densitometric analysis of Rac GTP

versus total Rac1 was performed using Image Studio Lite. Data represents mean ± SEM. p-values were

calculated using One-way ANOVA with Tukey's posthoc analysis, **p<0.01, ****p<0.0001. **(E)** Cells

transfected with either pEGFP (top) or pEGFP-PLPPR1 (bottom) were immunostained for Arp2/3, and

phalloidin to visualize F-actin. White arrows indicate localization of Arp2/3. Inset is a higher

magnification of boxed area. Scale bar, 10 µm. All experiments were conducted in triplicate.

Figure 8. PLPPR1 modulates RhoA and Rac1 activity. Schematic showing the modulatory effect of

PLPPR1 on RhoA and Rac1 activation through association with RhoGDI in response to CSPGs and LPA,

to influence cell adhesion, cell migration and neurite outgrowth.

Movie S1. "Trailing fibers" observed in migrating PLPPR1 cells with Interference Reflection

Microscopy.

Time lapse images of Neuro2a cells transfected with either pEGFP or pEGFP-PLPPR1 were acquired in

bright field using Interference Reflection Microscopy (IRM) on Zeiss 780 LSM confocal microscope

system using a 10×/0.45 NA objective lens. Movie shows 3 min/frame.

Movie S2. Focal adhesions remain nascent in cells overexpressing PLPPR1 compared to cells

expressing EGFP.

Time lapse images of Neuro2a cells co-expressing mCherry-paxillin with either EGFP (left) or EGFP-

PLPPR1 (right) were acquired on Delta Vision OMX microscope system using a 60×/1.49 NA TIRF

objective lens. Movie highlight the size and nature of focal adhesions, including formation of trailing

fibers in cells co-expressing mCherry-paxillin with EGFP-PLPPR1. Movie shows 3 min/frame.

Movie S3. LPA induces neurite retraction and cell rounding in Neuro2a cells. Serum starved

Neuro2a cells were exposed to either FAFBSA (left) or 16 µM LPA (right). Time lapse images were

acquired on a Nikon A1R microscope with a 20×/0.75 dry objective lens with no time delay between

frames for 20 min. Movie shows 1 min/frame.

Movie S4. PLPPR1 impedes LPA induced neurite retraction in Neuro2a cells. Serum starved

Neuro2a cells transfected with either pEGFP (left) or pEGFP-PLPPR1 (right) were exposed to 16 µM

LPA. Time lapse images were acquired on a Nikon A1R microscope with a 20×/0.75 dry objective lens

with no time delay between frames for 20 min. Movie shows 1 min/frame.

A

EGFP/DAPI   EGFP-PLPPR1/DAPI

0 min

15 min

B

% Adherent Cells

■ EGFP
■ EGFP-PLPPR1

****   ****   ****   ****

0   15   30   45   60

C

EGFP/F-actin

EGFP-PLPPR1/F-actin

High

Low

D

% Area attached to substrate

**

EGFP   EGFP-
            PLPPR1

E

EGFP/DAPI   EGFP-PLPPR1/DAPI

F

EGFP   EGFP-PLPPR1

G

EGFP/F-tractin   EGFP

                                  F-tractin

EGFP-PLPPR1/F-tractin   PLPPR1

                                 F-tractin

Figure S1. **Cells expressing PLPPR1 are resistant to detachment. (A)** Representative images of Neuro2a cells transfected either pEGFP (top) or pEGFP-PLPPR1 (bottom) and subjected to trypsin/EDTA treatment over time. Scale bar, 100μm. **(B)** Quantification of the number of adherent cells resistant to detachment was determined as a percentage of total unwashed transfected cells. Data represent mean ± SEM. p-values were calculated using Two-way ANOVA repeated measures with Tukey posthoc analysis. *p<0.05, **p<0.01. Experiment was conducted in triplicate.

Figure S2. **Expression of PLPPR1 reduces phosphorylation levels of RhoA downstream effectors after LPA treatment.** Neuro2a cells transfected with either pEGFP or pEGFP-PLPPR1 were serum starved and treated with either FAFBSA (F), serum (S), or LPA for 5min. **(A)** Immunoblots of cells transfected with either pEGFP or pEGFP-PLPPR1 and treated with FAFBSA, serum, or LPA were probed for pMLC and MLC, pMYPT1 and MYPT1, pERM and ERM. **(B)** Immunoblot for EGFP to show equal expression. Densitometry analysis of phosphorylated **(C)** MLC, **(D)** MYPT1, and **(E)** ERM versus total protein was performed using Image Studio Lite. Data represents mean ± SEM. p-values were calculated using Two-way ANOVA with Tukey's posthoc analysis, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 respectively. All experiments were conducted in triplicate.

Figure S3. **Effect of PLPPR1 expression on non-ROCK target phosphorylation after LPA treatment and all targets without LPA. (A)** Immunoblots showing phosphorylated levels of several ROCK targets Tau, AKT, and GSK3ß in serum starved Neuro2a cells expressing EGFP or EGFP-PLPPR1 and exposed to FAFBSA (F), LPA or serum (S) for 5 min. **(B)** All membranes were stripped and reprobed for each total protein and GFP. **C)** Immunoblots showing phosphorylated levels of several ROCK targets, MYPT1, ERM, MLC, AKT, Tau, GSK3ß in Neuro2a cells expressing EGFP or EGFP-PLPPR1. All membranes were stripped and reprobed for each total protein and GFP. ß-actin was used as loading control.

A

EGFP/HA/DAPI   EGFP/PLPPR1-HA/DAPI   EGFP-DN Rac1/HA/DAPI   EGFP-CA Rac1/HA/DAPI   EGFP-CA Rac1/ PLPPR1-HA/DAPI

B

% cells with trailing fibers

****
***
**
***
**
**

EGFP + HA
EGFP- PLPPR1 + HA
EGFP-DN Rac1 + HA
EGFP-CA Rac1 + HA
EGFP-CA Rac1 + PLPPR1-HA

C

MW (kDa)

EGFP + HA
EGFP + PLPPR1-HA
EGFP-DN Rac1 + HA
EGFP-CA Rac1 + HA
EGFP-CA Rac1+PLPPR1-HA

58 — Rac1-GTP

58 — EGFP-Rac1
46 —

Total Rac1
32 —

58 — EGFP-Rac1
46 —
32 — GFP

32 — PLPPR1-HA HA

D

Relative density (a.u)

**
****
**

EGFP + HA
EGFP-PLPPR1 + HA
EGFP-DN Rac1 + HA
EGFP-CA Rac1 + HA
EGFP-CA Rac1 + PLPPR1-HA

E

EGFP   Arp2/3   F-actin   Merge

EGFP   EGFP/F-actin/Arp2/3

EGFP-PLPPR1   EGFP-PLPPR1/F-actin/Arp2/3

Table 1. Antibodies Used

Primary Antibodies	Dilution	Source
Mouse anti-βIII tubulin	1:1000	Sigma Aldrich (SAB4700544)
Chicken anti-GFP	1:2000	Abcam (ab13970)
Rabbit anti-phospho-myosin light chain II (Ser19)	1:1000	Cell Signaling Technology (3671)
Rabbit anti-phospho-myosin light chain phosphatase I (Thr853)	1:1000	Cell Signaling Technology (4563)
Rabbit anti-phospho-GSK3β (Tyr216)	1:1000	Abcam (ab75745)
Rabbit anti-pERM (Thr567, Thr564, Thr558)	1:1000	Cell Signaling Technology (3141)
Rabbit anti-phospho-AKT (Ser473)	1:1000	Cell Signaling Technology (9271)
Rabbit anti-phospho-Tau (Ser396)	1:1000	Abcam (ab109390)
Rabbit anti-myosin light chain II	1:1000	Cell Signaling Technology (3672)
Rabbit anti-myosin light chain phosphatase I	1:1000	Cell Signaling Technology (2634)
Rabbit anti-GSK3β	1:1000	Cell Signaling Technology (27C10)
Rabbit anti-ERM	1:1000	Cell Signaling Technology (3142)
Rabbit anti-AKT	1:1000	Cell Signaling Technology (4685)
Rabbit anti-RhoGDI	1:1000	ThermoFisher Scientific (51-1000Z)
Anti-p34-Arc/ARPC2	1:100	Sigma Aldrich (07-227)
Mouse anti-Rac1	1:500	Cytoskeleton, Inc. (ARC03)
Mouse anti-Tau	1:2000	ThermoFisher Scientific (MA5-12808)
Mouse anti-RhoA	1:500	Cytoskeleton, Inc. (ARH03)
Mouse anti-β-actin	1:4000	Sigma Aldrich (A1978)
Rabbit anti-phospho-paxillin (Tyr31)	1:1000	ThermoFisher Scientific (44-720G)
Mouse anti-paxillin	1:1000	BD Biosciences (612405)
Rabbit anti-phospho FAK (Tyr576)	1:500	Epitomics (1700-1)

Rabbit anti-FAK   1:500   Epitomics (2103-1)

Secondary Antibodies	Dilution	Source
Alexa Fluor 568, Goat anti-Mouse IgG	1:1000	Molecular Probes (A-11019)
Alexa Fluor 488, Goat anti-Chicken	1:1000	ThermoFisher Scientific (A-11039)
Goat anti-Chicken HRP	1:2000	Abcam (ab6877)
Veriblot for IP Detection Reagent (HRP)	1:4000	Abcam (ab131366)
Donkey anti-Rabbit IgG HRP-linked F(ab')$_2$ fragment	1:2000	Amersham-GE (NA9310)
Sheep anti-Mouse IgG HRP-linked F(ab')$_2$ fragment	1:2000 or 1:4000	Amersham-GE (NA9340)
Texas Red®-X phalloidin	1:500	ThermoFisher Scientific (T7471)