Combined use of *Candida Utilis* and *Idesia polycarpa* var. *Vestita* Fruit Improve the Production Performance of Laying Quail

Xinanbei Liu[*,†]; Na Li[*,†]; Yueyue Shu[*,†]; Yiran Sun[*,†]; Yu Li[*,†]; Wan Hua[*,†];

Yang Ye[*,†]; Fang Chen[*,†]; Lin Tang[*,†,1]

* Key laboratory of Bio-Resources and Eco-Environment of Ministry of Education, College of Life sciences, Sichuan University, Chengdu 610064, P.R.China;

† National and Local Joint Engineering Laboratory for Energy Plant Bio-oil Production and Application, Chengdu 610064, China

[1]**Corresponding Author:** Key laboratory of Bio-resources and Eco-environment of Ministry of Education, college of Life sciences, Sichuan University, Chengdu 610064, P. R. China. Email: tanglin@scu.edu.cn (L. Tang). Tel: (028) 8541 7281

Abbreviated Title： A new feeding raw material applying on laying quails.

ABSTRACT: *Idesia polycarpa* Maxim. var. *vestita* Diels (*Idesia polycarpa*), which is widely distributed in south China, is still underexplored. This study applied *Idesia polycarpa* defatted fruit (IPF) and *Candida utilis* to the feed of laying quails using solid-state fermentation. In comparison to the standard diet group, birds fed with mixture of IPF and *candida utilis* (MIC) showed better production capacity, and the ML group (1% MIC added) achieved the greatest egg mass (9.77 on average; $P<0.01$) and laying rate (87.7% on average; $P<0.01$). Compared to the standard diet group, the cholesterol content was lower in both the ML (1% MIC addition groups) and IL (1% IPF addition groups), and 5% MIC added group had higher n-3 polyunsaturated fatty acid content. Furthermore, birds given the MIC dietary supplement showed a thicker jejunum wall than the standard diet group. In addition, the related mRNA expression of SRBEP-1, SREBP-2, ATGL, APOVLDL-II which are involved in the fatty acid and cholesterol biosynthesis suggested that the addition of *candida utilis* could effectively improve the production capacity of laying quails while decrease the negative effects of IPF. This work also demonstrates how MIC can be applied to improve the production of laying quails.

KEYWORDS: *Candida utilis, Idesia polycarpa fruit, Laying quails, Egg production,.*

Introduction

Idesia polycarpa (*Idesia polycarpa* Maxim. var. *vestita* Diels) is widely distributed in south China, and it is known for its high-quality edible oil, which has a high content of conjugated linoleic acid (CLA). Moreover, the application of *Idesia polycarpa* oil in biodiesel production has been demonstrated (1, 2). There have been several studies on *Idesia polycarpa* fruit, including functional research on *Idesia polycarpa* fruit extraction and studies on the improvement of the extraction process. The remarkable antioxidant qualities, anti-skin-aging ability (3) and anti-adipogenic ability (4) of *Idesia polycarpa* extracts have been demonstrated, and an optimized extraction method has also been established (5). However, reports on the application of *Idesia polycarpa* fruit for the other purposes are still limited. In this study, we are tried to apply *Idesia polycarpa* defatted fruit (IPF) in laying quail's breeding, and for purpose of effective use, solid-fermentation was selected.

From a historical perspective, phytogenic materials fermented with specific microbe are recognized to benefit farm animals and are described as far back as the beginning of the last century (6, 7). Even now, fermentation is still gaining interest because of its important role in the production of quality feed and the utilization of unexploited feed materials (8, 9). Moreover, based on existing studies, the addition of probiotic could effectively decrease anti-nutritive factors and allergenic proteins (10, 11). Furthermore, fermented feed represents a plausible alternative to traditional antibiotics, as it enhances a number of important processes in animals (12, 13).

Candida utilis is an important member of edible yeast, which has safely been

used as a food or feed additive in food and feed industry for years. *Candida utilis* is

able to robust growth on inexpensive nitrogen and carbon sources, such as pentoses

(14). The main product of *Candida utilis* is single-cell protein, which has excellent

nutritional properties and has an agreeable odor and flavor, and this odor and flavor

can also be conferred to product by fermentation (15, 16). Therefore, *Candida utilis*

is a good choice in our study.

As an important product in the poultry industry, quail eggs are gaining increasing

interest thanks to their good taste, health benefits and nutrient richness(17, 18). In

addition, as a sensitive experimental animal (19), *Coturnix japonica* and its eggs are

often used in research (20).The effects of various kinds of materials have been tested

on the production of quail eggs, for instance, soybean (21), cassava meal (22), and

sunflower meal (23). However, no reports are available regarding the effects of *Idesia*

polycarpa products on the quality of quail eggs.

In this study, mixture of IPF and *candida utilis* (MIC) was added to the feed of

laying quails as a probiotc additive. Production capacity, health performance and egg

quality of laying quails were analysed during or after the feeding phase. Moreover, the

relative mRNA expression level of SRBEP-1, SREBP-2, ATGL, APOVLDL-II which

are involved in the fatty acid and cholesterol biosynthesis are measured.

Materials and methods

Probiotic and Feed

Candida utilis was purchased from China Center of Industrial Culture Collection

(31170, CICC).

Idesia polycarpa defatted fruits (IPF) which obtained by physical cold pressing

process were kindly supplied by Agriculture Bureau of Jianyang, China, and

processed into a dried powder before it be used.

Mixture of *Idesia polycarpa* defatted fruits and *Candida utilis* (MIC) was

produced by a solid fermentation method (solid-liquid ration 1:1.2; ammonium nitrate

8.5%; inoculum size 10.5% (*Candida utilis* in suspension form, 1.0×10^7 colony

forming units (cfu)/ml); temperature 30℃; time 4d;), and processed into a dried

powder before it be used.

Basal feed (BD, Tequ302, Sichuan) in powder form which contains all nutrients

necessary for animals was purchased in the local market. Nutrients of MIC, IPF and

basal diet are shown in **Table 1**.

Birds and Experimental Design

All animal protocols were approved by the Sichuan University animal Care and

Use committee. A total of 250 layer quails (*Coturnix coturnix japonica*, 90 days old)

from a local breeding farm (Chengdu, China) were randomly divided into 5 groups:

Control, base diet only; MH, basal diet supplemented with 5% MIC; ML, basal diet

supplemented with 1% MIC; IH, basal diet supplemented with 5% IPF; IL, basal diet

supplemented with 1% IPF. All birds were kept in a room with a relatively stable

temperature (24-30 ℃) and humidity (65%). During the 8-week experimental period,

birds had access to water *ad libitum*, but with a fixed amount of diet (1,000 g/d,

equivalent to a dose of 20 g/d/bird), which based on the local farm feeding

management for this season. In addition, from week 0 to 2, a trace amount of MIC

(0.5g (0.05%) per day) was added into the daily diet of each group to help quails

adaptation to the peculiar smell of the feed，except the control group. The quail's

differentiated diet was supplied during the 6 weeks feeding test period from week 3 to

week 8. The feeding experimental design is shown in **Table 2**.

Sample collection

The number of eggs was counted every day, and the results are represented as the

mean value of each week. Eggs were randomly collected (five eggs from each group,

30 eggs in total) each week to measure the egg quality during the experimental period.

The egg weight, density, and axis ratio, as well as the egg yolk weight and the ratio

between the egg weight and egg yolk weight were measured. The Haugh unit values

were calculated using the Haugh unit formula based on egg weight and albumen

height, as determined using a Vernier caliper. The eggshell thickness and membrane

thickness were determined as the mean value of measurements taken at three locations

on the egg (sharp end, blunt end, and middle section).

The egg yolk samples were separated from the rest of the egg for triglyceride and

cholesterol determination on weeks 6 to 8. The yolk triglycerides and cholesterol were

determined using ELISA (SMP500-15859-SBRE, Molecular Devices, USA) using

commercial kits according to the manufacturer's protocol (Jiancheng, Nanjing, China).

On week 8, three additional eggs/each group were collected for yolk lipid extraction.

At the end of the Phase 2 (week 8), three quails in each group were randomly

sacrificed by cervical dislocation for sample collection after weighing. Tissue samples,

including those from the intestine, interclavicular fat pad and liver, were immediately

collected from all 15 quails after they were sacrificed. The same segment of the

intestine was collected and stored in a prepared fixative (10% formalin) for histology

analysis, the fat pads were immediately weighed, and both the fat pad and liver

samples were snap-frozen in liquid nitrogen and stored at -80 ℃ until the subsequent

RNA isolation.

Yolk lipid extraction

112 Yolk samples were separated from boiled eggs. Yolk lipids were extracted from

113 three yolks in each group using hexane extraction (AOAC 2003. 06). The extracted

114 yolk lipids were stored in -20 ℃ until the subsequent fatty acids analysis.

Methylation and GC/MS

116 The methylation of the yolk lipids was determined using the alkali catalysis

117 method (24). The peak area and fatty acid percentages were calculated using the

118 GCMS-solution software (Shimadzu, Japan), and the fatty acid methyl esters are

119 expressed as the percentage of the total fatty acids.

Jejunum morphology

After being fixed in 10% formalin for 24 h, samples of the jejunum tissues were

selected for 5 μm-thick paraffin sections and stained with hematoxylin-eosin (HE).

The analysis included well thickness, villus height and crypt depth and was performed

using a microscope equipped with a camera and connected to a computer with the

appropriate software (Olympus, BX53, Cell^B software, Japan). Data were collected

from at least five randomly selected visible regions under the microscope. Mean

values of three birds per group were calculated for the subsequent statistical analysis.

RNA isolation

The total RNA was extracted from the fat pad and liver tissue using TRIzol

reagent (TakaRa, USA). The RNA quality and quantity were determined using a

Nanodrop 2000 (Thermo Scientific, USA). The 260/280 absorption ratios of all

extracted samples were between 1.8 and 2.0, and the RNA integrity was further

analyzed using gel electrophoresis. The first-strand cDNA was immediately

synthesized using the PrimeScript RT reagent kit (TakaRa, USA) following the

manufacturer's instructions. The obtained cDNA was stored at -20 ℃ until the

subsequent real-time PCR analysis.

qRT-PCR analysis

The primers used for target gene amplification were either designed using primer

3.0 based on quail-related gene sequences or reported by specific references (**Table 3**).

β-actin was selected as the housekeeping gene to normalize the target gene expression.

All primers in this study were synthesized by Tsingke Biotech (Chengdu, China). The

specificity of each of the designed primers was checked via gel electrophoresis

analysis and melting curve analysis during quantitative real-time PCR.

The relative quantification of all transcripts was performed using qRT-PCR with

the Bio-Rad CFX96 Real-Time PCR System (Bio-Rad Laboratories, USA). Real-time

quantitative PCR was performed with SsoFast EvaGreen Supermix (Bio-Rad

Laboratories, USA). A total volume of 10 μl containing 5 μl of SsoFast EvaGreen

Supermix, 3 μl of RNA-free water, 0.5 μl of forward primer, 0.5 μl of reverse primer,

and 1 μl of template cDNA was used. PCR amplification was carried out as follows:

denaturation at 95 ℃ for 30 s, followed by 40 cycles at 95 ℃ for 5 s and a specific

annealing temperature of 55 ℃ for 30 s. The $2^{-\triangle\triangle CT}$ method was used to evaluate the

mRNA expression levels (25).

Statistical analysis

The statistical analysis was conducted using the SPSS software (SPSS, IBM,

version 20). The standard error of the means (SEM) was obtained from the analysis of

variance (ANOVA) that was conducted. The significance of the differences among

means was determined using one-way ANOVA and compared using Dunnett's

multiple range tests. The results were considered statistically significant at $P>0.05$.

Results

Jejunum morphology

In this study, all birds purchased from the local farm at the age of 90 days were

no different from the normal farmed birds. The jejunum morphology analysis was

performed in week 8 (**Table 4; Fig. 1-I**). Compared to control, the intestinal wall

thickness was significantly increased in the MIC groups; it was approximately 79 μm

and thus extremely thicker than that in the other groups ($P<0.01$). Meanwhile, The

villus height showed a significant increase both in MIC and IPF groups. Group MH

and ML showed a clear enhancement of the intestinal wall, villus height and crypt

depth. In addition, the intestinal wall thickness was 43 μm in the IPF groups, similar

to the control group (41 μm), and in contrast to the increased villus height, the crypt

depth was decreased.

Yolk cholesterol and fatty acids

The results of the yolk cholesterol and triglycerides determination are shown n

Table 5. In this study, compared to the control, the groups with lower amount of

additive showed a decrease in the yolk cholesterol content, but there were no effects

on the triglycerides. A decrease in yolk cholesterol was observed in groups ML and IL

compared to the control ($P<0.01$), while the 5% groups (MH and IH) showed no

significant change. In addition, the most significant yolk cholesterol decline was

observed in group ML.

In our study, both the MIC and IPF contained few lipids (**Table 1**). The

conjugated linoleic acid content was in agreement with the results of Yang et al. (2):

68.64% in IPF oil and 72.21% in MIC oil (**Table 6**). The composition of the yolk fatty

acids is shown in **Table 7.**

As shown in **Table 7,** the n6/n3 (n3-polyunsaturated fatty acids/

n6-polyunsaturated fatty acids) ratio was significantly higher in IPF groups than in the

control ($P<0.01$). Conversely, compared to the control, the decrease in the n6/n3 ratio

in group MH was significant ($P<0.01$), and group MH also showed an increased n3

(EPA: eicosapentaenoic acid; DHA: docosahexaenoic acid) content and decreased n6

content, which are beneficial to the absorption of unsaturated fatty acid (26).

Egg production and quality

The effects of MIC and IPF dietary supplements on quail eggs were measured

(**Table 8**). The yolk weight, shell thickness, Haugh unit and diameter ratio were

similar among the groups. The yolk weight was approximately 0.29-0.33% of the total

weight of the egg, the shell thickness was 0.16-0.18 mm, the Haugh unit was 79-84,

and the diameter ratio was 0.77-0.82.

Egg production in the MIC groups generally increased during the whole

experiment period compared to the control (**Table 9**). Group ML achieved a

significant increase in egg mass and laying ratelaying rate on average, higher than

control group during the same period of time. The average egg mass in group ML was

9.77 g/d/bird, which was significantly higher than that in the other groups. In contrast,

the egg mass decreased in the IPF groups; group IH (5% IPF added) exhibited a

significant production decrease that was more clear in the last week compared to the

control (6.06 g/d/bird; $P<0.01$). Meanwhile, laying ratelaying rate in the MIC groups

generally increased during the whole experiment period compared to the control, and

decreased laying ratelaying rate was observed in the IPF groups (**Table 9**). Similarly,

group ML achieved a significant increase in laying ratelaying rate, at 87.7% on

average, which is approximately 12% above the control. In group ML, nearly 44 eggs

per day were produced on average, while the control group produced 38 eggs per day

during the same period of time.

Genes

The relative mRNA expression level changes in SRBEP-1, SREBP-2, ATGL,

APOVLDL-II, which are involved in fatty acid and cholesterol biosynthesis were

measured. The relative mRNA expression of group ML and IL were showed in **Fig.**

1(II, III, IV). Group ML and group IL are showed similar relative mRNA expression

changes. Compared to the control, both ML and IL groups showed significant

up-regulation of APOVLDL-II in liver (**Fig. 1-III**, $P<0.01$). SREBP-1 was

up-regulated in the fat pad ($P<0.01$) but did not significantly change in the liver, while

SREBP-2 was significantly down-regulated in the fat pad and up-regulated in the liver.

However, the up-regulation of ATGL was found in ML.

Discussion

In the poultry industry, the wide use of antibiotics usually results in a thinner

intestinal wall, smaller total villus area, and shorter villus height and crypt depth (27).

In this study, all birds purchased from the local farm at the age of 90 days were no

different from the normal farmed birds. The intestinal wall thickness was significantly

increased in the MIC groups; it was approximately 79 μm and thus extremely thicker

than that in the other groups ($P<0.01$). The villus height of both MIC and IPF groups

showed a significant increase, and this increase may related to the activation of

intestinal villi growth (28). MIC groups showed a clear enhancement of the intestinal

wall, villus height, suggesting that the birds of MIC groups have better digestive and

absorptive capacity, and the appropriate amount of MIC additive was beneficial for

the intestinal health of laying quails. However, IPF showed no effect on the intestinal

wall, the intestinal wall thickness was 43 μm in the IPF groups, similar to the control

group (41 μm).

MIC and IPF supplementation also changed the cholesterol content of egg.

Cholesterol is important for many physiological functions (29-31), and as a

triglyceride, cholesterol is also an essential material for yolk formation and egg

production. However, high cholesterol intake must be avoided in our diet because

excess cholesterol is related to cardiovascular disease and nonalcoholic steatohepatitis

which contributing to hepatocellular carcinoma, thus increasing the risk to human

health (32-34). Moreover, cholesterol content is used in the evaluation of egg quality

in the poultry industry. From the perspective of human health, eggs with lower

cholesterol are considered to have superior quality, and more attention is currently

paid to reducing egg cholesterol (35, 36). In this study, compared to the control, the

groups with lower amount of additive showed a decrease in the yolk cholesterol

content, but there were no effects on the triglycerides. The most significant yolk

cholesterol decline was observed in group ML. These results indicate that both MIC

and IPF were effective in terms of affecting the yolk cholesterol, and the optimal

addition is1%.

Ouyang et al. found that a different oil supplement given to hens was able to

change the cholesterol and polyunsaturated fatty acid content in the yolk (37).

Furthermore, in a study conducted by Aydin et al., dietary conjugated linoleic acid (CLA) added to hen diets significantly changed the fatty acid composition of the yolk (38). In our study, both the MIC and IPF contained few lipids (**Table 1**). The conjugated linoleic acid content was in agreement with the results of Yang et al (2), and this small amount of CLA might be involved in the change of bird partly (**Table 6**).

n-3 polyunsaturated fatty acids and n-6 polyunsaturated fatty acids is indispensable fatty acid for human(39). It is well known that a lower n6/n3 ratio is better according to the guidelines of the WHO regarding healthy food for humans (26). As shown in **Table 7**, compared to the control, the decrease in the n6/n3 ratio in group A was significant ($P<0.01$), and group MH also showed an increased n3 (EPA: eicosapentaenoic acid; DHA: docosahexaenoic acid) content and decreased n6 content, which are beneficial to the absorption of unsaturated fatty acid (26). Conversely, the n6/n3 ratio was significantly higher in IPF groups than in the control ($P<0.01$).

As we know that both fatty acid and cholesterol metabolism are involved in the regulation of egg formation (40, 41), and lipid mobilization from the adipose tissue is one of the main sources supporting the onset of egg production. As a result of lipid mobilization, the increase in the fat pad is the most intuitive indication of a supply of raw material available for egg production (42). The relative mRNA expression level changes in SRBEP-1, SREBP-2, ATGL, APOVLDL-II, which are involved in fatty acid and cholesterol biosynthesis (43-45), were measured at the end of the study.

Adipose triglyceride lipase (ATGL) is highly expressed in adipose tissue and

catalyzes the initial step of triglyceride hydrolysis (46). In a study conducted by Chen

et al., G0/G1 switch gene 2 (G0S2) overexpression carried out the inhibition of ATGL

expression, which resulted in a delayed laying onset and reduced egg production (40).

Sterol regulatory element-binding protein-1 and -2 (SREBP-1 and SREBP-2) are

SREBPs encoding genes that mainly regulate the homeostasis of lipids and

cholesterol, respectively (43, 47). SREBP-1 up-regulation preferentially enhances

fatty acid and triglyceride synthesis, while increased SREBP-2 expression promotes

cholesterol synthesis (48-50). Apo very low density lipoprotein II (APOVLDL-II) is

related to the secretion of VLDL, which plays a pivotal role in egg formation (19).

The described changes in the fat pad and ATGL expression found in the study by

Chen et al (40) were also observed in our study (**Fig. 1**-III).

In this study, the interclavicular fat pat weight significantly decreased in both the

MIC-added and IPF-added groups. In group ML, for example, while the fat pad

decreased, the ATGL was up-regulated, that is, the lipid supply for egg formation was

enhanced. Meanwhile, as the results show in **Fig. 1**(III, IV), we found that fatty acid

synthesis was strengthened in the fat pad with the up-regulation of SREBP-1 and that

the synthesis and transport of cholesterol was enhanced in the liver with the

up-regulation of SREBP-2 and APOVLDL-II. In addition, the described jejunum

morphology results indicate that the birds in group ML had better intestine health,

which allowed better digestion and absorption. These changes indicate that the birds

in group ML have enhanced energy requirement and material supplementation for egg

production during the feeding period (**Fig. 2**). Hence, in the last week, MIC promoted

the egg quantity, but the single egg weight was lighter due to the fixed diet, just as the results showed in group ML. These results indicate that while the raw-material supplement supplied for the production of eggs was enhanced in the MIC groups, with an improvement in product performance, the fixed diet cannot meet the requirements of gradually increasing performance.

Moreover, the same tendency in gene expression was found in both groups ML and IL, combined with the change in the fat pad, we assumed that IPF facilitated several processes involved in egg production, although it ultimately resulted in some damage to body health. However, based on the results, we believe that *candida utilis* could help to reduce the disadvantage of IPF which was harmful to the laying quails and that MIC, a product of fermentation, was beneficial to intestinal health and could enhance the production capacity of laying quails. Meanwhile, the groups with a lower MIC amount (1%) showed that these amounts were more effective in promoting laying ratelaying rate and egg mass. Furthermore, the results also suggest that through solid-state fermentation, IPF, which is not suitable for direct feeding, could be a good feed additive.

Overall, our study showed that MIC improved egg production and quality in laying quails. This study is showing the application of *Idesia polycarpa* fruit and *candida utilis* in feeding. MIC at 1% is the most efficient dosage not only in terms of an increase in egg production but also a decrease in yolk cholesterol. Meanwhile, adding 5% MIC could result in a higher EPA and DHA content and a better n3/n6 ratio. Furthermore, MIC was beneficial to the health of the quail's intestine and

316   reduced the potential negative effects of industrial feeding.

■ Acknowledgements

This study was funded by Miaozi project in Science and Technology Innovation Program of Sichuan Province (2015MZGC001); Sichuan Science and Technology plan project (2016NZYZF0102).

Notes

The authors declare no competing financial interest.

■ REFERENCE

1. Wang QY, Li X, Du KF, Yao S, Song H. Conjugated Linoleic Acid Production by Alkali Isomerization of Linoleic Acid from Idesia polycarpa Maxim. var. vestita Diels Oil. Asian Journal of Chemistry. 2013;25(7):3744-8.

2. Yang F-X, Su Y-Q, Li X-H, Zhang Q, Sun R-C. Preparation of biodiesel from Idesia polycarpa var. vestita fruit oil. Industrial Crops and Products. 2009;29(2-3):622-8.

3. Ye Y, Jia RR, Tang L, Chen F. In Vivo Antioxidant and Anti-Skin-Aging Activities of Ethyl Acetate Extraction from Idesia polycarpa Defatted Fruit Residue in Aging Mice Induced by D-Galactose. Evidence-based complementary and alternative medicine : eCAM. 2014;2014:185716. PubMed PMID: 24971146. Pubmed Central PMCID: 4058274.

4. Lee M, Lee HH, Lee JK, Ye SK, Kim SH, Sung SH. Anti-adipogenic activity of compounds isolated from Idesia polycarpa on 3T3-L1 cells. Bioorganic & medicinal chemistry letters. 2013 Jun 1;23(11):3170-4. PubMed PMID: 23628332.

5. Ye Y, Tang X-S, Chen F, Tang L. Optimization of Phenolics Extracted from Idesia polycarpa Defatted Fruit Residue and Its Antioxidant and Depigmenting Activity In Vitro and In Vivo. Evidence-Based Complementary and Alternative Medicine. 2014;2014:1-12.

6. Thom C, Church MB. Aspergillus flavus, A. oryzae, and Associated Species. American Journal of Botany. 1921;8(2):103-26.

7. Curtis RS. Feeding fermented cotton-seed meal to hogs. 1909.

8. Hati S, Patel N, Mandal S. Comparative Growth Behaviour and Biofunctionality of Lactic Acid Bacteria During Fermentation of Soy Milk and Bovine Milk. Probiotics and antimicrobial proteins. 2018 Jun;10(2):277-83. PubMed PMID: 28455690.

9. Wang J, Cao F, Su E, Zhao L, Qin W. Improvement of Animal Feed Additives of Ginkgo Leaves through Solid-state Fermentation using Aspergillus niger. International journal of biological sciences. 2018;14(7):736-47. PubMed PMID: 29910684. Pubmed Central PMCID: 6001676.

10. Dwivedi M, Yajnanarayana VK, Kaur M, Sattur AP. Evaluation of anti nutritional factors in fungal fermented cereals. Food Science and Biotechnology. 2015;24(6):2113-6.

11. Li C-Y, Lu J-J, Wu C-P, Lien T-F. Effects of probiotics and bremelain fermented soybean meal

replacing fish meal on growth performance, nutrient retention and carcass traits of broilers. Livestock
Science. 2014;163:94-101.

12. Karásková K, Suchý P, Straková E. Current use of phytogenic feed additives in animal nutrition: a
review. Czech Journal of Animal Science. 2016;60(No. 12):521-30.

13. Hong KJ, Lee CH, Kim SW. Aspergillus oryzae GB-107 fermentation improves nutritional quality of
food soybeans and feed soybean meals. Journal of Medicinal Food. 2004;7(4):430-5.

14. Buerth C, Tielker D, Ernst JF. Candida utilis and Cyberlindnera (Pichia) jadinii: yeast relatives with
expanding applications. Applied microbiology and biotechnology. 2016 Aug;100(16):6981-90. PubMed
PMID: 27357226.

15. Boze HLN, Moulin G, Galzy P. Production of Food and Fodder Yeasts. Critical Reviews in
Biotechnology. 1992;12(1-2):65-86.

16. Lawford GR, Kligerman A, Williams T, Lawford HG. Production of high‐quality edible protein
from Candida yeast grown in continuous culture. Biotechnology & Bioengineering.
2010;21(7):1163-74.

17. Aba P, Eneasato C, Onah J. Effect of Quail Egg Pretreatment on the Lipid Profile and
Histomorphology of the Liver of Acetaminophen-Induced Hepatotoxicity in Rats. Journal of Applied
Biology and Biotechnology. 2016:034-8.

18. Ibukun EO, Oladipo GO. Lipidomic Modulation in Stressed Albino Rats Is Altered by Yolk and
Albumen of Quail (Coturnix japonica) Egg and Poultry Feed. Biochemistry Research International.
2016;2016(2):1-7.

19. . !!! INVALID CITATION !!!

20. Chah CC, Nelson RA, Carlson CW, Semeniuk G, Palmer IS, Hesseltine CW. Fungus-fermented
soybeans benefit the life cycle of Japanese quail (Coturnix coturnix japonica). Poultry Science.
1976;55(3):975-81.

21. Hong H, Abanto OD, Kim KH, Nam KT, Son JY, Jung WS, et al. The Effects of Dietary Soybean
Fermented with Aspergillus oryzae or Bacillus natto on Egg Production and Egg Lipid Composition in
Layer. Hangug chugsan sigpum haghoeji = Korean journal for food science of animal resources.
2010;30(4):609-16.

22. Kanyinji F, Moonga M. Effects of replacing maize meal with rumen filtrate-fermented cassava
meal on growth and egg production performance in Japanese quails (Cortunix japonica). Journal of
Advanced Veterinary & Animal Research. 2014;1(3):100-6.

23. Alagawany M, Farag MR, El-Hack MEA, Dhama K. The Practical Application of Sunflower Meal in
Poultry Nutrition. Advances in Animal & Veterinary Sciences. 2015;19:323-7.

24. Ulberth F. One-step extraction/methylation method for determining the fatty acid composition of
processed foods. Journal of the American Oil Chemists' Society. 1992;69(2):174-7.

25. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative
PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25(4):402-8.

26. Kris-Etherton PM, Innis S, Assocition AD, Canada DO. Position of the American Dietetic
Association and Dietitians of Canada: dietary fatty acids. Journal of the American Dietetic Association.
2007;107(9):1599-611.

27. Miles RD, Butcher GD, Henry PR, Littell RC. Effect of antibiotic growth promoters on broiler
performance, intestinal growth parameters, and quantitative morphology. Poultry Science.
2006;85(3):476-85.

28. Afsharmanesh M, Sadaghi B. Effects of dietary alternatives (probiotic, green tea powder, and

Kombucha tea) as antimicrobial growth promoters on growth, ileal nutrient digestibility, blood parameters, and immune response of broiler chickens. Comparative Clinical Pathology. 2013;23(3):717-24.

29.   Naviglio D, Gallo M, Grottaglie LL, Scala C, Ferrara L, Santini A. Determination of cholesterol in Italian chicken eggs. Food chemistry. 2012;132(2):701-8.

30.   Schroeder F, Jefferson JR, Kier AB, Knittel J, Scallen TJ, Wood WG, et al. Membrane cholesterol dynamics: cholesterol domains and kinetic pools. Proceedings of the Society for Experimental Biology & Medicine Society for Experimental Biology & Medicine. 1991;196(3):235-52.

31.   Simons K, Ikonen E. How cells handle cholesterol. Science. 2000;290(5497):1721-6.

32.   Liang JQ, Teoh N, Xu L, Pok S, Li X, Chu ESH, et al. Dietary cholesterol promotes steatohepatitis related hepatocellular carcinoma through dysregulated metabolism and calcium signaling. Nature communications. 2018 Oct 26;9(1):4490. PubMed PMID: 30367044. Pubmed Central PMCID: 6203711.

33.   Ahn J, Lim U, Weinstein SJ, Schatzkin A, Hayes RB, Virtamo J, et al. Prediagnostic Total and High-Density Lipoprotein Cholesterol and Risk of Cancer. Cancer Epidemiology Biomarkers & Prevention. 2009;18(11):2814-21.

34.   Detection EPO. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). Jama. 2001;285(19):2486-97.

35.   Lamas A, Anton X, Miranda JM, Roca-Saavedra P, Cardelle-Cobas A, Ibarra IS, et al. Technological Strategies for the Development of Egg-Derived Products with Reduced Content of Cholesterol. Food and Bioprocess Technology. 2016;9(1):81-90.

36.   Mattioli S, Bosco AD, Martino M, Ruggeri S, Marconi O, Sileoni V, et al. Alfalfa and flax sprouts supplementation enriches the content of bioactive compounds and lowers the cholesterol in hen egg. Journal of Functional Foods. 2016;22:454-62.

37.   Ouyang K, Wang W, Xu MS, Jiang Y, Sangguan X. Effects of different oils on the production performances and polyunsaturated fatty acids and cholesterol level of yolk in hens. Asian Australasian Journal of Animal Sciences. 2004;17(17):843-7.

38.   Aydin R, Cook ME. The effect of dietary conjugated linoleic acid on egg yolk fatty acids and hatchability in Japanese quail. Poultry Science. 2004;83(12):2016-22.

39.   Bird JK, Calder PC, Eggersdorfer M. The Role of n-3 Long Chain Polyunsaturated Fatty Acids in Cardiovascular Disease Prevention, and Interactions with Statins. Nutrients. 2018;10(6).

40.   Chen PR, Shin S, Choi YM, Kim E, Han JY, Lee K. Overexpression of G0/G1 Switch Gene 2 in Adipose Tissue of Transgenic Quail Inhibits Lipolysis Associated with Egg Laying. International journal of molecular sciences. 2016 Mar 15;17(3):384. PubMed PMID: 26999108. Pubmed Central PMCID: 4813241.

41.   Wu J, Fu W, Huang Y, Ni Y, Zhao R. Kisspeptin-10 enhanced egg production in quails associated with the increase of triglyceride synthesis in liver. Asian-Australasian journal of animal sciences. 2013 Aug;26(8):1080-8. PubMed PMID: 25049888. Pubmed Central PMCID: 4093219.

42.   Yang S, Suh Y, Choi YM, Shin S, Han JY, Lee K. Loss of Fat with Increased Adipose Triglyceride Lipase-Mediated Lipolysis in Adipose Tissue During Laying Stages in Quail. Lipids. 2013;48(1):13-21.

43.   Brown MS, Goldstein JL. The SREBP Pathway: Regulation of Cholesterol Metabolism by Proteolysis of a Membrane-Bound Transcription Factor. Cell. 1997;89(3):331–40.

44.   Hermansson M, Hanninen S, Hokynar K, Somerharju P. The PNPLA-family phospholipases involved in glycerophospholipid homeostasis of HeLa cells. Biochimica et biophysica acta. 2016

Sep;1861(9 Pt A):1058-65. PubMed PMID: 27317427.

45.   Zheng W, Mast N, Saadane A, Pikuleva IA. Pathways of cholesterol homeostasis in mouse retina

responsive to dietary and pharmacologic treatments. Journal of Lipid Research. 2014;56(1):81-97.

46.   Zimmermann R, Zechner R. Fat Mobilization in Adipose Tissue Is Promoted by

AdiposeTriglyceride Lipase. Science. 2004;306(5700):1383-6.

47.   Eberlé D, Hegarty B, Bossard P, Ferré P, Foufelle F. SREBP transcription factors: master regulators

of lipid homeostasis. Biochimie. 2004;86(11):839-48.

48.   Dong X, Xu H, Mai K, Xu W, Zhang Y, Ai Q. Cloning and characterization of SREBP-1 and

PPAR-alpha in Japanese seabass Lateolabrax japonicus, and their gene expressions in response to

different dietary fatty acid profiles. Comparative biochemistry and physiology Part B, Biochemistry &

molecular biology. 2015 Feb;180:48-56. PubMed PMID: 25448051.

49.   Li X, Li Y, Yang W, Xiao C, Fu S, Deng Q, et al. SREBP-1c overexpression induces triglycerides

accumulation through increasing lipid synthesis and decreasing lipid oxidation and VLDL assembly in

bovine hepatocytes. The Journal of steroid biochemistry and molecular biology. 2014 Sep;143:174-82.

PubMed PMID: 24565561.

50.   Park J, Yeom M, Hahm DH. Fucoidan improves serum lipid levels and atherosclerosis through

hepatic SREBP-2-mediated regulation. Journal of pharmacological sciences. 2016 Jun;131(2):84-92.

PubMed PMID: 27094367.

Tables

Table 1. The Main Nutrients of MIC, IPF and Basal diet (%)

	MIC	IPF	Basal diet
Crude protein	16.4	8.49	21
Crude lipid	6.34	13.34	2.5
Moisture content	5.04	7.20	14
Ash	4.62	4.49	12
Free Amino acid	6.81	5.01	
Lysine	0.13	0.09	
Methionine	0.11	0.05	0.46
Phenylalanine	0.36	0.27	
Threonine	0.31	0.23	
Leucine	0.57	0.44	
Isoleucine	0.34	0.26	
Arginine	0.35	0.15	
Histidine	0.30	0.23	
Valine	0.42	0.33	

MIC: means Mixture of *Idesia polycarpa* defatted fruit and candida utilis**; IPF**: means *Idesia polycarpa* defatted fruit.

Table 2. Daily Diet Arrangement of MIC, IPF and Control Groups (g)

	Week 0 to 2			Week 3 to 8		
	Basal diet	MIC	IPF	Basal diet	MIC	IPF
MH	1000	0.5	-	950	50	-
ML	1000	0.5	-	990	10	-
IH	1000	0.5	-	950	-	50
IL	1000	0.5	-	990	-	10
Control	1000	0.5	-	1000	-	-

[1]**MIC**: means Mixture of *Idesia polycarpa* defatted fruit and candida utilis**; IPF**: means *Idesia polycarpa* defatted fruit.

[2]**MH**: Basal diet supplemented with 5% MIC, **ML**: basal diet supplemented with 1% MIC, **IH**: basal diet supplemented with 5% IPF, **IL**: basal diet supplemented with 1% IPF, **Control**: Control group, base diet only.

[4]Superscripts **a, b, c, d, e**: Means within the same superscripts without the same row are significantly different (P<0.05).

Table 3. Primer Sequences for the mRNA Expression Analysis of Genes

Genes	primer sequence (5'-3')	product size (bp)	Gene Bank
APOVLDL-II	F: CAGTTCTTGCTGGATGTTTCCCAGAC	430	S82591.1
	R: CAATGGCCAAGTCATTCAGGAGGA		
SREBP-1	F: CTACCGCTCATCCATCAACG	181	NC_029529.1
	R: CTGCTTCAGCTTCTGGTTGC		
SREBF-2	F: CCCAGAACAGCAAGCAAGG	108	XM_416222
	R: GCGAGGACAGGAAAGAGAGTG		
ATGL	F: CAGCAGGACGGTTGGGTATTTC	154	GQ221783.1
	R: CCACGCAAGGTTGGAGGTATCA		
β-actin	F: TGATGGTTGGTATGGGTCAGAAAG	92	NC_029529
	R: ATGTTCAATGGGGTATTTCAAGGT		

Table 4. Jejunum Morphology Analysis of Laying Quails Fed a Basal Diet, MIC Added Diet or IPF Added Diet

	MIC		IPF				
	MH (5%)	ML (1%)	IH (5%)	IL (1%)	Control	SEM	*P*
Intestine wall thickness (μm)	78.5[a]	79.2[a]	43.7[b]	43.4[b]	41.3[b]	6.09	<0.01
villous height (μm)	835[b]	908[a]	818[bc]	801[c]	610[d]	26.9	<0.01
Crypt depth (μm)	112.8[b]	189.5[a]	72.9[d]	77.0[d]	91.5[c]	4.35	<0.01
V/C	7.40[b]	4.79[d]	11.22[a]	10.41[a]	6.67b[c]	0.50	<0.01

[1]N=9.

[2]**MIC**: means Mixture of *Idesia polycarpa* defatted fruit and candida utilis**; IPF**: means *Idesia polycarpa* defatted fruit.

[3]. **V/C** means the ratio of villous length/crypt depth.

[4]**MH**: Basal diet supplemented with 5% MIC, **ML**: basal diet supplemented with 1% MIC, **IH**: basal diet supplemented with 5% IPF, **IL**: basal diet supplemented with 1% IPF, **Control**: Control group, base diet only.

[4]Superscripts **a, b, c, d, e**: Means within the same superscripts without the same row are significantly different (P<0.05).

Table 5. Yolk Cholesterol and Triglyceride Content of Laying Quails Fed a Basal Diet, MIC Added Diet or IPF Added Diet

	MIC		IPF				
	MH	ML	IH	IL	Control	SEM	*P*-value
	(5%)	(1%)	(5%)	(1%)			
Yolk Weight							
(g)	3.61	3.10	3.60	3.29	3.39	0.39	0.73
Cholesterol							
(mg/g yolk)	13.1^{ab}	12.1^{b}	14.0^{a}	11.2^{b}	14.5^{a}	0.25	<0.01
(mg/ yolk)	46.54^{ab}	37.0^{c}	48.5^{ab}	40.3^{b}	49.1^{a}	4.73	<0.01
Triglyceride							
(mg/g yolk)	183	175	197	183	190	8.57	0.10
(mg/ yolk)	660.2^{a}	536.0^{b}	682.9^{a}	661.4^{b}	644.4^{a}	69.2	0.01

[1]N=9.

[2]**MIC**: means Mixture of *Idesia polycarpa* defatted fruit and candida utilis**; IPF**: means *Idesia polycarpa* defatted fruit.

[3]**MH**: Basal diet supplemented with 5% MIC, **ML**: basal diet supplemented with 1% MIC, **IH**: basal diet supplemented with 5% IPF, **IL**: basal diet supplemented with 1% IPF, **Control**: Control group, base diet only.

[4]Superscripts **a, b, c, d, e**: Means within the same superscripts without the same row are significantly different (P<0.05).

Table 6. Fatty Acid Composition of MIC Oil and IPF Oil (%)

Fatty Acid		MIC	IPF
Palmitic	C16:0	15.51	17.36
Stearic	C18:0	1.79	1.66
Oleic	C18:1	3.5	4.54
Palmitoleic	C16:1	6.41	7.55
Linoleic	C18:2	72.21	68.64

[1]**MIC**: means Mixture of *Idesia polycarpa* defatted fruit and candida utilis**; IPF**: means *Idesia polycarpa* defatted fruit.

Table 7. Fatty Acid Composition of Yolk Oil extracted from each groups

Fatty acid		Control	MH	ML	IH	IL
Myristic	C14:0	0.45[b]	0.57[a]	0.54[a]	0.43[b]	0.39[b]
Palmitic	C16:0	29.53	30.55	31.88	30.92	32.72
Palmitoleic	C16:1;n-7	2.16[c]	3.64[a]	2.39[bc]	2.82[b]	2.20[bc]
Stearic	C18:0	12.8[a]	11.6[b]	10.5[c]	9.23[d]	10.6[c]
Oleic	C18:1;n-9	33.8[b]	35.3[b]	34.6[b]	37.2[a]	32.2[c]
Linoleic	C18:2;n-6	17.1[b]	14.1[c]	16.7[b]	16.8[b]	18.2[a]
α-Linolenic	C18:3;n-3	0.22[a]	0.21[a]	0.25[a]	0.17[b]	0.22[a]
Arachidonic	C20:4;n-6	2.56[a]	2.38[a]	1.88[b]	1.65[b]	1.64[b]
Eicosapentaenoic	C20:5;n-3	0.27[b]	0.37[a]	0.29[ab]	0.14[c]	0.25[b]
Docosahexaenoic	C22:6;n-3	0.89[b]	1.02[a]	0.76[c]	0.51[d]	0.48[d]
SFA		42.8[a]	42.7[a]	42.9[a]	40.6[b]	43.7[a]
MUFA		36.0[c]	38.9[b]	37.0[bc]	40.0[a]	34.4[d]
PUFA		21.0[a]	18.1[c]	19.8[b]	19.3[b]	20.8[ab]
n-3PUFA		1.38[b]	1.60[a]	1.30[b]	0.82[c]	0.95[bc]
n-6PUFA		19.6[a]	16.5[c]	18.5[b]	18.4[b]	19.9[a]
n6/n3		14.2[c]	10.3[d]	14.3[c]	22.5[a]	20.9[b]

[1]N=3.

[2]**MIC**: means Mixture of *Idesia polycarpa* defatted fruit and candida utilis**; IPF**: means *Idesia polycarpa* defatted fruit.

[3]**MH**: Basal diet supplemented with 5% MIC, **ML**: basal diet supplemented with 1% MIC, **IH**: basal diet supplemented with 5% IPF, **IL**: basal diet supplemented with 1% IPF, **Control**: Control group (base diet);

[4]Superscripts **a, b, c, d, e** Means within the same superscripts without the same row are significantly different (P<0.05).

[5]**SFA**: short chain fatty acid; **MUFA**: monounsaturated fatty acid; **PUFA**: polyunsaturated fatty acids.

[6]All results in percentage terms.

Table 8. Egg Quality of Laying Quails Fed a Basal Diet, MIC Added Diet or IPF Added Diet

| | MIC | | IPF | | | | |
| | MH | ML | IH | IL | Control | SEM | P-VALUE |
	(5%)	(1%)	(5%)	(1%)			
Yolk Weight (%)							
initial	0.30	0.30	0.31	0.30	0.31	0.187	0.980
last week	0.31	0.30	0.33	0.31	0.29	0.029	0.935
average	0.31	0.30	0.32	0.31	0.30	0.119	0.754
Shell Thickness(mm)							
Initial	0.16	0.16	0.16	0.17	0.16	0.007	0.785
Last week	0.17	0.18	0.19	0.18	0.18	0.008	0.285
Average	0.18	0.18	0.18	0.18	0.17	0.008	0.563
Haugh Unit							
Initial	80.59	79.84	80.75	79.59	79.15	2.760	0.827
Last week	79.25	81.18	79.86	80.42	80.78	1.881	0.924
Average	82.92	81.51	84.75	81.26	80.81	1.437	0.458
Diameter Ratio							
Initial	0.77	0.80	0.82	0.79	0.80	0.346	0.805
Last week	0.77	0.78	0.78	0.78	0.79	0.082	0.285
Average	0.78	0.77	0.79	0.78	0.77	0.007	0.269

[1]N=9.

[2]**MIC**: means Mixture of *Idesia polycarpa* defatted fruit and candida utilis**; IPF**: means *Idesia polycarpa* defatted fruit.

[3]**MH**, basal diet supplemented with 5% MIC; **ML**, basal diet supplemented with 1% MIC; **IH**, basal diet supplemented with 5% IPF; **IL**, basal diet supplemented with 1% IPF; **Control**, Control group (base diet).

[4]**Initial**: Mean of week 0 to 2; **Last week**: Mean of week 8; **Average**: Mean of week 3 to 8.

Table 9. Egg Mass and Egg Production of Laying Quails Fed a Basal Diet, MIC Added Diet or IPF Added Diet

	MIC		IPF				
	MH (5%)	ML (1%)	IH (5%)	IL (1%)	Control	SEM	*P*-value
Egg mass (g/d/bird)							
Initial	9.44[a]	9.41[a]	9.31[a]	9.14[a]	9.18[a]	0.30	0.88
Last week	9.18[a]	9.31[a]	6.04[c]	7.82[b]	9.27[ab]	0.30	<0.01
Average	9.41[a]	9.77[a]	6.06[d]	7.86[c]	8.87[b]	0.38	<0.01
Laying rate (%)							
Initial	81.0[a]	81.3[a]	80.7[a]	80.7[a]	79.3[a]	2.57	0.97
Last week	80.1[b]	90.2[a]	56.3[e]	68.1[d]	78.4[c]	2.91	<0.01
Average	82.2[a]	87.7[a]	62.4[c]	72.3[b]	76.0[b]	3.15	<0.01

[1]N=50.

[2]**MIC**: means Mixture of *Idesia polycarpa* defatted fruit and candida utilis; **IPF**: means *Idesia polycarpa* defatted fruit.

[3] **MH**, basal diet supplemented with 5% MIC; **ML**, basal diet supplemented with 1% MIC; **IH**, basal diet supplemented with 5% IPF; **IL**, basal diet supplemented with 1% IPF; **Control**, Control group (base diet).

[4]Superscripts **a, b, c, d, e** Means within the same superscripts without the same row are significantly different (P<0.05).

[5]**Initial**: Mean of week 0 to 2; **Last week**: Mean of week 8; **Average**: Mean of week 3 to 8;

Figures

Fig. 1

Fig. 1 Results of jejunum morphology, fat pad, and relative mRNA expression.

I: Tissue sections of jejunum. **ML**: the Tissue section figure of group ML; **IL**: the Tissue section figure of group IL; collected in 40×amplifications and 200×amplifications respectively.

II: Ratio of Fat Pad to Body Weight（%）. **MH**, basal diet supplemented with 5% MIC; **ML**, basal diet supplemented with 1% MIC; **IH**, basal diet supplemented with 5% IPF; **IL**, basal diet supplemented with 1% IPF; **Control**, Control group (base diet). ******Means significant relative to control (P<0.01).

III: Relative mRNA expression of 1% MIC added group (**ML**) and 1%IPF added group (**IL**) in fat pad. ******Means significant relative to control (P<0.01); *****Means significant relative to Control (P<0.05).

IV: Relative mRNA expression of 1% MIC added group (**ML**) and 1%IPF added group (**IL**) in liver. ******Means significant relative to control (P<0.01).

Fig. 2

Fig. 2 The influence of MIC in laying quails. Compared to the control, the intestine statues of laying quails was significantly changed with the addition of MIC, as well as the relative mRNA expression in the fat pad and liver, ultimately, these changes result in an enhanced egg production.