1   **Natural strain variation reveals diverse biofilm regulation in squid-colonizing *Vibrio***
2   ***fischeri***
3
4

5   Ella R. Rotman[1,†], Katherine M. Bultman[2,†], John F. Brooks II[1,4], Mattias C. Gyllborg[1], Hector L.
6   Burgos[2], Michael S. Wollenberg[3], Mark J. Mandel[1,2,*]
7
8
9   [1] Department of Microbiology-Immunology, Northwestern University Feinberg School of
10   Medicine, Chicago, IL USA
11
12   [2] Department of Medical Microbiology and Immunology, University of Wisconsin-Madison,
13   Madison, WI USA
14
15   [3] Department of Biology, Kalamazoo College, Kalamazoo, MI USA
16
17   [4] Current address: Department of Immunology, The University of Texas Southwestern Medical
18   Center, Dallas, TX USA
19
20   [†] Authors contributed equally
21
22
23   Short title: *Vibrio fischeri* biofilm regulatory evolution
24
25   Keywords: Biofilm, phosphorelay, RscS, BinK, Vibrio fischeri, Aliivibrio fischeri
26
27
28   * Correspondence to:
29
Mark J. Mandel
University of Wisconsin-Madison
Department of Medical Microbiology and Immunology
1550 Linden Drive
Madison, WI 53706
Phone: (608) 261-1170
Fax: (608) 262-8418
Email: mmandel@wisc.edu

ABSTRACT

The mutualistic symbiont *Vibrio fischeri* builds a symbiotic biofilm during colonization of squid

hosts. Regulation of the exopolysaccharide component, termed Syp, has been examined in

strain ES114, where production is controlled by a phosphorelay that includes the inner

membrane hybrid histidine kinase RscS. Most strains that lack RscS or encode divergent RscS

proteins cannot colonize a squid host unless RscS from a squid symbiont is heterologously

expressed. In this study, we examine *V. fischeri* isolates worldwide to understand the landscape

of biofilm regulation during beneficial colonization. We provide a detailed study of three distinct

evolutionary groups of *V. fischeri* and find that while the RscS-Syp biofilm pathway is required in

one of the groups, two other groups of squid symbionts require Syp independent of RscS.

Mediterranean squid symbionts, including *V. fischeri* SR5, colonize without an RscS homolog

encoded in their genome. Additionally, Group A *V. fischeri* strains, which form a tightly-related

clade of Hawaii isolates, have a frameshift in *rscS* and do not require the gene for squid

colonization or competitive fitness. These same strains have a frameshift in *sypE*, and we

provide evidence that this Group A *sypE* allele leads to an upregulation in biofilm activity. This

work thus describes the central importance of Syp biofilm in colonization of diverse isolates, and

demonstrates that significant evolutionary transitions correspond to regulatory changes in the

syp pathway.

IMPORTANCE

Biofilms are surface-associated, matrix-encased bacterial aggregates that exhibit enhanced

protection to antimicrobial agents. Previous work has established the importance of biofilm

formation by a strain of luminous *Vibrio fischeri* bacteria as the bacteria colonize their host, the

Hawaiian bobtail squid. In this study, expansion of this work to many natural isolates revealed

that biofilm genes are universally required, yet there has been a shuffling of the regulators of

those genes. This work provides evidence that even when bacterial behaviors are conserved,

dynamic regulation of those behaviors can underlie evolution of the host colonization

phenotype. Furthermore, this work emphasizes the importance of investigating natural diversity

as we seek to understand molecular mechanisms in bacteria.

INTRODUCTION

A fundamental question in studying host-associated bacterial communities is understanding how

specific microbial taxa assemble reproducibly in their host. Key insights into these processes

were first obtained by studying plant-associated microbes, and the discovery and

characterization of Nod factors in Rhizobia was valuable to understand how partner choice

between microbe and host could be mediated at the molecular level (1, 2). There are complex

communities in humans and other vertebrate animals, yet metagenomic and imaging analyses

of these communities have revealed striking reproducibility in the taxa present and in the spatial

arrangement of those taxa (3–5). Invertebrate animal microbiomes provide appealing systems in

which to study microbiome assembly in an animal host: the number of taxa are relatively small,

and examination and manipulation of these organisms have yielded abundant information about

processes underlying host colonization (6). For this work we focused on the binary symbiosis

between *Vibrio fischeri* and bobtail squids, including the Hawaiian bobtail squid, *Euprymna*

scolopes. Bobtail squid have an organ for the symbiont termed the light organ, and passage of

specific molecules between the newly-hatched host and the symbiont leads to light organ

colonization specifically by planktonic *V. fischeri* and not by other bacteria (7–9). The

colonization process involves initiation, accommodation, and persistence steps, resulting in light

organ crypt colonization by *V. fischeri*. Upon colonization of the squid light organ, bacteria

accumulate to high density and produce light. The bacterial light is modulated by the host to

camouflage the moonlight shadow produced by the nighttime foraging squid in a cloaking

process termed counter-illumination (10, 11). A diel rhythm leads to a daily clearing of 90-95%

of the bacteria from the crypts and regrowth of the remaining cells (12). However, the initial

colonization process, including biofilm-based aggregation on the host ciliated appendages,

occurs only in newly-hatched squid. This work examines regulation of biofilm formation in

diverse squid-colonizing *V. fischeri* strains.

In the well-studied *V. fischeri* strain ES114, biofilm formation is required to gain entry into the

squid host. RscS is a hybrid histidine kinase that regulates *V. fischeri* biofilm formation through

a phosphorelay involving the hybrid histidine kinase SypF and the response regulator and σ^{54}-

dependent activator SypG (13–15). This pathway regulates transcription of the symbiosis

polysaccharide (Syp) locus, which encodes regulatory proteins (SypA, SypE, SypF, and SypG),

glycosyltransferases, factors involved in polysaccharide export, and other biofilm-associated

factors (14, 16). The products of the ES114 *syp* locus direct synthesis and export of a biofilm

exopolysaccharide that is critical for colonization. Additional pathways have been identified to

influence biofilm regulation in ES114, including the SypE-SypA pathway and inhibition of biofilm

formation by BinK and HahK (17–21).

V. fischeri biofilm regulation is connected to host colonization specificity. In the Pacific Ocean,

the presence of *rscS* DNA is strongly correlated to the ability to colonize squid (22). As one

example, while the fish symbiont MJ11 encodes a complete *syp* locus, it lacks RscS and does

not robustly colonize squid. Heterologous expression of ES114 RscS in MJ11 activates the

biofilm pathway and is sufficient to enable squid colonization (22). Similarly, addition of ES114

RscS to *mjapo*.8.1--a fish symbiont that encodes a divergent RscS that is not functional for

squid colonization--allows the strain to colonize squid (22). RscS has also been shown to be

necessary for squid colonization in certain strains. In addition to ES114, interruption of *rscS* in

V. fischeri strains KB1A97 and MJ12 renders them unable to colonize squid. Previous

phylogenetic analysis revealed that ancestral *V. fischeri* do not encode *rscS*, and that it was

acquired once during the organism's evolution, likely allowing for an expansion in host range.

From this analysis, it was concluded that strains with *rscS* can colonize squid, with the only

exception being the fish symbionts that harbor the divergent RscS, including *mjapo*.8.1 (22).

There are similar *Vibrio*-squid associations worldwide, yet only *V. fischeri* and the closely-

related *Vibrio logei* have been isolated from light organs (23–26). Our 2009 study revealed that

although most symbionts have *rscS* DNA, there are Mediterranean *V. fischeri* (e.g., SR5) that

do not have *rscS* yet can colonize squid (22, 24, 27). This unexpected finding prompted the

current work to examine whether strains such as SR5 colonize with the known biofilm pathway

or with a novel pathway. Here, we show that all *V. fischeri* strains tested require the *syp* locus to

colonize a squid host, and we identify two groups of isolates that colonize with novel regulation.

Given the exquisite specificity by which *V. fischeri* bacteria colonize squid hosts, this work

reinforces the importance of biofilm formation and reveals different regulatory modes across the

evolutionary tree.

RESULTS

Most *V. fischeri* strains synthesize biofilm in response to RscS overexpression. Biofilm

formation is required for squid colonization, and overexpression of the biofilm regulator RscS in

strain ES114 stimulates a colony biofilm on agar plates (15). Our previous work demonstrated

that *V. fischeri* strain MJ11 synthesizes a colony biofilm under similar inducing conditions, which

is notable because MJ11 does not encode RscS in its chromosome (22). While the ancestral

strain MJ11 did not encode RscS, it had what seemed to be an intact *syp* locus, and

overexpression of the heterologous RscS from ES114 was sufficient to enable robust squid

colonization (22). We examined a phylogenetic tree of *V. fischeri* isolates (Fig. 1), and in this

study we expand our analysis of RscS-Syp biofilm regulation in a wider group of *V. fischeri*

strains.

Initially, we asked whether responsiveness to RscS overexpression would yield a similar colony

biofilm in this diverse group of strains. We took the same approach as our previous study and

introduced plasmid pKG11, which overexpressed ES114 RscS, into strains across the

evolutionary tree (22, 28). We observed that almost all strains tested, including those that lack

rscS, were responsive to overexpression of ES114 RscS (Fig. 2). The morphology of the colony

biofilms differed across isolates; but in most cases colony biofilm was evident at 24 h and

prominent at 48 h. All of the strains exhibited some wrinkled colony morphology at 48 h with the

exception of CG101, which was isolated from the pineapplefish *Cleidopus gloriamaris (25)*.

These results demonstrated that most *V. fischeri* strains can produce biofilm in response to

RscS overexpression, and this includes strains that presumably have not encountered *rscS* in

their evolutionary history.

One unexpected observation was that there was a subset of *rscS*-encoding strains that were

reproducibly delayed in their colony biofilm, and had only a mild wrinkled colony phenotype at

48 h (strains MB11B1, ES213, KB2B1; Fig. 2). We considered whether this was due to

differential growth of the strains, but resuspension of spots and dilution plating to determine

CFU/spot demonstrated no significant growth difference between these strains and ES114

under these conditions. The strains are closely-related (Fig. 1) and a previous study had noted

that this group shared a number of phenotypic characteristics, e.g. reduced motility in soft agar

(29). Those authors termed this tight clade as "Group A" *V. fischeri* (30). Our results in Figure 2

argue that Group A strains do not respond to RscS in the same manner as other *V. fischeri*

strains, which prompted us to investigate the evolution of the RscS-Syp signaling pathway. We

have maintained the Group A nomenclature here, and furthermore we introduce the

nomenclature of Group B (a paraphyletic group of strains that contain *rscS*; this group includes

the common ancestor of all *rscS*-containing strains) and Group C (a paraphyletic group of

strains that contains the common ancestor of all *V. fischeri* - these strains do not contain *rscS*),

as shown in Figure 1.

Ancestral Group C squid isolates colonize *E. scolopes* independent of RscS and

dependent on Syp. Group C strains generally cannot colonize squid, yet there are

Mediterranean squid isolates that appear in this group (Fig. 1; (22)). The best-studied of these

strains, SR5, was isolated from *Sepiola robusta*, is highly luminous, and colonizes the Hawaiian

bobtail squid *E. scolopes* (24). Nonetheless, this strain lacks *rscS* (27). We first asked whether

the strain can colonize in our laboratory conditions, and we confirmed that it colonizes robustly,

consistent with the result result previously published by Fidopiastis et al. (24) (Fig. 3). Next, we

asked whether it uses the Syp biofilm to colonize. To address this question, we deleted the 18

kb *syp* locus (i.e., *sypA* through *sypR*) in strains SR5 and ES114. Deletion of *rscS* or the *syp*

locus in ES114 led to a substantial defect in colonization, consistent with a known role for these

factors (Fig. 3). Similarly, deletion of the *syp* locus in SR5, a strain that does not encode *rscS*,

led to a dramatic reduction in colonization (Fig. 3). Therefore, even though strain SR5 does not

encode *rscS*, it can colonize squid, and it requires the *syp* locus to colonize normally.

RscS is dispensable for colonization in Group A strains. We noted in the wrinkled colony

biofilm assays shown in Figure 2 that Group A strains exhibited a more modest response to

overexpression of RscS. Sequencing of the native *rscS* gene in these strains revealed a

predicted -1 frameshift (ΔA1141) between the PAS domain and the histidine kinase CA domain.

Whereas ES114 and other Group B strains have nine adenines at this position, the Group A

strains have eight, leading to a frameshift and then truncation at an amber stop codon, raising

the possibility that Group A strains have a divergent biofilm signaling pathway (Fig. 4A). Given

the importance of RscS in the Group B strains including ES114, we considered the possibility

that this apparent frameshift encoded a functional protein, either through ribosomal

frameshifting or through the production of two polypeptides that together provided RscS

function; there is precedent for both of these concepts in the literature (31, 32). We first

introduced a comparable frameshift into a plasmid-borne overexpression allele of ES114 *rscS*,

and this allele did not function with the deletion of the single adenine (Fig. 4B). This result

suggested to us that the frameshift in the Group A strains may not be functional. Therefore, we

proceeded to delete *rscS* in two Group A strains (MB11B1, ES213) and two Group B strains

(ES114, MB15A4). The Group B strains required RscS for squid colonization (Fig. 5A).

However, the Group A strains exhibited no deficit in the absence of *rscS* (Fig. 5A). We next

attempted a more sensitive assay in which a Group A strain was competed against MB15A4.

Previous studies have demonstrated that in many cases Group A strains outcompete Group B

strains (30, 33). We competed Group A strain MB11B1 against Group B strain MB15A4 and

observed a significant competitive advantage for the Group A strain, as was observed

previously (30). Deletion of *rscS* in the Group A strain did not affect competitive fitness,

demonstrating that MB11B1 can outcompete a Group B strain even if MB11B1 lacks RscS (Fig.

5B).

The *syp* locus is broadly required for squid colonization. Given that Group A strains

seemed to represent a tight phylogenetic group in which RscS was not required for colonization

or competitive fitness, we next asked whether this group requires the Syp biofilm for

colonization. We proceeded to delete the entire *syp* locus in two Group A and two Group B

strains and to conduct single-strain colonization analysis. In each strain assayed, the *syp* locus

was required for full colonization, and we observed a 2-4 log reduction in CFU per animal in the

absence of the *syp* genes, pointing to a critical role for Syp biofilm in these strains (Fig. 6). In

Group A strains in particular, no colonization was detected in the absence of the *syp* locus.

Additionally, we isolated a transposon insertion in SR5 *sypJ*, and this strain did not colonize

squid well, arguing that this effect is due to the Syp biofilm and not due to regulation of a distinct

phenotype by regulators within the locus.

Group A strains encode an alternate allele of SypE. It seemed curious to us that Group A

strains do not encode a functional RscS and do not require *rscS* for colonization, yet in many

cases Group A strains can outcompete Group B strains (e.g. MB11B1 in Fig. 5B; and Refs. (30,

33)). We reasoned that if the Syp biofilm had a different regulatory architecture in Group A

strains--e.g., constitutively activated or activated by a different regulatory protein--then this could

explain the Syp regulation independent of RscS. Genome sequencing of SR5 and MB11B1 did

not identify a unique histidine kinase that was likely to directly substitute for RscS (27, 33).

Given that the *syp* locus encodes biofilm regulatory proteins, we examined *syp* conservation.

We used TBLASTN with the ES114 Syp proteins as queries to determine amino acid

conservation in the other *V. fischeri* Group A strain MB11B1, Group C strain SR5, and the *Vibrio*

vulnificus type strain ATCC 27562 (34, 35). As shown in Figure 7, ES114 SypE, a response

regulator and serine kinase/phosphatase that is a negative regulator of the Syp biofilm (17, 36),

exhibited the lowest level of conservation among *syp* locus products. *V. vulnificus* does not

encode a SypE ortholog (37), as the syntenic (but not homologous) RbdE encodes a predicted

ABC transporter substrate-binding protein. The closest hit for SypE was AOT11_RS12130 (9%

identity), compared to 7% identity for the RbdE. Due to the reduced conservation at both the

strain and species levels, we analyzed *V. fischeri* MB11B1 SypE in greater detail. Examination

of the *sypE* coding sequence revealed an apparent -1 frameshift mutation in which the position

33 (guanine in ES114 and adenine in other Group B and C strains examined) is absent in Group

A strains (Fig. 7B). We therefore considered the hypothesis that SypE is nonfunctional in Group

A, and that these strains can colonize because they are lacking a functional copy of this

negative regulator that is itself regulated by RscS.

To test this hypothesis, we relied on knowledge of the biofilm regulatory pathway from ES114, in

which overexpression of SypG produces a wrinkled colony phenotype, but only in strains lacking

SypE activity (38, 39). Therefore, we introduced the SypG-overexpressing plasmid pEAH73 into

strains as a measure of whether the SypE pathway was intact. In the ES114 strain background,

we observed cohesive wrinkled colony formation at 48 h in an ES114 Δ*sypE* strain, but not in

the wild-type parent (Fig. 8A). If the *sypE* frameshift observed in MB11B1 led to a loss of

function, then introduction of that frameshift into ES114 would lead to a strain that is equivalent

to the Δ*sypE* strain. We constructed this strain and upon SypG overexpression we observed

wrinkled colony formation. Surprisingly, the biofilm phenotype was observed earlier (i.e., by 24

h) and leads to more defined colony biofilm architecture at 48 h. While the lack of SypE leads to

increased and more rapid biofilm formation, in this assay we observed an even greater increase

as a result of the frameshift in *sypE* (Fig. 8A).

We proceeded to conduct a similar assay in the MB11B1 strain background. The colony biofilm

phenotypes were muted compared to the ES114 background, but the pattern observed is the

same. Strains lacking the additional nucleotide at position 33 (i.e., the native MB11B1 allele)

exhibited the strongest cohesion, whereas strains with the nucleotide to mimic ES114 *sypE* (i.e.,

added back in MB11B1 *sypE*(nt::33G)) were not cohesive (Fig. 8B). These results argue that a

novel allele of *sypE* is found in Group A strains and this allele results in more substantial biofilm

formation than in a Δ*sypE* strain.

Our finding that the MB11B1 *sypE* allele promotes biofilm formation bolstered the model that

this allele contributes to the ability of MB11B1 to colonize squid independent of RscS. To test

this model, we introduced the frameshift into ES114 or "corrected" the frameshift in MB11B1.

We then conducted single-strain colonization assays, and in each case the *sypE* allele alone

was not sufficient to alter the overall colonization behavior of the strain (Fig. 9). Therefore, these

data suggest that the frameshift in the MB11B1 *sypE* is not sufficient to explain its ability to

colonize independent of RscS, and therefore other regions of SypE and/or other loci in the

MB11B1 genome contribute to its ability to colonize independent of RscS.

BinK is active in Group A, B, and C strains. We recently described the histidine kinase, BinK,

which negatively regulates *syp* transcription and Syp biofilm formation (18). In ES114,

overexpression of BinK impairs the ability of *V. fischeri* to colonize. We therefore reasoned that

if BinK could function in Group A strains and acted similarly to repress Syp biofilm, then

overexpression of BinK would reduce colonization of these strains. We introduced the pBinK

plasmid (i.e., ES114 *binK* (18)) and asked whether multicopy *binK* would affect colonization. In

strain MB11B1, BinK overexpression led to a dramatic reduction in colonization (Fig. 10A).

Therefore, there is a clear effect for BinK overexpression on the colonization of the Group A

strain MB11B1.

We attempted to ask the same question in Group C strain SR5, but the pES213-origin plasmids

were not retained during squid colonization. Therefore, we instead asked whether deletion of

the BinK, a negative regulator of ES114 colonization, has a comparable effect in SR5 (18). We

deleted *binK* and observed a 2.4-fold competitive advantage during squid competition (Fig.

10B), arguing that BinK in this Group C strain is active and performs an inhibitory function

similar to that in ES114.

We next examined the colony biofilm phenotype for strains lacking BinK. MB11B1 Δ*binK*

exhibited a mild colony biofilm phenotype at 48 h, as evidenced by the cohesiveness of the spot

when disrupted with a toothpick (Fig. 10C). The colonies also exhibited an opaque phenotype.

In a minority of experimental replicates, wrinkled colony morphology was evident at 48 h, but in

all samples wrinkled colony morphology was visible at 7 d (data not shown). The SR5 Δ*binK*

strain also exhibited slightly elevated biofilm morphology at 48 h, though the cells were not as

cohesive as those of MB11B1 Δ*binK* (Fig. 10C). Together, the results in Figure 10 argue that

BinK, a factor that has been characterized as a negative regulator of Syp biofilm, plays similar

roles in Group A and Group C strains and has a widely-conserved function across the *V. fischeri*

evolutionary tree.

DISCUSSION

This study examines regulation of a beneficial biofilm that is critical to host colonization

specificity in *V. fischeri.* The Syp biofilm was discovered thirteen years ago and has been

characterized extensively for its role in facilitating squid colonization by *V. fischeri.* This work

establishes that the *syp* locus is required broadly across squid symbionts, and it uncovers three

groups of *V. fischeri* that use different regulatory programs upstream of the *syp* locus. A

simplified phylogenetic tree showing key features of squid symbionts in these three groups is

shown in Figure 11.

There are three nested evolutionary groups of *V. fischeri* that have been described separately in

the literature and here we formalize the nomenclature of Groups C, B, and A. Group A is a

monophyletic group, as are Groups AB and ABC (Fig. 1). This work provides evidence that

squid symbionts in each group have a distinct biofilm regulatory architecture. Most *V. fischeri*

isolates that have been examined from the ancestral Group C cannot colonize squid; however,

those that can colonize do so without the canonical biofilm regulator RscS. We show that the

known targets of RscS regulation—genes in the *syp* biofilm locus—are nonetheless required for

squid colonization by this group. Group B strains include the well-characterized ES114 strain,

which requires RscS and the *syp* locus to colonize squid. Group A strains differ phenotypically

and behaviorally from the sister Group B strains (30), and we demonstrate that these strains

have altered biofilm regulation. Group A strains have a frameshift in *rscS* that renders it

nonfunctional, and a 1 bp deletion in *sypE*, and we provide evidence that the *sypE* allele

promotes biofilm development in the absence of RscS. Additionally, we note that the *sypE*

frameshift is not present in SR5, arguing for distinct modes of biofilm regulation in Groups A, B,

and C.

At the same time, this study provides evidence that some aspects of biofilm regulation are

conserved in diverse squid symbionts, such as the effects of the strong biofilm negative

regulator BinK. Published data indicate that evolved BinK alleles can alter colonization of H905

(Group B) and MJ11 (Group C), and that a deletion of MJ11 *binK* leads to enhanced

colonization (20). Our experiments in Figure 10 show a clear effect for BinK in all three

phylogenetic groups. We also observed responsiveness to RscS overexpression in all squid

symbionts examined (Fig. 2). CG101 was the only *V. fischeri* strain examined that did not exhibit

a colony biofilm in response to RscS overexpression. CG101 was isolated from the Australian

fish *Cleidopus gloriamaris*; based on these findings, we suspect that the strain does not have an

intact *syp* locus or otherwise has divergent biofilm regulation.

It remains a formal possibility that the entire *syp* locus is not required in Group A or Group C,

but instead that only one or a subset of genes in the locus are needed. We have constructed

Campbell-type (insertion duplication) alleles to interrupt *sypG* in MB11B1 and SR5, and

additionally have isolated a transposon insertion in SR5 *sypJ*, and none were able to colonize

well. Additionally, aggregation in squid mucus has been observed for the Group A strain

MB13B2, and this aggregation is dependent on *sypQ* (40). In our data we note that Group A

strains were completely unable to colonize in the absence of the *syp* locus, unlike the tested

Group B & C strains that exhibited reduced colonization in their respective mutants (Figs. 3, 6).

Therefore, the simplest explanation is that the *syp* locus is required in divergent strains in a

manner similar to how it is used in ES114. We think that the ability to completely delete the *syp*

locus is a clean way to ask whether the locus is required for specific phenotypes, and our

strains are likely to be useful tools in probing Syp protein function in diverse *V. fischeri* isolates.

It is intriguing to speculate as to how the two frameshifts in the Group A strains arose, and why

the nonfunctional RscS is tolerated in this group. One possible scenario is that the Group A

strains acquired a new regulatory input into the Syp pathway, and that the presence of this new

regulator bypassed the requirement for RscS. We note that comparative genomic analysis of

Hawaiian D (dominant)-type strains--which largely overlap with Group A--revealed an additional

250 kb of genomic DNA compared to other isolates, yielding a large cache of genes that could

play a role in this pathway (33). A related possibility is that *rscS*-independent colonization

results from altered regulation of the *syp* locus, either due to changes in regulators (e.g. SypF)

or sites that are conserved with Group B. An additional possibility is that the *sypE* frameshift

arose, enabling Group A strains to colonize independent of *rscS*. Given that correction of this

frameshift in MB11B1 does not significantly affect colonization ability (Fig. 9), this sequence of

events seems less likely, and we expect that another regulator in MB11B1 is required for the

RscS-independent colonization phenotype. There is evidence that under some conditions LuxU

can regulate the *syp* biofilm (41), and as this protein is conserved in *V. fischeri* it may play an

important role in Group A or Group C.

Results from two experimental conditions suggest that the Group A strains may have an

elevated baseline level of biofilm formation. Our data indicate that in the absence of BinK or

upon SypG overexpression, MB11B1 colonies exhibit strong cohesion under conditions in which

ES114 does not (Figs. 8, 10). Furthermore, we note that the Group A strain MB11B1, when

lacking BinK, also exhibits a darker, or more opaque, colony phenotype (Fig. 10). This

phenotype has been observed in some ES114 mutants (16) but not in the corresponding ES114

Δ*binK* strain (Fig. 10). The entire colonization lifecycle likely requires a balance between biofilm

formation/cohesion and biofilm dispersal, and these data argue that Group A strains may be

more strongly tilted toward the biofilm-producing state. There is evidence that strains lacking

BinK exhibit a colonization advantage in the laboratory (18, 20), suggesting that this strategy of

more readily forming biofilms may provide a fitness advantage in nature. At the same time, the

biofilm negative regulator BinK is conserved among *V. fischeri* strains examined (including

MB11B1; Fig. 10), arguing that there is a benefit to reducing biofilm formation under some

conditions.

Our study provides hints as to the role of SypE in MB11B1 and other Group A strains. In ES114,

the C-terminus is a PP2C serine kinase domain, whereas the N-terminus of SypE is an RsbW

serine phosphatase domain. SypE acts to phosphorylate and dephosphorylate SypA Ser-56,

with the unphosphorylated SypA being the active form to promote biofilm development (17). The

balance between SypE kinase and phosphatase is modulated by a central two-component

receiver domain (17). Our data that the MB11B1 *sypE* allele promotes biofilm formation suggest

that the protein is tilted toward the phosphatase activity. In MB11B1, the frameshift early in *sypE*

suggests that there is a different start codon and therefore a later start codon. An alternate GTG

start codon in MB11B1 occurs corresponding to codon 18 in ES114 *sypE* (Fig. 7), and this is

likely the earliest start for the MB11B1 polypeptide. We attempted to directly identify the SypE

N-terminus by mass spectrometry, yet we could not identify the protein from either strain.

Additional study is required to elucidate how MB11B1 SypE acts to promote biofilm formation.

V. fischeri strains are valuable symbionts in which to probe the molecular basis to host

colonization specificity in animals (22, 25, 26). A paradigm has emerged in which biofilm

formation through the RscS-Syp pathway is required for squid colonization but not for fish

colonization. This study affirms a role of the Syp biofilm, but at the same time points out

divergent (RscS-independent) regulation in Group C and Group A isolates. In another well-

studied example of symbiotic specificity, Rhizobial Nod factors are key to generating specificity

with the plant host, yet strains have been identified that do not use this canonical pathway (42,

43). Future work will elaborate on these RscS-independent pathways to determine how non-

canonical squid colonization occurs in diverse natural isolates.

MATERIALS & METHODS

Bacterial strains and growth conditions. *V. fischeri* and *E. coli* strains used in this study can

be found in Table 1. *E. coli* strains, used for cloning and conjugation, were grown in Luria-

Bertani (LB) medium (25 g Difco LB Broth [BD] per liter). *V. fischeri* strains were grown in Luria-

Bertani salt (LBS) medium (25 g Difco LB Broth [BD], 10 g NaCl, and 50 ml 1 M Tris buffer pH

7.0, per liter). Growth media were solidified by adding 15 g Bacto agar (BD) per liter. When

necessary, antibiotics (Gold Biotechnology) were added at the following concentrations:

tetracycline, 5 µg/ml for *V. fischeri*; erythromycin, 5 µg/ml for *V. fischeri*; kanamycin, 50 µg/ml for

E. coli and 100 µg/ml for *V. fischeri*; and chloramphenicol, 25 µg/ml for *E. coli*, 2.5 -5 µg/ml for

Group B *V. fischeri*, and 1 - 2.5 µg/ml for Group A *V. fischeri*. The two MB11B1 / pKV69 strains

listed reflect two separate constructions of this strain, though we have not identified any

differences between them.

Phylogenetic analysis. Phylogenetic reconstructions assuming a tree-like topology were

created with three methods: maximum parsimony (MP); maximum likelihood (ML); and

Bayesian inference (Bayes) as previously described (22, 30). Briefly, MP reconstructions were

performed by treating gaps as missing, searching heuristically using random addition, tree-

bisection reconnection with a maximum of 8 for swaps, and swapping on best only with 1000

repetitions. For ML and Bayesian analyses, likelihood scores of 1500+ potential evolutionary

models were evaluated using both the corrected and uncorrected Akaike Information Criterion,

the Bayesian Information Criterion, and Decision Theory (Performance Based Selection) as

implemented by jModelTest2.1 (44). For all information criteria, the most optimal evolutionary

model was a symmetric model with a proportion of invariable sites and a gamma distribution of

rate heterogeneity (SYM+I+Γ).

ML reconstruction was implemented via PAUP*4.0a163 (45) by treating gaps as missing,

searching heuristically using random addition, tree-bisection reconnection for swaps, and

swapping on best only with 1000 repetitions. Bayesian inference was done by invoking the

'nst=6' and 'rates=invgamma' and 'statefreqpr=fixed(equal)' settings in the software package

MrBayes3.2.6 (46). The Metropolis-coupled Markov chain Monte Carlo (MCMCMC) algorithm

used to estimate the posterior probability distribution for the sequences was set up with

'temp=0.2' and one incrementally 'heated' chain with three 'cold' chains; these four chains were

replicated two times per analysis to establish convergence of the Markov chains (i.e.,

'stationarity' as defined by (47) and interpreted previously in (30)). For this work, stationarity was

achieved after approximately 50,000 samples (5,000,000 generations) were collected, with 25%

discarded. The ~37,500 samples included were used to construct a 50% majority-rule

consensus tree from the sample distribution generated by MCMCMC and assess clades'

posterior probabilities. For ML and MP analyses, the statistical confidence in the topology of

each reconstruction was assessed using 1000 bootstrap replicates. Phylogenetic trees were

visualized with FigTree 1.4.3 (http://tree.bio.ed.ac.uk/software/figtree); the final tree was edited

for publication with Inkscape 0.91 (http://inkscape.org/) and GIMP 2.8.22 (http://www.gimp.org/).

DNA synthesis and sequencing. Each of the primers listed in Table 3 was synthesized by

Integrated DNA Technologies (Coralville, IA). Full inserts from all cloned constructs were

verified by Sanger DNA sequencing through ACGT, Inc via the Northwestern University

Feinberg School of Medicine NUSeq Core Facility; or the University of Wisconsin-Madison

Biotechnology Center. Sequence data was analyzed with SeqMan Pro (DNAStar software),

SnapGene (GSL Biotech), and Benchling.

Construction of gene deletions. Deletions in *V. fischeri* strains ES114 and MB11B1 were

made according to the lab's gene deletion protocol: doi:10.5281/zenodo.1470836. In brief, 1.6

kb upstream and 1.6 kb downstream of the targeted gene or locus were cloned into linearized

plasmid pEVS79 (amplified with primers pEVS79_rev_690/pEVS79_for_691) using Gibson

Assembly (NEBuilder HiFi DNA Assembly cloning kit) with the primer combinations listed in

Table S1. The Gibson mix, linking together the upstream and downstream flanking regions, was

transformed into *E. coli* on plates containing X-gal, with several white colonies selected for

further screening by PCR using primers flanking the upstream/downstream junction (Tables 3

and S1). The resulting plasmid candidate was confirmed by sequencing and conjugated into the

V. fischeri recipient by tri-parental mating with helper plasmid pEVS104, selecting for the

chloramphenicol resistance of the plasmid backbone. *V. fischeri* colonies were first screened for

single recombination into the chromosome by maintaining antibiotic resistance in the absence of

selection and then screened for double recombination by the loss of both the antibiotic resistant

cassette and the gene/locus of interest. Constructs were verified by PCR (Table 3) and

sequencing.

Deletion of SR5 *binK* was conducted using Splicing by Overlap Extension PCR (SOE-PCR) and

natural transformation (method modified from (48)). Oligos binK-F1 and binK-R1-LUH, and

oligos binK-F2-RUH and binK-R2 were used in a PCR with MJM1125 (SR5) genomic DNA as

the template to amplify DNA fragments containing ~1 kb of sequence upstream and

downstream relative to *binK*, respectively. Using SOE-PCR, these fragments were fused on

either side to a third DNA fragment containing an Erm[R] cassette, which was amplified using

pHB1 as template and oligos HB41 and HB42. We then used natural transformation with

pLostfoX (49) to insert this mutagenic DNA into MJM1125, where the flanking sequences guide

the Erm[R] cassette to replace *binK*, generating the desired gene deletion. Candidate SR5 Δ*binK*

mutants were selected after growth on LBS-Erm5 plates. Oligos binK-F1 and binK-R2, and HB8

and binK-FO were used to screen candidates for the correct deletion scar by PCR, and oligos

KMB_036 and KMB_037 were used to confirm the absence of *binK* in the genome. The deletion

was verified by Sanger sequencing with primers HB8, HB9, HB42, and HB146. The base

plasmid pHB1 contains an erythromycin resistance cassette flanked by FRT sites, and was

constructed using oligos HB23 and HB39 with gBlock gHB1 (sequence in Supplementary File

S1; Integrated DNA Technologies, Inc.) as template to amplify the Erm[R] cassette flanked by

HindIII and BamHI sites, which was then cloned into the corresponding site in pUC19.

For most constructs, the deleted genetic material was between the start codon and last six

amino acids (50), with two exceptions: the Δ*sypE* in MJM1130 included the ATG that is two

amino acids upstream of the predicted start codon, but not the canonical start codon; and the

Δ*binK* alleles in MJM1117, MJM1130, and MJM2114, which were constructed to be equivalent

to MJM2251 (Δ*binK* in ES114) (18). The Δ*binK* alleles in these strains include the start codon,

the next six codons, two codons resulting from ATCGAT (ClaI site), and the last three codons

for a predicted 12 amino acid peptide.

Construction of *sypE* alleles. To create *sypE*(ntG33Δ) in MJM1100 and *sypE*(nt33::G) in

MJM1130, the single point mutation was created by amplifying the gene in two halves, with the

N-terminal portion consisting of approximately 300 bp upstream of *sypE* up through nucleotide

33 and the C-terminal portion consisting of nucleotide 33 and the remaining *sypE* gene. The

overlap between the two halves contained the single nucleotide polymorphism in the primers

that connected them. The altered *sypE* alleles were initially cloned into plasmid pEVS107

(linearized with primers pEVS107_3837/pEVS107_3838) using Gibson Assembly and then the

entire altered *sypE* allele was subcloned into pEVS79 with Gibson Assembly (Table S1). After

double recombination of the vector into *V. fischeri*, candidate colonies for the altered *sypE* in

MJM1100 were screened with primers ES114_indel_for/ES114_indel_rev. The primer set

anneals more strongly to the wildtype *sypE* sequence than to *sypE*(ntG33::Δ). Candidates in the

MJM1100 background with a fainter PCR band were sequenced and confirmed to have the

sypE(ntG33::Δ) allele. For MJM1130, the primer set MB11B1_indel_for/MB11B1_indel_rev

anneals more strongly to the *sypE*(nt33::G) allele than to the naturally occurring *sypE* allele and

candidates in MJM1130 that contained a more robust PCR band were selected for sequencing

to be confirmed as being *sypE*(nt33::G).

Construction of pKG11 *rscS1*(ntA1141::Δ). Plasmid pKG11 encodes an overexpression allele

of RscS, termed *rscS1 (15, 28)*. *rscS* nucleotide A1141 was deleted on the plasmid using the

Stratagene Quikchange II Site-Directed Mutagenesis Kit with primers rscS_del1F and

rscS_del1R. The resulting plasmid, pMJM33, was sequenced with primers MJM-154F and

MJM-306R to confirm the single base pair deletion.

Squid colonization. Hatchling *E. scolopes* were colonized by exposure to approximately 3 x

10^3 CFU/ml (ranging from 5.2 x 10^2 - 1.4 x 10^4 CFU/ml; as specified in figure legends) of each

strain in a total volume of 40 ml of FSIO (filter-sterilized Instant Ocean) for 3 hours. Squid were

then transferred to 100 ml of FSIO to stop the inoculation and then transferred to 40 ml FSIO for

an additional 45 hours with a water change at 24 hours post inoculation. For Figure 10A,

kanamycin was added to the FSIO to keep selective pressure on the plasmid. After 48 hours of

colonization, the squid were euthanized and surface sterilized by storage at -80 ºC, according to

standard practices (51). For determination of CFU per light organ, hatchlings were thawed,

homogenized, and 50 µl of homogenate dilutions was plated onto LBS plates. Bacterial colonies

from each plate were counted and recorded. Mock treated, uncolonized hatchlings ("apo-

symbiotic") were used to determine the limit of detection in the assay. The competitive index

(CI) was calculated from the relative CFU of each sample in the output (light organ) versus the

input (inoculum) as follows:

Log_{10} ((Test strain[light organ] / Control strain[light organ]) / (Test strain[inoculum] / Control

strain[inoculum])). For competitions of natural isolates, the Group A strain (or its Δ*rscS*

derivative) was the test strain and the Group B strain was the control strain. Colony color was

used to enumerate colonies from each--white for Group A strains MB11B1 and ES213; yellow

for Group B strains ES114 and MB15A4--along with PCR verification of selected colonies. For

competition between SR5 and SR5 Δ*binK*, 100 colonies per squid were patched onto LBS-Erm5

and LBS.

Colony biofilm assays. Bacterial strains were grown in LBS media (Fig. 10C) or LBS-Cam2.5

media (Figs. 2, 8) for approximately 17 hours, then 10 µl (Fig. 2) or 8 µl (Fig. 8, 10C) was

spotted onto LBS plates (Fig. 10C) or LBS-Tet5 plates (Figs. 2, 8). Spots were allowed to dry

and the plates incubated at 25 °C for 48 hours. Images of the spots were taken at 24 and 48 h

post-spotting using a Leica M60 microscope and Leica DFC295 camera. After 48 h of growth,

the spots were disrupted using a flat toothpick and imaged similarly.

Analysis of DNA and protein sequences *in silico*. Amino acid sequences for *V. fischeri*

ES114 *syp* genes were obtained from RefSeq accession NC_006841.2. Local TBLASTN

queries were performed for each protein against nucleotide databases for the following strains,

each of which were derived from the RefSeq cds_from_genomic.fna file: *V. fischeri* SR5

(GCA_000241785.1), *V. fischeri* MB11B1 (GCA_001640385.1) and *V. vulnificus* ATCC27562

(GCA_002224265.1). Percent amino acid identity was calculated as the identity in the BLAST

query divided by the length of the amino acid sequence in ES114. Domain information is from

the PFAM database (52).

FIGURE LEGENDS

Figure 1. *Vibrio fischeri* phylogeny, highlighting the source of each strain. Bayesian

phylogram (50% majority-rule consensus) inferred with a SYM+I+Γ model of evolution for the

concatenated gene fragments *recA*, *mdh*, and *katA*. In this reconstruction, the root connected to

a clade containing the four non-*V. fischeri* outgroup taxa. Statistical support is represented at

nodes by the following three numbers: upper left, Bayesian posterior probability (of

approximately 37,500 non-discarded samples) multiplied by 100; upper right, percentage of

1000 bootstrap Maximum Likelihood pseudo-replicates; bottom middle center, percentage of

1000 bootstrap Maximum Parsimony pseudo-replicates. Statistical support values are listed only

at nodes where more than 2 methods generated support values ≥ 50%. Strains sharing identical

sequences for a given locus fragment are listed next to a vertical bar at a leaf; because of a lack

of space, some support values have been listed either immediately to the right of their

associated nodes and are marked with italicized lower-case Roman numerals in the phylogram.

The isolation habitat and geography of each strain are indicated by symbol and color,

respectively. The black bar represents 0.01 substitutions/site.

Figure 2. Most *V. fischeri* strains tested form colony biofilm in response to RscS

overexpression. Spot assays of the indicated *V. fischeri* strains carrying pKV69 (vector) or

pKG11 (*rscS1*; overexpressing ES114 *rscS*) after 24 and 48 h. Strains are MJM1268,

MJM1269, MJM1246, MJM1247, MJM1266, MJM1267, MJM1219, MJM1221, MJM1238,

MJM1239, MJM1104, MJM1106, MJM1276, MJM1277, MJM1270, MJM1271, MJM1258,

MJM1259, MJM1254, MJM1255, MJM1242, MJM1243, MJM1240, MJM1241, MJM1272,

MJM1273, MJM1274, MJM1275, MJM1278, MJM1279, MJM1109, MJM1111, MJM1280,

MJM1281, MJM1260, MJM1261, MJM1244, MJM1245, MJM1256, and MJM1257. Different

phenotypes were observed in the isolates examined; in most cases we observed wrinkled

colonies, but in some cases we observed only a subtle pocked pattern (EM30), and in other

cases we did not observe any change in colony morphology compared to the vector control

(noted by *). The black bar is 5 mm in length.

Figure 3. Squid colonization in Group C strain SR5, which does not encode RscS, is

dependent on the *syp* polysaccharide locus. Single-strain colonization experiments were

conducted and circles represent individual animals. The limit of detection for this assay,

represented by the dashed line, is 7 CFU/LO, and the horizontal bars represent the median of

each set. Hatchling squid were inoculated with 1.5-3.2 × 10^3 CFU/ml bacteria, washed at 3 h

and 24 h, and assayed at 48 h. Each dot represents an individual squid. Strains are: MJM1100,

MJM3010, MJM3062, MJM1125, and MJM3501. Statistical comparisons by the Mann-Whitney

test, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure 4. Group A strains have a frameshift in *rscS*. (A) ES114 RscS protein domains.

Nucleotides 1114-1173 in ES114 RscS (AF319618) and their homologous sequences in the

other Group B and Group A strains are listed. The -1 frameshift is present in the Group A *rscS*

alleles. The ES114 reading frame is noted on the top of the alignment and the Group A reading

frame on the bottom, which is predicted to end at the amber stop codon. (B) Deletion of

nucleotide A1141 in ES114 to mimic this frameshift in pKG11 renders it unable to induce a

colony biofilm in a spot assay at 48 h. Strains are MJM1104, MJM1106, and MJM2226.

Figure 5. Group A strains MB11B1 and ES213 do not require RscS for squid colonization.

Wild-type (WT) and Δ*rscS* derivatives of the indicated strains were assayed in (A) a single-strain

colonization assay and (B) competitive colonization against Group B strain MB15A4. Hatchling

squid were inoculated at 3.5-14 × 10^3 CFU/ml bacteria, washed at 3 h and 24 h, and assayed at

48 h. Each dot represents an individual squid. (A) Strains: MJM1100, MJM3010, MJM2114,

MJM3042, MJM1130, MJM3046, MJM1117, and MJM3017. The limit of detection is represented

by the dashed line, and the horizontal bars represent the median of each set. In both panels,

open dots are wild type and filled dots are Δ*rscS*. (B) The competitive index (CI) is defined in the

methods and is shown on a Log_{10} scale. Strains: MJM1130 and MJM3046, each competed

against MJM2114. Values greater than 1 indicate more MB11B1. Statistical comparisons by the

Mann-Whitney test, ns not significant, **** $p<0.0001$.

Figure 6. Group B and Group A strains require the *syp* locus for robust squid

colonization. Wild type (WT) and Δ*syp* derivatives of the indicated strains were assayed in a

single strain colonization assay. Hatchling squid were inoculated with 6.7-32 × 10^2 CFU/ml

bacteria (ES114 and MB15A4 backgrounds) or 5.2-8.9 × 10^2 CFU/ml bacteria (MB11B1 and

ES213 backgrounds), washed at 3 h and 24 h, and assayed at 48 h. Each dot represents an

individual squid. The limit of detection is represented by the dashed line and the horizontal bars

represent the median of each set. Strains are MJM1100, MJM3062, MJM2114, MJM3071,

MJM1130, MJM3065, MJM1117, and MJM3068. Statistical comparisons by the Mann-Whitney

test, **** $p<0.0001$.

Figure 7. Group A strains have a frameshift in *sypE*. (A) Amino acid identity in the Syp locus.

Results show the identity from TBLASTN query using the *V. fischeri* ES114 protein sequences

as queries against genes in the homologous loci in *V. fischeri* strains or *V. vulnificus* ATCC

27562. The identity for SypE against *V. vulnificus* is plotted for the syntenous RbdE, although

this is not the highest TBLASTN hit, as described in the text. (B) ES114 SypE protein domains.

Nucleotides 1-60 in ES114 *sypE* and their homologous sequences in the other Group C, B, and

A strains are listed. A -1 frameshift is present in the Group A *sypE* alleles. The ES114 reading

frame is noted on the top of the alignment and the Group A reading frame on the bottom, which

is predicted to end at the amber stop codon. A possible GTG start codon for the resumption of

translation in the ES114 reading frame is present at the position corresponding to the 18th

codon in ES114 *sypE*.

Figure 8. The MB11B1 *sypE* frameshift leads to an enhanced biofilm phenotype upon

SypG overexpression. Spot assays of strains carrying the pKV69 vector or pEAH73 SypG

overexpression plasmid. (A) ES114 strain background. Strains lacking SypE produce a wrinkled

colony phenotype upon SypG overexpression. Deletion of nucleotide 33 in *sypE* to mimic the

Group A frameshift led to earlier wrinkling and a more pronounced colony biofilm at 48 h.

Strains: MJM1104, MJM3455, MJM3418, MJM3419, MJM3364, and MJM3365. (B) Group A

strain MB11B1, which naturally carries a -1 frameshift in *sypE*, exhibits a cohesive phenotype at

48 h with overexpression of SypG. Deletion of *sypE* reduces this phenotype, and repairing the

frameshift by addition of a guanosine at nucleotide 33 further reduces the cohesiveness of the

spot. Strains: MJM3370, MJM3371, MJM3411, MJM3412, MJM3398, and MJM3399.

Figure 9. The *sypE* -1 frameshift allele is not sufficient to affect colonization ability. The

indicated strains were assayed in a single-strain colonization assay. Gray boxes denote alleles

distinct from their wild-type background. Frameshift "fs" refers to alleles--relative to an ES114

reference--that lack *rscS* nucleotide A1141, or that lack *sypE* nucleotide G33. The wild-type

MB11B1 strain contains natural frameshifts in these loci, and the ES114 nt33::ΔG allele was

constructed. Addition back of the nucleotide in MB11B1 *sypE* is denoted as "(+)". Hatchling

squid were inoculated with 6.8-8.4 × 10^2 CFU/ml bacteria (MB11B1 background) or 4.0-5.4 ×

10^3 CFU/ml bacteria (ES114 background), washed at 3 h and 24 h, and assayed at 48 h. Each

dot represents an individual squid. The limit of detection is represented by the dashed line and

the horizontal bars represent the median of each set. Strains are MJM1100, MJM3010,

MJM4323, MJM3394, MJM1130, and MJM3397. Statistical comparisons by the Mann-Whitney

test, ns not significant.

Figure 10. BinK is active in Groups A, B, and C. (A) Overexpression of pBinK inhibits

colonization in Group A strain MB11B1. Hatchling squid were inoculated with 3.6-6.8 × 10^3

CFU/ml bacteria, washed at 3 h and 24 h, and assayed at 48 h. Each dot represents an

individual squid. The limit of detection is represented by the dashed line and the horizontal bars

represent the median of each set. The vector control is pVSV104. Strains are MJM1782,

MJM2386, MJM2997, and MJM2998. (B) Deletion of *binK* confers a colonization defect in Group

C strain SR5. Strains are MJM1125 and MJM3571; mean inoculum of 7.2 × 10^3 CFU/ml;

median competitive index (CI) was 0.38 (i.e., 2.4-fold advantage for the mutant). (C) Deletion of

the native *binK* in MB11B1 yielded opaque and cohesive spots, which are stronger phenotypes

than we observe in ES114. Strains are MJM1100, MJM2251, MJM1130, MJM3084, MJM2997,

and MJM2998. Statistical comparisons by the Mann-Whitney test, **** p<0.0001.

Figure 11. Summary model of distinct modes of biofilm formation in squid-colonizing *V.*

fischeri. Phylogenetic tree is simplified from Figure 1, and illustrates key features of squid

symbionts in the three groups. Shown are divergent aspects (RscS, SypE) and conserved

regulation (BinK). In all groups, the *syp* exopolysaccharide locus is required for squid

colonization.

SUPPLEMENTAL FILES

Table S1 (PDF). Primer pairs for construction of the deletion mutants. Detailed

oligonucleotide and construction details for deletions and the MJM1100 sypE(ntG33Δ) and

MJM1130 *sypE*(*nt33*::*G*) strains.

File S1 (PDF). Sequence of the synthetic dsDNA, gBlock_erm. The sequence is provided in

FASTA format printed as PDF.

Table 1. Bacterial strains.

Strain	Genotype	Source/Reference
V. fischeri		
MJM1059	MJ11	(25, 53)
MJM1100	ES114	(54)
MJM1104	ES114 (MJM1100) / pKV69	This study
MJM1106	ES114 (MJM1100) / pKG11	This study
MJM1109	MJ11 (MJM1059) / pKV69	This study
MJM1111	MJ11 (MJM1059) / pKG11	This study
MJM1114	MJ12	(53)
MJM1115	CG101	(25)

MJM1117	ES213	(55)
MJM1119	EM18	(25, 53)
MJM1120	EM24	(53, 56)
MJM1121	EM30	(53)
MJM1122	WH1	(57)
MJM1125	SR5	(24)
MJM1126	SA1	(24)
MJM1127	KB1A97	(29)
MJM1128	KB2B1	(29)
MJM1129	KB5A1	(29)
MJM1130	MB11B1	(29)
MJM1136	EM17	(56)
MJM1147	*mjapo*.6.1	(22)
MJM1149	*mjapo*.7.1	(22)
MJM1151	*mjapo*.8.1	(22)
MJM1153	*mjapo*.9.1	(22)
MJM1219	*mjapo*.8.1 / pKV69	This study

MJM1221	*mjapo*.8.1 / pKG11	This study
MJM1238	MJ12 (MJM1114) / pKV69	This study
MJM1239	MJ12 (MJM1114) / pKG11	This study
MJM1240	SR5 (MJM1125) / pKV69	This study
MJM1241	SR5 (MJM1125) / pKG11	This study
MJM1242	SA1 (MJM1126) / pKV69	This study
MJM1243	SA1 (MJM1126) / pKG11	This study
MJM1244	MB11B1 (MJM1130) / pKV69	This study
MJM1245	MB11B1 (MJM1130) / pKG11	This study
MJM1246	EM17 (MJM1136) / pKV69	This study
MJM1247	EM17 (MJM1136) / pKG11	This study
MJM1254	KB1A97 (MJM1127) / pKV69	This study
MJM1255	KB1A97 (MJM1127) / pKG11	This study
MJM1256	KB2B1 (MJM1128) / pKV69	This study
MJM1257	KB2B1 (MJM1128) / pKG11	This study
MJM1258	KB5A1 (MJM1129) / pKV69	This study
MJM1259	KB5A1 (MJM1129) / pKG11	This study

MJM1260	ES213 (MJM1117) / pKV69	This study
MJM1261	ES213 (MJM1117) / pKG11	This study
MJM1266	EM18 (MJM1119) / pKV69	This study
MJM1267	EM18 (MJM1119) / pKG11	This study
MJM1268	EM24 (MJM1120) / pKV69	This study
MJM1269	EM24 (MJM1120) / pKG11	This study
MJM1270	EM30 (MJM1121) / pKV69	This study
MJM1271	EM30 (MJM1121) / pKG11	This study
MJM1272	*mjapo*.6.1 (MJM1147) / pKV69	This study
MJM1273	*mjapo*.6.1 (MJM1147) / pKG11	This study
MJM1274	*mjapo*.7.1 (MJM1149) / pKV69	This study
MJM1275	*mjapo*.7.1 (MJM1149) / pKG11	This study
MJM1276	*mjapo*.9.1 (MJM1151) / pKV69	This study
MJM1277	*mjapo*.9.1 (MJM1151) / pKG11	This study
MJM1278	CG101 (MJM1115) / pKV69	This study
MJM1279	CG101 (MJM1115) / pKG11	This study
MJM1280	WH1 (MJM1122) / pKV69	This study

MJM1281	WH1 (MJM1122) / pKG11	This study
MJM1782	ES114 (MJM1100) pVSV104	(18)
MJM2114	MB15A4	(29)
MJM2226	ES114 (MJM1100) / pMJM33	This study
MJM2251	ES114 (MJM1100) Δ*binK*	(18)
MJM2386	ES114 (MJM1100) / pBinK	This study
MJM2997	MB11B1 (MJM1130) / pVSV104	This study
MJM2998	MB11B1 (MJM1130) / pBinK	This study
MJM2999	ES213 (MJM1117) / pVSV104	This study
MJM3000	ES213 (MJM1117) / pBinK	This study
MJM3010	ES114 (MJM1100) Δ*rscS*	This study
MJM3017	ES213 (MJM1117) Δ*rscS*	This study
MJM3042	MB15A4 (MJM2114) Δ*rscS*	This study
MJM3046	MB11B1 (MJM1130) Δ*rscS*	This study
MJM3062	ES114 (MJM1100) Δ*syp*	This study
MJM3065	MB11B1 (MJM1130) Δ*syp*	This study
MJM3068	ES213 (MJM1117) Δ*syp*	This study

MJM3071	MB15A4 (MJM2114) Δ*syp*	This study
MJM3084	MB11B1 (MJM1130) Δ*binK*	This study
MJM3354	ES114 (MJM1100) *sypE*(ntG33Δ)	This study
MJM3364	ES114 (MJM1100) *sypE*(ntG33Δ) / pKV69	This study
MJM3365	ES114 (MJM1100) *sypE*(ntG33Δ) / pEAH73	This study
MJM3370	MB11B1 (MJM1130) / pKV69	This study
MJM3371	MB11B1 (MJM1130) / pEAH73	This study
MJM3394	ES114 (MJM1100) Δ*rscS* *sypE*(ntG33Δ)	This study
MJM3397	MB11B1 (MJM1130) *sypE*(nt33::G)	This study
MJM3398	MB11B1 (MJM1130) *sypE*(nt33::G) / pKV69	This study
MJM3399	MB11B1 (MJM1130) *sypE*(nt33::G) / pEAH73	This study
MJM3410	MB11B1 (MJM1130) Δ*sypE*	This study
MJM3411	MB11B1 (MJM1130) Δ*sypE* / pKV69	This study
MJM3412	MB11B1 (MJM1130) Δ*sypE* / pEAH73	This study
MJM3417	ES114 (MJM1100) Δ*sypE*	This study
MJM3418	ES114 (MJM1100) Δ*sypE* / pKV69	This study
MJM3419	ES114 (MJM1100) Δ*sypE* / pEAH73	This study

MJM3423	ES114 (MJM1100) Δ*rscS* Δ*sypE*	This study
MJM3455	ES114 (MJM1100) / pEAH73	This study
MJM3501	SR5 (MJM1125) Δ*syp*	This study
MJM3751	SR5 (MJM1125) ΔbinK::*erm*	This study
E. coli		
MJM534	CC118 λpir / pEVS104	(58)
MJM537	DH5α λpir	Lab stock
MJM570	DH5α / pEVS79	(58)
MJM580	DH5α λpir / pVSV104	(59)
MJM581	DH5α / pKV69	(60)
MJM583	DH5α / pKG11	(15)
MJM639	XL1-Blue / pMJM33	This study
MJM658	BW23474 / pEVS107	(61)
MJM2384	DH5α λpir / pBinK	(18)
MJM2540	KV5264 / pEAH73	(39)
MJM3008	DH5α / pEVS79-Δ*rscS*[MJM1100]	This study
MJM3014	DH5α λpir / pEVS79-Δ*rscS*[MJM1117]	This study

MJM3039	DH5α λpir / pEVS79-Δ*rscS*[MJM2114]	This study
MJM3043	DH5α λpir / pEVS79-Δ*rscS*[MJM1130]	This study
MJM3060	NEB5α / pEVS79-Δ*syp*[MJM1100]	This study
MJM3063	NEB5α / pEVS79-Δ*syp*[MJM1130]	This study
MJM3066	DH5α λpir / pEVS79-Δ*syp*[MJM1117]	This study
MJM3069	DH5α λpir / pEVS79-Δ*syp*[MJM2114]	This study
MJM3082	NEB5α / pEVS79-Δ*binK*[MJM1130]	This study
MJM3287	NEB5α / pHB1	This study
MJM3338	DH5α λpir / pEVS107-*sypE*[MJM1130](nt33::G)	This study
MJM3340	DH5α λpir / pEVS107-*sypE*[MJM1100](ntG33Δ)	This study
MJM3351	NEB5α / pEVS79-*sypE*[MJM1130](nt33::G)	This study
MJM3352	NEB5α / pEVS79-*sypE*[MJM1100](ntG33Δ)	This study
MJM3409	NEB5α / pEVS79-Δ*sypE*[MJM1130]	This study
MJM3416	NEB5α / pEVS79-Δ*sypE*[MJM1100]	This study

702 **Table 2. Plasmids.**

Plasmid	Relevant genotype	Source/Reference
pEVS79	Vector backbone (Cam^R) for deletion	(58)

	construction	
pKV69	Vector backbone (Cam^R/Tet^R)	(60)
pKG11	pKV69 carrying *rscS1*	(15)
pMJM33	pKG11 *rscS1*(ntA1141::Δ)	This study
pEVS104	Conjugation helper plasmid (Kan^R)	(58)
pEVS107	Mini-Tn7 mobilizable vector (Erm^R/Kan^R)	(61)
pEAH73	pKV69 carrying *sypG* from ES114	(39)
pVSV104	Complementation vector (Kan^R)	(59)
pBinK	pVSV104 carrying *binK* from MJM1100	(18)
pHB1	pUC19 FRT-*erm*-FRT	This study
pEVS79-Δ*rscS*[MJM1100]	pEVS79 carrying 1.6 kb US/1.6 kb DS of *rscS* from MJM1100	This study
pEVS79-Δ*rscS*[MJM1117]	pEVS79 carrying 1.6 kb US/1.6 kb DS of *rscS* from MJM1117	This study
pEVS79-Δ*rscS*[MJM2114]	pEVS79 carrying 1.6 kb US/1.6 kb DS of *rscS* from MJM2114	This study
DH5α λpir / pEVS79-Δ*rscS*[MJM1130]	pEVS79 carrying 1.6 kb US/1.6 kb DS of *rscS* from MJM1130	This study
pEVS79-Δ*syp*[MJM1100]	pEVS79 carrying 1.6 kb US of *sypA*/1.6 kb DS of *sypR* from MJM1100	This study
pEVS79-Δ*syp*[MJM1130]	pEVS79 carrying 1.6 kb US of *sypA*/1.6 kb DS of *sypR* from MJM1130	This study
pEVS79-Δ*syp*[MJM1117]	pEVS79 carrying 1.6 kb US of *sypA*/1.6 kb DS of *sypR* from MJM1117	This study

pEVS79-Δ*syp*[MJM2114]	pEVS79 carrying 1.6 kb US of *sypA*/1.6 kb DS of *sypR* from MJM2114	This study
pEVS79-Δ*binK*[MJM1130]	pEVS79 carrying 1.6 kb US/1.6 kb DS of *binK* from MJM1130	This study
pEVS107-*sypE*[MJM1130](nt33::G)	pEVS107 carrying the *sypE*(nt33::G) allele from MJM1130	This study
pEVS107-*sypE*[MJM1100](ntG33Δ)	pEVS107 carrying the *sypE*(ntG33Δ) allele from MJM1100	This study
pEVS79-*sypE*[MJM1130](nt33::G)	pEVS79 carrying the *sypE*(nt33::G) allele from MJM1130	This study
pEVS79-*sypE*[MJM1100](ntG33Δ)	pEVS79 carrying the *sypE*(ntG33Δ) allele from MJM1100	This study
pEVS79-Δ*sypE*[MJM1130]	pEVS79 carrying 1.6 kb US/1.6 kb DS of *sypE* from MJM1130	This study
pEVS79-Δ*sypE*[MJM1100]	pEVS79 carrying 1.6 kb US/1.6 kb DS of *sypE* from MJM1100	This study

705   **Table 3. DNA oligonucleotides for PCR amplification and sequencing.**

Primer name	Sequence (5' to 3')
DAT_015F	ACCAAGAAGCAGTACGACGATTAT
ES114_DS_ver	GGATGTTTTAGATGTTGCGG
ES114_indel_for	TTACTTTTTTCAGATACAAAGCCC
ES114_indel_rev	GTTGTTCTGATAGTGCGTGA
ES114_US_ver	ATCAACTCAAGAAACTCCCC

for_ver_sypE	CCGGCTCAAACTATTGCAG
Gib_ES114_binK_DS_for	attaatcgatGCGTATACATAAATAATGATTCATATATAC
Gib_ES114_binK_DS_rev	gcaggaattcgatatcaagcTTTCAATACTGTGTTTTTATGC
Gib_ES114_binK_US_for	gaggtcgacggtatcgataaGAGCCTTTTAAATCCCCTAAC
Gib_ES114_binK_US_rev	atgtatacgcATCGATTAATGACATATTATTATTCATAAAAAAC
Gib_ES114_rscS_DS_for	taatgcaatgGAGAAGTATGAAACACAATAAAC
Gib_ES114_rscS_DS_rev	gcaggaattcgatatcaagcAAAAATACATTGTTGCACTTG
Gib_ES114_rscS_US_for	gaggtcgacggtatcgataaGACGTCTAAAACTGAATCG
Gib_ES114_rscS_US_rev	catacttctcCATTGCATTAGCTCCTATAAAATAG
Gib_ES114_syp_DS_for	gcttattatgATATTTGCTCGAGGCCAATAAAAAC
Gib_ES114_syp_DS_rev	gcaggaattcgatatcaagcTGGTGAATGTAGGATCCAC
Gib_ES114_syp_US_for	gaggtcgacggtatcgataaCAACCGTAGCGCCAAATG
Gib_ES114_syp_US_rev	gagcaaatatCATAATAAGCTCCTAGGGAATAATC
Gib_ES114_sypE_C_for	cagatacaaaCCCACATCACTAGAGTCG
Gib_ES114_sypE_C_rev	ctagtggccaggtacctcgaAATTAAGCTTCCATCTTCAC
Gib_ES114_sypE_DS_for	tgtaatcatgCTGTTAATTGAGAATCAATAAAAAG
Gib_ES114_sypE_DS_rev	caactcttttttccgaaggtaTTGAGTAACCGGCATAATTTAG

Gib_ES114_sypE_N_for	tagagggccctaggcgcgccTGTTTCACAACTCAATACC
Gib_ES114_sypE_N_rev	gtgatgtgggTTTGTATCTGAAAAAAGTAAAGTAG
Gib_ES114_sypE_US_for	gaggtcgacggtatcgataaTGGTCAGATGAAATGTCATTTTTAG
Gib_ES114_sypE_US_rev	caattaacagCATGATTACACCACTGTTG
Gib_ES213_rscS_US_rev	catacttctcCATTGTATTAGCTCCTATAAAATAG
Gib_MB11B1_syp_DS_for	gcttattatgATATTTGCTCGAGGTCAATAAAAG
Gib_MB11B1_syp_US_for	gaggtcgacggtatcgataaGCACACTGATAACTAAATTATTAC
Gib_MB11B1_syp_US_rev	gagcaaatatCATAATAAGCTCCTAGGG
Gib_MB11B1_sypE_C_for	cagatacaaaGCCAACATCACTAGAATC
Gib_MB11B1_sypE_C_rev	ctagtggccaggtacctcgaTCAACAATTAAGCTTCCATC
Gib_MB11B1_sypE_DS_for	cagtggtatgCTGTTAATTGAAAACCAATAGC
Gib_MB11B1_sypE_DS_rev	gcaggaattcgatatcaagcATTTAGGATGTTTTTAATAACAATTTG
Gib_MB11B1_sypE_N_for	tagagggccctaggcgcgccAGTTTCACAACTCAATACTAATAATATTC
Gib_MB11B1_sypE_N_rev	tgatgttggcTTTGTATCTGAAAAAAGCAAAATAG
Gib_MB11B1_sypE_US_for	gaggtcgacggtatcgataaGAATGGTCAGATGAAATGTC
Gib_MB11B1_sypE_US_rev	caattaacagCATACCACTGTTGATAAAAATC
Gib_pEVS79_ES_sypE_for	gaggtcgacggtatcgataaTGTTTCACAACTCAATACC

Gib_pEVS79_ES_sypE_rev	gcaggaattcgatatcaagcAATTAAGCTTCCATCTTCAC
Gib_pEVS79_MB_sypE_for	gaggtcgacggtatcgataaAGTTTCACAACTCAATACTAATAATATTC
Gib_pEVS79_MB_sypE_rev	gcaggaattcgatatcaagcTCAACAATTAAGCTTCCATC
Gib_SR5_syp_DS_for	gcttattatgATATTTGCTCGAGGACAATAAAAAG
Gib_SR5_syp_DS_rev	gcaggaattcgatatcaagcTGGTGAGTGTAGAATCCATTC
Gib_SR5_syp_US_for	gaggtcgacggtatcgataaAACCGTAGCGCCAAATGG
Gib_SR5_syp_US_rev	gagcaaatatCATAATAAGCTCCTAGGGAATAATCC
HB8	ACAAAATTTTAAGATACTGCACTATCAACACACTCTTAAG
HB9	GGGAGGAAATAATCTAGAATGCGAGAGTAGG
HB23	TTGGAGAGCCAGCTGCGTTCGCTAA
HB39	TAGGAAGCTTACGAGACGAGCTTCTTATATATGCTTCGCCAGGAAGTTCCTATTCTCTAGAAAGTATAGGAACTTCCTTAGAAGCAAACTTAAGAGTGTG
HB41	CGATCTTGTGGGTAGAGACATCCAGGTCAAGTCCAGCCCCGCTCTAGTTTGGGAATCAAGTGCATGAGCGCTGAAG
HB42	ACGAGACGAGCTTCTTATATATGCTTCGCCAG
HB146	CGATCTTGTGGGTAGAGACATC
binK-F1	GAAATTACCATGGAGCCAACAGCAAGAC
binK-R1-LUH	ctggcgaagcatatataagaagctcgtctcgtCATAAAAAACCTAGCGCTTTATTTGTAGATATAATTATTAACTATAATCGC

binK-F2-RUH	gacttgacctggatgtctctacccacaagatcgCGCTCATTGTATCTATAGAGTATGTACTGAGTTACG
binK-R2	GGCATCATTATGGCAACCATTAAAGACG
binK-FO	CCGTTAATACTGGATTATTCGCTTGAATTTGAACG
KMB_036	CCACAATAGCAGAATACAAATTCGCTG
KMB_037	CTCAAAATGACAGTCAGAGTATCGTAGGC
JFB_287	ATGGAGTTTCTACGTCAACCAGAA
JFB_287_MB11B1	ATGGAGTTTTTACGTCAACCAGAG
JFB_288	TGTTATAACGATTACATGGCAGCG
JFB_365	GGAAAGAGAATGATTAAG
M13for	GTAAAACGACGGCCAG
M13rev	CAGGAAACAGCTATGAC
MB11B1_indel_for	GCTTTTTTCAGATACAAAGCCA
MB11B1_indel_rev	ATACCTGATGGAAACGACCT
MJM-154F	TAAAAAGGGAATTAATCCGC
MJM-306R	AACTCTAACCAAGAAGCA
pEVS107_3837	GGCGCGCCTAGGGCCCTC
pEVS107_3838	TCGAGGTACCTGGCCACTAG

pEVS79_for_691	GCTTGATATCGAATTCCTG
pEVS79_rev_690	TTATCGATACCGTCGACC
rev_ver_sypE	TTCACCATGAGTGCCAAATC
rscS_del1F	CTTATCTTCTAGTTCTTTTTTTTAGTGATGTCTCTTTCTACGGC
rscS_del1R	GCCGTAGAAAGAGACATCACTAAAAAAAAGAACTAGAAGATAAG
rscS_ver_1	GTAATTCAGTAATGCTACC
rscS_ver_2	GTCGCACCGTCAGGTATA
rscS_ver_3	AAGAAATTATTCGCTACC
rscS_ver_4	AGTTAGTAGGCCATTACG
SR5_syp_ver_for	TAGGCGTATCAAAAACCACCT
SR5_syp_ver_rev	TCAGGAATGTCGATGGCAG
Syp_ver_DS_rev	ATCGAGCATATTTTGCCAATC
Syp_ver_US_for	ACCTATCAACTCTTAAGTCGATTC
syp4F	TGAGGATCCCATCGTGCCATA
syp4R	AGCTCCTTTGCAATGTTTGCTT
syp5F	TATTAGGCCGTTTCCACCAGG
syp5F-B	TATTAGGTCGTTTCCATCAGG

sypA_out	AACAGGAATTGCGTTTTCAA
US_syp_flank_for	ACCACTGTGATAACTTGCAC
US_syp_flank_rev	ATGAGGCATAACCTGTTCCA

For Gibson assembly primers, capital letters indicate homology to the template. All primers were

designed for this study except MJM-154F, MJM-306R (22); JFB_287, JFB_288, and JFB_365

(18); and M13 for, M13 rev.

ACKNOWLEDGMENTS

The authors thank Elizabeth Bacon, Jacklyn Duple, Cheeneng Moua, Lynn Naughton, Olivia

Sauls, and Denise Tarnowski for assistance with experiments.

FUNDING INFORMATION

This work was funded by NIH grants R35GM119627 (to M.J.M.) and R21AI117262 (M.J.M.) and

NSF grant IOS-1757297 (M.J.M.). Support for trainees was provided on NIGMS grants

T32GM008061 (J.F.B.) and T32GM008349 (K.M.B.). This work was funded by the Chicago

Biomedical Consortium with support from the Searle Funds at The Chicago Community Trust

(supporting E.R.R.).

REFERENCES

1.   Long SR. 1996. *Rhizobium* symbiosis: nod factors in perspective. Plant Cell 8:1885–1898.

2.   Roche P, Maillet F, Plazanet C, Debellé F, Ferro M, Truchet G, Promé JC, Dénarié J. 1996.

The common *nodABC* genes of *Rhizobium meliloti* are host-range determinants. Proc Natl

Acad Sci U S A 93:15305–15310.

3.   Costello EK, Lauber CL, Hamady M, Fierer N, Gordon JI, Knight R. 2009. Bacterial

community variation in human body habitats across space and time. Science 326:1694–

1697.

4.   Grice EA, Kong HH, Conlan S, Deming CB, Davis J, Young AC, NISC Comparative

Sequencing Program, Bouffard GG, Blakesley RW, Murray PR, Green ED, Turner ML,

Segre JA. 2009. Topographical and temporal diversity of the human skin microbiome.

Science 324:1190–1192.

5.   Mark Welch JL, Rossetti BJ, Rieken CW, Dewhirst FE, Borisy GG. 2016. Biogeography of a

human oral microbiome at the micron scale. Proc Natl Acad Sci U S A 113:E791–E800.

6.   Ruby EG. 2008. Symbiotic conversations are revealed under genetic interrogation. Nat Rev

Microbiol 6:752–762.

7.   Nyholm SV, McFall-Ngai MJ. 2004. The winnowing: establishing the squid-vibrio symbiosis.

Nat Rev Microbiol 2:632–642.

8.   Visick KL, Ruby EG. 2006. *Vibrio fischeri* and its host: it takes two to tango. Curr Opin

Microbiol 9:632–638.

9.   Mandel MJ, Schaefer AL, Brennan CA, Heath-Heckman EAC, Deloney-Marino CR, McFall-

Ngai MJ, Ruby EG. 2012. Squid-derived chitin oligosaccharides are a chemotactic signal

during colonization by *Vibrio fischeri*. Appl Environ Microbiol 78:4620–4626.

10. Jones BW, Nishiguchi MK. 2004. Counterillumination in the Hawaiian bobtail squid,

Euprymna scolopes Berry (Mollusca: Cephalopoda). Mar Biol 144:1151–1155.

11. Ruby EG, McFall-Ngai MJ. 1992. A squid that glows in the night: development of an animal-
bacterial mutualism. J Bacteriol 174:4865–4870.

12. Wier AM, Nyholm SV, Mandel MJ, Massengo-Tiassé RP, Schaefer AL, Koroleva I, Splinter-
Bondurant S, Brown B, Manzella L, Snir E, Almabrazi H, Scheetz TE, Bonaldo M de F,
Casavant TL, Soares MB, Cronan JE, Reed JL, Ruby EG, McFall-Ngai MJ. 2010.
Transcriptional patterns in both host and bacterium underlie a daily rhythm of anatomical
and metabolic change in a beneficial symbiosis. Proc Natl Acad Sci U S A 107:2259–2264.

13. Visick KL. 2009. An intricate network of regulators controls biofilm formation and
colonization by *Vibrio fischeri*. Mol Microbiol 74:782–789.

14. Yip ES, Grublesky BT, Hussa EA, Visick KL. 2005. A novel, conserved cluster of genes
promotes symbiotic colonization and σ^{54}-dependent biofilm formation by *Vibrio fischeri*. Mol
Microbiol 57:1485–1498.

15. Yip ES, Geszvain K, DeLoney-Marino CR, Visick KL. 2006. The symbiosis regulator RscS
controls the *syp* gene locus, biofilm formation and symbiotic aggregation by *Vibrio fischeri*.
Mol Microbiol 62:1586–1600.

16. Shibata S, Yip ES, Quirke KP, Ondrey JM, Visick KL. 2012. Roles of the Structural
Symbiosis Polysaccharide (*syp*) Genes in Host Colonization, Biofilm Formation, and
Polysaccharide Biosynthesis in *Vibrio fischeri*. J Bacteriol 194:6736–6747.

17. Morris AR, Visick KL. 2013. The response regulator SypE controls biofilm formation and
colonization through phosphorylation of the syp-encoded regulator SypA in *Vibrio fischeri*.
Mol Microbiol 87:509–525.

18. Brooks JF 2nd, Mandel MJ. 2016. The Histidine Kinase BinK Is a Negative Regulator of
Biofilm Formation and Squid Colonization. J Bacteriol 198:2596–2607.

771 19. Tischler AH, Lie L, Thompson CM, Visick KL. 2018. Discovery of Calcium as a Biofilm-

772   Promoting Signal for Vibrio fischeri Reveals New Phenotypes and Underlying Regulatory

773   Complexity. J Bacteriol 200:e00016–18.

774 20. Pankey MS, Foxall RL, Ster IM, Perry LA, Schuster BM, Donner RA, Coyle M, Cooper VS,

775   Whistler CA. 2017. Host-selected mutations converging on a global regulator drive an

776   adaptive leap towards symbiosis in bacteria. Elife 6:e24414.

777 21. Thompson CM, Tischler AH, Tarnowski DA, Mandel MJ, Visick KL. 2018. Nitric oxide

778   inhibits biofilm formation by *Vibrio fischeri* via the nitric oxide-sensor HnoX. Mol Microbiol

779   doi:10.1111/mmi.14147.

780 22. Mandel MJ, Wollenberg MS, Stabb EV, Visick KL, Ruby EG. 2009. A single regulatory gene

781   is sufficient to alter bacterial host range. Nature 458:215–218.

782 23. Nyholm SV, Nishiguchi MK. 2008. The evolutionary ecology of a sepiolid squid-vibrio

783   association: from cell to environment. Vie Milieu Paris 58:175–184.

784 24. Fidopiastis PM, von Boletzky S, Ruby EG. 1998. A new niche for *Vibrio logei*, the

785   predominant light organ symbiont of squids in the genus *Sepiola*. J Bacteriol 180:59–64.

786 25. Ruby EG, Lee KH. 1998. The *Vibrio fischeri-Euprymna scolopes* Light Organ Association:

787   Current Ecological Paradigms. Appl Environ Microbiol 64:805–812.

788 26. Mandel MJ. 2010. Models and approaches to dissect host-symbiont specificity. Trends

789   Microbiol 18:504–511.

790 27. Gyllborg MC, Sahl JW, Cronin DC 3rd, Rasko DA, Mandel MJ. 2012. Draft genome

791   sequence of *Vibrio fischeri* SR5, a strain isolated from the light organ of the Mediterranean

792   squid *Sepiola robusta*. J Bacteriol 194:1639.

28. Geszvain K, Visick KL. 2008. Multiple factors contribute to keeping levels of the symbiosis

regulator RscS low. FEMS Microbiol Lett 285:33–39.

29. Wollenberg MS, Ruby EG. 2009. Population structure of *Vibrio fischeri* within the light

organs of *Euprymna scolopes* squid from Two Oahu (Hawaii) populations. Appl Environ

Microbiol 75:193–202.

30. Wollenberg MS, Ruby EG. 2012. Phylogeny and fitness of *Vibrio fischeri* from the light

organs of *Euprymna scolopes* in two Oahu, Hawaii populations. ISME J 6:352–362.

31. Elliott KT, DiRita VJ. 2008. Characterization of CetA and CetB, a bipartite energy taxis

system in *Campylobacter jejuni*. Mol Microbiol 69:1091–1103.

32. Antonov I, Coakley A, Atkins JF, Baranov PV, Borodovsky M. 2013. Identification of the

nature of reading frame transitions observed in prokaryotic genomes. Nucleic Acids Res

41:6514–6530.

33. Bongrand C, Koch EJ, Moriano-Gutierrez S, Cordero OX, McFall-Ngai M, Polz MF, Ruby

EG. 2016. A genomic comparison of 13 symbiotic *Vibrio fischeri* isolates from the

perspective of their host source and colonization behavior. ISME J 10:2907–2917.

34. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997.

Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.

Nucleic Acids Res 25:3389–3402.

35. Rusch DB, Rowe-Magnus DA. 2017. Complete Genome Sequence of the Pathogenic

Vibrio vulnificus Type Strain ATCC 27562. Genome Announc 5:e00907–17.

36. Morris AR, Darnell CL, Visick KL. 2011. Inactivation of a novel response regulator is

necessary for biofilm formation and host colonization by *Vibrio fischeri*. Mol Microbiol

82:114–130.

37. Guo Y, Rowe-Magnus DA. 2011. Overlapping and unique contributions of two conserved

polysaccharide loci in governing distinct survival phenotypes in Vibrio vulnificus. Environ

Microbiol 13:2888–2990.

38. Morris AR, Visick KL. 2013. Inhibition of SypG-induced biofilms and host colonization by the

negative regulator SypE in *Vibrio fischeri*. PLoS One 8:e60076.

39. Hussa EA, Darnell CL, Visick KL. 2008. RscS functions upstream of SypG to control the

syp locus and biofilm formation in *Vibrio fischeri*. J Bacteriol 190:4576–4583.

40. Koehler S, Gaedeke R, Thompson C, Bongrand C, Visick KL, Ruby E, McFall-Ngai M.

2018. The model squid-vibrio symbiosis provides a window into the impact of strain- and

species-level differences during the initial stages of symbiont engagement. Environ

Microbiol doi:10.1111/1462–2920.14392.

41. Ray VA, Visick KL. 2012. LuxU connects quorum sensing to biofilm formation in *Vibrio*

fischeri. Mol Microbiol 86:954–970.

42. Giraud E, Moulin L, Vallenet D, Barbe V, Cytryn E, Avarre J-C, Jaubert M, Simon D,

Cartieaux F, Prin Y, Bena G, Hannibal L, Fardoux J, Kojadinovic M, Vuillet L, Lajus A,

Cruveiller S, Rouy Z, Mangenot S, Segurens B, Dossat C, Franck WL, Chang W-S,

Saunders E, Bruce D, Richardson P, Normand P, Dreyfus B, Pignol D, Stacey G, Emerich

D, Verméglio A, Médigue C, Sadowsky M. 2007. Legumes symbioses: absence of Nod

genes in photosynthetic bradyrhizobia. Science 316:1307–1312.

43. Bonaldi K, Gargani D, Prin Y, Fardoux J, Gully D, Nouwen N, Goormachtig S, Giraud E.

2011. Nodulation of *Aeschynomene afraspera* and *A. indica* by photosynthetic

Bradyrhizobium Sp. strain ORS285: the Nod-dependent versus the Nod-independent

symbiotic interaction. Mol Plant Microbe Interact 24:1359–1371.

44. Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new

heuristics and parallel computing. Nat Methods 9:772.

45. Swofford DL. 2003. PAUP*: Phylogenetic Analysis Using Parsimony (* and Other Methods)

4th edn. Sinauer.

46. Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L,

Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic

inference and model choice across a large model space. Syst Biol 61:539–542.

47. Ronquist F, van der Mark P, Huelsenbeck JP. 2009. Bayesian phylogenetic analysis using

MrBayes, 2nd edn. Cambridge University Press.

48. Visick KL, Hodge-Hanson KM, Tischler AH, Bennett AK, Mastrodomenico V. 2018. Tools

for Rapid Genetic Engineering of *Vibrio fischeri*. Appl Environ Microbiol 84:e00850–18.

49. Pollack-Berti A, Wollenberg MS, Ruby EG. 2010. Natural transformation of *Vibrio fischeri*

requires *tfoX* and *tfoY*. Environ Microbiol 12:2302–2311.

50. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M,

Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene

knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

51. Naughton LM, Mandel MJ. 2012. Colonization of *Euprymna scolopes* squid by *Vibrio*

fischeri. J Vis Exp e3758.

52. Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M,

Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The Pfam protein

families database: towards a more sustainable future. Nucleic Acids Res 44:D279–85.

53. Lee K-H. 1994. Ecology of *Vibrio fisheri*: the light organ symbiont of the Hawaiian sepiolid

squid *Euprymna scolopes*. University of Southern California.

54. Boettcher KJ, Ruby EG. 1990. Depressed light emission by symbiotic *Vibrio fischeri* of the

sepiolid squid *Euprymna scolopes*. J Bacteriol 172:3701–3706.

55. Boettcher KJ, Ruby EG. 1994. Occurrence of plasmid DNA in the sepiolid squid symbiont

Vibrio fischeri. Curr Microbiol 29:279–286.

56. Nishiguchi MK, Ruby EG, McFall-Ngai MJ. 1998. Competitive dominance among strains of

luminous bacteria provides an unusual form of evidence for parallel evolution in Sepiolid

squid-vibrio symbioses. Appl Environ Microbiol 64:3209–3213.

57. Nishiguchi MK, Nair VS. 2003. Evolution of symbiosis in the Vibrionaceae: a combined

approach using molecules and physiology. Int J Syst Evol Microbiol 53:2019–2026.

58. Stabb EV, Ruby EG. 2002. RP4-based plasmids for conjugation between *Escherichia coli*

and members of the *Vibrionaceae*. Methods Enzymol 358:413–426.

59. Dunn AK, Millikan DS, Adin DM, Bose JL, Stabb EV. 2006. New *rfp*- and pES213-derived

tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux* expression in

situ. Appl Environ Microbiol 72:802–810.

60. Visick KL, Skoufos LM. 2001. Two-component sensor required for normal symbiotic

colonization of *Euprymna scolopes* by *Vibrio fischeri*. J Bacteriol 183:835–842.

61. McCann J, Stabb EV, Millikan DS, Ruby EG. 2003. Population dynamics of *Vibrio fischeri*

during infection of *Euprymna scolopes*. Appl Environ Microbiol 69:5928–5934.

Group C

i 100 99
  100

ii 100 99
   100

iii 100 84
    89

iv 100 85
   89

Group B

Group A

Acquisition
of rscS

Bayes ML
MP

Isolation
Habitat

Squid
Fish
Ocean

Geography
Mediterranean
Japan
Australia
Atlantic
Hawaii

0.01

A

TM   TM   RscS

1   PAS   DHp   CA   REC   HPT   927

                       277   381 402 467   514   630   661   772   834   920

```
EM17    Group B    TACATTGCCGTAGAAAGAGACATCACTAAAAAAAAAGAACTAGAAGATAAGTTAATAAAA
ES114         B    TACATTGCCGTAGAAAGAGACATCACTAAAAAAAAAGAACTAGAAGATAAGTTAATAAAA
MB15A4        B    TACATTGCCGTAGAAAGAGACATCACTAAAAAAAAAGAACTAGAAGATAAGTTAATAAAA
MB11B1  Group A    TACATTGCCGTAGAAAGAGACATCACT-AAAAAAAAAGAACTAGAAGATAAGTTAATAAAA
ES213         A    TACATTGCCGTAGAAAGAGACATCACT-AAAAAAAAAGAACTAGAAGATAAGTTAATAAAA
KB2B1         A    TACATTGCCGTAGAAAGAGACATCACT-AAAAAAAAAGAACTAGAAGATAAGTTAATAAAA
```

B

Vector   *rscS1*   *rscS1^{FS}*

Table S1. Primers pairs for construction of mutants.

Strain	Note	Forward Primer	Reverse Primer	PCR Product Size	Description of PCR product	E. coli with clone
MJM3010 =		Gib_ES114_rscS_US_for	Gib_ES114_rscS_US_rev	1.6 kb	Gibson assembly cloning of upstream *rscS* region in MJM1100 into pEVS79	**MJM3008**
MJM1100 Δ*rscS* and		Gib_ES114_rscS_DS_for	Gib_ES114_rscS_DS_rev	1.6 kb	Gibson assembly cloning of downstream *rscS* region in MJM1100 into pEVS79	
MJM3394 =		ES114_US_ver	ES114_DS_ver	586 bp	Verification of Δ*rscS* by flanking the upstream/downstream junction	
MJM3354 Δ*rscS*		ES114_US_ver	DAT015F	520 bp	Verification of Δ*rscS*; no product due to one primer binding within *rscS*.	
		Gib_ES114_rscS_US_for	Gib_ES213_rscS_US_rev	1.6 kb	Gibson assembly cloning of upstream *rscS* region in MJM1117 into pEVS79	**MJM3014**
MJM3017 =		Gib_ES114_rscS_DS_for	Gib_ES114_rscS_DS_rev	1.6 kb	Gibson assembly cloning of downstream *rscS* region in MJM1117 into pEVS79	
MJM1117 Δ*rscS*		ES114_US_ver	ES114_DS_ver	586 bp	Verification of Δ*rscS* by flanking the upstream/downstream junction	
		ES114_US_ver	DAT015F	520 bp	Verification of Δ*rscS*; no product due to one primer binding within *rscS*.	
		Gib_ES114_rscS_US_for	Gib_ES114_rscS_US_rev	1.6 kb	Gibson assembly cloning of upstream *rscS* region in MJM1130 into pEVS79	**MJM3043**
MJM3046 =		Gib_ES114_rscS_DS_for	Gib_ES114_rscS_DS_rev	1.6 kb	Gibson assembly cloning of downstream *rscS* region in MJM1130 into pEVS79	
MJM1130 Δ*rscS*		ES114_US_ver	ES114_DS_ver	586 bp	Verification of Δ*rscS* by flanking the upstream/downstream junction	
		ES114_US_ver	DAT015F	520 bp	Verification of Δ*rscS*; no product due to one primer binding within *rscS*	
		Gib_ES114_rscS_US_for	Gib_ES114_rscS_US_rev	1.6 kb	Gibson assembly cloning of upstream *rscS* region in MJM2114 into pEVS79	**MJM3039**
MJM3042 =		Gib_ES114_rscS_DS_for	Gib_ES114_rscS_DS_rev	1.6 kb	Gibson assembly cloning of downstream *rscS* region in MJM2114 into pEVS79	
MJM2114 Δ*rscS*		ES114_US_ver	ES114_DS_ver	586 bp	Verification of Δ*rscS* by flanking the upstream/downstream junction	
		ES114_US_ver	DAT015F	520 bp	Verification of Δ*rscS*; no product due to one primer binding within *rscS*	
		Gib_ES114_syp_US_for	Gib_ES114_syp_US_rev	1.6 kb	Gibson assembly cloning of upstream *sypA* region in MJM1100 into pEVS79	**MJM3060**
MJM3062 =		Gib_ES114_syp_DS_for	Gib_ES114_syp_DS_rev	1.6 kb	Gibson assembly cloning of downstream *sypR* region in MJM1100 into pEVS79	
MJM1100 Δ*sypA-R*		Syp_ver_US_for	Syp_ver_DS_rev	503 bp	Verification of Δ*sypA-R* by flanking the upstream/downstream junction	
		Syp_ver_US_for	sypA_out	490 bp	Verification of Δ*sypA-R*; no product due to one primer binding within *sypA*	
		Gib_ES114_syp_US_for	Gib_ES114_syp_US_rev	1.6 kb	Gibson assembly cloning of upstream *sypA* region in MJM1117 into pEVS79	**MJM3066**
MJM3068 =		Gib_MB11B1_syp_DS_for	Gib_ES114_syp_DS_rev	1.6 kb	Gibson assembly cloning of downstream *sypR* region in MJM1117 into pEVS79	
MJM1117 Δ*sypA-R*		Syp_ver_US_for	Syp_ver_DS_rev	503 bp	Verification of Δ*sypA-R* by flanking the upstream/downstream junction	
		Syp_ver_US_for	sypA_out	490 bp	Verification of Δ*sypA-R*; no product due to one primer binding within *sypA*	
		Gib_SR5_syp_US_for	Gib_SR5_syp_US_rev	1.6 kb	Gibson assembly cloning of upstream *sypA* region in MJM1125 into pEVS79	**MJM3500**
MJM3501 =		Gib_SR5_syp_DS_for	Gib_SR5_syp_DS_rev	1.6 kb	Gibson assembly cloning of downstream *sypR* region in MJM1125 into pEVS79	
MJM1125 Δ*sypA-R*		SR5_syp_ver_for	SR5_syp_ver_rev	506 bp	Verification of Δ*sypA-R* by flanking the upstream/downstream junction	
		Gib_SR5_syp_US_for	sypA_out	1.9 kb	Verification of Δ*sypA-R*; no product due to one primer binding within *sypA*	
		Gib_MB11B1_syp_US_for	Gib_MB11B1_syp_US_rev	1.6 kb	Gibson assembly cloning of upstream *sypA* region in MJM1130 into pEVS79	**MJM3063**
MJM3065 =		Gib_MB11B1_syp_DS_for	Gib_ES114_syp_DS_rev	1.6 kb	Gibson assembly cloning of downstream *sypR* region in MJM1130 into pEVS79	
MJM1130 Δ*sypA-R*		Syp_ver_US_for	Syp_ver_DS_rev	503 bp	Verification of Δ*sypA-R* by flanking the upstream/downstream junction	
		Syp_ver_US_for	sypA_out	490 bp	Verification of Δ*sypA-R*; no product due to one primer binding within *sypA*	
		Gib_ES114_syp_US_for	Gib_ES114_syp_US_rev	1.6 kb	Gibson assembly cloning of upstream *sypA* region in MJM2114 into pEVS79	**MJM3069**
MJM3071 =		Gib_ES114_syp_DS_for	Gib_ES114_syp_DS_rev	1.6 kb	Gibson assembly cloning of downstream *sypR* region in MJM2114 into pEVS79	
MJM2114 Δ*sypA-R*		Syp_ver_US_for	Syp_ver_DS_rev	503 bp	Verification of Δ*sypA-R* by flanking the upstream/downstream junction	
		Syp_ver_US_for	sypA_out	490 bp	Verification of Δ*sypA-R*; no product due to one primer binding within *sypA*	
		Gib_ES114_binK_US_for	Gib_ES114_binK_US_rev	1.6 kb	Gibson assembly cloning of upstream *binK* region in MJM1130 into pEVS79	**MJM3082**
MJM3084 =		Gib_ES114_binK_DS_for	Gib_ES114_binK_DS_rev	1.6 kb	Gibson assembly cloning of downstream *binK* region in MJM1130 into pEVS79	
MJM1130 Δ*binK*		JFB_287_MB11B1	JFB_288	767 bp	Verification of Δ*binK* by flanking the upstream/downstream junction	
		JFB_287_MB11B1	JFB_365	624 bp	Verification of Δ*binK*; no product due to one primer binding within *binK*	
MJM3417 =		Gib_ES114_sypE_US_for	Gib_ES114_sypE_US_rev	1.6 kb	Gibson assembly cloning of upstream *sypE* region in MJM1100 into pEVS79	**MJM3416**
MJM1100 Δ*sypE* and		Gib_ES114_sypE_DS_for	Gib_ES114_sypE_DS_rev	1.6 kb	Gibson assembly cloning of downstream *sypE* region in MJM1100 into pEVS79	
MJM3423 =		syp4F	syp4R	780 bp	Verification of Δ*sypE* by flanking the upstream/downstream junction	
MJM3010 Δ*sypE*		syp5F	syp4R	772 bp	Verification of Δ*sypE*; no product due to one primer binding within *sypE*	
		Gib_MB11B1_sypE_US_for	Gib_MB11B1_sypE_US_rev	1.6 kb	Gibson assembly cloning of upstream *sypE* region in MJM1130 into pEVS79	**MJM3409**
MJM3410 =		Gib_MB11B1_sypE_DS_for	Gib_MB11B1_sypE_DS_rev	1.6 kb	Gibson assembly cloning of downstream *sypE* region in MJM1130 into pEVS79	
MJM1130 Δ*sypE*		for_ver_sypE	rev_ver_sypE	732 bp	Verification of Δ*sypE* by flanking the upstream/downstream junction	
		MB11B1_indel_for	MB11B1_indel_rev	779 bp	Verification of Δ*sypE*; no product due to both primers binding within *sypE*	
		Gib_ES114_sypE_N_for	Gib_ES114_sypE_N_rev	332 bp	Gibson assembly cloning of N-terminal *sypE* in MJM1100 into pEVS107	**MJM3340**
MJM3354 =		Gib_ES114_sypE_C_for	Gib_ES114_sypE_C_rev	1786 bp	Gibson assembly cloning of C-terminal *sypE* in MJM1100 into pEVS107	**and**
MJM1100 *sypE* (ntG33Δ)		Gib_pEVS79_ES_sypE_for	Gib_pEVS79_ES_sypE_rev	2118 bp	Gibson assembly cloning of *sypE* (ntG33Δ) into pEVS79	**MJM3352**
		ES114_indel_for	ES114_indel_rev	802 bp	Verification of *sypE* (ntG33Δ); stronger band with original allele.	
		Gib_MB11B1_sypE_N_for	Gib_MB11B1_sypE_N_rev	332 bp	Gibson assembly cloning of N-terminal *sypE* in MJM1130 into pEVS107	**MJM3338**
MJM3397 =		Gib_MB11B1_sypE_C_for	Gib_MB11B1_sypE_C_rev	1786 bp	Gibson assembly cloning of C-terminal *sypE* in MJM1130 into pEVS107	**and**
MJM1130 *sypE* (nt33::G)		Gib_pEVS79_MB_sypE_for	Gib_pEVS79_MB_sypE_rev	2118 bp	Gibson assembly cloning *sypE* (nt33::G) into pEVS79	**MJM3351**
		MB11B1_indel_for	MB11B1_indel_rev	779 bp	Verification of *sypE* (nt33::G); stronger band with corrected allele.	

>gBlock_erm
GGTCAGCCTCTAATGGCTCGTAAGATAGTGTAGGAAGCTTTATCGAACTGCGCGAAAGATCCCGAAGTTCCTATTCTCT
AGAAAGTATAGGAACTTCCTTAGAAGCAAACTTAAGAGTGTGTTGATAGTGCAGTATCTTAAAATTTTGTATAATAGGA
ATTGAAGTTAAATTAGATGCTAAAAATTTGTAATTAAGAAGGAGTGATTACATGAACAAAAATATAAAATATTCTCAAA
ACTTTTTAACGAGTGAAAAAGTACTCAACCAAATAATAAAACAATTGAATTTAAAAGAAACCGATACCGTTTACGAAAT
TGGAACAGGTAAAGGGCATTTAACGACGAAACTGGCTAAAATAAGTAAACAGGTAACGTCTATTGAATTAGACAGTCAT
CTATTCAACTTATCGTCAGAAAAATTAAAACTGAATACTCGTGTCACTTTAATTCACCAAGATATTCTACAGTTTCAAT
TCCCTAACAAACAGAGGTATAAAATTGTTGGGAGTATTCCTTACCATTTAAGCACACAAATTATTAAAAAAGTGGTTTT
TGAAAGCCATGCGTCTGACATCTATCTGATTGTTGAAGAAGGATTCTACAAGCGTACCTTGGATATTCACCGAACACTA
GGGTTGCTCTTGCACACTCAAGTCTCGATTCAGCAATTGCTTAAGCTGCCAGCGGAATGCTTTCATCCTAAACCAAAAG
TAAACAGTGTCTTAATAAAACTTACCCGCCATACCACAGATGTTCCAGATAAATATTGGAAGCTATATACGTACTTTGT
TTCAAAATGGGTCAATCGAGAATATCGTCAACTGTTTACTAAAAATCAGTTTCATCAAGCAATGAAACACGCCAAAGTA
AACAATTTAAGTACCGTTACTTATGAGCAAGTATTGTCTATTTTTAATAGTTATCTATTATTTAACGGGAGGAAATAAT
CTAGAATGCGAGAGTAGGGAACTGCCGAAGTTCCTATTCTCTAGAAAGTATAGGAACTTCAGCGCTCATGCACTTGATT
CCGGATCCCTAATTAGCGAACGCAGCTGGCTCTCCAA