Validation of a Miniaturized Permeability Assay Compatible with CRISPR-Mediated Genome-Wide Screen

by

Claire Simonneau[1][§], Junning Yang[1], Xianguo Kong[1], Robert Kilker[1], Leonard Edelstein[1], Paolo Fortina[2], Eric Londin[3], and Arie Horowitz[*]

1 – Cardeza Center for Hematology Research, 2 – Department of Cancer Biology, 4 – Sidney Kimmel Cancer Center Genomics Laboratory, 3 – Computational Medicine Center, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, Pennsylvania, USA.

* – Corresponding author
§ – Current address: Roche Innovation Center Basel, Switzerland

Corresponding author: Arie Horowitz; email address: arie2006@gmail.com; telephone +1-617-784-2927

ABSTRACT

The impermeability of the luminal endothelial cell monolayer is crucial for the normal performance of the vascular and lymphatic systems. A key to this function is the integrity of the monolayer's intercellular junctions. The known repertoire of junction-regulating genes is incomplete. Current permeability assays are incompatible with high-throughput genome-wide screens that could identify these genes. To overcome these limitations, we designed a new permeability assay that consists of cell monolayers grown on ~150 μm microcarriers. Each microcarrier functions as a miniature individual assay of permeability (MAP). We demonstrate that false-positive results can be minimized, and that MAP sensitivity to thrombin-induced increase in monolayer permeability is similar to the sensitivity of the measurement of impedance. We validate the assay by showing that the expression of single guide RNAs (sgRNAs) that target genes encoding known thrombin signaling proteins blocks effectively thrombin-induced junction disassembly, and that MAPs carrying such cells can be separated effectively by a fluorescent probe from those that carry cells expressing non-targeting sgRNAs. These results indicate that MAPs are suitable for high-throughput experimentation and for genome-wide screens for genes that mediate the disruptive effect of thrombin on endothelial cell junctions.

INTRODUCTION

One of the new opportunities afforded by the sequencing of the human genome, as well as of many others, is the ability to run genome-wide reverse screens to identify genes that underlie diverse phenotypes. The repurposing of CRISPR/deactivated (d) Cas9 for targeting either repressive or activating modules to genomic loci provides a more versatile and specific tool than RNAi, becoming the screen method of choice[1]. In the few short years since this tool became available, it has already been applied to numerous ends, including cancer drug discovery[2], analysis of unfolded protein response[3], and the identification of cancer immunotherapy targets[4]. CRISPR/dCas9-mediated screens take advantage of the accurate sgRNA-dependent targeting of protein modules that either repress or activate transcription to proximal upstream and downstream regions near the transcription start site. Transcriptional activation has been achieved by various combinations of a range of modules, including NF-κB trans-activating subunit p65[5], tetrameric VP16 transactivation domain (VP64), heat shock factor 1, the catalytic histone acetyltransferase core domain of the human E1A-associated protein p300, and the viral replication and transcription activator [reviewed in[6]]. In contrast, transcriptional repression has been produced by a single module, the Krüppel-associated box (KRAB) domain of the zinc-finger transcription factor Kox[7]. The efficacy of the regulatory dCas9-fused module is highly dependent on the selection of the sgRNA genomic targets relative to the transcription start sites, based on rules that were derived and optimized empirically[5,8]. Activation on the order of 10^3-fold, as measured by quantifying mRNA levels of selected genes, was achieved by some of the above complexes[6], whereas transcriptional repression measured in the same way was on the order of 10^2-fold, and rarely 10^3-fold[9].

Genome-wide screens employing activating or repressive dCas9 constructs and sgRNA libraries were deployed to identify toxin or drug resistance and vulnerability-conferring genes[9-14], genes that mediate cell immune and inflammatory responses[15-18], and genes required for cell survival, proliferation, and metastasis[10,19-22]. The majority of screens performed to date phenotyped cell enrichment or depletion, though some studies measured the rates of differentiation[23], cell motility[24], and the abundance of fluorescent protein[25]. In all cases, the quantified phenotypes were individual-cell attributes. Such assays are suitable for the pooled format that, with rare exceptions[26,27], characterizes the vast majority of currently available sgRNA libraries, and enables a relatively facile and economical quantification of phenotypes. Though effective and fruitful, the current limitation to individual-cell functional assays leaves out collective cell functions which underlie physiologically essential processes. Here we validate a new functional assay we designed to facilitate adaptation of the CRISPR-mediated screen for the identification of genes that regulate the permeability of endothelial cell (EC) monolayers. The integrity of this monolayer is critical to the normal function of the vascular system because it confers the barrier function of the vessels. Since administration of thrombin produces large gaps in EC monolayers that can be detected with relative ease, we chose to validate the assay by testing if sgRNAs from an existing repressive CRISPR library[8] designed to knockdown the transcription of thrombin signaling pathway genes, become enriched in unresponsive cells. Thrombin-induced permeability is not manifest in vertebrates under normal conditions, but does occur under trauma, sepsis, and other inflammation-inducing conditions in both animal models[28-30] and patients[31,32].

Currently available permeability assays are not compatible with the requirement of genome-wide screens because they cannot accommodate the large number of samples, and because their multi-well format is unsuitable for the pooled sgRNA libraries used in practically all screens. To our knowledge, the highest sampling capacity currently available for the measurement of paracellular permeability is provided by the 96-well plate filter insert assay format. A comprehensive screen of the human genome by 10 sgRNAs and 5 technical replicates per gene

using this assay would require approximately 10^4 multi-well plates, a number that is beyond the
capacity of most research laboratories. To design a more feasible assay, we took advantage of
the microcarrier-based cell culture technique developed to maximize the number of adherent
cells grown per volume of medium. Typically, microcarriers consist of spherical beads of a
diameter in the range of 100-200 μm, made of cross-linked polymers. Commercially available
microcarriers come in a variety of polymeric matrices and surface treatments or coatings[33].
Microcarriers carrying EC monolayers had been used to study the effect of various agonists on
cell monolayer permeability by measuring the absorbance of dye into microcarriers, or in a
chromatographic format by measuring the elution time of a dye[34-37]

RESULTS
EC monolayers grown on single microcarriers serve as individual permeability assays.
We sought an in vitro permeability assay that would be compatible with the high throughput
format required for genome-wide screens to interrogate signaling pathways that regulate the
integrity of endothelial cell junctions. In the new assay, each microcarrier is treated as an
individual MAP. This facilitates high throughput testing of a large number of samples in a
relatively small volume of growth medium (approximately 6.2×10^5 microcarriers per 100 mL).
We envisioned that ECs will be transduced by repressive (CRISPRi) sgRNA libraries and grown
as clonal populations on each microcarrier. When treated by agonists that disrupt cell junctions,
e.g. vascular endothelial growth factor (VEGF) or thrombin, junctions among ECs expressing
sgRNAs that target genes required the for the cellular response to the administered agonist
would fail to disassemble. To distinguish between microcarriers carrying responsive or non-
responsive EC monolayers, we selected a probe that would bind to the microcarrier surface
exposed through the gaps between responsive cells. Since the microcarrier type that supported
EC growth optimally is coated with gelatin, we selected the collagen-binding fluorescently-
conjugated fragment of fibronectin (FN_c)[38] as a probe because of its high binding affinity to
gelatin[39]. The fluorescence of MAPs carrying responsive EC monolayers would increase upon
thrombin treatment because of the binding of fluorescently-conjugated FN_c (FN_{cf}) to the newly
formed gaps between ECs. The darker MAPs that carry non-responsive monolayers would then
be separated from the brighter ones by fluorescence-assisted sorting (Fig. 1).

Thrombin increases the permeability of telomerase-immortalized human primary EC
monolayers
Among a variety of known permeability factors, thrombin induces relatively large openings
between confluent ECs[40], on the same type of microcarriers used in this study[35]. To gauge
thrombin's effects on monolayers of telomerase-immortalized (human dermal) microvascular
endothelial (TIME) cells, we measured its impact on permeability by two approaches. In the first,
we measured the penetration of a fluorescently-conjugated dextran probe through an EC
monolayer. In the second, we measured thrombin-induced change in monolayer impedance.
Both approaches yielded large changes in the measured attributes, indicating substantial
increases in monolayer permeability. In the first type of assay, the fluorescence of the probe in
the lower chamber of the assay almost plateaued after a more than a 3-fold increase (Fig. 2a).
In impedance measurements, thrombin concentrations of 0.5, 1.0, and 2.0 U/mL reduced the
impedance by approximately one unit, although their rate of recovery to the pre-thrombin level
became slower when thrombin concentration was increased (Fig. 2b). The impedances
recovered their pre-thrombin levels and stabilized in approximately 120 min. To visualize the
structural effects of thrombin on the EC monolayer, we immunolabeled the cells to track the
localization of vascular endothelial cadherin (VEcad) and used phalloidin to observe thrombin-
induced changes in the actin cytoskeleton. Thrombin induced opening of large gaps, which
reached in some locations a width of approximately 10 μm, and formation of denser and thicker
stress fibers (Fig. 2c), similar to previous observations[41].

103
Single TIME cells grow to confluence on gelatin-coated microcarriers
Following tests with several types of microcarrier polymers (e.g. polystyrene, diethylaminoethyl dextran)**,** and coatings (e.g. fibronectin, collagen I), we determined that gelatin-coated microcarriers consisting of cross-linked dextran matrix are optimal for TIME cell growth. To observe monolayer integrity, we stained them with calcein, visualizing both the cell body and intercellular junctions (Fig. 3a). All the cells grown on each microcarrier in a genome-wide screen should express a single sgRNA to avoid mixed effects of silencing the expression of potentially antagonistic genes or additive effects of potentially cooperative genes. Consequently, the TIME cell monolayer on each microcarrier has to be monoclonal, having evolved from a single cell. To achieve a one-to-one cell-microcarrier ratio, we used a seeding ratio of one cell per two microcarriers. Starting from a single cell, TIME cells reached confluence in nine days, forming tightly sealed monolayers similar to cells grown from a larger initial number (Fig. 3b).

TIME cells grown on microcarriers form intact tight and adherens junctions
Formation and maintenance of a cobblestone monolayer with integral intercellular junctions is essential for the repurposing of cell-coated microcarriers as individual permeability assays. Evidence for the establishment of such monolayers on the same type of microcarriers used here, including at the ultrastructural level, had been reported 30 years ago[35,42]. To find if junction-specific proteins are present at TIME cell intercellular junctions, we probed the cells for the tight junction single-pass transmembrane protein JAMA, and for the adaptor protein ZO1, which bind each other[43]. Both proteins were abundantly present at the borders of all the cells that were visible in the field of view, appearing as narrow undisrupted bands (Fig. 3c). The adherens junction marker vascular endothelial (VE)-cadherin was similarly abundant and patterned (ibid.). We conclude that the intercellular junctions formed by MAPs do not differ from junctions formed by the same (Fig. 2c) or other[44] ECs on flat surfaces.

Thrombin increases attachment of FN_c to gelatin-coated surfaces
To assess the performance of FN_{cf} as a probe (Appendix SI, Fig. S1a), we tested if the extent of its binding to a gelatin-coated substrate decreases as EC confluence increases. The fluorescence signal of the probe attached to a glass coverslip harboring a confluent TIME cell monolayer was miniscule in comparison to the signal emanating from a sub-confluent monolayer (Appendix SI, Fig. S1b). Thrombin treatment of confluent TIME cell monolayers generated large gaps between adjacent cells (Appendix SI, Fig. S1c). Bare, gelatin-coated microcarriers bound FN_{cf} avidly, whereas microcarriers coated by a sub-confluent EC monolayer bound only a fraction of the latter (Appendix SI, Fig. S1d).

Likelihood of false discovery is reduced by removal of non-confluent microcarriers
Though the majority of TIME cell monolayers reached confluence even when grown from a single cell per microcarrier, it is practically impossible to ensure that the surface of all microcarriers is fully coated by cells. Elimination of microcarriers that have residual exposed surface patches is essential because the FN_{cf} probe would bind to the exposed surface, causing misidentification of the cells as intransigent to the sgRNA they had been transduced with, and the overlooking of potentially relevant genes. To eliminate sub-confluent microcarriers from the population, we sorted MAPs that were incubated with FN_{cf} in the absence of an agonist. MAP extinction profiles were frequently uniform across the imaged face of the microcarrier but peaked at the edges of individual MAPs (Fig. 4a). The peaks corresponded probably to the gelatin and TIME cell layers. The initial gating of the MAP population excluded debris or MAP clusters (Fig. 4b). The residual MAP population was gated again to remove MAPs whose peak fluorescence intensity was at least two-fold higher than the intensity of the highest counts (ibid.), assuming this would eliminate most of the non-confluent MAPs. The removed fraction amounted

154 to 12 percent of the MAP population. Consequently, the peak-height histogram of the residual
155 MAPs was confined to a narrow range (Fig. 4c). Visual inspection of samples from the initial and
156 residual populations confirmed the presence of exceptionally bright MAPs in the former (Fig.
157 4d), but not in the latter (Fig. 4e).
158
sgRNAs repress target gene expression level, protein abundance, and response to
thrombin
To effectively test if the microcarrier-based permeability assay can identify genes that affect EC
response to thrombin, we selected sgRNAs that target *F2R*, the gene that encodes the main
thrombin receptor in ECs[45], proteinase-activated receptor (PAR) 1, as a positive control. We
chose *KDR*, the gene that encodes VEGF receptor 2, the main VEGF receptor in blood vessel
ECs[46], as a negative control. We included sgRNAs that target *GNAQ*, a gene encoding $G\alpha_q$, a
trimeric GTPase. $G\alpha_q$ mobilizes calcium by activating phospholipase C-β, but, unlike $G\alpha_{12/13}$,
there is no consensus in regard to its ability to increase EC monolayer permeability[47,48]. We
selected it, therefore, as an unknown. Before testing the effects of the selected sgRNAs on EC
barrier function, we validated the presumed capacity of each of the five individual sgRNAs
designed to target these genes[8] by measuring how effectively they repressed the transcription
of their target genes and, consequently, reduced the abundance of the proteins these genes
encode. Using quantitative real-time polymerase chain reaction (qRT-PCR), we detected a large
variability among the repressive effects within each set of five sgRNAs, which reached more
than 13-fold in the case of *F2R*-targeting sgRNAs (Fig. 5a). The most effective sgRNA
repressed *F2R* transcription to approximately 6 percent relative to its expression level in TIME
cells treated by non-targeting sgRNAs (ibid.). Based on immunoblotting, the pattern of PAR1
abundance in each cell group expressing a single sgRNA fit closely the qRT-PCR expression
pattern of *F2R* (which encodes PAR1) once the weight of each band was normalized relative to
the weight of the corresponding loading-control β-actin band (Fig. S2a). The single sgRNA-
specific expression levels of *GNAQ* and *KDR* fit well the abundances of $G\alpha_q$ and VEGFR2 (Fig.
5b, 5c, Fig. 2b, 2c), confirming that two to three sgRNAs out of each set of five were capable of
repressing their target gene by at least 60 percent.
183
Efficient repression of *F2R* is expected to diminish the effect of thrombin on endothelial
cell junctions, which typically lose their interconnections and retract, producing large intercellular
gaps (Appendix SI, Fig. S1c). The effect of *GNAQ* repression, on the other hand, remains to be
revealed in the planned permeability assays. To test these predictions, we measured the
impedance of TIME cell populations expressing each of the most repressive sgRNAs of *F2R*,
GNAQ, and *KDR*, grown in multi-well plates before and after the application of thrombin. The
impedance of TIME cells expressing sgRNAs for *GNAQ*, *KDR*, and a validated negative-control
(NC) sgRNA[8] fell by close to one impedance unit (Fig. 5d). The impedance of TIME cells
expressing *F2R* sgRNA, on the other hand, fell only by 0.2 units and returned to the basal level
earlier than the other cell groups.
194
Fluorescence-assisted sorting separates MAPs carrying thrombin-treated and untreated
TIME cells
Execution of a genome-wide screen requires physical separation between the phenotypes that
ensue during the functional assay. In the current study, these are thrombin-responsive and non-
responsive MAPs. FN_{cf} would enable separation by fluorescence-assisted sorting. After
thrombin treatment and before sorting, we fixed the microcarrier-attached TIME cells by EtOH,
an optimal fixative for the preservation of nucleic acids[49], to prevent changes in cell and junction
morphology. Due to potential heterogeneity in cell response to thrombin, the light-intensity
distribution of among the sorted MAPs was continuous over an interval of intensity values,

rather than binary (Fig. 6a). To produce two distinct sorted populations, we defined two gates,
one for the low-responding microcarriers, referred to as 'negative', and another for the high-
responders that we refer to as 'positive' (ibid.). The fraction of untreated negative MAPs fell from
approximately 40 percent to only 10 percent of the thrombin-treated microcarrier population
because of the upward shift in MAP fluorescence intensity (Fig. 6b). Complementarily, the
percentage of positive microcarriers increased from approximately 15 to 40 percent of the
treated microcarrier population. These measurements demonstrate a robust and reproducible
response to thrombin that can be readily resolved into positive and negative populations. A
portion of approximately 45 percent of the total microcarrier population, encompassing the
medium range of response, remained between the upper and lower gates. This sub-population
is least likely to contain microcarriers carrying cells that express the most repressive sgRNAs.
Sidestepping this population is not likely, therefore, to preclude the identification of genes that
have major roles in mediating the disassembly of cell junctions in response to thrombin.
217
MAP sensitivity to *F2R* knockdown is similar to the sensitivity of impedance
measurement
We chose impedance measurement as a benchmark for assay sensitivity because this
technique is more sensitive than that of probe flux through cell monolayers cultured on
permeable substrates[50], the main alternative technique. We tested the sensitivity of impedance
measurement as a detector of sgRNA-mediated knockdown by comparing the thrombin
response of TIME cells transduced by the most effective *F2R*-targeting sgRNA, *F2R*-2, to cells
transduced by a negative-control sgRNA. Whereas the impedance of cells transduced by the
latter sgRNA fell by 2.6 units (denoted as I_{NC} in Fig. 7a), the impedance of TIME cells
transduced by *F2R*-2 fell only by 0.75 unit (I_{F2R}).
228
As a first step in the test of MAP sensitivity to *F2R* knockdown, we repeated the procedure
described above for the removal of particles that were not whole single MAPs and of non-
confluent single MAPs (Fig. 4b). The mean peak fluorescence intensity of thrombin-treated
MAPs carrying cells transduced by the most effective *F2R*-targeting sgRNA (denoted as F_{F2RT} in
Fig. 7b) was higher by approximately 18 percent than that of the same type of untreated MAPs
(F_{F2RU}). On the other hand, MAPs carrying cells transduced by a mixture of three negative
control sgRNAs responded to thrombin by a four-fold increase in mean fluorescence intensity
compared to an untreated group (compare F_{NCT} to F_{NCU} in Fig. 7c). Because impedance
measurement sensitivity is a dimensionless variable given by the ratio between I_{NC} and I_{F2R} (Fig.
7a), we defined MAP sensitivity as the ratio between the response of cells which express a
mixture of three negative control sgRNAs and cells that express *F2R*-2, i.e. the ratio between
$F_{NCT}-F_{NCU}$ (Fig. 7b) and $F_{F2RU}-F_{F2RT}$ (Fig. 7c). The respective sensitivities were 3.5 and 7.6,
indicating that MAP sensitivity was comparable or better than the sensitivity of impedance
measurement for detecting the repressive effect of the *F2R*-2 targeting sgRNA.
243
***F2R* sgRNA suppresses thrombin-induced hyperpermeability**
All the sgRNAs tested above repress, albeit to varying degrees, the expression of their target
genes. Therefore, we gauged the combined effect of these sgRNAs on MAP permeability by
comparing the fraction of MAPs sorted from a mixed population carrying cells that express on
each MAP one of the 15 sgRNAs employed in the study (five each for *F2R*, *GNAQ*, and *KDR*) to
the fraction of MAPs carrying cells that express negative control sgRNAs. The sorting of the first
group (Fig. 7d) showed that sgRNA expression almost abolished the population of untreated
positive MAPs and reversed the distribution of thrombin-treated MAPs shown in Fig. 6b, i.e.
reduced the number of positive MAPs while increasing the number of negative MAPs. The
former was more than halved, and the latter increased by close to seven-fold. These large
effects demonstrate the repressive efficacy of each gene-specific group of five sgRNAs. To

visualize the individual effect of the most repressive *F2R*, *GNAQ* and *KDR*-targeted sgRNAs on
MAP permeability in comparison to three validated NC sgRNAs[8], we imaged the fluorescence of
thrombin-treated MAPs expressing each of these sgRNAs. Cells expressing either *GNAQ*, or
KDR-targeted sgRNAs responded to thrombin similarly to negative-control sgRNA-expressing
MAPs (Fig. 7e). In contrast, the response of MAPs carrying cells that expressed the most
effective *F2R* sgRNA was similar to that of MAPs carrying untreated cells that expressed one of
the negative-control sgRNAs (top row of images in Fig. 7e). These results are consistent with
the other functional effects of the sgRNAs shown above. The persistence of low fluorescence
intensity emitted by MAPs carrying cells that express a *F2R* sgRNA indicates that thrombin
treatment did not cause an artifactual signal resulting from cell detachment from the microcarrier
surface. Furthermore, an assay with MAPs that express a repressive sgRNA library requires
sorting of microcarriers that emit weak light signals. Therefore, artifactually bright microcarriers
would not cause false gene identification.
268
DISCUSSION
Permeability is not amenable to existing functional assays, because it is inherently a collective
cell function. The new assay repurposes a pre-existing device – microcarriers that had been
developed over fifty years ago to increase the surface available for the growth of adherent
cells[51]. Microcarriers had already been used, in fact, to measure EC monolayer permeability,
albeit in bulk format[35]. In contrast, the new assay deploys each cell-coated microcarrier as a
single sample, thus taking advantage of its small diameter to pack as many as 6000
microcarriers per mL of medium. At this sample concentration, 166 mL of growth medium would
contain 10^6 microcarriers. As estimated above, this is a sufficiently large number of samples for
performing a genome-wide screen with this functional assay.
279
Screens based on cell growth can measure a continuum of quantitative phenotypes by
counting the numbers of cells that express sgRNAs of interest[52]. These measurements can be
used for further analysis of the targets identified in the first screen, such as construction of
genetic interaction (GI) maps[53]. Though the phenotypes of the MAPs (i.e., their fluorescence
intensities), also vary continuously within a certain range, the sorting converts that range into an
either 'positive' or 'negative' binary output, precluding the attribution of a continuous range of
quantitative phenotypes to each sample, and of further analysis. This drawback can be
overcome, however, by performing a set of assays where the position of the gate is changed
incrementally so that a larger range of phenotypes is sampled. Replicating the assay is
straightforward because it can be run in as little as 30 min using a volume of less than 200 mL
microcarriers in growth medium.
291
In addition to compactness, the format of the assay facilitates physical separation between
samples that are either responsive or refractory to the stimulus, by using the phenotype as a
sorting input. The separation is crucial for the identification of the genes that produce the
measured phenotype. Counting the number of cells on the microcarriers, and determining if the
cell monolayer reached confluence are straightforward because the cells on the microcarrier
surface can be readily visualized by a dye and by other imaging modalities[35,42]. The visibility
enables growth of a clonal cell population from a single cell per MAP and makes it possible to
verify that the cell monolayer coating the MAPs is confluent, preventing spurious signal as a
result of probe binding to the exposed surface of the microcarrier. The duration of the assay, i.e.
the time necessary for eliciting a measurable phenotype, depends on the nature of the agonist,
but it is not likely to allow sufficient time for the emergence of multiple phenotypes, such as
change in cell density. This narrows the scope of the assay to its intended purpose of
measuring intercellular permeability and prevents interference from other potential effects of the
agonists which are often pleiotropic, such as VEGF.

306
As is standard practice in pioneering proof-of-principle studies, we favored permissive
experimental conditions. Accordingly, we chose an agonist that induces relatively large gaps
that did not strain the detection limits of our probe. Admittedly, other agonists produce subtler
remodeling of intercellular junctions, requiring higher sensitivity for distinguishing between the
signals emitted by monolayers of quiescent and stimulated ECs. As long as the agonist induces
sufficiently large openings between adjacent cells to permit binding of the probe to the
microcarrier surface, our approach should be feasible as long as the binding affinity between the
probe and the surface is sufficiently high. VEGF, an obvious agonist of interest, generates gaps
on the order of several micrometers[54]. Its signaling pathway should be amenable, therefore, to
MAP screening. The MAP configuration limits to some extent the range of testable conditions
because it does not simulate abluminally-acting agonists, including, in some cases, VEGFR2-
downstream effects[55]. Genes that are required for cell proliferation and for adhesion to gelatin
will not be represented in the screened cell population because cells that express sgRNAs
which target these genes will be lost during cell growth on microcarriers. It is unlikely, however,
that such genes would be directly involved in regulating intercellular junction integrity.
322
323 **MATERIALS AND METHODS**
324
325 **Endothelial cell culture**
326 TIME cells (American Type Cell Culture) were grown at 37°C in a humidified 5 percent CO_2
327 incubator in Vascular Cell Basal Medium supplemented with Microvascular Endothelial growth
328 Kit-VEGF (American Type Cell Culture) and 12.5 μg/mL blasticidin (ThermoFisher Scientific).
329
Immunofluorescence
TIME cells grown to confluence on gelatin-coated round 12-mm glass coverslips (GG-12-
Gelatin, Neuvitro) were treated with 2 U/mL thrombin (human α-thrombin, Enzyme Research
Laboratories) for 30 min at 37°C, and 5 percent CO_2, or with PBS as a negative control. After
washing in ice-cold PBS, fixation in 4 percent paraformaldehyde/PBS for 30 min at 23°C, and
permeabilization in 0.1 percent Triton X-100 (Sigma-Aldrich) in PBS for 5 min, cells were
incubated in a blocking solution of 5 percent BSA (Sigma-Aldrich) in PBS for 1 h.
Immunolabeling by primary antibodies diluted according to the manufacturer's instructions in 0.1
percent Triton X-100 and1 percent BSA in PBS overnight at 4 °C was followed by a 3×PBS
wash and incubation with 2.5 μg/mL fluorescently-conjugated secondary antibodies in PBS for
30 min at 23°C. After a 3×PBS wash, coverslips were mounted on glass slides (ProLong Gold
Antifade Mountant, ThermoFisher Scientific). Images of TIME cells grown either as a two-
dimensional culture or on microcarriers were acquired by epi-fluorescence (EVOS FL Auto Cell
Imaging System, ThermoFisher Scientific) or by laser-scanning confocal microscopy (Nikon
A1R+).
345
Immunoblotting
Cells were scraped off plates, extracted in RIPA lysis buffer (25 mM Tris-HCl pH 7.6, 150mM
NaCl, 1 percent NP-40, 1 percent sodium deoxycholate, 0.1 percent SDS; ThermoFisher
Scientific) supplemented with phosphatase (Fisher Scientific) and protease (Sigma-Aldrich)
inhibitor cocktails, and clarified by centrifugation (20,000 g, 10 min). Total protein concentration
was determined by BCA protein assay (ThermoFisher Scientific). Equal amounts of total protein
were separated by 10 percent SDS-PAGE and transferred to a PVDF membrane. Membranes
were blocked by 5 percent casein in TBS with 0.1 percent Tween 20 (Sigma-Aldrich), probed
with the indicated primary antibodies, and incubated with HRP-conjugated secondary antibodies
for 1 h at 23°C. Protein-antibody complexes were visualized by chemiluminescence

(SuperSignal™ West Pico PLUS, ThermoFisher Scientific) and imaged by a charge-coupled
device camera (Chemidoc Imaging System, BioRad). Band densitometry was done with the
Gels submenu of ImageJ 2.0.0.
359
Impedance measurement
Impedance was measured on xCELLigence Real-Time Cell Analyzer (ACEA Biosciences)
according to the manufacturer's instructions. Briefly, 30000 TIME cells were seeded in 200 μL
growth medium per well of an electronic 16-well plate (E-Plate View 16, ACEA Biosciences) for
30 min at 23°C to allow cell sedimentation. The cells were grown to confluence at 37°C, 5
percent CO_2. On the day of the experiment, cells were treated by thrombin or by the same
volume of carrier (PBS), and placed in the analyzer for continuous recording of impedance. The
relative impedance output of the instrument is expressed as $(Z_i-Z_0)/15$ Ω, where Z_i is the
impedance at the i^{th} time point, and Z_0 is the initial impedance before agonist addition.
369
Macromolecular permeability
TIME cells seeded at 1×10^5 per filter insert (0.4 μm pores, 6.4 mm diameter, polyethylene
terephthalate membrane, Transwell®, Corning) were grown for 72 h in 24-well plates
(Transwell®, Corning) filled with 800 μL growth medium per well. The cells in the inserts were
supplemented at the same time by either 2 U/mL thrombin or carrier, and by 1 μg/mL
fluorescein isothiocyanate (FITC)-dextran (2000 kDa, Sigma-Aldrich) was added in the upper
chamber. Ten μL samples were removed after 0, 5, 15, 30, 60, 90 and 120 min from the lower
compartment, diluted in 90 μL of PBS, transferred in a 96 well black plate, clear bottom.
Fluorescence in each well was measured with excitation at 485 nm and readout at 535 nm
(VICTOR2, PerkinElmer). The results are expressed as light intensity counts.
380
TIME cell culture on microcarriers in spinner flasks
Cytodex-3 microcarriers (GE Health Sciences) were hydrated in PBS at 50 mL/g for 3 h at 23°C,
washed twice in a 30 mL/g PBS, and autoclaved (121°C, 20 min). A quantity of 92 mg cytodex-3
microcarriers were readied for cell growth by incubation overnight at 37°C, 5 percent CO_2, and
stirred at 35 rpm (Troemner) in 50 mL growth medium in siliconized (Sigmacote, Sigma-Aldrich)
spinner flasks (Bellco Glass, model 1965-95010) equipped with a glass-ball plunger. TIME cells
grown to 80 percent confluence in T75 flasks were briefly detached with 0.05 percent trypsin,
0.02 percent EDTA (American Type Cell Culture). The cells were then resuspended in growth
medium supplemented with 15 percent FBS and 4.5 g/L glucose. Microcarriers were inoculated
with $6X10^6$ cells in 50 mL same growth medium to achieve a density of 24000 cells/cm². The
microcarrier culture was incubated at 37°C, 5 percent CO_2 and stirred intermittently at 35 rpm for
1 min once every 15 min for a total of 1 hour, then stirred continuously at the same speed.
Medium was replenished once every two days. Cells reached confluence in 3-5 days. In order to
seed a single cell per microcarrier, the above amount of microcarriers was inoculated with
$1.5X10^5$ cells to achieve a density of 600 cells/cm². To monitor cell growth on microcarriers, a
small microcarrier sample was removed every day, treated with 2 μM of Calcein AM
(ThermoFisher Scientific), and visualized by epifluorescence microscopy. Under these
conditions, cells reached confluence in 9 days.
399
Detection of barrier disruption by fluorophore-conjugated fragment of fibronectin
To detect disruption of the confluent TIME cell monolayer on microcarriers, we used a 45 kDa
FN_c (Sigma-Aldrich) purified from human plasma (Sigma-Aldrich) conjugated to Dylight 550 with
a labeling kit (ThermoFisher Scientific). The cells were grown to confluence on gelatin-coated
round 12-mm diameter glass coverslips (GG-12-Gelatin, Neuvitro). On the day of the
permeability assay, the cells were treated with thrombin for 30 min at 2 U/mL and with 1 μg/mL

Dylight 550-conjugated FN_{cf} for 5 min. The cells were then washed with ice cold PBS, fixed in 4
percent paraformaldehyde/PBS for 30 min, and washed again with PBS. The coverslips were
mounted on glass slides with Prolong Gold (Invitrogen), a DAPI-containing medium.
Permeability was characterized by quantifying dye fluorescence (DyLight 550, Ex/Em 562⁄576
nm, ThermoFisher Scientific) in epifluorescence images (EVOS FL, ThermoFisher Scientific).

Permeability in response to thrombin on cytodex-3 microcarriers
413 TIME cells grown to confluence on microcarriers were incubated with 2 U/mL of Thrombin for 25
414 min in spinner flasks spun at 35 rpm at 37°C, 5 percent CO_2. The medium was then
415 supplemented with 1 μg/mL FN_{cf} and stirred for 5 min. MAPs were transferred to a glass-bottom
416 plate for visualization and quantification of the Dylight 550 fluorescence (EVOS FL).

Microcarrier sorting
419 MAPs treated as required were fixed in 50 percent EtOH/PBS and 1 percent FBS for 10 min at
420 23°C, washed twice in PBS for 10 min at 4°C, and transferred to PBS with 1 percent FBS.
421 MAPs were sorted at 561 nm on a Biosorter instrument (Union Biometrica) equipped with a FP-
422 1000 nozzle.

Expression of *dCas9-KRAB* and repressive sgRNAs in TIME cells
Lentiviri were produced and infected into TIME cells as described[56]. The cells were transduced
first by lentivirus expressing dCas9 fused to the KRAB repressor module[9] (pHR-SFFV-dCas9-
BFP-KRAB, Addgene). Lentivirus-expressing cells were sorted (FACSAria II, BD Biosciences)
using blue fluorescent protein (BFP) as marker. Oligonucleotides encoding five optimized
sgRNAs targeting the transcription start sites of *F2R*, *GNAQ*, and *KDR*, and NC sgRNAs were
synthesized based on sequences from the CRISPRi V2 library designed by Horlbeck et al.[8].
Each sgRNA was subcloned into the backbone plasmid of the CRISPRi V2 library (pU6-sgRNA
EF1Alpha-puro-T2A-BFP, Addgene) between the BstXI and BlpI restriction sites. Groups of
dCas9-KRAB-expressing TIME cells were each infected separately by lentiviri expressing a
single sgRNA and selected by puromycin resistance conferred by the sgRNA-expressing
plasmid. sgRNA sequences are provided in Appendix SI, Table S1.

Statistics
438 The significance of the difference between means was determined by two-tailed Student's *t*-test.
439 The null hypothesis was considered untrue if the probability satisfied the condition $P \leq 0.05$.

Non-standard Abbreviations
BFP – blue fluorescent protein
CRISPRi –inhibitory clustered regularly interspaced short palindromic repeats sgRNA library
dCas9 – deactivated CRISPR-associated protein 9
DAPI - 4′,6-diamidino-2-phenylindole
EC – endothelial cell
FITC – fluorescein isothiocyanate
FN_c – collagen-binding fragment of fibronectin
FN_{cf} – fluorophore-conjugated FN_c
KRAB – Krüppel-associated box (KRAB)
MAP – microcarrier assay of permeability
PAR – proteinase-activated receptor
qRT-PCR – quantitative real-time polymerase chain reaction
sgRNA – single guide RNA
TIME – telomerase-immortalized microvascular endothelial
UT – untreated

VE – vascular endothelial
VEGF – vascular endothelial growth factor
VP64 – tetrameric VP16 transactivation domain
460
461 **DECLARATIONS**
462 **Ethics approval and consent to participate**
463 No animals and no humans participated in this study.
464
465 **Consent for publication**
466 All authors consented for the publication of this study.
467
468 **Availability of data and material**
469 All data and materials generated in this study will be made available to the scientific community.
470

471 **Competing Interests**
472 The authors have no competing financial or other interests.
473
474 **Funding**
475 This study was supported by the National Heart Lung and Blood Institute (R01 grant HL119984
476 to AH and R01 grant HL128234 to LE) and by the National Cancer Institute (P30 grant CA56036
477 to PF).
478
Author Contributions
CS designed and performed experiments, analyzed the results, and wrote the manuscript; JY
and XK performed experiments, analyzed the results, and prepared reagents; RK performed
experiments, LE supervised the preparation of reagents and designed experiments; PF
supervised the preparation of reagents and DNA sequencing, and analyzed data; EL designed
experiments, analyzed data, and wrote the manuscript; AH conceived the study, coordinated its
performance, designed experiments, analyzed the results, and wrote the manuscript.
486
Acknowledgments
We thank Dr. Florin Tuluc, Flow Cytometry Core Laboratory of the Children's Hospital of
Philadelphia, for assistance in sorting; Drs. Luke Gilbert and Max Horlbeck, and Ryan Pak, for
assistance in the use of the CRISPR/dCas9 library developed in the lab of Dr. Jonathan
Weissman, University of California, San Francisco. We thank Dr. Jonathan Weissman for
depositing the CRISPRi library and dCas9-KRAB plasmid at Addgene.

REFERENCES

1   Qi, L. S. *et al.* Repurposing CRISPR as an RNA-guided platform for sequence-specific
control of gene expression. *Cell* **152**, 1173-1183, doi:10.1016/j.cell.2013.02.022 (2013).
2   Shi, J. *et al.* Discovery of cancer drug targets by CRISPR-Cas9 screening of protein
domains. *Nat Biotechnol* **33**, 661-667, doi:10.1038/nbt.3235 (2015).
3   Adamson, B. *et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables
Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867-1882 e1821,
doi:10.1016/j.cell.2016.11.048 (2016).
4   Manguso, R. T. *et al.* In vivo CRISPR screening identifies Ptpn2 as a cancer
immunotherapy target. *Nature* **547**, 413-418, doi:10.1038/nature23270 (2017).
5   Konermann, S. *et al.* Genome-scale transcriptional activation by an engineered
CRISPR-Cas9 complex. *Nature* **517**, 583-588, doi:10.1038/nature14136 (2015).
6   Chavez, A. *et al.* Comparison of Cas9 activators in multiple species. *Nat Methods* **13**,
563-567, doi:10.1038/nmeth.3871 (2016).
7   Gilbert, L. A. *et al.* CRISPR-mediated modular RNA-guided regulation of transcription in
eukaryotes. *Cell* **154**, 442-451, doi:10.1016/j.cell.2013.06.044 (2013).
8   Horlbeck, M. A. *et al.* Compact and highly active next-generation libraries for CRISPR-
mediated gene repression and activation. *Elife* **5**, doi:10.7554/eLife.19760 (2016).
9   Gilbert, L. A. *et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression and
Activation. *Cell* **159**, 647-661, doi:10.1016/j.cell.2014.09.029 (2014).
10 Chen, S. *et al.* Genome-wide CRISPR screen in a mouse model of tumor growth and
metastasis. *Cell* **160**, 1246-1260, doi:10.1016/j.cell.2015.02.038 (2015).
11 Gayle, S. *et al.* Identification of apilimod as a first-in-class PIKfyve kinase inhibitor for
treatment of B-cell non-Hodgkin lymphoma. *Blood* **129**, 1768-1778, doi:10.1182/blood-
2016-09-736892 (2017).
12 Kurata, M. *et al.* Using genome-wide CRISPR library screening with library resistant
DCK to find new sources of Ara-C drug resistance in AML. *Sci Rep* **6**, 36199,
doi:10.1038/srep36199 (2016).
13 Tao, L. *et al.* Frizzled proteins are colonic epithelial receptors for C. difficile toxin B.
Nature **538**, 350-355, doi:10.1038/nature19799 (2016).
14 Yamauchi, T. *et al.* Genome-wide CRISPR-Cas9 Screen Identifies Leukemia-Specific
Dependence on a Pre-mRNA Metabolic Pathway Regulated by DCPS. *Cancer Cell*,
doi:10.1016/j.ccell.2018.01.012 (2018).
15 Napier, B. A. *et al.* Complement pathway amplifies caspase-11-dependent cell death
and endotoxin-induced sepsis severity. *J Exp Med* **213**, 2365-2382,
doi:10.1084/jem.20160027 (2016).
16 Parnas, O. *et al.* A Genome-wide CRISPR Screen in Primary Immune Cells to Dissect
Regulatory Networks. *Cell* **162**, 675-686, doi:10.1016/j.cell.2015.06.059 (2015).
17 Shi, J. *et al.* Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell
death. *Nature* **526**, 660-665, doi:10.1038/nature15514 (2015).
18 Zimmermann, S. *et al.* ALPK1- and TIFA-Dependent Innate Immune Response
Triggered by the Helicobacter pylori Type IV Secretion System. *Cell Rep* **20**, 2384-2395,
doi:10.1016/j.celrep.2017.08.039 (2017).
19 Burr, M. L. *et al.* CMTM6 maintains the expression of PD-L1 and regulates anti-tumour
immunity. *Nature* **549**, 101-105, doi:10.1038/nature23643 (2017).
20 Fei, T. *et al.* Genome-wide CRISPR screen identifies HNRNPL as a prostate cancer
dependency regulating RNA splicing. *Proc Natl Acad Sci U S A* **114**, E5207-E5215,
doi:10.1073/pnas.1617467114 (2017).
21 Wang, T. *et al.* Identification and characterization of essential genes in the human
genome. *Science* **350**, 1096-1101, doi:10.1126/science.aac7041 (2015).

22 Xu, C. *et al.* piggyBac mediates efficient in vivo CRISPR library screening for tumorigenesis in mice. *Proc Natl Acad Sci U S A* **114**, 722-727, doi:10.1073/pnas.1615735114 (2017).

23 Liu, Y. *et al.* CRISPR Activation Screens Systematically Identify Factors that Drive Neuronal Fate and Reprogramming. *Cell Stem Cell* **23**, 758-771 e758, doi:10.1016/j.stem.2018.09.003 (2018).

24 Stoletov, K. *et al.* Quantitative in vivo whole genome motility screen reveals novel therapeutic targets to block cancer metastasis. *Nat Commun* **9**, 2343, doi:10.1038/s41467-018-04743-2 (2018).

25 Zhou, P. *et al.* Alpha-kinase 1 is a cytosolic innate immune receptor for bacterial ADP-heptose. *Nature* **561**, 122-126, doi:10.1038/s41586-018-0433-3 (2018).

26 Cong, L. *et al.* Multiplex genome engineering using CRISPR/Cas systems. *Science* **339**, 819-823, doi:10.1126/science.1231143 (2013).

27 Erard, N., Knott, S. R. V. & Hannon, G. J. A CRISPR Resource for Individual, Combinatorial, or Multiplexed Gene Knockout. *Mol Cell* **67**, 348-354 e344, doi:10.1016/j.molcel.2017.06.030 (2017).

28 Aman, J. *et al.* Effective treatment of edema and endothelial barrier dysfunction with imatinib. *Circulation* **126**, 2728-2738, doi:10.1161/CIRCULATIONAHA.112.134304 (2012).

29 Kaneider, N. C. *et al.* 'Role reversal' for the receptor PAR1 in sepsis-induced vascular damage. *Nat Immunol* **8**, 1303-1312, doi:10.1038/ni1525 (2007).

30 Klei, L. R. *et al.* MALT1 Protease Activation Triggers Acute Disruption of Endothelial Barrier Integrity via CYLD Cleavage. *Cell Rep* **17**, 221-232, doi:10.1016/j.celrep.2016.08.080 (2016).

31 Fisher, J. *et al.* Is Heparin-Binding Protein Inhibition a Mechanism of Albumin's Efficacy in Human Septic Shock? *Crit Care Med* **46**, e364-e374, doi:10.1097/CCM.0000000000002996 (2018).

32 Rahbar, E. *et al.* Endothelial glycocalyx shedding and vascular permeability in severely injured trauma patients. *J Transl Med* **13**, 117, doi:10.1186/s12967-015-0481-5 (2015).

33 Martin, Y., Eldardiri, M., Lawrence-Watt, D. J. & Sharpe, J. R. Microcarriers and their potential in tissue regeneration. *Tissue Eng Part B Rev* **17**, 71-80, doi:10.1089/ten.TEB.2010.0559 (2011).

34 Eaton, B. M., Toothill, V. J., Davies, H. A., Pearson, J. D. & Mann, G. E. Permeability of human venous endothelial cell monolayers perfused in microcarrier cultures: effects of flow rate, thrombin, and cytochalasin D. *J Cell Physiol* **149**, 88-99, doi:10.1002/jcp.1041490112 (1991).

35 Killackey, J. J., Johnston, M. G. & Movat, H. Z. Increased permeability of microcarrier-cultured endothelial monolayers in response to histamine and thrombin. A model for the in vitro study of increased vasopermeability. *Am J Pathol* **122**, 50-61 (1986).

36 Park, J. H. *et al.* Hypoxia/aglycemia increases endothelial permeability: role of second messengers and cytoskeleton. *Am J Physiol* **277**, C1066-1074 (1999).

37 Waters, C. M. Flow-induced modulation of the permeability of endothelial cells cultured on microcarrier beads. *J Cell Physiol* **168**, 403-411, doi:10.1002/(SICI)1097-4652(199608)168:2<403::AID-JCP19>3.0.CO;2-3 (1996).

38 Owens, R. J. & Baralle, F. E. Mapping the collagen-binding site of human fibronectin by expression in Escherichia coli. *EMBO J* **5**, 2825-2830 (1986).

39 Engvall, E., Ruoslahti, E. & Miller, E. J. Affinity of fibronectin to collagens of different genetic types and to fibrinogen. *J Exp Med* **147**, 1584-1595 (1978).

40 Laposata, M., Dovnarsky, D. K. & Shin, H. S. Thrombin-induced gap formation in confluent endothelial cell monolayers in vitro. *Blood* **62**, 549-556 (1983).

41   Rabiet, M. J. *et al.* Thrombin-induced increase in endothelial permeability is associated
with changes in cell-to-cell junction organization. *Arterioscler Thromb Vasc Biol* **16**, 488-
496 (1996).

42   Haselton, F. R., Mueller, S. N., Howell, R. E., Levine, E. M. & Fishman, A. P.
Chromatographic demonstration of reversible changes in endothelial permeability. *J Appl*
Physiol (1985) **67**, 2032-2048, doi:10.1152/jappl.1989.67.5.2032 (1989).

43   Ebnet, K., Schulz, C. U., Meyer Zu Brickwedde, M. K., Pendl, G. G. & Vestweber, D.
Junctional adhesion molecule interacts with the PDZ domain-containing proteins AF-6
and ZO-1. *J Biol Chem* **275**, 27979-27988, doi:10.1074/jbc.M002363200 (2000).

44   Ngok, S. P. *et al.* VEGF and Angiopoietin-1 exert opposing effects on cell junctions by
regulating the Rho GEF Syx. *J Cell Biol* **199**, 1103-1115, doi:10.1083/jcb.201207009
(2012).

45   Kataoka, H. *et al.* Protease-activated receptors 1 and 4 mediate thrombin signaling in
endothelial cells. *Blood* **102**, 3224-3231, doi:10.1182/blood-2003-04-1130 (2003).

46   Quinn, T. P., Peters, K. G., De Vries, C., Ferrara, N. & Williams, L. T. Fetal liver kinase 1
is a receptor for vascular endothelial growth factor and is selectively expressed in
vascular endothelium. *Proc Natl Acad Sci U S A* **90**, 7533-7537 (1993).

47   McLaughlin, J. N. *et al.* Functional selectivity of G protein signaling by agonist peptides
and thrombin for the protease-activated receptor-1. *J Biol Chem* **280**, 25048-25059,
doi:10.1074/jbc.M414090200 (2005).

48   Singh, I. *et al.* Galphaq-TRPC6-mediated Ca2+ entry induces RhoA activation and
resultant endothelial cell shape change in response to thrombin. *J Biol Chem* **282**, 7833-
7843, doi:10.1074/jbc.M608288200 (2007).

49   Srinivasan, M., Sedmak, D. & Jewell, S. Effect of fixatives and tissue processing on the
content and integrity of nucleic acids. *Am J Pathol* **161**, 1961-1971, doi:10.1016/S0002-
9440(10)64472-0 (2002).

50   Bischoff, I. *et al.* Pitfalls in assessing microvascular endothelial barrier function:
impedance-based devices versus the classic macromolecular tracer assay. *Sci Rep* **6**,
23671, doi:10.1038/srep23671 (2016).

51   van Wezel, A. L. Growth of cell-strains and primary cells on micro-carriers in
homogeneous culture. *Nature* **216**, 64-65 (1967).

52   Kampmann, M., Bassik, M. C. & Weissman, J. S. Integrated platform for genome-wide
screening and construction of high-density genetic interaction maps in mammalian cells.
Proc Natl Acad Sci U S A **110**, E2317-2326, doi:10.1073/pnas.1307002110 (2013).

53   Bassik, M. C. *et al.* A systematic mammalian genetic interaction map reveals pathways
underlying ricin susceptibility. *Cell* **152**, 909-922, doi:10.1016/j.cell.2013.01.030 (2013).

54   Wisniewska-Kruk, J. *et al.* A novel co-culture model of the blood-retinal barrier based on
primary retinal endothelial cells, pericytes and astrocytes. *Exp Eye Res* **96**, 181-190,
doi:10.1016/j.exer.2011.12.003 (2012).

55   Hudson, N. *et al.* Differential apicobasal VEGF signaling at vascular blood-neural
barriers. *Dev Cell* **30**, 541-552, doi:10.1016/j.devcel.2014.06.027 (2014).

56   Kampmann, M., Bassik, M. C. & Weissman, J. S. Functional genomics platform for
pooled screening and generation of mammalian genetic interaction maps. *Nat Protoc* **9**,
1825-1847, doi:10.1038/nprot.2014.103 (2014).

FIGURE LEGENDS

Figure 1: Microcarriers are repurposed as miniature individual permeability assays. a. Gelatin-coated microcarriers composed of cross-linked dextran carry a confluent EC monolayer (green, to designate calcein-loaded ECs), incubated in medium containing the collagen-binding proteolytic fragment of fibronectin conjugated to a fluorophore (FN_{cf}). Once treated by a junction-disrupting agonist (thrombin in this study), FN_{cf} binds to the exposed gelatin surface between cells whose junctions disassembled in response to the agonist. **b.** Microcarriers carrying untreated ECs bind a minimal amount of FN_{cf}. Microcarriers carrying agonist-treated ECs bind varying amounts of $FN_{cf,}$, depending on the identity of the sgRNA expressed by the clonal cell population on each microcarrier. Microcarriers carrying ECs that express sgRNAs targeting genes that encode proteins required for the induction of the disassembly of cell-cell junctions bind a low amount $FN_{cf,}$ similar to untreated microcarriers. ECs expressing sgRNAs that are unrelated to the signaling pathway of the junction-disrupting agonist respond by disassembling their junctions. FN_{cf} binds to the gelatin surface exposed between the responsive ECs, rendering the microcarriers that carry these cells fluorescent. The fluorescent microcarriers and the dark microcarriers are separated by fluorescence-assisted sorting. The gates of the sorting machine can be set up to capture any group of interest in this population, based on microcarrier fluorescence intensity.

Figure 2: Thrombin increases the paracellular permeability of TIME cell monolayers. a. The permeability of TIME cell confluent monolayers grown on filter inserts to 2000 kDa FITC-dextran increased in response to treatment by 2.0 U thrombin applied at t=0 min for the duration of the experiment, as indicated by the augmentation of the fluorescence signal from the bottom well under. **b.** The impedance of TIME cell monolayers grown on electronic 16-well plates fell sharply after treatment by the indicated dosages of thrombin (applied as above), indicating that the integrity of the monolayer deteriorated. **c.** Epifluorescence images of confluent TIME cells in monolayers grown on gelatin-coated coverslips. Intercellular junctions disassembled upon treatment by 2 U/mL thrombin (Thr), as indicated by the change in VEcad signal. The cells formed thicker stress fibers and detached from each other. Magnified images of the regions marked by white frames are shown on the left and right sides. The double-pointed arrow marks a gap between thrombin-treated cells. Scale bars, 50 μm.

Figure 3: TIME cells form a clonal confluent cobble-stone monolayer on microcarriers. a. Images of calcein AM-stained TIME cells seeded on microcarriers at a density of 1.5×10^4 cells/cm^2 microcarrier surface area and grown for the indicated durations in spinner flasks. The cells reached confluence after 4 days and formed a tightly-sealed monolayer (see cell junctions marked by arrowheads in inset). Scale bars, 200 (top row) and 100 (bottom row) μm. **b.** When initiated from a single TIME cell per microcarrier, a confluent monolayer was formed in 9 days. The inset visualizes intercellular junctions marked by arrowheads. Scale bar, 100 μm. **c.** Microcarriers coated by confluent TIME monolayers immunolabeled with the indicated antibodies. Scale bar, 100 μm.

Figure 4. Binding of FN_{cf} removes non-confluent MAPs. a. Gallery of emission and extinction intensity profiles of individual MAPs. **b**. Left side: density plot of MAP extinction per time-of-flight of untreated MAP population that was incubated with FN_{cf} prior to sorting; right side: histogram of the MAP population shown on the density plot; the gate is placed at approximately 2-fold intensity of the median peak MAP fluorescence. **c.** Left side: density plot of the MAP population gated above; right side: histogram of the MAP population shown on the density plot. **d.** Image of a group of presorted MAPs showing the fluorescence of bound FN_{cf}. **e.** A similar image of a group of sorted MAPs (scale bar, 1 mm).

688
Figure 5. Highly effective sgRNAs reduce substantially the expression and abundance of
the targeted gene and the encoded protein, respectively. a. Quantification of qRT-PCR
measurements of the expression levels of *F2R* in TIME cells transduced by each one of the five
CRISPRi library sgRNAs that target *F2R* (mean±SD, n=3). The immunoblot on the right shows
PAR1 abundance in TIME cells transduced by the same sgRNAs. The first lane on the left was
loaded with lysate of TIME cells that do not express sgRNAs. β-actin was used as a loading
control for all immunoblots. **b** and **c**. The same measurements as above for the sgRNAs that
target *GNAQ* or *KDR*, and the corresponding immunoblots. **d.** Impedance traces of TIME cells
expressing each of the indicated sgRNAs. Thrombin was applied to all the cell groups at the
time point marked by an arrow. The traces were normalized relative to the impedance of
untreated (UT) cells expressing negative control (NC-2) sgRNA.
700
Figure 6. Dark and fluorescent microcarriers are over-abundant in untreated or thrombin-
treated populations, respectively. a. Confluent TIME cells grown on microcarriers were either
untreated or treated by 2 U/mL thrombin for 30 min followed by 5 min incubation with 1 μg/mL
FN_{cf} for 5 min. The microcarriers were sorted according to the indicated gates into positive (Pos)
and negative (Neg) subgroups. **b**. Histogram summarizing the negative and positive untreated
or thrombin-treated populations of MAPs gated as shown in panel a (mean±SD, n=5, **
P<0.001).
708
Figure 7. *F2R* sgRNA suppresses thrombin-induced hyperpermeability. a. Time course of
impedance measurements of thrombin-treated TIME cells normalized relative to the impedance
of untreated (UT) cells expressing negative control (NC-2) sgRNA. **b.** Gating of peak-height
fluorescence density plot of MAPs carrying cells transduced by *F2R*-targeting sgRNA; righthand
panel: display of the counts per peak height and gate of the same MAP population shown on the
density plot. **c.** Counts per peak-height fluorescence of sorted MAPs that carry either untreated
or thrombin-treated cells transduced by *F2R*-targeting sgRNA. **d.** Effects of thrombin treatment
on the fluorescence intensity of MAPs carrying TIME cell transduced by the indicated sgRNAs.
e. Triplicate fluorescence images of MAPs carrying cells transduced by the indicated sgRNAs
under the indicated conditions. Scale bar, 100 μm.
719

a

Microcarrier Scale

Basal conditions

Treatment with junction-disrupting agonist

b

Sample Scale

Fluorescence Sorting

Dark MCs

Fluorescent MCs

Microcarrier

Microcarrier matrix

Fluorophore

Gelatin coat   EC

Collagen-binding Fn domain

SUPPLEMENTAL MATERIAL

TABLE LEGEND

Table S1. Sequences of the sgRNAs from the CRISPRi V2 library used in the study, the sgRNA-specific reverse primers used in qRT-PCR experiments, and the forward primer based on the sequence of the backbone plasmid of the sgRNA library, pU6-sgRNA EF1Alpha-puro-T2A-BFP.

FIGURE LEGENDS

Figure S1: FN$_{cf}$ identifies gaps between thrombin-treated confluent TIME cells. a. Scheme of human fibronectin showing the location of the collagen-binding domain. Numbers indicate amino acid positions. **b**. TIME cells grown on gelatin-coated glass coverslip until reaching the indicated confluences were incubated with 1 μg/mL FN$_{cf}$ (red) for 5 min. Nuclei were stained by DAPI (blue). Scale bar, 100 μm. **c.** Confluent TIME cells grown as described above were either untreated (UT) or treated by 2 U/mL thrombin (Thr) for 30 min, followed by incubation with 1 μg/mL FN$_{cf}$ for 5 m. **d.** Microcarriers devoid of cells (top) or carrying TIME cells at the indicated confluence (bottom) were incubated with 1 μg/mL FN$_{cf}$ for 30 minutes. Scale bar, 200 μm.

Figure S2: The pattern of sgRNA-induced decrease in protein abundance is similar to the pattern of the decrease in gene expression. a. Densitometric quantification of PAR1 abundance in cells expressing each of the 5 F2R-targeting sgRNAs shown in Fig. 5a, after normalization by the weights of the corresponding β-actin loading-control bands. **b** and **c.** Densitometric quantification similar to PAR1 of Gα_q and VEGFR2 abundances, respectively.

Figure S3. Full-length immunoblots of the bands shown in Fig. 5. The panel to which each immunoblot corresponds is indicated.

Table S1

KDR	Protospacer	5' forward	3' reverse complement		
	GCTGCCAGACGGACTTTCTG	TTGGCTGCCAGACGGACTTTCTGGTTTAAGAGC	TTAGCTCTTAAACCAGAAAGTCCGTCTGGCAGCCAACAAG		
	GCGCAAGTGATGCCCGGCGC	TTGGCGCAAGTGATGCCCGGCGCGTTTAAGAGC	TTAGCTCTTAAACGCGCCGGGCATCACTTGCGCCAACAAG		
	GTCTAGAGAAGGAGGCGCGG	TTGGTCTAGAGAAGGAGGCGCGGGTTTAAGAGC	TTAGCTCTTAAACCCGCGCCTCCTTCTCTAGACCAACAAG		
	GAGGCAGACAGGTCGGGTGA	TTGGAGGCAGACAGGTCGGGTGAGTTTAAGAGC	TTAGCTCTTAAACTCACCCGACCTGTCTGCCTCCAACAAG		
	GGCGGCGCGCAAGTGATGCC	TTGGGCGGCGCGCAAGTGATGCCGTTTAAGAGC	TTAGCTCTTAAACGGCATCACTTGCGCGCCGCCCAACAAG		
SRC					
	Gggcagacggacgcacggga	TTGGggcagacggacgcacgggaGTTTAAGAGC	TTAGCTCTTAAACTCCCGTGCGTCCGTCTGCCCCAACAAG		
	Ggcgggacgtccagcgggac	TTGGgcgggacgtccagcgggacGTTTAAGAGC	TTAGCTCTTAAACGTCCCGCTGGACGTCCCGCCCAACAAG		
	Gcggaccgcgggacgtccag	TTGGcggaccgcgggacgtccagGTTTAAGAGC	TTAGCTCTTAAACCTGGACGTCCCGCGGTCCGCCAACAAG		
	Gcaccggcagacggacgcac	TTGGcaccggcagacggacgcacGTTTAAGAGC	TTAGCTCTTAAACGTGCGTCCGTCTGCCGGTGCCAACAAG		
	Gggcgggcgggcgggctcac	TTGGggcgggcgggcgggctcacGTTTAAGAGC	TTAGCTCTTAAACGTGAGCCCGCCCGCCCGCCCCAACAAG		
PAR1					
	GGTGCCCACGGGTAAGATCA	TTGGGTGCCCACGGGTAAGATCAGTTTAAGAGC	TTAGCTCTTAAACTGATCTTACCCGTGGGCACCCAACAAG		
	GGGACCCTGATCTTACCCGT	TTGGGGACCCTGATCTTACCCGTGTTTAAGAGC	TTAGCTCTTAAACACGGGTAAGATCAGGGTCCCCAACAAG		
	GTCAGGGTCCAAGCGACCCT	TTGGTCAGGGTCCAAGCGACCCTGTTTAAGAGC	TTAGCTCTTAAACAGGGTCGCTTGGACCCTGACCAACAAG		
	GTGGCTTTCGGAGGAAACTG	TTGGTGGCTTTCGGAGGAAACTGGTTTAAGAGC	TTAGCTCTTAAACCAGTTTCCTCCGAAAGCCACCAACAAG		
	GTCGGCGAGCGCTGTGTCTG	TTGGTCGGCGAGCGCTGTGTCTGGTTTAAGAGC	TTAGCTCTTAAACCAGACACAGCGCTCGCCGACCAACAAG		
GNAQ					
	GACTGGAGCACAGATCCggg	TTGGACTGGAGCACAGATCCgggGTTTAAGAGC	TTAGCTCTTAAACCCCGGATCTGTGCTCCAGTCCAACAAG		
	GGCTCCGCGCCTGGCAATCG	TTGGGCTCCGCGCCTGGCAATCGGTTTAAGAGC	TTAGCTCTTAAACCGATTGCCAGGCGCGGAGCCCAACAAG		
	GGAGCACAGATCCgggaggg	TTGGGAGCACAGATCCgggagggGTTTAAGAGC	TTAGCTCTTAAACCCCTCCCGGATCTGTGCTCCCAACAAG		
	GCTCCTTCCCCGGGAACAGG	TTGGCTCCTTCCCCGGGAACAGGGTTTAAGAGC	TTAGCTCTTAAACCCTGTTCCCGGGGAAGGAGCCAACAAG		
	GCGCCAGGCGCACGGCGTAG	TTGGCGCCAGGCGCACGGCGTAGGTTTAAGAGC	TTAGCTCTTAAACCTACGCCGTGCGCCTGGCGCCAACAAG		
negative control					
	GCTGCATGGGGCGCGAATCA	TTGGCTGCATGGGGCGCGAATCAGTTTAAGAGC	TTAGCTCTTAAACTGATTCGCGCCCCATGCAGCCAACAAG		
	GTGCACCCGGCTAGGACCGG	TTGGTGCACCCGGCTAGGACCGGGTTTAAGAGC	TTAGCTCTTAAACCCGGTCCTAGCCGGGTGCACCAACAAG		
	GTGTGCAACCTCCGCCGTTG	TTGGTGTGCAACCTCCGCCGTTGGTTTAAGAGC	TTAGCTCTTAAACCAACGGCGGAGGTTGCACACCAACAAG		
	GCCCGAGCAGTGGCTCGCTA	TTGGCCCGAGCAGTGGCTCGCTAGTTTAAGAGC	TTAGCTCTTAAACTAGCGAGCCACTGCTCGGGCCAACAAG		
	GGTCAGGTAGAGGGATTGAG	TTGGGTCAGGTAGAGGGATTGAGGTTTAAGAGC	TTAGCTCTTAAACCTCAATCCCTCTACCTGACCCAACAAG		
Forward primer	gcacaaaaggaaactcaccct				

a Fibronectin

306 602

1 Collagen Binding Domain 2386

b

~40% conf

~100% conf

c

FN$_{cf}$ UT DAPI

Thrombin

d

No Cells

~70% conf

Figure S1

Figure S2

Fig 5a

β-actin loading controls

PAR1

Fig 5b

G$_{\alpha q}$

Fig 5c

VEGFR2

Figure S3