Genome-wide co-localization of RNA-DNA interactions and fusion RNA pairs

Zhangming Yan[a,1], **Norman Huang**[a,1], **Weixin Wu**[a,1], **Weizhong Chen**[a], **Yiqun Jiang**[a], **Jingyao Chen**[a], **Xuerui Huang**[b], **Xingzhao Wen**[a], **Jie Xu**[c], **Qiushi Jin**[c], **Kang Zhang**[d], **Zhen Chen**[e], **Shu Chien**[a,f,2], and **Sheng Zhong**[a,2]

[a]Department of Bioengineering and Institute of Engineering in Medicine, University of California San Diego, 9500 Gilman Dr., La Jolla, CA 92093, USA; [b]Division of Biological Sciences, University of California San Diego, 9500 Gilman Dr., La Jolla, CA 92093, USA; [c]Department of Biochemistry and Molecular Biology, College of Medicine, The Pennsylvania State University, Hershey, PA, USA; [d]Department of Ophthalmology, University of California San Diego, 9500 Gilman Dr., La Jolla, CA 92093, USA; [e]Department of Diabetes Complications and Metabolism, Beckman Research Institute, City of Hope, 1500 Duarte Rd., Duarte, CA 91010, USA; [f]Department of Medicine, University of California San Diego, 9500 Gilman Dr., La Jolla, CA 92093, USA

This manuscript was compiled on November 18, 2018

Fusion transcripts are used as biomarkers in companion diagnoses. Although more than 15,000 fusion RNAs have been identified from diverse cancer types, few common feature has been reported. Here, we compared 16,410 fusion transcripts detected in cancer (from a published cohort of 9,966 tumor samples of 33 cancer types) with genome-wide RNA-DNA interactions mapped in two normal, non-cancerous cell types (using iMARGI, an enhanced version of the MARGI [Mapping RNA-Genome Interactions assay]). Among the top 10 most significant RNA-DNA interactions in normal cells, 5 co-localized with the gene pairs that formed fusion RNAs in cancer. Furthermore, throughout the genome, the frequency of a gene pair to exhibit RNA-DNA interactions is positively correlated with the probability of this gene pair to present documented fusion transcripts in cancer. To test whether RNA-DNA interactions in normal cells are predictive of fusion RNAs, we analyzed in a validation cohort of 96 lung cancer samples using RNA-seq. We found 34 out of 42 fusion transcripts in the validation cohort exhibited RNA-DNA interactions in normal cells. Finally, combining RNA-seq, single-molecule RNA FISH, and DNA FISH, we detected a cancer sample with Eml4-Alk fusion RNA without forming the Eml4-Alk fusion gene. Collectively, these data suggest a novel RNA-poise model, where spatial proximity of RNA and DNA could poise for creation of fusion transcripts.

fusion transcript | RNA-DNA interaction | RNA-poise model

Fusion transcripts are associated with diverse cancer types and have been proposed as diagnostic biomarkers (1–3). Companion tests and targeted therapies have been developed to identify and treat fusion-gene defined cancer subtypes (2, 4). Recent efforts of detection of fusion transcripts primarily relied on analyses of RNA sequencing (RNA-seq) data (1, 3, 5–8). A recent study analyzed 9,966 RNA-seq datasets across 33 cancer types from The Cancer Genome Atlas (TCGA) and identified more than 15,000 fusion transcripts (4).

Despite the large number of gene pairs in identified fusion transcripts, it remains formidable to predict what unreported pair of genes may form a new fusion transcript. Recent analyses could not identify any distinct feature of fusion RNA forming gene pairs (9). Here, we report a characteristic pattern of the genomic locations of the gene pairs involved in fusion transcripts.

Chromatin-associated RNAs (caRNAs) provide an additional layer of epigenomic information in parallel to DNA and histone modifications (10). The recently developed MARGI (Mapping RNA-Genome Interactions) technology enabled identification of diverse caRNAs and the respective genomic interacting locations of each caRNA (6). In this work, we improved the MARGI experimental pipeline and mapped RNA-DNA interactions in human embryonic kidney (HEK) and human foreskin fibroblast (HFF) cells. The detected RNA-DNA interactions often appeared on the gene pairs involved in cancer-derived fusion transcripts. The wide-spread RNA-DNA interactions on the gene pairs involved in fusion transcripts suggest a model, where the RNA of gene 1 by interacting with the genomic sequence of gene 2 is poised for being spliced into gene 2's nascent transcript and thus creating a fusion transcript. Consistent with this model, we identified an RNA fusion in a new cancer sample that does not involve the creation of a fusion gene.

Results

Characteristics of genome-wide RNA-DNA interaction maps. To systematically characterize caRNAs and their genomic interaction locations, we developed iMARGI, an enhanced version of the MARGI assay (10). The main difference between iMARGI and MARGI is that iMARGI carries out the ligation steps *in situ* (Fig. 1A), whereas MARGI performs these ligation steps on streptavidin beads. We applied iMARGI to HEK and HFF cells to yield 361.2 and 355.2 million 2×100 bp paired-end sequencing read pairs, respectively. These resulted in 36.3 (HEK) and 17.8 (HFF) million valid RNA-DNA interaction read

Author contributions:Conceptualization, Z.Y., N.H., and S.Z.; Methodology, Z.Y., N.H., W.W., and W.C.; Investigation, Z.Y., N.H., W.W., W.C., Y.J., J.C., X.H., Z.C., K.Z., S.C., and S.Z.; Writing – Original Draft, Z.Y., N.H. W.W., and S.Z.; Writing – Review & Editing, Z.Y., Z.C., S.C., and S.Z.; Funding Acquisition, S.C. and S.Z.; Resources, K.Z. and S.Z.; Supervision, S.C. and S.Z.

[1]Z.Y., N.H. and W.W. contributed equally to this work.

[2]To whom correspondence should be addressed. E-mail: shuchien@ucsd.edu or szhong@ucsd.edu

Fig. 1. Overview of iMARGI method and data. (A) Schematic view of iMARGI experimental procedure. (B) An example of inter-chromosomal read pairs (horizonal lines), where the RNA ends (red bars) were mapped to the Malat1 gene on chromosomal 11 and the DNA ends (blue bars) were mapped to chromosome 14 near the Ktn1 gene (blue bars). (C) Proportions of proximal, distal, and inter-chromosomal read pairs in collection of valid RNA-DNA interaction read pairs. M: million read pairs. (D) Ratios of inter- and intra-chromosomal read pairs in HEK and HFF cells after removal of proximal read pairs.(E) Heatmap of an RNA-DNA interaction matrix in HEK cells. The numbers of iMARGI read pairs are plotted with respect to the mapped positions of the RNA-end (row) and the DNA-end (column) from small (blue) to large (red) scale, normalized in each row.

Fig. 2. Summary of iMARGI data. (A) The number of genomic bin pairs with 10 or more iMARGI read pairs (y axis) is plotted against the genomic distance between the pair of genomic bins (x axis), in HEK (red dot) and HFF cells (blue dot). For comparison, the number of Futra pairs (y axis) are plotted against the genomic distance (x axis) between the genes involved in the fusion (purple dot). (B) Histogram of the number of iMARGI read pairs of every gene pair (x axis) across all the intra-chromosomal gene pairs with genomic distance > 200 kb. Arrows: the gene pairs with the largest and the second largest numbers of iMARGI read pairs.

pairs, in which approximately 35%, 10%, and 55% were proximal, distal, and inter-chromosomal interactions, respectively (Fig. 1C and *SI Appendix*, Fig. S1A). The proximal and distal interactions were defined as the intra-chromosomal interactions where the RNA-end and DNA-end were mapped to within and beyond 200 kb, respectively. Following Sridhar et al. (10), we removed proximal read pairs from further analysis because proximal interactions likely represent interactions between nascent transcripts and their neighboring genomic sequences. Hereafter, we refer the union of distal and inter-chromosomal interactions as *remote interactions*. The rest of this paper only deal with remote interactions.

Among the remote RNA-DNA interactions, both cell types exhibited approximately 1:5 ratio of intra- and inter-chromosomal interactions (Fig. 1D). The two-dimensional map of RNA-DNA interactions exhibited more interactions near the diagonal line (Fig. 1E and *SI Appendix*, Fig. S1D). Within each chromosome, the number of iMARGI read pairs negatively correlated with their genomic distances (red and blue dots, Fig. 2A).

The most significant RNA-DNA interactions co-localized with the gene pairs forming fusion transcripts in cancer. We set off to identify the most significant distal RNA-DNA interactions from the iMARGI data. Excluding extremely abundant non-coding RNAs, such as XIST, the top gene pair with the largest amount of inter-chromosomal and distal iMARGI read pairs in HEK cells was FHIT-PTPRG (Fig. 2B and *SI Appendix*, Fig. S2A). Investigating this gene pair, we found FHIT-PTPRG fusion transcripts were reported from kidney, liver, head and neck, lung, and prostate cancers (7). The second largest amount of inter-chromosomal and distal iMARGI read pairs was GPC5-GPC6 (Fig. 2B, *SI Appendix*, Fig. S4 and Fig. S2B). Fusion transcripts from this 2[nd] ranked gene pair were reported from liver and prostate cancers (7). Notably, 5 of the top 10 gene pairs were reported as fusion transcripts in cancers (1, 7). These findings incited us to systematically analyze the relationship between RNA-DNA interactions and fusion transcripts.

Non-uniform distribution of the RNA pairs contributing to fusion transcripts. We asked whether there is any global characteristic of genome-wide distribution of the RNA pairs that contributed to fusion transcripts. To this end, we subjected the previously reported 16,410 fusion transcripts that were derived from 9,966 samples across 33 cancer types from the The Cancer Genome Atlas (TCGA) project to further analysis (4). On average there were two fusion transcripts per sample (Fig. 3A). The 16,410 fusion transcripts corresponded to 15,144 unique RNA pairs. Hereafter we call these gene pairs "Futra pairs" (fusion transcript contributing RNA pairs). More than 95% (14,482 out of 15,144) of Futra pairs occurred only in 1 sample out of the 9,966 samples analyzed (Fig. 3B). These data confirmed the scarcity of recurrent gene pairs in fusion transcripts (11, 12).

We visualized the frequency of Futra pairs in a two-dimensional heatmap, which we call "fusion map" (Fig. 3C and *SI Appendix*, Fig. S3A). The two-dimensional distribution was not uniform, with more intra-chromosomal than inter-chromosomal gene pairs (odds ratio = 27.91, p-value $< 2.2 \times 10^{-16}$, Chi-squared test). A total of 8,891 and 6,253 Futra pairs were intra- and inter-chromosomal, respectively. Chromosomes 1, 12, and 17 harbored the largest amounts of intra-chromosomal gene pairs (*SI Appendix*, Fig. S3B). Chromosomes 1, 12, 11, 17, 19 contribute to the largest amounts of inter-chromosomal gene pairs (*SI Appendix*, Fig. S3C). Higher density of gene pairs appeared on the diagonal line of the fusion map, suggesting enrichment of gene pairs within chromosomes or large chromosomal domains (Fig. 3C and Fig. 4A). We quantified the relative distances between the Futra pairs. The number of the intra-chromosomal Futra pairs negatively correlated with their chromosomal distance (purple dots, Fig. 2A). Taken together, Futra pairs exhibited non-uniform distribution in the genome, characterized

Fig. 3. Overview of the 15,144 Futra pairs derived from 33 cancer types. (A) Distribution of the number of detected Futra pairs in each cancer sample across the 9,966 cancer samples. (B) Pie chart of the numbers of Futra pairs that appeared in only one cancer sample (non-recurring Futra pairs, red) and in multiple cancer samples (recurring Futra pairs, green). (C) The genomic distribution of Futra pairs. Genomic coordinates of all chromosomes are ordered on rows and columns. The numbers of Futra pairs of corresponding genomic positions are shown in a blue (small) to red (large) color scheme. Bin size: 10 Mb.

by enrichment of intra-chromosomal pairs and preference to smaller genomic distances.

Differences between the genomic locations of Futra pairs and genome interactions. We asked to what extent Futra pairs may
correlate with genome interactions. Forty-one percent (6,253 out of 15,144) of Futra pairs were inter-chromosomal, whereas less
than 15% chromosomal conformation capture-derived read pairs were inter-chromosomal (13, 14). The intra-chromosomal
Futra pairs exhibited enrichment in large chromosomal domains (Fig. 3C, Fig. 4A and *SI Appendix*, Fig. S4A). These enriched
chromosomal domains ranged from approximately 1/10 to 1/3 of a chromosome in lengths, approximately 10 - 20 times of the
typical sizes of topologically associated domains (TADs) (15). Taken together, Futra pairs exhibited different global distribution
characteristics from that of genome interactions.

Genome-wide co-localization of Futra pairs and RNA-DNA interactions. We asked to what extent Futra pairs may coincide with
genome-wide RNA-DNA interactions. We carried out a visualized comparison the two-dimensional distribution of Futra pairs
with that of RNA-DNA interactions (Fig. 3C, Fig. 1E, and *SI Appendix*, Fig. S1D) and observed pronounced similarities (Fig.
4 A-C and *SI Appendix*, Fig. S4 A-C). For example, a set of 34 Futra pairs were enriched in a ~7 Mb region on chromosome 9
(chr9: 32Mb – 39Mb, Fig. 4 A and D). This Futra pair enriched region co-localized with a chromosomal region enriched of
RNA-DNA interactions (Fig. 4 B-D). Such co-localizations were observed in multi-scale analyses using different resolutions,
including 10 Mb (*SI Appendix*, Fig. S4 A-C), 1Mb (Fig. 4 A-C), 100 kb (Fig. 4D) resolutions, as well as at the resolution
of individual fusion pairs and read pairs (*SI Appendix*, Fig. S5). Four fusion transcripts, KMT2C-Auts2, KMT2C-Caln1,
KMT2C-Clip2, and KMT2C-Gtf2ird1 were formed between the KMT2C mRNA near the 152 Mb location of chromosome
7 (chr7:152,000,000) and four mRNAs that were approximately 80 Mb away (chr7:66,000,000 – 78,000,000) (Futra pairs, *SI
Appendix*, Fig. S5). Correspondingly, a total of 73 RNA-DNA iMARGI read pairs were mapped to KMT2C and the four fusion
partners in HEK cells (MARGI, *SI Appendix*, Fig. S5).
We quantified the overlaps between Futra pairs and iMARGI identified RNA-DNA interactions. Among the 6,253 inter-

Fig. 4. Comparison of Futra pairs and iMARGI read pairs. (A) Distribution of intra-chromosomal Futra pairs on chromosome 9. Genomic coordinates are plotted from top to bottom (rows) and from left to right (columns). (B-C) RNA-DNA interaction matrices in HEK (B) and HFF cells (C). The numbers of iMARGI read pairs are plotted with respect to the mapped positions of the RNA-end (rows) and the DNA-end (columns) on chromosomal 9 from small (blue) to large (red) scale. (D) Detailed view of a 30 Mb region (yellow boxes in A-C). This genomic region and the contained genes are plotted twice in the top and the bottom lanes. In the track of Futra pairs, each blue line linking a gene from the top lane to a gene at the bottom indicates a Futra pair. In the track of iMARGI read pair, each green line indicates a iMARGI read pair with the RNA-end mapped to the genomic position in the top lane and the DNA-end mapped to the bottom lane.

chromosomal Futra pairs, 3,014 (48.2%) overlapped with RNA-DNA interactions in either HEK or HFF cells (odds ratio = 14.1, p-value $< 2.2 \times 10^{-16}$, Chi-squared test). Among the 8,891 intra-chromosomal Futra pairs, 7,427 (83.5%) overlapped with RNA-DNA interactions in either HEK or HFF cells (odds ratio = 35.44, p value $< 2.2 \times 10^{-16}$, Chi-squared test). These data pointed to a common feature of cancer-derived Futra pairs, which is their co-localization with RNA-DNA interactions in normal cells.

Cancer-derived Futra pairs that co-localize with RNA-DNA interaction in normal cells do not form fusion transcripts in normal cells. A model that may explain the co-localization of RNA-DNA interactions and Futra pairs is that RNA-DNA interactions in the normal cells poise for creation of fusion transcripts. Recognizing that this model cannot be tested by perturbation due to the very small chance for a fusion transcript to occur in a cancer sample, we carried out two other tests. First, we tested whether the cancer-derived Futra pairs were detectable in normal cells. We re-analyzed merged RNA-seq datasets of more than 75 million 2×100 bp paired-end read pairs from HEK293T cells (16). We ran STAR-Fusion (17) on these datasets which reported a total of 8 Futra pairs. None of the previously derived 15,144 Futra pairs from TCGA RNA-seq data were detected in HEK293T cells. In addition, we specifically tested for Eml4-Alk fusion transcripts, which were reported in non-small cell

lung carcinoma (NSCLC) (18) and there were RNA-DNA interactions between Eml4 RNA and Alk genomic locus in HEK
and HFF cells (Fig. 6A). Neither PCR or quantitative PCR analysis detected Eml4-Alk fusion transcripts in HEK293T cells
(*SI Appendix*, Fig. S6), whereas in both assays detected fusion transcripts in a NSCLC cell line (H2228) (*SI Appendix*, Fig.
S6). Taken together, these data suggest that although cancer-derived Futra pairs co-localized with RNA-DNA interactions in
normal cells, the fusion transcripts found in cancer are not present in the normal cells.

Fig. 5. Fusion transcripts detected from the new lung cancer samples. (A) The number of RNA-seq read pairs (light blue bar) and the uniquely mapped read pairs (dark blue bar) of each sample (column). (B) Number of detected fusion transcripts (y axis) in each sample (columns). (C). Numbers of intra- and inter-chromosomal Futra pairs detected from these 69 cancer samples. (D-E) Intersections of these intra- (D) and inter-chromosomal (E) Futra pairs to RNA-DNA interactions in HEK (green) and HFF (pink). ∗: The 15 intra-chromosomal Futra pairs that overlapped with RNA-DNA interactions in both HEK and HFF (yellow-green) are ALK:EML4, RP11-557H15.4:SGK1, FRS2:NUP107, ACTN4:ERCC2, LRCH1:RP11-29G8.3, CUX1:TRRAP, CEP70:GSK3B, NIPBL:WDR70, KCTD1:SS18, CHST11:NTN4, RP1-148H17.1:TOP1, LMO7:LRCH1, NAV3:RP1-34H18.1, LPP:PPM1L, and MTOR-AS1:RERE. #: The 10 inter-chromosomal Futra pairs that overlapped with RNA-DNA interactions in both HEK and HFF (yellow-green) are LIN52:PI4KA, LPP:OSBPL6, FCGBP:MT-RNR2, KMT2B:MALAT1, ETV6:TTC3, KTN1:MALAT1, FLNA:MALAT1, COL1A2:MALAT1, COL1A1:MALAT1, and MALAT1:TNFRSF10B.

RNA-DNA interactions in normal cells are predictive of fusion transcripts in new cancer samples. Next, we tested whether the
RNA-DNA interactions in normal cells are predictive of fusion transcript formation in cancer. To this end, we analyzed a
validation cohort comprising 96 new lung cancer samples from patients who were not part of TCGA cohorts. We also analyzed
a NSCLC cell line (H2228). We extracted RNA and carried out targeted RNA sequencing (RNA-seq) with Illumina's TruSight
RNA Pan-Cancer Panel. Of these 96 samples, 27 samples did not yield enough RNA for sequencing, whereas the other 69
samples produced a sequencing library and yielded on average 3.9 million uniquely aligned read pairs per sample (Fig. 5A).
The TruSight data analysis package reported a total of 42 fusion transcripts from these 69 samples (Fig. 5 B and C). These 42
fusion transcripts included Eml4-Alk and Frs2-Nup107 fusion, which were also reported from the 9,966 TCGA cancer samples,
as well as 36 new fusion transcripts that were not documented before. The small amount of reoccurring Futra pairs between
these additional cancer samples and TCGA samples is expected from the small fraction of recurring Futra pairs across the
TCGA samples (Fig. 3B).
    Among these 42 Futra pairs detected from the validation cohort, 37 (88.1%) co-localized with RNA-DNA interactions in the
assayed normal cells, supporting the idea that RNA-DNA interactions in the already assayed normal cells are predictive of

Fig. 6. RNA fusion and DNA break. (A) Middle tracks: RNA-seq read pairs (purple bars) aligned to Alk (left) and Eml4 (right). Each pair of paired-end reads is linked by a horizontal line. Lower tracks: iMARGI read pairs aligned to the two genes. Red bars: RNA-end. Blue bars: DNA-end. Thin grey lines: pairing information of iMARGI read pairs. (B) Positive (red) and negative (black) outcomes of detection of Eml4-Alk fusion transcripts based on RNA-seq and FuseFISH, and DNA FISH (ALK Break Apart FISH, a DNA re-arrangement test) in each cancer sample (column). White box: no measurement. ∗: Detected a partial deletion of Alk gene without re-arrangement. (C) FuseFISH images of the #37 cancer sample from the Eml4 and Alk channels. Arrows: co-localized FISH signals indicating fusion RNA. Scale Bar: 2 μm. (D) A representative image of ALK Break Apart FISH, with co-localized red and yellow signals that indicate integral Alk gene without re-arrangement.

Futra pairs in cancer (odds ratio = 106.51, p-value $< 2.2\times10^{-16}$, Chi-squared test). We asked if only intra-chromosomal Futra pairs co-localized with RNA-DNA interactions. Nineteenout of the 36 (45%) detected Futra pairs were inter-chromosomal (Fig. 5C), comparable to the proportion (41%) of inter-chromosomal Futra pairs detected from TCGA samples. Eighty-three percent (19 out of 23) intra- and 95% (18 out of 19) inter-chromosomal Futra pairs overlapped with RNA-DNA interactions (Fig. 5 D and E), suggesting the co-localization of RNA-DNA interactions and Futra pairs were not restricted to intra-chromosomal interactions. Taken together, the co-localization of Futra pairs and RNA-DNA interactions, the lack of cancer-derived fusion transcripts in normal cells, and the predictability to additional Futra pairs in new cancer samples support the model where RNA-DNA interactions in normal cells poise for creation of fusion transcripts in cancers. Hereafter, we refer to this model as the *RNA-poise model*. We call the gene pairs with RNA-DNA interactions in normal cells as *fusion susceptible pairs*.

RNA-DNA interaction between Eml4 and Alk correlates with an RNA fusion without fusion gene in tumor. We tested whether genome re-arrangement is a prerequisite step for creation of fusion transcripts from fusion susceptible pairs. We chose Eml4-Alk fusion transcripts for this test because Eml4-Alk is a fusion susceptible pair (Fig. 6A), Eml4-Alk fusion transcripts are detected in one of our new tumor samples (Sample #44) (Fig. 6B) and there is an FDA approved diagnosis kit (Vysis ALK Break Apart FISH) based on DNA FISH detection of the Eml4-Alk fusion gene. We subjected the remaining tissue from Sample #44 for DNA FISH analysis. None of our 8 attempts produced any DNA FISH signal in the remaining tissue from either control or Alk probes. We therefore could not tell whether there was genome re-arrangement in the only sample with detectable Eml4-Alk fusion transcripts.

In order to identify other cancer samples that express Eml4-Alk fusion transcripts, we re-analyzed our collection of 96 lung

cancer samples with FuseFISH, a single-molecule fluorescence in situ hybridization (sm-FISH) based method for detection of fusion transcripts (19, 20). We carried out quantum dot-labelled sm-FISH (21) by labelling Eml4 and Alk transcripts with quantum dots at 705 nm and 605 nm, respectively (*SI Appendix*, Fig. S7). The FISH probes were designed to hybridize to the consensus exons shared among all the 28 variants of Eml4-Alk fusion transcripts that have been identified to date (22). Following prior literature (19, 20), fusion transcripts were detected by co-localized sm-FISH signals targeting Eml4 and Alk transcripts. In a validation test, an average of 12 co-localized sm-FISH signals per cell were detected in the total of 22 H2228 cells (*SI Appendix*, Fig. S8), which was known to express Eml4-Alk fusion transcripts (*SI Appendix*, Fig. S6) (23). In contrast, HEK293T cells exhibited on average zero co-localized signals per cell from 19 cells (*SI Appendix*, Fig. S8), consistent with the lack of such a fusion transcript in HEK293T cells (*SI Appendix*, Fig. S6).

In our collection of 96 tumor samples, only 57 samples had remaining tissues for FuseFISH analysis. These 57 samples included 39 samples that yielded RNA-seq data and 18 samples that did not yield RNA-seq data (Fig. 6B). The FuseFISH analysis detected Eml4-Alk fusion transcripts in two samples, including Sample #44 which was also analyzed by RNA-seq and Sample #37 which did not yield RNA-seq data (Fig. 6 B and C). To test whether Sample #37 had Alk- related fusion genes, we subjected Sample #37 together with other 6 randomly selected samples (#18, #51, #56 #57, #63, #65) for DNA re-combination analysis using Vysis ALK Break Apart FISH (15). None of these 7 samples exhibited Alk- related fusion genes. More specifically, one sample (#57) failed to generate DNA FISH signals from 4 attempts; one sample (#56, negative for Eml4-Alk fusion transcripts by RNA-seq and FuseFISH analyses) exhibited a partial deletion of the Alk gene but no sign of Alk- related fusion genes (star, Fig. 6B). The other 5 samples including Sample #37 exhibited integral Alk genes (Fig. 6 B and D). Taken together, the lung cancer Sample #37 expressed Eml4-Alk fusion transcripts without having an Eml4-Alk fusion gene. These data suggest that genome re-arrangement is not a prerequisite step for creation of fusion transcripts from fusion susceptible pairs. In other words, the RNA-poise model does not require alterations of the DNA.

Discussion

Abundance of genome re-arrangement independent fusion transcripts. Our RNA FISH and DNA FISH analyses revealed a cancer sample that contained a fusion transcript without the corresponding fusion gene. Such an example, although not often seen in literature may not be a rare case (11). The lack of reports are likely attributable to the research attentions paid to the other side of the coin, that is the fusion transcripts created by fusion genes (2). Indeed, approximately $36 - 65\%$ of fusion transcripts derived from cancer RNA-seq data were attributable to genome rearrangement (Low Pass bars, Figure S1A of (1)). However, this is likely an overestimate because when low quality whole genome sequencing (WGS) data were removed, only approximately $30 - 45\%$ fusion transcripts had corresponding WGS reads (High Pass bars, Figure S1A of (1)). These published results are consistent with the notion that fusion genes do not account for all observed fusion transcripts and suggest gene re-arrangement independent fusion transcripts.

The RNA-poise model allows for splicing errors. Fusion transcripts can be created by two means. The better recognized means is through transcription of a fusion gene, which was created by genome-rearrangement. The less recognized means is by RNA splicing errors, where two separate transcripts were spliced together (trans-splicing) (24). Trans-splicing does not involve genome-rearrangement. A theoretical gap in the splicing error model is that trans-splicing can only happen to two RNA molecules that are close to each other in the 3-dimensional (3D) space, however except for neighboring genes (11) the chances for two RNA molecules transcribed from distant chromosomal locations to meet in space are small. Therefore, it remains difficult to perceive a biophysical process in which fusion transcripts are created by splicing errors.

The RNA-poise model fills this theoretical gap. The pre-installation of Gene 1's transcripts on Gene 2's genomic sequence puts Gene 2's nascent transcripts spatially close to Gene 1's transcripts, allowing for the chances of trans-splicing. Furthermore, the majority of splicing events are co-transcriptional. The availability of transcripts of Gene 1 during Gene 2's transcription allows for the chances of making co-transcriptional trans-splicing.

Breaking down the RNA-poise model by the means of RNA-DNA interactions. Remote RNA-DNA interactions could be created by at least two means. First, the caRNA can target specific genomic sequences, which could be mediated by tethering molecules (RNA targeting, Fig. 7). Second, the spatial proximity of the genomic sequences in 3D space could bring the nascent transcripts of one gene to the genomic sequence of another gene (RNA confinement, Fig. 7). Both means of RNA-DNA interactions provide spatial proximity between two RNA molecules and thus allows for splicing errors. In addition, the spatial proximity of two genes in the RNA confinement model could enhance the chances of genome re-arrangement of the spatially close genomic sequences and thus creating fusion genes (25). Thus, the RNA-poise model can be regarded as a union of two sub-models depending on the means of RNA-DNA interaction. One sub-model (Targeting-poise, Fig. 7) could only create fusion transcripts by trans-splicing. The other sub-model (Confinement-poise, Fig. 7) could create fusion transcripts by either trans-splicing or creation of fusion genes.

Materials and Methods

Reference genome and gene annotations. Human genome assembly hg38/GRCh38 and Ensembl gene annotation release 84 (GRCh38.84) were used throughout all data analyses.

Fig. 7. RNA-poise model. In this model, the transcripts of one gene (RNA 1, purple bar) can exhibit spatial proximity to another gene (RNA 2, blue bar) due to tethering (RNA targeting) or spatial proximity of the two genes (RNA confinement). Both cases could facilitate splicing errors (grey arrows), whereas the proximity of genomic sequences may also facilitate gene fusion (grey arrow on the right), which subsequently produce fusion RNA.

Public RNA-seq data. HEK293T RNA-seq datasets were downloaded from NCBI BioProject database (Accession numbers: SRR2992206, SRR2992207, SRR2992208 under the project number PRJNA305831) (16). The three datasets were merged. ₁₉₄ ₁₉₅

TCGA derived fusion transcripts. The TCGA RNA-seq derived fusion transcripts were downloaded from Tumor Fusion Gene Data Portal (http://www.tumorfusions.org) (26). Tier 1 and tier 2 fusion transcripts were used in our analysis. Genomic coordinates were converted to hg38 by liftOver. Following Davidson et al. (8), Futra pairs within 200 kb on hg38 were removed. The data of Futra pairs were mainly processed using R (27) with Bioconductor packages GenomicRanges (28) and InteractionSet (29).

Visualization of Futra pairs. Heatmaps of the count matrix were plotted using Bioconductor package ComplexHeatmap (30). Genomic plots of Futra pairs were created with GIVE (31).

Constructing iMARGI sequencing libraries.

Nuclei preparation and chromatin digestion. Approximately 5×10^6 cells were used for construction of an iMARGI sequencing library. Cells were crosslinked in 1% formaldehyde at room temperature (RT) for 10 minutes with rotation. Crosslinking reaction was quenched with glycine at 0.2 M concentration and incubated at RT for 10 minutes. Cells were pelleted, washed using 1×PBS and aliquoted into ~5×10^6 in each tube. To prepare nuclei, crosslinked cells was incubated in 1 mL of cell lysis buffer (10 mM Tris-HCl pH 7.5, 10 mM NaCl, 0.2% Nonidet P-40, 1× protease inhibitor) on ice for 15 minutes and homogenized with dounce homogenizer pestle A for 20 strokes on ice. Nuclei were pelleted and weighed to estimate the pellet volume (10 mg of nuclei pellet was estimated to be about 10 μL). Nuclei pellet were incubated with SDS buffer (0.5×Cutsmart buffer, 0.5% SDS) with 1:3 volume ratio at 62 ℃ for 10 minutes with mixing and immediately quenched with a final of 1% Triton X-100. To digest chromatin, washed nuclei pellet was re-suspended in AluI chromatin digestion mix (2.3 U/μL AluI (NEB), 0.3 U/μL RNasinPlus, 1× Cutsmart buffer) and incubated at 37 ℃ overnight with mixing. After chromatin digestion, 1 μL of RNase I (1:10 diluted in 1×PBS) was directly added to the reaction mixture and incubated at 37 ℃ for 3 minutes to fragment RNA. Nuclei were pelleted and washed twice using PNK wash buffer (20 mM Tris-HCl pH 7.5, 10 mM MgCl2).

Ligations. To prepare the RNA and DNA ends for linker ligation, nuclei were incubated in 200 μL RNA 3´ end dephosphorylation reaction mix (0.5 U/μL T4 PNK (NEB), 0.4 U/μL RNasinPlus, 1× PNK phosphatase buffer pH 6.5) at 37 ℃ for 30 minutes with mixing. Nuclei were washed twice with PNK buffer, resuspended in 200 μL DNA A-tailing mix (0.3 U/μL Klenow fragment 3´-5´ exo- (NEB), 0.1 mM dATP, 0.1% Triton X-100, 1NEB buffer 2) and incubated at 37 ℃ for 30 minutes with mixing. The same linker sequence as described in the previous MARGI paper was used (10). For *in situ* RNA-linker ligation, nuclei were resuspended in 200 μL ligation mix (38 μL adenylated and annealed linker, 10 U/ μL T4 RNA ligase 2-truncated KQ (NEB), 1× T4 RNA ligase reaction buffer, 20% PEG 8000, 0.1% Triton X-100, 0.4 U/μL RNasinPlus) and incubated at 22 ℃ for 6 hour and then 16 ℃ overnight with mixing. After ligation, the nuclei were washed five times with PNK buffer to remove excess free linker. For *in situ* RNA-DNA proximity ligation, nuclei were resuspended in 2 mL of proximity ligation mixture (4 U/μL T4 DNA ligase (NEB), 1× DNA ligase reaction buffer, 0.1% Triton X-100, 1 mg/mL BSA (NEB), 0.5 U/μL RNasinPlus) and incubated at 16 ℃ overnight.

Library construction. To reverse crosslinking, nuclei were washed twice with 1× PBS, resuspended in 250 μL of extraction buffer (1 mg/mL Proteinase K [NEB], 50 mM Tris-HCl pH 7.5, 1% SDS, 1 mM EDTA, 100 mM NaCl) and incubated at 65 ℃ for 3 hours. DNA and RNA were extracted by adding an equal volume of Phenol:Chloroform:Isoamyl Alcohol (pH 7.9, Ambion) followed by ethanol precipitation. The subsequent steps including removal of biotin from non-proximity ligated linkers, pulldown of RNA-DNA chimera, reverse transcription of RNA, DNA denaturation, circularization, oligo annealing and BamHI (NEB) digestion and sequencing library generation were performed as previously described (10). iMARGI libraries were subsequently subject to pair-end 100-cycle sequencing on Illumina HiSeq 4000. The circularization and library construction strategy can phase RNA and DNA ends into Read 1 and Read 2 as shown Fig. 1A, which is the same with MARGI library configuration (10). Since AluI restriction enzyme recognizes "AGCT" sequence and leaves "CT" at the 5' end of the cut, we expect the first two bases of DNA end (Read 2) to be enriched with "CT".

Analysis of iMARGI sequencing data.

Mapping iMARGI read pairs. The detailed iMARGI data processing methods can be found in our GitHub repository
https://github.com/Zhong-Lab-UCSD/iMARGI_methods. Briefly, it includes three main steps. First, the read pairs were cleaned by
in-house scripts. According to the library construction design, read pairs were filtered out if the 5' most two bases of their DNA end (Read
2) were not "CT". Besides, the first two bases of RNA end (Read 1) were removed as they are random nucleotides. Then, the cleaned read
pairs were mapped to the human genome (hg38) using bwa mem (version 0.7.17) (32) with parameters "-SP5M". Finally, pairtools (version
v0.2.0, https://github.com/mirnylab/pairtools) and in-house scripts were used to parse, deduplicate and filter the mapped read pairs. The
non-redundant read pairs that 5' most ends of both Read 1 and Read 2 were uniquely mapped to the human genome were retained for
further analysis. The read pairs that were mapped to genomic locations within 200 kb to each other were defined as proximal interactions,
which were excluded from our analysis. GenomicRanges (28) and InteractionSet (29) were used for further analysis of iMARGI analysis.

Visualization of iMARGI read pairs. Heatmap of the count matrix were plotted using Bioconductor package ComplexHeatmap (30) . Genomic
plots of iMARGI read pairs were created with Bioconductor package Gviz (33) and GIVE (31).

Intersection of iMARGI read pairs and Futra pairs. An iMARGI read pair was regarded as overlapping with a Futra pair when the RNA-end
was strand-specifically mapped to the gene body of one gene in the Futra pair and the DNA-end was mapped to the gene body $\pm$ 100 kb
flanking regions of the other gene in the Futra pair.

fuseFISH analysis.

Probe design. Oligonucleotide probes were designed to hybridize to exons 2-6 of Eml4 RNA and exons 20-23 of Alk RNA. These exons
were chosen because they were present in all the observed variations of Eml4-Alk fusion genes. These probes were 35-40 nt in lenghs, with
similar GC contents and melting temperatures.

Conjugation of quantum dots to oligonucleotide probes. Oligos were modified on the 5' end with a primary amino group and a spacer of
30 carbons to minimize steric hindrance of probe-RNA hybridization. These probes were conjugated with quantum dots through the
amino group using EDC reaction (34). The probes were subsequently purified with .2 μm membrane filtration and 100,000 molecular
weight cut-off (MWCO). The retentate of the 100,000 MWCO was subjected with dynabeads MyOne SILANE purficiation to remove any
remaining unconjugated probes. A subsequent .2 μm membrane filtration was used to remove any final aggregates.

Hybridization of adherent cell lines. Probe hybridization in H2228 cells was carried out as previously described (19, 21). Briefly, probes
were added to the hybridization solution and incubated with the cells at 37 ℃ overnight. Cells were washed and resuspended in 1× PBS
for imaging.

Hybridization of tissue samples. Probe hybridization in tissue were carried out as previously described (35). Briefly, hybridization solution
with probes were added to the surface of parafilm to form droplets. A tissue slice (5-10 μm in thickness) fixed on a glass cover slip was
gently placed over the hybridization solution. The mixture was incubated at 37 ℃ overnight. The tissue was subsequently washed with
wash buffer and resuspended in 1x PBS for imaging.

Imaging and analysis. Cells or tissues were imaged in 1× PBS through wide field fluorescence imaging using an Olympus IX83 inverted
microscope at 60× oil immersion objective (NA=1.4). Image processing was carried out as previously described (36). Briefly, single
transcripts were detected using an automated thresholding algorithm that searches for robust thresholds. where counts do not change
within a range. Fusion transcripts were determined by searching for co-localization of detection transcripts by overlap between predicted
centers within a radius.

RNA sequencing and analysis. RNA was extracted with Trizol from H2228 cell line and lung cancer tissue samples of the approximate size
3 mm×3 mm×30 μm per sample. RNA-sequencing was carried out using the TruSight RNA Pan-Cancer Panel (Illumina) following the
manufacterer's protocol. All the RNA-seq data, including HEK293T public data, H2228 cell line and lung cancer sample sequencing data,
were mapped to the human genome (hg38) using STAR (v2.5.4b) with default parameters (37). Fusion transcripts were called using
STAR-Fusion (v0.8.0) (17) requiring both numbers of supporting discordant read pair and junction spanning read larger than zero and the
sum of them larger than 2.

ACKNOWLEDGMENTS. Male hTert-immortalized human foreskin fibroblasts (HFF-hTert-clone 6) are a cell line of the 4D Nucleome
Tier 1 cells, provided by Job Dekker lab (https://www.4dnucleome.org/cell-lines.html). This work is funded by NIH R00122368 (Z.C.),
HL106579 (S.C.), HL108735 (S.C.), DP1HD087990 (S.Z.), and NIH 4D Nucleome U01CA200147 (S.C. and S.Z.).

1. Gao Q, et al. (2018) Driver fusions and their implications in the development and treatment of human cancers. *Cell Rep* 23(1):227–238 e3.
2. Mertens F, Johansson B, Fioretos T, Mitelman F (2015) The emerging complexity of gene fusions in cancer. *Nat Rev Cancer* 15(6):371–81.
3. Kumar-Sinha C, Kalyana-Sundaram S, Chinnaiyan AM (2015) Landscape of gene fusions in epithelial cancers: seq and ye shall find. *Genome Med* 7:129.
4. Dai X, Theobard R, Cheng H, Xing M, Zhang J (2018) Fusion genes: A promising tool combating against cancer. *Biochim Biophys Acta Rev Cancer* 1869(2):149–160.
5. Stransky N, Cerami E, Schalm S, Kim JL, Lengauer C (2014) The landscape of kinase fusions in cancer. *Nat Commun* 5:4846.
6. Yoshihara K, et al. (2015) The landscape and therapeutic relevance of cancer-associated transcript fusions. *Oncogene* 34(37):4845–54.
7. Torres-Garcia W, et al. (2014) Prada: pipeline for rna sequencing data analysis. *Bioinformatics* 30(15):2224–6.
8. Davidson NM, Majewski IJ, Oshlack A (2015) Jaffa: High sensitivity transcriptome-focused fusion gene detection. *Genome Med* 7(1):43.
9. Lai J, et al. (2015) Fusion transcript loci share many genomic features with non-fusion loci. *BMC Genomics* 16:1021.
10. Sridhar B, et al. (2017) Systematic mapping of rna-chromatin interactions in vivo. *Curr Biol* 27(4):610–612.
11. Varley KE, et al. (2014) Recurrent read-through fusion transcripts in breast cancer. *Breast Cancer Res Treat* 146(2):287–97.
12. Kim J, et al. (2015) Recurrent fusion transcripts detected by whole-transcriptome sequencing of 120 primary breast cancer samples. *Genes Chromosomes Cancer* 54(11):681–91.
13. Nagano T, et al. (2015) Comparison of hi-c results using in-solution versus in-nucleus ligation. *Genome Biol* 16:175.
14. van de Werken HJ, et al. (2012) Robust 4c-seq data analysis to screen for regulatory dna interactions. *Nat Methods* 9(10):969–72.
15. Dixon JR, et al. (2012) Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* 485(7398):376–80.
16. Wynne JW, et al. (2017) Comparative transcriptomics highlights the role of the activator protein 1 transcription factor in the host response to ebolavirus. *J Virol* 91(23).
17. Haas B, et al. (2017) Star-fusion: Fast and accurate fusion transcript detection from rna-seq. *bioRxiv* (120295).
18. Vendrell JA, et al. (2017) Detection of known and novel alk fusion transcripts in lung cancer patients using next-generation sequencing approaches. *Sci Rep* 7(1):12510.
19. Semrau S, et al. (2014) Fusefish: robust detection of transcribed gene fusions in single cells. *Cell Rep* 6(1):18–23.
20. Markey FB, Ruezinsky W, Tyagi S, Batish M (2014) Fusion fish imaging: single-molecule detection of gene fusion transcripts in situ. *PLoS One* 9(3):e93488.
21. Nguyen TC, et al. (2016) Mapping rna-rna interactome and rna structure in vivo by mario. *Nat Commun* 7:12023.
22. Forbes SA, et al. (2017) Cosmic: somatic cancer genetics at high-resolution. *Nucleic Acids Res* 45(D1):D777–D783.
23. Martelli MP, et al. (2009) Eml4-alk rearrangement in non-small cell lung cancer and non-tumor lung tissues. *Am J Pathol* 174(2):661–70.

24. Li H, Wang J, Mor G, Sklar J (2008) A neoplastic gene fusion mimics trans-splicing of rnas in normal human cells. *Science* 321(5894):1357–61.   301

25. Gu W, Zhang F, Lupski JR (2008) Mechanisms for human genomic rearrangements. *Pathogenetics* 1(1):4.   302

26. Hu X, et al. (2018) Tumorfusions: an integrative resource for cancer-associated transcript fusions. *Nucleic Acids Res* 46(D1):D1144–D1149.   303

27. R Core Team (2018) *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria).   304

28. Lawrence M, et al. (2013) Software for computing and annotating genomic ranges. *PLOS Computational Biology* 9(8):e1003118.   305

29. Lun AT, Perry M, Ing-Simmons E (2016) Infrastructure for genomic interactions: Bioconductor classes for hi-c, chia-pet and related experiments. *F1000Research* 5.   306

30. Gu Z, Eils R, Schlesner M (2016) Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* 32(18):2847–2849.   307

31. Cao X, Yan Z, Wu Q, Zheng A, Zhong S (2018) Give: portable genome browsers for personal websites. *Genome Biol* 19(1):92.   308

32. Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with bwa-mem. *arXiv preprint arXiv:1303.3997*.   309

33. Hahne F, Ivanek R (2016) *Visualizing genomic data using Gviz and bioconductor.* (Springer), pp. 335–351.   310

34. Choi Y, et al. (2009) In situ visualization of gene expression using polymer-coated quantum-dot-dna conjugates. *Small* 5(18):2085–91.   311

35. Lyubimova A, et al. (2013) Single-molecule mrna detection and counting in mammalian tissue. *Nat Protoc* 8(9):1743–58.   312

36. Raj A, van den Bogaard P, Rifkin SA, van Oudenaarden A, Tyagi S (2008) Imaging individual mrna molecules using multiple singly labeled probes. *Nat Methods* 5(10):877–9.   313

37. Dobin A, et al. (2013) Star: ultrafast universal rna-seq aligner. *Bioinformatics* 29(1):15–21.   314