A splice donor variant in *CCDC189* is associated with asthenospermia in Nordic Red

dairy cattle

Terhi Iso-Touru[*], Christine Wurmser[†], Heli Venhoranta[‡], Maya Hiltpold[§], Tujia Savolainen[‡],

Anu Sironen[*], Konrad Fischer[**], Krzysztof Flisikowski[**], Ruedi Fries[†], Alejandro Vicente-

Carrillo[††], Manuel Alvarez-Rodriguez[††], Szabolcs Nagy[‡‡], Mervi Mutikainen[*], Jaana

Peippo[*], Juhani Taponen[‡], Goutam Sahana[§§], Bernt Guldbrandtsen[§§], Henri Simonen[***],

Heriberto Rodriguez-Martinez[††], Magnus Andersson[‡,1], Hubert Pausch[§,1]

[*]Natural Resources Institute Finland (Luke), 31600 Jokioinen, Finland

[†]Chair of Animal Breeding, Technische Universität München, 85354 Freising-
Weihenstephan, Germany

[‡] University of Helsinki, 04920 Saarentaus, Finland

[§]Animal Genomics, ETH Zurich, 8001 Zurich, Switzerland

[**]Chair of Livestock Biotechnology, Technische Universität München, 85354 Freising-

Weihenstephan, Germany

[††]Department of Clinical and Experimental Medicine, Linköping University, 58183

Linköping, Sweden

[‡‡]University of Pannonia, 8360 Keszthely, Hungary

[§§]Department of Molecular Biology and Genetics, Aarhus University, 8830 Tjele, Denmark

[***]VikingGenetics, 15870 Hollola, Finland

[1]joint senior authorship

Authors and e-mail addresses:
Terhi Iso-Touru: terhi.iso-touru@luke.fi
Christine Wurmser: christine.wurmser@tum.de
Heli Venhoranta: heli.venhoranta@googlemail.com
Maya Hiltpold: maya.hiltpold@usys.ethz.ch
Tujia Savolainen: tuiperi@gmail.com
Anu Sironen: anu.sironen@luke.fi
Konrad Fischer: konrad.fischer@wzw.tum.de
Krzysztof Flisikowski: flisikowski@wzw.tum.de
Ruedi Fries: ruedi.fries@tum.de
Alejandro Vicente-Carrillo: alejandro.vicentecarrillo@gmail.com

Manuel Alvarez-Rodriguez: manuel.alvarez-rodriguez@liu.se
Szabolcs Nagy: szabio@hotmail.com
Mervi Mutikainen: mervi.mutikainen@luke.fi
Jaana Peippo: jaana.peippo@luke.fi
Juhani Taponen: juhani.taponen@helsinki.fi
Goutam Sahana: goutam.sahana@mbg.au.dk
Bernt Guldbrantsen: bernt.guldbrandtsen@mbg.au.dk
Henri Simonen: hesim@vikinggenetics.com
Heriberto Rodriguez-Martinez: heriberto.rodriguez-martinez@liu.se
Magnus Anderson: magnus.andersson@helsinki.fi
Hubert Pausch: hubert.pausch@usys.ethz.ch

47
48
49

66 **Short running title**: Recessive asthenospermia in cattle

68 **Key words**: male infertility, sterility, immotile sperm, cattle, CCDC189

70 **Corresponding author:**

71 Hubert Pausch, ETH Zürich, Animal Genomics, Eschikon 27, 8315 Lindau, Switzerland,
72 Phone: +41 52 354 92 34, E-mail: hubert.pausch@usys.ethz.ch

Abstract

Cattle populations are highly amenable to the genetic mapping of male reproductive traits because longitudinal data on ejaculate quality and dense microarray-derived genotypes are available for many artificial insemination bulls. Two young Nordic Red bulls delivered sperm with low progressive motility (i.e., asthenospermia) during a semen collection period of more than four months. The bulls were related through a common ancestor on both their paternal and maternal ancestry. Thus, a recessive mode of inheritance of asthenospermia was suspected. Both bulls were genotyped at 54,001 SNPs using the Illumina BovineSNP50 Bead chip. A scan for autozygosity revealed that they were identical by decent for a 2.98 Mb segment located on bovine chromosome 25. This haplotype was not found in the homozygous state in 8,557 fertile bulls although five homozygous haplotype carriers were expected (P=0.018). Whole genome-sequencing uncovered that both asthenospermic bulls were homozygous for a mutation that disrupts a canonical 5' splice donor site of *CCDC189* encoding the coiled-coil domain containing protein 189. Transcription analysis showed that the derived allele activates a cryptic splice site resulting in a frameshift and premature termination of translation. The mutated CCDC189 protein is truncated by more than 40%, thus lacking the flagellar C1a complex subunit C1a-32 that is supposed to modulate the physiological movement of the sperm flagella. The mutant allele occurs at a frequency of 2.5% in Nordic Red cattle. Our study in cattle uncovered that CCDC189 is required for physiological movement of sperm flagella thus enabling active progression of spermatozoa and fertilization.

Background

Impaired male fertility is a well-recognized and highly heterogeneous condition in many species including humans, mice and cattle (*e.g.,* [1],[2],[3],[4],[5]). Males with reduced reproductive ability may produce ejaculates with anomalous characteristics such as low sperm count, impaired progressive motility or morphological aberrations of the spermatozoa [6]. However, standard semen parameters are only poor predictors of male reproductive ability because many fertility disorders are caused by aberrations of the spermatozoa that are not immediately apparent or regularly monitored (e.g., [7],[8],[4]).

The normal movement of the sperm flagellum, i.e., activated and hyperactivated motility, facilitates active progression of spermatozoa through the oviduct. Morphological

aberrations of the sperm tail such as stump tail, short tail, dysplasia of the fibrous sheath or disorganized flagellar microtubules may compromise sperm motility and fertilization success [9]. Genetic variants causing morphological abnormalities of the sperm tail have been identified in several species including humans [10],[11], pigs [12], cattle [13], and mice [14] where they impede motility. The fertilization success of ejaculates containing a high proportion of spermatozoa with low progressive motility (i.e., asthenospermia) is often reduced, both *in vivo* and *in vitro* [15],[16], albeit fertilization anecdotally being reported as possible *in vivo* [17].

Due to the low heritability of male reproductive performance, detecting genetic variants that underpin impaired male fertility requires large cohorts with phenotypic records that, moreover, have been genotyped [18]. Comprehensive data on male fertility have become available for many cattle populations due to the wide-spread use of artificial insemination (AI) [19]. The availability of both longitudinal data on semen quality and dense microarray-derived genotypes makes cattle populations highly amenable to the mapping of genetic determinants that underpin male reproductive performance [4],[13],[20],[21].

Prospective bulls for AI are selected at an early age and reared at AI centers under controlled conditions. When the bulls reach an age of approximately one year, their semen is collected once or twice per week. Immediately after semen collection, macroscopic, microscopic and computer-assisted analyses are carried out to assess whether the ejaculates are suitable for AI [22]. Up to 20% of all collected ejaculates are rejected because they do not fulfill the requirements for AI [23]. The most common pathological findings in ejaculates of insufficient quality include absence or low number of spermatozoa, reduced sperm motility, impaired sperm viability or aberrant morphology of heads and/or tails in many spermatozoa [24]. Usually, it takes a few weeks until young bulls produce semen that is considered suitable for AI, owing to individual differences in epididymal sperm maturation [22]. Young bulls that permanently produce ejaculates that do not fulfill the requirements for AI are sporadically noticed at semen collection centers. It is plausible that some of these bulls carry deleterious genetic variants that are responsible for the low semen quality they deliver (e.g., [4],[13]).

Here we present the phenotypic and genetic analysis of a recessively inherited form of sterility in Nordic Red bulls, which primarily manifests itself through asthenospermia. The availability of dense genotype data of two affected bulls enabled us to localize the disorder within a 2.98 Mb interval on bovine chromosome 25. Whole-genome sequencing of

138   affected bulls revealed that a mutation disrupting a canonical splice donor site of the

139   *CCDC189* gene is the most likely cause of the observed condition.

Material and Methods

Ethics statement

Ejaculates of two asthenospermic bulls were collected by employees of an AI center in

Finland as part of their routine practice under veterinary supervision. Both bulls were

eventually culled because their impaired sperm motility precluded the use of the collected

ejaculates for artificial inseminations. The decision to cull the bulls was at the sole

discretion of representatives of the AI center. Testicular tissue was collected immediately

after slaughter. Oocytes used for *in vitro* fertilization experiments were collected from offal

cow ovaries at a commercial slaughterhouse. No animal experiments were performed in

this study, and, therefore, approval from an ethics committee was not required.

Animals

Two fifteen-months old bulls from the Nordic Red dairy cattle breed that were born in 2012

and 2013 were housed at an AI center in order to collect their semen for AI. Both bulls

were apparently healthy and their andrological status was considered as normal, i.e., they

showed no signs of disease, the size of their testicles was within the normal range and

their libido was normal. During a period of more than four months, 11 and 27 ejaculates,

respectively, of both bulls were collected once or twice a week by technicians at the AI

center. Semen from bulls of documented normal sperm quality and fertility, of the same

breed and age interval, was likewise collected, processed and examined. Testicular tissue

samples were collected immediately after slaughter, frozen in liquid nitrogen and stored at

-80°C.

Analysis of fresh semen

Immediately after semen collection, sperm motility, viability and morphology of fresh

ejaculates were subjectively assessed by technicians at the AI center. Sperm motility was

evaluated using a microscope equipped with phase contrast optics. Sperm concentration

and plasma membrane viability were assessed using a flow-cytometer [25]. None of the

collected ejaculates fulfilled the requirements for AI because most spermatozoa had

reduced or missing progressive motility. Upon our request, one ejaculate from each of the

asthenospermic bulls was frozen using the commercial method routinely used by the AI
center.

Analysis of frozen-thawed semen samples

Velocity (μm/sec), and percentages of spermatozoa with total and progressive motility
were assessed using frozen-thawed ejaculates. Semen droplets (2.4 x 10^5 spermatozoa in
10 μL) were placed on a pre-warmed 76 x 26 mm Menzel-Gläser microscope slide
(ThermoFisher Scientific, Waltham, MA, USA) covered by a pre-warmed coverslip on a
thermal plate (38 ºC). Slides were analyzed using an upright Zeiss Axio Scope A1 light
microscope equipped with a 10x phase contrast objective (Carl Zeiss, Stockholm,
Sweden) connected via a CMOS camera (UEye, IDS Imaging Development Systems
GmbH, Obersulm, Germany) to a computer running the Qualisperm^{TM} software (Biophos
SA, Lausanne, Switzerland). The Qualisperm^{TM} technology is based on fluorescence
correlation spectroscopy-analysis of single particles (spermatozoa) in confocal volume
elements. Individual spermatozoa are projected on a pixel grid of the CMOS camera and
the algorithm calculates the number of fluctuations in each pixel using a correlation
function of sperm numbers and translation classes, thus determining the speed (velocity)
distribution of spermatozoa, from which proportions of motility classes are calculated [26].

Sperm viability (membrane integrity) and early membrane destabilization changes (YO-
PRO-1+/PI-), degree of mitochondrial activity (Mitotracker Deep Red+) and superoxide
production (MitoSOX+), production of reactive oxygen species (ROS; CellROX+) and lipid
peroxidation (BODIPY+) were assessed using a Gallios™ flow cytometer (Beckman
Coulter, Bromma, Sweden) equipped with standards optics. In all investigations, we
assessed 25,000 events per sample with a flow rate of 500 cells per second.

The total ATP concentration of frozen-thawed spermatozoa was analyzed using the ATP
Determination Kit A22066 (Molecular Probes TM Invitrogen detection technologies)
according to the manufacturer's protocol. Samples of an asthenospermic and
normospermic bull were analyzed using a Kikkoman Lumitester C-100 luminometer
(Kikkoman Corporation, Nishishinbashi, Tokyo, Japan).

Transmission electron microscopy on spermatozoa of two asthenospermic and one
normospermic bull was performed using frozen-thawed semen samples. After thawing,
semen samples were washed twice in phosphate buffered saline (PBS), fixed with 2.5%
glutaraldehyde and 2% paraformaldehyde in PBS for 2 hours on ice and pelleted at 6000g.

Pellets were washed in PBS, post-fixed in 1% osmium tetroxide, washed in ddH_2O,
stained in 1% uranyl acetate, dehydrated in graded series of ethanol (25%, 50%, 75%,
90%, and 100%), and embedded in Epoxy resin through increasing concentrations (25%,
50%, 75%, and 100%) using PELCO Biowave+ tissue processor, and then cured at 60°C
for 3 days. Embedded blocks were sectioned using Leica FC6 microtome and a DIATOME
diamond knife with 45° angle into 60nm sections and mounted on Quantifoil copper grids
with formvar carbon films. Sections were post-stained with 2% uranyl acetate followed by
lead citrate. Grids were imaged using FEI Morgani 268 electron microscope operated at
100 kV at 20k magnification.

In vivo and in vitro fertilization success

Fertility of two asthenospermic bulls was tested *in vivo* at a conventional Finnish cattle
herd within one breeding season; ninety-eight cows in oestrus were inseminated by the
farmer either with semen from one of two bulls with asthenospermia (n=9) or a control bull
with normal semen quality (n=89). Insemination success was controlled by a trained
veterinarian.

In vitro fertilization (IVF) success, with embryo cleavage rate at 42 hours post insemination
and embryo yield at 9 days post insemination as end-points, was tested for an
asthenospermic and a normospermic bull, using 470 offal oocytes. Ovaries of slaughtered
cows were collected at a commercial slaughterhouse and transported to the laboratory in
saline at room temperature. In the laboratory, follicles of 2-8 mm in diameter were
aspirated with an 18 G needle and a 20 ml syringe. Cumulus-oocyte complexes (COC)
were collected from the aspirated follicular fluid and 470 good quality COC were selected
for maturation. Unless stated otherwise, all the chemicals were purchased from Sigma-
Aldrich (St. Louis, MO, USA). COC were matured for 24h in TCM199 with glutamax-I
(Gibco™; Invitrogen Corporation, Paisley, UK) supplemented with 0.25mM Na-pyruvate,
100 IU/ml penicillin, 100 µg/ml streptomycin, 2 ng/ml FSH (Puregon, Organon, Oss,
Netherlands), 1 µg/ml β-estradiol (E-2257) and 10% heat inactivated fetal bovine serum
(FBS) (Gibco™, New Zealand) at 38.5°C in maximal humidity in 5% CO_2 in air. Following
maturation, 320 and 150 COC were inseminated using washed frozen–thawed semen
from an asthenospermic and a normospermic control bull of proven fertility in IVF,
respectively, at a concentration of 1 million spermatozoa per ml at 38.5°C in maximal
humidity in 5% CO_2 in air for 20h. During IVF spermatozoa capacitation and movement
were stimulated with 10 µg/ml heparin and PHE (20 µM penicillamine - 10 µM hypotaurine

234 – 1 μM epinephrine), respectively. After fertilization, presumptive zygotes were cultured in

235 G1/G2 media (Vitrolife, Göteborg, Sweden) supplemented with fatty-acid-free bovine

236 serum albumin (4 mg/ml) at 38.5°C in maximal humidity in an atmosphere of 5% O_2 and

237 5% CO_2. Cleavage rates of oocytes were recorded at 30 and 42 hours post insemination.

238 The number and quality of blastocysts were recorded at day 8 and 9 (IVF=day 0).

Genotyping, quality control and haplotype inference

The two asthenospermic bulls were genotyped at 54,001 SNPs using the Illumina

BovineSNP50 Bead chip using DNA purified from semen samples following standard DNA

extraction protocols. To facilitate haplotype inference, we also considered genotypes of

119 fertile bulls from the Nordic Red dairy cattle breed, including the sires (n=2) and 17

paternal half-sibs of the affected bulls. Genotypes for these 119 bulls were also

determined using the Illumina BovineSNP50 Bead chip comprising genotypes at 54,001

(version 1) or 54,609 SNPs (version 2). The chromosomal positions of the SNPs

corresponded to the University of Maryland reference sequence of the bovine genome

(UMD3.1) [27]. Following quality control, we retained 41,594 autosomal SNPs with a call-

rate greater than 95%. The per-individual call-rate was between 96.07% and 100% with an

average call-rate of 99.73%. The *Beagle* [28] software (version 3.2.1) was used to impute

sporadically missing genotypes and infer haplotypes. Genotype data are available in *plink*

binary format [29] (.bed/.bim/.fam) as **Supplementary File 1**.

Detection of runs of homozygosity

Runs of homozygosity (ROH) were identified in both asthenospermic bulls using the

homozyg-function that was implemented in the *plink* (version 1.9) software tool [29]. Due

to the relatively sparse SNP coverage (approximately 1 SNP per 60 kb), we considered

segments with a minimum length of 500 kb *(--homozyg-kb 500)* and at least 20 contiguous

homozygous SNPs (*--homozyg-snp 20*) as ROH. ROH were identified within sliding

windows of 40 SNPs (*--homozyg-window-snp 40*). For the identification of ROH, no

heterozygous SNP were allowed within ROH (*--homozyg-het 0*). The R script that was

used to identify and visualize ROH is available as **Supplementary File 2**.

Identification of segments of autozygosity in two bulls with asthenospermia

263 Genotypes contained in ROH in both asthenospermic bulls were manually inspected in

264 order to determine whether the bulls were identical by descent (IBD) for the same

265 haplotype (i.e., autozygous). In regions where both affected bulls were IBD, the haplotypes

of 119 fertile bulls (see above) were also inspected. Assuming recessive inheritance of a deleterious mutation, we considered IBD haplotypes as associated with asthenospermia if they were not detected in the homozygous state in 119 fertile bulls.

Generation of whole genome sequencing data and sequence variant detection

Genomic DNA was prepared from semen samples of the two asthenospermic bulls according to standard DNA extraction protocols. Paired-end libraries were prepared using the paired-end TruSeq DNA sample preparation kit (Illumina) and sequenced using a HiSeq 2500 instrument (Illumina). We collected 252 and 270 million paired-end reads (125 base pairs) corresponding to an average (raw) genome coverage of 12- and 13-fold, respectively. The raw sequencing data of both asthenospermic bulls are available at the European Nucleotide Archive (ENA) (http://www.ebi.ac.uk/ena) under primary accession number PRJEB29487. The raw sequencing reads were trimmed and filtered using *fastp* (version 0.19.4) [30] and subsequently aligned to the UMD3.1 reference sequence (GCA_000003055.3) of the bovine genome [27] using the *BWA mem* algorithm (version 0.7.13) [31]. Individual files in SAM format were converted into BAM format using *SAMtools* (version 1.8) [32]. Duplicate reads were marked with the *MarkDuplicates* command of the *Picard tools* software suite (https://broadinstitute.github.io/picard/). Polymorphic sequence variants were identified using the *Genome Analysis Toolkit* (*GATK*, version 4.0) [33]. Sequencing coverage along the genome was calculated based on the aligned reads using the *Mosdepth* software [34].

Identification of candidate causal variants

Assuming that both bulls with asthenospermia were homozygous for the same recessively inherited variant, we screened the sequence data of the affected bulls for variants that were: a) located within the 2.98-Mb segment of autozygosity, b) homozygous for the non-reference allele in both bulls and c) never observed in the homozygous state in a control group of 237 animals from breeds other than Nordic Red dairy cattle that were obtained from Schwarzenbacher *et al.* [35]. 126 variants that fulfilled these three criteria were considered as candidate causal variants. The functional consequences of the candidate causal variants were predicted according to the Ensembl transcripts database (release 91) using the *Variant Effect Predictor* (VEP) [36] tool. Genotypes of candidate causal variants were subsequently inspected in whole-genome sequencing data of 2,669 animals that have been sequenced in Run6 of the 1000 Bull Genomes Project [37].

Validating the CCDC189 splice donor variant in Nordic Red cattle

We estimated the frequency of the asthenospermia-associated haplotype in 8,557 Nordic
Red cattle from the Nordic genomic selection reference population that had been
genotyped at more than 50,000 SNPs [38]. Using Sanger sequencing, we genotyped the
splice donor variant (GCA_000003055.3:Chr25:g.27138357C>T) in *CCDC189* (transcript-
ID: ENSBTAT00000045037) in 118 sires that were either identified as heterozygous
carriers of the asthenospermia-associated haplotype (N=30) or have been used frequently
in AI (N=88). PCR reactions for genotyping were done with the DyNAzyme II DNA
Polymerase (Thermo Fisher, MA, US) in a 10µl volume of 1x PCR buffer, 0.2mM dNTPs,
10pmol primer mix (forward 5'- CAAGGTCCTGCCCTTAAGAA -3') and reverse primer:
(reverse 5'- ATGCCATCACTCTGGACCTC-3') and 150pg of DNA. The cycling conditions
were the following: a) an initial denaturation at 95°C for 3 min, b) 29 cycles of 30 sec
denaturation (94°C), 30 sec hybridization (58°C), 30 sec elongation (72°C), and c) a final 3
min elongation (72°C). PCR products were purified and directly sequenced using the
BigDye Terminator Cycle Sequencing Kit (Applied Biosystems, CA, US). Electrophoresis
of sequencing reactions was performed on 3500xL Genetic Analyzers (Applied
Biosystems, CA, US), and sequences were visualized with Sequencher 5.4.6 (Gene
Codes Corporation, MI, US).

Transcription analysis

Total RNA from two bulls homozygous for the *CCDC189* splice donor variant
(GCA_000003055.3:Chr25:g.27138357C>T) and two control bulls that were homozygous
for the wild type allele was extracted from testis tissues using the RNeasy Mini Kit
(Qiagen) following manufacturer's instructions. The quality and concentration of total RNA
was analysed using Agilent's 2100 Bioanalyzer and the Nanodrop ND-2000
spectrophotometer (Thermo Scientific). An amount of 500 ng of total RNA from each
sample was reversely transcribed using QuantiTech Rev. Transcription kit (Qiagen). PCR
reactions were done with the DyNAzyme II DNA Polymerase (Thermo Fisher, MA, US) in a
30 µl volume of 1 x PCR buffer, 0.2 mM dNTPs, 10 pmol primer mix (forward 5'-
ACTGAGGAGATGAGGGAGGT-3' and reverse primer: reverse 5'-
TAGAGCTTGAAGTGGCGGAA-3') and 50 ng of cDNA. The cycling conditions were the
following: 1) an initial denaturation at 95°C for 3 min, 2) 35 cycles of 30 sec denaturation
(94°C), 30 sec hybridization (61.8°C), 1 min elongation (72°C) and 3) a final 3 min
elongation (72°C). PCR products were separated on a 4% agarose gel and the length of

the products was analysed using the Image Lab version 5.0 software (Bio-Rad

Laboratories). PCR products were purified with QIAquick Gel Extraction kit (QIAgen) and

sequenced using the BigDye Terminator Cycle Sequencing Kit (Applied Biosystems, CA,

US). Electrophoresis of sequencing reactions was performed on 3500xL Genetic Analyzer

(Applied Biosystems, CA, US), and sequences were visualized with Sequencher 5.4.6

(Gene Codes Corporation, MI, US).

Results

Aberrant mitochondrial activity in ejaculates of asthenospermic bulls

Two young bulls from the Nordic Red dairy cattle breed producing ejaculates with immotile

spermatozoa were noticed at a semen collection center. The average ejaculate volumes

(3.16 ml) and sperm concentrations (1.94 Mio/µl) were within physiological ranges for bulls

of their age. More than 80% of the spermatozoa in the ejaculates of both bulls were

classified as viable and had typical morphology. However, less than 5% of the

spermatozoa were progressively motile, thus not fulfilling the requirements for AI, whose

threshold is at least 40%.

Frozen semen samples from both asthenospermic bulls were thawed and subjected to

computer-assisted motility analysis and flow cytometric studies. Post-thawing, the velocity

of the spermatozoa was between 10 and 14 µm/s. The velocity of spermatozoa in

normospermic control bulls was between 38 and 48 µm/s (sperm with progressive motility

typically move at a velocity faster than 25 µm/s). In two normospermic control bulls, more

than 50% of the spermatozoa showed progressive motility after thawing while in the

affected bulls, the proportion of motile spermatozoa was less than 10%. The discrepancy

between computer-assisted and subjective motility evaluations (10 vs 5%) results from

spermatozoa displaying minor movement, i.e., the sperm tails bended peculiarly which

was not classified as progressive motility in subjective microscopic sperm evaluation

(**Figure 1**, **Supplementary File 3 & 4**). The ATP content was numerically higher, though

not significant, in semen of asthenospermic than normospermic bulls. As expected, flow

cytometry analysis revealed that viable post-thawed spermatozoa examined in either bull

type had a tendency for destabilizing changes in the plasmalemma. However, the

asthenospermic bulls had more spermatozoa with high mitochondrial membrane potential

indicating increased mitochondrial activity. This leads to more spermatozoa producing

superoxide and reactive oxygen species. This in turn causes a higher proportion of

spermatozoa with lipid peroxidation in asthenospermic than normospermic bulls (**Table 1**).

Transmission electron microscopy of the sperm flagella from asthenospermic bulls showed

the typical arrangement of nine outer microtubule doublets surrounding the central pair

(9x2 + 2). No ultrastructural abnormalities of the axonemes were immediately apparent

(**Figure 1**).

Figure 1 – Phenotypic manifestation of asthenospermia. Microscopic view of a representative
sample of frozen-thawed spermatozoa from an asthenospermic (A) and normospermic (b) bull.
Bending of many spermatozoa of the asthenospermic bull and the lack of motility projected as
clear pictures of spermatozoa (A). Foggy (double pictured) representation of many spermatozoa in
the normospermic bull indicate actively moving spermatozoa (B). Both figures were photographed
using the same settings. Transmission electron microscopy of spermatozoa from two
asthenospermic (C, D, F, G) and an normospermic (E, H) bull. Representative cross-sections of
the midpiece (C-E) and principal piece (F-H) of the flagella revealed the typical arrangement of
nine outer microtubule doublets surrounding the central pair (9x2 + 2) in asthenospermic and
normospermic bulls. No aberrations of the axonemal ultrastructures including outer dense fiber
(ODF), radial spokes (RS), inner (ID) and outer dynein arms (OD) were apparent in cross-sections
of asthenospermic bulls.

    Table 1 - Results of the flow cytometric analysis of frozen thawed semen samples.

Animal	Early membrane destabilization	Increased mitochondrial activity	Increased mitochondrial superoxide production	Increased ROS production	Lipid peroxidation degree
Asthenospermic-1	2.0	72.3	24.9	39.0	22.0
Asthenospermic-2	4.2	71.9	25.9	17.8	23.3
Normospermic-1	10.7	47.3	4.9	5.1	3.1
Normospermic-2	8.3	54.3	1.8	6.7	3.2

Proportions (%) of viable spermatozoa post-thawing from two asthenospermic and two normospermic bulls of the Nordic Red breed depicting sperm early membrane destabilization (YO-PRO-1+/PI-), degree of mitochondrial activity (Mitotracker+/PI-), and increased production of mitochondrial superoxide (MitoSOX+/PI-), reactive oxygen species (ROS, CellROX+) and lipid peroxidation (Bodipy-C11 + (green)) assessed using flow cytometry.

Bulls with asthenospermia are sterile *in vivo* and *in vitro*

None of the 9 cows inseminated with semen of an asthenospermic bull conceived, whereas 42 of the 89 cows inseminated with normospermic semen did.

In vitro fertilization success was tested using 470 offal oocytes. Using post-thaw semen from an asthenospermic bull, the embryo cleavage rate at 42 hours post-insemination was only 11% (36 out of 320) compared to 70% (105 out of 150) using spermatozoa from a normospermic control bull. At day nine post-insemination, the embryo yield after embryo culture was only 0.9% (3 out of 320) for the asthenospermic bull compared to 23% (34 out of 150) for the normospermic bull. Moreover, the three embryos obtained using semen from the asthenospermic bull reached blastocyst stage only at day 9, indicating aberrantly slow development kinetics.

Because the semen analyses revealed consistent sperm deviations and deficient fertilization capacity both *in vivo* and *in vitro* in both asthenospermic bulls, we suspected that this type of sterility may represent a genetic condition. The analysis of pedigree records revealed a common male ancestor (born in 1990) that was present either as great-great-great-grandfather or great-great-grandfather in both the maternal and paternal ancestry of both affected bulls (**Supplementary File 5**). The sires of the two asthenospermic bulls had more than 300 and 2,000 daughters, respectively, indicating that they were frequently used for AI. According to the AI centers consulted, their semen quality was normal. Seven and twelve paternal half-sibs of the asthenospermic bulls were used for 6,257 and 36,689 artificial inseminations, respectively. According to the AI center,

their semen qualities were normal and their average calving rates (number of births per insemination) were 0.39 and 0.44, i.e., similar to average values in Nordic Red cattle. Because a common male ancestor was in the maternal and paternal ancestry of both asthenospermic bulls and fertility was normal in their first- and second-degree relatives, we suspected that the disorder might follow an autosomal recessive mode of inheritance.

Asthenospermia maps to a 2.98 Mb segment on bovine chromosome 25

To identify the genomic region associated with asthenospermia, the two affected bulls were genotyped using the BovineSNP50 Bead chip. Following quality control, genotypes of 41,594 autosomal SNPs were available for genetic analyses. Assuming recessive inheritance of a deleterious mutation, the genotypes of both asthenospermic bulls were screened for the presence of long ROH (**Supplementary Files 1 & 2**). We identified 49 and 28 ROH in the two bulls that covered 258 and 210 Mb of their genomes (**Figure 2**). The average length of the ROH was 6.1 Mb ranging from 1.4 to 30.4 Mb. ROH at chromosomes 2, 4, 9, 10, 14 and 25 were present in both asthenospermic bulls. However, the ROH at chromosomes 4, 9, 10 and 14 were homozygous for different haplotypes and thus not compatible with the presumed inheritance of a recessive allele from a common ancestor. Two haplotypes on chromosomes 2 (between 90,155,450 and 92,350,004 bp) and 25 (between 25,490,468 and 28,470,779 bp) were identical-by-descent (IBD) in both bulls with asthenospermia. The IBD haplotype located on chromosome 2 consisted of 28 adjacent SNPs of the BovineSNP50 Bead chip. In 100 randomly chosen fertile bulls of the Nordic Red cattle population, the haplotype had a frequency of 28%. Moreover, the haplotype occurred in the homozygous state in four bulls with normal fertility. Considering its high frequency and occurrence in the homozygous state in fertile bulls, it is unlikely that this haplotype was causal for the asthenospermia of two bulls. The IBD haplotype located on chromosome 25 consisted of 45 adjacent SNPs of the BovineSNP50 Bead chip (**Supplementary File 6**). It was not detected in the homozygous state in normospermic bulls. Both the sires and ten (out of 17) half-sibs of the asthenospermic bulls carried the haplotype in the heterozygous state. The average calving rate was 0.42 and 0.41 in half-sibs that were haplotype carriers and non-carriers, respectively, indicating that fertility is normal in heterozygous bulls. In 8,557 fertile sires from the Nordic Red cattle genetic evaluation reference population, the haplotype was observed in the heterozygous state in 426 animals corresponding to a frequency of 2.5%. Homozygous haplotype carriers were absent among the fertile bulls although five were expected assuming Hardy-Weinberg

Equilibrium proportions (P=0.018), corroborating recessive inheritance of a deleterious

allele.

Figure 2 – Homozygosity mapping in two asthenospermic bulls. Homozygosity mapping in
two asthenospermic bulls. Blue and green color represent autosomal runs of homozygosity (ROH)
that have been detected in the two asthenospermic bulls. The red frames highlight six ROH that
had been detected in both affected bulls. Only the 2.98-Mb segment located on chromosome 25
was never found in the homozygous state in normospermic bulls.
453
454

A variant disrupting a splice donor site in *CCDC189* is associated with

asthenospermia

In order to detect the causal mutation for the asthenospermia, we sequenced both affected

bulls at approximately 12-fold genome coverage. The sequencing depth in the interval

between 25,490,468 and 28,470,779 bp on chromosome 25, where both bulls shared an

extended segment of autozygosity, did not differ from the average sequencing coverage

(**Supplementary File 7**). Visual inspection of the aligned reads in the 2.98 Mb interval

revealed no evidence for a large deletion or structural variant that might segregate with the

haplotype. We identified 3,683 single nucleotide and short insertion and deletion

polymorphisms that were homozygous for the non-reference variant in both bulls in the

2.98 Mb IBD interval. Out of 3,683 sequence variants, 3,657 occurred in the homozygous

state in at least one out of 237 male control animals from breeds other than Nordic Red

that were not affected by the disorder. Such variants are not compatible with the presumed

recessive inheritance of a fully penetrant allele that causes asthenospermia, thus retaining

126 candidate causal variants. Variants annotation using the *Variant Effect Predictor* tool

revealed that 120 of 126 candidate causal variants were located in non-coding regions of

the genome (**Supplementary File 8**). Six variants were predicted to affect protein-coding

regions or splice sites including four synonymous mutations, a missense variant

(GCA_000003055.3:Chr25:g.26184191C>G) in *RABEP2*

(ENSBTAT00000008595:p.P413R, SIFT-score [39] 0.26) and a variant disrupting a

canonical 5' splice donor site (GCA_000003055.3:Chr25:g.27138357C>T) in *CCDC189*

(transcript-ID: ENSBTAT00000045037) encoding the coiled-coil domain containing protein

189. Of the 126 candidate causal variants, only the splice donor site mutation in *CCDC189*

was predicted to be deleterious to protein function.

The polymorphism Chr25:27138357C>T alters the first position of the fourth intron of

bovine *CCDC189*. The derived T-allele disrupts a canonical splice site (GT>AT; *CCDC189*

is transcribed from the reverse strand, **Figure 3**). The T-allele did not segregate in 237

healthy animals from cattle breeds other than Nordic Red that were sequenced in

Schwarzenbacher *et al.* [35]. In 2'669 animals that had been sequenced in the framework

of the 1000 Bull Genomes Project, the Chr25:27138357 T allele occurred in the

heterozygous state only in four animals of the Finnish Ayrshire and Swedish Red cattle

breeds (**Supplementary File 8**), i.e., two breeds that were used to form the Nordic Red

dairy cattle breed [38]. The T allele was never observed in the homozygous state in cattle

other than the two bulls with asthenospermia.

The effect of the Chr25:27138357C>T variant on *CCDC189* transcription was examined by

RT-PCR using RNA extracted from testicular tissues of the two asthenospermic and two

normospermic control bulls that were homozygous for the wild type allele. Using primers

located at the junction of exons 3 and 4 (forward) and in exon 6 (reverse) we obtained 358

and 368 bp RT-PCR products in asthenospermic (mt/mt) and normospermic (wt/wt) bulls,

respectively (**Figure 3D**). Sequence analysis of the RT-PCR product of mt/mt bulls

revealed that the Chr25:27138357C>T substitution activates a cryptic splice site in exon 4.

It alters the resulting CCDC189 protein sequence from amino acid residue 155 onwards

and creates a premature stop codon at amino acid residue 200, thus truncating 40% of the

332-amino acid CCDC189 protein. The truncated protein lacks domains that are supposed

to be essential for physiological protein function including the flagellar C1a complex

subunit C1a-32, which modulates ciliary and flagellar beating [40].

Prevalence of the splice donor variant in Nordic Red cattle

In order to confirm the association between the candidate causal splice site mutation in

CCDC189 and the asthenospermia-associated haplotype, we used Sanger sequencing to

obtain genotypes for the polymorphism in 118 normospermic control bulls of the Nordic

Red cattle breed. The variant was in perfect linkage disequilibrium with the 2.98 Mb

haplotype corroborating that it might be causative for the disorder. All bulls carrying the

associated haplotype in the heterozygous state were also heterozygous for the

Chr25:27138357 T allele. The T allele was not found in bulls that did not carry the

asthenospermia-associated haplotype.

Figure 3 – Schematic overview of bovine *CCDC189*. Structure of the bovine *CCDC189* gene
(transcript-ID: ENSBTAT00000045037). The bovine *CCDC189* gene encoding the coiled-coil
domain containing protein 189 is transcribed from the reverse strand. It consists of nine exons
(grey and dark-grey boxes) (A). A mutant allele (red triangle) at 27,138,357 basepairs disrupts a
canonical 5' splice donor variant in the fourth intron. Schematic view of the cDNA sequence of
CCDC189 (B). The coiled-coil domain containing protein 189 contains the flagellar C1a complex
subunit C1a-32 between amino acids 127 and 221 and a coiled coil domain between amino acids
280 and 308. Red arrows indicate the positions of the primers used for RT-PCR. The mutant allele
activates a cryptic splice site in exon 4 that truncates 10 basepairs from exon 4 (red colors) and
introduces a frameshift in translation that eventually creates a premature stop codon (C). The
mutant protein, if retained, lacks a large fraction of the flagellar C1a complex subunit C1a-32 and
the coiled coil domain. Effect of the mutant allele on CCDC189 transcription (D). The RT-PCR
products from testis tissue were shorter from two mt/mt (358 bp) than wt/wt (368 bp) bulls. Lane 5
(nc) is the negative PCR control. The length of the bands was defined using the Image Lab version
5.0 software (Bio-Rad Laboratories).
527

Discussion

A splice donor variant in bovine *CCDC189* is the likely cause of asthenospermia in the

Nordic Red cattle breed. An autosomal recessive mode of inheritance was suspected

because the two affected bulls were related through a common ancestor on both their

paternal and maternal paths, the spermatological findings were very similar in their

ejaculates, and their first- and second-degree relatives were fertile. Increasing rates of

inbreeding and the widespread use of individual sires in artificial insemination make cattle

populations susceptible to the propagation and phenotypic manifestation of recessive

alleles [41],[42]. Considering that only two bulls of the breed were affected by

asthenospermia, explanations other than recessive inheritance were also plausible

including, e.g., allelic heterogeneity [43],[44] and non-genetic causes of disease. However,

both bulls were housed at the AI center under controlled environmental conditions at the

same time as many fertile bulls and their health and andrological parameters were normal

indicating that non-genetic factors were less likely to cause this condition. Previous studies

showed that autozygosity mapping is a powerful approach to identify genomic regions

associated with recessive conditions even if only few affected individuals with genotype

information are available [13],[45]. Using autozygosity mapping in the two affected bulls,

we identified a 2.98 Mb segment of extended homozygosity located at chromosome 25

that was identical by descent in both bulls. This segment, encompassing a variant

disrupting a canonical 5' splice donor site in *CCDC189*, was never found in the

homozygous state in fertile bulls, in agreement with the presumed recessive inheritance of

an allele with detrimental phenotypic consequences.

Verifying the effect of candidate causal mutations in independent populations is critical

evidence to supporting variant causality [46],[35]. The splice donor variant in *CCDC189* did

not segregate in breeds distantly related to Nordic Red cattle indicating that it likely arose

after the formation of breeds thus precluding an immediate validation of the mutation in an

unrelated cattle population. Our study thus exemplifies difficulties inherent in validation and

functional investigation of deleterious sequence variants that manifest severe phenotypic

consequences such as impaired male fertility in livestock populations. Due to purifying

selection, deleterious sequence variants are rare in natural populations [46] unless they

either show pleiotropic effects on desired traits [43], are in linkage disequilibrium with

quantitative trait loci [47], or are propagated through the widespread use of heterozygous

mutation carriers in AI [48],[4]. The splice donor variant in *CCDC189* is rare in the Nordic

Red dairy cattle breed. We detected the derived allele in its homozygous state in only two

bulls. Assuming an allele frequency of the derived allele of 2.5% and random mating, only

1 out of 1,600 males is expected to be homozygous and thus sterile due to spermatozoa

with impaired motility. The disorder remains undetected in most homozygous animals because sterility becomes evident only in breeding bulls. Apart from producing asthenospermic ejaculates, both bulls homozygous for the derived T allele were healthy. However, genetic variants that cause asthenospermia may also compromise the function of motile cilia thus manifesting themselves in chronic detrimental diseases [11],[49],[35]. Further research is warranted to determine if DNA variants in *CCDC189* are associated with ciliopathies particularly in animals that are kept in conventional farms under normal husbandry conditions. A direct gene test will facilitate detection of and selection against carriers in order to prevent the birth of homozygous males that are sterile. Due to the low frequency of the deleterious allele, negative effects on genetic gain and genetic diversity are likely negligible when carrier animals are excluded from breeding in the Nordic Red cattle breed. However, sophisticated genome-based mating strategies are required to take into account the steadily increasing number of deleterious sequence variants discovered in breeding populations [50].

Functional and pathophysiological investigations were possible in two AI bulls only. Sperm analysis revealed an identical pattern in both bulls homozygous for the deleterious allele; sperm motility was impaired in both fresh ejaculates and post-thaw, e.g., the process of handling the semen, even with dilution of the seminal plasma did not ameliorate the low progressive motility. Slightly elevated ATP concentrations in post-thaw semen of the affected bulls might result from lower sperm metabolism or accumulation of ATP due to impaired progressive motility. However, elevated ATP concentrations might also derive from the inactive sperm tails. Moreover, a normal proportion of viable albeit asthenospermic spermatozoa resulted in high superoxide and reactive oxygen species production indicating defective mitochondria [51]. This further impairs sperm motility and eventually leads to lipid peroxidation and DNA damage and/or cell death, thus precluding fertilization [52]. Our study showed that the affected bulls were infertile *in vivo* and *in vitro*, precluding their use in AI. Fertilization might be possible *in vitro* using intracytoplasmic sperm injection, though DNA damage could impair embryo development. We have not explored this possibility, since the sperm phenotypic deviations were decisive to cull breeding sires from AI.

Most likely the causal variant for asthenospermia disrupts a canonical 5' splice donor site of bovine *CCDC189* encoding the coiled-coil domain containing protein 189. To the best of our knowledge, this is the first time that a phenotypic consequence has been associated

with a variant of *CCDC189* in any organism. CCDC189 (alias Spergen-4) is a sperm tail-associated protein that is mainly located in the principal piece and acrosome region of mature spermatozoa indicating that it may be required for their physiological activated progressive movement and acrosome reaction [53]. The velocity of spermatozoa was slower and their progressive motility was severely compromised in bulls that were homozygous for the Chr25:27138357 T allele indicating that flagellar movement was defective. Mutations disrupting components of the dynein arms or radial spokes of axonemal dyneins cause disorganized axonemes of motile cilia and flagella, thus compromising their physiological movement [11],[54],[55]. In spite of an impaired motility, the ultrastructural organization of the sperm flagella including the assembly of the dynein arms was normal in both asthenospermic bulls, indicating that disorganized axonemes were less likely to be responsible for the lack of progressive movement of the sperm. We showed that the Chr25:27138357 T allele introduces a cryptic splice site in the fourth exon of *CCDC189* leading to a frameshift in translation and premature translation termination. The resulting CCDC189 protein, if not degraded via nonsense-mediated mRNA decay, lacks a large part of the flagellar C1a complex subunit (C1a-32), that is an integral constituent of the central pair of flagellar microtubules. Aberrations of the small C1a microtubule projection of the central pair impair the physiological beating of flagella whose axonemes are normally organized otherwise [56],[57]. It has been shown that the C1a-32 subunit mediates the beating of cilia and flagella by modulating the activity of both the inner and outer dynein arms [40],[58]. Possibly, the resolution of the transmission electron microscopy was too low to reveal potential aberrations of the small C1a-32 subunit of the C1a projection [40]. Both bulls homozygous for the splice donor variant were sterile because they produced asthenospermic ejaculates corroborating that CCDC189 is important for physiological movement of spermatozoa and fertilization. However, spermatozoa of both asthenospermic bulls bended slightly which is not classified as progressive motility in automated semen analysis. The minor movement of affected spermatozoa resembles uncoordinated twitching of flagella in *Chlamydomonas* mutants lacking constituents of the flagellar C1a projection [57].

Conclusions

Our study in cattle revealed that a naturally occurring deleterious allele of *CCDC189* encoding the coiled-coil domain containing protein 189 causes immotile sperm with

630 defective mitochondria thus resulting in male sterility both *in vivo* and *in vitro*. The mutation

631 occurs in Nordic Red dairy cattle at a low frequency. Direct gene testing facilitates the

632 efficient monitoring of the deleterious allele in the population.

Acknowledgements

635 Genotype data of the control bulls were kindly provided by Viking Genetics. The authors

636 thank the Scientific Center for Optical and Electron Microscopy (ScopeM) of ETH Zurich,

637 particularly Dr. Cecilia Bebeacua for support in transmission electron microscopy.

Availability of data

640 Genotype data of two asthenospermic bulls are available in *plink* binary format as

641 supporting information. Whole-genome sequence data of both asthenospermic bulls have

642 been deposited at the European Nucleotide Archive (ENA) of EMBL-EBI under primary

643 accession PRJEB29487.

Author contributions

Sample collection: HV, MA; Analysis of genotyping data: TIT HV AS GS BG HP; Analysis

of whole-genome sequencing data: CW RF HP; RNA analysis: TIT AS; Sperm functional

analyses: AVC MAR SN HS HRM MA MH HV KFi KFI TS; In vitro fertilization experiment:

JP MM; In vivo fertilization experiment: JT; Transmission electron microscopy: MH; Wrote

the paper: TIT MA HRM HP; Conceived and designed the experiments: HV MA HP; Read

and approved the final version of the manuscript: all authors

References
653
1. Irvine DS. Epidemiology and aetiology of male infertility. Hum Reprod. 1998;13 Suppl
1:33–44.

2. Singh P, Schimenti JC. The genetics of human infertility by functional interrogation of
SNPs in mice. Proc Natl Acad Sci USA. 2015;112:10431–6.

3. Shorter JR, Odet F, Aylor DL, Pan W, Kao C-Y, Fu C-P, et al. Male Infertility Is
Responsible for Nearly Half of the Extinction Observed in the Mouse Collaborative Cross.
Genetics. 2017;206:557–72.

4. Pausch H, Kölle S, Wurmser C, Schwarzenbacher H, Emmerling R, Jansen S, et al. A
Nonsense Mutation in TMEM95 Encoding a Nondescript Transmembrane Protein Causes
Idiopathic Male Subfertility in Cattle. PLOS Genetics. 2014;10:e1004044.

5. Charlier C, Li W, Harland C, Littlejohn M, Coppieters W, Creagh F, et al. NGS-based
reverse genetic screen for common embryonic lethal mutations compromising fertility in
livestock. Genome Res. 2016;26:1333–41.

6. Silber SJ. The relationship of abnormal semen parameters to male fertility. Hum
Reprod. 1989;4:947–53.

7. Gredler B, Fuerst C, Fuerst-Waltl B, Schwarzenbacher H, Sölkner J. Genetic
parameters for semen production traits in Austrian dual-purpose Simmental bulls. Reprod
Domest Anim. 2007;42:326–8.

8. Ren D, Navarro B, Perez G, Jackson AC, Hsu S, Shi Q, et al. A sperm ion channel
required for sperm motility and male fertility. Nature. 2001;413:603–9.

9. Lim CC, Lewis SEM, Kennedy M, Donnelly ET, Thompson W. Human sperm
morphology and in vitro fertilization: sperm tail defects are prognostic for fertilization
failure. Andrologia. 30:43–7.

10. Ben Khelifa M, Coutton C, Zouari R, Karaouzène T, Rendu J, Bidart M, et al. Mutations
in DNAH1, which Encodes an Inner Arm Heavy Chain Dynein, Lead to Male Infertility from
Multiple Morphological Abnormalities of the Sperm Flagella. Am J Hum Genet.
2014;94:95–104.

11. Merveille A-C, Davis EE, Becker-Heck A, Legendre M, Amirav I, Bataille G, et al.
CCDC39 is required for assembly of inner dynein arms and the dynein regulatory complex
and for normal ciliary motility in humans and dogs. Nat Genet. 2011;43:72–8.

12. Sironen A, Thomsen B, Andersson M, Ahola V, Vilkki J. An intronic insertion in KPL2
results in aberrant splicing and causes the immotile short-tail sperm defect in the pig. Proc
Natl Acad Sci USA. 2006;103:5006–11.

13. Pausch H, Venhoranta H, Wurmser C, Hakala K, Iso-Touru T, Sironen A, et al. A
frameshift mutation in ARMC3 is associated with a tail stump sperm defect in Swedish Red
(Bos taurus) cattle. BMC Genetics. 2016;17:49.

14. Zhang Z, Kostetskii I, Tang W, Haig-Ladewig L, Sapiro R, Wei Z, et al. Deficiency of
SPAG16L causes male infertility associated with impaired sperm motility. Biol Reprod.
2006;74:751–9.

15. Pusch HH. The importance of sperm motility for the fertilization of human oocytes in
vivo and in vitro. Andrologia. 1987;19:514–27.

16. Donnelly ET, Lewis SE, McNally JA, Thompson W. In vitro fertilization and pregnancy
rates: the influence of sperm motility and morphology on IVF outcome. Fertil Steril.
1998;70:305–14.

17. Hellander JC, Samper JC, Crabo BG. Fertility of a stallion with low sperm motility and
a high incidence of an unusual sperm tail defect. Vet Rec. 1991;128:449–51.

18. de Kretser D. Male infertility. The Lancet. 1997;349:787–90.

19. Thibier M, Wagner H-G. World statistics for artificial insemination in cattle. Livestock
Production Science. 2002;74:203–12.

20. Taylor JF, Schnabel RD, Sutovsky P. Review: Genomics of bull fertility. animal.
2018;12:s172–83.

21. Ferenčaković M, Sölkner J, Kapš M, Curik I. Genome-wide mapping and estimation of
inbreeding depression of semen quality traits in a cattle population. J Dairy Sci.
2017;100:4721–30.

22. Rodríguez-Martínez H. State of the art in farm animal sperm evaluation. Reprod Fertil
Dev. 2007;19:91–101.

23. Patrick V, L US, Catherine D, Nadine B, Tom K, Patrick B. Bovine Semen Quality
Control in Artificial Insemination Centers. In: Bovine Reproduction. Wiley-Blackwell; 2014.
p. 685–95. doi:10.1002/9781118833971.ch74.

24. Tanghe S, Van Soom A, Sterckx V, Maes D, de Kruif A. Assessment of different sperm
quality parameters to predict in vitro fertility of bulls. Reprod Domest Anim. 2002;37:127–
32.

25. Christensen P, Stenvang JP, Godfrey WL. A flow cytometric method for rapid
determination of sperm concentration and viability in mammalian and avian semen. J
Androl. 2004;25:255–64.

26. Tejerina F, Buranaamnuay K, Saravia F, Wallgren M, Rodriguez-Martinez H.
Assessment of motility of ejaculated, liquid-stored boar spermatozoa using computerized
instruments. Theriogenology. 2008;69:1129–38.

27. Zimin AV, Delcher AL, Florea L, Kelley DR, Schatz MC, Puiu D, et al. A whole-genome
assembly of the domestic cow, Bos taurus. Genome Biology. 2009;10:R42.

28. Browning BL, Browning SR. A Unified Approach to Genotype Imputation and
Haplotype-Phase Inference for Large Data Sets of Trios and Unrelated Individuals. Am J
Hum Genet. 2009;84:210–23.

29. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. GigaScience. 2015;4:7.

30. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018;34:i884–90.

31. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv:13033997 [q-bio]. 2013. http://arxiv.org/abs/1303.3997. Accessed 25 May
2018.

32. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

33. DePristo MA, Banks E, Poplin RE, Garimella KV, Maguire JR, Hartl C, et al. A
framework for variation discovery and genotyping using next-generation DNA sequencing
data. Nat Genet. 2011;43:491–8.

34. Pedersen BS, Quinlan AR. Mosdepth: quick coverage calculation for genomes and
exomes. Bioinformatics. 2018;34:867–8.

35. Schwarzenbacher H, Burgstaller J, Seefried FR, Wurmser C, Hilbe M, Jung S, et al. A
missense mutation in TUBD1 is associated with high juvenile mortality in Braunvieh and
Fleckvieh cattle. BMC Genomics. 2016;17:400.

36. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The Ensembl
Variant Effect Predictor. Genome Biology. 2016;17:122.

37. Bouwman AC, Daetwyler HD, Chamberlain AJ, Ponce CH, Sargolzaei M, Schenkel
FS, et al. Meta-analysis of genome-wide association studies for cattle stature identifies
common genes that regulate body size in mammals. Nat Genetics. 2018;50:362–7.

38. Makgahlela ML, Mäntysaari EA, Strandén I, Koivula M, Nielsen US, Sillanpää MJ, et
al. Across breed multi-trait random regression genomic predictions in the Nordic Red dairy
cattle. J Anim Breed Genet. 130:10–9.

39. Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants
on protein function using the SIFT algorithm. Nature Protocols. 2009;4:1073–81.

40. Goduti DJ, Smith EF. Analyses of functional domains within the PF6 protein of the
central apparatus reveal a role for PF6 sub-complex members in regulating flagellar beat
frequency. Cytoskeleton (Hoboken). 2012;69:179–94.

41. Charlier C, Coppieters W, Rollin F, Desmecht D, Agerholm JS, Cambisano N, et al.
Highly effective SNP-based association mapping and management of recessive defects in
livestock. Nat Genet. 2008;40:449–54.

42. Pausch H, Schwarzenbacher H, Burgstaller J, Flisikowski K, Wurmser C, Jansen S, et
al. Homozygous haplotype deficiency reveals deleterious mutations compromising
reproductive and rearing success in cattle. BMC Genomics. 2015;16:312.

43. Sartelet A, Klingbeil P, Franklin CK, Fasquelle C, Géron S, Isacke CM, et al. Allelic
heterogeneity of Crooked Tail Syndrome: result of balancing selection? Anim Genet.
2012;43:604–7.

44. Medugorac I, Seichter D, Graf A, Russ I, Blum H, Göpel KH, et al. Bovine Polledness –
An Autosomal Dominant Trait with Allelic Heterogeneity. PLoS One. 2012;7.
doi:10.1371/journal.pone.0039477.

45. Pausch H, Ammermüller S, Wurmser C, Hamann H, Tetens J, Drögemüller C, et al. A
nonsense mutation in the COL7A1 gene causes epidermolysis bullosa in Vorderwald
cattle. BMC Genetics. 2016;17:149.

46. MacArthur DG, Manolio TA, Dimmock DP, Rehm HL, Shendure J, Abecasis GR, et al.
Guidelines for investigating causality of sequence variants in human disease. Nature.
2014;508:469–76.

47. Fujii J, Otsu K, Zorzato F, de Leon S, Khanna VK, Weiler JE, et al. Identification of a
mutation in porcine ryanodine receptor associated with malignant hyperthermia. Science.
1991;253:448–51.

48. Sartelet A, Druet T, Michaux C, Fasquelle C, Géron S, Tamma N, et al. A Splice Site
Variant in the Bovine RNF11 Gene Compromises Growth and Regulation of the
Inflammatory Response. PLOS Genetics. 2012;8:e1002581.

49. Zariwala MA, Knowles MR, Omran H. Genetic defects in ciliary structure and function.
Annu Rev Physiol. 2007;69:423–50.

50. Cole JB. A simple strategy for managing many recessive disorders in a dairy cattle
breeding program. Genet Sel Evol. 2015;47. doi:10.1186/s12711-015-0174-9.

51. Rodriguez-Martinez H. Semen evaluation and handling: emerging techniques and
future development. In: Chenoweth P, Lorton S, editors. Animal andrology: theories and
applications. Wallingford: CABI; 2014. p. 509–49. doi:10.1079/9781780643168.0509.

52. Koppers AJ, De Iuliis GN, Finnie JM, McLaughlin EA, Aitken RJ. Significance of
mitochondrial reactive oxygen species in the generation of oxidative stress in
spermatozoa. J Clin Endocrinol Metab. 2008;93:3199–207.

53. Howida A, Salaheldeen E, Iida H. Molecular Cloning of Spergen-4, Encoding a
Spermatogenic Cell-Specific Protein Associated with Sperm Flagella and the Acrosome
Region in Rat Spermatozoa. Zoological Science. 2016;33:195–203.

54. Alsaadi MM, Erzurumluoglu AM, Rodriguez S, Guthrie PAI, Gaunt TR, Omar HZ, et al.
Nonsense mutation in coiled-coil domain containing 151 gene (CCDC151) causes primary
ciliary dyskinesia. Hum Mutat. 2014;35:1446–8.

55. Bower R, Tritschler D, Mills KV, Heuser T, Nicastro D, Porter ME. DRC2/CCDC65 is a
central hub for assembly of the nexin-dynein regulatory complex and other regulators of
ciliary and flagellar motility. Mol Biol Cell. 2018;29:137–53.

56. Rupp G, O'Toole E, Porter ME. The Chlamydomonas PF6 locus encodes a large
alanine/proline-rich polypeptide that is required for assembly of a central pair projection
and regulates flagellar motility. Mol Biol Cell. 2001;12:739–51.

57. Wargo MJ, Dymek EE, Smith EF. Calmodulin and PF6 are components of a complex
that localizes to the C1 microtubule of the flagellar central apparatus. J Cell Sci. 2005;118
Pt 20:4655–65.

58. Nakazawa Y, Ariyoshi T, Noga A, Kamiya R, Hirono M. Space-Dependent Formation
of Central Pair Microtubules and Their Interactions with Radial Spokes. PLoS One. 2014;9.
doi:10.1371/journal.pone.0110513.

Supplementary Information

Supplementary File 1 – Genotypes of two asthenospermic bulls

Format: .zip-archive

Genotypes at 41,594 SNPs for two asthenospermic bulls in *plink* binary format. The positions of the SNP correspond to the UMD3.1 assembly of the bovine genome.

Supplementary File 2 – R script to identify IBD segments in two asthenospermic bulls

Format: .R-file

R script to visualize runs of homozygosity in the genotype data of two asthenospermic bulls. This program uses as input the files provided as Supplementary File 1. Please note that this R script calls the fast algorithm implemented in *plink* (version 1.9) to identify runs of homozygosity.

Supplementary File 3 – Post-thaw sperm motility in an asthenospermic bull

Format: .mov

Video of frozen-thawed spermatozoa of an asthenospermic bull. Please note the minor movement of some spermatozoa that is typically not classified as progressive motility during the subjective eye-evaluation of ejaculates.

Supplementary File 4 – Post-thaw sperm motility in a normospermic control bull

Format: .mov

Video of frozen-thawed spermatozoa of an normospermic bull.

Supplementary File 5 – Pedigree of two asthenospermic bulls

Format: .png

Pedigree of two affected bulls. Ovals and boxes represent female and male ancestors, respectively. The grey box highlights the most recent common ancestor (MRCA) that was present on both the maternal and paternal ancestry of both asthenospermic bulls.

Supplementary File 6 – Asthenospermia-associated haplotype

Format: .csv

ID, position and asthenospermia-associated allele of 45 adjacent SNPs of the

BovineSNP50 Bead chip. The coding of the alleles is based on the Illumina TOP/BOT

format. The chromosomal position of the SNP corresponds to the UMD3.1 assembly of the

bovine genome.

Supplementary File 7 – Sequencing depth along chromosome 25 in two

asthenospermic bulls

Format: .png

The depth of sequencing coverage in two asthenospermic bulls was calculated at bovine

chromosome 25 in sliding windows of 5,000 base pairs using the aligned reads. Grey

background indicates the segment of extended homozygosity. The red line represents the

position of the candidate causal variant-

Supplementary File 8 – Candidate causal variants compatible with recessive

inheritance of asthenospermia

Format: .txt

Functional consequences of 126 candidate causal variants that were compatible with

recessive inheritance of asthenospermia, i.e., they were homozygous for the alternate

allele in two asthenospermic bulls and either heterozygous or homozygous for the

reference allele in 237 healthy control animals. Variants were annotated using the *Variant*

Effect Predictor tool. The quality of the sequence variant genotypes for both

asthenospermic bulls is shown in columns 7 and 8. The number of heterozygous control

animals (out of 237) is shown in column 9. The last column indicates the genotype

distribution of 126 candidate causal variants (homozygous for the reference allele |

heterozygous | homozygous for the alternate allele) in 2,669 animals of the 1000 bull

genomes dataset.