1 **Tryptophan fluorescence quenching in β-lactam-interacting proteins is modulated by**
2 **the structure of intermediates and final products of the acylation reaction**
3
4

5 Sébastien Triboulet[a1], Zainab Edoo[a1], Fabrice Compain[a,b], Clément Ourghanlian[a], Adrian Dupuis[a],
6 Vincent Dubée[a], Laetitia Sutterlin[a], Heiner Atze[a], Mélanie Ethève-Quelquejeu[c], Jean-Emmanuel
7 Hugonnet[a2], and Michel Arthur[a2]
8
9
10 [1]Equal contribution
11 [2]Equal contribution and corresponding authors
12
13
14 [a] Dr. S Triboulet, Ms Z Edoo, Dr. F Compain, Dr. C Ourghanlian, Mr A Dupuis, Dr. V Dubée, Dr. L
15   Sutterlin, Dr. H Atze, Dr. JE Hugonnet, Dr. M Arthur, Sorbonne Université, Sorbonne Paris Cité,
16   Université Paris Descartes, Université Paris Diderot, INSERM, Centre de Recherche des Cordeliers,
17   CRC,   F-75006   Paris,   France.   E-mail: jean-emmanuel.hugonnet@crc.jussieu.fr;
18   michel.arthur@crc.jussieu.fr
19
20 [b] Dr. F Compain, Service de Microbiologie, Assistance Publique-Hôpitaux de Paris, Hôpital Européen
21   Georges Pompidou, Paris, France.
22
23 [c] Dr. M Ethève-Quelquejeu, Laboratoire de Chimie et de Biochimie Pharmacologiques et
24   Toxicologiques, Université Paris Descartes, UMR 8601, Paris, F-75005 France; CNRS UMR 8601,
25   Paris, F-75006 France.
26
27

28 Key words: acylenzyme, β-lactamase, fluorescence quenching, L,D-transpeptidase, tryptophan
29 fluorescence.
30

In most bacteria, β-lactam antibiotics inhibit the last cross-linking step of peptidoglycan synthesis by
acylation of the active-site Ser of D,D-transpeptidases belonging to the penicillin-binding protein
(PBP) family. In mycobacteria, cross-linking is mainly ensured by L,D-transpeptidases (LDTs), which
are promising targets for the development of β-lactam-based therapies for multidrug-resistant
tuberculosis. For this purpose, fluorescence spectroscopy is used to investigate the efficacy of LDT
inactivation by β-lactams but the basis for fluorescence quenching during enzyme acylation remains
unknown. In contrast to what has been reported for PBPs, we show here using a model L,D-
transpeptidase (Ldt$_{fm}$) that fluorescence quenching of Trp residues does not depend upon direct
hydrophobic interaction between Trp residues and β-lactams. Rather, Trp fluorescence was
quenched by the drug covalently bound to the active-site Cys residue of Ldt$_{fm}$. Fluorescence
quenching was not quantitatively determined by the size of the drug and was not specific of the
thioester link connecting the β-lactam carbonyl to the catalytic Cys as quenching was also observed
for acylation of the active-site Ser of β-lactamase BlaC from *M. tuberculosis*. Fluorescence quenching
was extensive for reaction intermediates containing an amine anion and for acylenzymes containing
an imine stabilized by mesomeric effect, but not for acylenzymes containing a protonated β-lactam
nitrogen. Together, these results indicate that the extent of fluorescence quenching is determined
by the status of the β-lactam nitrogen. Thus, fluorescence kinetics can provide information not only
on the efficacy of enzyme inactivation but also on the structure of the covalent adducts responsible
for enzyme inactivation.

50
51 **INTRODUCTION**
52
Peptidoglycan is an essential constituent of bacterial cell walls since it prevents cell swelling and lysis
by mechanically sustaining the osmotic pressure of the cytoplasm (1). This protective function depends
upon synthesis and maintenance during the entire cell cycle of the net-like peptidoglycan
macromolecule, which completely surrounds the bacterial cell. Peptidoglycan is made of glycan strands
cross-linked by short peptide stems. In most bacteria, the cross-linking step is performed by D,D-
transpeptidases, which are the essential targets of β-lactam antibiotics and are often referred to as
penicillin-binding proteins (PBPs) (2). In mycobacteria (3, 4) and in *Clostridium difficile* (5, 6), the cross-
links found are mainly (70% to 80%) formed by a second class of enzymes, the L,D-transpeptidases
(LDTs). Since LDTs are not inhibited by β-lactams belonging to the penam class, such as ampicillin, (7,
8), these enzymes are responsible for high-level resistance to these drugs in mutants of *Enterococcus*
faecium (9, 10) and *Escherichia coli* (11) selected in laboratory conditions.
PBPs and LDTs are structurally unrelated (12-15) and proceed through different catalytic
mechanisms for activation of Ser and Cys nucleophiles (16), which are part of Lys-Ser (17) and Cys-His-
Asp (18) catalytic diad and triad, respectively. PBPs and LDTs also differ by the structure of the stem
peptide used as an acyl donor, a pentapeptide for PBPs (17) and a tetrapeptide for LDTs (16), except
for *Enterococcus faecalis* Ldt$_{fs}$ (19). PBPs cleave the D-Ala^4-D-Ala^5 peptide bond of the acyl donor, hence
the D,D designation for these transpeptidases, and link the carbonyl of D-Ala^4 to the side-chain amino
group of the 3rd residue thereby generating 4→3 cross-links (17) (Supplementary Fig. S1). In contrast,
LDTs cleave the L-Lys^3-D-Ala^4 bond of the donor (L,D designation) and form 3→3 cross-links (16).
The first substrate of the transpeptidation reaction, the acyl donor, forms an acylenzyme
intermediate that subsequently reacts with the second substrate, the acceptor, leading to the final
cross-linked product. For PBPs, nucleophilic attack of the carbonyl of D-Ala^4 by the active-site Ser
results in an acylenzyme containing an ester bond and release of D-Ala^5 (17). For the acylenzyme

formed by LDTs, a thioester bond connects the carbonyl of L-Lys[3] to the γ sulfur of the active-site Cys
(16). β-lactams are structure analogues of the D-Ala[4]-D-Ala[5] extremity of stem pentapeptides and act
as suicide substrates of the PBPs (20). Nucleophilic attack of the carbonyl of the β-lactam ring by the
active-site Ser of PBPs leads to formation of an acylenzyme, which is only hydrolyzed very slowly,
leading in practice to "irreversible" inactivation in the time scale of a bacterial generation
(Supplementary Fig. S1) (2). LDTs are also acylated by β-lactams although efficacious enzyme
inactivation and antibacterial activity occur only for a single class of β-lactams, the carbapenems, such
as imipenem (7, 8). β-lactams of the cephem (cephalosporin) class, such as ceftriaxone, are only active
at high concentrations since acylation is slower and the resulting thioester bond is prone to hydrolysis
(7). The kinetic parameters are even less favorable for penams (ampicillin and penicillin) leading to
equilibrium between the acylated and unacylated (functional) forms, which accounts for the lack of
antibacterial activity (7).

Fluorescence kinetics revealed that the acylation reaction of LDTs by β-lactams of the
carbapenem class comprises two limiting steps (Fig. 1A and B). In the first step, nucleophilic attack of
the β-lactam carbonyl by the negatively charged sulfur atom of the catalytic Cys was thought to lead
to formation of a covalent sulfur-carbon bond and to the development of a negative charge on the
drug (21). Initially, this negative charge was proposed to be located on the β-lactam oxygen (21). This
hypothesis was based on analogies with PBPs, which contain an oxyanion hole for stabilization of the
negative charge developing on the β-lactam oxygen (17). However, hybrid potential simulation of the
acylation reaction indicated that the path to an oxyanion is energetically disfavored in the case of LDTs
and that an oxyanion could not correspond to a stabilized reaction intermediate (22). Rather, hybrid
potential simulation indicated that the most energetically favored reaction path involves a concerted
mechanism leading to the concomitant formation of the thioester bond, rupture of C-N β-lactam bond,
and formation of an amine anion (22). In the second step, the final acylenzyme is formed by
protonation of the amine anion. The origin of this proton remains elusive (18, 22). Formation of a non-
covalent complex is not considered in this reaction scheme as the affinity of the Ldt$_{fm}$ for carbapenems
is very low, in the order of 60 to 80 mM (7).

Fluorescence kinetics proved useful to assess the efficacy of inactivation of LDTs from
Mycobacterium tuberculosis (23-25) and contributed to the choice of the drugs that were tested in a
phase II clinical trial (26). The assays were also useful in evaluating series of synthetic carbapenems
(27) and non-β-lactam LDT inhibitors (28). However, the basis for the variation in fluorescence intensity
detected during the two steps of the acylation reaction remained elusive. Initially, we developed
fluorescence kinetics since determination of the crystal structure of Ldt$_{fm}$ revealed a Trp residue at the
entrance of the catalytic cavity (15), which appeared ideally located for fluorescence quenching upon
drug binding due to changes in its environment (21). However, this naïve explanation was subsequently
found to be insufficient to account for the various behaviors observed for acylation of LDTs by
representatives of the β-lactam classes (7, 22-25). This prompted us to investigate here the basis for
variations in fluorescence intensity occurring upon acylation of LDTs using the well-characterized L,D-
transpeptidase Ldt$_{fm}$ from *E. faecium* as a model. We show that variations in fluorescence intensity are
not due to modification of the environment of Trp residues but to the formation of a β-lactam-derived
fluorescence quencher during the acylation reaction.

RESULTS

Trp to Phe and His to Ala substitutions do not abolish the biphasic behavior of fluorescence kinetics observed for Ldt_{fm} inactivation by imipenem. We have previously proposed that the acylation of Ldt_{fm} by carbapenems is a two-step reaction involving reversible formation of an amine anion (EI^{An}, step 1) followed by irreversible formation of an acylenzyme (EI*, step 2) (Fig. 1A) (21, 22). Fluorescence kinetics also display two phases enabling the determination of kinetic parameters for the two steps of the reaction (Fig. 1B) (21). In the first phase, fluorescence intensity decreases because EI^{An} is rapidly formed to the detriment of the free enzyme (E) and the fluorescence intensity of EI^{An} is lower than that of E. In the second phase, fluorescence intensity increases since the acylenzyme (EI*) is formed to the detriment of EI^{An} (acylation step) and the fluorescence intensity of EI* is greater than that of EI^{An}. Our first aim was to determine whether variations in the fluorescence intensity could be correlated to the modification of the environment of a specific Trp residue of Ldt_{fm}. To address this question, Trp residues of the catalytic domain of Ldt_{fm} were replaced by Phe residues in various combinations. For the sake of simplicity, the six Trp residues at positions 355, 385, 410, 415, 425, and 434 of Ldt_{fm} were designated as residues a to f, respectively. Fluorescence kinetics were determined for Ldt_{fm} derivatives lacking 2 to 5 Trp residues (Fig. 1C). Fluorescence kinetics remained bi-phasic in all cases, except for the Ldt_{fm} derivative only retaining the "a" residue (*i.e.* Trp^{355}). Thus, the variations in the fluorescence intensity associated with the two steps of the acylation reaction do not depend upon the modification of the environment of a specific Trp residue of Ldt_{fm}. This prompted us to investigate fluorescence quenching by His residues of Ldt_{fm}. Fluorescence kinetics obtained with the His^{353}Ala and His^{440}Ala derivatives of Ldt_{fm} indicated that neither residue is essential for biphasic fluorescence kinetics (Fig. 1C). His at position 421 could not be analyzed since this residue is part of the catalytic triad and is essential for Ldt_{fm} activity.

Impact of Trp to Phe and His to Ala substitutions on the efficacy of the inactivation reaction and on the relative fluorescence intensity of the three enzyme forms. Fitting simulations to experimental data for four concentrations of imipenem was used to determine the inactivation kinetic parameters (Supplementary Fig. S2). None of the Trp residues was essential for the acylation of Ldt_{fm} by imipenem although important variations were observed in the k_1 (formation of the amine anion) and k_2 (acylation step) parameters (Fig. 2A and 2B, respectively). The Trp to Phe substitutions produced large variations in the intrinsic fluorescence of the three forms of the enzyme (apo Ldt_{fm}, amine anion, and acylenzyme; Fig. 2C). The formation of the acylenzyme (EI*) was confirmed by mass spectrometry for all Ldt_{fm} derivatives (mass increment of 299.3 Da; data not shown).

Conservation of Trp residues in Ldt_{fm} homologues. The Trp residues were poorly conserved in L,D-transpeptidases from Gram-positive and Gram-negative bacteria (Supplementary Fig. S3), as expected from the substantial high residual acylation activity of Ldt_{fm} derivatives containing Trp to Phe substitutions (above, Fig. 2A and B). However, residue e (Trp^{425}) was relatively conserved in L,D-transpeptidases from *M. tuberculosis*, being present in the Mt1, Mt2, Mt3, and Mt4 enzymes as well as in the corresponding orthologues of *M. abscessus*. These enzymes are known to display bi-phasic fluorescence kinetics upon inactivation by carbapenems (23, 24, 29) and (unpublished results). Since Trp^{425} was the only conserved Trp residue in the LDTs from mycobacteria we investigated the impact of a Trp to Phe substitution at this position in Ldt_{fm}. As shown in Supplementary Fig. S4A and B, derivatives of Ldt_{fm} lacking all Trp residues except Trp^{425} or lacking only this residue displayed biphasic kinetics. These results indicated that the biphasic nature of fluorescence kinetics does not depend upon any conserved Trp residue in the L,D-transpeptidase protein family.

Fluorescence kinetics with a single ectopic Trp residue at position 383. All six Trp residues of Ldt$_{fm}$
were replaced by Phe residues. As expected, the variant devoid of Trp residues was very weakly
fluorescent (data not shown). No modification of this weak fluorescence was observed upon acylation
by imipenem. This observation ruled out the formal possibility that the catalytic Cys acylated by
imipenem could be a fluorophore. Replacement of Tyr at position 383 by a Trp residue in this Ldt$_{fm}$
derivative restored biphasic fluorescence kinetics (Supplementary Fig. S4C). This result confirmed that
biphasic fluorescence kinetics are not dependent upon the presence of any of the six Trp residues of
Ldt$_{fm}$ in their original positions.

Investigation of the opened β-lactam ring linked to the catalytic Cys residue as a quencher. Since the
variations in the fluorescence observed during the two steps of the acylation reaction could not be
accounted for by modification of the environment of any specific Trp residue, our next objective was
to investigate whether quenching could result from energy transfer from Trp residues to the Cys-β-
lactam covalent adduct generated upon enzyme inactivation. To investigate this possibility, we first
examined acylation of Ldt$_{fm}$ by the chromogenic cephalosporin nitrocefin. Fluorescence kinetics with
this compound displayed a monophasic behavior with a large decrease (72%) in the fluorescence
intensity (Fig. 3A). In the Ldt$_{fm}$-nitrocefin acylenzyme, the negative charge developing on the β-lactam
nitrogen is stabilized by a mesomeric effect that prevents protonation of the amine (30). Thus, the
amine anion formed as an end product with nitrocefin (Fig. 3A) or as an intermediate with imipenem
(Fig. 1) was associated with extensive fluorescence quenching. As found for nitrocefin, acylation of
Ldt$_{fm}$ by ceftriaxone led to a monophasic decrease in the fluorescence intensity (Fig. 3B). Acylation of
Ldt$_{fm}$ by ceftriaxone involves a concerted mechanism leading to the elimination of the side chain and
formation of a conjugated imine (7). The extent of fluorescence quenching observed for the
acylenzyme formed with ceftriaxone (40%; imine) or by the intermediate formed with imipenem (45%;
amine anion) were more important than that observed for the final Ldt$_{fm}$-imipenem adduct (20%)
suggesting that quenching was decreased by protonation of the β-lactam nitrogen.

Fluorescence quenching associated with inactivation of β-lactamases by β-lactamase inhibitors. In
order to enrich the structural diversity of molecules investigated in the fluorescence quenching assay,
we aimed to test β-lactamase inhibitors. Most β-lactams could not be tested with β-lactamases since
the fluorescence assay requires relatively high enzyme concentrations (typically 10 µM) leading to
rapid hydrolysis of the drugs even if the turnovers are low. For this reason, we focused on inactivation
of BlaC from *Mycobacterium tuberculosis* by clavulanate since the corresponding acylenzyme is not
prone to hydrolysis at the timescale of our experiments (31). Acylation of BlaC by clavulanate led to a
large decrease in the fluorescence intensity (37%) (Fig. 3C). Previous studies have shown that acylation
of BlaC by clavulanate leads to formation of imine and *trans*-enamine tautomers (depicted in Fig. 3C)
following drug decarboxylation (32). Since Ldt$_{fm}$ and BlaC contain Cys and Ser as the active-site
nucleophile, respectively, these results indicate that fluorescence quenching is observed with
acylenzymes containing both thioester and ester bonds.

Avibactam offered the possibility to test a non-β-lactam inhibitor that inhibits BlaC by formation
of a carbamoylenzyme (33). Formation of the adduct (Supplementary Fig. S5) was not associated with
any significant modification of the fluorescence intensity of the enzyme (Fig. 3D). Protonation of the
amino sulfate nitrogen of avibactam in the avibactam-BlaC adduct was thus associated with limited
quenching as found for the protonation of the β-lactam nitrogen in the final Ldt$_{fm}$-imipenem and BlaC-
clavulanate adducts (Fig. 1 and 3C).

Acylation of Ldt$_{fm}$ and BlaC by faropenem. Faropenem, a β-lactam belonging to the penem class, is poorly hydrolyzed by BlaC and rapidly inactivates L,D-transpeptidases (24). Thus, faropenem provided the opportunity to study the acylation of the two enzyme types, *i.e.* a β-lactamase (BlaC) and an L,D-transpeptidase (Ldt$_{fm}$), by the same β-lactam. Acylation of BlaC by faropenem led to extensive fluorescence quenching (44%) (Fig. 3E). The resulting acylenzyme contained an unprotonated β-lactam nitrogen. In contrast, acylation of Ldt$_{fm}$ by faropenem, which resulted in the rupture of the C^5-C^6 bond and the loss of a large portion of the drug including the β-lactam nitrogen (24), was associated with a moderate decrease in the fluorescence intensity (17%) (Fig. 3F).

Acylation of BlaC by the meropenem carbapenem. Although BlaC displays moderate carbapenemase activity, previous studies have shown that a BlaC-meropenem adduct is the predominant enzyme form upon incubation of the enzyme with low drug concentrations (34, 35). As expected, the control experiment depicted in Supplementary Fig. S6 shows that the concentration of meropenem remains saturating for at least 80 min during hydrolysis of meropenem (40 – 200 μM) by BlaC (10 μM). Fluorescence kinetics at a shorter time scale (6 min) showed a transitory decrease in fluorescence intensity (40%) (Fig. 3G). This observation suggests that formation of an amine anion initially leads to an important fluorescence quenching. Then, fluorescence returned to the initial level suggesting that protonation of the amine anion fully suppressed quenching. Thus, formation of an amine anion following nucleophilic attack of the carbonyl carbon of carbapenems by Ldt$_{fm}$ (Fig. 1) and BlaC (Fig. 3G) similarly led to extensive fluorescence quenching despite the presence of different catalytic residues (Cys *versus* Ser) and different modes of activation of these nucleophiles involving His and Lys residues.

DISCUSSION

The main aim of the present study was to determine the basis for the variations in the fluorescence intensity observed upon acylation of LDTs by β-lactam antibiotics. Previous analyses of the D,D-transpeptidase R61 (PBP family) have shown that acylation of the enzyme by penicillin G leads to a fluorescence quenching, which specifically depends upon one of two Trp residues of the protein (36, 37). This residue, W^{233}, is essential for the D,D-carboxypeptidase activity of the protein, as determined by the release of the terminal D-Ala residue from a model peptide ending in D-Ala-D-Ala (37). The crystal structure of R61 in complex with a peptide mimicking the acyl donor revealed hydrophobic interactions between the side-chain of W^{233} and methylene groups of this substrate (38). Likewise, the structure of a penicillin-R61 complex revealed that W^{233} is in hydrophobic interaction with the phenylacetyl side chain of the drug (39) indicating that fluorescence quenching may depend upon this direct interaction. In contrast, we show here that the variations in fluorescence during acylation of LDTs by β-lactam antibiotics require at least one Trp residue but that the position of this residue is not determinant. This was established by deleting Trp residues from Ldt$_{fm}$ in various combinations (Fig. 1 and 2), by introducing a single Trp residue in an ectopic position of Ldt$_{fm}$ (Fig. S4), and by showing that large variations in fluorescence intensity occur in distantly related LDTs and unrelated β-lactamases that do not share any conserved Trp residue (Fig. S3). Since fluorescence quenching was not mainly due to changes in the environment of Trp residues, we investigated the possibility that the nucleophilic attack of the β-lactam ring could itself generate a fluorescence quencher. If this is the case, the fluorescence profile should depend upon the structural features of the drug bound to the active-site residue. Accordingly, extensive quenching was not mainly determined by the size of the side chains of β-lactams or by the nature of the enzyme nucleophile (Cys *versus* Ser in Ldt$_{fm}$ and BlaC, respectively) (Fig .3).

Acylation of Ser or Cys *per se* was not sufficient for fluorescence quenching since a modest or no decrease in fluorescence intensity was observed for the final acylation products of Ldt$_{fm}$ by faropenem, of BlaC by avibactam, and of both enzymes by carbapenems. By elimination, these results pointed to the status of the β-lactam nitrogen as the only source of variation that could account for the main variations in fluorescence intensity. A large fluorescence quenching occurred if the β-lactam nitrogen was negatively charged, as transiently found during acylation of Ldt$_{fm}$ and BlaC by carbapenems. A large fluorescence quenching also occurred if the nitrogen atom was engaged in a double bond as observed for the acylenzymes formed by Ldt$_{fm}$ with ceftriaxone or nitrocefin and by BlaC with faropenem or clavulanate. In agreement with the critical role of the double bond, fluorescence quenching was not observed for acylation of BlaC by avibactam, which results in the formation of a carbamoyl-enzyme with a protonated nitrogen, or for acylation of Ldt$_{fm}$ by faropenem, which results in the formation of an acylenzyme lacking the nitrogen atom following rupture of the C^5-C^6 bond. These results indicate that fluorescence kinetics can provide information not only on the efficacy of enzyme inactivation but also on the structure of the covalent adducts, including for example the presence of Δ1- or Δ2-pyrroline ring in enzyme-carbapenem adducts. Our results also indicate that nitrocefin provides a very sensitive assay to titrate the active site of enzymes that do not hydrolyze this cephalosporin. This property could be exploited to identify β-lactams and non-β-lactam inhibitors that reversibly or irreversibly bind to LDTs.

MATERIALS AND METHODS

Enzyme production and purification. The catalytic domain of Ldt$_{fm}$ (residues 341 to 466) and soluble BlaC (residues 39 to 306) were produced in *E. coli* and purified by metal affinity and size exclusion chromatography as previously described (30). Synthetic genes were purchased for production of Ldt$_{fm}$ derivatives with Trp to Phe and His to Ala substitutions (GeneCust). All experiments were performed with a fixed concentration of Ldt$_{fm}$ and BlaC (10 μM).

Spectrofluorimetry. Fluorescence kinetics were studied in 100 mM sodium phosphate (pH 6.0) at 20°C by using a stopped-flow apparatus (RX-2000, Applied Biophysics) coupled to a spectrofluorimeter (Cary Eclipse; Varian). Excitation was performed at 224 nm with a 5 nm slit and a 2 mm optical path. Fluorescence emission was determined at 335 nm with a 5 nm slit and a 10 mm optical path. Fluorescence data were collected exactly in the same conditions for comparison of the relative fluorescence of wild-type Ldt$_{fm}$ and derivatives with Trp to Phe and His to Ala substitutions. These conditions included the photomultiplier voltage, which was set to 600 V.

Mass spectrometry. Formation of the proposed Ldt$_{fm}$-β-lactam and BlaC-β-lactam adducts in the conditions described in the text was systematically checked by mass spectrometry (data not shown). Enzymes and β-lactams were incubated for appropriate time periods at 20°C. Five μl of acetonitrile and 1 μl of 1% formic acid were extemporaneously added. Injection in the mass spectrometer (Qstar Pulsar I; Applied Biosystem) was performed at a flow rate of 0.05 ml/min (acetonitrile, 50%, water, 49.5%, and formic acid, 0.5%; per volume). Spectra were acquired in the positive mode, as previously described (8). Detailed analyses of the mass spectra of the β-lactam adducts considered in the current study have been previously published (7, 8, 24, 29, 40).

ACKNOWLEDGEMENTS

We thank L. Dubost and A. Marie for technical assistance in the collection of mass spectra. This work was supported by the French National Research Agency (grant MycWall, ANR-17-CE18-0010) and the Fondation pour la Recherche Médicale (grant ECO20160736080 to ZE).

REFERENCES

1. **Turner RD, Vollmer W, Foster SJ.** 2014. Different walls for rods and balls: the diversity of peptidoglycan. Mol Microbiol **91:**862-874.

2. **Zapun A, Contreras-Martel C, Vernet T.** 2008. Penicillin-binding proteins and beta-lactam resistance. FEMS Microbiol Rev **32:**361-385.

3. **Lavollay M, Arthur M, Fourgeaud M, Dubost L, Marie A, Veziris N, Blanot D, Gutmann L, Mainardi JL.** 2008. The peptidoglycan of stationary-phase *Mycobacterium tuberculosis* predominantly contains cross-links generated by L,D-transpeptidation. J Bacteriol **190:**4360-4366.

4. **Lavollay M, Fourgeaud M, Herrmann JL, Dubost L, Marie A, Gutmann L, Arthur M, Mainardi JL.** 2011. The peptidoglycan of *Mycobacterium abscessus* is predominantly cross-linked by L,D-transpeptidases. J Bacteriol **193:**778-782.

5. **Peltier J, Courtin P, El Meouche I, Lemee L, Chapot-Chartier MP, Pons JL.** 2011. *Clostridium difficile* has an original peptidoglycan structure with a high level of *N*-acetylglucosamine deacetylation and mainly 3-3 cross-links. J Biol Chem **286:**29053-29062.

6. **Sutterlin L, Edoo Z, Hugonnet JE, Mainardi JL, Arthur M.** 2018. Peptidoglycan cross-linking activity of L,D-transpeptidases from *Clostridium difficile* and inactivation of these enzymes by beta-lactams. Antimicrob Agents Chemother **62**.

7. **Triboulet S, Dubee V, Lecoq L, Bougault C, Mainardi JL, Rice LB, Etheve-Quelquejeu M, Gutmann L, Marie A, Dubost L, Hugonnet JE, Simorre JP, Arthur M.** 2013. Kinetic features of L,D-transpeptidase inactivation critical for beta-lactam antibacterial activity. PLoS One **8:**e67831.

8. **Mainardi JL, Hugonnet JE, Rusconi F, Fourgeaud M, Dubost L, Moumi AN, Delfosse V, Mayer C, Gutmann L, Rice LB, Arthur M.** 2007. Unexpected inhibition of peptidoglycan LD-transpeptidase from *Enterococcus faecium* by the beta-lactam imipenem. J Biol Chem **282:**30414-30422.

9. **Mainardi JL, Legrand R, Arthur M, Schoot B, van Heijenoort J, Gutmann L.** 2000. Novel mechanism of beta-lactam resistance due to bypass of DD-transpeptidation in *Enterococcus faecium*. J Biol Chem **275:**16490-16496.

10. **Mainardi JL, Morel V, Fourgeaud M, Cremniter J, Blanot D, Legrand R, Frehel C, Arthur M, Van Heijenoort J, Gutmann L.** 2002. Balance between two transpeptidation mechanisms determines the expression of beta-lactam resistance in *Enterococcus faecium*. J Biol Chem **277:**35801-35807.

11. **Hugonnet JE, Mengin-Lecreulx D, Monton A, den Blaauwen T, Carbonnelle E, Veckerle C, Brun YV, van Nieuwenhze M, Bouchier C, Tu K, Rice LB, Arthur M.** 2016. Factors essential for L,D-transpeptidase-mediated peptidoglycan cross-linking and beta-lactam resistance in *Escherichia coli*. Elife **5:**e19469.

12. **Both D, Steiner EM, Stadler D, Lindqvist Y, Schnell R, Schneider G.** 2013. Structure of LdtMt2, an L,D-transpeptidase from *Mycobacterium tuberculosis*. Acta Crystallogr D Biol Crystallogr **69:**432-441.

13. **Correale S, Ruggiero A, Capparelli R, Pedone E, Berisio R.** 2013. Structures of free and inhibited forms of the L,D-transpeptidase Ldt_{Mt1} from *Mycobacterium tuberculosis*. Acta Crystallogr D Biol Crystallogr **69:**1697-1706.

14. **Li WJ, Li DF, Hu YL, Zhang XE, Bi LJ, Wang DC.** 2013. Crystal structure of L,D-transpeptidase
Ldt$_{Mt2}$ in complex with meropenem reveals the mechanism of carbapenem against
Mycobacterium tuberculosis. Cell Res **23:**728-731.
15. **Biarrotte-Sorin S, Hugonnet JE, Delfosse V, Mainardi JL, Gutmann L, Arthur M, Mayer C.** 2006.
Crystal structure of a novel beta-lactam-insensitive peptidoglycan transpeptidase. J Mol Biol
359:533-538.
16. **Mainardi JL, Fourgeaud M, Hugonnet JE, Dubost L, Brouard JP, Ouazzani J, Rice LB, Gutmann
L, Arthur M.** 2005. A novel peptidoglycan cross-linking enzyme for a beta-lactam-resistant
transpeptidation pathway. J Biol Chem **280:**38146-38152.
17. **Goffin C, Ghuysen JM.** 2002. Biochemistry and comparative genomics of SxxK superfamily
acyltransferases offer a clue to the mycobacterial paradox: presence of penicillin-susceptible
target proteins versus lack of efficiency of penicillin as therapeutic agent. Microbiol Mol Biol
Rev **66:**702-738.
18. **Lecoq L, Dubee V, Triboulet S, Bougault C, Hugonnet JE, Arthur M, Simorre JP.** 2013. Structure
of *Enterococcus faecium* L,D-transpeptidase acylated by ertapenem provides insight into the
inactivation mechanism. ACS Chem Biol **8:**1140-1146.
19. **Magnet S, Arbeloa A, Mainardi JL, Hugonnet JE, Fourgeaud M, Dubost L, Marie A, Delfosse
V, Mayer C, Rice LB, Arthur M.** 2007. Specificity of L,D-transpeptidases from gram-positive
bacteria producing different peptidoglycan chemotypes. J Biol Chem **282:**13151-13159.
20. **Tipper DJ, Strominger JL.** 1965. Mechanism of action of penicillins: a proposal based on their
structural similarity to acyl-D-alanyl-D-alanine. Proc Natl Acad Sci U S A **54:**1133-1141.
21. **Triboulet S, Arthur M, Mainardi JL, Veckerle C, Dubee V, Nguekam-Nouri A, Gutmann L, Rice
LB, Hugonnet JE.** 2011. Inactivation kinetics of a new target of beta-lactam antibiotics. J Biol
Chem **286:**22777-22784.
22. **Bhattacharjee N, Field MJ, Simorre JP, Arthur M, Bougault CM.** 2016. Hybrid potential
simulation of the acylation of *Enterococcus faecium* L,D-transpeptidase by carbapenems. J
Phys Chem B **120:**4767-4781.
23. **Dubee V, Triboulet S, Mainardi JL, Etheve-Quelquejeu M, Gutmann L, Marie A, Dubost L,
Hugonnet JE, Arthur M.** 2012. Inactivation of *Mycobacterium tuberculosis* L,D-transpeptidase
LdtMt(1) by carbapenems and cephalosporins. Antimicrob Agents Chemother **56:**4189-4195.
24. **Dhar N, Dubée V, Ballell L, Cuinet G, Hugonnet J-L, Signorino-Gelo F, Barros D, Arthur M,
McKinney J.** 2015. Rapid cytolysis of *Mycobacterium tuberculosis* by faropenem, an orally
bioavailable beta-lactam antibiotic. Antimicrob Agents Chemother **59:**1308-1319.
25. **Gupta R, Lavollay M, Mainardi JL, Arthur M, Bishai WR, Lamichhane G.** 2010. The
Mycobacterium tuberculosis protein Ldt(Mt2) is a nonclassical transpeptidase required for
virulence and resistance to amoxicillin. Nat Med **16:**466-469.
26. **Diacon AH, van der Merwe L, Barnard M, von Groote-Bidlingmaier F, Lange C, Garcia-Basteiro
AL, Sevene E, Ballell L, Barros-Aguirre D.** 2016. beta-lactams against tuberculosis-new trick for
an old dog? N Engl J Med **375:**393-394.
27. **Iannazzo L, Soroka D, Triboulet S, Fonvielle M, Compain F, Dubee V, Mainardi JL, Hugonnet
JE, Braud E, Arthur M, Etheve-Quelquejeu M.** 2016. Routes of synthesis of carbapenems for
optimizing both the inactivation of L,D-transpeptidase Ldt$_{Mt1}$ of *Mycobacterium tuberculosis*
and the stability toward hydrolysis by beta-Lactamase BlaC. J Med Chem **59:**3427-3438.
28. **Edoo Z, Iannazzo L, Compain F, Li de la Sierra Gallay I, van Tilbeurgh H, Fonvielle M, Bouchet
F, Le Run E, Mainardi JL, Arthur M, Etheve-Quelquejeu M, Hugonnet JE.** 2018. Synthesis of
avibactam derivatives and activity on beta-lactamases and peptidoglycan biosynthesis
enzymes of mycobacteria. Chemistry doi:10.1002/chem.201800923.
29. **Cordillot M, Dubee V, Triboulet S, Dubost L, Marie A, Hugonnet JE, Arthur M, Mainardi JL.**
2013. In vitro cross-linking of *Mycobacterium tuberculosis* peptidoglycan by L,D-
transpeptidases and inactivation of these enzymes by carbapenems. Antimicrob Agents
Chemother **57:**5940-5945.

30. **Edoo Z, Arthur M, Hugonnet JE.** 2017. Reversible inactivation of a peptidoglycan
transpeptidase by a beta-lactam antibiotic mediated by beta-lactam-ring recyclization in the
enzyme active site. Sci Rep **7:**9136.

31. **Hugonnet JE, Blanchard JS.** 2007. Irreversible inhibition of the *Mycobacterium tuberculosis*
beta-lactamase by clavulanate. Biochemistry **46:**11998-12004.

32. **Tremblay LW, Hugonnet JE, Blanchard JS.** 2008. Structure of the covalent adduct formed
between *Mycobacterium tuberculosis* beta-lactamase and clavulanate. Biochemistry **47:**5312-
5316.

33. **Xu H, Hazra S, Blanchard JS.** 2012. NXL104 irreversibly inhibits the beta-lactamase from
Mycobacterium tuberculosis. Biochemistry **51:**4551-4557.

34. **Hugonnet JE, Tremblay LW, Boshoff HI, Barry CE, 3rd, Blanchard JS.** 2009. Meropenem-
clavulanate is effective against extensively drug-resistant *Mycobacterium tuberculosis*. Science
323:1215-1218.

35. **Tremblay LW, Fan F, Blanchard JS.** 2010. Biochemical and structural characterization of
Mycobacterium tuberculosis beta-lactamase with the carbapenems ertapenem and
doripenem. Biochemistry **49:**3766-3773.

36. **Frere JM, Ghuysen JM, Iwatsubo M.** 1975. Kinetics of interaction between the exocellular DD-
carboxypeptidase-transpeptidase from *Streptomyces* R61 and beta-lactam antibiotics. A
choice of models. Eur J Biochem **57:**343-351.

37. **Bourguignon-Bellefroid C, Wilkin JM, Joris B, Aplin RT, Houssier C, Prendergast FG, Van
Beeumen J, Ghuysen JM, Frere JM.** 1992. Importance of the two tryptophan residues in the
Streptomyces R61 exocellular DD-peptidase. Biochem J **282 (Pt 2):**361-367.

38. **McDonough MA, Anderson JW, Silvaggi NR, Pratt RF, Knox JR, Kelly JA.** 2002. Structures of
two kinetic intermediates reveal species specificity of penicillin-binding proteins. J Mol Biol
322:111-122.

39. **Silvaggi NR, Josephine HR, Kuzin AP, Nagarajan R, Pratt RF, Kelly JA.** 2005. Crystal structures
of complexes between the R61 DD-peptidase and peptidoglycan-mimetic beta-lactams: a non-
covalent complex with a "perfect penicillin". J Mol Biol **345:**521-533.

40. **Soroka D, Ourghanlian C, Compain F, Fichini M, Dubée V, Mainardi J, Hugonnet J, Arthur M.**
2017. Inhibition of β-lactamases of mycobacteria by avibactam and clavulanate. J Antimicrob
Chemother **72:**1081-1088.

41. **Chow C, Xu H, Blanchard JS.** 2013. Kinetic characterization of hydrolysis of nitrocefin, cefoxitin,
and meropenem by beta-lactamase from M*ycobacterium tuberculosis*. Biochemistry **52:**4097-
4104.

Figure 1. Impact of amino acid substitutions on fluorescence kinetics. (**A**) Inactivation reaction scheme. E, free Ldt$_{fm}$; EI^{An}, amine anion; EI*, acylenzyme; I, imipenem. (**B**) Fluorescence variations during Ldt$_{fm}$ inactivation. The charts show the variations in the relative concentrations of the three enzyme forms (left panel) and the relative fluorescence intensity of the three enzyme forms (middle panel), which account for the observed fluorescence kinetics (right panel). (**C**) Fluorescence kinetics of Ldt$_{fm}$ inactivation by imipenem. Trp residues were replaced by Phe residues in various combinations. Designation of the remaining Trp residues: a, Trp^{355}; b, Trp^{385}; c, Trp^{410}; d, Trp^{415}; e, Trp^{425}; f, Trp^{434}. Insets present the initial decreases of fluorescence over a short timescale.

Figure 2. Inactivation efficacy and relative fluorescence intensity of Ldt$_{fm}$ and derivatives with Trp to
Phe and His to Ala substitutions. (**A**) Kinetic parameter k_1 for formation of the amine anion. (**B**) Kinetic
parameter k_2 for the acylation step. (**C**) Fluorescence intensity of the three enzyme forms relative to
that of the free form of the wild-type enzyme. E, Ldt$_{fm}$; EI^{An}, amine anion; EI*, acylenzyme. All, presence
of the full complement of Trp residues (a, d, c, d, e, and f). Trp residues were replaced by Phe residues
in various combinations. Designation of the remaining Trp residues: a, Trp^{355}; b, Trp^{385}; c, Trp^{410}; d,
Trp^{415}; e, Trp^{425}; f, Trp^{434}.

Figure 3. Variations in fluorescence intensity resulting from acylation of L,D-transpeptidase Ldt$_{fm}$ and β-lactamase BlaC by various β-lactams and β-lactamase inhibitors. Stopped-flow fluorescence kinetics were obtained with final concentrations of 10 μM for the enzymes and 100 μM for β-lactams and β-lactamase inhibitors (**A**) The negative charge developing on the amine nitrogen upon acylation of Ldt$_{fm}$ by nitrocefin is stabilized by a mesomeric effect involving the electron withdrawing dinitrobenzene (30, 41). (**B**) Transfer of the negative charge to a leaving group in the concerted mechanism leading to acylation of Ldt$_{fm}$ by ceftriaxone (7). (**C**) Tautomeric forms of the acylenzyme resulting from acylation of BlaC by clavulanate and decarboxylation of the β-lactamase inhibitor (32, 40). (**D**) BlaC-avibactam carbamoylenzyme (33). (**E** and **F**) Acylation of BlaC and Ldt$_{fm}$ by faropenem, respectively (24). (**G**) Accumulation of an amine anion in the onset of the reaction of hydrolysis of meropenem by BlaC (BlaC^{An}). This is followed by protonation of the β-lactam nitrogen (35).