Pioneer and nonpioneer factor cooperation drives lineage specific chromatin opening

Alexandre Mayran[1,2], Kevin Sochodolsky[1], Konstantin Khetchoumian[1], Juliette Harris[1], Yves Gauthier[1], Amandine Bemmo[1] Aurelio Balsalobre[1] and Jacques Drouin[1,2,3,4]

[1] Laboratoire de génétique moléculaire, Institut de Recherches Cliniques de Montréal (IRCM), Montreal, Québec, Canada,

[2] Department of Biochemistry, McGill University, Montreal, Québec, Canada

[3] Département de Biochimie, Université de Montréal, Montreal, Québec, Canada

[4]Correspondence: jacques.drouin@ircm.qc.ca

Main text: 4021 words /26674 ch

Summary

Pioneer transcription factors are coined as having the unique property of "opening closed chromatin sites" for implementation of cell fates. We previously showed that Pax7 pioneers pituitary melanotrope cell identity through deployment of an enhancer repertoire: this allows binding of Tpit, a nonpioneer factor that drives terminal differentiation of melanotropes and the related corticotropes. Here, we used scRNAseq and chromatin accessibility profiling to define shared and cell-specific gene expression programs and chromatin landscapes in these lineages. *Pax7-* and *Tpit*-deficient pituitaries fail to deploy the melanotrope enhancer repertoire indicating that both factors Pax7 and Tpit are required for chromatin opening. Finally in cells, binding of heterochromatin targets by Pax7 is independent of Tpit but Pax7-dependent chromatin opening requires Tpit. The present work shows that pioneer core properties are limited to the ability to recognize heterochromatin targets and facilitate nonpioneer binding. Chromatin opening per se may be provided through cooperation with nonpioneer factors.

Introduction

In development, cell differentiation leads to establishment of cell identities through the sequential action of transcription factors (TFs) acting as specification or determination factors. These processes involve master TFs that primarily act on the epigenome to open new regulatory chromatin landscapes. This is achieved by a unique class of TF, pioneer TFs. Unlike canonical TFs, pioneers bind "closed" chromatin, trigger chromatin opening and thus allow binding of other TFs.

For example, Foxa1 is essential for liver fate and binds regulatory sequences before gene activation (Cirillo et al., 2002; Lee et al., 2005). Ebf1 also acts as a pioneer factor during B-cell development and Neurod1 and Ascl1 (Mash1) were suggested to function as pioneer factors during neural development (Pataskar et al., 2016; Wapinski et al., 2017). Ectopic expression of pioneer factors is sufficient to drive trans-differentiation: thus, C/EBPα can direct trans-differentiation of pre-B cells into macrophages (Di Tullio et al., 2011). The most extreme example is the reprogramming of fibroblasts into induced pluripotent stem cells through action of the pioneer factors Oct4, Klf4 and Sox2 (Okita et al., 2007; Soufi et al., 2015). Similarly, C/EBPα can direct trans-differentiation of pre-B cells into macrophages (Di Tullio et al., 2011).

The ability to open chromatin has been implicitly considered as a core property of pioneer TFs and this would allow nonpioneer binding to newly accessible sites. Whether nonpioneers play a role in pioneer-driven chromatin opening has nor been assessed. Here, we used normal and perturbed pituitary differentiation to investigate the pioneer model and establish the specific and/or overlapping functions of the differentiation regulators Pax7 (pioneer) and Tpit (nonpioneer).

Two pituitary lineages express the hormone precursor pro-opiomelanocortin (POMC),

the melanotropes and corticotropes. Both require Tpit for POMC expression and cell fate

determination (Lamolet et al., 2001; Pulichino et al., 2003). Indeed, Tpit implements a

secretory cell transcriptional program by activation of scaling factors for translation and

secretory organellogenesis (Khetchoumian et al., 2018). While sharing the secretory

POMC identity, corticotropes and melanotropes differ by their functions as these POMC-

expressing lineages control corticosteroidogenesis and pigmentation, respectively

(Drouin, 2017). The pioneer Pax7 drives melanotropes specification (Budry et al., 2012)

through deployment of a melanotrope-specific enhancer repertoire (Mayran et al., 2018).

Here, we first establish that the two POMC lineages share a transcriptional program that

is distinct from other pituitary cells and that in addition, they each have a unique program

of gene expression. We then show that these two layers of identity (shared and lineage-

specific) are reflected at the level of chromatin accessibility and that the shared POMC

chromatin landscape requires Tpit. Further, Tpit is required for the opening of the Pax7-

dependent melanotrope chromatin landscape indicating that Pax7 and Tpit act together

during melanotrope differentiation. Finally, Pax7 and Tpit have complementary roles as

only Pax7 can bind heterochromatin while Tpit binding is needed for chromatin opening.

In summary, we propose that the essence of pioneer action is the ability to recognize and

bind DNA sites in closed chromatin whereas cooperating nonpioneer TFs, such as Tpit,

may drive chromatin opening.

Results

Diversity of pituitary cell lineages

The pituitary is a highly specialized organ where each lineage serves as a hormone producing factory. Each cell type is dedicated to the regulation of a specific endocrine organ and responds to specific signals from the hypothalamus and body. We used single cell RNAseq to decipher the transcriptional complexity of the different pituitary lineages. For each lineage, hormone-coding mRNAs are so abundant that they appear as peaks in cDNA libraries (Figure S1A). Profiling of adult mouse male pituitary cells (Figure S1B) was achieved by plotting single cell data using t-distributed stochastic neighbor embedding (t-SNE), a common method (Maaten, 2008) that uses dimensionality reduction to cluster together cells with similar transcriptional profiles (Figure 1A). This showed that cells expressing the same pituitary hormone cluster together and that they all express Pitx1, a marker of the oral ectoderm origin of the pituitary (Drouin, 2017) (Figure S1C and 1B). We identified 12 clusters composed of endocrine and non-endocrine cells (Figure 1C). Cluster 1 corresponds to somatotropes as they express the growth hormone (Gh) gene. Lactotropes that produce Prolactin (Prl) are found in cluster 2. Clusters 4 and 5 correspond to melanotropes and corticotropes, respectively; both express the *POMC* gene, yet only melanotropes express *Pcsk2*. Gonadotropes that express the *Lhβ* gene are in cluster 8. We also detected thyrotropes as *Tshβ*-expressing cells; however they did not appear as a separate cluster (Figure S1C). The pituitary stem cells that express Sox2 (Fauquier et al., 2008) are found in cluster 7. Although our study aimed at defining the transcriptome of the different pituitary lineages, we also uncovered several non-endocrine cells within the pituitary tissue that do not express Pitx1 (Figure 1B). We identified endothelial cells (cluster 9), macrophages (cluster 10), posterior

pituicytes (cluster 11) and pericytes (cluster 12). Cluster 3 is fragmented in three different

groups of cells that express either GH, Prolactin, POMC or Lhβ. We performed differential

expression analysis between each sub-cluster 3 and its matching cell type in order to

define these subsets. In all cases, cells of these subsets are specifically depleted of

ribosomal proteins and enriched for mitochondrial RNA. This was shown to be an artefact

of tissue dissociation (Ilicic et al., 2016) and to represent cells affected by the preparation:

we excluded cluster 3 from following analyses.

We then compared the transcriptomes of the different pituitary lineages by performing

differential expression analysis between clusters 1, 2, 4, 5, 7 and 8. Genes with two-fold

differential expression between each cluster (p value <0.05 and minimum 0.2 UMI in at

least one pooled cluster) are shown as a heatmap (Figure 1D). The Sox2+ stem cell niche

is the most transcriptionally divergent from other pituitary lineages based on correlation

clustering analyses. The two Pit1-dependent lineages, the lactotropes and somatotropes,

are the most different compared to the two POMC lineages and gonadotropes that cluster

together. Within the latter group, corticotropes and melanotropes are more

transcriptionally correlated together than with gonadotropes. Thus, the two POMC

lineages, melanotropes and corticotropes, have both a shared and a specific

transcriptional program.

Lineage-specific chromatin landscapes

We next aimed to identify cis-regulatory elements that regulate the transcriptional identity

of pituitary lineages. We used ATACseq (Buenrostro et al., 2013) to identify putative

regulatory elements that are accessible in each lineage. We complemented our

previously published datasets of purified melanotrope and corticotrope ATACseq with

accessibility profiles for gonadotropes and anterior lobe (AL) cells. Gonadotropes were

FACS-purified from transgenic pituitaries expressing the LHβ-Cerulean transgene (Budry

et al., 2011). As control, we isolated the remaining AL cells that are mostly composed of

a combination of Pit1-dependent somatotropes and lactotropes. Interestingly, we found

that the promoters of hormone genes *POMC*, *αGSU*, *Gh* and *Prl* as well as lineage

specifiers *Tpit* (*Tbx19*), *Pax7*, *SF-1* (*Nr5a1*) and *Pit1* show lineage-specific accessibility

(Figure 2A and S2A-C). However, the *Pcsk2* promoter is accessible in all pituitary

lineages but its numerous distal accessible sites (putative enhancers) are only accessible

in melanotropes. Globally, we identified 98926 open chromatin regions across the

pituitary lineages (Figure 2B). Segregation of lineage-specific accessibility yielded 33451

regions opened in all lineages, 14025 regions opened in a combination of three lineages,

20374 in two lineages and finally 31076 opened in only one lineage. Thus, there are

regions specifically accessible in melanotropes, corticotropes, gonadotropes or in the AL.

In accordance with the close transcriptional correlation between melanotropes and

corticotropes, we also found shared regions accessible in both melanotropes and

corticotropes (POMC-specific) but closed in gonadotropes and AL. Finally, we identified

13130 pituitary-specific sites that are closed in embryonic stem cells, as well as a set of

20321 regions accessible in both pituitary and ES cells (Figure S2D). These ubiquitous

peaks are for the majority (58%) composed of promoter elements (Figure S2E) while

regions opened in two or in only one pituitary lineage are mostly distal elements (94%

and 96%, respectively). This reinforces the idea that promoter accessibility is established

early during differentiation. Further, this suggests that lineage specific opening of

promoters tends to be an exception and may be involved in restricting appropriate

expression of critical genes such as hormone-coding genes and lineage specifiers.

Comparison of the spatial relationship between the POMC-specific transcriptional

program (Figure 2C) and lineage-specific accessibility (Figure 2D) shows that POMC-

specific gene promoters tend to be closer to both POMC- and melano-specific open

regions (Figure 2E) and are enriched for the motif of the lineage specifier Tpit (Figure 2F).

Melanotrope genes are also close to their lineage-specific open chromatin regions and

their lineage specific chromatin landscape is enriched for the motif of their lineage

specifier Pax7 (Figure 2G-J). Similar relationships were found for the Pit1-dependent

lineages, the corticotropes and the gonadotropes (Figure S2F-G).

Pax7 is required for phenotypical features of melanotropes

The two POMC lineages have shared and specific transcriptional programs and open

chromatin landscapes. Their most obvious similarity is expression of the hormone

precursor POMC as well as expression of the terminal differentiation factor Tpit. However,

POMC expression varies between these two lineages with low (blue) and high expression

(red) cells in scRNAseq analyses (Figure 3A). Indeed, POMC is more highly expressed

in Pax7-expressing melanotropes compared to GR (nr3c1)-expressing corticotropes as

shown in t-SNE and differential expression Volcano plots (Figure 3A,B). Analysis of the

POMC-EGFP (Lavoie et al., 2008) transgenic pituitaries (Figure 3C) also showed high

and low-expressing cells. Indeed, IL (melanotropes) transgenic pituitary cells have much

higher EGFP levels compared to most EGFP-positive corticotrope cells of the AL (Figure

3D). This indicates that the transgene expressed in both melanotropes and corticotropes

is subject to cell-type specific regulation and can be used as a surrogate phenotypical

readout. Strikingly in a *Pax7-/-* background, melanotropes express the EGFP transgene

at the same level as in AL corticotropes (Figure 3D, E). Melanotropes are also typically

larger and have a more complex organelle content than corticotropes as assessed by

forward and side scatter distributions in FACS profiles and this is also lost in Pax7-

deficient mice (Figure S3A, B). Thus, all three melanotrope features (high POMC

expression, large cell size and granularity) are fully dependent on Pax7 and virtually all

melanotropes switch to a corticotrope phenotype in *Pax7-/-* mice (Figure 3F). However,

the proportion of total EGFP-positive cells in AL or IL is not affected by loss of Pax7

(Figure 3G). Thus, Pax7 implements melanotrope but not the shared POMC cell identity

which is under the control of Tpit (Pulichino et al., 2003).

Pax7 and Tpit are required for opening cognate lineage-specific enhancer
landscape

We next sought to uncouple the specific roles of Pax7 and Tpit for implementation of the

shared and melanotrope-specific chromatin landscapes. To do so, we performed

ATACseq on IL cells from mice of genetic backgrounds lacking one or both alleles of *Pax7*

and/or *Tpit*. We used *Pax7* and *Tpit* double heterozygote littermates as control; these

show minimal differences compared to wild-type animals (Figure S4). We found that

melanotrope-specific peaks (Figure 2B) are lost in *Pax7-/-;Tpit+/-* IL whereas they are

present in wild-type or double heterozygotes (Figure 4A) in agreement with previous data

(Mayran et al., 2018) that showed Pax7 requirement for accessibility of melanotrope

regulatory modules. Also, we found that Tpit is required for the open status of POMC-

specific open chromatin (Figure 4B). This suggests that Tpit is involved in the pioneering

process of the shared POMC lineage enhancers. For example, POMC gene expression

critically relies on Tpit and accordingly, both its promoter and enhancer accessibility

strongly depend on Tpit but not on Pax7 (Figure 4C). This is also true for the promoter of

POMC-specific gene *Tnxb* and the Tpit-dependent enhancer (Khetchoumian et al., 2018)

of the *Creb3l2* gene (Figure 4C). Thus, Tpit is required for opening the shared POMC

regulatory modules while Pax7 is only required for melanotrope enhancers.

Tpit is required for Pax7-dependent chromatin opening

In order to define and compare the Pax7 and Tpit-dependent ATACseq landscapes, we

performed differential enrichment analysis (p<0.05 and fold changes >2) and found 16024

sites altered in *Tpit-/-* IL (Figure 5A). Most changes are decreased accessibility (12573

sites) and these show greater quantitative differences compared to sites with increased

accessibility (3451 sites). Analysis of these Tpit-dependent sites revealed two subsets,

one dependent and one independent of Pax7 (Figure 5B). In both Tpit-dependent

subsets, chromatin accessibility in *Pax7-/-;Tpit-/-* IL is the same as in *Tpit-/-*. In order to

ascertain the reliability of these analyses, we validated dependence on Pax7, Tpit or both

by qPCR analyses of three independent ATACseq libraries of each genotype (Figure S5).

Principal component analysis using all Tpit-regulated sites shows that most of the

variance between samples (77%) is explained by component 1 (Figure 5C). Consistent

with the heatmaps of Figure 5B, wild-type and *Pax7+/-;Tpit+/-* cluster together while *Pax7*

knockout are between wild-type and *Tpit* knockout samples. Thus, a subset of Tpit

chromatin targets are also dependent on Pax7. We then focused on Pax7-dependent

chromatin access to assess whether co-dependency on Tpit and Pax7 is a general

feature of Pax7-dependent sites. We found 7057 Pax7-dependent sites (Figure 5D): most

changes are decreased accessibility (6112 sites) and these are greater effects compared

to increased sites (945 sites). Strikingly, and unlike Tpit-dependent accessibility, virtually

all Pax7-dependent sites are also dependent on Tpit (Figure 5E). Accordingly, principal

component analysis of Pax7-dependent sites showed that Pax7 knockout samples cluster

with *Tpit* knockout and Pax7 on component 1 (79% of variance). This indicates that *in*

vivo, Tpit is absolutely required for the establishment of the Pax7-dependent chromatin

landscape. Thus in both cases, *Pax7* and *Tpit* double knockout samples cluster closely

with the *Tpit* single knockout (Figure 5C, F) showing that loss of both factors does not

have a greater impact on accessibility than the loss of Tpit alone.

To assess the biological basis of this co-dependency, we focused on the *Pcsk2* gene,

encoding the PC2 protein, a hallmark of melanotrope identity. We previously showed

strong Pax7 dependency for chromatin opening of an upstream enhancer (Budry et al.,

2012) together with multiple distal elements that are ATACseq sensitive in a lineage-

specific manner. Interestingly, this melanotrope-specific chromatin accessibility is

confined within a topologically associated domain (TAD) and stops at the border of this

TAD (Figure 5G); similar TAD-wide opening was observed at the *Drd2* and *Grik1* loci

(Figure S5G). In accordance with the role of Pax7 for *Pcsk2* expression in melanotropes,

we found that this TAD-wide chromatin opening doesn't occur in *Pax7-/-* animals.

Consistent with our previous observation that Pax7-dependent access also depends on

Tpit, chromatin opening within this TAD does not take place in *Tpit-/-* and in *Pax7-/-;Tpit-*

/- animals. Thus, Tpit is required for opening of the *Pcsk2* TAD. It is noteworthy that *Pcsk2*

is also expressed, albeit at low levels, in gonadotropes (Figure 5H). This gonadotrope

pattern of *Pcsk2* expression is also found in the *Tpit-/-* IL cells that have switched fate

(Budry et al., 2012); these cells occupy the dorsal side of the *Tpit-/-* mutant IL (Figure 5I).

In contrast, the ventral side of the same *Tpit-/-* IL harbors Pax7-positive cells that fail to

express Pcsk2 (Figure 5I). This confirms that Pax7 functionally requires Tpit to establish

melanotrope identity.

Productive Pax7 pioneer action in Tpit-positive cells

In order to evaluate the chromatin binding properties of Pax7 and Tpit, we compared their

binding properties to those of a pituitary nonpioneer factor Pitx1 and to two pioneer

factors, Neurod1 that is expressed in pituitary corticotropes (Lamolet et al., 2004; Poulin

et al., 1997), and the reprogramming factor Sox2 (Okita et al., 2007). The heatmaps of

ATACseq signals for all binding sites of these five TFs in AtT20 cells reveal striking

differences between the profiles of Tpit and Pitx1, compared to those of the pioneers,

Neurod1, Pax7 and Sox2 (Figure 6A). Whereas Tpit and Pitx1 mostly (~ 95% of all sites)

bind to sites that are open (ie. have a detectable ATACseq signal), a significant proportion

of the pioneer binding sites have little or no ATACseq; thus for Pax7 and Neurod1, ~ 30

% of binding sites are in inaccessible chromatin whereas this is true for ~ 60% of all Sox2

binding sites. The ability to bind closed chromatin (ATAC-negative) appears to be a

common property of the three pioneer factors in contrast to the non-pioneers Tpit and

Pitx1.

In order to reveal the contribution of Tpit to chromatin pioneering by Pax7, we compared

the impact of Pax7 expression in two different pituitary cell lines, one that expresses Tpit

(AtT20) and one that does not (αT3). In agreement with expression data, the Tpit locus

exhibits ATACseq signals in AtT20 cells but not in αT3 cells (Figure 6B) whereas the

locus for the gonadotrope-specific regulator SF1 presents ATACseq signals in αT3 but

not in AtT20 cells (Figure 6C).

We looked at the chromatin accessibility by ATACseq before/after Pax7 expression

(Budry et al., 2012; Mayran et al., 2018) in both lineages and found that in AtT-20 cells,

Pax7 binding leads to opening of 6607 regions. In contrast, only 661 sites are opened

following Pax7 expression in αT3 cells and this opening is of greater magnitude in AtT-

20 cells (Figure 6D). Thus, expression of Pax7 in αT3 cells does not lead to efficient

chromatin opening. We have previously identified steps of pioneering as follow: initial

weak binding to closed enhancers (less than 30 minutes), stabilization of pioneer binding

followed by chromatin opening (Mayran et al., 2018). Failure to perform any of these three

steps should impede Pax7 pioneering. To identify which specific step limits Pax7

pioneering in αT3 cells, we compared Pax7 binding in AtT-20 and αT3 cells (Figure S6A).

Pax7 has similar ability to bind closed chromatin in both AtT-20 and αT3 cells (Figure 6A

and S6). We extracted all Pax7 sites bound to closed chromatin in AtT-20 and in αT3

(Figure 6E). In both AtT-20 and αT3 cells, there are sites with strong and low binding

signals. This suggest that Pax7 is able to bind strongly to closed chromatin in both cell

context. However in AtT-20 cells, Pax7 binding strength correlates (r^2=0.41) with

accessibility after Pax7 expression whereas in αT3 cells, there is no correlation (r^2=0.09)

between Pax7 binding and post-Pax7 chromatin opening (Figure 6E). We identified a

subset of heterochromatin Pax7-bound sites based on their ability for Tpit binding after

Pax7 expression; these sites are not bound by Tpit in absence of Pax7 (Figure 6F) and

become accessible in presence of Tpit and Pax7 (Figure 6G). It is noteworthy that only

these newly accessible sites contain the Tpit DNA binding motif (Figure 6H). This

suggests that Pax7-dependent Tpit binding also depends on the Tpit DNA motif.

Notwithstanding, Pax7 and Tpit interact directly in vitro as shown in pull-down assays

(Figure 6I). The interaction between the two factors may also contribute to their

cooperation. This suggests that Pax7`s ability to bind strongly to closed chromatin is not

dependent on cell specific factors and that stable binding is not sufficient to drive

chromatin opening. During melanotrope differentiation, this pioneer-dependent chromatin

opening also requires Tpit.

Discussion

Pioneer factors are coined as "factors that can open closed chromatin". This label implied

that pioneers were expected to directly provide this ability. The present work shows that

there can be division of labor between pioneer and nonpioneer factors: in the case of

Pax7 and Tpit, the former recognizes and engages pioneering sites and the latter

provides the chromatin opening ability. Indeed, we show that Tpit is required for chromatin

opening at sites that must first be pioneered by Pax7, and we did not find evidence for

the reverse. While we envision that this is made possible by Tpit-dependent recruitment

of chromatin remodeling machineries, the finding clearly circumscribes the unique

features of the Pax7 pioneer function. Namely, the unique aspect of pioneer action is the

ability to bind DNA sites within closed chromatin that are inaccessible to probing by

techniques such as ATACseq; this appears to be a property shared with other pioneers

such as Sox2, NeuroD1 (Figure 6A) in contrast to nonpioneer factors such as Tpit or

Pitx1.

For Pax7, we have shown that this initial binding to closed heterochromatin (Figure 7A)

is rapid (within 30 min, Figure 7B) and that the first detectable change in chromatin

structure at these binding sites is revealed by stabilization (within 24 h, Figure 7C) of Pax7

binding (Mayran et al., 2018). This stabilization precedes chromatin opening (revealed by

ATACseq). Here, we find that strong Pax7 binding occurs even in absence of Tpit (ie in

αT3 cells, Figure 6E): this excludes Tpit`s involvement in Pax7 stabilisation of binding. In

contrast, the next step of pioneer action, namely chromatin opening, requires Tpit as it

does not occur in αT3 cells. The recruitment of Tpit at a subset of Pax7 closed chromatin

binding sites (Figure 7D) that have a Tpit DNA binding site in the vicinity results in

chromatin opening (Figure 7E). It would thus be the combined interaction of Pax7 with

Tpit together with the latter's ability to bind its DNA site exposed through the initial action

of Pax7 that altogether would lead to completion of pioneer action with Tpit bringing in the

chromatin remodeling ability (Figure 7E). With these interdependent functions, the

cooperation between pioneer and nonpioneer factors provides robust stringency and a

fail safe mechanism for triggering chromatin opening at a very specific subset of Pax7

sites.

The requirement for DNA binding sites for both Pax7 and Tpit at the Pax7 pioneered

enhancers is reminiscent of the action of other pioneers where two different pioneers

cooperate for chromatin opening. For example, the pluripotency factors cooperate with

each other to facilitate the binding of other and binding of multiple pioneers is required at

some sites to drive chromatin opening (Chronis et al., 2017; Soufi et al., 2015). Also, there

is cooperation between FoxA1 and GATA4 for chromatin binding (Donaghey et al., 2018).

Foxa1 also facilitates the binding of the estrogen receptor by opening chromatin (Hurtado

et al., 2011; Laganiere et al., 2005). Reciprocally, another study showed that steroid

receptors can facilitate the binding of Foxa1 (Swinstead et al., 2016). This suggest that in

the case of steroid receptors and Foxa1, both can play the role of pioneer at specific

subsets of their targets. To our knowledge, the required participation of a nonpioneer for

pioneer-driven chromatin opening has not been shown so far.

Prior data indicated that pioneer binding precedes gene activation, for example the

binding of FoxA at liver targets precedes their activation (Cirillo et al., 1998). More

recently, Grainy head binding at pioneer sites was shown to determine chromatin opening

(ATAC signal) but not enhancer activity (Jacobs et al., 2018).

The required cooperation between pioneer and nonpioneer factors for implementation of

a specific genetic program ensures specificity of action. For example, Pax7 is also

involved in cell fate specification in muscle and neural tissues in addition to pituitary but

its interdependent action with Tpit in the pituitary ensures that Pax7 access to pituitary-

unrelated sites may not result in chromatin opening in absence of Tpit. This essential

cooperation between specification and determination factors provides robustness for

lineage identity by preventing mis-activation of inappropriate gene regulatory networks.

This limitation to the capacity of pioneer factors would allows specific combinations of

pioneers and nonpioneers to activate different regulatory networks and explain the wide

variability in targets depending on the same pioneer factor in different contexts.

ACKNOWLEDGMENTS

We are very grateful to our colleague Nicole Francis for comments on the manuscript, to

Odile Neyret for NGS analyses, Eric Massicotte for FACS sorting, Dominic Fillon for

microscopy, Simone Terouz for histology preparations, Dimitar Dimitrov for mouse

management and to Evelyne Joyal for her expert secretarial assistance. Data analyses

were possible thanks to the support of Compute Canada. This work was supported by a

Foundation grant to J.D. from the Canadian Institutes of Health Research.

AUTHOR CONTRIBUTIONS

A.M. and J.D. conceived the study, A.M., Au.B. and J.D. conceived and designed the

experiments, A.M., K.S. and K.K. performed mouse tissue collection, A.M. performed

ATACseq assays, A.M. and Y.G. performed ChIP experiments, K.S. performed

immunofluorescence assays, J.H. did the pull-down assays, A.M. and Am.B. analyzed

the single cell RNAseq, A.M. performed analysis of ATACseq and ChIPseq, A.M. and

J.D. wrote the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

Legends to Figures

Figure 1. Transcriptional complexity of the pituitary adult gland

    A. t-SNE map (t-distributed stochastic neighbor embedding) plot of the 9269 profiled pituitary cells colored by the 12 clusters identified using unsupervised k-means clustering. Cluster identification included expression of the hallmarks gene(s) indicated between parenthesis and the markers shown in Figure 1C.

    B. t-SNE map showing color-coded Pitx1 expression.

    C. t-SNE map showing color-coded expression of indicated markers for the major pituitary lineages.

    D. Heatmap showing normalized expression of the 1000 most differentially expressed genes (p value<0.05, FC>2 and minimum number of UMI >0.3) in clusters representing the different endocrine and progenitor cells (cluster 1, 2, 4, 5, 7, 8). Rows are centered; unit variance scaling is applied to rows. Both rows and columns are clustered using correlation distance and average linkage.

Figure 2. Lineage-specific chromatin access reveals lineage regulators

    A. Genome browser view (IGV) of ATACseq profiles at genes marking pituitary lineages: shared POMC markers (green), melanotrope (red) or gonadotrope (orange). The *SF1* promoter is indicated by an arrow.

    B. Heatmaps showing ATACseq signals (RPKM) across the different pituitary lineages in a 4kb window around the ATACseq peak center indicated by an arrow. Colored boxes indicate peaks specifically enriched in the indicated lineage.

C. Boxplot showing expression (UMI) of gene markers of the POMC lineages across

the different lineages. Center lines show medians; box limits indicate the twenty-

fifth and seventy-fifth percentiles; whiskers extend to 1.5 times the interquartile

range from the twenty-fifth to seventy-fifth percentiles.

D. Average profiles of ATACseq signals at POMC-specific ATACseq peaks.

E. Box-plot of distances between the TSS of POMC-specific genes and the closest

ATACseq peak in POMC, melanotropes and gonadotrope cells. Box plot features

as in C.

F. Motif enriched (assessed by HOMER) under POMC-specific ATACseq peaks and

not found in other subsets.

G. Box-plot showing expression (UMI) of melanotrope gene markers across the

different lineages. Box plot features as in C.

H. Average profiles of ATACseq signals at melanotrope specific ATACseq peaks.

I. Box-plot of distances between the TSS of melanotrope-specific genes and the

closest ATACseq peak in POMC, melanotrope and gonadotrope cells. Box plot

features as in C.

J. Motif enriched (assessed by HOMER) under melanotrope specific ATACseq peaks

and not found in other subsets.

Figure 3. Pax7 implements melanotrope features onto a shared POMC cell identity

A. t-SNE map of adult pituitary cells colored for expression of the POMC lineage

regulator Tpit (Tbx19). Enlarged panels showing two clusters of POMC cells with

high and low POMC levels, melanotropes expressing Pax7 and corticotropes

expressing GR.

B. Volcano plot showing differential transcription factor gene expression (p value vs log2 FC) between the high versus low POMC expressing cells. GR and Pax7 are highlighted.

C. Experimental scheme to assess Pax7 dependence of melanotrope phenotypical features. Transgenic mice expressing POMC-EGFP were crossed into *Pax7-/-* mice and compared to WT. The two pituitary lobes were dissected for each genotype and analyzed by FACS.

D. Representative FACS profiles showing cell populations with different POMC-EGFP transgene levels in intermediate (IL) and anterior lobes (AL) of *WT* (n=5) and *Pax7 KO* (n=3) pituitaries. Blue and red shading labels low and high EGFP levels, respectively.

E. Bar graph showing ratios of EGFP signals in *WT* IL, *Pax7 KO* AL, *Pax7 KO* IL compared to *WT* AL. The analyses included 5 *WT* and 3 *Pax7 KO* replicates. P values were computed using unpaired two-sided t-test.

F. Bar graph showing the proportion of high versus low EGFP expressing cells based on fluorescence signals detected by FACS. P values were computed using unpaired two-sided t-test.

G. Bar graph showing the proportion of EGFP expressing and non-expressing cells based on fluorescence signals detected by FACS. P values were computed using unpaired two-sided t-test.

Figure 4. Pax7 and Tpit are required for opening cognate enhancer landscapes

A. Read density heatmaps showing ATACseq signals (RPKM) across the different pituitary lineages in a 4kb window centered at melanotrope specific ATAC peaks

(left panel). Right panel shows corresponding ATACseq heatmaps in the ILs of

WT, *Pax7$^{+/-}$;Tpit$^{+/-}$* (labeled Pax7$^{+/-}$) and *Pax7$^{-/-}$;Tpit$^{+/-}$* (labeled Pax7$^{-/-}$) mice.

B. Read density heatmaps showing ATACseq signals (RPKM) across the different

pituitary lineages in a 4kb window centered at POMC-specific ATAC peaks (left

panel). Right panel shows corresponding ATACseq heatmaps in the intermediate

lobe of *WT*, *Pax7$^{+/-}$;Tpit$^{+/-}$* (labeled Pax7$^{+/-}$) and *Pax7$^{-/-}$;Tpit$^{+/-}$* (labeled Pax7$^{-/-}$) mice.

C. Genome browser view (IGV) of ATACseq profiles at POMC-specific ATACseq

peaks in the different pituitary lineages and in ILs of *Pax7$^{+/-}$;Tpit$^{+/-}$*, *Pax7$^{-/-}$;Tpit$^{+/-}$*

and *Pax7$^{+/-}$;Tpit$^{-/-}$* mice.

Figure 5. Pax7 dependent chromatin landscape requires Tpit.

A. Dispersion plot showing average ATACseq rlog values (assessed by Deseq2) over

the log2 fold changes of *Tpit* heterozygote versus *Tpit* knockout IL at all accessible

regions. Differentially accessible regions (p-value <0.05 and log2 FC > +/- 1 as

computed by Deseq2) are shown as red circles.

B. Read density heatmaps showing ATACseq signals at Tpit-dependent chromatin

opening (log2 FC<-2) in the indicated mouse genotypes.

C. Principal component analysis of the ATAC signals at all 16024 Tpit regulated

chromatin opening across the tested genotypes.

D. Dispersion plot showing average ATACseq rlog values (assessed by Deseq2) over

the log2 fold changes of *Pax7* heterozygote versus *Pax7* knockout IL at all

accessible regions. Differentially accessible regions (p value <0.05 and log2 FC >

+/- 1 as computed by Deseq2) are shown as red circles.

E. Read density heatmap showing ATACseq signals at Pax7-dependent chromatin
opening (log2 FC<-2) in the indicated mouse genotypes.

F. Principal component analysis of the ATAC signals at all 7058 Pax7 regulated
chromatin opening across the tested genotypes.

G. Hi-C interaction map (top) from mouse ES cells (Bonev et al., 2017) around the
Pcsk2 locus showing the boundaries of the *Pcsk2* TAD. Genome browser views
(bottom) of the ATACseq profiles in purified pituitary cells and ILs of the indicated
genotypes at the corresponding genome location.

H. t-SNE map colored for single cell *Pcsk2* expression showing no Pcsk2 expression
in corticotropes, highest expression in melanotropes and weak expression in
gonadotropes.

I. Co-staining immunofluorescence for Pax7 (red) and PC2 (green) of *Tpit*
heterozygote and *Tpit* knockout pituitaries from 5 days postnatal mice.

Figure 6. Pax7 binding on closed chromatin is only productive in Tpit-expressing cells.

A. Read density heatmaps of ATACseq signal density in a 4kb window centered on
binding sites for the indicated factors. The heatmaps are ranked by their
decreasing ATACseq central (200bp) read densities.

B,C. Genome browser views of ATACseq profiles in AtT-20 and αT3-cells at the Tpit
(B) and SF1 (C) loci.

D. Dispersion plots of central (200 bp) ATACseq read densities in Neo (x-axis) versus
Pax7 (y-axis) expressing AtT-20 (left) and αT3 cells (right) at all Pax7 binding sites

in the indicated cell lines. Colored dots represent sites with significantly stronger

signals after Pax7 expression.

E. Dispersion plots of ATACseq read densities in Pax7-expressing cells (x-axis) over

Pax7 ChIPseq read densities (y-axis) in AtT-20 (left) and αT3 cells (right) at Pax7

binding sites with no ATACseq signal before Pax7 expression.

F,G. Boxplots of Tpit ChIPseq (F) and ATACseq (G) read densities in Neo and Pax7

expressing AtT-20 cells at Pax7 sites no ATACseq signal before Pax7 expression

subdivided into Tpit-bound (>1 RPKM in Pax7 expressing cells, blue) and not bound

by Tpit (<1 RPKM in Pax7 expressing cells, grey). Center lines show medians; box

limits indicate the twenty-fifth and seventy-fifth percentiles; whiskers extend to 1.5

times the interquartile range from the twenty-fifth to seventy-fifth percentiles.

H. Tpit motif density at Pax7 sites no ATACseq signal before Pax7 expression

subdivided into Tpit-bound (>1 RPKM in Pax7 expressing cells, blue) and not

bound by Tpit (<1 RPKM in Pax7 expressing cells, grey).

I. Pull-down assay of in vitro translated Tpit interaction with MBP-Pax7 but not with

MBP-βGal.

Figure 7. Stepwise model of pioneer and nonpioneer cooperation for chromatin opening

A. Inactive nucleosomal heterochromatin with binding sites for Pax7 (red) and Tpit

(blue).

B. Weak binding of pioneer factor Pax7 to heterochromatin site.

C. Stabilized Pax7 binding with altered heterochromatin.

D. Recruitment of nonpioneer Tpit through interaction with Pax7 and its DNA binding

site.

E. Accessible chromatin with both factors bound at enhancer.

Methods

Mice, tissues, and cell culture.

Mice genotypes and strains used in this study. Foe single cell RNAseq, a 4 month old male C57Bl/6 mouse pituitary was used; cells were dissociated as described below. For ATACseq, FACS-purified pituitary cells are isolated from 3-5 month old adult LH-cerulean (Budry et al., 2011) or POMC EGFP (Lavoie et al., 2008) C57Bl/6 transgenic pituitaries. ATACseq was performed in duplicates for each genotypes of *Pax7* (Mansouri and Gruss, 1998) and *Tpit* (Pulichino et al., 2003) knockouts. Each replicate used a pool of four dissected intermediate pituitaries from 8-20 days old mice in mixed Balb/c and 129sv backgrounds. FACS analyses of intermediate and anterior pituitaries used 15-20 days-old *Pax7* knockout mice harboring the *POMC-EGFP* transgene in a mixed 129sv and Balb/c background. All animal experimentation was approved by the IRCM Animal Ethics Committee in accordance with Canadian regulations.

Cell lines used in this study. AtT-20 cells (obtained from the late E. Herbert in 1981 and subsequently maintained in our laboratory, with yearly negative mycoplasma tests) and αT3 cells were cultured in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum and antibiotics (penicillin/streptomycin).

Generation of stable Neo, Pax7 and Sox2 expressing cells. Expression vectors constructed in the pLNCX2 vector were described previously (Budry et al., 2012). Retroviruses were packed using the EcoPack 2-293 cells (Clontech, Mountain View, CA) and infections were performed as described (Budry et al., 2012). Selection of retrovirus-infected cell populations was achieved with 400 μg/ml Geneticin (Gibco, 11811-031).

530 Resistant colonies were pooled to generate retrovirus-infected populations of more than

531 1,000 independent colonies.

Pituitary intermediate and anterior lobe cell dissociation

After dissection, mouse pituitary intermediate or anterior lobes were dissociated as

described (Budry et al., 2011). Briefly, dissected pituitaries were separated into

intermediate and anterior lobes and kept during the dissections in 300µl of dissection

buffer (DMEM, 10% FBS, HEPES 10mM and DNase 100 U/ml). Anterior lobes were cut

in pieces using a scalpel to facilitate tissue dissociation and digested at 37°C using

5mg/ml Trypsin for 10 minutes. We then added 2mM EDTA and incubated 5 minutes

more. 10% FBS was then added to stop the dissociation and samples were centrifuged

and then resuspended into 150µl of PBS 1X, 0.1% BSA, 10mM HEPES and for FACS

analyses.

3' end single cell RNAseq

Dissociated pituitary cells were diluted at 500 cells per µl and processed using Chromium

Single Cell 3' v2 Reagent (10x Genomics, Pleasanton, CA) following the manufacturer

recommendation. Briefly, cells were passed on the channel and 9269 cells were

recovered. Pituitary cells were partitioned into gel beads in emulsion for cell lysis and

barcoded with oligo-dT priming and reverse transcribed. cDNA library was amplified

fragmented and size selected. Samples were controlled at multiple steps during the

procedure by running on BioAnalyzer. Libraries were sequenced on Hiseq 4000 with

100bp paired-end reads.

Purification of pituitary lineages by FACS

Dissociated anterior pituitary cells from 62 LH-cerulean mice (Budry et al., 2011) were

sorted using FACSAria instrument (BD) and the gate used to define cerulean positive

versus negative cells were defined by first assessing auto fluorescence of WT mice of the

same strain, C57/Bl6. Cerulean positive and negative cells constituted the gonadotrope

and anterior lobe (AL) samples, respectively, that were used in this study.

FACS analyses

Dissociated anterior or intermediate pituitaries from *WT* or *Pax7-KO* mice crossed with

the POMC-EGFP transgene were analysed using the FACSCalibur cell analyser (BD

Bioscience). EGFP levels were quantified together with forward scatter (FSC) as an

indicator of cell size and side scatter (SSC) as an indicator of granularity for both high

and low EGFP-expressing cells. Experiments were repeated on 3 *Pax7* knockout and 5

wild-type litter mates in a mixed C57/Bl6 and 129sv background.

Immunohistofluorescence

Immunohistofluorescence was performed on PFA-fixed paraffin sections as described

(Bilodeau et al., 2009). Briefly, 5 days old *Tpit* knockout and WT pituitaries were

dissected, fixed in 4% PFA, embedded in paraffin and cut into 5 μm thick sections. The

following antibodies used for immunohistofluorescence: Pax7 (DSHB AB_528428), PC2

(a gift of Dr Nabil Seidah, IRCM, Montreal).

ATACseq

All ATACseq samples were processed as previously described (Mayran et al., 2018).

Briefly, 50 000 cells were washed in PBS and incubated on ice for 30 minute in a

hypotonic cell lysis buffer (0.1% $^w/_v$ sodium citrate tribasic dehydrate and 0.1% $^v/_v$ Triton

X100) and centrifuged (5 minutes at 2000g at 4°C). Cells were then incubated 30 minutes

on ice in cell lysis buffer (10mM Tris-HCl, pH7.4, 10mM NaCl, 3mM MgCl$_2$, 0.1% $^v/_v$

IGEPAL CA-630. After centrifugation (5 minutes at 2000g at 4°C), the nuclei pellets were

resuspended in transposase Master Mix (1.25 µl 10x TD buffer, 5 µl H$_2$O and 6.5 µl of

Tn5: Illumina Nextera Kit; FC-121-1031) and incubated for 30 minutes at 37°C. Samples

were purified using the DCC purification columns (Zymo). The eluted DNA was barcoded

for multiplexing of samples using Nextera barcodes and PCR-enriched using the Phusion

kit. Libraries were recovered with GeneRead Purification columns. Samples were then

evaluated by qPCR to test enrichments and sequenced on Illumina Hiseq 2500 with 50bp

or 125bp paired-end reads according to Illumina's recommendation.

ChIPseq

ChIPseq were performed as previously described (Langlais et al., 2012). At least 3

immuno-precipitations were pooled per ChIP experiments. Library and flow cells were

prepared by the IRCM Molecular Biology Core Facility according to Illumina's

recommendations and sequenced on Illumina Hiseq 2500. The following antibodies were

used for ChIPseq: FlagM2 (Sigma F3165), Neurod1 (Poulin et al., 2000), Sox2 (Ab59776,

Abcam).

Pull-down assay

MBP fusion proteins coupled to maltose amylose beads were produced as described

(Bilodeau et al., 2006). ^{35}S labeled proteins were synthesized *in vitro* using the TNT T7

Quick for PCR DNA kit (Promega, L5540). Labeled proteins were incubated with MBP-

tagged proteins in TNEN$_{50}$ (50mM Tris pH7.5, 5mM EDTA, 50mM Nacl, 0.1% NP-40) with

1mM PMSF and 2% BSA for 4 hours at 4°C. Beads were washed three times with 1ml

TNEN$_{125}$. Bound proteins were resolved by SDS-Page and visualized by autoradiography.

Data Analyses

Single Cell RNAseq analyses. Using Cell Ranger v2.1.1 (10X Genomics), reads were aligned on the mm10 mouse reference genome using the default parameters of Cell Ranger to generate unique molecular identifier counts for each genes across the 9269 cells that were profiled. We obtained an average of 25220 reads per cell and we detected 1807 median genes per cell. Using the Cell Ranger pipeline with default parameters, we generated a gene-barcode matrix, principal component analysis and dimensionality reduction using the t-SNE algorithm. Unbiased clustering of single cells was performed using Cell Ranger which combines K-means clustering and graph-based clustering uncovering 12 clusters. We used Loupe Cell Browser (10X Genomics) to visualize the t-SNE plot with colored cell according to their assigned cluster or colored by gene UMI. Loupe Cell Browser was also used to perform local differential expression analyses of clusters 1, 2, 4, 5, 7, 8. Cluster corresponds to 4 islands of cells that each express a different hormone gene, Gh, Prolactin, Lh and POMC. Differential analysis of the four sub-clusters against their matching hormone-expressing cell cluster showed that mitochondrial RNAs are down-regulated in each case and this is an indicator of low quality cells (Ilicic et al., 2016). Thus to avoid confounding effects, cluster 3 was not included in all further analyses.

ATACseq. Reads were trimmed, if required, to obtain a read length of 50bp and aligned to the mm10 mouse reference genome using Bowtie v2.3.1 (Langmead et al., 2009) with the following parameters: --fr --no-mixed --no-unal. Sam files were converted into tag

directories using HOMER v4.9.1 (Heinz et al., 2010) and into bam files using Samtools

v1.4.1 (Li et al., 2009) view function. Tag directories were used to generate the normalized

BigWig files with Homer using the command makeUCSCfile with the parameters: -fsize

1e20 -res 5 -fragLength 100. Peaks were identified by comparing each sample replicate

to sequenced input DNA from pituitary using MACS v2.1.1.20160309 (Zhang et al., 2008)

callpeak function using the parameters: -f BAMPE --bw 250 -g mm --mfold 10 30 -p 1e-5.

Peaks with an associated pvalue less than 10^{-5} were kept. First we compared ATACseq

profiles of purified pituitary cells: melanotrope (2 replicates), corticotropes (2 replicates)

gonadotropes (1 replicate) and whole AL (1 replicate). Peaks from all datasets from

purified pituitary cells were merge using HOMER v4.9.1 mergePeaks tool to obtain a file

with all unique positions from the ATACseq datasets. This list was clustered by k-means

in 2 clusters for each samples giving the 16 combinations of ATAC clustering as

represented in a heatmap in Figure 2B. Peaks from all datasets from the various

genotypes of *Pax7* and *Tpit* knockout ILs were merged together using HOMER v4.9.1

mergePeaks tool to obtain a file with all unique IL positions from all ATACseq datasets.

ATACseq signals were quantified in these different datasets using the analyzeRepeats.pl

HOMER command and differential accessibility analyses was performed using

getDiffExpression.pl with default parameters which uses Deseq2. Peaks showing a

differential p-value less than 0.05 and a fold change of 2 fold or more were considered

differentially accessible.

ChIPseq. We mapped ChIPseq reads on the mouse genome assembly mm10 by using

Bowtie v1.1.2 with the following settings: bowtie -t -p 4 --trim5 1 --best mm10 –S. Sam

files were converted into tag directories using HOMER v4.9.1 and into bam files using

Samtools v1.4.1 view function. Peaks were identified by comparing each sample to its

control (IP Flag for Pax7, IP IgG for others) using MACS v2.1.1.20160309 callpeak

function using the parameters: --bw 250 -g mm --mfold 10 30 -p 1e-5. Peaks with an

associated p value less than 10^{-5} were kept.

Data presentation. Heatmaps and average profiles were generated using Easeq (Lerdrup

et al., 2016), We used IGV (Thorvaldsdottir et al., 2013) to visualize the BigWig files on

the genome. Principal component analysis , clustered Heatmap associated with

dendrograms from Figures 1D,5C,F were generated using ClustVis (Metsalu and Vilo,

2015).

References

Bilodeau, S., Roussel-Gervais, A., and Drouin, J. (2009). Distinct developmental roles of cell cycle
inhibitors p57Kip2 and p27Kip1 distinguish pituitary progenitor cell cycle exit from cell cycle re-entry of
differentiated cells. Mol Cell Biol *29*, 1895-1908.
Bilodeau, S., Vallette-Kasic, S., Gauthier, Y., Figarella-Branger, D., Brue, T., Berthelet, F., Lacroix, A.,
Batista, D., Stratakis, C., Hanson, J*., et al.* (2006). Role of Brg1 and HDAC2 in GR trans-repression of the
pituitary POMC gene and misexpression in Cushing disease. Genes Dev *20*, 2871-2886.
Bonev, B., Mendelson Cohen, N., Szabo, Q., Fritsch, L., Papadopoulos, G.L., Lubling, Y., Xu, X., Lv, X.,
Hugnot, J.P., Tanay, A*., et al.* (2017). Multiscale 3D Genome Rewiring during Mouse Neural
Development. Cell *171*, 557-572.e524.
Budry, L., Balsalobre, A., Gauthier, Y., Khetchoumian, K., L'honore, A., Vallette, S., Brue, T., Figarella-
Branger, D., Meij, B., and Drouin, J. (2012). The selector gene Pax7 dictates alternate pituitary cell fates
through its pioneer action on chromatin remodeling. Genes Dev *26*, 2299-2310.
Budry, L., Lafont, C., El Yandouzi, T., Chauvet, N., Conéjero, G., Drouin, J., and Mollard, P. (2011). Related
pituitary cell lineages develop into interdigitated 3D cell networks. Proc Natl Acad Sci USA *108*, 12515-
12520.

Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y., and Greenleaf, W.J. (2013). Transposition of native
chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and
nucleosome position. Nature methods *10*, 1213-1218.
Chronis, C., Fiziev, P., Papp, B., Butz, S., Bonora, G., Sabri, S., Ernst, J., and Plath, K. (2017). Cooperative
Binding of Transcription Factors Orchestrates Reprogramming. Cell *168*, 442-459.e420.
Cirillo, L.A., Lin, F.R., Cuesta, I., Friedman, D., Jarnik, M., and Zaret, K.S. (2002). Opening of compacted
chromatin by early developmental transcription factors HNF3 (FoxA) and GATA-4. Mol Cell *9*, 279-289.
Cirillo, L.A., McPherson, C.E., Bossard, P., Stevens, K., Cherian, S., Shim, E.Y., Clark, K.L., Burley, S.K., and
Zaret, K.S. (1998). Binding of the winged-helix transcription factor HNF3 to a linker histone site on the
nucleosome. Embo j *17*, 244-254.
Di Tullio, A., Vu Manh, T.P., Schubert, A., Castellano, G., Mansson, R., and Graf, T. (2011).
CCAAT/enhancer binding protein alpha (C/EBP(alpha))-induced transdifferentiation of pre-B cells into
macrophages involves no overt retrodifferentiation. Proc Natl Acad Sci U S A *108*, 17016-17021.
Donaghey, J., Thakurela, S., Charlton, J., Chen, J.S., Smith, Z.D., Gu, H., Pop, R., Clement, K., Stamenova,
E.K., Karnik, R.*, et al.* (2018). Genetic determinants and epigenetic effects of pioneer-factor occupancy.
Nat Genet.
Drouin, J. (2017). Chap 1 - Pituitary Development. In The Pituitary (Fourth Edition), S. Melmed, ed.
(Academic Press), pp. 3-22.
Fauquier, T., Rizzoti, K., Dattani, M., Lovell-Badge, R., and Robinson, I.C. (2008). SOX2-expressing
progenitor cells generate all of the major cell types in the adult mouse pituitary gland. Proc Natl Acad Sci
USA *105*, 2907-2912.
Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and
Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory
elements required for macrophage and B cell identities. Mol Cell *38*, 576-589.
Hurtado, A., Holmes, K.A., Ross-Innes, C.S., Schmidt, D., and Carroll, J.S. (2011). FOXA1 is a key
determinant of estrogen receptor function and endocrine response. Nat Genet *43*, 27-33.
Ilicic, T., Kim, J.K., Kolodziejczyk, A.A., Bagger, F.O., McCarthy, D.J., Marioni, J.C., and Teichmann, S.A.
(2016). Classification of low quality cells from single-cell RNA-seq data. Genome Biol *17*, 29.
Jacobs, J., Atkins, M., Davie, K., Imrichova, H., Romanelli, L., Christiaens, V., Hulselmans, G., Potier, D.,
Wouters, J., Taskiran, II*, et al.* (2018). The transcription factor Grainy head primes epithelial enhancers
for spatiotemporal activation by displacing nucleosomes. Nat Genet *50*, 1011-1020.
Khetchoumian, K., Balsalobre, A., Mayran, A., Christian, H., Chénard, V., St-Pierre, J., and Drouin, J.
(2018). Cell-autonomous transcriptional mechanism for enhancement of translation capacity in
secretory cells BioRxiv.
Laganiere, J., Deblois, G., Lefebvre, C., Bataille, A.R., Robert, F., and Giguere, V. (2005). From the Cover:
Location analysis of estrogen receptor {alpha} target promoters reveals that FOXA1 defines a domain of
the estrogen response. Proc Natl Acad Sci USA *102*, 11651-11656.
Lamolet, B., Poulin, G., Chu, K., Guillemot, F., Tsai, M.J., and Drouin, J. (2004). Tpit-independent function
of NeuroD1(BETA2) in pituitary corticotroph differentiation. Mol Endocrinol *18*, 995-1003.
Lamolet, B., Pulichino, A.M., Lamonerie, T., Gauthier, Y., Brue, T., Enjalbert, A., and Drouin, J. (2001). A
pituitary cell-restricted T-box factor, Tpit, activates POMC transcription in cooperation with Pitx
homeoproteins. Cell *104*, 849-859.
Langlais, D., Couture, C., and Drouin, J. (2012). The Stat3/GR interaction code: predictive value of
direct/indirect DNA recruitment for transcription outcome. Molecular Cell *47*, 38-49.
Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. Genome Biol *10*, R25.
Lavoie, P.L., Budry, L., Balsalobre, A., and Drouin, J. (2008). Developmental Dependence on NurRE and
EboxNeuro for Expression of Pituitary POMC. Mol Endocrinol *22*, 1647-1657.

Lee, C.S., Friedman, J.R., Fulmer, J.T., and Kaestner, K.H. (2005). The initiation of liver development is
dependent on Foxa transcription factors. Nature *435*, 944-947.
Lerdrup, M., Johansen, J.V., Agrawal-Singh, S., and Hansen, K. (2016). An interactive environment for
agile analysis and visualization of ChIP-sequencing data. Nature structural & molecular biology *23*, 349-
357.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R.
(2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079.
Maaten, v.d. (2008). Visualizing Data using t-SNE. Journal of Machine Learning Research JMLR *9*, 2579-
2605.
Mansouri, A., and Gruss, P. (1998). Pax3 and Pax7 are expressed in commissural neurons and restrict
ventral neuronal identity in the spinal cord. Mech Dev *78*, 171-178.
Mayran, A., Khetchoumian, K., Hariri, F., Pastinen, T., Gauthier, Y., Balsalobre, A., and Drouin, J. (2018).
Pioneer factor Pax7 deploys a stable enhancer repertoire for specification of cell fate. Nat Genet *50*,
259–269
Metsalu, T., and Vilo, J. (2015). ClustVis: a web tool for visualizing clustering of multivariate data using
Principal Component Analysis and heatmap. Nucleic Acids Res *43*, W566-570.
Okita, K., Ichisaka, T., and Yamanaka, S. (2007). Generation of germline-competent induced pluripotent
stem cells. Nature *448*, 313-317.
Pataskar, A., Jung, J., Smialowski, P., Noack, F., Calegari, F., Straub, T., and Tiwari, V.K. (2016). NeuroD1
reprograms chromatin and transcription factor landscapes to induce the neuronal program. Embo j *35*,
24-45.
Poulin, G., Lebel, M., Chamberland, M., Paradis, F.W., and Drouin, J. (2000). Specific protein:protein
interaction between basic Helix-Loop-Helix transcription factors and homeoproteins of the Pitx family.
Mol Cell Biol *20*, 4826-4837.
Poulin, G., Turgeon, B., and Drouin, J. (1997). NeuroD1/BETA2 contributes to cell-specific transcription of
the POMC gene. Mol Cell Biol *17*, 6673-6682.
Pulichino, A.M., Vallette-Kasic, S., Tsai, J.P., Couture, C., Gauthier, Y., and Drouin, J. (2003). Tpit
determines alternate fates during pituitary cell differentiation. Genes Dev *17*, 738-747.
Soufi, A., Garcia, M.F., Jaroszewicz, A., Osman, N., Pellegrini, M., and Zaret, K.S. (2015). Pioneer
transcription factors target partial DNA motifs on nucleosomes to initiate reprogramming. Cell *161*, 555-
568.
Swinstead, E.E., Miranda, T.B., Paakinaho, V., Baek, S., Goldstein, I., Hawkins, M., Karpova, T.S., Ball, D.,
Mazza, D., Lavis, L.D.*, et al.* (2016). Steroid Receptors Reprogram FoxA1 Occupancy through Dynamic
Chromatin Transitions. Cell *165*, 593-605.
Thorvaldsdottir, H., Robinson, J.T., and Mesirov, J.P. (2013). Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. Brief Bioinform *14*, 178-192.
Wapinski, O.L., Lee, Q.Y., Chen, A.C., Li, R., Corces, M.R., Ang, C.E., Treutlein, B., Xiang, C., Baubet, V.,
Suchy, F.P.*, et al.* (2017). Rapid Chromatin Switch in the Direct Reprogramming of Fibroblasts to
Neurons. Cell Rep *20*, 3236-3247.
Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers, R.M.,
Brown, M., Li, W.*, et al.* (2008). Model-based analysis of ChIP-Seq (MACS). Genome Biol *9*, R137.

Figure 1

A

Single cell RNAseq of 9269 pituitary cells

- 1 Somatotropes (Gh)
- 2 Lactotropes (Prl)
- 4 Melanotropes (POMC/Pc2)
- 5 Corticotropes (POMC)
- 7 Stem cells (Sox2)
- 8 Gonadotropes (Lhb)
- 9 Endothelium (Plvap)
- 10 Macrophages (C1qa)
- 11 Posterior lobe (Nkx2.1)
- 12 Pericytes (Dcn)
- 3 } Other
- 6

B

Low UMI High UMI

Pitx1

Pituitary lineage
Non-Pituitary lineage

C

Somatotropes (37%) — *Pappa2*

Corticotrope (9%) — *Crhr1*

Lactotropes (19%) — *Myoc*

Gonadotropes (3%) — *SF1/Nr5a1*

Melanotropes (9%) — *Oacyl*

Stem cells (5%) — *Sox2*

Low UMI High UMI

D

Stem cell genes (Sox2, Sox9)

Lacto genes (Prl, Myoc)
Somato genes (Gh, Ghrhr)

Gonado genes (Lhb, Fshb, SF-1)

Cortico genes (Crhr1, Avpr1b)
POMC genes (Tpit, POMC)
Melano genes (Pcsk2, Pax7)

Stem cells (7) Somato (1) Lacto (2) Gonado (9) Melano (4) Cortico (5)

Figure 2

Figure 3

Figure 4

A

FACS-purified | IL

Cortico   Melano   Gonado   AL | *Pax7*[+/+]   *Pax7*[-/+]   **Pax7**[-/-]

13708 melanotrope accessible sites

B

FACS-purified | IL

Cortico   Melano   Gonado   AL | *Tpit*[+/+]   *Tpit*[-/+]   **Tpit**[-/-]

8672 POMC-lineages accessible sites

C

FACS-purified

Melano   30   20   20
Cortico   30   20   20
Gonado   30   20   20
AL   30   20   20

POMC   *Tnxb*   *Creb3l2*

Pax7   *Tpit*

IL

-/+   -/+   30   20   20
-/-   -/+   30   20   20
-/+   **-/-**   30   20   20

Figure 5

A Tpit-regulated chromatin opening

3451 up
12573 down

Log2FC (*Tpit⁻/⁻ vs Tpit⁻/⁺*)
rLog(mean accessibility)

B Tpit-dependent chromatin opening (log2FC <-2)

| *Pax7* | +/+ | -/+ | -/- | -/+ | -/- |
| *Tpit* | +/+ | -/+ | -/+ | -/- | -/- |

C Tpit-dependent opening (n=16024)

PC2(9%)
PC1(77%)

	Pax7	*Tpit*
WT		
Het		
Tpit KO		
dKO		
Pax7 KO		
▲	+/+	+/+
■	-/+	-/+
●	-/-	-/+
●	-/+	-/-
●	-/-	-/-

D Pax7-regulated chromatin opening

945 up
6112 down

Log2FC (*Pax7⁻/⁻ vs Pax7⁻/⁺*)
rLog(mean accessibility)

E Pax7-dependent chromatin opening (log2FC <-2)

| *Pax7* | +/+ | -/+ | -/- | -/+ | -/- |
| *Tpit* | +/+ | -/+ | -/+ | -/- | -/- |

F Pax7-dependent opening (n=7057)

PC2 (9%)
PC1 (79%)

	Pax7	*Tpit*
Het		
WT		
Tpit KO		
dKO		
Pax7 KO		
▲	+/+	+/+
■	-/+	-/+
●	-/-	-/+
●	-/+	-/-
●	-/-	-/-

G *Pcsk2* TAD

Melano
Cortico
Gonado
AL

Pcsk2

Pax7	*Tpit*
+/-	+/-
-/-	+/-
+/-	-/-
-/-	-/-

H *Pcsk2*

gonado Cortico Melano

I Pax7 PC2

Tpit -/+
Tpit -/-

Figure 6

Figure 7

A

B

Pioneer factor

Weak binding of pioneer factor

C

Pioneer factor

Stabilization of pioneer factor binding

D

Nonpioneer   Pioneer factor

Recruitment of **non-pioneer** factor

E

Nonpioneer   Pioneer factor

Chromatin **opening**.

Legends to Supplementary Figures

Figure S1. Validation of single cell RNAseq data.

(A) Bioanalyzer profile showing size distribution of cDNAs from dissected pituitary cells.
(B) Single cell RNAseq plot showing barcode numbers (x-axis) over the number of UMI per barcode (y-axis). The threshold (colored) used for selecting the 9269 cells analysed.
(C) t-SNE map showing color-coded expression of hormone genes Gh, Lhb, POMC, Prl and Tshb.

Figure S2. Chromatin landscapes in other pituitary lineages.

(A-C) Genome browser views (IGV) of ATACseq profiles from the indicated pituitary lineages at genes marking the identity of somatotropes (A), lactotropes (B) and both (C).
(D) Read density heatmaps showing ATACseq signals at sites of accessibility common to all four pituitary lineages clustered by ATACseq signals in mouse embryonic stem cells (GSE64058).
(E) Genomic distribution of the distances between the indicated category of ATACseq peaks and the closest TSS.
(F) Average profiles of ATACseq signals for the four indicated lineages at cortico- (blue), anterior lobe (purple) and gonado- (orange) specific ATACseq sites.
(G) Motif enrichments (assessed by HOMER) under the indicated category of ATACseq peaks.

Figure S3. FACS analyses of wild-type (WT) and *Pax7-/- POMC-EGFP* pituitary cells.

(A) Representative FACS profiles showing forward (left) and side (right) scatter for cells from wild type and $Pax7^{-/-}$ IL and AL.
(B) Bar graphs showing the ratios of forward (right) and side (right) scatter for EGFP positive cells from wild type (n=5) and $Pax7^{-/-}$ (n=3) IL and AL.

Figure S4. Dispersion plot comparing ATACseq signals in WT and Pax7+/-;Tpit+/- IL cells.

Dispersion plot showing rLog values of accessibility (ATACseq, x-axis) in wild type and $Pax7^{+/-};Tpit^{+/-}$ IL over log2 fold changes in wild type versus $Pax7^{+/-};Tpit^{+/-}$ IL. Red circles identify the differentially accessible regions (p value<0.05, Log2 FC >+/-1).

Figure S5. Q-PCR validation of Pax7 and/or Tpit-dependent accessibility (ATACseq).

(A-C) Genome browser views (IGV) of ATACseq profiles from the indicated genotypes at unaffected (A), Tpit-only dependent (B) and Pax7 dependent (C) sites. Regions amplified in the qPCR measurements of Figure S5D-F are highlighted in yellow.

(D-F) Relative enrichments over a negative region, measured by qPCR of ATACseq libraries at unaffected (D), Tpit-only dependent (E) and Pax7 dependent (F) sites.

(G) Hi-C interaction map (top) from mouse ES cells (Bonev et al., 2017) around the *Drd2 (left)* and *Grik1 (right)* loci showing the boundaries of their respective TADs. Genome browser views (bottom) of the ATACseq profiles in purified pituitary cells and ILs of the indicated genotypes at the corresponding genome location.

Figure S6. Comparison of Pax7 ChIPseq data in AtT-20 and αT3 cells.

Average profiles of Pax7 ChIPseq in AtT-20 (blue) and αT3 (orange) cells at the best 2000 Pax7 peaks from each lineage showing similar enrichments in both ChIPseqs.

Figure S1

A

B

C

Figure S3

A

B

Figure S4

Figure S6

Pax7 ChIPseq

read density

bp from Pax7 peak center

— AtT-20 — aT3