A spatial map in the somatosensory cortex

Xiaoyang Long and Sheng-Jia Zhang*

Department of Neurosurgery, Xinqiao Hospital, Army Medical University,

Chongqing, China

*To whom correspondence should be addressed.

Correspondence and requests for materials should be addressed to:

sheng-jia.zhang@tmmu.edu.cn or sheng-jia.zhang@outlook.com

Abstract

Spatially selective firing in the forms of place cells, grid cells, boundary vector/border cells and head direction cells are the basic building blocks of a canonical spatial navigation system centered on the hippocampal-entorhinal complex. While head direction cells can be found throughout the brain, spatial tuning outside the hippocampal formation are often non-specific or conjunctive to other representations such as a reward. Although the precise mechanism of spatially selective activities is not understood, various studies show sensory inputs (particularly vision) heavily modulate spatial representation in the hippocampal-entorhinal circuit. To better understand the contribution from other sensory inputs in shaping spatial representation in the brain, we recorded from the primary somatosensory cortex in foraging rats. To our surprise, we were able to identify the full complement of spatial activity patterns reported in the hippocampal-entorhinal network, namely, place cells, head direction cells, boundary vector/border cells, grid cells and conjunctive cells. These newly identified somatosensory spatial cell types form a spatial map outside the hippocampal formation and support the hypothesis that location information is necessary for body representation in the somatosensory cortex, and may be analogous to spatially tuned representations in the motor cortex relating to the movement of body parts. Our findings are transformative in our understanding of how spatial information is used and utilized in the brain, as well as functional operations of the somatosensory cortex in the context of rehabilitation with brain-machine interfaces.

Introduction

Spatial representation in the brain has been largely associated with the hippocampus and the entorhinal cortex [1, 2]. Place cells [3] were originally discovered in the hippocampus, with boundary vector/border cells [4, 5], grid cells [6] and conjunctive cells [7] discovered later in the entorhinal cortex. Unlike the location-specific spatial cells in the hippocampal-entorhinal circuit, head direction cells [8] can be found in areas ranging from the diencephalon to thalamic, striatal and cortical regions [9].

Despite the focus on hippocampal-entorhinal contributions to spatial representation, location-dependent firing conjunctive to other sensory/cognitive (e.g. reward) information has been reported in sensory areas [10-12] and cortical regions heavily interconnected with the hippocampal-entorhinal system such as the septum, subicular complex, cingulate cortex, parietal cortex and retrosplenial cortex [13-20]. In rodents, the only extra-parahippocampal region exhibiting apparent location-dependent firing restricted to space alone is the anterior claustrum, where stable, visually-anchored place cell- and border cell-like activities can be recorded [21]. A preliminary report also suggests the dorsal geniculate nucleus may contain place cells similar to those found in the hippocampus [22]. In humans, hexadirectional grid-like signals have been observed in many brain regions outside the hippocampal formation including the posterior cingulate cortex, the medial prefrontal cortex, the retrosplenial cortex, the medial parietal cortex and the frontal cortex [23-26] while grid-cell-like neuronal representations were identified with intracranial electroencephalography recordings on presurgical epilepsy patients and fMRI studies in the human entorhinal cortex [27-29]. Computational modelling also predicted the existence of grid cell-like neurons in all cortical regions throughout the neocortex [30]. These results therefore indicate that spatial tuning is much more distributed into multiple cortical domains beyond the classical hippocampal formation than we previously thought.

The exact physiological mechanism of spatial selectivity in the hippocampal-entorhinal

system is still incompletely understood. Lesions to either area severely affect, but do not abolish spatially selective activities in the other [31, 32]. Past studies have shown manipulation of visual cues [33, 34], olfaction [35] and vibrissae inputs [36] all impact on hippocampal place cell activity, yet none is necessary. Recent theoretical work has suggested the somatosensory area may contain higher-order presentations of the body itself beyond elementary sensory information, of which location presentation is integral in a "body simulation" model of somatosensation [37]. While there are place cell-like activities in the wheelchair-seated monkey sensorimotor cortices [38], it is unclear if these spatial representations are comparable to their freely moving rodent equivalents. Based on the assumption that somatosensation should play a major role in sensory inputs shaping parahippocampal spatial activity [39, 40] and spatial presentation has been theorized to be crucial for body representation [37], we sought to identify spatially selective activities in the primary somatosensory cortex (S1). Remarkably, we were able to detect all spatial cell types reported in the hippocampal-entorhinal system [1, 2] within the S1. We show these spatial cell types within this newly discovered somatosensory spatial representation system share similar properties to their parahippocampal counterparts in terms of stability and response to environmental manipulations [41].

Results

Extracellular recording from the somatosensory cortex

We obtained neurophysiological recordings in freely behaving rats by using tetrodes mounted on movable microdrives. We recorded 2025 putative single units from eight implanted rats across 287 recording sessions. Recordings were obtained mainly from layers IV to VI within the S1 area (**Fig. 1a** and **Fig. S1**), as confirmed by the histological reconstruction. The tetrodes were generally lowered in steps of either 25 μm or 50 μm per day. The recording sessions lasted between 10 to 30 minutes to ensure sufficient coverage of the whole arena, which was facilitated by foraging food pellets intermittently thrown into the arena. Spike sorting was performed manually offline

using TINT (Axona, St. Albans, U.K.). The separation of the clusters and waveforms

was quantitatively measured with an isolation distance and the L-ratio [42] (**Fig. S2**).

We applied the same analysis described in previous work [7, 17, 43] on 2025 somatosensory

cells and 392 met the criteria for various types (e.g. place, head direction, boundary

vector/border and grid cells) of spatial tuning (**Fig. S3**). The number of each

functionally distinct somatosensory spatial cell type from individual implanted animals

was summarized in the supplementary **Table S1.** No significant differences were

observed among spatial information, mean vector length and grid score across different

animals (ANOVA test, $P = 0.38$, 0.08 and 0.18, respectively, **Figs. S4a, b** and **d**). Both

border score and grid spacing remained similar across interindividual animals except

that they are slightly different in one of eight animals (ANOVA test, * $P < 0.05$, **Figs.**

S4c-d). The distribution of different spatial cell types across different animals suggested,

at least between layers IV and VI, there was a tendency for head direction cells to avoid

layer VI and the presence of spatially tuned somatosensory cells was sparse in

superficial layer V (**Figs. S3b-f**). Place cells appeared to predominate deeper layers

(**Figs. S3a-c**).

Place cells in the somatosensory cortex

As illustrated in **Fig. 1**, somatosensory cells possessing place cell-like properties were

the most abundant spatial cell type encountered (**Figs. S3a-c**). Spatially selective firing

patterns similar to those originally described for hippocampal place cells [3] can be

observed (**Fig. 1b** and **Fig. S5**). These S1 place cells were classified if their spatial

information (SI) score [44] exceeds a stringent 99th percentile threshold (SI > 1.50 cut-off)

from the population-based distribution of shuffled rate maps (**Fig. 1d**). 195 out of the

2025 S1 neurons (9.63%) met criterion and thus are referred to as "place cells" from

here on. These place cells also passed the spatial threshold for the within-cell shuffling

(**Fig. 1c** and **Fig. S6a**), which is less stringent than the population shuffling. The within-

cell shuffling outcome is also true of other types of somatosensory spatial cells (**Figs.**

S6b-d). Accordingly, we used the population shuffling for defining all different types

of somatosensory spatial cells throughout this study. Furthermore, we applied the

maximum-likelihood approach [45, 46] to evaluate inhomogeneous sampling biases and

found that somatosensory place cells showed a robust locational effect, which was not

altered by the correction algorithm (**Figs. S7a-c**). The average SI in bits per spike of all

195 classified S1 place cells was 2.19 ± 0.04 (mean $\pm$ s.e.m., all data are shown as mean

$\pm$ s.e.m. unless otherwise indicated). This percentage was significantly higher than

expected by random selection from the entire shuffled population (**Fig. 1d**; $Z = 39.03$,

$P < 0.001$; binomial test with expected P_0 of 0.01 among large samples).

We noted that if a 95th percentile criterion was applied, 511 out of the 2025 S1 neurons

(24.00%) would be classified as place cells (average SI of 1.59 ± 0.03; cut-off = .81).

This percentage was substantially higher than expected with a random selection from

the entire shuffled population (**Fig. 1d**; $Z = 109.61$, $P < 0.001$; binomial test with

expected P_0 of 0.01 among large samples). We found that the percentage of identified

S1 place cells is lower within the somatosensory cortex than that of hippocampal place

cells [47]. The average peak firing rate and the mean firing rate of identified

somatosensory location-specific place cells were 10.32 ± 0.37 Hz and 0.70 ± 0.04 Hz

(**Fig. S8a**), respectively. The histograms of peak-to-peak amplitude and peak-to-trough

spike width were shown in **Fig. S8e**. Like a mixture of a single place field and multiple

place fields identified from the hippocampal place cells [48], we found that the proportion

of multi-field place cells was 25.64%. The average somatosensory spatial firing field

size was 891.82 ± 32.65 cm^2 (**Fig. 1f**). To quantify the smoothness of recorded

somatosensory place fields, we calculated the spatial coherence by computing the mean

correlation between the firing rate of each bin and the averaged firing rate of the 8

neighboring bins from unsmoothed spatial firing rate maps [34]. For 195 recorded

somatosensory place cells, the average spatial coherence was 0.61 ± 0.01 (**Fig. 1g**).

Furthermore, we also calculated spatial sparsity to evaluate the extent of spatial

information in each spike discharged by S1 place cells [44]. The spatial sparsity measure

of S1 place cells was 0.12 ± 0.01 (**Fig. 1h**). Besides, S1 place cells can be found at any

position within the arena for the uniformity testing based on the Friedman-Rafsky's

MST test (**Fig. 1e,** Friedman-Rafsky's MST test, $P = 0.42$, two-tailed z-test) as in the

hippocampus [49].

Place cell activity was defined not only by its location-specific firing but also by the

stability of its spatial representation. To this end, we next evaluated the spatial stability

of somatosensory place fields by computing the spatial correlations between single

intra-trial behavioral sessions. The average spatial correlation for S1 place cells was

0.64 ± 0.01 (**Fig. S9d**) between the two halves of a recording session (**Figs. S9a-c**),

which remain stable in extended recording sessions up to 30 min (**Figs. S10a-c**).

Hippocampal place cell activities were known to be heavily modulated by visual cues

[33, 34], and were known to rotate with the positioning of visual cues [34]. We performed

similar experiments to test if S1 place cell activities were modulated by visual cues by

moving it 90° from its original position and back. Remarkably, the relative position of

S1 place fields did indeed rotate with the placement of the visual cue and remained

stable after the cue was moved back to its original position (**Figs. S11a-c**). Furthermore,

as hippocampal place codes were reorganized to remap under different geometric

environments [50, 51], we also found that S1 place fields remapped between a square box

and a circular box in the same recording room with changed firing rate and firing

location (**Figs. S12a-c**).

To measure the animal's running speed within and outside the firing fields of S1 place

cells, we calculated the average "in-field" and "out-field" running speed (> 2.5 cm/s)

for 195 identified place cells in S1. We quantified the average running speed across all

the spatial bins within the firing field of the place cell ("in-field" running speed) as well

as outside the firing field of the place cell ("out-field" running speed) (**Fig. S13a**). We

found that there was no significant difference between the average "in-field" and "out-

field" running speed (14.58 ± 0.25 cm/s vs 15.00 ± 0.18 cm/s, $n = 195$, $P = 0.10$, two-

tailed paired t-test, **Fig. S13b**). To evaluate the effect of the animal's running speed on

spatial responses, we calculated the spatial information scores of S1 place cells for

running speed above 2.5 cm/s and below 2.5 cm/s, respectively. We found that the place

fields during inactive mobility are less prevalent with lower spatial information score

than those for active exploring (**Figs. S14a-c**). Similar results were observed for S1

border cells (**Figs. S27a-c**) and S1 grid cells (**Figs. S30a-c**). As for S1 head direction

cells, it appeared that head directional responses were less influenced with immobility

than other three S1 spatial cell types (**Figs. S21a-c**).

Fig. 1 Place cells in the somatosensory cortex.

(a) A Nissl-stained coronal section (upper) showing tetrode tracking trajectory

(arrowheads) through all of six layers across the rat primary somatosensory cortex.

Broken dashes depict the boundaries of the hindlimb (S1HL) and shoulder (S1Sh)

regions of the primary somatosensory cortex (S1). The bottom panel shows the

schematic delimitation of the different layers and sub-regions of the primary

somatosensory cortex. Scale bar, 1 mm. **(b)** Trajectory (grey line) with superimposed

spike locations (red dots) (left column); heat maps of spatial firing rate (middle left

column) and autocorrelation (middle right column) are color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and spatial information (si) for each representative cell are labelled at the top of the panels. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 μV, 300 μs. (**c**) Distribution of within-cell shuffled spatial information for three representative somatosensory place cells. The orange and blue stippled lines mark the 99[th] and the 95[th] percentile significance level of each randomly shuffled distribution, respectively. The red line indicates the observed spatial information. (**d**) Distribution of spatial information for all isolated somatosensory units. The top panel shows the distribution for observed spatial information. The bottom panel shows the distribution for randomly shuffled data from the same sample. The orange and blue stippled lines mark the 99[th] and the 95[th] percentile significance level of each randomly shuffled distribution, respectively. A zoomed panel shows the magnification of the specified area marked by the red dashed rectangle. (**e**) The uniformly areal distribution of various place firing fields relative to the field center. (**f-h**) The population histograms of place firing field sizes (**f**), spatial coherence (**g**) and spatial sparsity (**h**) for all of 195 identified somatosensory place cells.

Head direction cells in the somatosensory cortex

Head direction cells discharge only when the animal's head is at a particular angle respective to the environment [8]. To classify the preferred firing angle of each cell, the mean vector length was used to compute head direction tuning [7]. Cells were categorized as head direction cells if the mean vector length exceeded the 99[th] percentile of the total distribution of the shuffling procedure from the entire pool of putative single cells (**Fig. 2c**). 80 cells (3.95% of all recorded cells) met the criterion, showing significant modulation with the animal's head direction (**Fig. 2a** and **Fig. S15**), with the average mean vector length found to be 0.53 ± 0.01 and the mean firing rate to be 0.96 ± 0.11 (**Fig. S8b**). The histograms of peak-to-peak amplitude and peak-to-trough spike width were displayed in **Fig. S8f**. Besides, the within-cell shuffling for those three

representative somatosensory head direction cells generated similar results to the population shuffling (**Fig. 2b**). This percentage is significantly higher than expected by random selection from the entire shuffled population (**Fig. 2c**; $Z = 13.34$, $P < 0.001$; binomial test with expected P_0 of 0.01 among large samples). Notably, 352/2025 identified S1 neurons (17.38%) fulfilled the lower 95th percentile criterion with the threshold of mean vector length being 0.26. To correct for the possible effects of direction and location for identified head direction cells, we applied the maximum-likelihood correction approach [45, 46] and found that the somatosensory head directional responses remain unaltered from the corrected directional firing rate histograms (**Figs. S16a-c**).

To measure the sharpness of the head directional tuning curve, we quantified the average full width at half maximum (FWHM) of somatosensory head direction cells. The average FWHM of the angular distributions was 77.58 ± 4.36 degrees (mean $\pm$ s.e.m.) (**Fig. 2d**). The distribution of head direction preferences was uniformly distributed across $360°$ (Rayleigh test for uniformity: $P = 0.17$, **Fig. 2e**). We next evaluated the intra-trial directional stability by correlating the distributed firing rate across all directional bins between the first and second halves of individual recording trials. We found head direction related firing to be stable within sessions (**Figs. S17a-c**), with the average angular correlation coefficient for S1 head direction cells being 0.58 ± 0.02 (**Fig. S17d**). Like hippocampal place cells, head direction cells also anchored to salient visual cues [52]. We further tested the specificity and stability of S1 head direction cells by rotating a salient visual cue $90°$ from its original position in the enclosure and back (**Fig. S18**, as in S1 place cells described above). As with S1 place cells, S1 head direction cells also rotated with the movement of the salient visual cue (**Figs. S18a-c**).

To measure the difference of the animal's angular velocity within and outside preferred firing directions for those identified head direction cells in S1, we calculated the average angular velocity in the "preferred firing directions" and "non-preferred firing

directions" for 80 identified head direction cells in S1 (**Fig. S19a**). We found that there was no significant difference between the average angular velocity in the "preferred firing directions" and "non-preferred firing directions" for head direction cells in S1 (52.97 ± 1.15 degrees/s vs 55.52 ± 1.34 degrees/s, $n = 80$, $P = 0.12$, two-tailed paired *t*-test, **Fig. S19b**).

Although the previous study had shown that food chasing had little effect on head directional firing in the rat anterior thalamic nuclei [53], we evaluated the firing patterns of head direction cells in the S1 without food pellet dispersion. We recorded the same head direction cell in two consecutive secessions with (bottom panels in **Fig. S20a**) and without (upper panels in **Fig. S20a**) food pellets, respectively. No significant differences were observed to suggest that food chasing did not modulate preferred firing angles in S1 head direction cells. Similar results were observed for place cells in the S1 without throwing food pellets within the running arena (upper panels in **Fig. S20b**).

Fig. 2 Head direction cells recorded from the somatosensory cortex.

(a) Three examples of somatosensory head direction cells. Trajectory (grey line) with superimposed spike locations (red dots) (left column); spatial firing rate maps (middle left column), autocorrelation diagrams (middle right column) and head direction tuning curves (black) plotted against dwell-time polar plot (grey) (right column). Firing rate is color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr), mean vector length (mvl) and angular peak rate for each representative head direction cell are labelled at the top of the panels. The directional plots show strong head direction tuning. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs. (**b**)

Distribution of within-cell shuffled mean vector length for three representative somatosensory head direction cells. The orange and blue stippled lines mark the 99[th] and the 95[th] percentile significance level of each randomly shuffled distribution, respectively. The purple line indicates the observed mean vector length. (**c**) Distribution of mean vector length for all identified somatosensory units. The top panel shows the distribution for observed values. The bottom panel shows the distribution for randomly shuffled data from the same sample. The orange and blue stippled lines mark the 99[th] and the 95[th] percentile significance level of each randomly shuffled distribution, respectively. A zoomed panel shows the magnification of the specified area marked by the purple dashed rectangle. (**d**) Distribution of head directional tuning width. (**e**) Preferred direction of recorded head direction cells from S1. The polar plot shows the distribution of the peak firing direction of all identified somatosensory head direction cells. The preferred head direction exhibits a uniform distribution ($P = 0.17$, Rayleigh's test).

Border/Boundary vector coding in the somatosensory cortex

Boundary vector cells and border cells discharge exclusively whenever an animal is physically close, at a specific distance and direction, to one or several environmental boundaries, for example, the enclosure walls of the recording arena [4, 5, 54]. S1 cells were defined as border cells if border scores (see Methods) were larger than the 99[th] percentile of population shuffled scores (**Fig. 3c**). A total of 86/2025 (4.25%) S1 cells were classified as border cells, and this percentage was significantly higher than expected by random selection from the entire shuffled population (**Fig. 3c**; $Z = 14.68$, $P < 0.001$; binomial test with expected P_0 of 0.01 among large samples). Similar results were observed by performing cell-specific shuffling for those three representative somatosensory border cells (**Fig. 3b**). Most border cells ($n = 58$, 67.44%) fired along a single boundary while others were active along with two ($n = 15$, 17.44%), three ($n = 6$, 6.98%) or even four ($n = 7$, 8.14%) boundaries within the enclosure (**Fig. 3a** and **Fig. S22**). The average firing rate and the border score were 1.00 ± 0.09 Hz (**Fig. S8c**) and 0.66 ± 0.01, respectively. The histograms of peak-to-peak amplitude and peak-to-trough

spike width were summarized in **Fig. S8g**. The number of S1 border cells seemed to be

higher in layer V than in layer IV or layer VI (**Fig. S3b** and **Fig. S3e**). At a lower 95th

percentile criterion, 234/2025 (11.56%) somatosensory cells were classified as border

cells.

Qualitatively, S1 border cells maintained their border-specific firing fields between

circular and square enclosures (**Figs. S23a-c**), comparable to boundary vector cells and

border cells from the presubiculum, parasubiculum and the medial entorhinal cortex

(MEC) [4, 5, 17, 54]. S1 border cells also responded to a new wall insert to a familiar

enclosure (**Figs. S24a-c**), as described in the MEC [5, 54]. As observed in the MEC and

the subiculum [4, 5], removal of walls from the enclosure on an elevated platform

preserved similar "border" specificity to that in the walled arena [54]. These results

confirmed that somatosensory border cells continued to code for geometric boundaries

instead of physical borders in the unwalled environments (**Figs. S25a-c**). Interestingly,

we also detected a certain number of boundary vector cells [4, 54] which discharged further

away from the edge of the enclosure (**Fig. S26**). Further quantitative characterization

of somatosensory boundary vector cells combined with the previously described

"perimeter" coding might reveal how somatosensory boundary vector cell closely

resemble their counterparts in the subiculum [1, 4, 54].

311

312 **Fig. 3 Border cells recorded from the somatosensory cortex.**

313 **(a)** Three representative examples of somatosensory border cells. Trajectory (grey line)

314 with superimposed spike locations (red dots) (left column); heat maps of spatial firing

315 rate (middle column) and autocorrelation diagrams (right column). Firing rate is color-

316 coded with dark blue indicating minimal firing rate and dark red indicating maximal

317 firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate

318 maps. Peak firing rate (fr), mean firing rate (fr) and border score (b) for each

319 representative border cell are labelled at the top of the panels. Spike waveforms on four

320 electrodes are shown on the right column. Scale bar, 150 µV, 300 µs. **(b)** Distribution

321 of within-cell shuffled border score for three representative somatosensory border cells.

The orange and blue stippled lines mark the 99[th] and the 95[th] percentile significance

level of each randomly shuffled distribution, respectively. The green line indicates the

observed border score. (**c**) Distribution of border scores for pooled somatosensory cells.

The top panel shows the distribution for observed values. The bottom panel shows the

distribution for randomly shuffled versions from the same sample. The orange and blue

stippled lines mark the 99[th] and the 95[th] percentile significance level of each randomly

shuffled distribution, respectively. A zoomed panel shows the magnification of the

specified area marked by the green dashed rectangle.

Grid cells in the somatosensory cortex

Hexagonal firing patterns identified in the MEC [6] can also be observed in the

somatosensory cortex. To quantify the regularity of hexagonal firing patterns of the

somatosensory cells (**Fig. 4a** and **Fig. S28**), we calculated grid scores based on rotated

autocorrelograms of rate maps. A cell was categorized as a grid cell if its grid score was

higher than the 99[th] percentile of the shuffled data. 72/2025 cells (3.55%) met this

criterion (**Fig. 4d**), and this percentage was significantly higher than expected by

random selection from the entire shuffled population (**Fig. 4d**; $Z = 12.90$, $P < 0.001$;

binomial test with expected P_0 of 0.01 among large samples). Similar results were

observed by performing within-cell shuffling (**Fig. 4b**). To eliminate the possible

influence of inhomogeneous sampling on the multiple firing fields of somatosensory

grid cells and verify S1 grid cells through spike shuffling, we performed additional

"field shuffling" [22] and found that these exampled grid cells also passed the gridness

threshold for the "field shuffling' procedure (**Fig. 4c**). The average firing rate and the

grid score were 1.18 ± 0.12 Hz (**Fig. S8d**) and 0.55 ± 0.02 (**Fig. S4d**), respectively. The

histograms of peak-to-peak amplitude and peak-to-trough spike width were presented

in **Fig. S8h**.

Grid spacing in somatosensory grid cells (**Fig. 4e**) was 42.72 ± 1.30 cm, and the average

grid size was 842.44 ± 54.22 cm^2 (**Fig. 4f**). The fluctuation in somatosensory grid

spacings is not a result of interindividual differences across different animals (**Fig. S4d**).

Moreover, the orientation of the somatosensory grid cells varied and was on the average found to be 32.63 ± 2.64 degrees (**Fig. 4g**). Again, with a lower threshold at the 95[th] percentile, 218/2025 S1 neurons (10.37%) were classified as grid cells, with the grid score threshold of 0.17 (blue line in **Fig. 4d**). When recorded in a larger environment (1.5 m x 1.5 m box), the discharging patterns of somatosensory grid cells showed similar multiple regular triangular structures in a larger enclosure to those within the smaller enclosure (1.0 m x 1.0 m) with a high grid score (**Figs. S29a-c**), confirming that those identified grid cells were not false-positive grid cells due to accidental triangular node structures. It appeared that somatosensory grid cells were less prevalent and a little noisier than place cells, head direction cells and grid cells identified in the S1. It remains to be determined whether multisensory inputs and intrinsic oscillatory dynamics within the somatosensory cortex might generate different rate-coded grid computations from those within the hippocampal-entorhinal cortex. Taken together, the local configuration of grid cells in the somatosensory cortex exhibited similar features as those in MEC. Furthermore, we detected irregular grid cells with negative grid scores in the S1 along with regular grid cells (**Fig. 4h** and **Fig. S31**).

Fig. 4 Grid cells in the somatosensory cortex.

(**a**) Three representative examples of somatosensory grid cells. Trajectory (grey line) with superimposed spike locations (red dots) (left column); heat maps of spatial firing rate (middle column) and autocorrelation diagrams (right column). Firing rate is color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and grid score (g) for each representative grid cell are labelled at the top of the panels. A crystal-like hexagonal firing pattern was observed. Spike waveforms on four electrodes are shown on the right column. Scale bar, 100 µV, 300 µs. (**b**) Distribution of within-cell shuffled grid score for three representative somatosensory grid cells. The orange and blue stippled lines mark the 99th and the 95th percentile significance level of each randomly shuffled distribution, respectively. The cyan line indicates the observed grid score. (**c**) The same as (**b**) but for field shuffle. (**d**) Distribution of grid scores for somatosensory cells. The top panel shows the distribution for observed values. The bottom panel shows the distribution for randomly shuffled versions from the same sample. The orange and blue stippled lines mark the 99th and the 95th percentile significance level of each randomly shuffled distribution, respectively. A zoomed panel shows the magnification of the specified area marked by the cyan dashed rectangle. (**e-f**) The raster plots show the distribution of grid spacing and grid size of the classified somatosensory grid cells. (**g**) The histogram of grid orientation from categorized somatosensory grid cells. (**h**) Three representative examples of irregular somatosensory grid cells. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Conjunctive cells in the somatosensory cortex

To establish the existence of S1 neurons coding more than one spatial correlate, we assessed the strength of head direction tuning for all somatosensory place cells, border cells and grid cells [7]. A certain number of somatosensory neurons were found to be conjunctive cells (**Fig. 5** and **Fig. S3a**). Specifically, 25/195 place cells (**Fig. 5a**), 11/86 border cells (**Fig. 5b**) and 5/72 grid cells (**Fig. 5c**) were significantly tuned by head

direction. Interestingly, we also detected irregular hexagonal grid cells modulated with

head direction (**Fig. 5d**).

The degree of directionality in conjunctive cells showed no obvious difference from

that of pure head direction cells (Watson's U^2 test, $P = 0.09$). These results suggested

that the somatosensory cortex carried similar conjunctive spatial x directional, border

x directional and gird x directional signal to those identified previously in the rat

subiculum [45] and the rat MEC [5, 7].

Fig. 5 Four different types of conjunctive cells in the somatosensory cortex.

(**a**) A representative conjunctive place x head direction cell. (**b**) A representative

conjunctive border x head direction cell. (**c**) A representative conjunctive grid x head

direction cell. (**d**) A representative conjunctive irregular grid x head direction cell.

Trajectory (grey line) with superimposed spike locations (red dots) (left column); heat

maps of spatial firing rate (middle left column), autocorrelation diagrams (middle right

column) and head direction tuning curves (black) plotted against dwell-time polar plot

(grey) (right column). Firing rate is color-coded with dark blue indicating minimal

firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation

maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate

(fr), spatial information (si), border score (b), grid score (g) and mean vector length

(mvl) for each representative cell are labelled at the top of the panels. Spike waveforms

on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Impact of the whisker trimming on the somatosensory spatial activities

Since rodents use their whiskers to actively explore the external environment for

whisker-based navigation [55, 56], we implanted two additional rats (**Fig. S1b**) and

recorded neuronal activity before and after repeated whisker trimming (**Fig. S32a** and

Fig. S32c) to gauge the involvement of the vibrissae. Two consecutive recording

sessions from the same rat before and right after whisker trimming showed that the

distribution of clusters and waveforms in the S1 was qualitatively similar (**Fig. S32b**

and **Fig. S32d**).

To further quantify the change of well-isolated spike clusters before and after whisker

trimming, we calculated the isolation distance and L-ratio [42]. The median isolation

distances were 122.35 and 132.9 before and after whisker trimming, respectively. There

was no significant difference in cluster separation and L-ratio in the S1 before and right

after whisker trimming (**Fig. S32e** and **Fig. S32f**, $n = 78$, $P = 0.73$ and 0.76,

respectively, two-tailed paired t-test). The average drift of center of mass for the total

cell sample before and after whisker trimming was 0.13 ± 0.01, indicating there was

minimal change in spike clusters before and after whisker trimming. Besides, both peak

and average firing rates of the same neurons before and after whisker trimming did not

change significantly (**Fig. S32g** and **Fig. S32h**, $n = 78$, $P = 0.75$ and 0.47, respectively,

two-tailed paired t-test).

To further assess whether whisker trimming affected the S1 spatial firing properties, we

were able to continuously record the same S1 border cell before and after whisker

trimming. The spatial firing patterns of the border cell persisted and responded to a new wall inserted into the square enclosure (**Fig. S33a** and **Fig. S33b**). Overall, both the spatial responses and firing properties in the S1 were not significantly affected by the whisker trimming.

When we pooled and analyzed a total of 461 cells recorded after whisker trimming, we were still able to identify all different spatial cell types including 43 place cells (**Fig. 6a**), 15 head direction cells (**Fig. 6b**), 29 border cells (**Fig. 6c**) and 17 grid cells (**Fig. 6d**). The percentage of all recorded spatially tuned cell types after whisker trimming were similar to those recorded from rats without whisker trimming. Furthermore, the cutoffs for defining place cells (**Fig. 6e**), head direction cells (**Fig. 6f**), border cells (**Fig. 6g**) and grid cells (**Fig. 6h**) after whisker trimming highly resembled those for different somatosensory spatial cell types without whisker trimming. Taken together, the persistence of recorded somatosensory spatial cells after whisker trimming demonstrated that our identified cells were not significantly influenced by the whisker trimming.

Fig. 6 Somatosensory spatial response after the whisker trimming.

(**a**-**d**) Four representative examples of somatosensory place cell (**a**), head direction cell (**b**), border cell (**c**) and grid cell (**d**) after whisker trimming. Trajectory (grey line) with superimposed spike locations (red dots) (left column); spatial firing rate maps (middle left column), autocorrelation diagrams (middle right column) and head direction tuning curves (black) plotted against dwell-time polar plot (grey) (right column). Firing rate is color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr), spatial information (si), mean vector length (mvl), border score (b) or grid score (g) and angular peak rate for each representative cell are labelled at the top of the panels. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs. (**e**-**h**) Distribution of spatial information (**e**), mean vector length (**f**), border score (**g**) or grid score (**h**) for somatosensory cells after whisker trimming. The top panel shows the distribution for observed values. The bottom panel shows the distribution for randomly shuffled data from the same sample. The orange and blue stippled lines mark the 99[th] and the 95[th] percentile significance level of each randomly shuffled distribution, respectively.

Discussion

This study provides the first demonstration and characterization of spatially tuned neuronal discharges in the primary somatosensory cortex. We have shown place, grid, boundary vector/border, head direction and conjunctive cell activities, as reported in parahippocampal and associated areas [39], to co-exist in the single-domain primary somatosensory cortex. These spatially tuned activities encoded within the newly identified somatosensory spatial navigation system appear to be specific, stable, anchored to external visual cues and robust against non-spatial environmental perturbations.

Based on the assumption that proprioception makes a significant contribution to spatial representation in the parahippocampal cortices [39, 40], we predicted spatially selective activities should exist in the somatosensory cortex. To our surprise, we were able to

record the full complement of spatially selective cells (place, grid, boundary

vector/border, head direction and conjunctive cells) in the primary somatosensory

cortex. Given the paucity of direct projections from the parahippocampal areas to the

S1 region [57, 58], we assume somatosensory spatial activities do not arise from the

hippocampal-entorhinal circuit. However, we cannot completely eliminate the

possibility that the somatosensory spatial positioning system may be at least partially

dependent on the hippocampal-entorhinal system. Whether the newly found

somatosensory spatial navigation system is independent of, parallel with or convergent

onto the classical one within the hippocampal-entorhinal microcircuit remains an

interesting question and needs to be addressed in future studies [1]. The availability of

single-cell transcriptomics and connectomics for the somatosensory cortex might help

to unravel the interaction or interdependence between these two genetically pre-

configured spatial maps [59, 60]. Since there is very little overlap between septal

projections to the hippocampal-entorhinal areas [61], and that septal inactivation severely

attenuates place and grid cell activities in both regions [62-64], how S1 responds to septal

inactivation may provide the necessary insight for hippocampal-entorhinal dependency.

The laminar organization of the S1 in relation to connectivity and function has been

well-characterized, largely arising from studies on the vibrissae barrel fields [65, 66]. There

is insufficient information on the specific connectivity patterns of S1 in the rat, but

mouse data suggest similar cortico-cortical, thalamic-cortical, cortical-thalamic,

cortico-striatal and corticofugal connection patterns between S1 and other parts of the

S1 [67]. Our current study is roughly limited to sampling layers IV through VI. There is a

pattern for head direction cells to be absent in layer VI and place cells to predominate

in deeper layers (**Figs. S3e-h**). From such distribution of spatial cell types across

cortical depth, we could infer that head direction information is not present in cortico-

thalamic communications, given the lack of head direction cells at depths

corresponding to layer VI in the somatosensory cortex. There also appears to be a gap

in the most superficial aspect of layer V (Va) where all spatial cell types appear to be

under-represented. Interestingly, layer Va projections are mostly recurrent [68] and appear

to serve as a trans-columnar and trans-laminar integration hub within the S1 [69-71]. Layer

Va may serve as an area that integrates spatial and proprioceptive information in the S1,

hence salient spatial activity is not readily observable in this layer. If true, this would

also suggest that the major extra-cortical output from S1 combines, but do not contain

conventional spatial information as characterized here. A full sampling of cortical

layers in S1 using high-density linear probes [72] is necessary to address how spatial

information is generated, integrated and distributed elsewhere (e.g. thalamic, striatal

and brainstem targets) in the brain.

As mentioned above, there is a lack of direct connections between S1 and areas that are

traditionally associated with spatial processing. Where do the spatial signals come from?

Are they more universal in cortical processing than previously thought? What is the

functional significance of spatial activities in the S1, or S1 in general? While extensive

intra and inter columnar/laminar exist in the S1 and may support *de novo* generation of

place cells (from integrating local grid cell activities) and conjunctive cells, it is

unlikely the full complement of known spatial representations are generated

independently within the S1. We have demonstrated that S1 place fields and preferred

head direction rotate with the varied positioning of salient visual cue as cells found in

the hippocampal-entorhinal circuit. This observation suggests the spatial activities

cannot solely be driven by proprioception arising from lemniscal/paralemniscal

pathways. While a sparse, direct and functional connection exists between the S1 and

the visual cortex [73], it is unlikely to account for visually anchored spatial tuning in the

somatosensory cortex. Instead, S1 spatial activities are more likely to be an efference

copy from elsewhere in the brain.

S1 functions and activities are intricately linked to motor structures, especially the

motor cortex. Functional coupling between S1 spindle activity and spontaneous muscle

twitches can be observed as early as two days after birth [74]. It is known that both primary

and secondary motor cortex are reciprocally connected with S1 [75, 76], and the

preparation, execution and dynamic changes in movements can all be predicted from

M1/S1 activities [77]. In fact, S1 has recently been shown to have an active role in directly

controlling motor output [78]. Position-dependent neuronal discharges (i.e. place field-

like representations) relating to hand movement through space are well-established in

the primate motor cortex [79]. In fact, considering movement of the hand or forelimb in

general: speed [79, 80], (hand) direction [79] and conjunctive [79, 80] coding in the motor cortices

can be seen as homologous to their spatial navigation counterparts in the limbic system.

Essentially, it appears the motor cortex contains spatially tuned activities for specific

body parts that are analogous to limbic activities that appear to represent whole

body/head movement. Therefore, whole body/head and limb movements through space

may require similar neurocomputation, of which is also present in the S1 in addition to

motor and limbic cortices. This proposal is also consistent with the idea that S1

simulates the body itself in addition to its representation, which requires spatial

information [37]. From this synthesis, we speculate that we would be able to record

spatially tuned activities in the limb region of the motor cortex, as well as other

sensorimotor cortices (e.g. the forelimb and trunk regions). Particularly, inactivation of

the reciprocally connected motor cortex (i.e. limb region of the motor cortex) can test

if S1 spatial activities are efference copy from the motor cortex. Recent reports of place

cell-like activities in the primate sensorimotor cortices [38] during quasi-active

movements through space support the notion that spatial representation of body

movement may exist throughout the cortex.

Our discussion so far favors somatosensory spatial activity may be an efferent copy

from the motor cortex. If true, then the question remains as to how an area such as the

motor cortex without direct connectivity with spatially tuned limbic structures (like the

S1) can acquire spatial activities. To the best of our knowledge, no grid- or boundary

vector-/border-like spatial representation has been reported in the motor cortex in

conventional reaching studies, although this discrepancy may be related to that motor

tasks in such studies are often limited to a single defined trajectory (insufficient spatial

coverage to detect grid cells) in open task space (without physical borders). Likewise,

the "body simulation" model [37] predicts the existence of body part location information

(which we interpret to be place cell-like representations) to exist in layer IV, but our data indicate the full complement of spatial activities can be detected from at least layers IV through VI. Our data is not fully compatible with currently available experimental and theoretical work in explaining why all currently recognized spatial cell activities can be found in S1 and how they are generated. Regardless, we propose it is likely spatial information is integral to the "body simulation" function in S1 as put forward by Brecht [37]. High-resolution kinematic studies [81] of body part (e.g. limb) movements through space would be required to test if spatial activities in the S1 do in fact represent body parts in space.

In this report, we have demonstrated place, grid, boundary vector/border, head direction and conjunctive cell activities in the primary somatosensory cortex. These activities are comparable to their counterparts in the parahippocampal cortices and are stable, specific, robust to non-spatial manipulations and anchored to salient spatial cues. We propose spatial tuning in the primary somatosensory cortex may be crucial for body representation/simulation in space, and the newly identified somatosensory spatial representation system may generalize to other sensorimotor cortices pertinent to behavioral coordination.

Materials and Methods

Subjects

Ten male Long-Evans rats (2-4 months old, 250-450 grams at the time of the surgery) were used for this study. All animals were singly housed in transparent cages (35 cm x 45 cm x 45 cm, W x L x H) and maintained on a 12-hour reversed light-dark cycle (lights on at 9 p.m. and off at 9 a.m.). Experiments were performed during the dark phase. Rats were maintained in a vivarium with controlled temperature (19-22°C), humidity (55-65%). and were kept at about 85-90% of free-feeding body weight. Food restriction was imposed 8-24 hours before each training and recording trial. Water was available *ad libitum*. All animal experiments were performed in accordance with the

596   National Animal Welfare Act under a protocol approved by the Animal Care and Use

597   Committee from both Army Medical University and Xinqiao Hospital.

Surgery and tetrode placement

Rats were anesthetized with isoflurane. Microdrives loaded with four tetrodes were

implanted to target the hindlimb (S1HL), forelimb (S1FL) and shoulder (S1Sh) regions

of the primary somatosensory cortex (anterior-posterior (AP): 0.2-2.2 mm posterior to

bregma; medial-lateral (ML): 2.2-3.4 mm lateral to midline, dorsal-ventral (DV):

0.4/0.6-3 mm below the dura.), secured with dental cement with 8-10 anchor screws. A

screw served as the ground electrode. Tetrodes were assembled with four 17 µm

Platinum/Iridium wires (#100167, California Fine Wire Company). Tetrodes had

impedances between 150 and 300 kΩ at 1 kHz through electroplating (nanoZ; White

Matter LLC). 8-10 jeweler screws were attached into the rat skull, and individual

microdrives were anchored to screws with several rounds of application of the dental

cement.

Training and data collection

Behavioral training, tetrode turning and data recording started one-week post-surgery.

Rats were trained to run around in a 1 m x 1 m square box with a white cue card (297

mm x 210 mm) mounted on one side of the wall. Food pellets were scattered into the

arena intermittently to encourage exploration.

Each recording session lasted between 10 and 30 min to facilitate full coverage of the

testing arena. Tetrodes were lowered in steps of 25 or 50 µm daily until well-separated

single units can be identified. Data were acquired by an Axona system (Axona Ltd., St.

Albans, U.K.) at 48 kHz, band-passed between .8-6.7 kHz and a gain of x5-18k. Spikes

were digitized with 50 8-bit sample windows. Local field potentials were recorded from

one of the electrodes with a low-pass filter (500 Hz).

Spike sorting, cell classification and rate map

Spike sorting was manually performed offline with TINT (Axona Ltd, St. Albans,

U.K.), and the clustering was primarily based on features of the spike waveform (peak-
to-trough amplitude and spike width), together with additional autocorrelations and
cross-correlations [44, 82]. During our manual cluster cutting, we always counted neurons
with similar or identical waveform shapes only once whenever similar or identical
individual neurons were recorded and tracked across two consecutive recording
sessions. To confirm the quality of cluster separation, we calculated L-ratio as well as
isolation distance between clusters.

Two small light-emitting diodes (LEDs) were mounted to the headstage to track the rats'
speed, position and orientation via an overhead video camera. Only spikes with
instantaneous running speeds > 2.5 cm/s were chosen for further analysis in order to
exclude confounding behaviors such as immobility, grooming and rearing.

To classify firing fields and firing rate distributions, the position data were divided into
2.5-cm x 2.5-cm bins, and the path was smoothed with a 21-sample boxcar window
filter (400 ms; 10 samples on each side) [7, 17, 43]. Cells with > 100 spikes per session and
with a coverage of >80% were included for further analyses. Maps for spike numbers
and spike times were smoothed with a quasi-Gaussian kernel over the neighboring 5 x
5 bins [7, 17, 43]. Spatial firing rates were calculated by dividing the smoothed map of spike
numbers with spike times. The peak firing rate was defined as the highest rate in the
corresponding bin in the spatial firing rate map. Mean firing rates were calculated from
the whole session data.

Analysis of place cells

644 Spatial information is a quantification of the extent to which a neuron's firing pattern
645 can predict the position of freely moving animals and is expressed in the unit of bits per
646 spike. The spatial information was calculated as:

$$\text{spatial information} = \sum_i p_i \frac{\lambda_i}{\lambda} \log_2 \frac{\lambda_i}{\lambda}$$

where λ_i is the mean firing rate of the cell in the i-th bin, λ is the overall mean firing

rate of the cell in the trial, and p_i is the probability for the animal being at the location

of the i-th bin.

Adaptive smoothing [44] was applied to optimize the trade-off between spatial resolution

and sampling error before the calculation of spatial information. The data were first

divided into 2.5-cm x 2.5-cm bins, and then the firing rate within each bin was

calculated by expanding a circle centered on the bin until

655
$$\gamma \geq \frac{\alpha}{n\sqrt{s}}$$

where γ is the circle's radius in bins, n is the number of occupancy of samples within

the circle, s is the total number of spikes fired within the circle, and α is a constant

parameter set to 10,000. With a position sampling frequency at 50 Hz, the firing rate

assigned to that bin was then set to $50 \cdot s/n$.

A place cell was classified as a cell with the spatial information above chance level,

which was computed by a random permutation process using all recorded cells. For

each round of the shuffling process, the entire sequence of spike trains from each cell

was time-shifted along the animal's trajectory by a random period between 20 s and the

trial duration minus 20 s, with the end wrapped to the beginning of the trial. A spatial

firing rate map was then constructed, and spatial information was calculated. This

shuffling process was repeated 100 times for each cell, generating a total of 202,500

permutations for the 2025 somatosensory neurons. This shuffling procedure preserved

the temporal firing characteristics in the unshuffled data while disrupting the spatial

structure at the same time.

Spatial information score was then measured for each shuffled rate map. The

distribution of spatial information values across all 100 permutations of all cells was

computed and finally, the 99th percentile of the significant level was determined. The

threshold values for categorizing cells into place cells were defined as the spatial

information scores above the 99th percentile of the distribution from shuffled

populations. In addition to the population shuffling, the within-cell shuffling was also

performed within the entire sequence of spike trains from each individual neuron, and

the shuffling process was also repeated 100 times for each single cell.

Spatial sparsity was used to measure how compact and selective the place field of each

place cell is relative to the recording enclosure. The spatial sparsity was calculated

using the formula as follows [44]:

$$681 \quad \text{Sparsity} = \frac{(\sum p_i \lambda_i)^2}{\sum p_i \lambda_i^2}$$

Where p_i is the occupancy probability for the animal being at the location of the i-th

bin in the map and λ_i is the mean firing rate of the cell in bin i.

Spatial coherence was estimated by calculating the mean correlation between the firing

rate of each bin in the map and the aggregate firing rate of the eight nearest bins [34]. We

used unsmoothed firing rate maps for computing the spatial coherence.

The spatial correlation across trials from the same recording arena was computed for

each cell by correlating the firing rates in corresponding paired bins of two smoothed

rate maps. The spatial stability within trials was estimated by calculating spatial (2D)

correlations between firing rate maps generated from the first and second halves of the

same trial. Place cells with spatial stability lower than 0.3 were excluded for further

analysis.

Analysis of grid cells

Spatial autocorrelation was calculated with smoothed rate maps [7, 17, 43].

Autocorrelograms were derived from Pearson's product-moment correlation

coefficient correcting for edge effects and unvisited locations.

With $\lambda(x, y)$ representing the average firing rate of a cell at coordinate (x, y), the

autocorrelation between the spatial firing field itself and the spatial firing field with lags

of τ_x and τ_y was calculated as:

$r(\tau_x, \tau_y)$

701
$$= \frac{n \sum \lambda(x,y)\, \lambda(x - \tau_x, y - \tau_y) - \sum \lambda(x,y) \sum \lambda(x - \tau_x, y - \tau_y)}{\sqrt{n \sum \lambda(x,y)^2 - [\sum \lambda(x,y)]^2}\, \sqrt{n \sum \lambda(x - \tau_x, y - \tau_y)^2 - [\sum \lambda(x - \tau_x, y - \tau_y)]^2}}$$

where the summation is over whole n pixels in $\lambda(x, y)$ for which firing rate was

calculated for both $\lambda(x, y)$ and $\lambda(x - \tau_x, y - \tau_y)$. Autocorrelations were not

calculated for spatial lags of τ_x, τ_y where $n < 20$.

The degree of spatial regularity ("gridness" or "grid score") was calculated for each

unit by using a circular sample centered on the central peak of the autocorrelogram but

excluding the central peak itself, and by comparing rotated versions of this circular

sample [5, 7]. The Pearson's correlations between this circular sample and its rotated

versions were calculated, with the angles of rotation of 60° and 120° in the first group,

and 30°, 90° and 150° in the second group. Gridness or the neuron's grid score was

defined as the minimal difference between any of the coefficients in the first group and

any of the coefficients in the second group. Shuffling was performed in the same

procedure used for defining place cells. Grid cells were categorized as cells with the

rotational symmetry-based grid scores exceeding the 99[th] percentile of the distribution

of grid scores for shuffled data from the entire population of identified somatosensory

cells.

Grid spacing was computed as the median distance from the grid center to the closest

peak among six neighboring firing fields in the autocorrelogram of the spatial firing

map. Since such analysis is sensitive to noise in the grid autocorrelogram, grid spacing

was computed only when the median distance to the six neighboring peaks for the

analyzed cell was comparable to the radius of the circle centered on the gird

autocorrelogram with the highest grid score. The radius of this circle around the center

of the autocorrelogram was also referred to as the grid field size.

Grid orientation was calculated by first computing vectors from the center of the

autocorrelogram to each of the three adjacent peaks among six neighboring firing

fields in the autocorrelogram of the spatial firing map in the counterclockwise

direction, beginning with a camera-based reference line of zero degree. The angle

between the minimal orientation of those three vectors and the camera-based

reference line was defined as the grid orientation.

Analysis of head direction cells

The rat's head direction was estimated by the relative position of the LEDs

differentiated through their sizes [7, 17, 43]. The directional tuning curve for each recorded

cell was drawn by plotting the firing rate as a function of the rat's head angle, which is

divided into bins of 3 degrees and then smoothed with a 14.5-degree mean window

filter (2 bins on each side). To avoid bias, data were only used if all head angle bins

contain data.

The strength of directionality was calculated by computing the mean vector length from

circular distributed firing rates. The chance values were determined by a shuffling

process simulated in the same way as for place cells, with the entire sequence of spike

trains time-shifted between 20 s and the whole trail length minus 20 s along the animal's

trajectory. Cells were defined as head direction cells if the mean vector lengths of the

recorded cells were larger than the 99th percentile of the mean vector lengths in the

shuffled distribution. Angular stability was computed by calculating the correlation of

firing rates across directional bins generated from the first and second halves of the

same trial. Head direction cells with angular stability lower than 0.3 were excluded for

further analysis.

Analysis of border cells

Border or boundary vector cells were identified by calculating, for each recorded cell,

the difference between the maximal length of any of the four walls touching on any

single spatial firing field of the cell and the average distance of the firing field to the

nearest wall, divided by the sum of those two values [5, 17, 43]. Border scores ranged from

−1 for cells with perfect central firing fields to +1 for cells with firing fields that exactly

line up with at least one entire wall. Firing fields were defined as summation of

neighboring pixels with total firing rates higher than 0.3 times the cell's maximum

firing rate that covered a total area of at least 200 cm^2.

Border cell classification was verified in the same way as for place cells, head direction

cells and grid cells. For each permutation trial, the whole sequence of spike trains was

time-shifted along the animal's trajectory by a random period between 20 s and 20 s

less than the length of the entire trial, with the end wrapped to the start of the trial. A

spatial firing rate map was then obtained, and a border score was estimated.

The distribution of border scores was calculated for the entire set of permutation trials

from all recorded cells, and the 99th percentile was then determined. Cells were defined

as border cells if the border score from the observed data was higher than the 99th

percentile for border scores in the entire distribution generated from the permutated

data.

Whisker trimming

To test how the removal of another salient somatosensation mediate by vibrissae input

impacts on FL/HL S1 spatial cells, we carefully trimmed all whiskers of each recoded

rat bilaterally using blunt surgical scissors to within 1-2 mm above the skin surface just

right before each recording session every day. Since bilateral whisker-trimmed rats did

not exhibit abnormal behaviors, we continuously recorded somatosensory units from 2

whiskers-trimmed rats.

Environmental manipulations

For visual landmark rotation, we first recorded neuronal activity in the standard session

followed by a 90° cue-card rotation in the clockwise or counterclockwise direction.

Then another standard session was performed with the cue-card rotated back to the
original position. For recording in the elevated platform without walls, we first recorded
the somatosensory spatial cells in the square box, followed by the recording in the
elevated platform without walls. Finally, the animals were returned to the original
square box for another recording session. For the recording of border cells in the
presence of inserted wall, we first identified the somatosensory border cells in the
square box. Then the recording session was followed by the insertion of a wall along
the center of the external wall. Another recording session was performed after removing
the inserted wall. For food/no food comparisons, we recorded the somatosensory spatial
cells in two consecutive recording sessions without and with throwing food pellets into
the running enclosure.

Histology and reconstruction of recording positions

At the end of the experiment, rats were euthanized with an overdose of sodium
pentobarbital and perfused transcardially with phosphate-buffered saline (PBS)
followed by 4% paraformaldehyde (PFA). Brains were removed and stored in 4% PFA
overnight. The brain was then placed in 10, 20 and 30% sucrose/PFA solution
sequentially across 72 hours before sectioning using a cyrotome. Thirty-micron
sections were obtained through the implant region. Sections were mounted on glass
slides and stained with cresyl violet (Sigma-Aldrich). The final recording positions
were determined from digitized images of the Nissl-stained sections. Positions of each
individual recordings were estimated from the deepest tetrode track, notes on tetrode
advancement with tissue shrinkage correction by dividing the distance between the
brain surface and electrode tips by the last advanced depth of the recording electrodes.
Electrode traces were confirmed to be located within the hindlimb region (S1HL) from
eight implanted rats, the forelimb region (S1FL) from one rat and the shoulder region
(S1Sh) from another rat but all electrode tracking paths from 10 implanted rats were
verified to be away from the barrel field (S1BF) of the primary somatosensory cortex
according to the Rat Brain Atlas [83].

Data Availability: Recording dataset will be prepared available in a forthcoming public domain, and inquiries into acquiring the recording dataset beforehand should be directed to the corresponding author.

Acknowledgments: We would like to thank five anonymous reviewers for their insightful comments and constructive suggestions during the peer reviews. We are indebted to H. Wu, W. Tang, H. Chen, S. Lv, H. Yang and J. Ni for their encouragement and generous help. We would like to acknowledge J. Cai, B. Deng, L. Hu and S. Wang for their technical assistance. We would like to express our gratitude to Neil Burgess and Caswell Barry for sharing their Matlab codes with us. We would like to appreciate Calvin Young for his critical comments on the manuscript. X.L. is supported by the Chongqing Municipality postdoctoral fellowship (Grant# cstc2019jcyj-bshX0035). We are grateful for two medical research funds from Xinqiao Hospital (Grant# 2017A034 and Grant# 2019XQY16) and a startup fund from Army Medical University (Grant# 2017R028) to S.-J.Z. This work was also supported by the National Natural Science Foundation of China through the Project Grant NSFC-31872775 to S.-J.Z. Original results published in this manuscript were previously uploaded on the preprint server for biology (https://www.biorxiv.org/content/10.1101/473090v1) at *bioRxiv* on November 19, 2018 [41].

Author Contributions: S.-J.Z. conceived and designed the study. X.L. and S.-J.Z. performed the experiments, collected the data and performed the analyses. X.L. made the figures. S.-J.Z. wrote the manuscript.

Additional Information

Supplementary information (supplementary **Figs. S1-S33** and **Table S1**) accompanies this paper.

Competing Interests: The authors declare no competing interests.

Fig. S1 Electrode track and recording locations in the primary somatosensory cortex.

(**a**) Cresyl violet-stained coronal brain sections show recording electrode tracks (arrowheads) and final recording positions in seven rats with tetrodes implanted in the rat primary somatosensory cortex. (**b**) Cresyl violet-stained coronal brain sections showing representative recording locations (arrowheads) for two additionally implanted rats in the whisker trimming experiment. Broken dashes depict the boundaries of the hindlimb region (S1HL), shoulder region (S1Sh) and forelimb (S1FL) region of the primary somatosensory cortex according to the rat brain atlas of Paxinos and Watson (2007). Scale bar, 1 mm.

Fig. S2 Cluster diagrams, waveforms and isolation quality of spike clusters recorded from the somatosensory cortex.

(**a**) Scatterplots show the relationship between peak-to-trough amplitudes for all spikes for each wire (e1-e4) in a tetrode. (**b**) Overlaid and mean waveforms from two separated green and orange clusters in the scatterplots are shown for four electrodes from the same tetrode. Scale bar, 150 µV, 200 µs. (**c**) The distribution of isolation distance for identified somatosensory units. (**d**) Same as (**c**) for the L-ratio.

Fig. S3 Layer distribution of four different somatosensory spatial cell types.

(**a**) Venn diagram displaying the distribution of four functionally distinct somatosensory spatial cell types. (**b**) The approximate distributed laminar locations of all recorded functionally distinct somatosensory spatial cells. (**c-f**) The layer distribution of average number of identified somatosensory place cells (**c**), head direction cells (**d**), border cells (**e**) and grid cells (**f**) across different implanted rats.

Fig. S4 Distribution of number and characteristics of four different somatosensory spatial cell types.

(**a-d**) Distribution of number (upper panels) and spatial characteristics (bottom panels) of identified somatosensory place cells (**a**), head direction cells (**b**), border cells (**c**) and grid cells (**d**) across different animals.

Fig. S5 More examples of somatosensory place cells recorded from the somatosensory cortex.

(**a-c**) Representative somatosensory place cells with bin coverage over 90%. Trajectory

(grey line) with superimposed spike locations (red dots) (left column); heat maps of

spatial firing rate (middle column) and autocorrelation (right column) are color-coded

with dark blue indicating minimal firing rate and dark red indicating maximal firing

rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps.

Peak firing rate (fr), mean firing rate (fr) and spatial information (si) for each

representative cell are labelled at the top of the panels. Spike waveforms on four

electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Fig. S6 Comparison of spatial threshold defined by population shuffling and
873

within-cell shuffling for four different somatosensory spatial cell types.
874

875 (**a-d**) Histograms showing the 99th percentile significance level of each randomly

876 shuffled distribution for population shuffling and within-cell shuffling of all 2025

877 identified single units for identified somatosensory place cells (**a**), head direction cells

878 (**b**), border cells (**c**) and grid cells (**d**).

Fig. S7 Quantification of spatial responses of somatosensory place cells using the maximum likelihood factorial model.

(**a-c**) Firing rate maps of three representative place cells from **Fig. 1b**. Left column shows the corrected locational and directional firing rate maps using the maximum-likelihood correction approach; right column shows the corresponding uncorrected

firing rate maps. Spatial firing rate maps in four directions are shown in the lower panels.

Note the similar locational responses in four different directions.

Fig. S8 Distribution of the average firing rate and summary of spike waveform for

four different somatosensory spatial cell types.

(**a-d**) Histograms showing the average firing rate of identified somatosensory place

cells (**a**), head direction cells (**b**), border cells (**c**) and grid cells (**d**). (**e-h**) Histograms

showing the peak-to-peak amplitudes (upper panels) and peak to trough width (bottom

panels) of spike waveforms on four electrodes of identified somatosensory place cells

(**e**), head direction cells (**f**), border cells (**g**) and grid cells (**h**).

Fig. S9 Spatial stability of somatosensory place cells.

(**a-c**) Intra-trial spatial stability between the first and second halves of three representative place cells from **Fig. 1b**. Trajectory (grey line) with superimposed spike locations (red dots) (left column); rate maps (middle column) and autocorrelation maps (right column) for the first half (**i**) and the second half (**ii**) of the trials. Firing rate is color-coded with blue indicating minimum firing rate and red indicating maximum firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and spatial information (si) are labelled at the top of the plots. Pearson's correlation coefficients of firing rate maps between the first and second halves are indicated with *r* at the top-left corner. (**d**) Distribution of spatial stability of firing rate maps between the first and second halves of all identified somatosensory place cells.

Fig. S10 Persistence of somatosensory place fields from longer recording sessions.
(**a-c**) Spatial stability of three representative somatosensory place cells between the short (top panels) and longer (bottom panels) recording sessions. Trajectory (grey line) with superimposed spike locations (red dots) (left column); spatial firing rate maps (middle column) and autocorrelation diagrams (right column). Firing rate is color-coded with blue indicating minimum firing rate and red indicating maximum firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and spatial information (si) for each recording session are labelled at the top of the panels. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Fig. S11 Cue-rotation control of somatosensory place cell coding.

(**a-c**) Each panel shows the response of the same somatosensory place cell during three

sessions of the cue-rotation condition. Top panels, before the cue-rotation; middle

panels, counterclockwise or clockwise 90° of cue-rotation; bottom panels, cue-rotation

back to the original condition. The cue card is represented by a white arc in each panel.

The experimental diagram (left column); trajectory (grey line) with superimposed spike

locations (red dots) (middle left column); spatial firing rate maps (middle right column)

and autocorrelation diagrams (right column). Firing rate is color-coded with blue

indicating minimum firing rate and red indicating maximum firing rate. The scale of

the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate

(fr), mean firing rate (fr) and spatial information (si) for each recording session are

labelled at the top of the panels. Spike waveforms on four electrodes are shown on the

right column. Scale bar, 150 µV, 300 µs.

Fig. S12 Remapping of somatosensory place cells in different environments.

(**a-c**) Spatial responses of three representative somatosensory place cells in different

boxes in a constant location. The experimental diagram (left column); trajectory (grey

line) with superimposed spike locations (red dots) (middle left column); spatial firing

rate maps (middle right column) and autocorrelation diagrams (right column). Firing

rate is color-coded with blue indicating minimum firing rate and red indicating

maximum firing rate. The scale of the autocorrelation maps is twice that of the spatial

firing rate maps. Peak firing rate (fr), mean firing rate (fr) and spatial information (si)

for each recording session are labelled at the top of the panels. Spike waveforms on

four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Fig. S13 Distribution of the average in-field and out-field running speed for place

cells in the somatosensory cortex.

(**a**) The distribution of the average running speed within and outside the firing fields of

all identified place cells in the somatosensory cortex. (**b**) The comparison of the average

in-field and out-field running speed. $n = 195$, $P = 0.10$, two-tailed paired t-test, n.s., not

significant.

Fig. S14 Comparison of the spatial response of somatosensory place cells between active running and slow mobility.

(**a-c**) Comparison of spatial response of representative place cells from **Fig. 1b** with instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s, respectively. Trajectory (grey line) with superimposed spike locations (red dots) (left column); rate maps (middle column) and autocorrelation maps (right column) for instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s, respectively. Firing rate is color-coded with blue indicating minimum firing rate and red indicating maximum firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and spatial

962   information (si) are labelled at the top of the plots. Pearson's correlation coefficients

963   between two firing rate maps when the instantaneous running speeds <2.5 cm/s and

964   instantaneous running speeds > 2.5 cm/s are indicated with r at the top-left corner.

Fig. S15 More examples of somatosensory head directions cells recorded from the somatosensory cortex.

(**a-c**) Representative somatosensory head direction cells with bin coverage over 90%. Trajectory (grey line) with superimposed spike locations (red dots) (left column); spatial firing rate maps (middle left column), autocorrelation diagrams (middle right column) and head direction tuning curves (black) plotted against dwell-time polar plot (grey) (right column). Firing rate is color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr), mean vector length (mvl) and angular peak rate for each representative head direction cell are labelled at the top of the panels. The directional plots show strong head direction tuning. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Fig. S16 Quantification of head direction selectivity of somatosensory head direction cells using the maximum likelihood factorial model.

(**a-c**) Comparison of head directionality of three representative head direction cells from

Fig. 2a using a maximum-likelihood approach. Top left two panels show the corrected rate maps and polar plots under the maximum likelihood factorial model. Top right two panels show the uncorrected histograms. Mean vector length (mvl) is indicated at the top right corner of the polar plot panel. The head direction responses in each quadrant of the running box are shown in the four panels below. Somatosensory HD cells preserve the same sharp head direction selectivity in different parts of the running box regardless of possible inhomogeneous sampling of the animal's locations and orientations.

Fig. S17 Angular stability of somatosensory head direction cells.

(**a-c**) Intra-trial angular stability between the first and second halves of three representative head direction cells from **Fig. 2a**. Trajectory (grey line) with superimposed spike locations (red dots) (left column); spatial firing rate maps (middle left column), autocorrelation diagrams (middle right column) and head direction tuning curves (black) plotted against dwell-time polar plot (grey) (right column) for the first half (**i**) and the second half (**ii**) of the trials. Firing rate is color-coded with blue indicating minimum firing rate and red indicating maximum firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr), mean vector length (mvl) and angular peak rate for each

representative head direction cell are labelled at the top of the panels. Correlation coefficients of the distributed firing rate across all directional bins between the first and second halves of individual recording trials are indicated with *r*. (**d**) Population histogram of angular stability of the distributed firing rate across all directional bins between the first and second halves of all identified somatosensory head direction cells.

1007

Fig. S18 Cue-rotation control of somatosensory head directional responses.

(**a-c**) Spatial responses of three representative somatosensory head direction cells

during cue-rotation. Each panel shows the response of the same S1 head direction cell during three recording trails during the cue-rotation condition. Top panels, before the cue-rotation (**a**); middle panels, clockwise 90° of cue-rotation (**b**); bottom panels, cue-rotation back to the original condition (**c**). The cue card is represented by a white line in each panel. The experimental diagram (left column); trajectory (grey line) with superimposed spike locations (red dots) (middle left column); spatial firing rate maps (middle column), autocorrelation diagrams (middle right column) and head direction tuning curves (black) plotted against dwell-time polar plot (grey) (right column) for each recording trial. Firing rate is color-coded with blue indicating minimum firing rate and red indicating maximum firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr), mean vector length (mvl) and angular peak rate for each representative head direction cell are labelled at the top of the panels. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Fig. S19 Distribution of the average angular velocity in the preferred and non-preferred firing directions for head direction cells in the somatosensory cortex.
(**a**) The distribution of the average angular velocity in the "preferred firing directions" and "non-preferred firing directions" for all identified head direction cells in the somatosensory cortex. (**b**) The comparison of the angular velocity in the "preferred firing directions" and "non-preferred firing directions". $n = 80$, $P = 0.12$, two-tailed paired t-test, n.s., not significant.

1032

Fig. S20 Somatosensory place and head direction cells in the presence or absence

of food pellets.

(**a**) The directional tuning of the same S1 head direction cell recorded with (top panels)

and without (bottom panels) food pellets. (**b**) The firing fields of the same S1 place cell

recorded with (top panels) and without (bottom panels) food pellets. Trajectory (grey

line) with superimposed spike locations (red dots) (left column); spatial firing rate maps

(middle left column), autocorrelation diagrams (middle right column) and head

direction tuning curves (black) plotted against dwell-time polar plot (grey) (right

column). Firing rate is color-coded with blue indicating minimum firing rate and red

indicating maximum firing rate. The scale of the autocorrelation maps is twice that of

the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr), mean vector length

(mvl) or spatial information (si) and angular peak rate for each representative head

direction cell are labelled at the top of the panels. Spike waveforms on four electrodes

are shown on the right column. Scale bar, 150 µV, 300 µs.

1047

Fig. S21 Comparison of the spatial response of somatosensory head direction cells between active running and slow immobility.

(**a-c**) Comparison of the spatial response of representative head direction cells from **Fig. 2a** with instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s, respectively. Trajectory (grey line) with superimposed spike locations (red dots) (left column); rate maps (middle left column), autocorrelation maps (middle right column) and head direction tuning curves (black) plotted against dwell-time polar plot (grey) (right column) for instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s, respectively. Firing rate is color-coded with blue indicating minimum firing rate and red indicating maximum firing rate. The scale of the

autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr),

mean firing rate (fr) and mean vector length (mvl) are labelled at the top of the plots.

Correlation coefficients of the distributed firing rate across all directional bins when the

instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s

are indicated with *r* at the top-left corner.

1063

Fig. S22 More examples of somatosensory border cells recorded from the somatosensory cortex.

(**a-c**) Representative somatosensory border cells with bin coverage over 90%. Trajectory (grey line) with superimposed spike locations (red dots) (left column); spatial firing rate maps (middle left column), autocorrelation diagrams (middle right column) and head direction tuning curves (black) plotted against dwell-time polar plot (grey) (right column). Firing rate is color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr), mean vector length (mvl) and angular peak rate for each representative head direction cell are labelled at the top of the panels. The directional plots show strong head direction tuning. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

1077

Fig. S23 Preserved firing across different geometric shapes of somatosensory

border cells.

(**a-c**) Three representative S1 border cells preserve the firing patterns in the square and

cylindrical box. Top panels, original square running box; middle panels, cylindrical box;

bottom panels, back to the original square box. The experimental diagram (left column);

trajectory (grey line) with superimposed spike locations (red dots) (middle left column);

rate maps (middle right column) and autocorrelation diagrams (right column) for each

recording trail. Firing rate is color-coded with blue indicating minimum firing rate and

red indicating maximum firing rate. The scale of the autocorrelation maps is twice that

of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and border score

(b) for each recording session are labelled at the top of the panels. Spike waveforms on

four electrodes are shown on the right column. Scale bar, 200 µV, 300 µs.

Fig. S24 Spatial responses to the external inserts of somatosensory border cells.

(**a-c**) Three representative somatosensory border cells recorded in the square enclosure with or without one external insert. Top panels, the square enclosure without insert; middle panels, the same square enclosure with an internal insert; bottom panels, the

same square enclosure after removing the internal insert. The experimental diagram

(left column); trajectory (grey line) with superimposed spike locations (red dots)

(middle left column); rate maps (middle right column) and autocorrelation diagrams

(right column) for each recording trail. Firing rate is color-coded with blue indicating

minimum firing rate and red indicating maximum firing rate. The scale of the

autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr),

mean firing rate (fr) and border score (b) for each recording session are labelled at the

top of the panels. Spike waveforms on four electrodes are shown on the right column.

Scale bar, 150 μV, 300 μs.

 Fig. S25 Somatosensory border cells recorded from the elevated platform without

walls.

(**a-c**) Spatial responses of three representative somatosensory border cells recorded in

the square box, in the elevated platform without walls and back to the square box.

Trajectory (grey line) with superimposed spike locations (red dots) (left column); heat

maps of firing rate (middle column) and autocorrelation maps (right column). Firing

rate is color-coded with blue indicating minimum firing rate and red indicating

maximum firing rate. The scale of the autocorrelation maps is twice that of the spatial

firing rate maps. Peak firing rate (fr), mean firing rate (fr) and border score (b) for each

representative border cell are labelled at the top of the panels. Spike waveforms on four

electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

1116

Fig. S26 Somatosensory boundary vector cells recorded from the somatosensory

cortex.

Examples of somatosensory boundary vector cells. Trajectory (grey line) with

superimposed spike locations (red dots) (left column); heat maps of firing rate (middle

column) and autocorrelation diagrams (right column). Firing rate is color-coded with

blue indicating minimum firing rate and red indicating maximum firing rate. The scale

of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate

(fr), mean firing rate (fr) and border score (b) for each representative boundary vector

cell are labelled at the top of the panels. Spike waveforms on four electrodes are shown

on the right column. Scale bar, 150 µV, 300 µs.

1127

Fig. S27 Comparison of the spatial response of somatosensory border cells between

active running and slow immobility.

(**a-c**) Comparison of the spatial response of representative border cells from **Fig. 3a**

with instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5

cm/s, respectively. Trajectory (grey line) with superimposed spike locations (red dots)

(left column); rate maps (middle column) and autocorrelation maps (right column) for

instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s,

respectively. Firing rate is color-coded with blue indicating minimum firing rate and

red indicating maximum firing rate. The scale of the autocorrelation maps is twice that

of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and border score

(b) are labelled at the top of the plots. Pearson's correlation coefficients between two

firing rate maps when the instantaneous running speeds <2.5 cm/s and instantaneous

running speeds > 2.5 cm/s are indicated with *r* at the top-left corner.

1141

Fig. S28 More examples of somatosensory grid cells recorded from the somatosensory cortex.

Representative somatosensory grid cells with bin coverage over 90%. Trajectory (grey line) with superimposed spike locations (red dots) (left column); spatial firing rate maps (middle column) and autocorrelation diagrams (right column). Firing rate is color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and grid score (g) for each representative head direction cell are labelled at the top of the panels. The directional plots show strong head direction tuning. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

1153

Fig. S29 Somatosensory grid cells recorded in the larger environment.

(**a-c**) Three examples of somatosensory grid cells recorded in 1 m x 1m, 1.5 m x 1.5 m

and 1 m x 1m square box. The experimental diagram (left column); trajectory (grey line)

with superimposed spike locations (red dots) (middle left column); spatial firing rate

maps (middle right column) and autocorrelation diagrams (right column). Firing rate is color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and grid score (g) for each representative head direction cell are labelled at the top of the panels. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Fig. S30 Comparison of the spatial response of somatosensory grid cells between active running and slow immobility.

(**a-c**) Comparison of the spatial response of representative grid cells from **Fig. 4a** with instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s,

respectively. Trajectory (grey line) with superimposed spike locations (red dots) (left column); rate maps (middle column) and autocorrelation maps (right column) for instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s, respectively. Firing rate is color-coded with blue indicating minimum firing rate and red indicating maximum firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and grid score (b) are labelled at the top of the plots. Pearson's correlation coefficients between two firing rate maps when the instantaneous running speeds <2.5 cm/s and instantaneous running speeds > 2.5 cm/s are indicated with r at the top-left corner.

1178

1179 **Fig. S31 More examples of irregular grid cells recorded from the somatosensory**

1180 **cortex.**

Representative somatosensory irregular grid cells with bin coverage over 90%. Trajectory (grey line) with superimposed spike locations (red dots) (left column); spatial firing rate maps (middle column) and autocorrelation diagrams (right column). Firing rate is color-coded with dark blue indicating minimal firing rate and dark red indicating maximal firing rate. The scale of the autocorrelation maps is twice that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and grid score (g) for each representative head direction cell are labelled at the top of the panels. The directional plots show strong head direction tuning. Spike waveforms on four electrodes are shown on the right column. Scale bar, 150 µV, 300 µs.

Fig. S32 Preserved firing properties of the isolated somatosensory units before and after whisker trimming.

(**a** and **c**) The diagram showing the rat before (**a**) and after (**c**) whisker trimming. (**b** and **d**) Scatterplots show preserved cluster diagrams before (**b**) and after (**d**) whisker trimming of the same tetrode from the same recording rat. Each individual dot

represents a single recorded spike. Waveforms from two separated blue and red clusters in the scatterplots are shown for four electrodes from the same tetrode. The waveforms remain unchanged after whisker trimming. Scale bar, 150 µV, 300 µs. (**e**) The comparison of isolation distance for identified S1 units right before and after whisker trimming. (**f**) Same as **e** for the L-ratio. (**g**) Same as in (**f**) for the mean firing rate. (**h**) Same as in (**g**) for the peak firing rate. $n = 78$, two-tailed paired t-test, n.s., not significant.

Fig. S33 Preserved spatial firing properties of the same somatosensory border cell before and after whisker trimming.

(**a** and **b**) The diagram showing the rat before (**a**) and after (**b**) whisker trimming. A representative somatosensory border cell preserves the firing patterns in the square box

with (top and bottom panels) and without (middle panels) the internal insert before and
after whisker trimming. The experimental diagram (left column); trajectory (grey line)
with superimposed spike locations (red dots) (middle left column); rate maps (middle
right column) and autocorrelation maps (right column) for each recording trial from the
same S1 border cell. Firing rate is color-coded with blue indicating minimum firing rate
and red indicating maximum firing rate. The scale of the autocorrelation maps is twice
that of the spatial firing rate maps. Peak firing rate (fr), mean firing rate (fr) and border
score (b) for each recording session are labelled at the top of the panels. Spike
waveforms on four electrodes are shown on the right column. Scale bar, 100 µV, 300
µs.

Table S1. Summary of somatosensory spatial cell types recorded in each rat.

Rat Number	Somatosensory Spatial Cell Types				
	Place Cells	Head Direction Cells	Border Cells	Grid Cells	Total Cells
1 (S1HL)	22	13	6	12	182
2 (S1HL)	27	15	14	13	312
3 (S1FL)	19	9	8	10	353
4 (S1HL)	20	7	12	9	275
5 (S1Sh)	20	6	8	12	263
6 (S1HL)	35	8	16	7	197
7 (S1HL)	36	8	12	5	301
8 (S1HL)	16	14	10	4	142
Total	195	80	86	72	2025

References

1 . Poulter S, Hartley T, Lever C. The Neurobiology of Mammalian Navigation.
    Current biology : CB 2018; **28**:R1023-R1042.
2 . Buzsaki G, Moser EI. Memory, navigation and theta rhythm in the hippocampal-
    entorhinal system. *Nat Neurosci* 2013; **16**:130-138.
3 . O'Keefe J, Dostrovsky J. The hippocampus as a spatial map. Preliminary
    evidence from unit activity in the freely-moving rat. *Brain Res* 1971; **34**:171-
    175.
4 . Lever C, Burton S, Jeewajee A, O'Keefe J, Burgess N. Boundary vector cells in
    the subiculum of the hippocampal formation. *J Neurosci* 2009; **29**:9771-9777.
5 . Solstad T, Boccara CN, Kropff E, Moser MB, Moser EI. Representation of
    geometric borders in the entorhinal cortex. *Science* 2008; **322**:1865-1868.
6 . Hafting T, Fyhn M, Molden S, Moser MB, Moser EI. Microstructure of a spatial

1232   map in the entorhinal cortex. *Nature* 2005; **436**:801-806.

.   Sargolini F, Fyhn M, Hafting T *et al.* Conjunctive representation of position, direction, and velocity in entorhinal cortex. *Science* 2006; **312**:758-762.

.   Taube JS, Muller RU, Ranck JB, Jr. Head-direction cells recorded from the postsubiculum in freely moving rats. I. Description and quantitative analysis. *J Neurosci* 1990; **10**:420-435.

.   Cullen KE, Taube JS. Our sense of direction: progress, controversies and challenges. *Nat Neurosci* 2017; **20**:1465-1473.

.   Town SM, Brimijoin WO, Bizley JK. Egocentric and allocentric representations in auditory cortex. *PLoS Biol* 2017; **15**:e2001878.

.   Fiser A, Mahringer D, Oyibo HK, Petersen AV, Leinweber M, Keller GB. Experience-dependent spatial expectations in mouse visual cortex. *Nat Neurosci* 2016; **19**:1658-1664.

.   Ji D, Wilson MA. Coordinated memory replay in the visual cortex and hippocampus during sleep. *Nat Neurosci* 2006.

.   Rancz EA, Moya J, Drawitsch F, Brichta AM, Canals S, Margrie TW. Widespread vestibular activation of the rodent cortex. *J Neurosci* 2015; **35**:5926-5934.

.   Alexander AS, Nitz DA. Spatially Periodic Activation Patterns of Retrosplenial Cortex Encode Route Sub-spaces and Distance Traveled. *Current biology : CB* 2017; **27**:1551-1560 e1554.

.   Leutgeb S, Mizumori SJ. Context-specific spatial representations by lateral septal cells. *Neuroscience* 2002; **112**:655-663.

.   Wilber AA, Clark BJ, Forster TC, Tatsuno M, McNaughton BL. Interaction of egocentric and world-centered reference frames in the rat posterior parietal cortex. *J Neurosci* 2014; **34**:5431-5446.

.   Boccara CN, Sargolini F, Thoresen VH *et al.* Grid cells in pre- and parasubiculum. *Nat Neurosci* 2010; **13**:987-994.

.   Weible AP, Rowland DC, Pang R, Kentros C. Neural correlates of novel object and novel location recognition behavior in the mouse anterior cingulate cortex. *J Neurophysiol* 2009; **102**:2055-2068.

.   Mao D, Kandler S, McNaughton BL, Bonin V. Sparse orthogonal population representation of spatial context in the retrosplenial cortex. *Nat Commun* 2017; **8**:243.

.   Jacob PY, Casali G, Spieser L, Page H, Overington D, Jeffery K. An independent, landmark-dominated head-direction signal in dysgranular retrosplenial cortex. *Nat Neurosci* 2017; **20**:173-175.

.   Jankowski MM, O'Mara SM. Dynamics of place, boundary and object encoding in rat anterior claustrum. *Front Behav Neurosci* 2015; **9**:250.

.   Barry C, Burgess N. To be a Grid Cell: Shuffling procedures for determining "Gridness". *bioRxiv* 2017.

.   Constantinescu AO, O'Reilly JX, Behrens TEJ. Organizing conceptual knowledge in humans with a gridlike code. *Science* 2016; **352**:1464-1468.

.   Kunz L, Maidenbaum S, Chen D, Wang L, Jacobs J, Axmacher N. Mesoscopic

1276 Neural Representations in Spatial Navigation. *Trends Cogn Sci* 2019; **23**:615-
1277 630.

25 . Epstein RA, Patai EZ, Julian JB, Spiers HJ. The cognitive map in humans:
spatial navigation and beyond. *Nat Neurosci* 2017; **20**:1504-1513.

26 . Stangl M, Achtzehn J, Huber K, Dietrich C, Tempelmann C, Wolbers T.
Compromised Grid-Cell-like Representations in Old Age as a Key Mechanism
to Explain Age-Related Navigational Deficits. *Current biology : CB* 2018;
28:1108-1115 e1106.

27 . Jacobs J, Weidemann CT, Miller JF *et al.* Direct recordings of grid-like neuronal
activity in human spatial navigation. *Nat Neurosci* 2013; **16**:1188-1190.

28 . Doeller CF, Barry C, Burgess N. Evidence for grid cells in a human memory
network. *Nature* 2010; **463**:657-661.

29 . Maidenbaum S, Miller J, Stein JM, Jacobs J. Grid-like hexadirectional
modulation of human entorhinal theta oscillations. *Proceedings of the National
Academy of Sciences of the United States of America* 2018; **115**:10798-10803.

30 . Hawkins J, Lewis M, Klukas M, Purdy S, Ahmad S. A Framework for
Intelligence and Cortical Function Based on Grid Cells in the Neocortex. *Front
Neural Circuits* 2018; **12**:121.

31 . Bonnevie T, Dunn B, Fyhn M *et al.* Grid cells require excitatory drive from the
hippocampus. *Nat Neurosci* 2013; **16**:309-317.

32 . Hales JB, Schlesiger MI, Leutgeb JK, Squire LR, Leutgeb S, Clark RE. Medial
entorhinal cortex lesions only partially disrupt hippocampal place cells and
hippocampus-dependent place memory. *Cell reports* 2014; **9**:893-901.

33 . Jezek K, Henriksen EJ, Treves A, Moser EI, Moser MB. Theta-paced flickering
between place-cell maps in the hippocampus. *Nature* 2011; **478**:246-249.

34 . Muller RU, Kubie JL. The effects of changes in the environment on the spatial
firing of hippocampal complex-spike cells. *J Neurosci* 1987; **7**:1951-1968.

35 . Zhang S, Manahan-Vaughan D. Spatial Olfactory Learning Contributes to Place
Field Formation in the Hippocampus. *Cerebral cortex (New York, NY : 1991)*
2013.

36 . Gener T, Perez-Mendez L, Sanchez-Vives MV. Tactile modulation of
hippocampal place fields. *Hippocampus* 2013.

37 . Brecht M. The Body Model Theory of Somatosensory Cortex. *Neuron* 2017;
94:985-992.

38 . Yin A, Tseng PH, Rajangam S, Lebedev MA, Nicolelis MAL. Place Cell-Like
Activity in the Primary Sensorimotor and Premotor Cortex During Monkey
Whole-Body Navigation. *Sci Rep* 2018; **8**:9184.

39 . McNaughton BL, Battaglia FP, Jensen O, Moser EI, Moser MB. Path integration
and the neural basis of the 'cognitive map'. *Nat Rev Neurosci* 2006; **7**:663-678.

40 . Ravassard P, Kees A, Willers B *et al.* Multisensory control of hippocampal
spatiotemporal selectivity. *Science* 2013; **340**:1342-1346.

41 . Long X, Zhang SJ. A novel somatosensory spatial navigation system outside the
hippocampal formation. *bioRxiv* 2018; **doi: 10.1101/473090**.

42 . Harris KD, Hirase H, Leinekugel X, Henze DA, Buzsaki G. Temporal

interaction between single spikes and complex spike bursts in hippocampal pyramidal cells. *Neuron* 2001; **32**:141-149.

. Zhang SJ, Ye J, Miao C *et al.* Optogenetic dissection of entorhinal-hippocampal functional connectivity. *Science* 2013; **340**:1232627.

. Skaggs WE, McNaughton BL, Wilson MA, Barnes CA. Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* 1996; **6**:149-172.

. Cacucci F, Lever C, Wills TJ, Burgess N, O'Keefe J. Theta-modulated place-by-direction cells in the hippocampal formation in the rat. *J Neurosci* 2004; **24**:8265-8277.

. Burgess N, Cacucci F, Lever C, O'Keefe J. Characterizing multiple independent behavioral correlates of cell firing in freely moving animals. *Hippocampus* 2005; **15**:149-153.

. Thompson LT, Best PJ. Place cells and silent cells in the hippocampus of freely-behaving rats. *J Neurosci* 1989; **9**:2382-2390.

. Jung MW, Wiener SI, McNaughton BL. Comparison of spatial firing characteristics of units in dorsal and ventral hippocampus of the rat. *J Neurosci* 1994; **14**:7347-7356.

. Rich PD, Liaw HP, Lee AK. Place cells. Large environments reveal the statistical structure governing hippocampal representations. *Science* 2014; **345**:814-817.

. Leutgeb S, Leutgeb JK, Barnes CA, Moser EI, McNaughton BL, Moser MB. Independent codes for spatial and episodic memory in hippocampal neuronal ensembles. *Science* 2005; **309**:619-623.

. Lever C, Wills T, Cacucci F, Burgess N, O'Keefe J. Long-term plasticity in hippocampal place-cell representation of environmental geometry. *Nature* 2002; **416**:90-94.

. Goodridge JP, Dudchenko PA, Worboys KA, Golob EJ, Taube JS. Cue control and head direction cells. *Behavioral neuroscience* 1998; **112**:749-761.

. Taube JS. Head direction cells recorded in the anterior thalamic nuclei of freely moving rats. *J Neurosci* 1995; **15**:70-86.

. Stewart S, Jeewajee A, Wills TJ, Burgess N, Lever C. Boundary coding in the rat subiculum. *Philos Trans R Soc Lond B Biol Sci* 2014; **369**:20120514.

. Sofroniew NJ, Svoboda K. Whisking. *Current biology : CB* 2015; **25**:R137-140.

. Chorev E, Preston-Ferrer P, Brecht M. Representation of egomotion in rat's trident and E-row whisker cortices. *Nat Neurosci* 2016; **19**:1367-1373.

. Insausti R, Herrero MT, Witter MP. Entorhinal cortex of the rat: cytoarchitectonic subdivisions and the origin and distribution of cortical efferents. *Hippocampus* 1997; **7**:146-183.

. Cenquizca LA, Swanson LW. Spatial organization of direct hippocampal field CA1 axonal projections to the rest of the cerebral cortex. *Brain Res Rev* 2007.

. Markram H, Muller E, Ramaswamy S *et al.* Reconstruction and Simulation of Neocortical Microcircuitry. *Cell* 2015; **163**:456-492.

. Li CL, Li KC, Wu D *et al.* Somatosensory neuron types identified by high-

coverage single-cell RNA-sequencing and functional heterogeneity. *Cell Res* 2016; **26**:83-102.

. Gaykema RP, Luiten PG, Nyakas C, Traber J. Cortical projection patterns of the medial septum-diagonal band complex. *J Comp Neurol* 1990; **293**:103-124.

. Bolding KA, Ferbinteanu J, Fox SE, Muller RU. Place cell firing cannot support navigation without intact septal circuits. *Hippocampus* 2019.

. Koenig J, Linder AN, Leutgeb JK, Leutgeb S. The spatial periodicity of grid cells is not sustained during reduced theta oscillations. *Science* 2011; **332**:592-595.

. Brandon MP, Bogaard AR, Libby CP, Connerney MA, Gupta K, Hasselmo ME. Reduction of theta rhythm dissociates grid cell spatial periodicity from directional tuning. *Science* 2011; **332**:595-599.

. Feldman DE, Brecht M. Map plasticity in somatosensory cortex. *Science* 2005; **310**:810-815.

. Petersen CCH. Sensorimotor processing in the rodent barrel cortex. *Nat Rev Neurosci* 2019; **20**:533-546.

. Killackey HP, Sherman SM. Corticothalamic projections from the rat primary somatosensory cortex. *J Neurosci* 2003; **23**:7381-7384.

. Yamawaki N, Shepherd GM. Synaptic circuit organization of motor corticothalamic neurons. *J Neurosci* 2015; **35**:2293-2307.

. Schubert D, Staiger JF, Cho N, Kotter R, Zilles K, Luhmann HJ. Layer-specific intracolumnar and transcolumnar functional connectivity of layer V pyramidal cells in rat barrel cortex. *J Neurosci* 2001; **21**:3580-3592.

. Larkum M. A cellular mechanism for cortical associations: an organizing principle for the cerebral cortex. *Trends in neurosciences* 2013; **36**:141-151.

. Lefort S, Tomm C, Floyd Sarria JC, Petersen CC. The excitatory neuronal network of the C2 barrel column in mouse primary somatosensory cortex. *Neuron* 2009; **61**:301-316.

. Buzsaki G, Stark E, Berenyi A *et al.* Tools for probing local circuits: high-density silicon probes combined with optogenetics. *Neuron* 2015; **86**:92-105.

. Sieben K, Roder B, Hanganu-Opatz IL. Oscillatory entrainment of primary somatosensory cortex encodes visual control of tactile processing. *J Neurosci* 2013; **33**:5736-5749.

. Khazipov R, Sirota A, Leinekugel X, Holmes GL, Ben-Ari Y, Buzsaki G. Early motor activity drives spindle bursts in the developing somatosensory cortex. *Nature* 2004; **432**:758-761.

. Kaas JH. What, if anything, is SI? Organization of first somatosensory area of cortex. *Physiological Reviews* 1983; **63**:206-231.

. Welker E, Hoogland PV, Van der Loos H. Organization of feedback and feedforward projections of the barrel cortex: a PHA-L study in the mouse. *Exp Brain Res* 1988; **73**:411-435.

. Umeda T, Isa T, Nishimura Y. The somatosensory cortex receives information about motor output. *Science advances* 2019; **5**:eaaw5388.

. Karadimas SK, Satkunendrarajah K, Laliberte AM *et al.* Sensory cortical control

of movement. *Nature Neuroscience* 2019.

79 .   Ashe J, Georgopoulos AP. Movement parameters and neural activity in motor

cortex and area 5. *Cerebral cortex (New York, NY : 1991)* 1994; **4**:590-600.

80 .   Moran DW, Schwartz AB. Motor cortical representation of speed and direction

during reaching. *J Neurophysiol* 1999; **82**:2676-2692.

81 .   Mathis A, Mamidanna P, Cury KM *et al.* DeepLabCut: markerless pose

estimation of user-defined body parts with deep learning. *Nat Neurosci* 2018;

21:1281-1289.

82 .   Wilson MA, McNaughton BL. Dynamics of the hippocampal ensemble code for

space. *Science* 1993; **261**:1055-1058.

83 .   Paxinos G, Watson C. The rat brain in stereotaxic coordinates. Amsterdam:

Elsevier 2007.