Ran-GTP is non-essential to activate NuMA for spindle pole focusing,

but dynamically polarizes HURP to control mitotic spindle length

Kenta Tsuchiya[1#], Hisato Hayashi[1#], Momoko Nishina[1#], Masako Okumura[1#],

Yoshikatsu Sato[1], Masato T. Kanemaki[2,3], Gohta Goshima[1], Tomomi Kiyomitsu[1,2,4*]

[1] Division of Biological Science, Graduate School of Science, Nagoya University, Chikusa-ku, Nagoya 464-8602, Japan.

[2] Precursory Research for Embryonic Science and Technology (PRESTO) Program, Japan Science and Technology Agency, 4-1-8 Honcho Kawaguchi, Saitama 332-0012, Japan.

[3] Department of Chromosome Science, National Institute of Genetics, Research Organization of Information and Systems (ROIS), and Department of Genetics, SOKENDAI (The Graduate University of Advanced Studies), Yata 1111, Mishima, Shizuoka 411-8540, Japan.

[4] Okinawa Institute of Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Kunigami-gun, Okinawa 904-0495, Japan

\# These authors contributed equally to this work.
* Corresponding author:
E-mail: tomomi.kiyomitsu@oist.jp
Phone & Fax: +81-98-966-1609
Characters: 5,173 words

Abstract

During mitosis, a bipolar spindle is assembled around chromosomes to efficiently capture chromosomes. Previous work proposed that a chromosome-derived Ran-GTP gradient promotes spindle assembly around chromosomes by liberating spindle assembly factors (SAFs) from inhibitory importins. However, Ran's dual functions in interphase nucleocytoplasmic transport and mitotic spindle assembly have made it difficult to assess its mitotic roles in somatic cells. Here, using auxin-inducible degron technology in human cells, we developed acute mitotic degradation assays to dissect Ran's mitotic roles systematically and separately from its interphase function. In contrast to the prevailing model, we found that the Ran pathway is not essential for spindle assembly activities that occur at sites spatially separated from chromosomes, including activating NuMA for spindle pole focusing or for targeting TPX2. In contrast, Ran-GTP is required to localize HURP and HSET specifically at chromosome-proximal regions. We demonstrated that Ran-GTP and importin-β coordinately promote HURP's dynamic microtubule binding-dissociation cycle near chromosomes, which results in stable kinetochore-fiber formation. Intriguingly, this pathway acts to establish proper spindle length preferentially during prometaphase, rather than metaphase. Together, we propose that the Ran pathway is required to activate SAFs specifically near chromosomes, but not generally during human mitotic spindle assembly. Ran-dependent spindle assembly is likely coupled with parallel pathways to activate SAFs, including NuMA, for spindle pole focusing away from chromosomes.

Highlights

・ Using auxin-inducible degron technology, we developed mitotic degradation assays for the Ran pathway in human cells.

・ The Ran pathway is non-essential to activate NuMA for spindle pole focusing.

・ The Ran pathway dynamically polarizes HURP and defines mitotic spindle length preferentially during prometaphase.

・ Ran-GTP is required to activate SAFs specifically near chromosomes, but not generally, in human mitotic cells.

Introduction

During cell division, a microtubule-based spindle structure is assembled around chromosomes to efficiently capture and segregate duplicated chromosomes into daughter cells [1, 2]. To assemble a spindle around chromosomes, chromosomes generate a gradient of Ran-GTP, a GTP-bound form of Ran, in animal cells [3, 4]. Ran-GTP is produced by regulator of chromosome condensation 1 (RCC1), a guanine nucleotide exchange factor for Ran [5], and is hydrolyzed to Ran-GDP by RanGAP1, a GTPase-activating protein for Ran [6]. Because RCC1 and RanGAP1 mainly localize on chromosomes and in cytoplasm, respectively, these opposing enzymes create a chromosome-derived Ran-GTP gradient after the nuclear envelope breaks down (Fig. 2A). During interphase, these enzymes generate different Ran-GTP concentrations in the nucleus and cytoplasm, which drives nucleocytoplasmic transport [4]. The Ran-GTP gradient has been best characterized in *Xenopus* egg extracts [7, 8], but is also found in other meiotic and mitotic cell types [9-11]. Recent studies indicate that Ran-GTP is essential for acentrosomal spindle assembly in female meiosis [9, 12, 13], but the significance of Ran-GTP in mitotic spindle assembly has been debated [10, 11, 14]. The dual functions of Ran in both interphase and mitosis have made it difficult to identify its mitotic roles in somatic cells.

As in mechanisms in nucleocytoplasmic transport, Ran-GTP binds to importin-β and releases inhibitory importins from SAFs, thereby activating SAFs near chromosomes (Fig. 2A) [15-18]. Once activated, most SAFs interact with microtubules and spatially regulate microtubule nucleation, dynamics, transport, and cross-linking, to create specialized local structures of the spindle [3, 4]. For instance, nuclear mitotic

apparatus protein (NuMA) recognizes minus-ends of microtubules and transports and

crosslinks microtubules in cooperation with cytoplasmic dynein, a minus-end-directed

motor, to focus spindle microtubules at the poles of mammalian cells [19-22]. The

targeting protein for Xklp2 (TPX2) is required for spindle pole organization [23, 24] and

stimulates microtubule nucleation in a Ran- and importin-α-regulated manner [25-27].

Kinesin-14 HSET/XCTK2 cross-links both parallel and anti-parallel microtubules near

chromosomes, but preferentially cross-links parallel microtubules near the spindle poles

[28-30]. Hepatoma upregulated protein (HURP) accumulates on microtubules near

chromosomes to form stabilized kinetochore-fibers (k-fibers) [31].

Most SAFs, including NuMA, TPX2, and HSET contain a nuclear localization

sequence/signal (NLS) [28, 32, 33]. The NLS is specifically recognized by importin-α,

which forms hetero-dimer with importin-β through an importin-β binding (IBB) domain

(Fig. 2A). On the other hand, some SAFs, such as HURP, are directly recognized by

importin-β (Fig. 2A) [31]. Because SAFs represent a small fraction of NLS-bearing

nuclear proteins and need to be regulated effectively and selectively in mitosis, each

SAF interacts with importins in specific ways to reduce competition with other nuclear

proteins and SAFs [32, 33].

In mitotic human cells, NuMA localizes to the spindle poles and the polar cell

cortex, where it facilitates spindle-pole focusing and astral microtubule capture/pulling,

respectively [19, 20, 34]. Recent structural and *in vitro* studies have demonstrated that

NuMA's microtubule-binding activities are inhibited by steric blockage of importin-β,

mediated by importin-α [32], but this model has not been rigorously tested in cells. In

addition, given that the Ran-GTP gradient diminishes with increasing distances from

chromosomes, it is unclear whether and how the Ran-GTP gradient activates NuMA at

the spindle poles.

To precisely understand mechanisms and significance of Ran-based regulation

of SAFs, it is critical to separate Ran's mitotic roles from its interphase

nucleocytoplasmic transport function. To achieve this, we developed mitotic depletion

assays for the Ran pathway in human cells by combining mitotic drugs with auxin-

inducible degron (AID) technology [35], which allows us to degrade mAID-tag fusion

proteins with a half-life of 20 min. In contrast to the prevailing model, we found that

degradation of RCC1, RanGAP1, or importin-β does not substantially affect localization

and function of NuMA at the spindle poles, even if these proteins were degraded during

mitosis. In sharp contrast, the Ran pathway polarizes both HURP and importin-β on k-

fibers near chromosomes, where HURP stabilizes k-fibers independently of importin-β.

Based on our results, we propose that the Ran-Importin pathway is required to activate

SAFs specifically near chromosomes, but not generally, in human mitotic cells.

Results

In human cells, NuMA focuses spindle microtubules at spindle poles using its C-terminal conserved microtubule-binding domain

NuMA functions in spindle microtubule focusing in cultured mammalian cells [19-22]. Silk et al. demonstrated that NuMA's C-terminal microtubule-binding domain (MTBD1) adjacent to a NLS is required for spindle pole focusing in mouse fibroblasts [22] (Fig. 1A). However, this domain is dispensable for spindle pole focusing in mouse keratinocytes [36]. In addition, NuMA has a second microtubule-binding domain (MTBD2) at the C-terminal end (Fig. 1A) [32, 37], which has stronger microtubule-binding activity and is sterically inhibited by importin-β *in vitro* [32]. To understand which domain of NuMA is required for spindle pole focusing in mitotic human cells, we replaced endogenous NuMA with C-terminal truncation mutants in HCT116 cells (Fig. 1A). Endogenous NuMA was visualized by integrating an mAID-mClover-FLAG (mACF) tag into both alleles of the NuMA gene [20]. NuMA-mACF was depleted using the auxin inducible degradation (AID) system following Dox and IAA treatment (S1A) [20, 35], and mCherry-tagged NuMA mutants were simultaneously expressed from the Rosa 26 locus by Dox treatment (Fig. 1B-C, S1B) [20]. Like endogenous NuMA, mCherry-tagged NuMA wild type (WT) accumulated in interphase nuclei (Fig. 1B) and at mitotic spindle poles (Fig. 1C #1) and was able to rescue pole-focusing defects caused by NuMA depletion (Fig. 1C-D #1). NuMA-ΔNLS mutants were unable to localize at nuclei in interphase (Fig. 1B, S1C), but were able to accumulate at spindle poles to rescue pole-focusing defects (Fig. 1C-D #2). As expected, NuMA ΔC-ter mutants, which lack both MTBDs, diffused into the cytoplasm during metaphase (Fig. 1C #5), and were unable to

rescue the spindle-pole focusing defect (Fig. 1C-D #5). In contrast, NuMA

Δ(NLS+MTBD2) mutants localized around spindle-poles to rescue the focusing defects

(Fig. 1C-D #4). However, NuMA Δex24 mutants, which lack NLS and the part of MTBD1

containing the well-conserved NLM motif (Fig. S1D) [38, 39], were unable to fully rescue

focusing defects, while localizing around the spindle-poles (Fig. 1C-D #3). These results

indicate that NuMA's MTBD1, but not MTBD2, is essential for spindle pole focusing in

human cells.

NuMA localizes at the spindle poles and participates in spindle pole focusing

independently of RCC1

NuMA's MTBD1 is located next to NLS, which is recognized by importin-α [32]. A recent

study indicated that the importin-α/β complex sterically inhibits NuMA's microtubule-

binding activity, but is released from NuMA by Ran-GTP *in vitro* [32] (Fig. 2A). To test

this model in cells, we next depleted RCC1 (RanGEF) by integrating mAID-mClover

(mAC) tag (Fig. 2B, S2A) [35]. In contrast to the model, NuMA accumulated normally

around the spindle poles, and spindle microtubules were properly focused in RCC1-

depleted cells (Fig. 2B, D, Fig. S2B-C), although metaphase spindle length diminished

(Fig. 2B-C), and mitotic duration was slightly delayed (Fig. S2D-F).

To further analyze functions of NuMA in RCC1-depleted cells, we next co-

depleted RCC1 and NuMA. Following treatment with Dox and IAA, both RCC1-mAC

and NuMA-mAID-mCherry were degraded, and spindle microtubules were not properly

focused (Fig. 2D-E, S2G-H). These results indicate that NuMA acts at spindle poles,

even in the absence of Ran-GTP in human mitotic cells.

Mitotic degradation of RCC1 does not affect localization and function of NuMA at

spindle poles

NuMA is transported into the nucleus during interphase (Fig. 1B) [32, 40], where it is

likely released from importins by nuclear Ran-GTP. Because we found that NuMA is

maintained in the nucleus following RCC1 degradation in interphase (Fig. S2E, t = -

0:10), the majority of NuMA may already have been liberated from importins by RCC1

before its degradation and may have been maintained in an active form in the nucleus,

thereby producing no aberrant phenotypes in the subsequent mitosis in RCC1-depleted

cells. To exclude this possibility, we next depleted RCC1 in nocodazole-arrested cells

and analyzed the behavior of NuMA following nocodazole washout (For procedure, see

Fig. S2I).

In RCC1-positive control cells, NuMA diffused into the cytoplasm during

nocodazole arrest (Fig. 2F, t = -90), but rapidly accumulated near chromosome masses

following nocodazole washout (Fig. 2F, t = 10). NuMA localized at the poles of

metaphase spindles within 60 min (Fig. 2F, t = 60) and entered the nucleus following

mitotic exit (Fig. 2F, t = 85). Importantly, NuMA accumulated similarly at focused spindle

poles, even if RCC1 was degraded during nocodazole arrest. RCC1-mAC signals were

initially detectable on chromosome masses during nocodazole-arrest (Fig. 2G, t = -90,

arrow), but were reduced to undetectable levels after addition of IAA (Fig. 2G, t = 0).

After nocodazole-washout, NuMA localized to focused spindle poles after ~60 min (Fig.

2G, t = 55), as observed in control cells. Cells entered anaphase with timing similar to

that of control cells (Fig. S2J), but NuMA was not recruited to the nucleus after mitotic

exit (Fig. 2G, t = 80).

As observed when RCC1 was degraded in asynchronous culture (Fig. 2C), the

metaphase spindle became shorter when RCC1 was depleted during nocodazole-arrest

(Fig. 2H). In addition, the metaphase spindle was not properly oriented to the attached

culture dishes (Fig. 2I) Taken together, these results indicate that RCC1 participates in

some fashion in spindle assembly in human mitotic cells, but is dispensable for NuMA

localization and function at spindle poles, even if RCC1 is degraded during mitosis.

NuMA localized at spindle poles is released from importins independently of Ran-

GTP

Our results suggest that NuMA is liberated from importins in the absence of Ran-GTP

(Fig. 3A). To confirm this, we next analyzed importin localization. Because importin-α

has several isoforms in human cells [41], we first examined localization of endogenous

importin-β in living cells by fusing it with mCherry (Fig. S3A). Unexpectedly, importin-β-

mCh accumulated on kinetochore-microtubules (k-fibers) near chromosomes, but not at

metaphase spindle poles (Fig. 3B top). Although this is inconsistent with the reported

spindle-pole localization of importin-β [42], this result was confirmed by other

visualization methods using a mAC tag and anti-importin-β antibodies (Fig. S3B,C).

Importantly, RCC1 depletion diminished importin-β from k-fibers, but did not cause

importin-β accumulation at the spindle poles (Fig. 3B, bottom) where NuMA localized

(Fig. 2B). This suggests that NuMA is released from importin-β at the spindle poles,

even in the absence of Ran-GTP (Fig. 3A).

To further test whether Ran-independent pathways exist for NuMA activation,

we next analyzed localizations of importin-α wild type (WT) and ΔIBB mutants, which

lack the importin-β-binding (IBB) domain. Importin-α ΔIBB mutants are insensitive to

Ran-GTP due to the lack of an IBB domain, but are still able to interact with NuMA and

partially inhibit NuMA's microtubule-binding activity *in vitro* (Fig. 3A) [32]. However,

importin-α ΔIBB diffused into cytoplasm similarly to importin-α WT, and neither affected

NuMA's spindle-pole localization nor colocalized with NuMA at the spindle poles in our

experimental conditions (Fig. 3C, S3D). These results suggest that NuMA is released

from the importin-α/β complex in a Ran-GTP-independent manner and that it localizes

at spindle poles.

Ran-GAP1 and importin-β degradation do not affect NuMA localization and

function at spindle poles

Although RCC1 depletion does not affect NuMA localization and functions, degradation

of Ran-GAP1 or importin-β may cause abnormal activation of NuMA throughout human

cells, resulting in spindle assembly defects. To test this, we next degraded either Ran-

GAP1 or importin-β using AID technology (Fig. 3D-E). Ran-GAP1 degradation caused

few mitotic phenotypes (Fig. 3D, S3E-I) and did not affect NuMA's spindle-pole

localization (Fig. 3D bottom). Similarly, importin-β degradation did not affect NuMA's

localization and function at spindle poles (Fig. 3E bottom), although importin-β

degradation caused short spindles and mitotic delay (Fig. S3J-N). These results indicate

that Ran-dependent spatial regulation is dispensable for NuMA localization and function

at spindle poles in cultured human cells.

RCC1 regulates chromosome-proximal localization of HURP and HSET

RCC1 depletion caused shorter mitotic spindles (Fig. 2B, C, F, H), suggesting that Ran-

GTP serves some function in spindle assembly in human cells. To identify spindle

assembly factors (SAFs) regulated by Ran-GTP, we next analyzed the localization of 3

other major SAFs: TPX2, HSET, and HURP. mCherry-tagged TPX2 colocalized with

SiR-tubulin signals in metaphase (Fig. 4A top, Fig. S4A), and its localization was

virtually unaffected in RCC1-depleted cells (Fig. 4A bottom), as observed for NuMA

(Fig. 2B). In contrast, mCherry-tagged HSET localized everywhere along spindle

microtubules (Fig. 4B top, Fig. S4B) [30], and its spindle localization was selectively

reduced near chromosomes following RCC1 depletion, although HSET still localized

along spindle fibers farther away from chromosomes (Fig. 4B bottom). On the other

hand, mCherry-tagged HURP accumulated at k-fibers near chromosomes, but localized

weakly on spindle microtubules following RCC1 depletion (Fig. 4C and Fig. S4C). These

results suggest that in human mitotic cells, the chromosome-derived Ran-GTP gradient

regulates SAF localization preferentially near chromosomes, regardless of the presence

of NLS (Fig. 4D).

HURP, but not importin-β, is required to stabilize k-fibers

Importin-β inhibits HURP's microtubule-binding activities by masking one of HURP's

microtubule-binding domains (MTBD2) [43] (Fig. 5J). To understand the relationship

between HURP and importin-β for k-fiber localization and function, we next sought to

degrade endogenous HURP using AID (Fig. 5A and Fig. S5A-B). Endogenous HURP-

mACF accumulated at k-fibers near chromosomes (Fig. 5A top), as observed with anti-

HURP antibodies [31]. HURP depletion resulted in diminished importin-β localization to

k-fibers (Fig. 5A-B, S5C) and reduced mitotic spindle length (Fig. 5C). Because k-fibers

are resistant to cold treatment [31], we next incubated cells with ice-cold medium for 20

min and analyzed cold-stable microtubules. HURP localized to cold-stable microtubules

(Fig. 5D, top), which were disrupted by HURP depletion (Fig. 4D bottom), consistent

with a previous study [31].

We next depleted importin-β and analyzed effects of this depletion on HURP and

k-fibers (Fig. 5E, S5D). Importin-β depletion caused a remarkable re-localization of

HURP from k-fibers near chromosomes to spindle microtubules (Fig. 5E-F). Although k-

fiber localization of HURP was unclear in importin-β-depleted cells due to the relatively

strong accumulation of HURP on spindle microtubules around spindle poles (Fig. 5E

bottom), HURP was clearly detected on cold-stable k-fibers in importin-β-depleted cells

(Fig. 5G bottom). These results suggest that HURP acts in k-fiber stabilization,

independently of importin-β.

HURP and importin-β localize throughout the spindle in RanGAP1-depleted cells

Whereas HURP and importin-β have different roles in k-fiber stabilization (Fig. 5D, G),

both proteins accumulate at k-fibers near chromosomes downstream of RCC1 (Fig. 3B,

4C). To better understand mechanisms of Ran-based spatial regulation of HURP and

importin-β, we next analyzed behavior of HURP and importin-β in RanGAP1-depleted

cells, in which Ran-GTP should exist throughout cells. Interestingly, both HURP and

importin-β localized throughout the spindle with increased intensities in RanGAP1-

depleted cells (Fig. 5H-I, S5E). These results suggest that HURP and importin-β act

together and interact with microtubules preferentially in the presence of Ran-GTP (Fig.

5J).

HURP dynamically associates with k-fibers in the presence of importin-β.

Based on our results, we developed a local cycling model for activation and polarization

of HURP (Fig. 5J). In this model, importin-β inhibits HURP globally, including at k-fibers,

by masking HURP's 2nd microtubule-binding domain (MTBD2). The resulting HURP-

importin-β complex binds weakly to microtubules through HURP's MTBD1 [43], but the

Ran-GTP gradient locally releases importin-β from HURP, resulting in full activation of

HURP near chromosomes (Fig. 5J). To test this model, we first performed fluorescence

recovery after photobleaching (FRAP) for HURP, and analyzed its dynamics on spindle

microtubules in the presence and absence of importin-β. In control cells, HURP was

quickly recovered at k-fibers after bleaching (Fig. 6A top, 6B black, S6A $t_{1/2}$ = 20.5 sec).

In contrast, HURP's fluorescent signals were hardly seen on the spindle in importin-β-

depleted cells (Fig. 6A bottom, 6B red, S6B). These results indicate that HURP

dynamically associates with k-fibers in the presence of importin-β, whereas HURP binds

tightly to spindle microtubules in the absence of importin-β.

HURP is dynamically maintained at k-fibers during metaphase

mAID-tag fusion proteins can be rapidly degraded with a half-life of 20 min [35]. To

confirm the dynamic regulation of HURP by importin-β and Ran-GTP, we next sought to

degrade importin-β during metaphase by combining AID-mediated degradation with

APC/C inhibitors [44] (Fig. 6C). Following treatment with the APC/C inhibitors, Apcin

and proTAME, cells arrested at metaphase, in which both importin-β and HURP

accumulated at k-fibers near chromosomes (Fig. 6 D, t = 0). Importantly, importin-β-

mAC signals diminished to undetectable levels 60-90 min after addition of IAA (Fig. 6D,

arrows), and HURP relocated from k-fibers to spindle microtubules in response to the

reduction of importin-β signals (Fig. 6D, E).

To confirm these results, we next acutely degraded RCC1 in metaphase-arrested

cells. As with importin-β degradation, HURP dissociated from k-fibers and localized

weakly on the spindle in response to degradation of RCC1 (Fig. 7A, B). Unexpectedly,

in contrast to the prometaphase degradation assay (Fig. 2H), spindle length appeared

normal when RCC1 was degraded in the metaphase-arrested condition (Fig. 7C).

Together, these results indicate that HURP is dynamically maintained at k-fibers near

chromosomes by the Ran-Importin pathway, even in metaphase, but HURP is critical for

spindle length regulation primarily during prometaphase.

Discussion

NuMA is liberated from importins independently of Ran-GTP for spindle-pole

focusing in human mitotic cells

In contrast to the prevailing model (Fig. 2A), we demonstrated that the Ran-Importin

pathway is dispensable for localization and functions of NuMA at the spindle poles in

human HCT116 cells (Fig. 2, 3, 7D right). This is consistent with the recent observation

that NuMA is less sensitive to Ran-GTP level than to HSET/XCTK2 [29]. Although we

do not exclude the possibility that Ran-GTP liberates NuMA from importin-α/β

complexes near chromosomes, we favor the idea that parallel pathways exist to activate

NuMA in mitotic human cells. In fact, recent studies indicate that importin-α/β-binding

TPX2 can be activated not only by Ran-GTP, but also by Golgi- or palmitoylation-

dependent sequestration of importin-α [45, 46]. In addition, mitotic spindles contain

centrosomes, which may generate special signals that liberate NuMA from inhibitory

importins (Fig. 7D). Interestingly, NuMA is broadly distributed on a bundle-like structure

between the poles in human acentrosomal cells [47]. It is necessary to analyze whether

NuMA is preferentially regulated by Ran-GTP in acentrosomal cells, especially in

oocytes, where Ran-GTP governs meiotic spindle assembly [12].

Although NLS-containing SAFs are recognized by importin-α, structural studies

indicate that importin-α binds to NuMA and TPX2 with slightly different binding patches

[32]. In addition, whereas TPX2-NLS and NLS-binding sites of importin-α are well

conserved in vertebrates, NLS of NuMA is not well conserved in fish (Fig. S1D-F).

Furthermore, NLS is lacking in other NuMA-like proteins in lower eukaryotes, such as

Caenorhabditis elegans LIN-5, *Drosophila* Mud, and yeast Num1 [38, 48, 49],

suggesting that NuMA acquired NLS in higher animals and is likely to be regulated

differently than TPX2. Future research should be undertaken to understand how the

NuMA-importin interaction is regulated in a Ran-independent manner, and why NLS-

dependent regulation of NuMA was acquired in higher animals.

The Ran-Importin pathway locally activates and polarizes HURP by promoting its

microtubule binding-dissociation cycle near chromosomes

In contrast to NuMA, we demonstrated that HURP is preferentially regulated by the

Ran-Importin pathway in mitotic human cells (Fig. 4C, 5E, H). Although HURP has been

identified previously as a downstream target of Ran-GTP [31], we unexpectedly found

that HURP also colocalizes with importin-β on k-fibers near chromosomes (Fig. S3C,

Fig. 5A, E), and stabilizes k-fibers independently of importin-β (Fig. 5D, G). In addition,

HURP's spindle distribution is sensitive to levels of Ran-GTP and importin-β (Fig. 4C,

5E, H), and is dynamically and spatially maintained during metaphase in a Ran-

pathway-dependent manner (Fig. 6A, B, D, 7A). Based on these results, we propose a

local cycling model for establishment and maintenance of HURP's polarized localization

to spindle microtubules (Fig. 5J, 7D left). After nuclear envelope breakdown (NEBD),

HURP strongly interacts with microtubules through its two microtubule-binding domains

(MTBD1 and MTBD2 in Fig. 5J) [31, 43]. Since importin-β is localized diffusely

throughout cells (Fig. 3E, 5E), it binds to HURP on microtubules, and then dissociates

HURP from the microtubules by masking HURP's MTBD2 domain [43]. However, in the

vicinity of chromosomes, Ran-GTP releases HURP from importin-β [31], and the

liberated HURP interacts strongly with microtubules around chromosomes. By repeating

this local binding-dissociation cycle, HURP, but not importin-β, stabilizes microtubules

and generates stable k-fibers near chromosomes (Fig. 5D, G). This dynamic regulation

is similar to that of HSET/XCTK2 [50] and would be suitable for bundling short

microtubules around kinetochores during prometaphase [51] and for coupling HURP's

polarized localization with microtubule flux on the metaphase spindle.

RCC1 is required to define proper spindle length during prometaphase

RCC1 depletion causes shortened bipolar spindles in human cells (Fig. 2B, C, F, H).

This is probably due to multiple defects in spindle assembly processes, including the

lack of HURP-based k-fiber formation (Fig. 5C, D) and HSET-dependent microtubule-

sliding (Fig. 4B) [28]. Intriguingly, our mitotic degradation assays indicate that Ran-GTP

controls spindle length primarily during prometaphase, rather than in metaphase (Fig.

2H, 7C). Once metaphase spindles are assembled, other k-fiber localized proteins, such

as clathrin, TACC3, and ch-TOG [52], may be able to maintain bundled-k-fibers during

metaphase in a Ran-independent manner.

In addition, our mitotic degradation assay revealed that RCC1-depletion does

not affect mitotic progression (Fig. S2J). This suggests that the mitotic delay observed

in RCC1 depletion in asynchronous culture (Fig. S2F) is a secondary defect caused by

loss of interphase RCC1 activity. In fact, ectopically expressed HSET-NLS mutants

localize in cytoplasm and causes abnormal cytoplasmic microtubule-bundling in

interphase [30]. Numerous similar defects would be created by RCC1 depletion in

interphase and would affect subsequent mitotic progression.

A new toolkit and mitosis-specific degradation assays to dissect mitotic roles of

the Ran-importin pathway

As discussed above, mitotic inactivation is critical to precisely analyze mitotic functions

of Ran-GTP and importins. Previously, tsBN2, a temperature-sensitive RCC1 mutant

hamster cell line [53, 54] and a small molecule inhibitor, importazole [55], have been

developed to acutely inhibit functions of RCC1 and importin-β, respectively. Here, we

established three human AID-cell lines for RCC1, RanGAP1, and importin-β [35], and

succeeded in degrading RCC1 and importin-β specifically in prometaphase (Fig. 2C-F)

or metaphase (Fig. 6C-E, 7A-B). Because these AID-cell lines and mitotic degradation

assays are applicable to other Ran-regulated SAFs/cortical proteins [4, 54, 56] and

other multi-functional proteins such as dynein and NuMA [20, 35], respectively, these

toolkits and assays will further advance our understanding of mechanisms and roles of

spindle assembly, maintenance, and positioning in animal cells.

Acknowledgments

We thank Iain M. Cheeseman for critical reading of the manuscript, and Yuki Tsukada, Rie Inaba and Kiyoko Murase for technical assistance. This work was supported by grants from the PRESTO program (JPMJPR13A3) of the Japan Science and Technology agency (JST) for T.K, a Career Development Award of the Human Frontier Science Program (CDA00057/2014-C) for T.K., KAKENHI (16K14721 and 17H05002 for T.K, 17H01431 for G.G.) of the Japan Society for Promotion of Science (JSPS), NIG-JOINT (2014B-B-3, 2015-A1-19, 2016-A1-22 for T.K.) of National Institute of Genetics (NIG), the Naito Foundation for T.K, and JSPS and DFG under the Joint Research Projects-LEAD with UKRI for G.G.

Author contributions

Conceptualization, TK; Investigation, TK, KT, HH, MN, and MO; Formal analysis, TK and KT; Methodology, TK, YS, and MK; Writing, TK; Supervision, TK and GG; Funding Acquisition, TK and GG.

Declaration of interests

The authors declare no competing interests.

Figure Legends

Figure 1. NuMA acts in spindle pole focusing using its conserved microtubule-binding domain in human cells. (A) Full length NuMA and tested NuMA truncation fragments. NLS and a microtubule-binding domain (MTBD) are shown in magenta and green, respectively. (B and C) Interphase (B) and metaphase (C) Metaphase NuMA-mACF cell lines showing live fluorescent images of NuMA-mACF, NuMA-mCh WT or mutants, SiR-DNA and SiR-700 tubulin (TUB) after 24 hr following treatment with Dox and IAA. Arrows in C indicate unfocused microtubules. (D) Quantification of cells with unfocused spindles in each condition from data in (C). Bars indicate means ± SEMs. N = 47 (-/-), 75 (-/+), 31 (#1/+), 30 (#2/+), 31 (#3/+), 30 (#4/+), and 48 (#5/+) from 3 independent experiments. p-values calculated using Dunnett's multiple comparisons test after one-way ANOVA (F (3,6) = 33.81, p = 0.0004).

Figure 2. NuMA functions in spindle pole focusing independently of RCC1. (A) The prevailing model of SAF inhibition and activation by importins and Ran-GTP. (B) Metaphase RCC1-mAC cells showing live fluorescent images of RCC1mAC, NuMA-mCherry (mCh), and SiR-TUB after 24 hr following Dox and IAA treatment. (C) Scatterplots of the ratio of spindle length and cell diameter in controls (0.54 ± 0.04, n = 32) and RCC1-depleted (0.47 ± 0.04, n = 23) cells. Bars indicate means ± SDs from >3 independent experiments. * indicates statistical significance according to Welch's t-test (p < 0.0001). (D) Quantification of cells with unfocused spindles in each condition from data in (C) and (E). Bars indicate means ± SEMs. N = 27, 34, 37, and 113 from >4 independent experiments. p-values calculated using Dunnett's multiple comparisons test after one-way ANOVA (F (3,14) = 36.40, p < 0.0001). * indicates p < 0.0001. (E) Live fluorescent images of SiR-DNA, RCC1-mAC, NuMA-mAID-mCherry, and SiR700-TUB in RCC1-mAC and NuMA-mAID-mCh double knock-in cells following 24 hr of Dox and IAA treatment. Two cells with or without RCC1 and NuMA signals were analyzed in the same field. Eight z-section images were acquired using 1.0-μm spacing. Maximum intensity projection images are shown. (F, G) Live fluorescent images of RCC1-mAC and NuMA-mCh in RCC1-positive control (F) and RCC1-negative cells (G) treated with nocodazole and IAA, as described in Fig. S2I. * indicates RCC1-undegraded cells. (H) Scatterplots of the ratio of spindle length and cell diameter in control (0.57 ± 0.05, n = 35) and RCC1-depleted (0.50 ± 0.06, n = 30) cells. Bars indicate means ± SDs from >3 independent experiments. * indicates statistical significance according to Welch's t-test (p < 0.0001). (I) Quantification of spindle orientation on the x-z plane in control (n = 43) and RCC1-depleted (n = 42) cells from 2 independent experiments. See Methods for the definition of parallel and tilted orientations. * indicates statistical significance according to Z-test (significance level 0.1). Scale bars = 10 μm.

Figure 3. NuMA is liberated from importins at spindle poles independently of Ran-GTP. (A) A model showing NuMA liberation from importins in RCC1 depleted cells. (B) Metaphase RCC1-mAC cells showing live fluorescent images of RCC1-mAC, importin-β-mCh, and SiR-TUB and after 24 hr following treatment with Dox and IAA. Right:

Quantification of cells with k-fiber localization of importin-β in control (n > 40) and RCC1-depleted (n > 40) cells from 3 independent experiments. * indicates statistical significance according to Z-test (significance level 0.0001). (C) Live fluorescent images of NuMA-mClover-FLAG (mCF, left) and importin-α (right) wild type (WT, top) and a ΔIBB mutant (bottom). (D-E) Metaphase RanGAP1-mAC (D) and importin-β-mAC (E) cells showing live fluorescent images of NuMA-mCherry (mCh), SiR-tubulin (SiR-TUB), and RanGAP1-mAC (D) or importin-β-mAC (E) after 24 hr following treatment with Dox and IAA. * in E indicates cells with importin-β signals in the presence of Dox and IAA. Scale bars = 10 μm.

Figure 4. RCC1 regulates chromosome-proximal localization of HURP and HSET. (A-C) Left: Metaphase RCC1-mAC cells showing live fluorescent images of RCC1-mAC, SiR-TUB and TPX2-mCh (A), mCh-HSET (B), and HURP-mCh (C) after 24 hr following treatment with Dox and IAA. Right: Quantification of throughout spindle or k-fiber localization of HSET or HURP in control (n > 30) and RCC1-depleted (n > 40) cells from 2 or 3 independent experiments. * indicates statistical significance according to Z-test (significance level 0.0001). (D) A list summarizing localization of SAFs in control and RCC1-depleted cells. Scale bars = 10 μm.

Figure 5. HURP, but not importin-β, is required to stabilize k-fibers. (A) Metaphase HURP-mACF cells showing live fluorescent images of HURP-mACF, importin-β-mCh and SiR-TUB after 24 hr following Dox and IAA treatment. (B) Quantification of k-fiber localization of importin-β in control (n = 49) and HURP-depleted (n = 46) cells from 3 independent experiments. (C) Scatterplots of the ratio of spindle length and cell diameter in control (0.64 ± 0.05, n = 49) and HURP-depleted (0.52 ± 0.06, n = 43) cells. * indicates statistical significance according to Welch's *t*-test ($p < 0.0001$). (D) Fluorescent images of HURP-mACF, TUB, and DNA (Hoechst 33342 staining) in metaphase fixed cells treated with ice-cold medium for 20 min. Two cells with or without HURP signals were analyzed in the same field. (E) Metaphase importin-β-mAC cells showing live fluorescent images of importin-β-mAC, HURP-SNAP and SiR-TUB after 24 hr following treatment with Dox and IAA. (F) Quantification of spindle localization of HURP in control (n = 49) and importin-β-depleted (n = 43) cells from 3 independent experiments. (G) Fluorescent images of importin-β-mAC, HURP-SNAP, TUB, and DNA (Hoechst 33342 staining) in metaphase fixed cells treated with ice-cold medium for 20 min. Five z-section images were obtained using 0.5-μm spacing and maximum intensity projection images are shown in (D) and (G). (H-I) Left: metaphase RanGAP1-mAC cells showing live fluorescent images of RanGAP1-mAC, SiR-TUB and HURP-mCh (H) or importin-β-mCh (I) after 24 hr following Dox and IAA treatment. Right: quantification of k-fiber localization of HURP or importin-β in control (n = 45) and RanGAP1-depleted (n > 45) cells from 3 independent experiments. * in (B), (F), (H) and (I) indicates statistical significance according to Z-test (significance level 0.0001). (J) A local cycling model of HURP on k-fibers regulated by Ran-GTP and importin-β. See text for details. Scale bars = 10 μm.

Figure 6. HURP dynamically accumulates on metaphase k-fibers in an importin-β-dependent manner. (A) Live fluorescent images of HURP-SNAP visualized with TMR-star (magenta) and SiR-tubulin (TUB) in control (top) and importin-β-depleted cells (bottom). Fluorescent signals were bleached in the indicated box region at t = 0, and the fluorescence recoveries were monitored for 120 sec. (B) A graph showing fluorescence recovery after photobleaching. An average of 7 samples was plotted. Bars indicate SDs. (C) Schematic diagram of the metaphase degradation assay. Following release from RO-3336-mediated G2 arrest, proTAME and Apcin were added to arrest cells in metaphase. Auxin (IAA) was added (indicated by the red line) to induce RCC1 degradation during metaphase. (D) Live fluorescent images of SiR-DNA, importin-β-mAC, HURP-mCh, and SiR-700-tubulin (TUB). IAA was added at t = 0. Arrows indicate a cell showing a reduction of importin-β signal during metaphase. (E) Enlarged images from (D) showing a re-localization of HURP-mCh from k-fibers (t = 0) to the spindle (t =90). Scale bars = 10 μm.

Figure 7. Models of local activation mechanisms for HURP and NuMA in mitosis. (A) Live fluorescent images of SiR-DNA, RCC1-mAC, HURP-mCh, and SiR-700-tubulin (TUB). IAA was added at t = 0. (B) Enlarged images of indicated regions in (A) showing a reduction of HURP-mCh from k-fibers in response to degradation of RCC1. (C) Spindle length measurement (n = 6) at t = -5 and 60 min in (A). (D) Left: in the vicinity of chromosomes, Ran-GTP and importin-β promote the microtubule binding and dissociation cycle of HURP, resulting in polarized HURP accumulation and stable k-fiber formation. Right: chromosome-derived Ran-GTP is not required to activate NuMA at the spindle poles in mitotic human cells. A Ran-independent, parallel pathway would exist to activate NuMA away from chromosomes. See text for details. Scale bars = 10 μm.

Materials and Methods

- Plasmid Construction

Plasmids for CRISPR/Cas9-mediated genome editing and auxin-inducible degron were constructed according to protocols of Natsume et al. [35] and Okumura et al., [20]. To construct donor plasmids containing homology arms for RCC1 (~500-bp homology arms), RanGAP1 (~500-bp), importin-β (~500-bp), HURP (~200-bp), TPX2 (~200-bp), and HSET (~200-bp), gene synthesis services from Eurofins Genomics K.K. (Tokyo, Japan) or Genewiz (South Plainsfield, NJ) were used. Plasmids and sgRNA sequences used in this study are listed in Supplementary Tables S1 and S2, and will be deposited in Addgene.

- Cell Culture, Cell Line Generation, and Antibodies

HCT116 cells were cultured as described previously [20]. Knock-in cell lines were generated according to procedures described in Okumura et al. [20]. To activate auxin-inducible degradation, cells were treated with 2 μg/mL Dox and 500 μM indoleacetic acid (IAA) for 20–24 hr. Cells with undetectable signals for mAID-fusion proteins were analyzed. Flip-In T-REx 293 cells were used in Figure 3C to express mCherry-tagged importin-α constructs. Cell lines were created according to procedures described in Kiyomitsu et al. [57]. To induce transgenes, cells were incubated with 1 μg/mL tetracycline (MP Biomedicals). Cell lines and primers used in this study are listed in Tables S1 and S3, respectively.

Antibodies against tubulin (DM1A, Sigma-Aldrich, 1:2,000), NuMA (Abcam, 1:1,000), RCC1 (Cell Signaling Technology, D15H6, Rabbit mAb, 1:100), RanGAP1 (Santa Cruz Biotechnology, H-180, 1:200), importin-β (GeneTex, 3E9 Mouse mAb, 1:100), and HURP (E. Nigg laboratory, 1 : 200) were used for western blotting. For RCC1 immunoblots, membranes were incubated with anti-RCC1 antibody overnight at 4 °C.

- Microscope System

Imaging was performed using spinning-disc confocal microscopy with a 60× 1.40 numerical aperture objective lens (Plan Apo λ, Nikon, Tokyo, Japan). A CSU-W1 confocal unit (Yokogawa Electric Corporation, Tokyo, Japan) with five lasers (405, 488, 561, 640, and 685 nm, Coherent, Santa Clara, CA) and an ORCA-Flash 4.0 digital CMOS camera (Hamamatsu Photonics, Hamamatsu City, Japan) were attached to an ECLIPSE Ti-E inverted microscope (Nikon) with a perfect focus system. DNA images in Figure 2A/B or Figure 4D/G were obtained using either a SOLA LED light engine (Lumencor, Beaverton, OR) or a 405-nm laser, respectively.

- Immunofluorescence and Live Cell Imaging

For immunofluorescence in Figure S1K, HURP-mACF cells were fixed with PBS containing 3% paraformaldehyde and 2% sucrose for 10 min at room temperature. Fixed cells were permeabilized with 0.5% Triton X-100™ for 5 min on ice, and pretreated with PBS containing 1% BSA for 10 min at room temperature after washing with PBS. Importin-β was visualized using anti-importin-β antibody (1:500). Images of multiple z-sections were acquired by spinning-disc confocal microscopy using 0.5-µm

spacing and camera binning 2. Maximally projected images from 3 z-sections are shown.

   For live cell imaging, cells were cultured on glass-bottomed dishes (CELLview™, #627860 or #627870, Greiner Bio-One, Kremsmünster, Austria) and maintained in a stage-top incubator (Tokai Hit, Fujinomiya, Japan) to maintain the same conditions used for cell culture (37° C and 5% CO_2). In most cases, three to five z-section images using 0.5-µm spacing were acquired and single z-section images are shown, unless otherwise specified. Microtubules were stained with 50 nM SiR-tubulin or SiR700-tubulin (Spirochrome) for >1 hr prior to image acquisition. DNA was stained with 50 ng/mL Hoechst® 33342 (Sigma-Aldrich) or 20 nM SiR-DNA (Spirochrome) for > 1 hr before observation. To visualize SNAP-tagged HURP in Fig. 4E, cells were incubated with 0.1 µM TMR-Star (New England BioLabs) for > 2 hr, and TMR-Star were removed before observation. To optimize image brightness, the same linear adjustments were applied using Fiji and Photoshop.

· Prometaphase degradation assay and nocodazole washout
To degrade mAID-tagged proteins during nocodazole arrest, cells were treated with 2 µg/mL Dox and 3.3 µM nocodazole at the indicated times (Fig. 2C). Five hours after addition of nocodazole, cell culture dishes were moved to the stage of a microscope equipped with a peristaltic pump (SMP-21S, EYELA, Tokyo Rikakikai). Two z-section images were acquired using 2-µm spacing at three different (X.Y) positions and at 5-min intervals, with 500 µM IAA added during the first interval. After 90 min, the nocodazole-containing medium was completely replaced with fresh medium using the peristaltic pump at a velocity of 20 sec/mL for 15 min. Images were acquired for a further 2 hr and maximum intensity projection images are shown in Figure 2D-F. To analyze spindle orientation in Figure 2I, we took five z-section images using 2-µm spacing. When both spindle poles are included within three z-section images, we judged the spindle as having parallel orientation.

· Metaphase degradation assay
To degrade mAID-tagged proteins in metaphase-arrested cells, cells were treated with 50 µM Apcin (I-444, Boston Biochem) and 20 µM proTAME (I-440, Boston Biochem) at the indicated times (Fig. 5A). Three z-section images were acquired using 1-µm spacing at six different (X,Y) positions and at 5-min intervals, with 500 µM IAA added during the first interval. Maximum intensity projection images are shown in Figure 5B.

· Cold treatment assay
To increase the number of cells in metaphase, cells were treated with 20 µM MG132 (C2211, Sigma-Aldrich) for 90 min. To visualize SNAP-tagged HURP (Fig. 4G), cells were incubated with 0.1 µM TMR-Star (S9105S, New England BioLabs) for at least 30 min. Before fixation, cells were incubated in ice-cold medium for 20 min [31] to depolymerize non-kinetochore microtubules.

· FRAP
FRAP was conducted with a microscope (LEM 780, Carl Zeiss MicroImaging, Inc.), using a 63 x objective lens. Images were acquired every 5 sec before and after

photobleaching. The bleached area (BA) was set as it covers half spindle and illuminated at t = 0 using 560 nm laser (20 mW) with the following setting: speed 4.0 and iteration 1. Metaphase cells that orient parallel to the bottom cover-glass were selected. HURP (TMR-Star) intensity of BA was normalized using the intensity of non-bleached area (NBA) that covers the remaining half spindle. Corrected relative intensity at time t_n was calculated as $(BA_n - BG_n) / (BA_{-1} - BG_{-1})$ x $(NBA_{-1} - BG_{-1}) / (NBA_n - BG_n)$, where t = -1 represents the first time point of image acquisition before bleaching. BG means background [58]. Curve fitting and analyses shown in Fig. S6 were performed using Fiji.

· Statistical Analysis

To determine the significance of differences between the mean values obtained for two experimental conditions, Welch's *t*-tests (Prism 6; GraphPad Software, La Jolla, CA) or a Z-test for proportions (epitools.ausvet.com.au/ztesttwo) were used, as indicated in figure legends.

Supplemental Information

Supplemental Figure Legends

Figure S1. Generation of cell lines that conditionally degrade endogenous NuMA and express NuMA mutants. (A) Schematic of the auxin-inducible degradation (AID) system. OsTIR1, an F-box protein expressed following Dox treatment, forms SCF E3 ubiquitin ligase complexes. Following auxin (IAA) treatment, mAID-fusion protein was poly-ubiquitinated by SCF^{OsTIR1} and degraded by proteasomes with a half-life of ~20 min. (B) Genomic PCR showing clone genotypes after hygromycin (Hygro) selection. Clones used in this study are listed in Table S1. (C) Interphase NuMA-mACF cell lines showing live fluorescent images of NuMA-mACF, NuMA-mCh mutants, and SiR-DNA after 24 hr following treatment with Dox and IAA. Endogenous NuMA-mACF signals were undetectable, whereas ectopically expressed NuMA mutants were detected in cytoplasm. NuMA-mCh Δ(NLS+MTBD2) appeared to accumulate on microtubules around centrosomes. (D) Amino acid sequence alignment of the NLS of NuMA proteins in *H. sapiens* (NP_006176), *R. norvegicus* (NP_001094161), *M. musculus* (NP_598708), *G. gallus* (NP_001177854), *X. laevis* (NP_001081559), *D. rerio* (NP_001316910), *O. latipes* (XP_020564048), and *A. ocellaris* (XM_023273896) aligned by ClustalWS. NLSs are not well conserved in fish, although NuMA clustering motif (shown in orange) and NLM motif (sky blue) are highly conserved in vertebrates. In the NLS alignment, key amino acids that interact with importin-α [32] are boxed in red, whereas positively charged amino acids in fish are boxed in green. (E) Amino acid sequence alignment of the major (i) and minor (ii) NLS-binding site of importin-α proteins in *H. sapiens* (NP_001307540), *M. musculus* (NP_034785), *G. gallus* (NP_001006209), *X. laevis* (NP_001080459), *D. rerio* (NP_001002335), and *O. latipes* (XP_023816136) aligned by ClustalWS. (F) Amino acid sequence alignment of the NLS of TPX2 proteins in *H. sapiens* (NP_036244), *M. musculus* (NP_001135447), *G. gallus* (NP_989768), *X. laevis* (AAH68637), *D. rerio* (NP_001314674), and *O. latipes* (XP_020557297) aligned by ClustalWS. Key amino acids interact with importin-α [33] are boxed in red.

Figure S2. Generation of cell lines for auxin-inducible degradation of endogenous RCC1. (A) Genomic PCR showing clone genotypes after neomycin (Neo) selection. Clone No.1 was used as a parental cell in subsequent selections. * indicates a non-specific band. (B) Genomic PCR showing clone genotypes after hygromycin (Hygro) selection. Clone No.1 was used in this study. (C) Immunoblotting for anti-NuMA, anti-RCC1, and anti-α-tubulin (TUB, loading control) showing bi-allelic insertion of the indicated tags. (D-E) Live fluorescent images of DNA (Hoechst 33342 staining), RCC1-mAC, NuMA-mCh, and SiR-TUB in control (D) and RCC1-depleted (E) cells. (F) Scatterplots of mitotic duration (NEBD to anaphase onset) in control (34.1 ± 7.6, n=32) and RCC1-depleted cells (47.2 ± 10.5, n=27). Bars indicate means ± SDs from >3 independent experiments. * indicates statistical significance according to Welch's *t*-test (p<0.0001). (G) Genomic PCR showing clone genotype after hygromycin (Hygro) selection. Clone No.3 was selected for further use. (H) Live fluorescent images of SiR-DNA, RCC1-mAC, NuMA-mAID-mCh, and SiR-TUB. A spindle-pole focusing defect

(indicated by the arrow in panel 2) and abnormal spindle formation (panel 3) were
observed in RCC1-mAC and NuMA-mAID-mCh co-depleted cells 20-24 hr after Dox
and IAA treatment. Five z-section images were acquired using 1.0-µm spacing and
maximum intensity projection images are shown. (I) (C) Schematic diagram of the
prometaphase degradation assay. Nocodazole was added to arrest the cells in
prometaphase, and then Auxin (IAA) was added to induce RCC1 degradation during
nocodazole-arrest. Nocodazole were washed out by changing medium for 15 min with
peristaltic pumps, while recording the cells. See Methods for details. (J) Scatterplots of
mitotic duration (from nocodazole wash-out to anaphase onset) in RCC1-positive
control (68.7 ± 2.1, n=46) and RCC1-depleted cells (61.5 ± 1.4, n=47). Bars indicate
means ± SDs from >3 independent experiments. * indicates statistical significance
according to Welch's *t*-test (p = 0.0061). Scale bars = 10 µm.

Figure S3. Generation of cell lines for auxin-inducible degradation of endogenous
Ran-GAP1 and importin-β. (A) Genomic PCR showing clone genotypes after
hygromycin (Hygro) selection. Clones No.6 were used. (B) Metaphase importin-β-mAC
cells showing live fluorescent images of importin-β-mAC, and SiR-TUB. Single z-section
images are shown. (C) Immunofluorescence images of fixed metaphase cells showing
k-fiber localization endogenous importin-β and mAID-tagged HURP (HURP-mACF).
Maximally projected images from 3 z-sections are shown. (D) Immunoblotting for anti-
importin-α and anti-α-tubulin (TUB, loading control) showing ectopic expression of the
importin-α wild type (WT, right) and a ΔIBB mutant (left) following Dox treatment. *
indicates endogenous importin-α. (E) Genomic PCR showing clone genotypes after
neomycin (Neo) selection. Clone No.9 was used as a parental cell in the second
selections. (F) Genomic PCR showing clone genotypes after hygromycin (Hygro)
selection. Clone No.3 (NuMA-mCh) was selected. (G) Immunoblotting for anti-NuMA,
anti-RanGAP1 and anti-α-tubulin (TUB, loading control) showing bi-allelic insertion of
the indicated tags. * and ** indicate RanGAP1 and SUMO-1 conjugated RanGAP1,
respectively. (H) Scatterplots of the ratio of spindle length and cell diameter in control
(0.54 ± 0.04, n=26) and RanGAP1-depleted (0.52 ± 0.07, n=19) cells. (I) Scatterplots of
mitotic duration (NEBD to anaphase onset) in control (35.5 ± 9.0, n=29) and RanGAP1-
depleted (39.1 ± 10.1, n=23) cells. Bars in (H) and (I) indicate means ± SDs from >3
independent experiments. Differences were not statistically significant based on Welch's
t-test in H (p=0.2108) and I (p=0.1851). (J) Genomic PCR showing clone genotypes
after neomycin (Neo) selection. Clone No.7 was used as a parental cell in the second
selections. (K) Genomic PCR showing clone genotype after hygromycin (Hygro)
selection. Clone No.1 was selected for further use. (L) Western blot detection using anti-
NuMA, anti-importin-β and anti-α-tubulin antibodies (TUB, loading control) showing bi-
allelic insertion of the indicated tags. (M) Scatterplots of the ratio of spindle length and
cell diameter in control (0.49 ± 0,05, n=26) and importin-β-depleted (0.44 ± 0.07, n=17)
cells. (N) Scatterplots of mitotic duration (NEBD to anaphase onset) in control (41.9 ±
16.3, n=27) and importin-β-depleted (66.7 ± 26.7, n=12) cells. Bars in (M) and (N)
indicate mean ± SD from >3 independent experiments. * indicates statistical significance
according to Welch's *t*-test (p<0.05) in (M) and (N). Scale bars = 10 µm.

Figure S4. Generation of double knock-in cell lines that express RCC1-mAC and
mCherry-fused TPX2, HSET, or HURP. (A-C) Genomic PCR showing clone genotypes
after hygromycin (Hygro) selections. Clones No.1 (A), No. 1 (B), and No.8 (C) were
used. The mCherry cassette was inserted into only one copy of TPX2 gene loci (A).

Figure S5. Generation of cell lines that degrade or visualize endogenous HURP.
(A) Genomic PCR showing the clone genotype after neomycin (Neo) selection. Clone
No.13 was used as a parental cell in subsequent selections. (B) Immunoblotting for anti-
HURP and anti-α-tubulin (TUB, loading control) showing bi-allelic insertion of the
indicated tags. (C-E) Genomic PCR showing clone genotypes after hygromycin (Hygro)
selection. Clone No.14 (C), No. 3 (D), No. 5 (E: HURP-mCh), and No. 12 (E: importin-β)
were used, respectively. The SNAP cassette was inserted into only one copy of HURP
gene loci (D).

Figure S6. Fluorescent recovery kinetics of HURP in the presence or absence of
importin-β. (A-B) Graphs showing a fitted curve or a straight line on each plot.
Formulas and parameters are also indicated.

 Table S1: Cell lines used in this study.

No.	Name	Description	Clone No.	Plasmids used	Parental cell	Reference
1	HCT116 tet-OsTIR1	AAVS1::PTRE3G OsTIR1 (Puro)		pAAVS1 T2 and MK243 (Addgene#72835)		[35]
2	NuMA-mACF + DHC-SNAP + mCh-NuMA WT	AAVS1::PTRE3G OsTIR1 (Puro), NuMA1:: NuMA-mAID-mClover-3FLAG (Neo), DHC1:: DHC-SNAP (BSD), Rosa26:: PTRE3G mCherry-NuMA WT (Hygro)	7	hROSA26 CRISPR-pX330 and pTK503		[20]
3	NuMA-mACF + DHC-SNAP + mCh-NuMA ΔNLS	AAVS1::PTRE3G OsTIR1 (Puro), NuMA1:: NuMA-mAID-mClover-3FLAG (Neo), DHC1:: DHC-SNAP (BSD), Rosa26:: PTRE3G mCherry-NuMA ΔNLS (Hygro)	1	hROSA26 CRISPR-pX330 and pTK699	2	This study
4	NuMA-mACF + DHC-SNAP + mCh-NuMAΔex24	AAVS1::PTRE3G OsTIR1 (Puro), NuMA1:: NuMA-mAID-mClover-3FLAG (Neo), DHC1:: DHC-SNAP (BSD), Rosa26:: PTRE3G mCherry-NuMA Δex24 (Hygro)	1	hROSA26 CRISPR-pX330 and pTK700	2	This study
5	NuMA-mACF + DHC-SNAP + mCh-NuMA Δ(NLS +MTBD2)	AAVS1::PTRE3G OsTIR1 (Puro), NuMA1:: NuMA-mAID-mClover-3FLAG (Neo), DHC1:: DHC-SNAP (BSD), Rosa26:: PTRE3G mCherry-NuMA Δ(NLS+MTBD2) (Hygro)	1	hROSA26 CRISPR-pX330 and pTK509	2	This study
6	NuMA-mACF + DHC-SNAP + mCh-NuMA ΔC-ter	AAVS1::PTRE3G OsTIR1 (Puro), NuMA1:: NuMA-mAID-mClover-3FLAG (Neo), DHC1:: DHC-SNAP (BSD), Rosa26:: PTRE3G mCherry-NuMA ΔC-ter (Hygro)	3	hROSA26 CRISPR-pX330 and pTK510	2	This study
7	Flip-In T-REx 293	Invitrogen				[57]
8	Flip-In T-REx 293 importin-α-WT mCherry	Flip-In:: importin-α-WT mCherry (hygro)	Poly clon al	pOG44 and pTK960	7	This study
9	Flip-In T-REx 293 importin-α-ΔIBB mCherry	Flip-In:: importin-α-ΔIBB mCherry (hygro)	Poly clon al	pOG44 and pTK961	7	This study
10	RCC1-mAC	AAVS1::PTRE3G OsTIR1 (Puro), RCC1:: RCC1-mAID-mClover (Neo)	1	pTK361+ pHH45	1	This study
11	RCC1-mAC + NuMA-mCh	AAVS1::PTRE3G OsTIR1 (Puro), RCC1:: RCC1-mAID-mClover (Neo), NuMA1:: NuMA-mCh (Hygro)	1	pTK372+ pTK435	10	This study
12	RanGAP1-mAC	AAVS1::PTRE3G OsTIR1 (Puro), RanGAP1:: RanGAP1-mAID-mClover (Neo)	9	pHH49 + pHH51	1	This study
13	RanGAP1-mAC + NuMA-mCh	AAVS1::PTRE3G OsTIR1 (Puro), RanGAP1:: RanGAP1-mAID-mClover (Neo), NuMA1:: NuMA-mCh (Hygro)	5	pTK372+ pTK435	12	This study
14	importin-β-mAC	AAVS1::PTRE3G OsTIR1 (Puro), importin-β:: importin-β-mAID-mClover (Neo)	7	pHH50 + pHH57	1	This study
15	importin-β-mAC + NuMA-mCh	AAVS1::PTRE3G OsTIR1 (Puro), importin-β:: importin-β-mAID-mClover (Neo), NuMA1:: NuMA-mCh (Hygro)	1	pTK372+ pTK435	14	This study
16	RCC1-mAC + importin-β-mCh	AAVS1::PTRE3G OsTIR1 (Puro), RCC1:: RCC1-mAID-mClover (Neo), NuMA1:: NuMA-mCh (Hygro)	6	pHH50 + pTK481	10	This study

17	RCC1-mAC + HURP-mCh	AAVS1::PTRE3G OsTIR1 (Puro), RCC1:: RCC1-mAID-mClover (Neo), HURP:: HURP-mCh (Hygro)	8	pTK532+ pTK541	10	This study
18	RCC1-mAC + TPX2-mCh	AAVS1::PTRE3G OsTIR1 (Puro), RCC1:: RCC1-mAID-mClover (Neo), TPX2:: TPX2-mCh (Hygro)	1	pTK527+ pTK502	10	This study
19	RCC1-mAC + mCh-HSET	AAVS1::PTRE3G OsTIR1 (Puro), RCC1:: RCC1-mAID-mClover (Neo), HSET:: mCh-HSET (Hygro)	1	pTK523+ pTK531	10	This study
20	RanGAP1-mAC + HURP-mCh	AAVS1::PTRE3G OsTIR1 (Puro), RanGAP1:: RanGAP1-mAID-mClover (Neo), HURP:: HURP-mCh (Hygro)	5	pTK532+ pTK541	12	This study
21	RanGAP1-mAC + importin-β-mCh	AAVS1::PTRE3G OsTIR1 (Puro), RanGAP1:: RanGAP1-mAID-mClover (Neo), importin-β:: importin-β-mCh (Hygro)	12	pHH50 + pTK481	12	This study
22	importin-β-mAC + HURP-SNAP	AAVS1::PTRE3G OsTIR1 (Puro), importin-β:: importin-β-mAID-mClover (Neo), HURP:: HURP-SNAP (Hygro)	3	pTK532+ pTK589	14	This study
23	HURP-mACF	AAVS1::PTRE3G OsTIR1 (Puro), HURP:: HURP-mAID-mClover-3FLAG (Neo)	13	pTK532+ pTK596	1	This study
24	HURP-mACF + importin-β-mCh	AAVS1::PTRE3G OsTIR1 (Puro), HURP:: HURP-mAID-mClover-3FLAG (Neo), importin-β:: importin-β-mCh (Hygro)	14	pHH50 + pTK481	23	This study

759
760

761 **Table S2: sgRNA sequences for CRISPR/Cas9-mediated genome editing**

Gene locus	sgRNA (5'-3')	PAM	Plasmid Name
NuMA1 (C-terminus)	gtgggggccactcactggtac	tgg	pTK372 [20]
RCC1 (C-terminus)	gactgtatgctggcccccgc	tgg	pTK361
RanGAP1 (C-terminus)	tctgctgcagacgctgtaca	agg	pHH49
importin-β (C-terminus)	agttcgagccgccgcccgaa	agg	pHH50
HURP	caaaattctcctggttgtag	agg	pTK532
TPX2	tgcggataccgcccggcaat	ggg	pTK527
HSET	tgcattcccccggcgcgtgt	ggg	pTK523

763 **Table S3: PCR primers used to confirm gene editing**

Gene	Primer sequence	Primer name	Figures
RCC1	gaatgccattccaggcag	oHH88	Figure S2A
RCC1	ttctgcacgttcctctgg	oHH89	Figure S2A
NUMA1	gagcctcaaagaaggccc	oTK542	Figure S2B, S2G, S3F, S3K
NUMA1	agcaggaaccagggcctac	oTK566	Figure S2B, S2G, S3F, S3K
RanGAP1	gctgccgcaggaccagggcttggtg	oHH93	Figure S3E
RanGAP1	attccctggcctatgtctgctggaa	oHH94	Figure S3E
HURP	ctcttgatggatactttactg	oTK749	Figure S4C, S5A, S5D, S5E
HURP	cccttgagaaagagtatatcta	oTK750	Figure S4C, S5A, S5D, S5E
importin-β	ggagtaaggagttttgagagtatcg	oHH97	Figure S3A, S3J, S5C, S5E
importin-β	aaatcttctctagagctaggcaacg	oHH98	Figure S3A, S3J, S5C, S5E
TPX2	tctgacatccctctcactg	oTK660	Figure S4A
TPX2	ggagtctaatcgagacattc	oTK661	Figure S4A
HSET	ggccctcggctgtggc	oTK766	Figure S4B
HSET	ctccccgggtgctctaag	oTK767	Figure S4B
Rosa 26	ggtgggaggcgcttgttc	oTK846	Figure S1B
mCherry-NuMA	ctgtggggtctgcaggat	oTK445	Figure S1B

References

1.   Reber, S., and Hyman, A.A. (2015). Emergent Properties of the Metaphase Spindle. Cold Spring Harb Perspect Biol *7*, a015784.

2.   Heald, R., and Khodjakov, A. (2015). Thirty years of search and capture: The complex simplicity of mitotic spindle assembly. J Cell Biol *211*, 1103-1111.

3.   Kalab, P., and Heald, R. (2008). The RanGTP gradient - a GPS for the mitotic spindle. J Cell Sci *121*, 1577-1586.

4.   Forbes, D.J., Travesa, A., Nord, M.S., and Bernis, C. (2015). Reprint of "Nuclear transport factors: global regulation of mitosis". Curr Opin Cell Biol *34*, 122-134.

5.   Bischoff, F.R., and Ponstingl, H. (1991). Catalysis of guanine nucleotide exchange on Ran by the mitotic regulator RCC1. Nature *354*, 80-82.

6.   Bischoff, F.R., Klebe, C., Kretschmer, J., Wittinghofer, A., and Ponstingl, H. (1994). RanGAP1 induces GTPase activity of nuclear Ras-related Ran. Proc Natl Acad Sci U S A *91*, 2587-2591.

7.   Kalab, P., Weis, K., and Heald, R. (2002). Visualization of a Ran-GTP gradient in interphase and mitotic Xenopus egg extracts. Science *295*, 2452-2456.

8.   Kalab, P., Pralle, A., Isacoff, E.Y., Heald, R., and Weis, K. (2006). Analysis of a RanGTP-regulated gradient in mitotic somatic cells. Nature *440*, 697-701.

9.   Dumont, J., Petri, S., Pellegrin, F., Terret, M.E., Bohnsack, M.T., Rassinier, P., Georget, V., Kalab, P., Gruss, O.J., and Verlhac, M.H. (2007). A centriole- and RanGTP-independent spindle assembly pathway in meiosis I of vertebrate oocytes. J Cell Biol *176*, 295-305.

10.   Moutinho-Pereira, S., Stuurman, N., Afonso, O., Hornsveld, M., Aguiar, P., Goshima, G., Vale, R.D., and Maiato, H. (2013). Genes involved in centrosome-independent mitotic spindle assembly in Drosophila S2 cells. Proc Natl Acad Sci U S A *110*, 19808-19813.

11.   Hasegawa, K., Ryu, S.J., and Kalab, P. (2013). Chromosomal gain promotes formation of a steep RanGTP gradient that drives mitosis in aneuploid cells. J Cell Biol *200*, 151-161.

12.   Holubcova, Z., Blayney, M., Elder, K., and Schuh, M. (2015). Human oocytes. Error-prone chromosome-mediated spindle assembly favors chromosome segregation defects in human oocytes. Science *348*, 1143-1147.

13.   Drutovic, D., Duan, X., Li, R., Kalab, P., and Solc, P. (2020). RanGTP and importin beta regulate meiosis I spindle assembly and function in mouse oocytes. EMBO J *39*, e101689.

14.   Furuta, M., Hori, T., and Fukagawa, T. (2016). Chromatin binding of RCC1 during mitosis is important for its nuclear localization in interphase. Mol Biol Cell *27*, 371-381.

15.   Nachury, M.V., Maresca, T.J., Salmon, W.C., Waterman-Storer, C.M., Heald, R., and Weis, K. (2001). Importin beta is a mitotic target of the small GTPase Ran in spindle assembly. Cell *104*, 95-106.

16.   Wiese, C., Wilde, A., Moore, M.S., Adam, S.A., Merdes, A., and Zheng, Y. (2001). Role of importin-beta in coupling Ran to downstream targets in microtubule assembly. Science *291*, 653-656.

17.   Gruss, O.J., Carazo-Salas, R.E., Schatz, C.A., Guarguaglini, G., Kast, J., Wilm, M., Le Bot, N., Vernos, I., Karsenti, E., and Mattaj, I.W. (2001). Ran induces spindle assembly by reversing the inhibitory effect of importin alpha on TPX2 activity. Cell *104*, 83-93.

18.   Stewart, M. (2007). Molecular mechanism of the nuclear protein import cycle. Nat Rev Mol Cell Biol *8*, 195-208.

19.   Hueschen, C.L., Kenny, S.J., Xu, K., and Dumont, S. (2017). NuMA recruits dynein activity to microtubule minus-ends at mitosis. Elife *6*.

20.   Okumura, M., Natsume, T., Kanemaki, M.T., and Kiyomitsu, T. (2018). Dynein-Dynactin-NuMA clusters generate cortical spindle-pulling forces as a multi-arm ensemble. Elife *7*.

21.   Gaglio, T., Saredi, A., and Compton, D.A. (1995). NuMA is required for the organization of microtubules into aster-like mitotic arrays. J Cell Biol *131*, 693-708.

22.   Silk, A.D., Holland, A.J., and Cleveland, D.W. (2009). Requirements for NuMA in maintenance and establishment of mammalian spindle poles. J Cell Biol *184*, 677-690.

23.   Wittmann, T., Wilm, M., Karsenti, E., and Vernos, I. (2000). TPX2, A novel xenopus MAP involved in spindle pole organization. J Cell Biol *149*, 1405-1418.

24.   Garrett, S., Auer, K., Compton, D.A., and Kapoor, T.M. (2002). hTPX2 is required for normal spindle morphology and centrosome integrity during vertebrate cell division. Curr Biol *12*, 2055-2059.

25.   Petry, S., Groen, A.C., Ishihara, K., Mitchison, T.J., and Vale, R.D. (2013). Branching microtubule
821   nucleation in Xenopus egg extracts mediated by augmin and TPX2. Cell *152*, 768-777.

822   26.   Roostalu, J., Cade, N.I., and Surrey, T. (2015). Complementary activities of TPX2 and chTOG constitute
823   an efficient importin-regulated microtubule nucleation module. Nat Cell Biol *17*, 1422-1434.

824   27.   King, M.R., and Petry, S. (2020). Phase separation of TPX2 enhances and spatially coordinates microtubule
825   nucleation. Nat Commun *11*, 270.

826   28.   Ems-McClung, S.C., Zheng, Y., and Walczak, C.E. (2004). Importin alpha/beta and Ran-GTP regulate
827   XCTK2 microtubule binding through a bipartite nuclear localization signal. Mol Biol Cell *15*, 46-57.

828   29.   Ems-McClung, S.C., Emch, M., Zhang, S., Mahnoor, S., Weaver, L.N., and Walczak, C.E. (2020).
829   RanGTP induces an effector gradient of XCTK2 and importin alpha/beta for spindle microtubule cross-
830   linking. J Cell Biol *219*.

831   30.   Cai, S., Weaver, L.N., Ems-McClung, S.C., and Walczak, C.E. (2009). Kinesin-14 family proteins
832   HSET/XCTK2 control spindle length by cross-linking and sliding microtubules. Mol Biol Cell *20*, 1348-
833   1359.

834   31.   Sillje, H.H., Nagel, S., Korner, R., and Nigg, E.A. (2006). HURP is a Ran-importin beta-regulated protein
835   that stabilizes kinetochore microtubules in the vicinity of chromosomes. Curr Biol *16*, 731-742.

836   32.   Chang, C.C., Huang, T.L., Shimamoto, Y., Tsai, S.Y., and Hsia, K.C. (2017). Regulation of mitotic spindle
837   assembly factor NuMA by Importin-beta. J Cell Biol *216*, 3453-3462.

838   33.   Giesecke, A., and Stewart, M. (2010). Novel binding of the mitotic regulator TPX2 (target protein for
839   Xenopus kinesin-like protein 2) to importin-alpha. J Biol Chem *285*, 17628-17635.

840   34.   Kiyomitsu, T. (2019). The cortical force-generating machinery: how cortical spindle-pulling forces are
841   generated. Curr Opin Cell Biol *60*, 1-8.

842   35.   Natsume, T., Kiyomitsu, T., Saga, Y., and Kanemaki, M.T. (2016). Rapid Protein Depletion in Human
843   Cells by Auxin-Inducible Degron Tagging with Short Homology Donors. Cell Rep *15*, 210-218.

844   36.   Seldin, L., Muroyama, A., and Lechler, T. (2016). NuMA-microtubule interactions are critical for spindle
845   orientation and the morphogenesis of diverse epidermal structures. Elife *5*.

846   37.   Gallini, S., Carminati, M., De Mattia, F., Pirovano, L., Martini, E., Oldani, A., Asteriti, I.A., Guarguaglini,
847   G., and Mapelli, M. (2016). NuMA Phosphorylation by Aurora-A Orchestrates Spindle Orientation. Curr
848   Biol *26*, 458-469.

849   38.   Siller, K.H., Cabernard, C., and Doe, C.Q. (2006). The NuMA-related Mud protein binds Pins and
850   regulates spindle orientation in Drosophila neuroblasts. Nat Cell Biol *8*, 594-600.

851   39.   Du, Q., Stukenberg, P.T., and Macara, I.G. (2001). A mammalian Partner of inscuteable binds NuMA and
852   regulates mitotic spindle organization. Nat Cell Biol *3*, 1069-1075.

853   40.   Tang, T.K., Tang, C.J., Chao, Y.J., and Wu, C.W. (1994). Nuclear mitotic apparatus protein (NuMA):
854   spindle association, nuclear targeting and differential subcellular localization of various NuMA isoforms. J
855   Cell Sci *107 (Pt 6)*, 1389-1402.

856   41.   Kohler, M., Speck, C., Christiansen, M., Bischoff, F.R., Prehn, S., Haller, H., Gorlich, D., and Hartmann,
857   E. (1999). Evidence for distinct substrate specificities of importin alpha family members in nuclear protein
858   import. Mol Cell Biol *19*, 7782-7791.

859   42.   Ciciarello, M., Mangiacasale, R., Thibier, C., Guarguaglini, G., Marchetti, E., Di Fiore, B., and Lavia, P.
860   (2004). Importin beta is transported to spindle poles during mitosis and regulates Ran-dependent spindle
861   assembly factors in mammalian cells. J Cell Sci *117*, 6511-6522.

862   43.   Song, L., Craney, A., and Rape, M. (2014). Microtubule-dependent regulation of mitotic protein
863   degradation. Mol Cell *53*, 179-192.

864   44.   Sackton, K.L., Dimova, N., Zeng, X., Tian, W., Zhang, M., Sackton, T.B., Meaders, J., Pfaff, K.L.,
865   Sigoillot, F., Yu, H., et al. (2014). Synergistic blockade of mitotic exit by two chemical inhibitors of the
866   APC/C. Nature *514*, 646-649.

867   45.   Wei, J.H., Zhang, Z.C., Wynn, R.M., and Seemann, J. (2015). GM130 Regulates Golgi-Derived Spindle
868   Assembly by Activating TPX2 and Capturing Microtubules. Cell *162*, 287-299.

869   46.   Brownlee, C., and Heald, R. (2019). Importin alpha Partitioning to the Plasma Membrane Regulates
870   Intracellular Scaling. Cell *176*, 805-815 e808.

871   47.   Chinen, T., Yamamoto, S., Takeda, Y., Watanabe, K., Kuroki, K., Hashimoto, K., Takao, D., and
872   Kitagawa, D. (2020). NuMA assemblies organize microtubule asters to establish spindle bipolarity in
873   acentrosomal human cells. EMBO J *39*, e102378.

48. Lorson, M.A., Horvitz, H.R., and van den Heuvel, S. (2000). LIN-5 is a novel component of the spindle
apparatus required for chromosome segregation and cleavage plane specification in Caenorhabditis elegans.
J Cell Biol *148*, 73-86.
49. Greenberg, S.R., Tan, W., and Lee, W.L. (2018). Num1 versus NuMA: insights from two functionally
homologous proteins. Biophys Rev *10*, 1631-1636.
50. Weaver, L.N., Ems-McClung, S.C., Chen, S.H., Yang, G., Shaw, S.L., and Walczak, C.E. (2015). The Ran-
GTP gradient spatially regulates XCTK2 in the spindle. Curr Biol *25*, 1509-1514.
51. Sikirzhytski, V., Renda, F., Tikhonenko, I., Magidson, V., McEwen, B.F., and Khodjakov, A. (2018).
Microtubules assemble near most kinetochores during early prometaphase in human cells. J Cell Biol *217*,
2647-2659.
52. Booth, D.G., Hood, F.E., Prior, I.A., and Royle, S.J. (2011). A TACC3/ch-TOG/clathrin complex stabilises
kinetochore fibres by inter-microtubule bridging. EMBO J *30*, 906-919.
53. Nishimoto, T., Eilen, E., and Basilico, C. (1978). Premature of chromosome condensation in a ts DNA-
mutant of BHK cells. Cell *15*, 475-483.
54. Kiyomitsu, T., and Cheeseman, I.M. (2012). Chromosome- and spindle-pole-derived signals generate an
intrinsic code for spindle position and orientation. Nat Cell Biol *14*, 311-317.
55. Soderholm, J.F., Bird, S.L., Kalab, P., Sampathkumar, Y., Hasegawa, K., Uehara-Bingen, M., Weis, K.,
and Heald, R. (2011). Importazole, a small molecule inhibitor of the transport receptor importin-beta. ACS
Chem Biol *6*, 700-708.
56. Kiyomitsu, T., and Cheeseman, I.M. (2013). Cortical dynein and asymmetric membrane elongation
coordinately position the spindle in anaphase. Cell *154*, 391-402.
57. Kiyomitsu, T., Murakami, H., and Yanagida, M. (2011). Protein interaction domain mapping of human
kinetochore protein Blinkin reveals a consensus motif for binding of spindle assembly checkpoint proteins
Bub1 and BubR1. Mol Cell Biol *31*, 998-1011.
58. Goshima, G., Nedelec, F., and Vale, R.D. (2005). Mechanisms for focusing mitotic spindle poles by minus
end-directed motor proteins. J Cell Biol *171*, 229-240.
900

Figure 1

Figure 1. NuMA acts in spindle pole focusing usign its conserved microtubule-binding domain in human cells.

Figure 2. NuMA functions in spindle pole focusing independently of RCC1.

Figure 3. NuMA is liberated from importins at spindle poles independently of Ran-GTP.

Figure 4. RCC1 regulates chromosome-proximal localization of HURP and HSET.

Figure 5. HURP, but not importin-β, is requierd to stabilize k-fibers.

Figure 6. HURP dynamically accumulates on metaphase k-fibers in an importin-β dependent manner.

Figure 7. Models of local activation mechanisms for HURP and NuMA in mitosis.

Figure S1. Generation of cell lines that conditionally degrade endogenous NuMA and express NuMA mutants.

Figure S2. Generation of cell lines for auxin-inducible degradation of endogenous RCC1.

Figure S3. Generation of cell lines for auxin-inducible degradation of endogenous Ran-GAP1 and importin-β.

Figure S4. Generation of double knock-in cell lines that express RCC1-mAC and mCherry-fused TPX2, HSET, or HURP.

Figure S5. Generation of cell lines that degrade or visualize endogenous HURP.

A Control (Exponential recovery)

B +Dox+IAA (Straight line)

Figure S6. Fluorescent recovery kinetics of HURP in the presence or absence of importin-β