Forest loss as a catalyst of population and biodiversity change

Gergana N. Daskalova[1]*, Isla H. Myers-Smith[1], Anne D. Bjorkman[3], Sarah R. Supp[4], Shane A.

Blowes[5], Anne Magurran[2], Maria Dornelas[2]

[1]School of GeoSciences, University of Edinburgh, West Mains Road, Edinburgh EH9 3FF, Scotland

[2]Centre for Biological Diversity, University of St Andrews, Greenside Place, St Andrews KY16 9TF,

Scotland

[3]Senckenberg Biodiversity and Climate Research Centre, Frankfurt D-60325, Germany

[4]Data Analytics Program, Denison University, Granville, OH 43023, USA

[5]German Centre for Integrative Biodiversity Research (iDiv), Leipzig 04103, Germany

* Corresponding author

Gergana Daskalova

Crew Building, School of GeoSciences

University of Edinburgh

West Mains Road, Scotland

gndaskalova@gmail.com

Keywords: forest cover change, forest loss, populations, species richness, turnover, biodiversity

change, attribution

Main text word count: 1,843

Methods word count: 4,272

Figures: 4

Extended Data Figures: 10

Extended Data Tables: 1

Summary

Accelerating human impacts are reshaping Earth's ecosystems. Populations[1], richness[2–4] and composition[4] of communities at sites around the world are being altered over time in complex and heterogeneous ways[5–7]. Land-use change is thought to be the greatest driver of this population and biodiversity change in terrestrial ecosystems[8–10]. However, a major knowledge gap is whether land-use change drivers, such as forest loss and habitat conversion, can indeed explain the high heterogeneity of temporal population and biodiversity trends[9,11]. Here, we fill this gap by analysing change in 6,667 time series of populations (species' abundance)[12] and biodiversity (species richness and turnover in ecological communities)[13] over one and a half centuries of forest cover change and habitat transitions. We revealed an acceleration in both increases and decreases in population size, species richness and turnover after peak forest loss at over 2,000 sites across the globe. We found that temporal lags in population and biodiversity change following forest loss can extend up to half of a century and were longer for species with longer generation times such as large mammals. Together, our results demonstrate that historic and contemporary forest cover change do not universally lead to population declines and biodiversity loss, though population declines were most pronounced during and immediately following peak forest loss. By explicitly quantifying multi-decadal temporal lags in population and biodiversity responses to land-use change, our findings inform projections of how life on Earth will be reshaped across the Anthropocene.

Main text

Earth's biodiversity is changing[3–5,14]. At sites across the planet, populations are increasing and decreasing[1,15,16], species are lost[17] and gained[18,19], yet synthesis studies across sites indicate no net change in local scale species richness[3,4] despite marked shifts in species composition (turnover)[2,4,5]. At present, we have only a limited understanding of how global change drivers

produce these complex population and biodiversity patterns over time[6,20,21]. Our current

knowledge of the mechanisms explaining the ongoing reorganisation of ecological communities

predominantly stems from space-for-time[8,22] and modelling[23,24] approaches that attribute

population and richness declines to land-use change. Yet, space-for-time methods can

overestimate the effects of global change drivers compared to long-term monitoring, because

they do not account for ecological lags[6,25,26] and community self-regulation[27]. Such temporal

discrepancies in the magnitude of effects have been observed when studying the impacts of

warming on community change[25], and of habitat fragmentation on populations and

biodiversity[28,29]. The integration of multi-century long reconstructions of past land cover[30] and

high-resolution remote-sensing observations[31,32] with recent compilations of over five million

population and biodiversity records[12,13] provides an unprecedented opportunity to test the *in-*

situ impacts of land-use change. Only now can we quantitatively attribute the heterogeneous

patterns of population and biodiversity change observed over time to land-use dynamics, thus

improving projections of human impacts on the world's biota.

Here, we asked how populations (trends in numerical abundance) and biodiversity (trends in

species richness and community composition) across vertebrate, invertebrate and plant taxa

vary according to the timing and magnitude of forest cover change and habitat conversions

(Figure 1, Extended Data Figures 1 and 3b). We assessed whether population and biodiversity

change were different after versus before contemporary peak forest loss (the timing of the

largest forest loss event across the duration of each time series). Additionally, we tested

population and biodiversity change versus overall forest cover gain and loss experienced across

the duration of each time series. In a *post-hoc* analysis, we categorized population time series

based on whether they were recorded before, during, or after the period of all-time historic peak

forest loss (the timing of the largest forest loss event at the location of each time series between

the years 850 and 2015), and then compared population trends among the three categories.

Finally, we investigated among-taxa variation in temporal lags of population and biodiversity

responses to peak forest (the time period between contemporary peak forest loss and maximum

change in populations and communities). We used a hierarchical Bayesian modelling

framework for all attribution analyses, with individual time series nested within biomes[33] to

account for the spatial and temporal structure of the data. We used the Living Planet Database

(133,092 records) and the BioTIME database (4,970,128 records), currently the two largest

databases of population and community time series, respectively. We calculated population

change using state-space models that account for observation error and random fluctuations[34].

We quantified turnover by partitioning Jaccard's dissimilarity measure into its nestedness

(change due to communities becoming reduced subsets of themselves or new species

colonising in addition to the original species) and turnover[35] components. We focused on

turnover because it quantifies compositional changes due to species replacement and is

independent of changes in species richness. Our data synthesis quantitatively tests the

attribution of change in populations and ecological communities to land-use change through

time across the world's woody biomes.

We predicted greater population and species richness declines with increased forest loss.

Forest degradation and land-use conversion reduce habitat and resource availability[8,19,36] and

are the most common global threats for our studied species[37] (Figure 1b, Extended Data Figure

4e). Conversely, we predicted greater increases in population abundance and species richness

with larger gains in forest cover. Forest restoration and natural regeneration, two examples of

forest cover gain, can lead to positive biodiversity responses[38,39]. We expected greater turnover

of species within ecological communities with greater change in forest cover (both loss and

gain), as these extremes of the forest cover change spectrum both create novel environmental

conditions prompting local extinctions but also colonisations by new species[27,36]. Secondly, we

predicted that the largest population declines will occur during the periods of all-time peak forest

loss across sites as that is the baseline for maximum intensity of forest cover change[26]. Finally,

we predicted that temporal lags will be greater for species with longer generation times, as they

typically respond more slowly to environmental change[40] and have more limited dispersal[41]. If

we find support for our overall prediction that population and biodiversity loss will be greater

with higher forest loss, this would indicate that forest loss is a key driver of rapid and pervasive

declines over time. Alternatively, if we find support for heterogeneous and temporally-delayed

responses of populations and ecological communities to forest loss, this would imply that the

effects of land-use change over time are more complex than previously thought, which has

implications for improved prediction of future ecological change and development of biodiversity

and conservation policy.

a Distribution of study sites

Duration ● 20 ● 50 ● 80

LPD ●
BioTIME ●

b Timeline of monitoring and forest loss

Number of time series — 3300, 0
Forest cover change — 0, -0.05

1000   1250   1500   1750   2000

Proportion of species threatened by forest cover change and land-use conversion — 90%   82%

c Questions

MAGNITUDE	TYPE	TIMING
Do populations and biodiversity respond to contemporary peak forest loss?	Which type of habitat conversion yields the most population and biodiversity change?	Which taxa exhibit the longest temporal lags in their responses to contemporary peak forest loss?

d Predictions

Baseline fluctuations → Directional change

Contemporary peak forest loss

ΔVariable: +, 0, -

Time

↑ Population change Richness change Turnover

Primary forest   Secondary forest   Agricultural and urban areas

Lags in forest loss responses increase with longer generation times of species.

Peak forest loss ↔ Peak population / biodiversity change

Time

Figure 1. Forest cover change can influence population and biodiversity change through multiple mechanisms. For 38% of population time series and 70% of biodiversity time series monitoring began after historic peak forest loss had already occurred (see Extended Data Figures 2-3, 4a-b). See Extended Data Figure 1 for workflow of analyses. **a,** Global map of locations and duration of 542 Living Planet Database (LPD) and 190 BioTIME studies, containing the 6,497 time series from 2,154 locations analysed here. From the total number of time series analysed, 2,852 had experienced historic or contemporary forest cover gains and losses. Black outlines on **a** indicate locations that were forested at the start of the population or biodiversity monitoring period (1,247 sites). **b**, Timeline of monitoring and forest loss across the LPD and BioTIME databases (for variation in monitoring duration among time series, see Extended Data Figures 2-4). Forest loss was quantified over 10-year periods across 96 km^2 cells as a proportion, with e.g., -0.05 meaning a 5% loss of forest cover. Inset on **b** shows the proportion of species, part of the LPD and BioTIME databases, that are threatened by land-use change, based on species' IUCN threat assessments (see Extended Data Figure 4e). **c-d**, Questions and predictions for this study outlined with respect to population change, richness change and turnover (species composition change). Habitat conversion refers to a change in the dominant land cover type. Photos in **d** are by G. N. Daskalova.

We found that forest loss and habitat transitions did not universally lead to population declines and biodiversity loss (Figures 2-3) and are instead reshaping populations and ecological communities in more complex ways than previously recognised[8,15,22,24]. Surprisingly, forest loss acted as a catalyst for both positive and negative change and intensified population declines, population increases and species richness losses over time, despite equally long monitoring periods before and after peak forest loss. In 72% of the populations which were in decline before peak forest loss, the declines became more acute after forest loss (slope = -0.04 CI = -0.04 to

-0.03, Figure 2a). Similarly, 66% of the increasing populations experienced even more positive

population change after peak forest loss (slope = 0.02, CI = 0.02 to 0.03, Figure 2b). In contrast,

among time series, we did not find directional relationships among greater forest loss and

population and species richness declines (Figure 3, Extended Data Figures 5-7). This

disconnect between the magnitude of land-use change and population and biodiversity change

could be due to a number of factors including temporal lags in population or community

responses[26,40] and/or less forest cover change having occurred during the monitoring period

relative to historic forest clearing[26,42,43] (Extended Data Figures 2-3). In further contrast to our

first prediction, larger magnitudes of forest loss often led to greater increases, rather than

decreases, in species richness over time, particularly among time series comparisons with

shorter durations (Extended Data Figure 6e). Forest loss is a key driver of habitat fragmentation

which can lead to rapid colonisation by new species due to increased landscape heterogeneity

and larger breadth of ecological niches across sites[5,28,36]. Our results highlight that the same

global change driver, forest loss, affects populations and ecological communities in

heterogeneous ways at different sites around the world, and accounting for this heterogeneity

is key when scaling from local impacts of human activities to global scale biodiversity patterns

and attribution of change[10].

Figure 2. At the site level, population and biodiversity change increase after contemporary peak forest loss. Though species richness tended to decrease after peak forest loss, the proportion of populations that experienced increases and declines in population abundance and turnover were similar. Numbers on plots indicate number of time series. Population, richness and turnover change increased across 55% of the 1,007 time series for

which baseline comparisons were possible. Distributions show posterior means from model fits

of **a**, population declines (μ), **b**, population increases (μ), **d**, richness losses (slopes), **e**, richness

gains (slopes) and **g**, turnover (in the final year of each period relative to the first year, measured

as Jaccard's dissimilarity, where zero indicates no changes in species composition and one

indicates a completely new set of species). Vertical lines indicate the mean for each category.

The y-axis on **a**, **b** and **c** refers to the probability density function for the kernel density estimation

per unit on the x-axis, and the distributions are relative to one another. Temporal change before

and after peak forest loss, **c**, **f**, **h** is indicated with lines for individual time series. The time

window duration varied among time series but was consistent at the time series level (*n* years

before disturbance = *n* years after disturbance at the time series level, n ≥ 5 years). Light grey

and dark grey points indicate the mean values across all times series, with error bars indicating

the 2.5 and 97.5% quantiles. See Extended Data Table 1 for model outputs.

Our results revealed that population declines were most pronounced during the period of all-

time peak forest loss and turnover was highest when primary forests were converted to

agricultural and urban areas (Figure 3). Following peak contemporary forest loss, turnover

increased by over 10% in 19% of the time series (Figure 2), further testifying to the high rates

of compositional change detected across the Anthropocene[4,5,14]. However, within 22% of the

time series, turnover declined by over 10% after forest loss, suggesting that biotic

homogenisation might also be occurring following human-induced environmental change[44].

Taken together, our findings suggest site-specific impacts that were stronger and more common

when the population and biodiversity monitoring captured the largest forest loss events and the

most dramatic habitat conversion events across time relative to when monitoring is mismatched

with forest cover change (Figures 2-3). A greater proportion of the planet is projected to

experience an unprecedented amount of land-use change in the coming decades[45], highlighting

the importance of improved biodiversity monitoring in current and future hotspots of forest loss

and habitat conversion.

Figure 3. Population declines were most acute when the all-time peak in deforestation for study sites occurred within the duration of population monitoring. a, Population declines are more likely to be detected when peak deforestation occurs during the population monitoring period (slope = -0.01, CI = -0.01 to -0.01). Density plots show distributions of μ values

(population trend). **b,** Among time series, there were no directional relationships among overall

forest loss and population declines and increases, species richness declines and turnover.

Standardised effect sizes in **b** were calculated by dividing the slopes from hierarchical models

testing population and biodiversity change as a function of forest loss by the standard deviation

of forest loss. For visualisations of each model, see Extended Data Figures 5-6. Low sample

size precluded an analysis of biodiversity change differences based on whether all-time peak

forest loss had occurred before, during or after the biodiversity monitoring. Distributions in **c**

show μ values for population change, posterior means (slopes) for richness change and

Jaccard's dissimilarity for turnover under different habitat conversions. Line thickness

corresponds with magnitude of detected change. Forest loss led to habitat conversion in 304 of

5795 (approximately 5%) of monitored population and biodiversity time series. There was only

one instance of a population time series experiencing habitat conversion with secondary forest

as the starting dominant land cover, thus no distribution is plotted for that category. The y-axis

in **c** refers to the probability density function for the kernel density estimation per unit on the x-

axis, and the distributions are relative to one another. Numbers in plots indicate number of time

series for each category. Small sample sizes of an average 10 time series per transition types

of interest for this analysis precluded statistical analysis and inferences on the effects of habitat

transitions were drawn by visually inspecting the density distributions. See Extended Data

Figures 5-6 and Extended Data Table 1 for models of forest cover gain and loss (GFC

database[12], 2000-2016) and forest cover loss (LUH database[11], across the time series) and

population and biodiversity change. See Extended Data Figure 7 for distributions of population

and biodiversity change following habitat transitions detected by the MODIS Land Cover

Database[32].

We found evidence for up to half-century ecological lags in changes in population abundance,

species richness and community composition following forest loss (Figure 4). On average, we

documented maximum change in populations and biodiversity six to 13 years after

contemporary peak forest loss across taxa. Yet, nearly half of population and biodiversity

change (40%) occurred within three years of peak forest loss, demonstrating that rapid shifts in

populations and ecological communities occur frequently (Figure 4a). As predicted, the period

between peak forest loss and peak change in populations and biodiversity was longer for taxa

with longer generation times (e.g., large mammals, Figure 4b, Extended Data Table 1), further

confirmed by a *post-hoc* analysis of lags in population change versus mammal generation time

(Extended Data Figure 8a). Population declines and increases occurred on similar time scales

(Extended Data Figure 8b-c), potentially explaining why previous temporal analyses of

population change have not found evidence for net population declines[1,20]. Losses in species

richness lagged behind richness gains only by approximately half a year (slope = 0.5, CI = 0.1

– 1.05), indicating that potential extinction debts and immigration credits accumulated at roughly

the same speed across taxa. The similar pace and temporal delay of richness gains and losses

could be the source of the previously observed findings of no net local richness change[3,4], yet

substantial compositional change[2,4] across sites and taxa. Such temporal lags in biodiversity

change have also been observed in post-agricultural forests[2,46] and fragmented grasslands[40],

where agricultural activity has ceased decades to centuries ago, yet richness and community

composition change continue to the modern-day. Overall, our results indicate that increasing

rates of land-use change in the Anthropocene[11,45] will alter ecosystems on both short- and long-

term timescales.

Figure 4. Population and community change after peak forest loss may be delayed by up to half a century across 3,187 time series from around the world, with species with long generation times and low mobility showing the largest temporal lags. a, Distributions of

temporal lag values (measured in years). **b,** Vertebrate population change lags by taxa (reptiles,

amphibians, birds and mammals). **c,** Richness change and **d**, turnover lags by taxa

(invertebrates, amphibians, birds, mammals, plants and trees). Peak forest loss refers to the

timing of the largest forest loss event across the duration of each time series. Lag was measured

as the number of years between peak forest loss and peak population/biodiversity change.

Turnover was measured as Jaccard's dissimilarity, where zero indicates to changes in species

composition and one indicates a completely new set of species. Numbers on plots indicate

number of time series. Some time series did not experience richness change following peak

forest loss, thus no lags were calculated for them. See Extended Data Table 1 for model outputs.

In summary, our analysis reveals an acceleration of increases and decreases of populations

and biodiversity after forest loss and habitat conversion at sites around the planet. Our findings

that all is not loss contrast with our hypothesis and challenge the widely-held assumption that

land-use change universally leads to population declines and species richness loss[8,15,23].

Nevertheless, the increased magnitude and likelihood of population declines during and

following peak forest loss highlight that human impacts are altering the biodiversity of the planet

and emphasize the importance of expanded biodiversity monitoring in current and future

hotspots of land-use change. A critical assumption underlying existing projections of biodiversity

responses to land-use change[8,23] is that space-for-time approaches accurately reflect longer-

term population and biodiversity dynamics[11]. On the contrary, we find that temporal lags in

population and biodiversity change following forest loss varied by taxa and generation time and

extended up to half of a century. Over the Anthropocene, ecosystems could be responding to a

suite of global change drivers, in addition to land-use change, and a key next research step is

to test the synergy and discord between the effects of multiple anthropogenic threats on Earth's

biota. Our results highlight the complex biological responses to habitat conversion across sites,

taxa and time scales that are leading to the reorganisation of ecological communities. Thus,

indicators used to assess biodiversity change regionally and globally[10], including progress

towards Aichi targets[47], must capture the full spectrum and temporal spread of population and

biodiversity responses to human impacts across the Anthropocene.

References

1. Daskalova, G. N., Myers-Smith, I. H. & Godlee, J. L. Rarity and conservation status do not predict vertebrate population trends. *bioRxiv* (2018). doi:https://doi.org/10.1101/272898

2. Baeten, L., Hermy, M., Van Daele, S. & Verheyen, K. Unexpected understorey community development after 30 years in ancient and post-agricultural forests: Land use and 30-year forest development. *J. Ecol.* **98**, 1447–1453 (2010).

3. Vellend, M. *et al.* Global meta-analysis reveals no net change in local-scale plant biodiversity over time. *Proc. Natl. Acad. Sci.* **110**, 19456–19459 (2013).

4. Dornelas, M. *et al.* Assemblage Time Series Reveal Biodiversity Change but Not Systematic Loss. *Science* **344**, 296–299 (2014).

5. Magurran, A. E. *et al.* Divergent biodiversity change within ecosystems. *Proc. Natl. Acad. Sci.* **115**, 1843–1847 (2018).

6. Yoccoz, N. G., Ellingsen, K. E. & Tveraa, T. Biodiversity may wax or wane depending on metrics or taxa. *Proc. Natl. Acad. Sci.* **115**, 1681–1683 (2018).

7. Hillebrand, H. *et al.* Biodiversity change is uncoupled from species richness trends: Consequences for conservation and monitoring. *J. Appl. Ecol.* **55**, 169–184 (2018).

8. Newbold, T. *et al.* Global effects of land use on local terrestrial biodiversity. *Nature* **520**, 45–50 (2015).

9. Maxwell, S. L., Fuller, R. A., Brooks, T. M. & Watson, J. E. Biodiversity: The ravages of guns, nets and bulldozers. *Nature* **536**, 143–145 (2016).

10. IPBES. Summary for policymakers of the regional assessment report on biodiversity and ecosystem services for Europe and Central Asia of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. M. Fischer, M. Rounsevell, A. Torre-Marin Rando, A. Mader, A. Church, M. Elbakidze, V. Elias, T. Hahn. P.A. Harrison, J.

Hauck, B. Martín-López, I. Ring, C. Sandström, I. Sousa Pinto, P. Visconti and N.E

Zimmermann (eds.). IPBES secretariat, Bonn, Germany. (2018).

11. De Palma, A. *et al.* Challenges With Inferring How Land-Use Affects Terrestrial

Biodiversity: Study Design, Time, Space and Synthesis. in *Advances in Ecological*

Research **58**, 163–199 (Elsevier, 2018).

12. LPI. Living Planet Index database. (2016).

13. Dornelas, M. *et al.* BioTIME: A database of biodiversity time series for the Anthropocene.

Glob. Ecol. Biogeogr. **27**, 760–786 (2018).

14. Blowes, S., Supp S. R. *et al.* Biodiversity trends are stronger in marine than terrestrial

assemblages. (2018). doi:10.1101/457424

15. Ceballos, G., Ehrlich, P. R. & Dirzo, R. Biological annihilation via the ongoing sixth mass

extinction signaled by vertebrate population losses and declines. *Proc. Natl. Acad. Sci.*

201704949 (2017). doi:10.1073/pnas.1704949114

16. Bowler, D. E., Heldbjerg, H., Fox, A. D., O'Hara, R. B. & Böhning-Gaese, K. Disentangling

the effects of multiple environmental drivers on population changes within communities. *J.*

Anim. Ecol. **87**, 1034–1045 (2018).

17. Biesmeijer, J. C. Parallel Declines in Pollinators and Insect-Pollinated Plants in Britain and

the Netherlands. *Science* **313**, 351–354 (2006).

18. Steinbauer, M. J. *et al.* Accelerated increase in plant species richness on mountain

summits is linked to warming. *Nature* **556**, 231–234 (2018).

19. Elahi, R. *et al.* Recent Trends in Local-Scale Marine Biodiversity Reflect Community

Structure and Human Impacts. *Curr. Biol.* **25**, 1938–1943 (2015).

20. Leung, B., Greenberg, D. A. & Green, D. M. Trends in mean growth and stability in

temperate vertebrate populations. *Divers. Distrib.* **23**, 1372–1380 (2017).

21. Bowler, D. *et al.* The geography of the Anthropocene differs between the land and the

sea. (2018). doi:10.1101/432880

22. Betts, M. G. *et al.* Global forest loss disproportionately erodes biodiversity in intact

landscapes. *Nature* **547**, 441–444 (2017).

23. Newbold, T. Future effects of climate and land-use change on terrestrial vertebrate

community diversity under different scenarios. *Proc. R. Soc. B Biol. Sci.* **285**, 20180792

(2018).

24. Newbold, T., Tittensor, D. P., Harfoot, M. B. J., Scharlemann, J. P. W. & Purves, D. W.

Non-linear changes in modelled terrestrial ecosystems subjected to perturbations. (2018).

doi:10.1101/439059

25. Elmendorf, S. C. *et al.* Experiment, monitoring, and gradient methods used to infer climate

change effects on plant communities yield consistent patterns. *Proc. Natl. Acad. Sci.* **112**,

448–452 (2015).

26. Mihoub, J.-B. *et al.* Setting temporal baselines for biodiversity: the limits of available

monitoring data for capturing the full impact of anthropogenic pressures. *Sci. Rep.* **7**,

41591 (2017).

27. Gotelli, N. J. *et al.* Community-level regulation of temporal trends in biodiversity. *Sci. Adv.*

3, e1700315 (2017).

28. Fahrig, L. Ecological Responses to Habitat Fragmentation Per Se. *Annu. Rev. Ecol. Evol.*

Syst. **48**, 1–23 (2017).

29. Haddad, N. M. *et al.* Experimental evidence does not support the Habitat Amount

Hypothesis. *Ecography* **40**, 48–55 (2017).

30. Hurtt, G. C. *et al.* Harmonization of land-use scenarios for the period 1500–2100: 600

years of global gridded annual land-use transitions, wood harvest, and resulting

secondary lands. *Clim. Change* **109**, 117–161 (2011).

31. Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change.

Science **342**, 850–853 (2013).

32. Channan, S., Collins, K. & Emanuel, W. R. Global mosaics of the standard MODIS land

cover type data. University of Maryland and the Pacific Northwest National Laboratory,

College Park, Maryland, USA. (2014).

33. Olson, D. M. & Dinerstein, E. The Global 200: Priority Ecoregions for Global Conservation.

Ann. Mo. Bot. Gard. **89**, 199 (2002).

34. Humbert, J.-Y., Scott Mills, L., Horne, J. S. & Dennis, B. A better way to estimate

population trends. *Oikos* **118**, 1940–1946 (2009).

35. Baselga, A. Partitioning the turnover and nestedness components of beta diversity:

Partitioning beta diversity. *Glob. Ecol. Biogeogr.* **19**, 134–143 (2010).

36. Sax, D. F. & Gaines, S. D. Species diversity: from global decreases to local increases.

Trends Ecol. Evol. **18**, 561–566 (2003).

37. IUCN. The IUCN Red List of Threatened Species. Version 2017-3. (2017).

38. Benayas, J. M. R., Newton, A. C., Diaz, A. & Bullock, J. M. Enhancement of Biodiversity

and Ecosystem Services by Ecological Restoration: A Meta-Analysis. *Science* **325**, 1121–

1124 (2009).

39. Chazdon, R. L. Beyond Deforestation: Restoring Forests and Ecosystem Services on

Degraded Lands. *Science* **320**, 1458–1460 (2008).

40. Krauss, J. *et al.* Habitat fragmentation causes immediate and time-delayed biodiversity

loss at different trophic levels: Immediate and time-delayed biodiversity loss. *Ecol. Lett.*

13, 597–605 (2010).

41. Bommarco, R. *et al.* Dispersal capacity and diet breadth modify the response of wild bees

to habitat loss. *Proc. R. Soc. B Biol. Sci.* **277**, 2075–2082 (2010).

42. Spooner, F. E. B., Pearson, R. G. & Freeman, R. Rapid warming is associated with

population decline among terrestrial birds and mammals globally. *Glob. Change Biol.* **24**,

4521–4531 (2018).

43. Kaplan, J. O., Krumhardt, K. M. & Zimmermann, N. The prehistoric and preindustrial

deforestation of Europe. *Quat. Sci. Rev.* **28**, 3016–3034 (2009).

44. Olden, J. D., LeRoy Poff, N., Douglas, M. R., Douglas, M. E. & Fausch, K. D. Ecological

and evolutionary consequences of biotic homogenization. *Trends Ecol. Evol.* **19**, 18–24

(2004).

45. Egli, L., Meyer, C., Scherber, C., Kreft, H. & Tscharntke, T. Winners and losers of national

and global efforts to reconcile agricultural intensification and biodiversity conservation.

Glob. Change Biol. (2018). doi:10.1111/gcb.14076

46. Vellend, M. *et al.* Extinction debt of forest plants persists for more than a century following

habitat fragmentation. *Ecology* **87**, 542–548 (2006).

47. CBD. Strategic plan for biodiversity 2011–2020 and the Aichi targets. In Report of the

Tenth Meeting of the Conference of the Parties to the Convention on Biological Diversity.

(2010).

Acknowledgements

We thank the WWF and ZSL for compiling the Living Planet Database, the BioTime team for

compiling the BioTime database (which was supported by ERC AdG BioTIME 250189 and

ERC PoC BioCHANGE 727440), the creators of the Land Use Harmonization Database, The

Hansen Lab for producing the Forest Cover Change Database and NASA for producing the

MODIS Landcover Database. We thank the Forest & Nature Lab at Ghent University for a

stimulating discussion on historic and contemporary land-use change and choosing appropriate

baselines for comparison of biodiversity change through time. We are grateful to Albert

Phillimore and Kyle Dexter for providing advice during the conceptualization of the study, and

to Laura Antão and Mark Vellend for providing feedback on the draft manuscript. We thank the

German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig and the sChange

working group for supporting the initial data synthesis work that has led to this study. G.N.D.

was funded by a Carnegie-Caledonian PhD Scholarship and supported by a NERC doctoral

training partnership grant (NE/L002558/1).

Author contributions

G.N.D., M.A.D. and I.M.S. conceptualised the study. G.N.D. integrated databases and

conducted statistical analyses with input from S.B., I.M.S., A.D.B. and M.A.D. G.N.D. created

the figures with input from co-authors. S.B. and S.S. wrote the code for the rarefaction of the

BioTIME studies. I.M.S. was the primary supervisor and M.A.D. the co-supervisor for G.N.D.

M.A.D. and A.M. funded the compilation of the BioTime database. G.N.D. wrote the first draft

and all authors contributed to revisions.

Methods

For an illustration of the workflow of our analyses of forest cover and population and biodiversity

change through time, see Extended Data Figures 1 and 3b. All data and statistical analyses are

described in detail below. We did not predetermine sample size and instead worked with all

available temporal population, biodiversity and forest cover change data that met our duration

criteria. For analyses of population change, we included time series with five or more survey

points. For analyses of biodiversity change, we included time series with five or more data points

when analysing the full time series, and time series with two or more data points when matching

the duration of time series comparisons to the 16-year duration of the Global Forest Change

Database from 2000 to 2016).

Databases

Forest cover change data

To quantify historic and contemporary forest cover change, we extracted historic forest loss

from the Land Use Harmonisation (LUH; 850 – 2015, forest loss and habitat transitions at a

0.25° degree resolution)[30] and contemporary forest cover change and habitat conversions from

the Global Forest Change (GFC, 2000 – 2016, forest loss and gain at a 30 m resolution)[31], and

MODIS Landcover (2000 – 2013, land-use transitions at 500m resolution)[32] datasets.

Land Use Harmonisation Database

To estimate forest cover change across a time period matching the full duration of the

biodiversity observations, we derived the change in primary forest cover from the Land Use

Harmonisation database (LUH)[30] for 96 km^2 cells around the location of each population in the

LPD database and for the standardised grid cells of the BioTIME database (~ 96 km^2 each).

LUH includes annual gridded fractions of land-use states for the period from 850 to 2013 at

0.25° x 0.25° resolution. The estimates are based on historical reconstructions using Earth

System models, with inputs such as regional and national rates of wood harvest and potential

biomass density. The accuracy and precision of LUH increases towards the modern day, when

there are more available data to inform the Earth System models. Note that unlike GFC, LUH

estimates forest cover as a proportion (bounded between zero and one). For our analyses, we

focused on time series from locations that have experienced at least 0.05 (equivalent to 5%)

forest loss. To calculate total forest cover change over the period of a given population or

biodiversity time series, we subtracted the proportion of forest cover in the first year of

biodiversity monitoring from the proportion of forest cover in the last year. The type of forest

cover change detected by the LUH database was predominantly forest loss, with forest gain

occurring infrequently and at very small magnitudes (<0.001 out of maximum 1), thus we focus

our analysis on forest loss.

To estimate the historic baseline of forest cover change, we calculated yearly change in % forest

cover in a study cell from one year to the next for each site from 850 to 2015 from the LUH data,

and determined the 10-year period when the most forest loss occurred (historic peak forest loss,

calculated by adding the yearly proportions of forest loss in each cell over standardised 10-year

blocks). Time since historic peak forest loss was a poor predictor of the variation in

contemporary population and biodiversity change (Extended Data Figure 10e-f). To determine

contemporary peak forest loss for each time series of monitoring data, we calculated yearly

changes in forest cover across the duration of each time series and determined the year when

the most change had occurred.

Global Forest Change Database

We derived overall forest loss and forest gain across the 2000-2016 period for 96 km^2 cells

around the location of each population in the LPD database and for the standardised grid cells

of the BioTIME database (~ 96 km^2 each) from the Global Forest Change (GFC)[31] database

using the Google Earth Engine[48]. The GFC database provides high resolution forest cover

change data, derived from Landsat satellite observations at a 30-meter spatial resolution. We

calculated the total area of forest cover gain and loss separately (measured in km^2) for each 96

km^2 cell on a yearly time step. We then summed the yearly values for the period that coincided

with population and biodiversity monitoring to estimate overall forest cover gain and loss (two

separate metrics). For example, for a biodiversity time series spanning 2002 – 2009, our forest

cover gain and loss metrics included the total amount of forest cover gained and lost during that

same period. For our analyses, we focused on time series from locations that have experienced

at least 0.5 km^2 of forest gain or loss. GFC does not distinguish between primary forest,

secondary forest and plantations, but it does provide a very high-resolution measure of general

forest cover. The drivers of the forest loss detected by GFC across our study sites are

predominantly forestry, changes in agricultural practices and wildfires[49]. Note that the GFC

database spans from 2000 to 2016, whereas the earliest terrestrial biodiversity record in

BioTIME is from 1858.

MODIS Landcover Database

We used the MODIS Landcover Database[32] to quantify habitat conversion for locations where

we had population and biodiversity monitoring data. The MODIS Database has a resolution of

500 m, and it uses satellite-derived reflectance data to classify land cover around the world. To

determine the types of habitat conversion between 2000 and 2013 (the time span of available

MODIS data) across all monitoring locations, we calculated the dominant land cover type at the

start and end of each population and biodiversity time series and split time series into categories

such as "no habitat conversion" and "grassland to woody savannah". We focused on the eight

most frequent types of habitat conversion (Extended Data Figure 7).

By synthesising information from scenario data based on Earth Dynamics Models (LUH) and

remote-sensing databases (GFC, MODIS), we were able to determine historic forest loss from

the start of the monitoring period to 2015, as well as contemporary forest cover change (gain

and loss) and habitat transitions from 2000 to 2016. GFC and MODIS detect forest cover, with

no distinction between primary and secondary forests, thus we derived information on

transitions from primary to secondary forest from the LUH database. We calculated overall

forest cover change because we considered total change in habitat to be more meaningful for

long-term population and biodiversity trends as opposed to an annual rate of forest cover

change which does not capture cumulative effects. Together, the three databases (GFC,

MODIS, LUH) encompass two different elements of land-use change: 1) land cover types and

long-term historical reconstructions of past land-use and habitat conversions and 2) high-

resolution satellite data from recent years of forest cover change and habitat conversion types.

Thus, the combined analysis allows for a comprehensive test of the effects of land-use change

on populations and biodiversity around the world.

Population time series data (Living Planet Database)

We analysed 4,228 population time series, with records distributed around the world.

Geographic representation is variable with, for example, an under-representation of tropical

regions in the population data (Figure 1). In the LPD, some populations have precise

coordinates, whereas the location of others are approximate. Because of the extent over which

we are calculating forest cover change (96 km^2), we included both types of populations in our

analysis. Duration varied across time series (Extended Data Figure 4c-d) and we only included

populations with at least five survey points. The overall range of the time series covered the

period between the years 1970 and 2014. We calculated population change using state-space

models which are particularly appropriate when quantifying change in data with varying

collection methodology, as they take into account observation error and process noise[50,51]. For

more details on state-space model calculations, see Humbert *et al.* 2009[34] and Daskalova *et al.*

2018[1]. We scaled the population size data to be between 0 and 1 to analyse within-population

relationships and to make sure that we were not conflating within-population relationships and

between-population relationships[52]. State-space models partition the variance in abundance

estimates into process error (σ^2) and observation or measurement error (τ^2) and estimate

population trends (μ):

$$X_t = X_{t-1} + \mu + \varepsilon_t, (1)$$

where X_t and X_{t-1} are the scaled (observed) abundance estimates (between 0 and 1) in the

present and past year, with process noise represented by $\varepsilon_t \sim gaussian(0, \sigma^2)$. We included

measurement error following:

$$Y_t = X_t + F_t,$$

where Y_t is the estimate of the true (unobserved) population abundance with measurement

error:

$$F_t \sim gaussian(0, \tau^2).$$

We substituted the estimate of population abundance (Y_t) into equation 1:

$$Y_t = X_{t-1} + \mu + \varepsilon_t + F_t.$$

Given $X_{t-1} = Y_{t-1} - F_{t-1}$, then:

$$Y_t = Y_{t-1} + \mu + \varepsilon_t + F_t - F_{t-1}.$$

For each time series, we calculated overall population change (μ) experienced 1) across the

periods before and after contemporary peak forest loss, 2) across the full duration of the time

series, 3) from 2000 to 2016 (matching the temporal scale of the GFC database), and 4) from

2000 to 2013 (matching the temporal scale of the MODIS database). We standardised the

number of years over which we calculated population change before and after peak forest loss

on the population-level, meaning that the number of years before and after was the same within

populations, but might differ among populations.

Biodiversity time series data (BioTIME Database)

We analysed 2,339 time series from 190 studies from terrestrial biomes across the globe, part

of the BioTIME database[13] (with the addition of 36 studies that are not yet a part of the public

database). Similarly to the LPD, tropical regions and some taxa such as amphibians and reptiles

were under-represented. Some of the study locations fall within protected areas (32%). Because

those studies only had one time series each, overall only 1% of analysed time series were from

inside protected areas. To account for the different spatial extents of the BioTIME database and

uneven sampling, studies with multiple locations and extents > 72 km^2 were partitioned into 96

km^2 grids, and then sample-based rarefaction was applied to standardise sampling within each

time series[14]. Duration varied across time series (Extended Data Figure 4c-d) and the overall

range of the time series covered the period between the years 1858 and 2016. For time series

with five or more years of monitoring records, we calculated overall richness change and

turnover experienced 1) across the periods before and after contemporary peak forest loss, 2)

across the full duration of the time series. For time series with two or more years of monitoring

records, we calculated overall richness change and turnover experienced 3) from 2000 to 2016

(matching the temporal scale of the GFC database), and 4) from 2000 to 2013 (matching the

temporal scale of the MODIS database). The GFC and MODIS databases cover shorter time

periods, thus we included biodiversity time series with shorter durations than the five-year cut

off point that was used in the rest of our analyses using datasets with longer durations (but note

that 76% of biodiversity time series had a duration of three or more years). To estimate richness

change, we modelled species richness versus time (year, mean centered) with random slopes

and intercepts for each rarefied cell and a Poisson error distribution with a log link.

564
$$log(\mu_{j,i,t}) = \beta_0 + \beta_{0j} + \beta_{0j,i} + (\beta_1 + \beta_{1j} + \beta_{1j,i})year_{j,i,t},$$

565
$$y_{j,i,t} \sim poisson(\mu_{j,i,t}),$$

where $year_{j,i,t}$ is the time in years, β_0 and β_1 are the global intercept and slope (fixed effects),

β_{0j} and β_{1j} are the biome-level departures from β_0 and β_1 (respectively; biome-level random

effects), $\beta_{0j,i}$ and $\beta_{1j,i}$ are the (nested) cell-level departures from β_0 and β_1 (cell-level random

effects); $y_{j,i,t}$ is the (rarefied) species richness within the jth biome in the ith cell in year t.

From the richness over time model, we extracted the posterior means for richness change for

each time series (i.e., the cell-level slope estimates), which then became the response variable

in the second stage of our analyses where we tested richness change versus forest cover

change (see Statistical analyses section).

To determine changes in community composition, we calculated the turnover component of

beta diversity (changes due to species replacement rather than changes in species

abundances[14,35]), at the end of each time period outlined above relative to the first year of

observation in the same period. Turnover is bound between zero and one, where zero is no

change in species composition and one indicates that all of the original species of a community

have been replaced with new species.

Statistical analyses

When testing for an attribution signal (i.e., evidence that a predictor variable is a potential driver

of population or biodiversity change), we always matched the temporal scales of the forest cover

change data and the population and biodiversity data. For example, when testing the effects of

forest cover change and land-use transitions as detected by GFC (2000 – 2016) and MODIS

(2000 – 2013), we calculated population and biodiversity change for the same time periods.

Because of the longer duration of the LUH database, we were also able to extract forest and

land cover information for the full duration of the LPD and BioTIME time series. For our analyses

of contemporary peak forest loss and overall forest loss (using the LUH database over a time

period matching the duration of each time series), we excluded locations which had less than

0.05 (out of maximum 1) forest cover change. We excluded locations which had no forest cover

across the duration of the time series in both the 96 km^2 cells and the 500 km^2 larger landscape

cells from our analyses of population and biodiversity change versus forest cover gain and loss

from 2000 to 2016 (using the GFC database). See Extended Data Table 1 for the outputs of all

statistical models and their respective sample sizes.

Population and biodiversity change after versus before contemporary peak forest loss

To test if temporal population and biodiversity change differed before and after peak forest loss

on the site-level, we split each time series into two periods – before and after peak deforestation

– and estimated population change, richness change and turnover for each period separately.

Then, to infer if population and biodiversity change differed following peak forest loss, we

modelled μ (population change), richness change (cell-level random slopes) and turnover as a

function of period (categorical with two levels – before or after forest loss) and time series

duration (numeric) as fixed effects, with a biome random effect to account for the spatial

clustering of the data. For population and richness change, we modelled the positive and

negative components of the distributions of change separately, e.g., one model for populations

with positive μ values and one model for populations with negative μ values. This approach

allowed us to test if the effects of forest loss differ across the positive and negative dimensions

of population and biodiversity change. The models were as follows:

612
$$\mu_{j,i,p} = \beta_0 + \beta_{0j} + \beta_1 * duration_{j,i,p} + \beta_2 * period_{j,i,p},$$

613
$$y_{j,i,p} \sim gaussian(\mu_{j,i,p}, \sigma^2),$$

where $duration_{j,i,p}$ is the duration of the time series in years of cell i within biome j for period p,

and $period_{j,i,p}$ is an indicator variable for the period (before or after forest loss); β_0, β_1 and β_2

are the global intercept and slope estimates for duration and the categorical period effect,

respectively (fixed effects), β_{0j} is the biome-level departures from β_0 (biome-level random

effects); $y_{j,i,p}$ is the estimate for change in population size or species richness for the *ith* cell in

the *jth* biome for the *pth* period.

To model the change in turnover before and after contemporary peak forest loss, we followed

the same conceptual framework as outlined above, but we used a zero one inflated beta

distribution to account for the properties of turnover (bounded between zero and one, inclusive,

where one is a complete change in species composition). The probability density function for

the zero one inflated beta distribution is:

$$betainf(y; \alpha, \gamma, \mu, \phi) = \begin{cases} \alpha(1 - \gamma), & y = 0 \\ \alpha\gamma, & y = 1 \\ (1 - \alpha)\gamma f(y; \mu, \phi), & 0 < y < 1, \end{cases}$$

where α is the probability that a zero or one occurs, γ is the probability that a one occurs (given

an observation is a zero or a one), and μ and ϕ are the mean and precision of the beta

distribution, respectively. In the parameterisation approach we used[53] ϕ is inversely related to

the variance. Beta parameterisation is also sometimes expressed through the parameters p and

q that can be derived from our framework following $\phi = p + q$[54]. Because only 7% of time series

did not experience any change in species composition (y = 0) in the time period after

contemporary forest loss, and less than 1% of time series had a completely new set of species

(y = 1) occupying the ecological communities, for y = 0 and y = 1, α and γ were modelled

assuming a Bernoulli distribution and logit-link function, and models were fit with only an

intercept. For $0 < y < 1$, we assumed a beta error distribution and a logit-link function:

$$logit(\mu_{j,i,p}) = \beta_0 + \beta_{0j} + \beta_1 * duration_{j,i,p} + \beta_2 * period_{j,i,p},$$

$$y_{j,i,p} \sim Beta(\mu_{j,i,p}, \phi),$$

where $duration_{j,i,p}$ is the duration of the time series in years of cell *i* within biome *j* for period *p*,

and $period_{j,i,p}$ is an indicator variable for the period (before or after forest loss); β_0, β_1 and β_2

are the global intercept and slope estimates for duration and the categorical period variable

respectively (fixed effects), and β_{0j} are the biome-level departures from β_0 (biome-level random

intercepts); $y_{j,i,p}$ is the estimate of turnover for the *ith* cell in the *jth* biome for the *pth* period.

Population change before, after and during the period of all-time historic peak forest loss

To determine if population change differed based on whether population time series were

recorded before, during, or after the period of all-time historic peak forest loss (the timing of the

largest forest loss event at the location of each time series between the years 850 and 2015),

we modelled μ (population change) as a function of when monitoring started (categorical with

three levels – before, during or after peak forest loss) and time series duration (numeric) as

fixed effects, with a biome random effect to account for the spatial clustering of the data. Low

sample size precluded a similar analysis for biodiversity change (Extended Data Figure 3). The

model was as follows:

654
$$\mu_{j,i,m} = \beta_0 + \beta_{0j} + \beta_1 * duration_{j,i,m} + \beta_2 * monitoring\ start_{j,i,m},$$

655
$$y_{j,i,m} \sim gaussian(\mu_{j,i,m}, \sigma^2),$$

where $duration_{j,i,m}$ is the duration of the time series in years of cell *i* within biome *j* for monitoring

start *m,* and $monitoring\ start_{j,i,m}$ is an indicator variable denoting when monitioring started; β_0,

β_1 and β_2 are the global intercept and slope estimates for duration and the categorical

monitoring start variable respectively (fixed effects), β_{0j} is the biome-level departures from β_0

(respectively; biome-level random effects); $y_{j,i,m}$ is the estimate for change in population size or

species richness for the *ith* cell in the *jth* biome for the *mth* monitoring start.

Habitat conversion and population and biodiversity change

To determine the influence of the type of forest cover change (i.e., land-use transitions) on

population and biodiversity change, we compared the distributions of population and biodiversity

change across transitions types (from primary forest to secondary forest, from primary forest to

non-natural habitat, and from secondary forest to non-natural habitat, to which we refer as

habitat conversion). Small sample sizes (on average 10 time series per transition type)

precluded statistical analysis, thus we report findings from a visual inspection of distributions of

population and biodiversity change across habitat conversion types.

To test the effect of forest cover change on population and biodiversity change among sites, we

modelled population and biodiversity change versus overall forest cover change (calculated as

forest cover gain and forest cover loss (GFC database, 2000-2016) and forest loss (LUH

database, across the duration of the time series). Models of population and richness change

versus forest cover change were fitted assuming Gaussian error.

$$\mu_{j,i} = \beta_0 + \beta_{0j} + \beta_1 * duration_{j,i} + \beta_2 * forest\ change_{j,i},$$

$$y_{j,i} \sim gaussian(\mu_{j,i}, \sigma^2),$$

where $duration_{j,i}$ is the duration of the time series in years of cell i within biome j, $forest\ change_{j,i}$

is the forest cover change in cell i within biome j; β_0, β_1 and β_2 are the global intercept and slope

estimates for duration and forest cover change respectively (fixed effects), and β_{0j} are the

biome-level departures from β_0 (biome-level random intercepts); $y_{j,i}$ is the population or

richness change metric (a separate model for population declines, population increases,

richness losses and richness gains) in the ith cell within the jth biome.

Models of turnover versus forest cover change were fit with a zero one inflated beta distribution

to account for the properties of turnover (bounded between zero and one). We used the same

probability density function for the zero one inflated beta distribution as in the model for turnover

before and after contemporary peak forest loss. For $y = 0$ and $y = 1$, α and γ were modelled

assuming a Bernoulli distribution and logit-link function, and we fit models with only an intercept.

For $0 < y < 1$, we assumed a beta error distribution and a logit-link function:

$$logit(\mu_{j,i}) = \beta_0 + \beta_{0j} + \beta_1 * duration_{j,i} + \beta_2 * forest\ change_{j,i},$$

$$y_{j,i} \sim Beta(\mu_{j,i}, \phi),$$

where $duration_{j,i}$ is the duration of the time series in years of cell i within biome j, $forest\ change_{j,i}$

is the forest cover change in cell i within biome j; β_0, β_1 and β_2 are the global intercept and slope

estimates for duration and forest cover change respectively (fixed effects), and β_{0j} are the

biome-level departures from β_0 (biome-level random intercepts); $y_{j,i}$ is turnover in the ith cell

within the jth biome.

Lags in population and biodiversity responses to contemporary peak forest loss

To test for temporal lags in population and biodiversity responses to contemporary peak forest

loss, we first calculated when population and biodiversity change were greatest following peak

forest loss for each time series. Rates of population change were calculated using state-space

models and a Kalman filter[20,34]. Peak richness change and peak turnover were calculated as

the maximum value of the absolute differences between consecutive observations of species

richness and turnover. We then quantified lag as the number of years between contemporary

peak forest loss and peak population/biodiversity change. We modelled lag as a function of

taxa, as we expect that species with longer generation times will respond to disturbance more

slowly.

$$\mu_{j,i} = \beta_{0j} + \beta_1 * taxa_{j,i},$$

$$y_{j,i} \sim gaussian(\mu_{j,i}, \sigma^2),$$

where $taxa_{j,i}$ is the taxa of the cell i in the biome j time series, β_1 is the slope for taxa effect

(fixed effect), and β_{0j} are the biome-level random intercepts; $y_{j,i}$ is the temporal lag in the

population or biodiversity change metric (a separate model for population change, richness

change and turnover) for the ith cell within the jth biome.

We conducted a *post-hoc* analysis where we tested our temporal lag and generation time

hypothesis in a more quantitative manner by modelling lag as a function of generation time in

mammals, the taxa for which generation time data were freely available[55].

$$\mu_g = \beta_0 + \beta_1 * generation\ time_g,$$

721
$$y_g \sim gaussian(\mu_g, \sigma^2),$$

where $generation\ time_g$ is the mammal generation time in years, β_0 and β_1 are the global

intercept and slope (fixed effect); y_g is the temporal lag in population change for a species with

generation time g.

Prior specification

For all models except the model of turnover versus overall forest cover change (which was a

zero one inflated model), we used weakly regularising normally-distributed priors for the global

intercept and slope:

$$\beta_0 \sim gaussian(0, 6),$$

731
$$\beta_1 \sim gaussian(0, 6).$$

732 For the turnover models that had a zero one inflated beta distribution, we used the following

733 priors:

$$\beta_0 \sim gaussian(0, 6),$$

735
$$\beta_1 \sim gaussian(0, 6),$$

$$zoi \sim gaussian(0, 0.5),$$

$$coi \sim gaussian(0, 0.5),$$

where zoi is the probability of being a zero or a one and coi is the conditional probability of being

a one (given an observation is a zero or a one).

Group-level parameters (the rarefied cell random effect in the species richness over time model,

i, and the biome random effect in all models, j) were all assumed to be $gaussian(0, \sigma)$, and priors

on the σ were the same for all models:

$$\sigma\beta_{0j} = \sigma\beta_{0j,i} \sim half\ Cauchy(0, 2).$$

All models were fitted in a Bayesian framework using the *brms* package v2.1.0[53] in R v3.5.1[56].

Models were run for 6000 iterations, with a warm up of 2000 iterations. Convergence was

assessed visually by examining trace plots and using *Rhat* values (the ratio of the effective

sample size to the overall number of iterations, with values close to one indicating convergence).

Sensitivity analyses

Our analyses were not sensitive to our calculation of turnover in the final year of the time series

relative to the first year, and previous examinations of the BioTIME database have found that

calculating turnover relative to the second year of observation produced similar results[4]. We

also quantified population change using the BioTIME database (following the same state-space

modelling framework as with the LPD) and found similar lack of directional patterns in the

relationships between population change and overall forest loss (Extended Data Figure 5f). We

found no distinct geographic or taxonomic patterning in the relationships between population

change, biodiversity change and forest cover change (Extended Data Figure 9). Furthermore,

the relationships between population decreases and increases and forest loss were not

influenced by whether species were tightly associated with forests or not (Extended Data Figure

5g-i). Similar *post-hoc* analysis was not possible for the biodiversity time series because habitat

preference data were not available for many of the species included in the BioTIME database.

The cell size over which we calculated forest cover change (from 10 km^2 to 500 km^2) did not

influence overall findings, as detected forest cover change scaled proportionately with cell size

across locations (Extended Data Figure 10a-b). Landscape context (forest cover in a 500 km^2

cell around sites) also did not influence the relationship between forest cover change and

population and biodiversity change (Extended Data Figure 10c-d). We did not find directional

patterns between population and biodiversity change and time since the largest forest loss event

(Extended Data Figure 10f-h). Our findings were not influenced by the type of forest cover

(primary vs secondary), as loss of secondary forest cover scaled proportionately to primary

forest loss (Extended Data Figure 10e).

Supplementary methods references

48. Gorelick, N. *et al.* Google Earth Engine: Planetary-scale geospatial analysis for everyone.

Remote Sens. Environ. **202**, 18–27 (2017).

49. Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A. & Hansen, M. C. Classifying drivers

of global forest loss. *Science* **361**, 1108–1111 (2018).

50. Knape, J., Jonzén, N. & Sköld, M. On observation distributions for state space models of

population survey data: Observation models for population data. *J. Anim. Ecol.* **80**, 1269–

1277 (2011).

51. Pedersen, M. W., Berg, C. W., Thygesen, U. H., Nielsen, A. & Madsen, H. Estimation

methods for nonlinear state-space models in ecology. *Ecol. Model.* **222**, 1394–1400

(2011).

52. van de Pol, M. & Wright, J. A simple method for distinguishing within- versus between-

subject effects using mixed models. *Anim. Behav.* **77**, 753–758 (2009).

53. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *J. Stat.*

Softw. **80**, (2017).

54. Ferrari, S. & Cribari-Neto, F. Beta Regression for Modelling Rates and Proportions. *J.*

Appl. Stat. **31**, 799–815 (2004).

55. Pacifici, M. *et al.* Generation length for mammals. *Nat. Conserv.* **5**, 89–94 (2013).

56. R Core Team. R: A language and environment for statistical computing. R Foundation   for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/. (2017).

Data and code availability

Code for the rarefaction of the BioTIME Database is available from

https://doi.org/10.5281/zenodo.1475218. Code for statistical analyses is available from

http://doi.org/10.5281/zenodo.1490144. Population and biodiversity data are freely available in

the Living Planet and BioTIME Databases[12,13]. The Living Planet Database can be accessed on

http://www.livingplanetindex.org/data_portal. The BioTIME Database can be accessed on

Zenodo   (https://doi.org/10.5281/zenodo.1211105)   or   through   the   BioTIME   website

(http://biotime.st-andrews.ac.uk/). Land-use change data are publicly available in the Land Use

Harmonization Database[30], the Forest Cover Change Database[31], and the MODIS Landcover

Database[32].