Title: Hunger for Knowledge: How the Irresistible Lure of Curiosity is Generated in the Brain

Authors: Johnny King L Lau[1,2]*, Hiroki Ozono[3], Kei Kuratomi[4], Asuka Komiya[5], Kou Murayama[1,2]*

Affiliations:

[1]School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK.

[2]Centre for Integrative Neuroscience and Neurodynamics, University of Reading, Reading, UK.

[3]Faculty of Law, Economics and Humanities, Kagoshima University, Kagoshima, Japan.

[4]Faculty of Psychology, Aichi Shukutoku University, Nagakute-city, Japan.

[5]Graduate School of Integrated Arts and Sciences, Hiroshima University, Higashi-Hiroshima city, Japan.

*Correspondence to: johnny.lau@reading.ac.uk (Johnny Lau), k.murayama@reading.ac.uk (Kou Murayama)

Abstract

Curiosity is often portrayed as a desirable feature of human faculty. However, curiosity may come at a cost that sometimes puts people in a harmful situation. Here, with a set of behavioural and neuroimaging experiments using stimuli that strongly trigger curiosity (e.g., magic tricks), we examined the psychological and neural mechanisms underlying the irresistible lure of curiosity. We consistently demonstrated that across different samples, people were indeed willing to gamble, subjecting themselves to physical risks (i.e. electric shocks) in order to satisfy their curiosity for trivial knowledge that carries no apparent instrumental value. Also, this seductive power of curiosity shares common neural mechanisms with that of extrinsic incentives (i.e. hunger for food). In particular, we showed that acceptance (compared to rejection) of curiosity/incentive-driven gambles was accompanied by enhanced activity in the ventral striatum (when curiosity was elicited), which extended into the dorsal striatum (when participants made a decision).

Curiosity is a fundamental part of human motivation that supports an enormous variety of human intellectual behaviours, ranging from early learning in children to scientific discovery [1–3]. A meta-analysis indicates that intellectual curiosity predicts academic performance over and above intelligence [4], and corroborating findings show the benefits of curiosity in enhancing long-term consolidation of learning and memory [5,6]. The critical importance of curiosity in human intellectual behaviour is succinctly expressed in Albert Einstein's famous quote "*I have no special talent. I am only passionately curious.*" Moreover, empirical literature has suggested a number of positive outcomes associated with curiosity over the life span [4,7–9]. However, in both historic and modern literature, the positive portrayal of curiosity is often compromised by its inherent negative aspect: strong seductive power [10]. In Greek mythology, for example, after losing his beloved *Eurydice* to the underworld, *Orpheus* convinced the gods to let him take her back to the world of the living on the condition that he would not look back until they had returned. *Orpheus* could not help but look back; he succumbed to curiosity and lost *Eurydice*. This theme, which appears repeatedly in classic ancient anecdotes (e.g., *Pandora*, *Psyche*, *Eve*), illustrates the irresistible *motivational* power of curiosity, which biases our decision making despite the knowledge of consequential negative outcomes.

How does curiosity for inconsequential knowledge motivate people to make risky decisions? Although there have been various theories on the construct and origin of curiosity in the literature [10–12], a recent body of research has seen an emerging consensus that, like food and other extrinsic rewards, curiosity can be understood as a reward-learning process of knowledge acquisition or information seeking [8,13–16]. According to this account, individuals are motivated to actively seek knowledge because knowledge acquisition serves as an inherent 'reward' (possibly due to uncertainty reduction and potential anticipatory utility[17]), reinforcing further information

seeking or behaviours that promote information gain. In fact, studies have repeatedly demonstrated that both animals [18–20] and humans [6,21–23] are willing to pay small amounts to satisfy their curiosity for knowledge about a future reward that cannot be changed. For example, people are willing to sacrifice parts of future monetary reward in exchange for immediate information just to learn about the outcome of a monetary lottery, even though that information cannot be used to alter prospective lottery outcomes[21].

Past work on how extrinsic incentives shape behaviour and decisions highlight the involvement of a strong, visceral motivational feeling, known as incentive salience ('wanting') [24,25]. Incentive salience can readily be triggered by encountering reward cues and by vivid imagery of reward. Also, it is hypothesised to integrate the current physiological neurobiological state and previously learned associations about the reward. For example, when you arrive hungry to a restaurant, you may start salivating and find every item on the menu to be more appealing than they usually are to you. The motivational urge of incentive salience may even prompt people towards irrational, impulsive and even obsessive behaviours[26,27]. Previous animal research suggested that incentive salience is mediated by the mesolimbic dopamine system as well as the surrounding relatively large brain network including both cortical and subcortical areas [28,29], whereas neuroimaging studies in human have identified the particular involvement of the ventral striatum (i.e. the nucleus accumbens) and the dorsal striatum (i.e. the caudate) in the processing of different extrinsic rewards [30–32]. For example, Lawrence et al. showed that the activation in the nucleus accumbens in response to food cues predicts participants' consumption of crisps when they are left alone in a room after the experiment [33].

Integrating these ideas and findings, we suggest that, like extrinsic incentives such as food, the rewarding property of knowledge may evoke incentive salience, energising people

toward knowledge acquisition behaviour even if it entails significant risks [15,34]. In fact, although the neurobiological mechanisms underpinning curiosity-based decisions have been under-examined in the literature, the investigations using brain imaging collectively suggest that the states of curiosity about information that does not signal any future rewards modulate neural regions in dopamine systems, including caudate nucleus, nucleus accumbens and midbrain structures (i.e. ventral tegmental area/substantia nigra [VTA/SN])[5,6]. As discussed previously, these areas, with the caudate nucleus and nucleus accumbens in particular, have been implicated in incentive salience in neuroimaging studies of cognitive processing related to food stimuli[30]. A recent study using a neurocomputational approach and a monetary gamble paradigm also found that the same patterns of activations in striatum responding to the values of potential rewards per se (i.e. the varying amount of monetary gain) can predict subjective value of information (i.e. how much a person would be willing to pay for extra information), which may be partially driven by non-instrumental, curiosity-based motives[17]. Taking these together, it is possible that decisions driven by a strong urge of knowledge acquisition rely on similar mechanisms that support motivated behaviours triggered by incentive salience of extrinsic incentives[35,36].

The current study examines whether and how people are willing to subject themselves to physical risks to satisfy their curiosity for trivial, inconsequential knowledge, by using a novel 'curiosity-based risky decision-making' paradigm in a series of behavioural and fMRI experiments. In the 'curiosity condition', participants first saw a curiosity-stimulating stimulus (i.e. video clips of magic tricks, or text-based trivia questions), followed by a wheel of fortune which visually depicted the probability of winning (versus losing) a lottery in each trial. Participants then had to make a decision regarding whether or not they would gamble on the lottery (risking electric shock) to have a chance of learning the solution to the trick. The

experiment also included food trials, in which participants were presented with food pictures and decided whether to gamble (risking electric shock) to have a chance of obtaining the food. The inclusion of the food condition allows us to directly compare the influence of curiosity and extrinsic incentives on risk-based decisions and to examine the common and differentiated mechanisms supporting them. We postulated that i) people are willing to subject themselves to physical risks (i.e. electric shocks) to satisfy their curiosity for trivial, inconsequential knowledge (i.e. magic tricks, trivia) or their hunger for foods, and ii) the motivational lure of curiosity and hunger share common neural correlates implicated in the incentive salience of food, namely the ventral and dorsal striatum [30].

Results

In the decision-making task, participants viewed short video clips of magic tricks (curiosity condition) performed by professional magicians (filmed specifically for this study; Supplementary videos 1-2), followed by a wheel of fortune visually depicting the outcome probability of winning (versus losing) a lottery (with a certain probability; between 16.7% - 83.3%; see Methods) in each trial (Fig. 1). Participants were then asked to make a decision regarding whether they would gamble on the lottery or not. They were told that, if they accepted to gamble and won, they might see the secret behind the magic trick after the experiment as a reward. The experiment also included a food condition, in which participants were presented with food pictures and were told that they might eat the food as a reward after the experiment if they accepted to gamble and won. Critically, for both types of trials, if participants gambled and lost, they would expect to receive electric shock after the experiment. The magnitude of electric shock was calibrated and demonstrated to participants before the experimental task. Stimulating electrodes were attached to the participants throughout the task and the shock was expected to be delivered at the very end. For each stimulus, participants gave a rating of how curious they were to see the solution to the trick (curiosity rating) or how much they would like to eat the food item (desirability rating). To maximise participants' desire for food during the experiment, they were required to refrain from food consumption shortly before attending the testing session. Additional details on the task are reported in the Methods section: see 'Procedure'.

[INSERT FIG 1]

Behavioural experiments

Curiosity and food desirability bias decision-making. In the initial behavioural

experiment (N = 32), generalised linear mixed-effects modelling (GLME) on all trials showed

that participants were more likely to reject the gamble as the presented outcome probability of

receiving electric shock increased [GLME: Z = -19.045, P < 0.001, Exp(β) = 0.455, β = -0.787,

95% C.I. = -0.868 – -0.706]. This indicates that electric shock worked effectively as an aversive

stimulus. Importantly, above and beyond the shock probability, stimulus rating (i.e. curiosity or

food desirability rating) positively predicted the 'accept' decision [GLME: Z = 20.253, P < 0.001,

Exp (β) = 2.110, β =0.747, 95% C.I. = 0.674 – 0.819]. There was no significant main effect of

'Incentive Category' (curiosity vs. food) on the decision [GLME: Z = 1.25, P = 0.211, Exp(β) =

1.067, β = 0.065, 95% C.I. = -0.037 – 0.167]. Also, none of the interaction effects between the

stimulus rating, presented probability, and incentive category (including a three-way interaction)

were statistically significant, suggesting that the observed effects of rating were consistent across

different categories and levels of shock probability (all Ps > 0.05). In fact, when we analysed

curiosity trials and food trials separately, in addition to outcome probability, ratings of curiosity

and food desirability were similarly associated with higher tendency to accept the gamble

[GLME - curiosity: Z=7.236, Exp(β) = 3.185, β = 1.159, 95% C.I. = 0.845 – 1.472; food:

Z=7.493, Exp(β) = 3.145, β = 1.146, 95% C.I. = 0.846 – 1.446; Ps < 0.001 in both conditions]

(Fig. 2; also see Supplementary Table 1).

19   [INSERT FIG 2]

21 ***Did curiosity distort the perception of the probability of outcome?*** As shown in the

22 initial experiment, people are willing to subject themselves to potential risks to satisfy their

23 curiosity for trivial, inconsequential knowledge, corroborating our hypothesis. However, an

alternative explanation may be that curiosity simply distorts one's perception of the probability

of risks. People tend to report higher probabilities for positive events and lower probabilities for

negative events when they are in a pleasant mood [37]. It could be that when people are curious,

feel more motivated and experience positive feelings, they overestimate the probability of

winning in the lottery (or underestimate the risk). To evaluate this possibility, we carried out a

follow-up behavioural experiment. In this experiment, after seeing a curiosity-/hunger-evoking

stimulus, participants (N = 29) estimated their subjective chance of winning(/losing) in the

lottery when presented with a wheel of fortune in each trial, before also making their actual

decision to gamble or not (see Supplementary Methods for the procedures of this modified task).

First, we applied the same GLME to predict participants' decision as an attempt to check

whether the results from the initial experiment could be replicated. In this follow-up experiment,

in addition to the presented shock probability [GLME: Z = -17.334, P < 0.001, Exp(β) = 0.251, β

= -1.381, 95% C.I. = -1.537 – -1.225], stimulus rating [GLME: Z = 19.568, P < 0.001, Exp (β) =

3.947, β = 1.373, 95% C.I. = 1.236 – 1.511] again positively predicted the decision to gamble.

There was no significant main effect of 'Incentive Category' (curiosity vs. food) on the decision

[GLME: Z = -0.505, P = 0.613, Exp (β) = 0.958, β = -0.043, 95% C.I. = -0.208 – 0.123]. Also,

none of the interaction effects between the presented probability, stimulus rating, and incentive

category (including a three-way interaction) was statistically significant (all Ps > 0.05). These

results are consistent with those from the initial experiment (see Supplementary Table 1, for

separate analysis of curiosity and food conditions).

As the main objective of the follow-up experiment, we analysed the predictors of

participants' subjective estimation of the chance to win/lose with a linear mixed-effects model

(LME). Participant's own estimation of win/loss probability was predicted by the presented

shock (outcome) probability that they actually viewed in each trial [LME - with all trials: t = -27.522, β = -1.762, 95% C.I. = -1.887 – -1.637; with curiosity trials only: t = -25.519, β = -1.798, 95% C.I. = -1.936 – -1.660; with food trials only: t = -26.27, β = -1.728, 95% C.I. = -1.857 – -1.599; Ps < 0.001 in all cases], which perhaps is not too surprising. In contrast, stimulus rating [LME - with all trials: t = 1.565, P = 0.128, β = 0.045, 95% C.I. = -0.011 – 0.102] did not predict the participant's subjective outcome estimation. There was also no significant interaction effect of 'Rating' x 'Incentive Category' [LME: t = -0.311, P = 0.756, β = -0.004, 95% C.I. = -0.028 – 0.021]. In fact, when investigating each category separately, both curiosity rating for magic tricks [LME: t = 1.618, P = 0.118, β = 0.033, 95% C.I. = -0.007 – 0.072] and desirability rating for food items [LME: t = 1.406, P = 0.17, β = 0.045, 95% C.I. = -0.018 – 0.108] were not associated with the subjective estimation of the outcome. Comparing the fits of two linear mixed-effects models, one with a 'stimulus rating' term and one without, the one without the rating term had a better fit [with all trials, Bayes Factor (BF01) = 0.00559; for magic condition only, BF01 = 0.00572; for food condition only, BF01 = 0.00661; very strong support for the null hypothesis in all cases]. This means one's subjective estimation of the chance to win/lose is not modulated by the curiosity for information or how desirable the food is.

Neuroimaging experiments

To examine the neural mechanisms underlying motivational biases on decision-making, we conducted a pair of neuroimaging experiments using fMRI, in which separate groups of participants performed a similar experimental task as in the initial behavioural investigation. The two versions of the fMRI experiment were constructed to be very comparable but differed in the materials used to induce curiosity, an attempt to optimise the generalisability of the findings. The

first version (N=31), used magic tricks as curiosity-evoking stimuli (ver1: 'magic version')

whereas the second version (N=30), used trivia questions (ver2: 'trivia version'). While magic

video clips trigger strong curiosity [38], a potential weakness is that they may exert extra demands

on visual/perceptual attention and processing for participants, which could be a confound. Trivia

questions, on the other hand, have often been used in past experiments of curiosity [5,6,39,40], and

the visual input is relatively minimal. By running this pair of experiments using different stimuli,

we are in a better position to ensure our investigation reflects the effect of curiosity other than

some stimulus-specific effect, which has recently been noted as a critical issue in fMRI research

[41]. In addition to the curiosity condition, both versions of the experiment included food trials. We

analysed the two versions altogether in order to statistically evaluate the commonalities and

differences in neural activation patterns across different types of experimental materials.

Behaviourally, participants from the fMRI experiments (N = 61) accepted the gamble

more often when the presented shock probability decreased [GLME: Z = -30.357, P < 0.001,

Exp(β) = 0.337, β = -1.087, 95% C.I. = -1.157 – -1.017]. Also, on top of the shock probability,

participants decided to take the gamble more often when they gave a higher rating to a stimulus

in the trial (i.e. higher curiosity or food desirability) [GLME: Z = 28.572, P < 0.001, Exp (β) =

2.121, β = 0.752, 95% C.I. = 0.700 – 0.804]. Unexpectedly, the fMRI experiments showed a

significant interaction between 'Rating' and 'Incentive Category' [GLME: Z = -8.937, P < 0.001,

Exp (β) = 0.816, β = -0.203, 95% C.I. = -0.248 – -0.159], indicating stronger effects of food

desirability than curiosity on decisions. Yet it is worth noting that the effect of rating was still

very significant when each category was investigated alone (See Supplementary Table 1). There

was also a significant 'Rating' x 'Probability' interaction effect [GLME: Z = -2.786, P = 0.0053,

Exp (β) = 0.942, β = -0.061, 95% C.I. = -0.103 – -0.018]. Separate analyses for curiosity and

food conditions indicated that this interaction effect appeared to be driven mainly by the food

condition, which also exhibited an interaction effect between 'Rating' and 'Probability'.

Specifically, the effect of food desirability seemed to become stronger at lower rates of shock

probability [GLME: $Z = -3.39$, $P < 0.001$, Exp (β) = 0.905, $\beta = -0.100$, 95% C.I. = -0.158 – -

0.042]. In contrast, the interaction effect was not significant in the curiosity condition ($P =$

0.397). Since these interaction effects are only specific to these fMRI participants and not

consistent with the previous experiments, we will not pursue them further in the current

investigation (also see Discussion for a potential interpretation). No other interaction effects

were significant ($Ps > 0.05$). Moreover, an 'Experiment Version' effect was also taken into

account in this GLME with data from the fMRI participants and this effect on decisions was not

significant ($P = 0.662$).

Our first fMRI analyses identified brain areas that are recruited at different stages of

motivated decision-making and focused on contrasting the trials in which participants accepted

the gamble with those that they chose not to (Fig 3a). Based on past studies of extrinsic reward

anticipation [32,42] and curiosity-guided behavioural change [5,6], we hypothesised that activity in the

caudate nucleus, NAcc, and VTA/SN would be modulated during the course of

curiosity/incentive-driven decision-making. As a number of separate region-of-interest (ROI)

masks (one for each of the 3 regions) were tested, we corrected for multiple comparisons across

them using a Bonferroni-adjusted alpha level of 0.0167 (familywise error corrected, FWE) per

ROI analysis. We focused on the brain activation during the elicitation phase (i.e. when stimuli

were presented) and decision phase (i.e. when participants made the decision) in a trial.

[INSERT FIG 3]

The activity of nucleus accumbens during elicitation of curiosity and food desirability.
Comparing all accepted trials with the rejected ones ('Decision' contrast) at the elicitation phase revealed enhanced activations specifically in bilateral NAcc ($P_{FWE} < 0.0167$, for ROI analysis) among all ROIs (Fig 3b; Supplementary Table 2). There was no significant variation in activation due to the interaction of 'Decision' (accept vs. reject) x 'Experimental Version' (magic version vs. trivia version), indicating limited evidence that the type of curiosity-triggering stimuli used had differential influence on neural responses. On the other hand, a 'Decision' x 'Incentive Category' (curiosity vs. food) interaction effect was shown, with the 'accept > reject' effect being potentially stronger for food than curiosity (in a small cluster of voxels in the left NAcc). (Supplementary Table 4).

It is also interesting to observe that while rejected gambles (when compared with accepted ones; i.e., reject > accept 'Decision' contrast) did not yield any enhanced responses in the ROIs, an exploratory whole-brain analysis revealed greater activations in the insula and dorsomedial prefrontal cortex encompassing anterior cingulate cortex ($P_{FWE} < 0.05$, for whole-brain analysis) (Supplementary Fig 1; Supplementary Table 3), an area previously implicated in value computation[43,44] and decision conflict[45,46].

Decision to gamble is associated with activity in the brain's reward network (neural activations at the decision phase). At the decision phase, a main 'Decision' contrast (i.e. accept vs. reject) on our anatomical ROIs showed that bilateral caudate nucleus, NAcc, and VTA/SN were indeed more activated for the accepted trials in comparison with the rejected trials (Fig 3b). An exploratory whole-brain analysis of the same contrast also indicated that these striatal and midbrain structures had extensive activations along with other medial brain areas including the cingulate gyrus and medial frontal cortex, as well as the right anterior insula, inferior frontal

gyrus and lateral prefrontal cortex (although the peaks were still located in the caudate nucleus) (Supplementary Fig 2; Supplementary Table 5). On the other hand, no significant activation was observed for the reversed contrast (i.e. reject > accept 'Decision') at decision phase (Supplementary Table 6). There were also no significant variations in brain activation within the ROIs for the interaction effect of 'Decision' x 'Experimental Version' (magic version vs. trivia version) or 'Decision' x 'Incentive Category' (curiosity vs. food), suggesting that neither the type of curiosity-inducing stimuli nor the category of incentive had significant differential influence on neural responses in the reward system at the decision phase (see Supplementary Tables 7 & 8).

Next, we examined whether the association between the motivated decision and the striatal activity would still exist after considering the potential effect of the shock (outcome) probability that was shown with the lottery during the decision phase. To this aim, we performed a parametric modulation analysis with another general linear model (GLM), in which the decision to gamble and the presented probability were included as simultaneous parametric modulators for each regressor (i.e. one regressor for curiosity condition; one for food condition) at the decision phase (see Experimental Procedures for details). Indeed, the association was still robustly observed in the parametric modulation analysis, even with the presented probability of winning/losing the lottery taken into consideration. Specifically, the activations in bilateral caudate nucleus, NAcc and VTA/SN were shown to be parametrically modulated by the decision of accepting (versus rejecting) to gamble with the ROI approach ($P_{FWE} < 0.0167$ in all ROIs; Supplementary Fig. 3), regardless of the incentive category. Moreover, a follow-up whole-brain analysis again revealed extensive neural responses in striatal and midbrain structures, along with the thalamus and right frontal cortex (Supplementary Table 9).

Striatal activities mediated the relationship between curiosity/food desirability and the gamble decision. Our behavioural data indicated that the decision to accept/reject a gamble was predicted by the rating of curiosity and food desirability. fMRI results also showed that the decision to accept/reject a gamble was associated with the changes in neural responses in the striatum. Given the literature that the ventral and the dorsal parts of the striatum have been related to different stages of decision-making (the ventral striatum for value computation and the dorsal striatum for action selection [47–49]; also elucidated in Discussion in more detail), we evaluated whether the link between rating and the decision was mediated by the activities of the ventral striatum (NAcc) at the elicitation phase and dorsal striatum (caudate nucleus) during decision-making on a trial-by-trial basis. The mediation model was tested using a multilevel structural equation model (with trials being nested within participants), and took into account the incentive category and presented shock probability in each trial as well as the experiment version. This multilevel mediation analysis confirmed that stimulus rating predicted choice through the phase-specific activities of nucleus accumbens (at elicitation) and caudate nucleus (at decision-making) (see Fig 4; Supplementary Table 10). The three-path indirect effect (Rating → NAcc at elicitation → caudate nucleus at decision → accept/reject decision response) was statistically significant ($P = 0.044$). These results support the idea that curiosity and food desirability induced by the stimuli prompted participants to make risky decisions through modulating phase-specific striatal activities.

[INSERT FIG 4]

Exploratory functional connectivity analysis

To further explore the modulation of functional interaction between brain regions during

decision-making, we conducted a functional connectivity analysis based on a beta-series

correlation approach using the caudate nucleus as the seed. This approach compared correlations

of trial-by-trial beta series variability (i.e. variability of series of parameter estimates) between

the seed and the rest of the brain in different conditions. We corrected for multiple comparisons

across the results using separate left and right caudate as the seed with a Bonferroni-adjusted

alpha level of 0.025 per analysis. Using the left caudate nucleus as a seed, this exploratory

analysis found weaker connectivity with the left sensorimotor area near precentral gyrus and

central sulcus, suggesting a decoupling effect (i.e. weaker functional correlation), when

accepting the gamble in comparison with rejecting it. Taking the right caudate nucleus as a seed,

we again found weaker connectivity with the left sensorimotor area (a cluster near central sulcus

and postcentral gyrus) in accepted (relative to rejected) trials ($P_{FWE} < 0.025$ in both analyses; Fig.

3, Supplementary Table 4). The sensorimotor cortex has been associated with virtual feeling and

anticipation of pain in action observation [50–52]. Although speculative, our results might suggest

that, when a participant accepted the risk of receiving electric shock, there could be a reduced

association between the motivation of satisfying their desire and expected fear of physical shock

in the brain. There was no significant variation in brain connectivity shown with the reversed

contrast (reject < accept 'Decision') and also no interaction effects ('Decision' x 'Experiment

Version' or 'Decision' x 'Incentive Category').

[INSERT FIG 5]

Discussion

Using a novel curiosity-based decision-making paradigm, the current study aimed to examine whether and how the seductive lure of curiosity overcomes the potential fear of significant risks. Consistent with our speculations, the results showed that i) both curiosity for inconsequential knowledge (magic tricks, trivia questions) and hunger for food indeed prompted participants to subject themselves to physical risks (i.e. electric shocks), and ii) this motivated decision-making was supported by the striatal reward areas. Specifically, the activation in the ventral striatum in response to curiosity-evoking stimuli predicted the decision to accept the risky option through the activation in the dorsal striatum during the decision phase. These results are consistent with the notion that incentive salience plays a role in impulsive behaviours caused by curiosity in a similar manner as it does with extrinsic incentives (e.g., food).

This study extends on the existing neuroscientific literature of curiosity[53], which primarily focuses on its influence on memory and learning [5,6,39], and our findings provide evidence that corresponds with the idea of a potential role for incentive salience in curiosity-driven risky decision. The seductive nature of curiosity in the decision-making process has been recognised in the literature [54]; this point has also been demonstrated in recent studies on 'morbid-curiosity', where people actively seek negative stimuli (e.g., negative pictures, sounds) in order to satisfy their curiosity for experiencing these uncertain but negative stimuli themselves[55–57]. However, the underlying mechanisms have still been a matter of debate. One of the predominant (but controversial) explanations is that curiosity arouses an aversive state, and it is this aversive state that urges people to acquire information to satisfy their curiosity [10,11]. However, there has been little empirical evidence that directly supports the assumption that curiosity involves a negative aversive state [12] (bar for a few exceptions [58]). Although we did not assess the emotional

valence of participants' feelings of curiosity and therefore cannot directly refute this hypothesis, our findings lend greater support to the incentive salience hypothesis of curiosity, which addresses the strong motivational power of curiosity without making such a restrictive assumption of the aversive nature of curiosity. The incentive salience hypothesis can also be reasonably incorporated into reinforcement learning models of decision-making [26], providing a more parsimonious and integrative view of the function of curiosity.

One particularly interesting finding was that curiosity-based decision-making was supported by the different patterns of striatal activation during the elicitation phase (i.e. the ventral striatum) and the decision phase (i.e. broader part of the striatum peaked at the dorsal striatum); our mediation analysis suggested that the ventral striatum and the dorsal striatum sequentially mediate the relationship between participants' subjective motivation (i.e. curiosity and hunger for foods) and risky decision-making. These findings correspond with the notion that the ventral striatum computes the value of the stimulus ('critic') while the dorsal striatum plays a role in selecting an action/response ('actor') [47]. Several empirical studies provided supportive evidence for the idea but they were all limited to extrinsic incentives [48,49,59]. Our findings suggest that the neural computation of curiosity-based decision-making can possibly be understood within the actor-critic model of reinforcement learning. Like many other functional connectivity studies, however, our mediation model is correlational (although there is a temporal separation between the elicitation phase and the decision phase); future research should design a more fine-grained experiment to allow for examination of the dissociable causal role of the ventral and the dorsal striatum in curiosity-based decision-making.

Another interesting observation was that we found very few differences in brain activation between different types of incentives (i.e. food and curiosity) and between different

types of curiosity-evoking stimuli (i.e. magic trick and trivia questions) overall as far as the decision-making process is concerned (i.e. when comparing accepted with rejected gambles). These findings are consistent with the view that, regardless of the types of incentives, the decision-making process can be portrayed with a common reward-learning framework[13,16,60]. Note that we in fact observed a few differences between the conditions. Food stimuli seemed to have stronger behavioural effects than curiosity-evoking stimuli in the fMRI experiments (but not in the two behavioural experiments), and this was also reflected in the enhanced activation in a small cluster of voxels in the ventral striatum (NAcc) during the elicitation phase in the food condition than in the curiosity condition. Also, when participants accepted a gamble (vs. rejected it), the magic trick version of the experiment seemed to induce stronger activation in a few voxels in the VTA and SN during the decision phase than the trivia question version; this result suggests a potentially stronger curiosity effect of magic tricks in comparison to trivia questions. However, such differences are all concerned with the *magnitude* of the motivational effect which is likely to reflect diverse factors irrelevant to *how* the decision has been made. For example, participants in the fMRI experiments had a prolonged session before the main task actually took place in comparison to those in the two behavioural experiments, due to extra time taken for preparation and mandatory paperwork to undergo fMRI scan. It is possible that this delay bolstered the hunger of participants (who already had a fasting period before coming to the study), causing the stronger effects of hunger both at the behavioural and neural levels. Importantly, however, the incentive salience hypothesis can easily explain the difference in the magnitude of the effects (i.e. food induced stronger incentive salience due to increased hunger) without supposing qualitatively different mechanisms.

Our current findings provide some new insights into how curiosity motivates risky behaviour; however, they should be interpreted with caution. Striatal activation has also been implicated in functions other than simple reward processing and incentive salience [61,62], thus it is still possible that the observed activation may reflect other decision-making mechanisms than incentive salience. Future studies are needed to examine the neurobiological mechanisms more comprehensively, for example through the use of different methodologies and experimental paradigms such as computational modelling and pharmacological intervention. Moreover, our exploratory analyses revealed some interesting observations that have not been thoroughly discussed in the literature. For example, despite being speculatively only, the decoupling of the dorsal striatum and sensorimotor areas might suggest a neural pathway through which the motivational force of curiosity modulates actual behaviour. The whole-brain fMRI analyses suggested several brain areas related to curiosity-based decision-making, including the anterior cingulate cortex, lateral prefrontal cortex, inferior frontal gyrus and anterior insula. These brain areas were also reported in previous literature on curiosity [5,6,53,57,63], as well as in the studies of intrinsically-motivated behaviour in general [64]. There is a possibility that curiosity is a part of broader autonomous self-regulatory functioning [15], which involves the coordination of wide range of brain areas beyond the reward network. These ideas need to be explicitly tested and confirmed in future studies with paradigms directly tapping into these processes.

Methods

Participants

The study was approved by the research ethics committees of the University of Reading,

UK (ethics approval number: UREC 16/36). Participants were recruited via mailing lists and a

research participation pool (managed through SONA Systems) at the University. Participants

provided informed consent, completed and passed a health and safety screening on the eligibility

of receiving electric stimulation to confirm that i) they did not have a cardiac pacemaker (or any

other devices that can be affected by electric stimulation); ii) they were free from

neurophysiological symptoms or conditions including peripheral vascular disease, vasculitis

cryoglobulinemia, lupus, tingling or numbness in hands and/or feet. To maximise participants'

desire for food during the experiment, they were required not to eat or drink anything (apart from

water) within 2 hours prior to the testing session.

Depending on personal preference, participants were compensated either with course

credits at a fixed rate of 1 unit per hour or cash payments for their participation. For behavioural

experiments, a fixed rate of £7 per hour was given, while for fMRI experiments the

compensation for participation was fixed at £10 per hour. In addition, participants were informed

that they may receive extra rewards (i.e. food and solutions to magic tricks or trivia questions)

according to their task performance in the experiment. Each participant took part in only one

version of the experiment.

Initial behavioural experiment: Initially, seventeen healthy individuals were recruited; an

additional fifteen participants were included later in order to achieve more reliable estimates in

data modelling. A final sample of 32 participants (8 males) were 21.90 years old (sd= ±2.66) on

average. When analysing these data, we controlled for the potential inflation of Type-1 error rate resulting from an interim analysis performed (see Methods: Behavioural analyses below).

Follow-up behavioural experiment (on subjective outcome estimation): Twenty-nine healthy individuals (7 males) with a mean age of 20.17 years (sd=±3.96) participated in this experiment.

fMRI experiments: The first version (magic version) of the experiment included thirty-two individuals. One of them accepted every single gamble (100% acceptance on all trials) in the decision-making task and was thus excluded prior to data analysis. In the second version (trivia version), thirty-two participants were also recruited. One of them accepted every single gamble (100% acceptance on all trials) and another participant had pronounced head movements during MRI scan (> 3mm displacement in a motion direction, see Supplementary Methods: fMRI preprocessing for further details). Both were excluded prior to data analysis. The final sample from both versions comprised a total of 61 participants (11 males) with a mean age of 19.92 years (sd= ± 2.11) and are all right-handed.

For further explanation of the decisions regarding our sample size, see Statement on Statistics and Reproducibility below.

Materials

Food images: All pictures were colour photographs selected from different sources on the Internet. They had a resolution of at least 512 x 384 pixels and were edited so that the single food item was presented in the centre against a white background using GNU Image Manipulation Program (GIMP) 2, a free open-source graphics editor. A selection of food was chosen based on the following criteria: first, the items would be familiar to participants to avoid hesitation due to

uncertainty during decision-making; second, there was a wide variety of items including fruits/vegetables, sweets, snacks and savoury bites (e.g. grapes, salad, chocolate, nuts, sausage roll, etc.), which might accommodate different individual preferences and tastes for food and elicit different levels of desirability in participants.

Magic trick videos: Magic tricks, performed by three professional magicians including a champion of an international magic competition, were recorded in a TV studio with a professional cameraman using high resolution video cameras. All videos were then edited using Adobe® Premiere Pro CC® (2015) software to a similar monotonic (dark) background, size (720 x 404 pixels) and viewing focus. The videos were muted (and subtitles were added in a few videos, when needed). The face of the magician was hidden to avoid potential distraction due to appearance and any facial expressions (see Supplementary videos 1-2). Out of the pool of 166 video clips, we selected magic tricks to be used in the current study by ensuring that they (1) included a range of different features (such as the use of cards, sleight of hands, optical illusions) and (2) likely elicited curiosity to different extents (based on curiosity ratings obtained in a different pilot study). Our initial behavioural experiment included 45 food items and 45 magic trick videos. These videos ranged between 8 – 46 s in length (mean=22.22 s; median=20 s). Due to time constraints (an increase in duration of each trial with an extra event), the follow-up 'subjective outcome probability estimation' experiment used only a subset of 36 food items and 36 magic tricks from the initial experiment (length of the videos: range= 8-46 s; mean=20.61 s; median=18.5 s). The magic version of the fMRI experiment used the same subset of stimuli as the follow-up experiment.

Trivia questions: Sixty trivia questions were selected from a publicly-available 244-item

database [65] (http://koumurayama.com/resources.php). The selected questions were not obviously

culture or age specific. Also, for all questions, the answers were likely to be unknown to the

majority of participants. The selection of items corresponded to different trivia categories that

might elicit curiosity among individuals to different extents, including art/music,

history/geography, movies/TV, nature/animals, science, space, sports, food, as well as other

miscellaneous facts. To ensure within-person variability in curiosity evoked across the

experiment, half of the chosen questions were picked among those with high mean curiosity

scores in the database and the other half among those with low mean scores (rated by a sample of

1498 respondents from a separate online study; for more information, refer to [65]). The trivia

version of the fMRI experiment used 60 trivia questions, as well as 60 food items. On average,

the chosen questions contained 10 (ranging between 6-16) words.

Experimental Procedure

Curiosity-driven Decision-Making Task

The main task of the study followed similar procedures (although there were slight

modifications for the follow-up behavioural and fMRI experiments, which are detailed in

Supplementary Methods). In brief, each trial started with a central fixation cross and then a brief

letter cue ('M' signified 'magic trick', or 'T' for 'trivia question' in the 'trivia version' fMRI

experiment; 'F' signified 'food') to prepare participants for the kind of stimulus they were about

to see. This was followed by the stimulus, which was either a video of a magic trick (or a trivia

question in the 'trivia version' fMRI experiment) in the curiosity condition or an image of a food

item. Participants then gave a rating to indicate their level of curiosity about the magic trick (i.e.

how curious they were to see the solution to the trick) or level of desirability of the food (i.e.

how much they would like to eat the food), on a 7-point scale (1=not at all, 7=very much). In

curiosity trials, they also had to report how confident they were that they knew the solution to a

magic trick, using also a 7-point scale (1=not at all, 7=very much). However, because rated

confidence was not associated with a participant's decision and the inclusion of this measure did

not change any main results, to make straightforward comparisons between curiosity and food

trials, it was not included in the reported analysis here. After rating the stimulus, participants

were presented with a wheel of fortune (WoF) representing a lottery which visualised the

probability of them winning (the chance of getting the reward) versus losing (the chance of

receiving electric shock) in that trial, and were asked to decide whether to gamble or not.

Participants were instructed that if they accepted the gamble and won, they would receive a

token that might allow them to see the secret behind the magic trick/get the food item after the

experiment. If they gambled and lost, they would get a token that might increase the duration of

shock they were to experience at the end of the experiment. Participants could also opt to reject

the gamble. At the end of each trial, the outcome of the lottery was presented. Participants were

informed that the final amounts of rewards and shock to be delivered would be probabilistic as

determined by a mathematical algorithm, but with more tokens collected, the likelihood of

getting more solutions, food, and shock would be augmented. Participants had an understanding

that, as a general rule, the more 'win' tokens they collected, the more likely they would get to see

more solutions to the tricks and obtain more food items (i.e. the rewards). Similarly, the more

'loss' tokens they got, the more likely they would receive more electric shock.

  There were 5 versions of WoF, each displaying a different combination of the

probabilities of winning and losing a gamble: i) 16.7% (1/6) win vs. 83.3% (5/6) loss; ii) 33.3%

(2/6) win vs. 66.7% (4/6) loss; iii) 50% (3/6) win vs. 50% (3/6) loss; iv) 66.7% (4/6) win vs.

33.3% (2/6) loss; v) 83.3% (5/6) win vs. 16.7% (1/6) loss. In the experiments, participants were

never shown the actual percentages but the relative win-to-loss contrast of probabilities was

illustrated visually by the relative sizes of the constituent slices on the WoF (Fig. 1a). To control

for the number of 'success' and 'loss' experiences, unbeknownst to the participants, there was an

equal chance of winning or losing in the lottery across all trials.

The curiosity and food trials were mixed and shown in a random order. The task was

programmed and presented using PsychoPy [66].

Program on the day of the testing session

Participants were asked not to consume any food and drinks (apart from water) within at

least 2 hours before attending their testing session, so as to maximise their desire for food during

the experiment. This was confirmed with the participants at the beginning of the study, including

asking them to indicate when they last ate and had their last meal. Following standard procedures

of informed consent and completing corresponding health and safety screening, participants

underwent calibration for electric stimulation to identify a maximum (uncomfortable yet non-

painful) threshold of electric shock they can endure (see Supplementary Methods for details).

Participants did not receive electric shock during the experiment although stimulating electrodes

were continuously attached to them. As informed by a pilot study, expectation of electric shock

would have more sustained effects on fear perception than receiving actual electric shock

multiple times. After the calibration procedures, participants were then given the instructions on

the decision-making task (also presented via PsychoPy). There was a practice task that used a

different set of stimuli (3 curiosity, 3 food) prior to the main task.

At the end of the experiment, participants were asked in a questionnaire the extent to which they expected to receive electric shock. In this post-experiment questionnaire, the majority of participants across the follow-up behavioural and fMRI experiments reported prospectively that they expected to receive the electric shock during the experiment - 89% of them gave a rating of 3 or above out of 5 (mean=3.67, mode=4), representing their belief that the shock would have happened (we do not have this information from the initial behavioural study). Wherever relevant, in our analysis this shock expectation rating was included as a covariate variable. After completing the questionnaire, rewards (i.e. solutions to some magic tricks and food items) were delivered. Participants did not actually receive any electric shock. They were told that as determined by a probabilistic equation programmed in the task, they were assigned with zero or negligible shock at the end.

fMRI acquisition

For participants in the fMRI experiments, whole-brain functional and anatomical images were acquired using a 3.0 Tesla Siemens MAGNETOM scanner with a 32-channel Head Matrix coil at The Centre for Integrative Neuroscience and Neurodynamics (CINN), University of Reading, UK, in a single one-hour scanning session.

Magic version fMRI experiment

Functional images were acquired using a T2*-weighted gradient-echo echo planar imaging (EPI) pulse sequence with 37 axial slices (in-plane resolution of 3 x 3 x 3mm, interslice gap: 0.75mm), interleaved from bottom to top (echo time (TE): 30 ms; repetition time (TR): 2000 ms; flip angle: 90˚; field of view (FOV): 1344 x 1344 mm^2; in-plane matrix: 64 x 64). A

high-resolution T1-weighted three-dimensional anatomical image was also collected, using an MPRAGE-gradient sequence with 176 x 1mm slices (in-plane resolution of 1 x 1 x 1 mm; TE: 2.52 ms; TR: 2020 ms; Inversion Time (TI):1100 ms; FOV: 250 x 250; flip angle: 9°), enabling optimal localisation of the functional effects.

Trivia version fMRI experiment

Functional images were acquired using the same sequence and parameters as in fMRI experiment 1. A high-resolution T1-weighted three-dimensional anatomical image was collected using an MPRAGE-gradient sequence with 192 x 1 mm slices (in-plane resolution of 1 x 1 x 1 mm; TE: 2.29 ms; TR: 2300 ms; TI:900 ms; FOV: 240 x 240; flip angle: 8°).

fMRI analysis

Preprocessing and data analyses of the imaging data were performed using the SPM12 software (www.fil.ion.ucl.ac.uk/~spm). The preprocessing procedures included spatial realignment of the EPI volumes, co-registration with the structural image, segmentation, group-wise normalisation using DARTEL, and smoothing (see Supplementary Methods for more details).

Region-of-Interest (ROI) Mask

Based on past findings, the striatal reward network is modulated by motivational states and plays an important role in influencing motivation-driven behaviour [5,6,67,68]. We examined bilateral striatal structures including caudate nucleus and NAcc, as well as VTA/SN, and performed in these anatomical masks the small volume correction [69], implemented in SPM. The

masks were anatomically defined using a high-resolution probabilistic atlas of human subcortical

brain nuclei by Pauli and colleagues [70]. Our ROI masks included 565 voxels (3.7 resels) for

caudate nucleus, 53 voxels (0.3 resels) for NAcc, and 66 voxels (0.3 resels) for VTA/SN. Results

were yielded with a height-defining threshold at voxel-level $P < 0.001$ and a cluster extent: k ≥ 5.

Also, to control for the risk of type I error due to multiple comparisons using several anatomical

masks, we focused on results that only survived an adjusted familywise error-corrected

significance threshold of $P < 0.0167$ at cluster level (i.e. 0.05 over 3 comparisons).

As an exploratory attempt to examine any other brain regions outside of the reward

system that are related to motivation-driven decisions, in addition to the ROI approach, we also

analyse the fMRI data with a whole-brain approach. Again, results were obtained using a height-

defining threshold at voxel-level $P < 0.001$, and we focused only on results that survived a

familywise error-corrected significance threshold of $P < 0.05$ at cluster level.

GLM - Activation predicting motivated decision-making

One of the aims of the current study was to test whether curiosity about knowledge and

desire for food influenced a participant's decision in a similar manner via the reward system and

also whether there were similarities and differences in effects depending on the type of stimuli

presented. To this aim, we implemented a general linear model (GLM 1) for each subject that

regressed brain activation depending on the dichotomous decision in the lottery – whether an

individual accepted or rejected it – in each incentive category. Specifically, four separate

regressors were specified for the accepted and rejected trials in curiosity and food categories (i.e.

4 conditions), at the onset of the decision phase of each trial (i.e. when a wheel of fortune

appeared and the participant had to make a choice). To account for the brain activation related to

the elicitation of curiosity and food desirability, another 4 regressors were specified to model the stimulus presentation onset (for magic trick trials specifically, this was time-locked to the moment of subjective surprise defined by the group; see Supplementary Methods for details) of the accepted and rejected trials in the two categories. Moreover, reaction time of the decision response was added for parametric modulation in the four 'decision phase' regressors and head motion parameters were included as regressors of no interest. To test for group-level effects, beta images for the four conditions from each subject in the fMRI experiments were entered as input data for second-level analysis. Using a flexible factorial design in the second-level models (separately for the elicitation phase and the decision phase), we specified the factors of 'Decision' (accept vs. reject), 'Incentive Category' (food vs. curiosity) and 'Experimental Version' (magic vs. trivia), and included each subject's gender and shock expectation rating (see above) as covariates of no interest. Our main focus was the main effect of 'Decision' (i.e. accepted vs. rejected trials) to examine if curiosity and food influenced decision-making in a similar way via the reward system. We also examined the interaction effects to probe any potential variations in neural activity specific to the type of materials used (i.e. stimuli of different incentive categories or different curiosity-triggering materials).

To run a multilevel mediation analysis (explained below), we also created a different linear model (GLM 2) in which each trial was modelled as a separate regressor. These models (implemented separately for the elicitation phase and the decision phase) enabled us to estimate a separate statistical map for each trial (i.e. single-trial activation patterns). We were then able to extract parameter estimates (beta values associated with 'brain activations') for each trial in these maps as inputs for the mediation analysis. As was the case in GLM 1, head motion parameters were included as regressors of no interest.

Parametric modulation analysis - Accounting for the presented outcome probability in

evaluating the relationship between decision and brain activation

To account for the effect of the presented probability of winning/losing on decisions, we implemented a further linear model (GLM 3) and performed parametric modulation analysis. This time, in first-level design, two regressors were specified to model the onsets of the decision phase of trials for the two incentive categories (curiosity, food). To each regressor, the participant's decision in the lottery was added as a parametric modulator (PM) (i.e. 'accept' choice was coded as +1; 'reject' choice as -1); the corresponding outcome probability presented in each trial was added as another PM. In addition, we included reaction time as another PM. The onsets of the elicitation phase (stimulus presentation) for the two incentive categories were specified as separate regressors, along with the head motion parameters as regressors of no interest. Our main focus of interest here was to examine whether the 'accept/reject' decision (i.e. the 'Decision' PM) still parametrically modulated activations in the pre-defined ROIs even after taking the presented outcome probability of into account.

Functional Connectivity Analysis

Functional connectivity was examined with the beta series correlation method [71] implemented in BASCO toolbox [72]. This method allowed us to use trial-by-trial variability to characterise dynamic inter-regional interactions. We specifically tried to explore the functional interaction between brain regions during decision-making and used the left and right caudate nuclei as the ROIs in this connectivity analysis (as the peaked activations at the decision phase were shown within the caudate in the main fMRI analysis; see Results). The anatomical masks used here were defined similarly as in the ROI approach of the main fMRI analysis above. Left

and right caudate was analysed separately because brain connectivity might be strongly

constrained by laterality [73,74]. We corrected for multiple comparisons across the results using

separate left and right caudate seeds with a Bonferroni-adjusted (FWE-corrected) alpha level of

0.025 per analysis.

At the first level of the analysis, a new GLM was constructed (implemented in BASCO),

in which BOLD response time-locked to the onset of the decision phase of each trial was

modelled individually by a separate regressor using a canonical haemodynamic response

function. This resulted in different parameter estimates for each trial for each participant. The six

motion parameters for each run were also included in this GLM. Next, seed-based correlations

were computed voxel-wise for each participant and for each of the experimental conditions of

interest. This procedure generated an individual's correlation map between each seed region's

beta series and the beta series of all other voxels in the brain separately for each condition of

interest, which was normalised using Fisher's r-to-z transformation. At the second level, the set

of correlation maps from each participant was subjected to random-effects analysis to identify

voxels that showed changes in functional connectivity with the seed (based on trial-by-trial

variability in parameter estimates) across different conditions at the group level. Specifically, we

used a flexible factorial design (implemented in SPM) and specified factors of 'Decision' as well

as 'Incentive Category' and 'Experiment Version'. We also included the participant's gender and

his/her shock expectation rating as covariates of no interest. Again, our main focus was the main

effect of 'Decision' (i.e. accepted vs. rejected trials), but we also examined the interaction effects

to probe any potential variations in functional connectivity that may be specific to the type of

materials used (i.e. stimuli of different incentive categories or different curiosity-triggering

materials).

Behavioural analysis

Linear mixed-effects modelling

1. To examine the main question of whether curiosity and food desirability influence

decisions on a trial-by-trial basis at the behavioural level, we performed a generalised linear

mixed-effects modelling (GLME) analysis. In the 'full' model, a participant's decision in the

lottery (i.e. accept or reject) was specified as the dichotomous outcome, in a logistic link

function. Stimulus rating (group-mean centred), the presented shock probability, and the

category of incentive (curiosity vs. food) as well as their interaction terms (including the three-

way interaction), were entered as predictors of the decision. To account for the nested structure

of the data, we specified random intercepts and slopes of participants.

Mixed-effects model with multiple random slopes is prone to convergence errors given

the complexity of the model [75]. We first modelled all the possible random slopes and then

dropped one random slope each time until model convergence (i.e. error-free). The results

reported here were based on the models acquired with this strategy. Importantly, we also carried

out a sensitivity check to make sure that the results were not biased by omitting specific random

slopes [76]. The sensitivity analysis also check for the random effects of stimulus as this effect

could potentially inflate Type-1 error rates [77]. All the reported results were demonstrated to be

very robust to the specification of random effects.

We analysed behavioural data from the initial experiment, the follow-up experiment and

the fMRI experiments. In the GLME for the fMRI experiments, we added the experiment version

(ver 1: magic vs. ver 2: trivia) as an extra fixed-effect term. Note that this 'experiment version'

term was not a significant predictor, indicating little differences in participants' performance (in

terms of acceptance rate) between different versions of the experiment. To facilitate the

understanding of the results further, Supplementary Table 1 present GLME results separately for curiosity and food conditions.

As noted earlier, for the initial behavioural experiment we carried out additional data collection in order to attain more reliable parameter estimates. We recognise this procedure would result in repeated analyses (as data accumulation continued), thereby increasing the potential risk of overestimation of any effects and type 1 error [78]. To control for this risk, we followed the Šidák approach[78,79] to adjust critical P-value for multiple interim analyses conducted (similar to multiple comparisons). Based on the Šidák adjustment [$P' = 1 - (1-P)^{1/k}$], with an ordinary critical P-value being 0.05 and the number of interim analyses k=2 in our case, the adjusted critical value P' would be 0.025. Any analyses using the data from the initial behavioural experiment would use this adjusted critical value as a cut-off in significance testing.

The follow-up 'subjective outcome estimation' experiment had a specific aim to investigate whether curiosity and food desirability distort a participant's subjective estimation of the probability of winning/losing the lottery (a continuous dependent variable). The main analysis in this experiment was based on a linear mixed-effects model (LME) that included the stimulus rating (curiosity/food desirability) and the presented probability of winning/losing as predictors of subjective outcome probability estimation. Again, we specified random intercepts and slopes of participants. We compared two models, one including the stimulus rating and one without, and computed Bayes factors based on the Bayesian Information Criterion (BIC) from each model (readily extracted using 'BIC' function in R) to evaluate the effect of the rating. A measure of BIC quantifies a formal model's goodness of fit to data, taking into account the number of free parameters in the model [80,81]. A lower BIC value indicates a better fit, and vice versa. As described by Wagenmakers[81], Bayes factor (BF) can be estimated using this

transformation (exponentiation of the average of the difference in BIC values for two competing

models, i & j): $BF \simeq \exp[(BIC_i - BIC_j)/2]$ (see also Masson's tutorial [82]). The resulting estimate

of the Bayes factor yields the odds favouring the null hypothesis, relative to the alternative

hypothesis.

Testing of GLME and LME was carried out using the package 'lme4' [83] in R.

Mediation analysis

To test whether the link between rated curiosity/food desirability and the decision to

gamble is mediated by the striatal activations observed in the main fMRI analysis, we performed

a multilevel mediation analysis using multilevel structural equation modelling with Mplus [84]

(version 7). Specifically, we attempted to test the involvement of the activities of the ventral

striatum during the elicitation phase and the dorsal striatum during decision-making in mediating

the link.

In this analysis, we specified the indirect paths as starting from Rating through Striatal

activations at elicitation phase and decision phase to Decision response (i.e. Rating → NAcc

activity at Elicitation → Caudate activity at Decision-Making → Decision response) as well as

the corresponding direct paths, and analysed data of curiosity and food conditions altogether

using trials as the unit of analysis (Fig 4). Incentive category (curiosity vs. food), the presented

shock probability, as well as the experiment version (ver 1: magic vs. ver 2: trivia) were included

as covariates in the model. Moreover, gender of the participant as well as their shock expectation

rating in the post-experiment questionnaire were accounted for in the model as subject-level

covariates. Decision here is a binary outcome variable (accept vs. reject).

To account for the multilevel nature of the data, we used person-mean centring [85] before

parameter estimation and computed cluster robust-standard error to make statistical inference [86].

After estimating the model, the standard error of the mediation effect was computed using the

delta method and statistical significance of the mediation effect was tested.

Statement on statistics and reproducibility

The study employed a within-subject design and participants were not assigned to

different experimental treatments. All statistical tests on behavioural data were two-tailed and

used an alpha level of 0.05 (unless otherwise stated). In the analysis with continuous dependent

variable, residual distribution was assumed to be normal but this was not formally tested. By

making reference to the estimated effect size of a separate pilot study (N = 34), the original

sample size of the initial behavioural experiment (N = 17) was sufficient to detect the estimated

effect at a statistical power of 95%. Additional participants were added later into this initial

experiment in order to achieve more reliable estimates in data modelling and also for better

comparison with other experiments (upon reviewers' suggestion). For the follow-up behavioural

experiment, the sample size (N = 29) was sufficient to detect a small to medium effect (of

curiosity rating on subjective probability estimation) with a statistical power of 80% [87]. The

sample sizes of the fMRI experiments were determined to be comparable to or larger than other

neuroimaging studies of motivational biases [88] and curiosity-based effects [5,6]. Also, such a

sample size should be sufficient to reliably detect medium effect sizes in task-based fMRI

analysis [89].

Code availability

The analyses in this study were performed in standard software and based on published

routine, as specified in detail in the Methods and the Supplementary Methods. The custom codes

are available from the corresponding author on request.

Data availability

The behavioural data that support the findings of the current study will be available in OSF repository. The unthresholded statistical maps of the fMRI results will be made available at NeuroVault.org before publication.

Acknowledgments

The study was supported by the Marie Curie Career Integration Grant (CIG630680 to K.M.), JSPS KAKENHI (15H05401; 16H06406 , 18H01102, and 18K18696 to K.M.), F. J. McGuigan Early Career Investigator Prize (to K.M.), and the Leverhulme Trust (RPG-2016-146 and RL-2016-030 to K.M.). We are grateful of the magicians (including Shota Irieda and Ohkubo Kohei) for producing magic tricks for our research. We thank E. Daveau and C. Ogulmus for helping with data collection, C. Inaltay, A. Firat, A. Haffey, J. Raw, and G. Fastrich for editing and pilot-testing magic video clips, A. Mihalik for providing advice on analysis of data, and C. McNabb for providing useful comments on the drafts of the article.

References

1. Gottlieb, J., Oudeyer, P.-Y., Lopes, M. & Baranes, A. Information-seeking, curiosity, and attention: computational and neural mechanisms. *Trends Cogn. Sci.* **17**, 585–593 (2013).

2. Jirout, J. & Klahr, D. Children's scientific curiosity: In search of an operational definition of an elusive concept. *Dev. Rev.* **32**, 125–160 (2012).

3. Kidd, C. & Hayden, B. Y. The psychology and neuroscience of curiosity. *Neuron* **88**, 449–460 (2015).

4. von Stumm, S., Hell, B. & Chamorro-Premuzic, T. The hungry mind: Intellectual curiosity is the third pillar of academic performance. *Perspect. Psychol. Sci.* **6**, 574–588 (2011).

5. Gruber, M. J., Gelman, B. D. & Ranganath, C. States of curiosity modulate hippocampus-dependent learning via the dopaminergic circuit. *Neuron* **84**, 486–496 (2014).

6. Kang, M. J. *et al.* The wick in the candle of learning: Epistemic curiosity activates reward circuitry and enhances memory. *Psychol. Sci.* **20**, 963–973 (2009).

7. Renninger, K. A. & Hidi, S. *The power of interest for motivation and engagement*. (Routledge, NY, 2016).

8. Sakaki, M., Yagi, A. & Murayama, K. Curiosity in old age: A possible key to achieving adaptive aging. *Neurosci. Biobehav. Rev.* **88**, 106–116 (2018).

9. Rozek, C. S., Svoboda, R. C., Harackiewicz, J. M., Hulleman, C. S. & Hyde, J. S. Utility-value intervention with parents increases students' STEM preparation and career pursuit. *Proc. Natl. Acad. Sci.* **114**, 909–914 (2017).

10. Loewenstein, G. The psychology of curiosity: A review and reinterpretation. *Psychol. Bull.* **116**, 75–98 (1994).

11. Berlyne, D. E. *Conflict, arousal, and curiosity.* (McGraw-Hill Book Company, 1960). doi:10.1037/11164-000

12. Silvia, P. J. *Exploring the Psychology of Interest*. (Oxford University Press, 2006). doi:10.1093/acprof:oso/9780195158557.001.0001

13. Marvin, C. B. & Shohamy, D. Curiosity and reward: Valence predicts choice and information prediction errors enhance learning. *J. Exp. Psychol. Gen.* **145**, 266–272 (2016).

14. Gottlieb, J. & Oudeyer, P.-Y. Towards a neuroscience of active sampling and curiosity. *Nat. Rev. Neurosci.* **19**, 758–770 (2018).

15. Murayama, K. A reward-learning framework of autonomous knowledge acquisition: An integrated account of curiosity, interest, and intrinsic-extrinsic rewards. Preprint at *OSF* (2019). doi:10.31219/osf.io/zey4k

16. Gruber, M. J. & Ranganath, C. How curiosity enhances hippocampus-dependent memory: The Prediction-Appraisal-Curiosity-Exploration (PACE) Framework. Preprint at *OSF* (2019). doi:10.31219/osf.io/5v6nm

17. Kobayashi, K., Ravaioli, S., Baranès, A., Woodford, M. & Gottlieb, J. Diverse motives for human curiosity. *Nat. Hum. Behav.* **3**, 587–595 (2019).

18. Blanchard, T. C., Hayden, B. Y. & Bromberg-Martin, E. S. Orbitofrontal cortex uses distinct codes for different choice attributes in decisions motivated by curiosity. *Neuron* **85**, 602–614 (2015).

19. Daddaoua, N., Lopes, M. & Gottlieb, J. Intrinsically motivated oculomotor exploration guided by uncertainty reduction and conditioned reinforcement in non-human primates. *Sci. Rep.* **6**, 20202 (2016).

20. Vasconcelos, M., Monteiro, T. & Kacelnik, A. Irrational choice and the value of information. *Sci. Rep.* **5**, 13874 (2015).

21. Bennett, D., Bode, S., Brydevall, M., Warren, H. & Murawski, C. Intrinsic valuation of information in decision making under uncertainty. *PLOS Comput. Biol.* **12**, e1005020 (2016).

22. Brydevall, M., Bennett, D., Murawski, C. & Bode, S. The neural encoding of information prediction errors during non-instrumental information seeking. *Sci. Rep.* **8**, 6134 (2018).

23. Eliaz, K. & Schotter, A. Paying for confidence: An experimental study of the demand for non-instrumental information. *Games Econ. Behav.* **70**, 304–324 (2010).

24. Berridge, K. C. Motivation concepts in behavioral neuroscience. *Physiol. Behav.* **81**, 179–209 (2004).

25. Anselme, P. & Robinson, M. J. F. Incentive Motivation: The Missing Piece between Learning and Behavior. in *The Cambridge Handbook of Motivation and Learning* (eds. Renninger, K. A. & Hidi, S.) 163–182 (Cambridge University Press, 2019). doi:10.1017/9781316823279.009

26. Berridge, K. C. From prediction error to incentive salience: mesolimbic computation of reward motivation. *Eur. J. Neurosci.* **35**, 1124–1143 (2012).

27. Robinson, T. E. & Berridge, K. C. The incentive sensitization theory of addiction: some current issues. *Philos. Trans. R. Soc. B Biol. Sci.* **363**, 3137–3146 (2008).

28. Kringelbach, M. L. & Berridge, K. C. Neuroscience of Reward, Motivation, and Drive. in *Recent Developments in Neuroscience Research on Human Motivation* (eds. Kim, S., Reeve, J. & Bong, M.) 23–35 (Emerald Group Publishing Limited, Bingley, 2016). doi:10.1108/S0749-742320160000019020

29. Berridge, K. C. 'Liking' and 'wanting' food rewards: Brain substrates and roles in eating disorders. *Physiol. Behav.* **97**, 537–550 (2009).

30. Tang, D. W., Fellows, L. K., Small, D. M. & Dagher, A. Food and drug cues activate similar brain regions: A meta-analysis of functional MRI studies. *Physiol. Behav.* **106**, 317–324 (2012).

31. O'Doherty, J. P., Deichmann, R., Critchley, H. D. & Dolan, R. J. Neural Responses during Anticipation of a Primary Taste Reward. *Neuron* **33**, 815–826 (2002).

32. Knutson, B., Adams, C. M., Fong, G. W. & Hommer, D. Anticipation of Increasing Monetary Reward Selectively Recruits Nucleus Accumbens. *J. Neurosci.* **21**, RC159–RC159 (2001).

33. Lawrence, N. S., Hinton, E. C., Parkinson, J. A. & Lawrence, A. D. Nucleus accumbens response to food cues predicts subsequent snack consumption in women and increased body mass index in those with reduced self-control. *Neuroimage* **63**, 415–422 (2012).

34. Litman, J. Curiosity and the pleasures of learning: Wanting and liking new information. *Cogn. Emot.* **19**, 793–814 (2005).

35. Gerdeman, G. L., Partridge, J. G., Lupica, C. R. & Lovinger, D. M. It could be habit forming: drugs of abuse and striatal synaptic plasticity. *Trends Neurosci.* **26**, 184–192 (2003).

36. Telzer, E. H. Dopaminergic reward sensitivity can promote adolescent health: A new perspective on the mechanism of ventral striatum activation. *Dev. Cogn. Neurosci.* **17**, 57–67 (2016).

37. Wright, W. F. & Bower, G. H. Mood effects on subjective probability assessment. *Organ. Behav. Hum. Decis. Process.* **52**, 276–291 (1992).

38. Moss, S. A., Irons, M. & Boland, M. The magic of magic: The effect of magic tricks on subsequent engagement with lecture material. *Br. J. Educ. Psychol.* **87**, 32–42 (2017).

39. Ligneul, R., Mermillod, M. & Morisseau, T. From relief to surprise: Dual control of epistemic curiosity in the human brain. *Neuroimage* **181**, 490–500 (2018).

40. Baranes, A., Oudeyer, P.-Y. & Gottlieb, J. Eye movements reveal epistemic curiosity in human observers. *Vision Res.* **117**, 81–90 (2015).

41. Westfall, J., Nichols, T. E. & Yarkoni, T. Fixing the stimulus-as-fixed-effect fallacy in task fMRI. *Wellcome Open Res.* **1**, 23 (2016).

42. Adcock, R. A., Thangavel, A., Whitfield-Gabrieli, S., Knutson, B. & Gabrieli, J. D. E. Reward-Motivated Learning: Mesolimbic Activation Precedes Memory Formation. *Neuron* **50**, 507–517 (2006).

43. Shenhav, A., Cohen, J. D. & Botvinick, M. M. Dorsal anterior cingulate cortex and the value of control. *Nat. Neurosci.* **19**, 1286–1291 (2016).

44. Kolling, N. *et al.* Value, search, persistence and model updating in anterior cingulate cortex. *Nat. Neurosci.* **19**, 1280–1285 (2016).

45. Pochon, J.-B., Riis, J., Sanfey, A. G., Nystrom, L. E. & Cohen, J. D. Functional Imaging of Decision Conflict. *J. Neurosci.* **28**, 3468–3473 (2008).

46. Botvinick, M. M., Cohen, J. D. & Carter, C. S. Conflict monitoring and anterior cingulate cortex: an update. *Trends Cogn. Sci.* **8**, 539–546 (2004).

47. Niv, Y. Reinforcement learning in the brain. *J. Math. Psychol.* **53**, 139–154 (2009).

48. O'Doherty, J. Dissociable Roles of Ventral and Dorsal Striatum in Instrumental Conditioning. *Science.* **304**, 452–454 (2004).

49. Marche, K., Martel, A.-C. & Apicella, P. Differences between Dorsal and Ventral

Striatum in the Sensitivity of Tonically Active Neurons to Rewarding Events. *Front. Syst. Neurosci.* **11**, (2017).

50. Morrison, I., Tipper, S. P., Fenton-Adams, W. L. & Bach, P. "Feeling" others' painful actions: The sensorimotor integration of pain and action information. *Hum. Brain Mapp.* **34**, 1982–1998 (2013).

51. Guo, X. *et al.* Empathic neural responses to others' pain depend on monetary reward. *Soc. Cogn. Affect. Neurosci.* **7**, 535–541 (2012).

52. Whitmarsh, S., Nieuwenhuis, I. L. C., Barendregt, H. P. & Jensen, O. Sensorimotor alpha activity is modulated in response to the observation of pain in others. *Front. Hum. Neurosci.* **5**, (2011).

53. Jepma, M., Verdonschot, R. G., van Steenbergen, H., Rombouts, S. A. R. B. & Nieuwenhuis, S. Neural mechanisms underlying the induction and relief of perceptual curiosity. *Front. Behav. Neurosci.* **6**, (2012).

54. Kruger, J. & Evans, M. The paradox of Alypius and the pursuit of unwanted information. *J. Exp. Soc. Psychol.* **45**, 1173–1179 (2009).

55. Oosterwijk, S. Choosing the negative: A behavioral demonstration of morbid curiosity. *PLoS One* **12**, e0178399 (2017).

56. Hsee, C. K. & Ruan, B. The Pandora effect: The power and peril of curiosity. *Psychol. Sci.* **27**, 659–666 (2016).

57. Oosterwijk, S., Snoek, L., Tekoppele, J., Engelbert, L. & Scholte, H. S. Choosing to view morbid information involves reward circuitry. Preprint at *bioRxiv* (2019). doi:10.1101/795120

58. Noordewier, M. K. & van Dijk, E. Curiosity and time: from not knowing to almost

knowing. *Cogn. Emot.* **31**, 411–421 (2017).

59. Schonberg, T. *et al.* Selective impairment of prediction error signaling in human dorsolateral but not ventral striatum in Parkinson's disease patients: evidence from a model-based fMRI study. *Neuroimage* **49**, 772–781 (2010).

60. Murayama, K., FitzGibbon, L. & Sakaki, M. Process Account of Curiosity and Interest: A Reward-Learning Perspective. *Educ. Psychol. Rev.* (2019). doi:10.1007/s10648-019-09499-9

61. Zink, C. F., Pagnoni, G., Martin-Skurski, M. E., Chappelow, J. C. & Berns, G. S. Human Striatal Responses to Monetary Reward Depend On Saliency. *Neuron* **42**, 509–517 (2004).

62. Krebs, R. M., Boehler, C. N., Roberts, K. C., Song, A. W. & Woldorff, M. G. The Involvement of the Dopaminergic Midbrain and Cortico-Striatal-Thalamic Circuits in the Integration of Reward Prospect and Attentional Task Demands. *Cereb. Cortex* **22**, 607–615 (2012).

63. van Lieshout, L. L. F., Vandenbroucke, A. R. E., Müller, N. C. J., Cools, R. & de Lange, F. P. Induction and Relief of Curiosity Elicit Parietal and Frontal Activity. *J. Neurosci.* **38**, 2579–2588 (2018).

64. Di Domenico, S. I. & Ryan, R. M. The Emerging Neuroscience of Intrinsic Motivation: A New Frontier in Self-Determination Research. *Front. Hum. Neurosci.* **11**, (2017).

65. Fastrich, G. M., Kerr, T., Castel, A. D. & Murayama, K. The role of interest in memory for trivia questions: An investigation with a large-scale database. *Motiv. Sci.* **4**, 227–250 (2018).

66. Peirce, J. W. Generating stimuli for neuroscience using PsychoPy. *Front. Neuroinform.* **2**, (2009).

67. Murayama, K., Matsumoto, M., Izuma, K. & Matsumoto, K. Neural basis of the undermining effect of monetary reward on intrinsic motivation. *Proc. Natl. Acad. Sci.* **107**, 20911–20916 (2010).

68. Murayama, K. *et al.* How self-determined choice facilitates performance: A Key role of the ventromedial prefrontal cortex. *Cereb. Cortex* **25**, 1241–1251 (2015).

69. Worsley, K. J. *et al.* A unified statistical approach for determining significant signals in images of cerebral activation. *Hum. Brain Mapp.* **4**, 58–73 (1996).

70. Pauli, W. M., Nili, A. N. & Tyszka, J. M. A high-resolution probabilistic in vivo atlas of human subcortical brain nuclei. *Sci. Data* **5**, 180063 (2018).

71. Rissman, J., Gazzaley, A. & D'Esposito, M. Measuring functional connectivity during distinct stages of a cognitive task. *Neuroimage* **23**, 752–763 (2004).

72. Göttlich, M., Beyer, F. & Krämer, U. M. BASCO: a toolbox for task-related functional connectivity. *Front. Syst. Neurosci.* **9**, (2015).

73. Tomasi, D. & Volkow, N. D. Laterality Patterns of Brain Functional Connectivity: Gender Effects. *Cereb. Cortex* **22**, 1455–1462 (2012).

74. Wang, D., Buckner, R. L. & Liu, H. Functional Specialization in the Human Brain Estimated By Intrinsic Hemispheric Interaction. *J. Neurosci.* **34**, 12341–12352 (2014).

75. Barr, D. J., Levy, R., Scheepers, C. & Tily, H. J. Random effects structure for confirmatory hypothesis testing: Keep it maximal. *J. Mem. Lang.* **68**, 255–278 (2013).

76. Brauer, M. & Curtin, J. J. Linear mixed-effects models and the analysis of nonindependent data: A unified framework to analyze categorical and continuous independent variables that vary within-subjects and/or within-items. *Psychol. Methods* **23**, 389–411 (2018).

77. Murayama, K., Sakaki, M., Yan, V. X. & Smith, G. M. Type I error inflation in the

traditional by-participant analysis to metamemory accuracy: A generalized mixed-effects model perspective. *J. Exp. Psychol. Learn. Mem. Cogn.* **40**, 1287–1306 (2014).

78. Ludbrook, J. Interim analyses of data as they accumulate in laboratory experimentation. *BMC Med. Res. Methodol.* **3**, 15 (2003).

79. Šidák, Z. Rectangular Confidence Regions for the Means of Multivariate Normal Distributions. *J. Am. Stat. Assoc.* **62**, 626–633 (1967).

80. Stone, M. Comments on Model Selection Criteria of Akaike and Schwarz. *J. R. Stat. Soc. Ser. B* **41**, 276–278 (1979).

81. Wagenmakers, E.-J. A practical solution to the pervasive problems ofp values. *Psychon. Bull. Rev.* **14**, 779–804 (2007).

82. Masson, M. E. J. A tutorial on a practical Bayesian alternative to null-hypothesis significance testing. *Behav. Res. Methods* **43**, 679–690 (2011).

83. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

84. Muthén, L. K. & Muthén, B. O. *Mplus User's Guide. Seventh Edition.* (2012).

85. Enders, C. K. & Tofighi, D. Centering predictor variables in cross-sectional multilevel models: A new look at an old issue. *Psychol. Methods* **12**, 121–138 (2007).

86. McNeish, D., Stapleton, L. M. & Silverman, R. D. On the unnecessary ubiquity of hierarchical linear modeling. *Psychol. Methods* **22**, 114–140 (2017).

87. Arend, M. G. & Schäfer, T. Statistical power in two-level models: A tutorial based on Monte Carlo simulation. *Psychol. Methods* **24**, 1–19 (2019).

88. Leong, Y. C., Hughes, B. L., Wang, Y. & Zaki, J. Neurocomputational mechanisms underlying motivated seeing. *Nat. Hum. Behav.* **3**, 962–973 (2019).

89. Geuter, S., Qi, G., Welsh, R. C., Wager, T. D. & Lindquist, M. A. Effect Size and Power in fMRI Group Analysis. Preprint at *bioRxiv* (2018). doi:10.1101/295048

90. Allen, M., Poggiali, D., Whitaker, K., Marshall, T. R. & Kievit, R. A. Raincloud plots: a multi-platform tool for robust data visualization. *Wellcome Open Res.* **4**, 63 (2019).

  Figure Captions

Fig. 1: Experimental task

In each experimental trial, participants viewed either a video of a magic trick (curiosity

condition) or an image of a food item (food condition). They then gave a rating to indicate their

level of curiosity about the magic trick or the level of desirability of the food. After rating the

stimulus, participants were shown a wheel of fortune (WoF) representing a lottery, which

visualised the probability of them winning/losing in that trial (in this example, as visually

presented in the WoF, there is a higher probability of winning compared with losing).

Participants had to decide whether to gamble or not. Participants were instructed that if they

accepted the gamble and won, they might see the secret behind the magic trick (or get the food,

for food condition) after the experiment as a reward. If they gambled and lost, they would expect

to receive electric shock after the experiment. Participants could also opt to reject the gamble.

Participants understood that, in general, the more gambles they won, the more likely they were to

see more solutions to the tricks and obtain more food items. Similarly, the more gambles they

lost, the more likely they were to receive electric shock for a longer duration. In the modified

1   version of the task for fMRI experiments, (due to time constraints in scanning) the rating phase

2   appeared in only 10% of all the trials during the task. In addition, for the 'trivia version' of the

3   fMRI experiment, participants were presented with trivia questions (instead of magic tricks) as

4   stimuli in the curiosity condition.

Fig. 2: Behavioural results of motivation-driven decision-making.

Both graphs include data of 32 participants from the initial behavioural experiment. a) As

illustrated by the modelled logistic curve, participants tended to reject the gamble more as the

probability of electric shock presented in a wheel of fortune increased, in curiosity (Z=-7.80,

P<0.001, β=-1.28; red curve) and food (Z=-6.29, P<0.001, β=-1.21; blue curve) conditions.

Acceptance rate on y-axis ranges from 0 (100% rejection) to 1 (100% acceptance). Each dot

represents the average acceptance rate of a participant at that level of shock probability presented

in each condition. b) As illustrated in this 'raincloud' plot [90], higher average ratings of curiosity

about magic tricks (Z=7.24, P<0.001, β=1.16) and food desirability (Z=7.49, P<0.001, β=1.15)

were shown for accepted (orange solid circle) compared with rejected (blue solid circle) trials.

Each dot represents the average rating per participant, for each condition. The split-half violin

plots show the probability density of the data at different ratings. Error bars indicate standard

error of the mean. In both graphs, dots are jittered for better visual display. Statistical

significance was assessed with generalised linear mixed-effects models (see 'Behavioural

analysis' in Methods).

Activations within ROIs: ■ CaudN ■ NAcc ■ VTA/SN

Fig. 3: Neural activity in the reward network is modulated by motivation-driven decision-making.

a) Brain activity associated with decision-making was analysed according to whether, in the trials, participants accepted or rejected the gamble to satisfy curiosity/obtain the food item at the risk of electric shock. b) Activity was stronger within (i) NAcc at elicitation phase and within (ii) all the dopaminergic circuit ROIs at decision phase for accepted over rejected trials. For visual illustration here, a voxel-wise threshold of P<0.001 (uncorrected) is applied, and all these clusters survived the ROI analysis with an adjusted FWE-corrected statistical significance of P<0.0167 (at cluster level, correcting for multiple comparisons due to the number of ROIs tested). A, anterior; L, left; P, posterior; R, right.

1
Fig. 4: Mediation path diagram.

Activations of NAcc and caudate nucleus at different phases of decision-making partially

mediated the relationship between the level of curiosity/food desirability (represented by rating)

and the decision to gamble. The mediation effect (paths *a* x *b* x *c*) was significant (Z=2.015,

P=0.044) and so was the direct path *d'* (controlling for mediators). The entire analysis also

accounted for the incentive category of the stimulus (curiosity-inducing or food), the stimulus

version of the experiment (v1 or v2), the presented outcome probability, as well as participants'

characteristics (including gender and subjective shock expectation). Trial-by-trial average striatal

activations were extracted within the anatomical masks of bilateral NAcc at the elicitation phase

and of bilateral caudate nuclei at the decision phase (see Methods: fMRI analysis). Path lines are

labelled with path coefficient (β), Z-value and P-value.

Weaker connectivity with the caudate ROI at decision phase in Accepted (relative to Rejected) trials

y= -1   y= -9   y= -14

R   L

Areas less connected to left caudate ROI

Areas less connected to right caudate ROI

Fig. 5: Functional connectivity of caudate nucleus at decision phase.

Whole-brain analysis of functional connectivity was based on the beta series correlation

approach using anatomical masks of left and right caudate nuclei as the ROIs. A significant main

effect of decision was revealed in the left sensorimotor areas (SMA) near central gyrus,

exhibiting less connectivity to the left and right caudate ROIs respectively when accepted trials

were compared with rejected trials. For visual illustration here, a voxel-wise threshold of

p<0.001 (uncorrected) is applied, and all clusters shown survived an adjusted FWE-corrected

statistical significance of p<0.025 (correcting for multiple comparisons due to the multiple ROIs

applied). L, left; R, right.