1

Error sensitivity and optimization of steady-state kinetic parameters using

multidimensional chemical kinetic analysis

Joseph S. Eskew[1], Christopher G. Connell[2], and Jared C. Cochran[1,]*

[1]Department of Molecular & Cellular Biochemistry, Indiana University, Bloomington, IN 47405,
USA.
[2]Department of Mathematics, Indiana University, Bloomington, IN 47405, USA.
14
15 [*]Corresponding author: Jared C. Cochran, Department of Molecular and Cellular Biochemistry,
16 Indiana University, 405C Simon Hall, 212 S. Hawthorne Dr., Bloomington, IN 47405 USA, Tel:
17 812-855-6935, Fax: 812-856-5710, Email: jcc6@indiana.edu

18
19 **Short Title:** Multidimensional chemical kinetics analysis
20
21 **Abbreviations:** [1] MM, Michaelis-Menten; E, enzyme; S, substrate; P, product; errA, additive
22 error; P_i, inorganic phosphate; C_{est}, estimated constant from nonlinear least squares fitting; C_{exp},
23 the expected constant; v_f, final instantaneous velocity; v_o, initial velocity; CI, confidence interval
24 (95%); AIC, Akaike information criterion; BIC, Bayesian information criterion; MT,
25 microtubule.
26

Abstract

Enzyme behavior has been described using the Michaelis-Menten mechanism. The analysis of extended time domains provides a means to extract the Michaelis-Menten constants through direct fitting of raw data. We have developed a scheme for determining Michaelis-Menten rate constants by appropriate fitting of multidimensional experimental data sets to the closed form of the Michaelis-Menten model. We considered how varying parameters in experimental data affect the accuracy of the remaining parameter estimates. We determine how to improve experimental design to achieve a given accuracy, relative to the amount of intrinsic or external error. We analyze this scheme on data sets built around 20 hypothetical and 2 natural enzymes (kinesin and apyrase) to test error sensitivity in different parameter regimes. Overall, we provide evidence that our data fitting regime will tolerate significant experimental error in the raw data and still converge on the four Michaelis-Menten constants.

Introduction

Enzymes are proteins that catalyze specific biochemical reactions in living cells to generate needed products for metabolic and/or replicative processes. Scientists have been studying enzymes for decades and a wealth of information is available for many different types of enzymes that exist in all domains of life (Eukarya, Bacteria, and Archaea). The Michaelis-Menten (MM)[1] mechanism has been the foundation for studying the steady-state kinetic behavior of enzymatic reactions for over 100 years [1]. In the basic MM scheme (Equation 1), a substrate (S) binds reversibly with a free enzyme (E) to form a non-covalent enzyme-substrate complex, which reacts irreversibly to yield free product (P) and the regenerated free enzyme. One can easily demonstrate the hyperbolic relationship between the initial reaction velocity and substrate concentration through mathematical derivation under a given set of basic assumptions [2]. However, these assumptions are not always valid for enzymatic systems and/or experimental design regimens. This is especially true for extended time domains that allow for both substrate exhaustion and product inhibition of enzyme activity through binding competition. Therefore, more complex mechanisms must be employed to consider the biochemical steps for chemistry (ES$\rightleftharpoons$EP) and product release and re-binding to free enzyme (EP$\rightleftharpoons$E + P).

MM steady-state experiments focus on capturing data for product formation during the initial velocity time regime with variant substrate concentrations. These raw data are subsequently fit to a linear function and the slopes that define the initial reaction velocities are plotted against the substrate concentration to be fit to the hyperbolic MM equation. This process leads to extraction of the $k_{cat}\ and\ K_S$ (also called the K_m) parameter for the enzyme-catalyzed reaction. Due to restrictions that are directly related to the experimental methodologies (e.g. linear range for product detection) or to the nature of the enzyme under investigation (e.g. an enzyme that binds its substrate very tightly but has a very slow maximal rate of substrate turnover), these experiments often do not accurately define the intrinsic rate and equilibrium constants. Also, when these experiments do work as intended, the enzymologist would only yield two of the multiple parameters that govern the overall enzymatic cycle.

A real challenge in obtaining reliable steady state values centers on designing experiments to minimize error and maximize information about the enzyme mechanism. Since

results from steady-state experiments are typically the first insight into the mechanistic character
of an unknown enzyme, having a more thorough analysis gives the enzymologist a better
understanding for designing more complicated kinetic and/or thermodynamic experiments.
Obtaining microscopic rate constants are typically validated by numerous orthogonal chemical
and physical studies, such as isotope and viscosity effects [3-9]. More elaborate single turnover
and transient state kinetic experiments can be employed to define intrinsic rate constants in the
minimal enzyme mechanism [10-15]. Together, data from multiple kinetic approaches can be
globally fit based on numerical integration of the rate equations using a host of kinetic simulation
software packages including KINSIM/FITSIM [16, 17], DYNAFIT [18], COPASI [6, 19],
KinTek Global Explorer [20, 21], and Berkeley Madonna™. Current computation speed has
made these algorithms very effective, especially in one and two dimensions, for recovering best-
fits for rate constants from a given sampling of data without the need to explicitly solve the
underlying model. However, as the number of simultaneously unknown rate constants increases,
corresponding to the dimension of numerical integration, e.g. via the Levenberg–Marquardt
algorithm, the relative speed gain from using explicit model solutions increases. Also by their
very nature, such integration fitting algorithms are unable to easily exploit the specific features
of analytic solutions to achieve the goal of minimizing the data collection needed to obtain
reasonable precision of the rate constants. We propose to improve optimized collection through
selection of data points based on direct, yet fully automated, analysis of features in analytical
solutions of the specific model. This allows us to achieve surprising accuracy in fitting even
three or more rate constants simultaneously from a sparing number of experimental data points.

Since the seminal work of MM [22], reaction kinetics have been monitored beyond the
initial linear phase into a phase where substrate deletion and product accumulation results in
slowing the instantaneous velocity of the reaction, approaching a velocity of zero at infinite time
(or at equilibrium). The analysis of these long time domains through fitting of progress curves
has provided an alternative method for ascertaining the parameters that govern the basic MM
steady-state mechanism [23-27]. Nevertheless, the sensitivity of the fitting regimes has not been
well defined [28], especially along the axes for experimental conditions that can be varied for
any given enzymatic system. We attempt to address these issues with a fitting scheme outlined in
this manuscript. We analyze for accuracy in recovering known rate constants from data in the
presence of various error regimes. Parameter redundancy is elucidated explicitly, allowing us to
both avoid redundant sampling on the one hand, and to test its effects on poorly chosen data as
well. Several statistical comparisons of the efficiency of different experimental designs are
performed. These especially focus on contrasting the effectiveness of finding unknown
parameters starting from different known parameters and how sample size and the dimension of
the data collected affects accuracy.

Materials and Methods

Experimental conditions and proteins used.

The DNA construct coding for the motor domain of human kinesin-5 (Eg5; residues 1–
368) was cloned into a modified pET16b (Novagen) vector for bacterial expression to achieve an
N-terminally 6x His tag and TEV cleavage sequence. (N-terminus: MGSSHHHHHH-
SSGENLYFQGSH-[1]MASQ). Expression and purification of nucleotide-free Eg5 was performed

as previously described [29-32]. Kinetic experiments for Eg5 were performed in assay buffer 1
(20 mM HEPES pH 7.2 with KOH, 5 mM magnesium acetate, 0.1 mM EGTA, 0.1 mM EDTA,
50 mM potassium acetate, 150 mM sucrose, 1 mM DTT) at 298 K as described previously [30,
31, 33]. Microtubules (MTs) were prepared using an aliquot of purified bovine brain tubulin that
was thawed and cycled, and the MTs were stabilized with 20 µM paclitaxel (Sigma-Aldrich).
Kinetic experiments for *Solanum tuberosum* apyrase (Apy; Grade VII; Sigma-Aldrich) were
performed in assay buffer 2 (10 mM HEPES pH 7.2 with KOH, 15 mM $MgCl_2$, 1 mM EGTA,
100 mM NaCl, 150 mM sucrose, 0.02% v/v TWEEN 20) at 298 K with the concentrations
reported as final after mixing. Protein concentrations were determined using the Bradford assay
(Sigma-Aldrich) with bovine serum albumin as standard.

Kinetic assays.

For microtubule-stimulated ATPase measurements, Eg5 at 0.2 µM was complexed with
paclitaxel-stabilized microtubules (MTs) at 2 µM and reacted with varying concentrations of
ATP substrate (95–310 µM) and added ADP product (0 or 1500 µM). Samples of the reaction
(10 µL each) were quenched in 1 N HCl (2.5 µL of 5 N stock) followed by addition of malachite
green reagent (150 µL; room temperature; [34]). After thorough mixing by vortexing, 150 µL of
this mixture was transferred to a 96-well plate for endpoint absorbance (λ=650 nm) using a
BioTek Epoch microplate spectrophotometer. A standard curve from 10–400 µM
Na_2HPO_4/NaH_2PO_4 (pH 7.2) was used to convert A_{650} to inorganic phosphate (P_i) concentration.
For each progress curve, P_i produced by Eg5 ATPase (total P_i minus background P_i) was plotted
against time.

For ADPase measurements, Apy at 0.13 µM was reacted with varying ADP substrate
(40–260 µM) and added AMP product (0 or 10 mM). P_i product was monitored using the
malachite green assay as described above. For each progress curve, P_i produced by Apy ADPase
was plotted against time. Stock ATP and ADP concentrations were experimentally determined
using absorbance (λ=260 nm) with molar extinction coefficient (15,400 L mol^{-1} cm^{-1}).

Mechanisms.

137 Expansions of the Michaelis-Menten (MM) mechanism establish irreversible product
138 formation/release (MM model; Equation 1) as well as competitive product inhibition (PI) of
139 enzyme activity with irreversible chemistry step (PI model; Equation 2) and the fully reversible
140 (FR) three-step mechanism (FR model; Equation 3).

141
$$E + S \underset{k_2}{\overset{k_1}{\rightleftharpoons}} ES \overset{k_3}{\rightarrow} E + P \tag{1}$$

143
$$E + S \underset{k_2}{\overset{k_1}{\rightleftharpoons}} ES \overset{k_3}{\rightarrow} EP \underset{k_6}{\overset{k_5}{\rightleftharpoons}} E + P \tag{2}$$

145
$$E + S \underset{k_2}{\overset{k_1}{\rightleftharpoons}} ES \underset{k_4}{\overset{k_3}{\rightleftharpoons}} EP \underset{k_6}{\overset{k_5}{\rightleftharpoons}} E + P \tag{3}$$

Closed form of the model.

Using *Mathematica* 10 (Wolfram), it was verified by differentiation in time that the differential equation solution takes the closed form of Scheme 1, where ω represents the Lambert ω-function and t is time [35-44].

Scheme 1

$$[P]_t = \alpha - \beta\,\omega\left(\frac{\alpha}{\beta}e^{\frac{\alpha-\gamma t}{\beta}}\right)$$

	Fully Reversible (FR)	Product Inhibition (PI)	Michaelis-Menten (MM)
α	$\dfrac{[S_o]k_{cat}^F K_P - [P_o]k_{cat}^R K_S}{k_{cat}^F K_P + k_{cat}^R K_S}$	$[S_o]$	$[S_o]$
β	$\dfrac{K_P K_S\{[P_o](k_{cat}^R + k_{cat}^F) + [S_o](k_{cat}^R + k_{cat}^F) + k_{cat}^F K_P + k_{cat}^R K_S\}}{(K_P - K_S)(k_{cat}^F K_P + k_{cat}^R K_S)}$	$\dfrac{([P_o] + [S_o] + K_P)K_S}{K_P - K_S}$	K_S
γ	$[E_o]\dfrac{k_{cat}^F K_P + k_{cat}^R K_S}{K_P - K_S}$	$[E_o]\dfrac{k_{cat}^F K_P}{K_P - K_S}$	$[E_o]k_{cat}^F$

Asymmetric confidence estimates.

Since our simulation analyses tested different magnitudes of total degrees of freedom, we utilized mean squared error (MSE) from nonlinear least squared fitting to construct 3D contour plots for asymmetric confidence estimates based on a similar method of FitSpace Explorer [21]. Briefly, pairs of parameters were fixed and varied around their best fit value, and remaining constants were floated in the fitting scheme. The best fit MSE was divided by the MSE from each fixed pair of constants $\left(\text{MSE}_{min}/\text{MSE}_{x,y}\right)$ within a 14x14 matrix centered on the best fit values. Contour plots were constructed in *Mathematica* using these arrays of MSE ratios to linearly interpolate values to provide smooth contours at 0.1 intervals. Bounds of the asymmetric confidence estimates were defined at 11% and 25% increases in MSE $\left(\text{MSE}_{min}/\text{MSE}_{x,y} = 0.9 \text{ (red) and } 0.8 \text{ (orange), respectively}\right)$ as previously reported [21].

Error simulation.

To generate variably 'noisy' data sets that realistically represent a range of experimental observations, normally distributed random additive error (errA) was added to the quantity of product formed during the simulations using Scheme 1 and the following equation:

(4)

$$[P]_t^\sigma = [P]_t + \text{errA}[S_o]\kappa$$

where κ represents the fraction of $[S_o]$ converted to product at equilibrium defined as:

(5)

$$\kappa = \frac{k_{cat}^F K_P}{\left(k_{cat}^F K_P + k_{cat}^R K_S\right)}$$

Time parameter threshold computation.

We used the closed form of the model to determine the near-optimal time parameter interval to maximize the efficiency of the fitting algorithms given a fixed sampling frequency. The most valuable portion of the curve is where the absolute value of the second derivative is bounded away from zero, since the remaining portions are nearly linear where the data points are quite redundant for the nonlinear parameter estimation. Toward this end, we compute the derivative of the product formula (Scheme 1). The derivative $\omega'(t)$ of the Lambert ω-function is:

(6)

$$\omega'(t) = \frac{1}{t\left(\dfrac{1}{\omega(t)} + 1\right)}$$

Hence $[P]'(t)$ simplifies to:

(7)

$$[P]'_t = \gamma - \frac{\gamma}{1 + \omega\left(\dfrac{\alpha}{\beta} e^{\frac{\alpha - \gamma t}{\beta}}\right)}$$

This quantity exponentially decays to 0 as $t \to \infty$. Given this derivative, we chose an appropriate scale to determine when it has become sufficiently small. Since the product $[P]_t$ is almost exactly linear at the beginning of the time regime, we use the scale of the derivative at 0 to set the appropriate scale.

(8)

$$[P]'_{t=0} = \frac{[E_o]\left([S_o]k_{cat}^F K_P - [P_o]k_{cat}^R K_S\right)}{[S_o]K_P + ([P_o] + K_P)K_S}$$

We set the stopping time t_{stop} to be the time when the derivative reaches $[P]'[t_{stop}] = \theta * [P]'_{t=0}$ for a given threshold parameter $\theta = v_f/v_o$ where v_f is the instantaneous final velocity and v_o is the initial velocity as $t \to 0$.

Design for data simulation.

Simulated progress curves were generated through calculating the product concentration (Scheme 1) with random error applied (Eqs. 4 and 5) over time under various conditions. SIM1 represents a data set at increasing $[S_o]$ with no added $[P_o]$ at $t = 0$. SIM2 represents a data set at constant $[S_o]$ with increasing $[P_o]$ that was added at $t = 0$. SIM3 represents a data set at increasing $[S_o]$ and increasing $[P_o]$ that was added at $t = 0$. All simulated data sets were designed for equal total degrees of freedom (i.e. equal total data points with 25 points per curve and 9 curves; Table A.1). All statistical evaluations were standard for NonlinearModelFit in *Mathematica.*

Scoring functions.

Judging the quality of the final nonlinear least squares fitting for our SIM1, SIM2, and SIM3 simulations required a scoring function that incorporated both accuracy (i.e. estimated

206 constant from curve fitting divided by the expected constant) and precision (i.e. symmetric 95%
207 confidence interval). In general, more meaningful fitted constants correlate with higher accuracy
208 (~1) and lower confidence intervals. A scoring function was developed to assess each fitted
209 constant:

210 (9)

$$\text{Score} = e^{-\rho \left| \frac{C_{est}}{C_{exp}} - 1 \right|^{\sigma} - \tau |CI_{95\%}|^{2\sigma}}$$

where C_{est} was the estimated constant from nonlinear least squares fitting, C_{exp} was the
simulated (i.e. expected) constant, and $CI_{95\%}$ was the confidence interval (95%) for each
measurement based on a simple multiple of the standard error. ρ, σ, and τ are weighting
constants chosen to provide a score of 0.5–1 when normalized constants (C_{est}/C_{exp}) are 0.5–1.5
with narrow confidence intervals ($< \pm 50\%$). For all score calculations presented in this
manuscript, $\rho = 1$, $\sigma = 0.75$, and $\tau = 0.5$ (Fig S2*b*).

Results

Analysis of experimental kinetic data for Apy and Eg5

Experimental data for kinesin-5 (Eg5) and apyrase (Apy) were fit to either one of three
mechanistic models, represented by the algebraic rate equations listed in Scheme 1. The three
models, in order of increasing complexity, are designated as "MM", "PI", and "FR",
respectively. Model MM represents the simplest Michaelis-Menten kinetic model, whereby the
reaction product does not rebind and inhibit the enzyme and the reaction proceeds only in the
forward direction (Equation 1). Model PI represents the product inhibition model, whereby the
reaction product rebinds and inhibits the enzyme but the overall reaction can only proceed in the
forward direction (Equation 2). Model FR represents the fully reversible kinetic model, whereby
the reaction product not only rebinds and inhibits the enzyme, but also the overall reaction can
proceed in either the forward or reverse direction (Equation 3).

Apy (type VII; EC 3.6.1.5) catalyzes the hydrolysis ADP to AMP and P_i [45]. To
measure the ADPase kinetics of Apy, we performed a series of reactions at different ADP
substrate (40–260 µM) and different AMP product (0 and 10 mM) concentrations and used the
malachite green colorimetric assay to quantify P_i produced over time. Figs 1*a*–*b* summarize the
results from fitting the Apy ADPase data using basic initial velocity analysis as well as to the
three independent models (Table 1). Using both AIC and BIC to assess the goodness of fit
relative to the complexity of the model, the apyrase data were best modeled to the FR
mechanism (Equation 3) given the lower AIC and BIC values compared to the PI model (Δ
AIC $=- 17$ and ΔBIC $=- 14$). However, asymmetric confidence analysis (Table 2; Fig S1*a*)
indicated that not all the parameters were well constrained. Specifically, we observed large upper
bounds for K_P and k_{cat}^R and no lower bound for k_{cat}^R. These results suggest that AMP product
weakly binds the enzyme yet will proceed in the reverse direction to reform ADP substrate (i.e.
the equilibrium constant for the chemistry step was 7 at the best fit values).

Fig 1. Global fitting of Apy and Eg5 progress curves. (*a*) P_i product formation as a function of
time for Apy (closed symbols) shown at different initial substrate (ADP) concentrations (as
indicated) with no added AMP product at the start of the reactions. (*b*) Time courses of Apy
product formation (open symbols) at different substrate concentrations (as indicated) plus added
initial AMP product (10 mM) at the start of the reactions. (*c*) P_i product formation as a function
of time for Eg5 (closed symbols) shown at different initial substrate (ATP) concentrations (as
indicated) with no added ADP product at the start of the reactions. (*d*) Time courses of Eg5
product formation (open symbols) at different substrate concentrations (as indicated) plus added
initial ADP product (1.5 mM) at the start of the reactions. The smooth lines in each panel
represent the best fit global FR model for Apy and PI model for Eg5 using NonlinearModelFit in
Mathematica with fit constants provided in Table 1.

Eg5 (EC 3.6.4.4) catalyzes the hydrolysis of ATP to ADP and P_i products, which can be
monitored using the malachite green colorimetric assay [34]. We designed a matrix of
microtubule-stimulated Eg5 ATPase reactions with varying ATP substrate (95–310 µM) and
added ADP product (0 and 1500 µM) concentrations for our analysis. Figs 1*c–d* summarize the
results from fitting the microtubule-stimulated Eg5 ATPase data using the three independent
models from above with a direct comparison with previously published results (Table 1) [30, 33].
We found good agreement with the steady-state constants, especially the comparison of the k_{cat}^F
values ($5.4 \pm 0.1 \text{ s}^{-1}$ in previous work [33]; $5.7 \pm 0.4 \text{ s}^{-1}$ here). The estimates for K_S were
consistently 3–8-fold higher than the previously determined MM value, yet the estimate from the
PI model was very similar to the $K_{d,ATP}$ determined from presteady-state acid quench
experiments at 20 ± 7 µM [33]. However, increasing the complexity of the model from PI to FR
led to a decrease in the goodness of fit with higher AIC and BIC values (Δ
AIC = 2 and ΔBIC = 5). These results are consistent with the known chemistry of Eg5, which
shows no evidence of ATP synthesis in the presence of microtubules [33]. Asymmetric
confidence analysis (Table 2; Fig S1*b*) showed no lower bounds for K_S and K_P. Thus, our
analysis provides progress curve fitting to extract the binding constants for substrate and product
as well as the forward catalytic rate constant with minimal data sets (i.e. 6–8 time courses).

Simulated enzyme mechanisms

To demonstrate the ability of our multidimensional chemical kinetics analysis of SIM1,
SIM2, and SIM3 simulations to extract the four Michaelis-Menten (MM) constants from 'noisy'
data, we established 2 enzymes based on the best fit constants of Apy and Eg5 as well as 20
hypothetical enzymes that show a wide range of kinetic mechanisms (E1-E20, Table 3, Fig S2*a*).
These enzyme mechanisms display 1) widely varying equilibrium constants that span
biologically relevant enzymes ($K_{eq}^* = k_{cat}^F K_P / k_{cat}^R K_S = 1.0 - 2.4\text{x}10^4$), 2) a wide variety of
substrate and product binding constants (K_S and K_P, respectively) including various combinations
seen in natural enzymes, 3) a combinatorial spectrum of forward and reverse catalytic constants (
k_{cat}^F and k_{cat}^R, respectively) as well as 4) a range of catalytic efficiencies (k_{cat}^F / K_S) from
extremely inefficient enzymes (Apy = $0.13 \text{ µM}^{-1}\text{s}^{-1}$) to a 'catalytically perfect' enzyme (E1
$= 1003 \text{ µM}^{-1}\text{s}^{-1}$). To generate 'noisy' data sets that realistically represent a range of
experimental observations, normally distributed random additive error was added to the quantity
of product formed during the simulations using Equations 4 and 5 (Fig S3). Given that the four

constants (K_S,K_P,k_{cat}^F,k_{cat}^R) were defined in these simulations, we could assess our nonlinear least
squares fitting for both accuracy and precision using a scoring algorithm (Equation 9), which is a
function of both the normalized constant (defined as the fitted constant divided by the actual
constant) and symmetric 95% confidence interval (Fig S4).

Data set design

For the SIM1 simulations, the concentration of substrate was varied over a range of
concentrations from 0.5*K_S to 5*K_S with no added product ($[P_o]$) at the start of the simulations.
For the SIM2 simulations, the substrate concentration was kept constant at 5*K_S and the
concentration of initial added product ($[P_o]$) at the start of each simulation was varied from 0 to
5*K_P (Table 2). The SIM3 simulations provided a matrix (3x3) using the variation of initial
substrate ($[S_o]$) and initial added product ($[P_o]$) concentrations as described for SIM1 and SIM2
simulations, respectively (see Table S1 for an example of simulation conditions). The calculated
time domain was optimized through the θ ratio of final instantaneous velocity (v_f) to initial
velocity (v_o) to normalize the data point density across each time domain (see Materials and
Methods for details). Each simulation provided a unique experimental data set with a
pseudorandom Gaussian error distribution and these data sets were fit using NonlinearModelFit
in *Mathematica*. For Apy and E1-E20, all four constants (K_S,K_P,k_{cat}^F,k_{cat}^R) were floated during
the analysis. Since Eg5 demonstrated a minimal PI mechanism, only three constants (K_S,K_P,k_{cat}^F)
were floated during the fitting. We assessed the sensitivity of the SIM1, SIM2, and SIM3 designs
against varying additive error for Apy and Eg5 (Fig 2). For Apy, SIM1 was significantly inferior
to the SIM2 and SIM3 designs based on overall fitting scores (Figs 2*a–d*). However, SIM3 was a
significantly better design compared to the SIM2 design (Fig 2*d*), suggesting that varying both
initial substrate and product concentrations provided better overall fitting scores. For Eg5, SIM1
and SIM3 were comparable for K_S and k_{cat}^F estimates (Figs 2*e–g*), but SIM3 provided a
significantly improved mean fitting score for K_P (Fig 2*h*).

**Fig 2. Error sensitivity for Apy and Eg5 progress curves under different simulation
designs.** (*a-c*) Fitting results for Apy simulations with the fitting score for each constant (K_S,
orange; K_P, blue; k_{cat}^F, red; k_{cat}^R, cyan) plotted against error using SIM1, SIM2, and SIM3
conditions, respectively. (*d*) At constant 3% error, the mean fitting score for each constant was
determined under different Apy simulation designs (as indicated). n=200. Error bars represent
the standard deviation of the mean. (*e-g*) Fitting results for Eg5 simulations with the fitting score
for each constant plotted against error using SIM1, SIM2, and SIM3 conditions, respectively. (*h*)
At constant 3% error, the mean fitting score for each constant was determined under different
Eg5 simulation designs (as indicated). n=200. N/A, not applicable.

We tested the sensitivity of all three simulation designs (SIM1, SIM2, SIM3) against
constant additive error for our hypothetical enzymes, E1-E20 (Fig S5). For the hypothetical
enzymes with $K_S \geq K_P$ (E1-E10), SIM2 provided very low (or indistinguishable from zero)
fitting scores for K_S and K_P. SIM3 provided comparable fitting scores for K_S, K_P and k_{cat}^F, but
consistently higher scores for k_{cat}^R. For the hypothetical enzymes with $K_S < K_P$ (E11-E20), SIM1
provided very low (or indistinguishable from zero) fitting scores for K_P and k_{cat}^R. SIM3 provided

consistently higher scores for all four constants compared to SIM2. There was no obvious
correlation to better fitting scores when comparing 1) k_{cat}^F relative to k_{cat}^R, 2) overall equilibrium
constant (K_{eq}^*), or 3) enzymatic efficiency (k_{cat}^F/K_S). Overall, varying both initial substrate and
added product was successful to attain higher fitting scores regardless of the enzyme mechanism.

We assessed the sensitivity of the SIM3 design against varying additive error for our
hypothetical enzymes, E1-E20 (Fig S6). All enzymes showed fitting scores that dropped below
0.5 at >3% error except for E10, which had fitting scores that dropped below 0.5 around 2%
error. Using asymmetric confidence estimate analysis, simulated Apy SIM3 designs were tested
at various additive errors (Fig S7). Since the asymmetric confidence interval was consistently
larger than the symmetric 95% confidence estimates obtained from nonlinear least squares
fitting, it was difficult to directly compare quality scores (Fig S8). Given the wide range of
different hypothetical enzymes tested in this analysis, the SIM3 design (where both initial
substrate and added product is varied) is likely to be successful at estimating the four MM
parameters from fairly 'noisy' data for any enzyme found in natural systems.

Correlations between the estimated constants during the fitting

We next examined the six possible correlations between the four fitted parameters: K_S, K_P
$, k_{cat}^F, k_{cat}^R$ (Fig 3). Using the standard asymptotically unbiased sample correlation matrix, we
computed the estimated correlations as we varied through E1-E20, Apy, and Eg5 with constant
error and fixed starting values. We observed that the correlations remain roughly the same, but
there was some variability as the parameters for the different enzymes vary. The relationship in
the model accounts for most of the nonzero correlation especially considering a given fitting
curve determined only three of the four parameters uniquely as discussed (Equations S18-S29).
The simplicity of the symmetry between K_S and K_P accounts for the near perfect correlation
between these two constants.
353
354 **Fig 3. Parameter correlations for SIM3.** All six parameter correlation coefficients (ρ) are
355 shown for E1-E20, Apy, and Eg5 for the SIM3 design. Error was held constant at 3%. E1-E20
356 and Apy were modeled using the FR mechanism whereas Eg5 was modeled using the PI
357 mechanism.

Time domain length effects on the parameter estimation

We investigated the sensitivity of the SIM3 design for the length of the time domain
based on the ratio of instantaneous velocity from the start of the reaction till the end ($\theta = v_f/v_o$;
Fig 4). We observed similar fitting scores for Apy and Eg5 over a broad range of time domains,
indicating a vast insensitivity of the SIM3 design on the short time domain for data simulation
(Figs 4*a-c* and S9). At θ ratios greater than 0.7, the fitting scores became markedly lower,
indicating that the fitting relies on a significant quantity of product formed during the reaction
(i.e. 60-85% of initial substrate converted to product during the time course). However, if the
time domain was extended beyond the optimal range and thus few data were found in the initial
velocity phase of the curve, the fitting scores steadily decreased for Apy and Eg5 (Figs 4*d-f*). For
the hypothetical enzymes with $K_S \geq K_P$ (E1-E10), fitting scores for K_S and K_P decreased rapidly
as the time domain was lengthened (Fig S10). For the hypothetical enzymes with $K_S < K_P$ (E11-
E20), the overall fitting scores were less sensitive to the longer time domains (Fig S10).

Fig 4. SIM3 fitting scores as a function of time domain. Apy and Eg5 simulations were performed by varying the time domain [q as the ratio of final instantaneous velocity (v_f) to initial velocity (v_o)]. (*a*) Representative Apy simulations with constant error (3%) are shown with increasing q ratios (as indicated). (*b-c*) Fitting scores as a function of the q ratio for Apy and Eg5, respectively. (*d*) Representative plots of Apy simulations at decreasing q ratios. (*e-f*) Fitting scores as a function of decreasing q values with constant error at 3% for Apy and Eg5, respectively.

Data point density effects on parameter estimation

The total degrees of freedom correspond to the length of each progress curve (i.e. 25 data points per curve) multiplied by the number of progress curves. The number of data points per reaction was easy to modify since the total time domain was divided evenly from 2–1024 data points (i.e. changing the time constant for the calculation of $[P]_t$). Figs 5*a-b* highlights the SIM3 results for Apy and Eg5 by varying these conditions. As expected, the fitting scores increased proportionally with the number of data points per time course. These results were not specific to Apy and Eg5, but similar observations were made for all hypothetical enzymes tested (Fig S11). We observed a similar increase of the fitting scores by holding the number of data points per curve constant at 25 and varying the number of progress curves (data not shown). However, using methods to solve the asymmetric confidence estimates, we observed nearly constant fitting score as a function of data points per curve (Fig 5*c*). Therefore, the asymmetric confidence estimates solved from fitting two constants while fixing the values of the remaining two constants provided a robust estimate of the confidence interval independent of the number of total data points in the data set.

Fig 5. SIM3 results as a function of data point density. (*a*) Apy and (*b*) Eg5 simulations were performed by increasing the number of data points per curve (2-1024 points). Fitting scores were plotted as a function of number of data points per curve. (*c*) Asymmetric confidence estimates for K_S in Apy simulations at 0.8 (triangles) and 0.9 (circles) contours. Data plotted with log scale x-axis and were fit to a linear function. Error was held constant at 3%.

Discussion

Different enzyme mechanisms

We have tested the sensitivity of three different experimental designs centered on the closed form (Scheme 1) of the expanded Michaelis-Menten (MM) mechanism for enzyme activity. To simultaneously fit the raw data from individual progress curves to attain the four MM constants ($K_S, K_P, k_{cat}^F, k_{cat}^R$), we fit real experimental data for two enzymes (Apy and Eg5) as well as simulated data for 20 different enzymes (E1–E20). These hypothetical enzymes have a wide range of equilibrium constants, substrate and product binding affinities, catalytic constants, and enzymatic efficiencies (Fig S2*a*, Table 3). Despite all the divergent mechanisms tested in this study, the SIM3 design was robust in accurately and precisely estimating all four MM constants from an experimentally feasible data set.

Scoring function provides standard assessment

Given the complex nature of the data analysis from this study, we developed a simple
exponential scoring function (Equation 9; Fig S2*b*) that yielded a score from 0 to 1 based on
nonlinear least squares fitting accuracy (defined as the fit estimate divided by the actual constant)
and fitting precision (defined as the symmetric 95% confidence interval for each estimate). In
general, more meaningful fitted constants correlate with higher accuracy and lower confidence
intervals. The weighting parameters that govern the scoring function were empirically
determined to provide a fitting score of 0.5–1 when accuracy and precision were less than ±50%.
The same weighting constants were used for the entire analysis, and depending on the nature of
the questions being addressed, these weighting factors can be scaled accordingly to provide more
stringent or more flexible scoring outcomes. The weighting factors employed in this study
provide a reasonably conservative approach to evaluating the fitting results.

Best experimental design of steady-state experiments

Our analysis took advantage of three experimental designs called SIM1 (increasing $[S_o]$
with no added $[P_o]$ at $t = 0$), SIM2 (constant $[S_o]$ with increasing added $[P_o]$ at $t = 0$), and SIM3
(matrix of increasing $[S_o]$ with increasing added $[P_o]$ at $t = 0$). Typically, steady-state
experiments are designed by only varying the substrate concentration as in SIM1. However, we
found that adding initial product in the beginning of the reaction will lead to better parameter
fitting results from both experimental and simulated kinetic data. Indeed, the best fitting regime
happened with varying both substrate and initial product concentrations in the 3x3 matrix (Fig
2). We found that >50% of initial substrate conversion to product was necessary for accurately
defining all four parameters (Fig 4). If experimental error remains constant, having greater than
24 data points per curve (assuming a 3x3 matrix) provided sufficiently accurate parameter
estimates, but more data narrowed the symmetric confidence limits (Fig 5). We also observed
that most hypothetical enzyme mechanisms tolerated a wide range of initial substrate and initial
added product concentrations for achieving accurate parameter estimates (data not shown).
Given the capabilities of *Mathematica*, future investigations aim to extend this analysis to
mechanisms involving two substrates and products during the reaction (i.e. bi-bi mechanisms).

Variable conditions in the design of steady-state experiments

When studying a new uncharacterized enzyme, the decisions made by the enzymologist
for initial substrate concentrations, total enzyme concentration, length of time domains, the
number of data points per time domain, and the number of data sets to collect are difficult.
Interestingly, global fitting under different simulation designs tolerated a wide spectrum of initial
substrate and/or added product concentrations (~2-orders of magnitude around the K_S and K_P
values; data not shown). The total enzyme concentration and the length of the time domain are
directly related, though limits are often encountered based on the solubility limit of the enzyme,
the dissociation of higher order oligomers of the enzyme, the minimum time constant for data
sampling, and/or the stability of the enzyme under the experimental conditions. Therefore, these
caveats make it difficult to achieve the optimal conditions to meet MM steady-state assumptions.
Nevertheless, we found that our SIM3 design converged on all four parameters for all enzymes
tested despite the length of time domain used.

As expected, we observed higher fitting scores when data sampling frequency increased
over the course of the progress curve (Figs 5*a-b* and S11), yet, surprisingly, the scores did not
linearly improve with the added number of data sets in the matrix (data not shown). These results

indicate that an enzymologist can attain accurate and precise information about the kinetic
mechanism with ~10-fold less time and materials. We also found the matrix of initial substrate
and added product concentrations need not be symmetrical for convergence on the four
parameters (data not shown). Even a minimum of 2 initial substrate concentrations and 2 added
product concentrations are sufficient to attain a rough sketch of the behavior of the enzyme,
which will rapidly guide the researcher to subsequent experimental conditions for the next round
of experiments. This approach will provide rapid feedback to appropriately hone on the
experimental conditions that provide the lowest range of confidence intervals and (hopefully)
more accurate estimates of the steady-state parameters.

Conclusions

Results presented here will impact how steady-state experiments will be designed and
how steady-state kinetic data will be analyzed in the future. In the past, MM steady-state
experiments focused on capturing data during the initial velocity time regime with variant
substrate concentrations. These data were subsequently fit to a linear function and the slopes that
define the initial reaction velocities were plotted against the substrate concentration to be fit to
the hyperbolic MM equation to extract the k_{cat} and K_S parameters. Due to restrictions that were
directly related to the experimental methodologies (e.g. linear range for product detection) or to
the nature of the enzyme under investigation (e.g. an enzyme that binds its substrate very tightly
but has a very slow maximal rate of substrate turnover), sometimes these experiments did not
accurately define the intrinsic rate and equilibrium constants. Also, when these experiments did
work as intended, the researcher would only yield two of the multiple parameters that govern the
enzymatic cycle. Therefore, to minimize the time and materials used in the laboratory as well as
maximize mechanistic insight through the multidimensional chemical kinetic analysis, the
development of a modified experimental methodology is required. Combining the variation of
both initial substrate and added product concentrations to establish a matrix of reaction
conditions provides a simple strategy to generate experimental data that can be fit to extract all
four MM parameters from a relatively simple data set. Since results from these experiments are
typically the first insight into the mechanistic character of an unknown enzyme, having a more
thorough analysis gives the enzymologist a better understanding for designing more complicated
kinetic and/or thermodynamic experiments. Although these results promote the advancement of
steady-state kinetic methodologies, data from presteady-state experiments are indispensible.
These advancements in steady-state methodologies can provide insights of the mechanistic
details that are derived through presteady-state methodologies and *vice versa*. Given the ability
of *Mathematica* to fit data with pseudo-infinite dimensionality, these strategies can be applied to
more complex mechanisms with multiple different substrates and products, activators, and small
molecule inhibitors. Also, the progress curve fitting allows for enzyme concentration being a
variable parameter, thus eliminating the need to normalize all reactions to a common enzyme
concentration for comparison purposes. This study sets a foundation for modeling more complex
mechanisms and will provide a new strategy for obtaining accurate and precise measurements of
basic MM parameters.

Acknowledgments

We thank Jeffrey Ewer for preliminary data collection as well as Kayla Bell and Benjamin Walker for insightful comments on the manuscript. We acknowledge financial support from the Simons Foundation (grant #210442 to C.G.C.), the IU Faculty Research Support Program-External Resubmission (J.C.C.) and the National Science Foundation (MCB 1614514 to J.C.C.).

References

1. Michaelis L, Menten ML, Johnson KA, Goody RS. The original Michaelis constant: translation of the 1913 Michaelis-Menten paper. Biochemistry. 2011;50(39):8264-9. Epub 2011/09/06. doi: 10.1021/bi201284u. PubMed PMID: 21888353; PubMed Central PMCID: PMC3381512.

2. Briggs GE, Haldane JB. A Note on the Kinetics of Enzyme Action. Biochem J. 1925;19(2):338-9. Epub 1925/01/01. PubMed PMID: 16743508; PubMed Central PMCID: PMC1259181.

3. Loveridge EJ, Tey LH, Allemann RK. Solvent effects on catalysis by Escherichia coli dihydrofolate reductase. J Am Chem Soc. 2010;132(3):1137-43. doi: 10.1021/ja909353c. PubMed PMID: 20047317.

4. Spies MA, Woodward JJ, Watnik MR, Toney MD. Alanine racemase free energy profiles from global analyses of progress curves. J Am Chem Soc. 2004;126(24):7464-75. doi: 10.1021/ja049579h. PubMed PMID: 15198593.

5. Toney MD, Castro JN, Addington TA. Heavy-enzyme kinetic isotope effects on proton transfer in alanine racemase. J Am Chem Soc. 2013;135(7):2509-11. doi: 10.1021/ja3101243. PubMed PMID: 23373756; PubMed Central PMCID: PMCPMC3579662.

6. Toney MD. Common enzymological experiments allow free energy profile determination. Biochemistry. 2013;52(34):5952-65. doi: 10.1021/bi400696j. PubMed PMID: 23906433; PubMed Central PMCID: PMCPMC4286415.

7. Gadda G, Fitzpatrick PF. Solvent isotope and viscosity effects on the steady-state kinetics of the flavoprotein nitroalkane oxidase. FEBS Lett. 2013;587(17):2785-9. doi: 10.1016/j.febslet.2013.04.021. PubMed PMID: 23660407; PubMed Central PMCID: PMCPMC3916838.

8. Waldauer SA, Stucki-Buchli B, Frey L, Hamm P. Effect of viscogens on the kinetic response of a photoperturbed allosteric protein. J Chem Phys. 2014;141(22):22D514. doi: 10.1063/1.4897975. PubMed PMID: 25494785.

9. Herdendorf TJ, Nelson SW. Catalytic mechanism of bacteriophage T4 Rad50 ATP hydrolysis. Biochemistry. 2014;53(35):5647-60. doi: 10.1021/bi500558d. PubMed PMID: 25137526.

10. Ankrett RJ, Walmsley AR, Bagshaw CR. Kinetic analysis of regulated myosin ATPase activity using single and limited turnover assays. J Cell Sci Suppl. 1991;14:1-5. PubMed PMID: 1832160.

11. Johnson KA. Transient-state kinetic analysis of enzyme reaction pathways. The Enzymes. 20. New York: Academic Press; 1992. p. 1-61.

538   12.   Gilbert SP, Mackey AT. Kinetics: a tool to study molecular motors. Methods.
539   2000;22(4):337-54. Epub 2001/01/03. doi: 10.1006/meth.2000.1086. PubMed PMID:
540   11133240.

541   13.   Francklyn CS, First EA, Perona JJ, Hou YM. Methods for kinetic and thermodynamic
542   analysis of aminoacyl-tRNA synthetases. Methods. 2008;44(2):100-18. doi:
543   10.1016/j.ymeth.2007.09.007. PubMed PMID: 18241792; PubMed Central PMCID:
544   PMCPMC2288706.

545   14.   Johnson KA. A century of enzyme kinetic analysis, 1913 to 2013. FEBS Lett.
546   2013;587(17):2753-66. doi: 10.1016/j.febslet.2013.07.012. PubMed PMID: 23850893.

547   15.   Maxwell BA, Suo Z. Recent insight into the kinetic mechanisms and conformational
548   dynamics of Y-Family DNA polymerases. Biochemistry. 2014;53(17):2804-14. doi:
549   10.1021/bi5000405. PubMed PMID: 24716482; PubMed Central PMCID:
550   PMCPMC4018064.

551   16.   Barshop BA, Wrenn RF, Frieden C. Analysis of numerical methods for computer
552   simulation of kinetic processes: development of KINSIM--a flexible, portable system.
553   Anal Biochem. 1983;130(1):134-45. PubMed PMID: 6688159.

554   17.   Zimmerle CT, Frieden C. Analysis of progress curves by simulations generated by
555   numerical integration. Biochem J. 1989;258(2):381-7. PubMed PMID: 2705989.

556   18.   Kuzmic P. Program DYNAFIT for the analysis of enzyme kinetic data: application to
557   HIV proteinase. Anal Biochem. 1996;237(2):260-73. PubMed PMID: 8660575.

558   19.   Hoops S, Sahle S, Gauges R, Lee C, Pahle J, Simus N, et al. COPASI--a COmplex
559   PAthway SImulator. Bioinformatics. 2006;22(24):3067-74. doi:
560   10.1093/bioinformatics/btl485. PubMed PMID: 17032683.

561   20.   Johnson KA. Fitting enzyme kinetic data with KinTek Global Kinetic Explorer. Methods
562   Enzymol. 2009;467:601-26. doi: 10.1016/S0076-6879(09)67023-3. PubMed PMID:
563   19897109.

564   21.   Johnson KA, Simpson ZB, Blom T. FitSpace explorer: an algorithm to evaluate
565   multidimensional parameter space in fitting kinetic data. Anal Biochem. 2009;387(1):30-
566   41. doi: 10.1016/j.ab.2008.12.025. PubMed PMID: 19168024.

567   22.   Michaelis L, Menten ML. Die kinetik der invertinwirkung. Biochem Z. 1913;49:333-69.

568   23.   Bates DJ, Frieden C. Treatment of enzyme kinetic data. 3. The use of the full time course
569   of a reaction, as examined by computer simulation, in defining enzyme mechanisms. J
570   Biol Chem. 1973;248(22):7878-84. Epub 1973/11/25. PubMed PMID: 4796073.

571   24.   Duggleby RG. Quantitative analysis of the time courses of enzyme-catalyzed reactions.
572   Methods. 2001;24(2):168-74. Epub 2001/06/01. doi: 10.1006/meth.2001.1177. PubMed
573   PMID: 11384191.

574   25.   Philo RD, Selwyn MJ. Use of progress curves to investigate product inhibition in
575   enzyme-catalysed reactions. Application to the soluble mitochondrial adenosine
576   triphosphatase. Biochem J. 1973;135(3):525-30. Epub 1973/11/01. PubMed PMID:
577   4272499; PubMed Central PMCID: PMC1165854.

578   26.   Orsi BA, Tipton KF. Kinetic analysis of progress curves. Methods Enzymol.
579   1979;63:159-83. Epub 1979/01/01. PubMed PMID: 502859.

580   27.   Cao W, De La Cruz EM. Quantitative full time course analysis of nonlinear enzyme
581   cycling kinetics. Sci Rep. 2013;3:2658. doi: 10.1038/srep02658. PubMed PMID:
582   24029878; PubMed Central PMCID: PMC3772379.

583   28.   Gutierrez OA, Danielson UH. Sensitivity analysis and error structure of progress curves.
584   Anal Biochem. 2006;358(1):1-10. Epub 2006/09/19. doi: 10.1016/j.ab.2006.07.007.
585   PubMed PMID: 16979133.

586   29.   Maliga Z, Kapoor TM, Mitchison TJ. Evidence that monastrol is an allosteric inhibitor of
587   the mitotic kinesin Eg5. Chem Biol. 2002;9(9):989-96. Epub 2002/09/27. PubMed
588   PMID: 12323373.

589   30.   Cochran JC, Sontag CA, Maliga Z, Kapoor TM, Correia JJ, Gilbert SP. Mechanistic
590   analysis of the mitotic kinesin Eg5. J Biol Chem. 2004;279(37):38861-70. Epub
591   2004/07/13. doi: 10.1074/jbc.M404203200. PubMed PMID: 15247293; PubMed Central
592   PMCID: PMC1356567.

593   31.   Cochran JC, Gilbert SP. ATPase mechanism of Eg5 in the absence of microtubules:
594   insight into microtubule activation and allosteric inhibition by monastrol. Biochemistry.
595   2005;44(50):16633-48. Epub 2005/12/14. doi: 10.1021/bi051724w. PubMed PMID:
596   16342954; PubMed Central PMCID: PMC2270472.

597   32.   Zhao YC, Kull FJ, Cochran JC. Modulation of the kinesin ATPase cycle by neck linker
598   docking and microtubule binding. J Biol Chem. 2010;285(33):25213-20. Epub
599   2010/06/19. doi: 10.1074/jbc.M110.123067. PubMed PMID: 20558732; PubMed Central
600   PMCID: PMC2919084.

601   33.   Cochran JC, Krzysiak TC, Gilbert SP. Pathway of ATP hydrolysis by monomeric kinesin
602   Eg5. Biochemistry. 2006;45(40):12334-44. Epub 2006/10/04. doi: 10.1021/bi0608562.
603   PubMed PMID: 17014086; PubMed Central PMCID: PMC2288585.

604   34.   Leonard M, Song BD, Ramachandran R, Schmid SL. Robust colorimetric assays for
605   dynamin's basal and stimulated GTPase activities. Methods Enzymol. 2005;404:490-503.
606   PubMed PMID: 16413294.

607   35.   Lambert JH. Observations variae in Mathesin Puram. Acta Helvitica, physico-
608   mathematico-anatomico-botanico-medica. 1758;3:128-68.

609   36.   Euler L. De serie Lambertina Plurimisque eius insignibus proprietatibus. Acta Acad
610   Scient Petropol. 1783;2:29-51.

611   37.   Wright EM. Solution of the equation ze^z=a. Bull Amer Math Soc. 1959;65:89-93.

612   38.   Fritsch FN, Shafer RE, Crowley WP. Algorithm 443: Solution of the transcendental
613   equation we^w=x. Comm ACM. 1973;16(2):123-4.

614   39.   Corless RM, Gonnet GH, Hare DEG, Jeffrey DJ, Knuth DE. On the Lambert W Function.
615   Adv Comput Math. 1996;5:329-59.

616   40.   Schnell S, Mendoza C. Closed form solution for time-dependent enzyme kinetics. J
617   Theor Biol. 1997;187(2):207-12. doi: Doi 10.1006/Jtbi.1997.0425. PubMed PMID:
618   ISI:A1997XL78400007.

619   41.   Goudar CT, Sonnad JR, Duggleby RG. Parameter estimation using a direct solution of
620   the integrated Michaelis-Menten equation. Biochim Biophys Acta. 1999;1429(2):377-83.
621   Epub 1999/02/16. PubMed PMID: 9989222.

622   42.   Goudar CT, Harris SK, McInerney MJ, Suflita JM. Progress curve analysis for enzyme
623   and microbial kinetic reactions using explicit solutions based on the Lambert W function.
624   J   Microbiol   Methods.   2004;59(3):317-26.   Epub   2004/10/19.   doi:
625   10.1016/j.mimet.2004.06.013. PubMed PMID: 15488275.

626   43.   Golicnik M. Explicit analytic approximations for time-dependent solutions of the
627   generalized integrated Michaelis-Menten equation. Anal Biochem. 2011;411(2):303-5.
628   doi: Doi 10.1016/J.Ab.2011.01.016. PubMed PMID: ISI:000288519300020.

629   44.   Schnell S, Mendoza C. Time-dependent closed form solutions for fully competitive
630   enzyme   reactions.   Bull   Math   Biol.   2000;62(2):321-36.   Epub   2000/05/29.   doi:
631   10.1006/bulm.1999.0156. PubMed PMID: 10824432.

632   45.   Kettlun AM, Uribe L, Calvo V, Silva S, Rivera J, Mancilla M, et al. Properties of two
633   apyrases   from   Solanum   tuberosum.   Phytochemistry.   1982;21(3):551-8.   doi:
634   http://dx.doi.org/10.1016/0031-9422(82)83139-7.
635
636

Table 1. Parameter estimates from experimental data for Apy and Eg5.

Apy Constant	Initial Velocity	MM Model Estimate ± SE	CI (95%)	PI Model Estimate ± SE	CI (95%)	FR Model Estimate ± SE	CI (95%)
K_S (µM)	75 ± 12	153 ± 22	110–197	218 ± 24	170–266	176 ± 20	137–214
K_P (µM)	ND [k]	NA	NA	25300 ± 2400	20600–30000	38500 ± 7000	24600–52300
k_{cat}^{F} (s^{-1})	18.7 ± 0.9	18.3 ± 1.5	15.3–21.2	22.7 ± 1.6	19.4–25.9	19.8 ± 1.3	17.2–22.4
k_{cat}^{R} (s^{-1})	NA	NA	NA	NA	NA	2.9 ± 1.1	0.7–5.1
AIC [i]	NA	NA		1620		1603	
BIC [j]	NA	NA		1634		1620	

Eg5 Constant	Initial Velocity [a]	MM Model [b] Estimate ± SE [e]	CI (95%) [f]	PI Model [c] Estimate ± SE	CI (95%)	FR Model [d] Estimate ± SE	CI (95%)
K_S (µM)	7.0 ± 0.4 [g]	58.2 ± 4.6	49–67	23.2 ± 10.3	2.9–43.5	23.1 ± 10.9	2–45
K_P (µM)	117 ± 86 [g]	NA	NA	124 ± 49	28–220	124 ± 49	27–221
k_{cat}^{F} (s^{-1})	5.4 ± 0.1 [g]	6.0 ± 0.1	5.7–6.3	5.6 ± 0.4	4.9–6.3	5.6 ± 0.4	4.8–6.4
k_{cat}^{R} (s^{-1})	NA [h]	NA	NA	NA	NA	1x10^{-13}	0–0.2
AIC	NA	NA		1119		1121	
BIC	NA	NA		1131		1136	

[a] Constants provided by the typical MM analysis of initial velocities. [b] Irreversible MM mechanism defined as in Equation 1. [c] Product inhibition (PI) mechanism defined as in Equation 2. [d] Fully reversible (FR) mechanism as in Equation 3. [e] Nonlinear least-squares fitting estimates with standard error for each fit parameter. [f] CI, confidence interval. [g] Steady-state and equilibrium binding constants obtained under similar experimental conditions [30, 33]. [h] NA, not applicable. [i] AIC, Akaike information criterion. [j] BIC, Bayesian information criterion. [k] ND, not determined.

Table 2. Asymmetric confidence limits for Apy and Eg5 experimental data.

Apy Constant	FR Model Estimate $\pm$ SE	Confidence $\mathrm{MSE}_{min}/\mathrm{MSE}_{x,y} = 0.9$	Confidence $\mathrm{MSE}_{min}/\mathrm{MSE}_{x,y} = 0.8$
K_S (µM)	176 ± 20	$105 - 324$	$83 - 483$
K_P (µM)	$38,500 \pm 7,000$	$19,410 - 238,360$	$15,105 - 345,170$
k_{cat}^{F} (s^{-1})	19.8 ± 1.3	$16.3 - 32.3$	$14.6 - 43.8$
k_{cat}^{R} (s^{-1})	2.9 ± 1.1	$0 - 50$	$0 - 74$
Eg5 Constant	**PI Model Estimate $\pm$ SE**	**Confidence** $\mathrm{MSE}_{min}/\mathrm{MSE}_{x,y} = 0.9$	**Confidence** $\mathrm{MSE}_{min}/\mathrm{MSE}_{x,y} = 0.8$
K_S (µM)	23.2 ± 10.3	$0 - 80$	$0 - 134$
K_P (µM)	124 ± 49	$0 - 334$	$0 - 457$
k_{cat}^{F} (s^{-1})	5.6 ± 0.4	$4.6 - 7.5$	$4.3 - 9.3$

Table 3. Simulated enzyme constants.

	Kin5	Apy	E1	E2	E3	E4
$K_S(\mu M)$	30	150	0.11	0.11	1.1	10.1
$K_P(\mu M)$	120	25000	0.12	0.12	1.2	10.2
$k_{cat}^F(s^{-1})$	6	20	110.3	5.3	5.3	10.3
$k_{cat}^R(s^{-1})$	0.001	3	2.4	0.4	0.4	0.4
K_{eq}^*	24000	1111	50.1	14.5	14.5	26
$k_{cat}^F/K_S(\mu M^{-1}s^{-1})$	0.2	0.13	1003	48.2	4.8	1.0

	E5	E6	E7	E8	E9	E10
$K_S(\mu M)$	0.11	1.1	10.1	100.1	1.1	10.1
$K_P(\mu M)$	0.12	1.2	10.2	100.2	0.62	1.2
$k_{cat}^F(s^{-1})$	20.3	2.3	15.3	100.3	10.3	5.3
$k_{cat}^R(s^{-1})$	20.4	2.4	15.4	100.4	4.4	0.4
K_{eq}^*	1.1	1.0	1.0	1.0	1.3	1.6
$k_{cat}^F/K_S(\mu M^{-1}s^{-1})$	185	2.1	1.5	1.0	9.4	0.5

	E11	E12	E13	E14	E15	E16
$K_S(\mu M)$	0.1	0.1	1.1	10.1	0.1	1.1
$K_P(\mu M)$	1.2	1.2	10.2	100.2	1.2	10.2
$k_{cat}^F(s^{-1})$	10.3	5.3	5.3	10.3	2.3	2.3
$k_{cat}^R(s^{-1})$	4.4	0.4	0.4	3.4	2.4	2.4
K_{eq}^*	28.1	159	123	30	11.5	8.9
$k_{cat}^F/K_S(\mu M^{-1}s^{-1})$	103	53	4.8	1.0	23	2.1

	E17	E18	E19	E20		
$K_S(\mu M)$	1.1	10.1	0.1	1.1		
$K_P(\mu M)$	100.2	100.2	1.2	100.2		
$k_{cat}^F(s^{-1})$	2.3	100.3	1.3	1.3		
$k_{cat}^R(s^{-1})$	2.4	100.4	5.4	10.4		
K_{eq}^*	87	9.9	2.9	11.4		
$k_{cat}^F/K_S(\mu M^{-1}s^{-1})$	2.1	9.9	13	1.2		

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5