Electrochemotherapy with bleomycin associated with doxorubicin induces tumor regression

and decreases the proliferative index in canine cutaneous squamous cell carcinomas

Electrochemotherapy in canine cutaneous squamous cell carcinomas

Denner S. Anjos[1#a], Cynthia Bueno[2], Larissa F. Magalhães[3], Georgia M. Magalhães[4], Ewaldo Mattos-

Junior[1], Marcela M.R. Pinto[5], Andrigo B. De Nardi[2], Carlos H.M. Brunner[6], Antonio F. Leis-Filho[7],

Carlos E. Fonseca-Alves[7¶], Sabryna G. Calazans[1#a¶]*

1. Veterinary Science Graduate Program, University of Franca (UNIFRAN), Franca, Brazil

2. Department of Veterinary Clinic and Surgery, São Paulo State University (UNESP), Jaboticabal, Brazil

3. Department of Veterinary Pathology, University of Franca (UNIFRAN), Franca, Brazil

4. Federal Institute of Education, Science and Technology of the South of Minas Gerais - Muzambinho, Minas

Gerais, Brazil

5. Pathologist veterinarian at the CEVEPAT Laboratory, Botucatu, SP

6. Department of Veterinary Clinic of University Paulista, São Paulo, Brazil.

7. Department of Veterinary Clinic, School of Veterinary Medicine and Animal Science, São Paulo State

University (UNESP), Botucatu, São Paulo, Brazil

[#a]Current address: Department of Veterinary Clinic and Surgery, São Paulo State University (UNESP),

Jaboticabal, Brazil

*Corresponding author:

Email: carlos.e.alves@unesp.br (CEFA)

[¶]These authors contributed equally to this work.

ABSTRACT

Canine cutaneous squamous cell carcinoma (cSCC) is the most common skin cancer in dogs, and due to its low metastatic rate, local treatments such as electrochemotherapy (ECT) promote disease control or even complete remission and increase the survival time in most cases. This study aimed to evaluate the expression of BAX, Bcl-2, and Ki67 and clinical parameters in dogs with cSCC subjected to ECT. A prospective clinical nonrandomized study was performed in dogs with naturally occurring cSCC treated with ECT. Eighteen lesions (from 11 dogs) were selected, independent of breed, sex and age. The ECT protocol consisted of bleomycin plus doxorubicin followed by electric pulses characterized by 8 biphasic electric pulses lasting 100 ms, 1 Hz and 1000 V/cm. Among the 18 lesions, the lesion volume significantly decreased after treatment (p=0.04). The tumor size at D0 had no impact on survival time or prognosis (P>0.05). A decreased mitotic index was observed at compared with D0 (P=0.019). We also observed more intratumoral necrosis at D21 compared to D0 (P=0.041). The median expression level of Ki67 was 277.96 at D0 and 193.92 at D21. Thus, tumor samples had a lower proliferative index after ECT (D21) (P=0.031). The survival times of subjects with Ki67 values lower and higher than the Ki67 median value were not significantly different (P>0.05). Regarding apoptotic markers, there was no significant difference in BAX or Bcl-2 expression between D0 and D21 (P>0.05) or in overall survival between subjects with different levels of apoptotic markers. Furthermore, a positive correlation was observed between BAX and Bcl-2 before ECT (D0) (P=0.0379, r=0.5067). In conclusion, there was no change in BAX and Bcl-2 protein expression levels in response to ECT at the time points evaluated, and ECT was able to reduce tumor volume and cellular proliferation in cSCC.

Key-words: Apoptotic markers, cutaneous carcinoma; electroporation, proliferative index, tumor size.

Introduction

Canine cutaneous squamous cell carcinoma (cSCC) is the most common skin cancer in

dogs, and the most important etiological factor is chronic sunlight exposure. On the other

hand, canine papillomavirus infection, chronic inflammation and immunosuppression might

be involved in the development of cSCC [1-4]. cSCC is the second most common cutaneous

tumor in dogs [5], with an incidence between 3-20% [6,7]; however, the incidence of cSCC

depends on the geographic area [5]. In humans and dogs, actinic keratosis is an SCC precursor

lesion with more than 80% of the SCC cases in humans being derived from previous actinic

keratosis [8, 9].

Generally, cSCCs are locally invasive tumors with a low metastatic rate (13%) [10-

13], with is similar to the behavior of SCCs humans, in whom the metastatic rate is 5%;

metastasis mainly occurs in locoregional lymph nodes [14-16]. Due to this low risk of

metastasis, local treatments such as surgery, cryotherapy, radiotherapy, photodynamic therapy

and electrochemotherapy (ECT) promote disease control and increase the survival time in

most cases [1,17].

ECT is able to induce an inflammatory response, necrosis, scar tissue and apoptosis in

tumors [18-20]. This process is a genetically programmed process for the elimination of

damaged cells, and its onset is controlled by numerous interrelating processes that are

influenced by extrinsic and intrinsic signals that converge in an effector pathway. Alterations

in these pathways are an important in the process of tumorigenesis, leading to the persistence

of neoplastic cells and the promotion of progression and metastasis [21,22].

BAX (Bcl-2 associated X protein) and Bcl-2 are important proteins in the BCL-2

family, which consists of pro- and anti-apoptotic proteins. The BAX/Bcl-2 cross-regulation

controls apoptosis, cell survival or cellular proliferation [23,24]. Several studies have

determined the expression level of a single apoptosis-associated protein, such as BAX and/or BCL-2, by immunohistochemistry or flow cytometry and have correlated its expression with the prognosis of mammary tumors and lymphoma [25,26]. However, conflicting data have been observed among the studies [25-27]. In feline cutaneous SCCs, neither BAX nor BCL-2 expression was detected. However, in basal cell tumors, BCL-2 expression was higher (23/24) than that of BAX, which was expressed only in seven out of 24 tumors. For the tumors that expressed both BAX and BCL-2, the BAX:BCL-2 ratio was low [28].

Another important factor to be evaluated in cSCC is cellular proliferation, which is measured by the level of Ki67. It is a nuclear protein that is used to detect proliferative cells because it is present in proliferating cells during the late G1, S, G2 and M phases of the cell cycle, and it is correlated with poor prognosis in dogs [29-32]. It is known that the high expression level of Ki67 is correlated with a poorer prognosis in dogs with several tumor types as well as in humans, such as mammary tumors, mast cell tumors, perianal tumors, oral and cSCCs [29-37].

ECT has gained popularity in recent years in veterinary medicine as well as in human medicine [38,39]. ECT is a combination of chemotherapy and the localized delivery of electric pulses to the tumor nodule. ECT aims to increase antineoplastic drug diffusion into the cell after cell membrane electroporation, thus increasing its cytotoxicity [17].

To the best of the authors' knowledge, this is the first study to evaluate the expression of BAX, Bcl-2, and Ki67 in dogs with cSCC that underwent ECT. This study aimed to evaluate the clinical parameters, proliferative index, and expression levels of BAX and Bcl-2 in dogs with cSCC that underwent ECT.

Materials and Methods

Ethical approval

This study was performed in accordance with the National and International Recommendations for the Care and Use of Animals. All procedures were performed after receiving the approval of the Institutional Animal Ethics Committee (CEUA/UNIFRAN, #033/15).

Study design

A prospective clinical nonrandomized study was performed in dogs that presented to the Veterinary Teaching Hospital of Sao Paulo State University (UNESP) and the Veterinary Teaching Hospital of University of Franca (UNIFRAN) with naturally occurring cSCC and that were treated with ECT. Consent was obtained from the dogs' owners to perform the treatment. The recommendations made by Campana et al. [40] for reporting clinical studies on ECT were followed.

Inclusion criteria and animal selection

All dogs enrolled in this study fulfilled the following characteristics: a histopathologically confirmed diagnosis of stage T1 cSCC (according to the World Health Organization) [41]; the absence of distant metastases; the compliance of the owner with follow-up after 21 days; the owner's permission to perform biopsies before (D0) and after ECT (D21).

127   Eighteen lesions (from 11 dogs) were selected, independent of breed, sex and age. The

128   eligibility inclusion criteria were as follows: dogs with clinically staged cSCC and complete

129   physical examinations, laboratory exams, and fine needle aspiration of regional lymph nodes.

130   Three-view thoracic radiography and abdominal ultrasonography were also performed.

Electrochemotherapy protocol

Bleomycin (Cinaleo®-Meizler, Barueri-SP) was diluted in 5 mL of saline solution and

administered intravenously (IV) at a dose of 15,000 UI/m^2. Five minutes after the

administration of the bleomycin, doxorubicin (Cloridrato de Doxorrubicina® - Eurofarma

Laboratórios S.A. Ribeirao Preto – SP) was diluted in 25 mL of saline solution and

administered intravenously at a dose of 30 mg/m², followed by sequences of 8 biphasic

electric pulses lasting 100 ms each, with a frequency of 1 Hz, 1000 V, generated by an LC

BK-100 portable electroporator and delivered by six needle electrodes with 0.3 mm distance

between them; the needle electrodes were arranged in rows (parallel array) until they achieved

complete coverage of the tumor. The procedure lasted until 28 minutes, according to a

previous report [42,43].

All ECT procedures were administered under general anesthesia using propofol (5

mg/kg) IV followed by endotracheal intubation; anesthesia was maintained with isoflurane.

All animals received postoperative analgesia, including the IV injection of meloxicam (0.2

mg/kg) and tramadol (2 mg/kg).

Tumor response

The total volume of the neoplasm was calculated by the following formula:

$\pi \; x \; length \; x \; width \; x \; height/6$; tumor volume was measured at D0 and D21 using a digital

pachymeter [29,30]. The same evaluator measured tumor volume to exclude measurement

bias. Complete remission (CR) was defined as a total reduction in measured tumor volume,

while partial remission (PR) was defined as ≥ 30% reduction in tumor volume. Progressive

disease (PD) was defined as ≥ 20% increase in tumor volume or new lesions, and stable

disease (SD) was defined as ☐ 30% reduction in tumor volume or ☐ 20% increase in tumor

volume [43]. CR and RP were considered "favorable" responses, while PD and SD were

considered "unfavorable" responses.

Histopathological evaluation

The first sample was collected immediately before the first session of ECT (D0). The

second sample was obtained on day 21 (D21) after ECT. A 6-mm punch biopsy was used to

obtain the tumor samples. All samples were immediately placed in 10% formalin for 24

hours, followed by 70% alcohol until paraffinized sections were prepared. All tumor samples

were classified according to Gross et al. [5], as follows: 1) well-differentiated SCC,

presenting centralized accumulatio of compact laminated keratin or keratin pearls, with

keratinization progressing through the granular cell layer as in the normal epidermis and the

keratinized centers of the lobules undergoing necrosis and becoming infiltrated by neutrophils

or 2) poorly differentiated SCC presenting smaller epithelial structures, cords and nests rather

than large islands of squamous epithelial cells, moderate to high mitotic activity, and no

keratin pearls.

Immunohistochemistry

Immunohistochemical staining for BAX, BCL-2 and Ki67 antibodies was performed

in all cSCCs in the original biopsy samples (D0) and in the biopsy specimens collected 21

days after the first ECT (D21). Immunohistochemical staining was performed using the

peroxidase method and 3,3' diaminobenzidine tetrachloride (DAB). The slides were dewaxed

in xylol and rehydrated in graded ethanol. For antigen retrieval, the slides were incubated in

citrate buffer (pH 6.0) in a pressure cooker (Pascal, Agilent Technologies, Santa Clara, CA,

USA). The endogenous peroxidase was blocked with a commercial solution (Protein Block,

Agilent Technologies, Santa Clara, CA, USA), and the samples were incubated with primary

antibodies overnight at 4ºC. Anti-BAX (mouse monoclonal, Santa Cruz Biotechnology,

Dallas, TX, USA) was detected with a monoclonal antibody at a 1:200 dilution overnight.

Bcl-2 (mouse monoclonal, Santa Cruz Biotechnology, Dallas, TX, USA) was a monoclonal

mouse antibody used at a 1:400 dilution overnight, and anti-Ki67 (MIB-1, Agilent

Technologies, Santa Clara, CA, USA) with a monoclonal mouse antibody used at a 1:50

dilution, applied overnight. After incubation with the above antibodies, the slides were placed

on an autostainer platform (Agilent Technologies, Santa Clara, CA, USA). Then, the sections

were counterstained with Harris´s hematoxylin, dehydrated, and mounted. The anti-Ki67

antibody cross-reactivity with canine tissue was provided by the manufacturer in the antibody

datasheet. For the anti-Bcl-2 and anti-BAX antibodies, we performed western blotting to

show the antibody reactivity.

Negative controls were run for all antibodies with a mouse universal negative control

(Dako, Carpinteria, CA, USA) according to the manufacturer's instructions. The positive

control consisted of normal lymph nodes for all antibodies according to Protein Atlas

guidelines (www.proteinatlas.org).

Immunohistochemical evaluation

The immunohistochemical slides were examined with a light microscope using an

ocular grid 26-mm in diameter (Leica Microscopy DMLB, HC, PLAN 10x/20, 4"x5") at

400X magnification. The immunoexpression levels of BAX, Bcl-2 and Ki67 were established

by counting the number of stained cells and considering the number of positive cells in

relation to the total number of cells inside the ocular grid per five random high-power fields

(400x). The samples were classified as 0 (absence of stained cells), 1 (< 10% stained cells), 2

(10-25% stained cells), 3 (26-50% stained cells) and 4 (> 50% stained cells) according to

Fonseca-Alves et al. [44]. For the evaluation of the scores, cytoplasmic staining for BAX and

Bcl-2 and nuclear staining for Ki67 were considered.

210 Western blotting

The western blot analysis was performed to validate the protein cross-reactivity with

canine tissue. The procedures were performed according to a previously published method

[45]. Briefly, two cSCC tissue biopsies were collected and homogenized (Polytron

homogenizer, Kinematica, Lucerne, Switzerland) for 30 seconds at 4°C in RIPA Buffer

(Millipore, Burlington, MA, USA). Total protein was extracted and quantified as described by

Bradford [46]. A total of 70 μg of protein was subjected to electrophoresis and then

transferred to nitrocellulose membranes (Sigma Chemical Co., St. Louis, MO). The blots

were blocked with 3% bovine serum albumin in TBS-T for 2 hours and incubated overnight

with BAX (mouse monoclonal, Santa Cruz Biotechnology, Dallas, TX, USA) and Bcl-2

(mouse monoclonal, Santa Cruz Biotechnology, Dallas, TX, USA) antibodies. Goat anti-β-

actin antibody (1:1,000; sc-1615, Santa Cruz Biotechnology, Santa Cruz, CA, USA) was used

as a control. A horseradish peroxidase-conjugated secondary antibody was incubated with the

blots for 1 hour, and the blots were visualized via chemiluminescence (Amersham ECL Select

Western Blotting Detection Reagent, GE Healthcare).

226 Statistical analysis

For statistical purposes, we calculated the median value of each clinical,

histopathological and immunohistochemical parameter and classified the value as "low" when

it was lower than median and "high" when the value was greater than the median. Then, we

compared the survival between dogs with low and high expression values of each parameter

through the Kaplan-Myer method. To evaluate the association between the tumor stage ($\leq$T2

x >T2) before treatment and after ECT, Fisher's exact test was applied. Furthermore,

Spearman's correlations between the expression levels of BAX, Bcl-2 and Ki67 were

performed to compare the levels at two time points, namely, before and after ECT.

Commercial software (GraphPad Prism® 6.0 – GraphPad Software, Inc. 2015) was used for

the statistical analysis. P values <0.05 were considered significant.

Results

Clinical data

Mixed breed dogs (n=5) were the dogs most commonly affected by cSCC, followed

by American Pit Bulls (n=3), Boxers (n=2) and English Pointers (n=1). All subjects had

sparse fur and lightly pigmented skin. The mean age of the animals was 7.5 ($\pm$ 2.29) years,

with females being the more common sex (n=10). Five animals had more than one lesion.

Thus, we investigated 18 lesions in 11 subjects.

The most commonly affected sites observed were the abdominal region (n=10),

followed by the thoracic region (n=5), axillar region (n=1), preputial region (n=1) and tibial

region (n=1). Among the 11 subjects, only three had regional lymph node involvement at the

time of diagnosis. There was no difference in survival time between subjects with and without

lymph node involvement (S Fig 1A) (P> 0.05).

Suppl Fig. 1. Survival analysis according to lymph node involvement, tumor size, and

apoptotic markers in dogs with cSCC. (A) There was no difference in survival time

between subjects with and without lymph node involvement ($P > 0.05$).

Among the 18 lesions, the volume before the treatment ranged from 0.14 to 112.9

cm³ (median of 4.64), and the volume decreased significantly after treatment (p=0.04),

ranging from 0.11 to 118.2 cm³ (median of 1.49) (Table 1) (Fig 1A).

Table 1. Records from the subjects that underwent ECT regarding the clinical data, tumor localization, stage and tumor size.

ID	Breed	Age	Sex	Number of tumors	Tumor localization	Histopathological evaluation	TNM	Tumor size at D0 (cm³)	Tumor size at D21 (cm³)	Outcome	Number of sessions
1	Pitbull	8	M	2 (a, b)	Axillar (a) Prepucial (b)	poorly differentiated well differentiated	T2N0M0 T2N0M0	3.75 19.23	0.91 6.44	PR PR	3 / S
2	Pitbull	9	F	1	Abdominal	well differentiated	T2N0M0	5.04	0.73	PR	2
3	Boxer	2	F	1	Thorax lateral	well differentiated	T3N0M0	68.49	65.94	SD	2 / S
4	Mixed	10	F	2 (a. b)	Thorax lateral (a) Abdominal (b)	poorly differentiated well differentiated	T2N0M0 T3N0M0	2.3 14.15	0.278 1.68	PR PR	1 / S
5	Mixed	7	F	1	Abdominal	well differentiated	T3N0M0	93.57	118.24	PD	2
6	Pitbull	9	F	4 (a, b, c, d)	Thorax lateral (a) Abdominal (b) Abdominal (c) Abdominal (d)	well differentiated poorly differentiated well differentiated well differentiated	T1N0M0 T3N0M0 T1N0M0 T2N0M0	1.14 112.92 0.81 22.36	0.175 48.2 0.533 5.014	PR PR PR PR	2 / S
7	Mixed	NA	F	1	Abdominal	well differentiated	T2N1M0	104.6	104.6	SD	2
8	Mixed	NA	F	2 (a. b)	Abdominal (a) Thorax lateral (b)	well differentiated well differentiated	T3N0M0 T2N0M0	4.24 0.22	10.36 1.046	PD PD	2
9	English Pointers	8	F	1	Abdominal	well differentiated	T3N1M0	76.93	30.03	PR	2 / S
10	Boxer	8	F	2 (a. b)	Thorax lateral (a) Tibial (b)	well differentiated well differentiated	T1N0M0 T1N0M0	0.314 0.141	0.113 0.28	PR PD	2
11	Mixed	7	F	1	Abdominal	well differentiated	T4N1M0	1.2	1.31	SD	1

259
M, male; F, female
PR: partial remission; SD: stable disease; PD: progressive disease
S: surgery
NR: death not related to cancer
NF: no follow-up
NA: not available

Fig. 1. Evaluation of tumor volume, clinical response according to tumor volume, mitotic

index, intratumoral necrosis and Ki67 expression before and after treatment with ECT

in dogs with cSCC. (A) The volume before the treatment ranged from 0.14 to 112.9 cm³

(median of 4.64), and the volume significantly decreased after treatment (p=0.04), ranging

from 0.11 to 118.2 cm³ (median of 1.49).

Based on the volume measurements, 11 (61.1%) lesions underwent PR, 4 (22.2%)

underwent PD, and 3 (16.6%) underwent SD. We evaluated the tumor size at D0 as a

prognostic factor according to the TNM recommendation for human SCCs, and we grouped

the subjects according to whether they had tumors smaller or larger than 5 cm^3. The volume at

D0 had no impact on survival time (Fig 1B) and had no prognostic value (P>0.05).

Fig. 1. Evaluation of tumor volume, clinical response by tumor volume, mitotic index,

intratumoral necrosis and Ki67 expression before and after treatment by ECT in dogs

with cSCC. (B) Evaluation of tumor size at D0 as prognostic factor in dogs with tumors

smaller and larger than 5 cm^3. The volume at D0 had no impact on survival time and had no

prognostic value (P>0.05), and the response to ECT was not significant (p=0.332).

In addition, the association between the clinical stage before treatment (D0) and the

response to ECT was not significant (p=0.332) (Fig 1B). The median survival time of the 11

dogs was 180 days (32 to 570 days) (Fig 2A).

Fig. 2. Overall median survival time of the 11 dogs and survival time based on mitotic

index. (A) The median survival time of the 11 dogs was 180 days (32 to 570 days).

The additional clinical results are shown in Table 1. The number of sessions that each

subject underwent ranged from one to three. Five subjects underwent a surgical procedure

after D21 (four due to PD and one due to the owner request).

Histopathological features

Regarding the histopathological grade, 16 lesions (83.3%) were classified as well-

differentiated SCC (Suppl Fig 2A), and three (16.6%) were classified as poorly differentiated

SCC (Suppl Fig 2B). Interestingly, only well-differentiated tumors progressed, and all (3/18)

poorly differentiated cSCC experienced PR.

The median mitotic index in the tumor group at D0 was 4.94 (0 to 34). We found a

relationship between the mitotic index and overall survival. Subjects with mitosis numbers

lower than 4.9 at D0 experienced a longer survival time (P=0.009) (Fig 2B).

Suppl Fig 2. Histopathological evaluation of squamous cell carcinoma (SCC) in dogs

that underwent electrochemotherapy (ECT). (A) Well-differentiated SCC with the

presence of mononuclear inflammatory infiltrate before ECT. (B) SCC before ECT. It is

possible to see tissue disorganization with the presence of apoptotic cells showing

basophilic nuclei.

Fig. 2. Overall median survival time of the 11 dogs, and the survival time based on the

mitotic index. (B) Subjects with mitotic index values lower than 4.9 at D0 experienced a

longer survival time than subjects with mitotic index values greater than 4.9 (P=0.009).

Additionally, we found decreased mitotic index values at D21 compared to D0

(median 1.5, ranging from 0 to 20) (P=0.019) (Fig 1C). We also assessed intratumoral

necrosis in the tumor samples at D0 and D21. It is not surprising that we found more necrosis

in tumors at D21 than at D0 (P=0.041) (Fig 1D). We evaluated necrosis at D0 and the survival

time, and we did not find significant differences (P>0.05).

Fig. 1. Evaluation of tumor volume, clinical response by tumor volume, mitotic index,

intratumoral necrosis and Ki67 expression before and after treatment by ECT in dogs

with cSCC. (C) A decreased mitotic index at D21 (median 1.5, ranging 0 to 20) (P=0.019)

was observed. (D) Greater intratumoral necrosis was observed in tumor samples at D21 compared

to those at D0 (P=0.041).

Protein expression

Immunohistochemical expression was evaluated in 15 tumors (15/18) at two different

time points (D0 and D21). Three out of 18 samples were excluded due to the absence of

neoplasticism in tumor samples from D0 or D21. The median expression of Ki67 in D0 was

277.96 (113.4 to 511.4), and D21 was 193.92 (15 to 494). We found decreased proliferative

expression between D0 and D21 (Fig 3 A and B). Thus, tumor samples after ECT (D21) had a

lower proliferative index than those before ECT (p=0.031) (Figure 1E). Using the

proliferative index at D0, we evaluated the survival time between subjects with Ki67 values

lower and higher than the median Ki67 value, and we did not find a significant difference

(P>0.05).

Figure 3. Photomicroscopy of cSCC in dogs with Ki67 immunolabeling. (A) High

nuclear immunostaining of neoplastic cells at D0 (score 3). (B) Low nuclear immunostaining

at D21 (score 1). The 4"x5" area, used for counting the cells, is observed in the images

(immunocytochemistry, Envision, DAB, counterstaining with Harris hematoxylin, x400).

341   **Fig. 1. Evaluation of tumor volume, clinical response by tumor volume, mitotic**

342 **index, intratumoral necrosis and Ki67 expression before and after treatment by ECT in**

343 **dogs with cSCC**. (E) Lower proliferative index (p=0.031) was observed at D21 when

344 compared to D0.

346   For BAX and Bcl-2, cytoplasmic expression was identified (Fig 4A). We found

347 median BAX expression levels of 21.02% (ranging from 8.34 to 70.56) at D0 and 24.53%

348 (ranging from 10 to 74.47) at D21. There was no significant difference in BAX expression

349 between D0 and D21 (P>0.05) (Fig 4B). There was no significant difference in overall

350 survival between subjects with low and high BAX expression levels (P>0.05) (Suppl Fig 1C).

352 **Fig. 4. BAX and Bcl-2 immunoexpression levels and the correlation between both**

353 **apoptotic markers in dogs with cSCC before and after ECT.** (A) Cytoplasmic expression

354 of BAX (arrows). (B) There was no significant difference in BAX expression between D0 and

355 D21 (P>0.05).

357 **Suppl Fig. 1. Survival analysis according to lymph node involvement, tumor size, and**

358 **apoptotic markers in dogs with cSCC.** (C) There was no significant difference in overall

359 survival between subjects with high and low levels of BAX (P>0.05).

361   Regarding Bcl-2 expression (Fig 4C), we found medians of 20.62% (ranging from

362 2.76% to 48.37) at D0 and 26.42% at D21 (ranging from 0.65 to 77.23). There was no

363 statistically significant difference in Bcl-2 expression between D0 and D21 (P>0.05) (Fig

364 4D). When we evaluated the survival time between samples with low Bcl-2 expression versus

365 those with high expression, we did not find a significant difference (P>0.05) (Suppl Fig 1D).

Furthermore, a positive correlation was observed between BAX/Bcl-2 expression

levels before ECT (D0) (p=0.0379, r=0.5067) (Fig 4E). However, after treatment (D21), we

did not find a significant Gaussian approximation (p=0.7370, r=0.0911) (Fig 4F). Regarding

the correlation between BAX/Ki67 and Bcl-2/Ki67, there was no significant difference among

them at different time points (p>0.05).

Fig. 4. BAX and Bcl-2 immunoexpression and the correlation between both apoptotic

markers in cSCC before and after ECT. (C) Cytoplasmic expression of Bcl-2 (arrows). (D)

There was no statistically significant difference in Bcl-2 expression between D0 and D21

(P>0.05). (E) Positive correlation between BAX and Bcl-2 before ECT (D0) (p=0.0379,

r=0.5067). (F) Twenty-one days after treatment, no significant Gaussian approximation was

observed (p=0.7370, r=0.0911)

Western blot analysis

We found a 26-kDa band for Bcl-2 and a 21-kDa band size for BAX. Both antibodies

showed only one band each in the western blot analysis (Fig 5).

Discussion

cSCC is a very common tumor in tropical regions, and one of the biggest therapeutic

challenges for tropical regions is removing the animal from exposure to sunlight during

treatment. Because cSCC has a low level response to chemotherapy and a low metastatic rate,

a local approach is required to achieve long-term disease-free interval and overall survival.

Surgery is the primary treatment for cSCC. However, many patients have multiple tumors at

the time of diagnosis, making surgical approaches difficult. Therefore, new techniques are

required to improve the outcomes of canine patients with cSCC.

In this study, we evaluated ECT as a primary therapy for cSCC. We determined the

association of different parameters with overall survival and tumor response to ECT. We

demonstrated a reduction of the tumor size after ECT (D21), even in subjects with poorly

differentiated tumors. Four lesions (in four different patients) presented progressive disease

(PD). Of these four lesions, two had large volumes (ID 5 and 8), indicating that ECT may be a

treatment option for small tumors. Unfortunately, we have no studies in Veterinary Medicine

indicating the association of a tumor size cutoff with prognosis. In a large series study about

the use of ECT in basal cell carcinoma (BCC) in humans, a 50% CR rate was observed after a

single ECT cycle in primary BCC. Interestingly, in the same study, a second ECT cycle

increased the CR rate from 50 to 63%, and retreatment was more advantageous in patients

with local BCC. Thus, retreatment with ECT seems a reasonable option in patients with small

BCCs to reduce the response duration [47]. In accordance with this human study, our results

indicated that ECT could be a good therapeutic option for small tumors, and large may need

more than one round of ECT. A previous case report by our study group observed complete

remission of a digital trichoblastoma after three sessions of ECT, reaffirming the usefulness of

repeated sessions of ECT for larger tumors [20].

Among the limiting factors of our study, there was an absence of a cutoff point for

tumor size in cutaneous neoplasms in dogs, unlike in human medicine. In addition, the staging

adopted in veterinary medicine does not take into account the size of the subject, which may

influence the behavior of a tumor of the same size in dogs of different sizes. The small sample

size is also a limitation of our study. However, we performed a G power analysis to guarantee

that our tumor group had a sufficient number of samples.

Interestingly, lymph node involvement was not a prognostic factor in dogs with cSCC.

The number and size of the lesions seems to be more important than the involvement of local

lymph nodes. This can be explained by the invasive behavior of cSCC [5]. Usually, these

tumors are more locally invasive than metastatic tumors [5,34,48]. On the other hand, the

number of mitoses was correlated with overall survival in our study. Because the mitotic

index is a routine evaluation, it will be important to evaluate the sensitivity and specificity of

these indices to predict outcomes in dogs with cSCC. However, a large sample size is

necessary. We also found more necrosis in samples after ECT, indicating that this therapy

induces both apoptosis and necrosis. Thus, the greater amount of necrosis identified at D21

was due to the therapy.

One of the most interesting results of our research was the decreased proliferative

index in tumors after ECT. Because ECT induces tumor apoptosis and necrosis, the number of

proliferating cells was reduced. Furthermore, we did not find any difference in cell apoptosis

between the two time points (D0 and D21), indicating that another mechanism may be

involved in inhibiting tumor proliferation after ECT. Because ECT is a recent technique in

Veterinary Medicine, a limited number of studies have demonstrated the ECT antitumoral

effect in cSCC.

In general, the high expression level of Ki67 observed in SCC, especially in poorly

differentiated SCC, is related to an aggressive potential [9,34,37,49]. However, in the lesions

studied, the proliferation index could also be considered high despite the predominance of

well-differentiated SCC (83.3%), suggesting the aggressive potential, despite the degree of

differentiation. Additionally, there is no established cutoff point for Ki67 to be considered an

indicator of recurrence or metastasis, as occurs in cases of cutaneous mast cell tumor, which is

23 positive cells in a 10 mm x 10 mm / 400x area [50]. It should be noted that the complete

evaluation of proliferation markers may be more effective in predicting the biological

behavior of the tumor. In this context, Auler et al. [49] reported the case of a dog with well-

differentiated SCC in the foreskin with inguinal lymph node metastasis. This animal showed

high Ki67 positivity in the tumor tissue, in addition to the high expression of the growth

factors HER-2 and EGFR (epidermal growth factor receptor).

A previous study observed the same pattern in mast cell tumors as the pattern

identified in this study with regard to the proliferative index, which was observed in fewer

positive cells for Ki67 among the neoplastic cells 28 days after treatment [19].

Regarding the apoptotic markers, all cSCC lesions showed cytoplasmic positivity for

BAX. These results differ from those of Madewell et al. [28], who observed less than 2%

BAX expression in cSCC in felines. However, other human studies have observed the

expression of BAX in cutaneous and pulmonary SCC, which ranged from 53-100% [51,52].

A recent study observed that the expression of BAX in normal keratinocytes in the epidermis

of dogs ranged from very weak to moderate, elucidating their importance in the pathogenesis

of diseases [53]. To avoid any unspecific cross-reaction with canine tissue and the primary

antibody, we performed western blotting for both antibodies using cSCC samples. Both

antibodies presented only one specific band, confirming the reactivity of both antibodies with

canine tissue.

Likewise, no significant difference was observed between the expression of the anti-

apoptotic Bcl-2 protein before and after treatment with ECT. This is the first study to evaluate

whether the immunohistochemical expression of Bcl-2 protein is altered by ECT in cSCC. A

previous report evaluated the expression in mast cell tumors and observed increased

expression of anti-Bcl-2 28 days after ECT and then decreased expression at D46 [19]. In our

study, all cSCC samples showed immunolabeling of Bcl-2. However, other studies have

shown different results and only one evaluated Bcl-1 by IHC in dogs with cSCC (n = 5),

among which only one showed positivity [54]. In humans, Puizina-Ivic et al. [55] observed an

absence of Bcl-2 positivity in SCC (n = 20), whereas Abu Juba et al. [56] observed positivity

in 50% of cSCC samples (n = 10).

It must be noted that the correlation between BAX/Bcl-2 markers was significant

before treatment, but no difference was observed afterwards, and no difference was observed

between BAX/Ki67 and Bcl-2/Ki67 at both time points. In human studies, the Bax/Bcl-2 ratio

can act as a rheostat that determines cell susceptibility to apoptosis [56], and lower levels of

this ratio may lead to resistance of human cancer cells to apoptosis. In colorectal tumors,

BAX and Bcl-2 expression levels were the most predictive of outcome when the Bax/Bcl-2

expression ratio was used [57]

Although the chemotherapeutic agents used have the capacity to cause cell damage

that induces cell death by apoptosis, the BAX and Bcl-2 markers in the samples were not

altered at the time point evaluated (D21). It is possible that other proteins involved in the

apoptosis pathways exert a greater influence on the process of cell death induced by ECT. In

addition, it cannot be ruled out that, in view of the dynamism and complexity of the apoptosis

process, the expression of BAX and Bcl-2 may have been influenced by the ECT prior to the

assessed moment, but not on the 21st day of treatment. A limitation in scoring a single protein

is that the expression of a single protein may not reflect the level of apoptosis because

apoptosis is a dynamic, complex process.

Regarding the proliferative index, 21 days after the application of ECT, it was possible

to observe a significant reduction in proliferative indices in the lesions sampled. The same

behavior was also observed in mammary neoplasms of human patients submitted to systemic

chemotherapy because there was a significant reduction in the level of Ki67 in the tumor at 21

days of treatment [58,59]. However, in the veterinary literature, no research was found

regarding the proliferative index before and after treatment in malignant neoplasms.

Conclusions

ECT was able to reduce tumor volume and cellular proliferation in cSCC.

Furthermore, our results showed that the BAX and Bcl-2 proteins did not show alterations in

their expression levels in response to ECT at the time points evaluated. Therefore, studies

involving the serial evaluation of these proteins, as well as the investigation of other proteins

involved in apoptosis, may contribute to the understanding of the effect of ECT on apoptosis

in canine cSCC.

Disclosure

496 All the authors have declared that they have no competing interests.

References

1. Muller WHM, and Kirk R. Neoplastic and Non-neoplastic tumors. In: Muller & Kirk.

Small Animal Dermatology, Editora Elsevier, 7[th] edition, 2013, p. 774-780.

2. Teifke JP, Lohr CV, and Shirasawa H (1998). Detection of canine oral papillomavirus-

DNA in canine oral squamous cell carcinomas and p53 overexpressing skin papillomas of

the dog using the polymerase chain reaction and non-radioactive in situ hybridization. *Vet*

Microbiol **60**, 119–130.

3. Schwegler K, Walter JH, and Rudolph R. (1997). Epithelial neoplasms of the skin, the

cutaneous mucosa and the transitional epithelium in dogs: an immunolocalization study

for papillomavirus antigen. *Zentralblatt fur Veterinarmedizin* **44**, 115–123.

4. Kirkham N. (1997) Tumors and cysts of the epidermis. In: *Lever's Histopathology of the*

Skin, 8th edition, 712–17. Lippincott Raven, Philadelphia.

5. Gross TL, Ihrke PJ, Walder EJ, and Affolter VK. Epidermal tumors. (2005). In: ______

Skin diseases of the dog and cat. Clinical and histopathologic diagnosis. Oxford:

Blackwell Science Ltd, 581-585.

6. Bevier DF, and Goldschmidt MH. (1981). Skin tumors in the dog. Part I: Epithelial

tumors and tumor-like lesions. *Comp Cont Ed 3,* 389–98.

7. Goldschmidt MH, and Shofer FS. (1992). *Skin Tumors of the Dog and Cat*, Pergamon

Press, Oxford, 37–49.

8. Mittelbronn MA, Mullins DL, Ramos-Caro FA, and Flowers FP. (1998) Frequency of pre-

existing actinic keratosis in cutaneous squamous cell carcinoma. *Int J Dermatol* **37**, 677–

81.

9. Poggiani SSC, Hatayde MR, Laufer-Amorim R, and Werner J. (2012) Expression of

Cyclooxygenase-2 and Ki-67 in Actinic Keratosis and Cutaneous Squamous Cell

Carcinoma in Dogs. *Open J Vet Med* **2**, 41-47.

10. Rogers KS, Helman RG, and Walker MA (1995). Squamous cell carcinoma of the canine

nasal planum: eight cases (1988-1994). *J Am Anim Hosp Assoc* **31**, 373–378.

11. Kirpensteijn J, Withrow SJ, and Straw RC (1994). Combined resection of the nasal

planum and premaxilla in three dogs. *Vet Surg* **23**, 341–346.

12. Lascelles BD, Parry AT, Stidworthy MF, Dobson JM, and White RA. (2000). Squamous

cell carcinoma of the nasal planum in 17 dogs. *Vet Rec* **147**, 473–476.

13. Withrow SJ, and Straw RC (1990). Resection of the nasal planum in nine cats and five

dogs. *J Am Anim Hosp Assoc* **26**, 219–222.

14. Brougham ND, Dennett ER, Cameron R, and Tan ST. (2012). The incidence of metastasis

from cutaneous squamous cell carcinoma and the impact of its risk factors. *J Surg Oncol*

106, 811-815.

15. Karia PS, Jambusaria-Pahlajani A, Harrington DP, Murphy GF, Qureshi AA, and
Schmults CD. (2014). Evaluation of American Joint Committee on Cancer, International
Union Against Cancer, and Brigham and Women's Hospital tumor staging for cutaneous
squamous cell carcinoma. *J Clin Oncol* **32**, 327-334.

16. Schmults CD, Karia PS, Carter JB, Han J, and Qureshi AA. (2013). Factors predictive of
recurrence and death from cutaneous squamous cell carcinoma: a 10-year, single-
institution cohort study. *JAMA Dermatology* **149**, 541-547.

17. Spugnini EP, Azzarito T, Fais S, Fanciulli M, and Baldi A. (2016). Electrochemotherapy
as First Line Cancer Treatment: Experiences from Veterinary Medicine in Developing
Novel Protocols. *Curr Cancer Drug Targets* **16**, 43-52, 2016.

18. Spugnini EP, Baldi F, Mellone P, Feroce F, D'avino A, Bonetto F, Vincenzi B, Citro G,
and Baldi, A. (2007). Patterns of tumor response in canine and feline cancer patients
treated with electrochemotherapy: preclinical data for the standardization of this treatment
in pets and humans. *J Transl Med* **7**, 48.

19. Salvadori C, Svara T, Rocchigiani G, Millanta F, Pavlin D, Cemazar M, et al. Effects of
electrochemotherapy with cisplatin and peritumoral IL-12 gene electrotransfer on canine
mast cell tumors: a histopathologic and immunohistochemical study. Radiol Oncol. 2017;
51(3): 286-294.

20. Dos Anjos DS, Rossi YA, Magalhaes LF, Calazans SG, Fonseca-Alves CE. Digital
trichoblastoma treated with electrochemotherapy in a dog. Vet Record. 2018; 6: e000671.
doi: 10.1136/vetreccr-2018-000671.

21. Groeger AM, Esposito V, De Luca A, Cassandro R, Tonini G, Ambrogi V, Baldi F,
Goldfarb R, Mineo TC, Baldi A, and Wolner E (2004). Prognostic value of
immunohistochemical expression of p53, BAX, BCL-2 and BCL-Xl in resected non small
cell lung cancer. *Histopathology* **44**, 54-63.

22. Karam JA, Lotan Y, Karakiewicz PI, Ashfaq R, Sagalowsky AI, Roehrborn CG, and

Shariat SF. (2007). Use of combined apoptosis biomarkers for prediction of bladder

cancer recurrence and mortality after radical cystectomy. *Lancet Oncol* **8**, 128 – 36.

23. Zeestraten ECM, Benard A, Reimers MS, Schouten PC, Liefers GJ, van de Velde CJH, et

al. The Prognostic Value of the Apoptosis Pathway in Colorectal Cancer: A Review of the

Literature on Biomarkers Identified by Immunohistochemistry. Biomarkers Cancer. 2013;

5: 13-29. doi: 10.4137/BIC.S11475.

24. De Bruin EC, Medema JP. Apoptosis and non-apoptotic deaths in cancer development and

treatment response. Cancer Treat Reviews. 2008; 34: 737-749.

doi:10.1016/j.ctrv.2008.07.001.

25. Meichner K, Fogle JE, English L, Suter SE. Expression of Apoptosis-regulating Proteins

Bcl-2 and Bax in Lymph Node Aspirates from Dogs with Lymphoma. J Vet Intern Med.

2016; 30(3): 819-826. doi: 10.1111/jvim.13937.

26. Dolka L, Król M, Sapierzynski R. Evaluation of apoptosis-associated protein (Bcl-2, Bax,

cleaved caspase-3 and p53) expression in canine mammary tumors: An

immunohistochemical and prognostic study. Res Vet Sci. 2016; 105: 124-133. doi:

10.1016/j.rvsc.2016.02.004

27. Yildirim F, Sonmez K, Ozyogurtçu H, Sennazli G, Gurel A, Gunduz MC, et al.

Evaluation of Bcl-2, Bcl-xl, and Bax expression and apoptotic index in canine mammary

tumours. Kafkas Univ Vet Fak Derg. 2014; 20(4): 513-520.

28. Madewell BR, Gandour-Edwards R, Edwards BF, Matthews, KR and Griffey (2001).

Bax/Bcl-2: Cellular modulator of apoptosis in Feline Skin and Basal Cell Tumours. *J

Comp Path* **124**, 115-121.

29. Scase TJ, Edwards D, Miller J, Henley W, Smith K, Blunden A, and Murphy S. (2006). Canine mast cell tumors: correlation of apoptosis and proliferation markers with prognosis. J Vet Inter Med 20, 151–158.

30. Bergkvist GT, Argyle DJ, Morrison L, Macintyre N, Hayes A, and Yool DA. (2011). Expression of epidermal growth factor receptor (EGFR) and Ki67 in feline oral squamous cell carcinomas (FOSCC). *Vet Comp oncol* **9**, 106–117.

31. Pereira RS, Schweigert A, DE Melo GD, Fernandes FV, Sueiro FAR, and Machado GF. (2013). Ki-67 labeling in canine perianal glands neoplasms: a novel approach of immunohistological diagnostic and prognostic. *BMC Vet Res* **9**, 83.

32. Brodzki A, Lopuszyñski W, Brodzki P, and Tatara M (2014). Diagnostic and prognostic value of cellular proliferation assessment with Ki-67 in dogs suffering from benign and malignant perianal tumors. *Folia Biologica (Kraków)* **62**, 235-241.

33. Peña L, Zarate AINR, Pérez-Alenza MD, Cuesta PL, Castaño M. (1998). Immunohistochemical Detection of Ki-67 and PCNA in Canine Mammary Tumors: *Relationship to Clinical and Pathologic Variables* **10**, 237-246.

34. Khodaeiani E, Fakhrjou A, Amirnia M, Babaei-Nezhad S, Taghvamanesh F, Razzagh-Karimi E, H (2013). Immunohistochemical Evaluation of p53 and Ki67 expression in Skin Epithelial Tumors. *Indian J Dermatology* **58**, 181-187.

35. Xie S, Liu Y, Qiao X, Hua R, Wang K, Shan XF, and Cai ZG (2016). What is the Prognostic Significance of Ki-67 Positivity in Oral Squamous Cell Carcinoma? *J Cancer* **7**, 758-767.

36. Batinac T, Zamolo G, Coklo M, Hadzisejdic I, Stemberger C,and Zauhar G (2006). Expression of cell cycle and apoptosis regulatory proteins in keratoacanthoma and squamous cell carcinoma. *Pathol Res Pract*. 202, 599–607.

37. Stratigos AJ, Kapranos N, Petrakou E, Anastasiadou A, Pagouni A, Christofidou E, et al.

(2005). Immunophenotypic analysis of the p53 gene in non-melanoma skin cancer and

correlation with apoptosis and cell proliferation. *J Eur Acad Dermatol Venereol* **19**:180–6.

38. Cadossi R, Ronchetti M and Cadossi M. (2014). Locally enhanced chemotherapy by

electroporation: Clinical experiences and perspective of use of electrochemotherapy.

Future Oncol **10**, 877–890.

39. Spugnini EP and Baldi A. (2014). Electrochemotherapy in veterinary oncology: From

rescue to first line therapy. *Methods Mol Biol* **1121**, 247–256

40. Campana LG, Clover AJP, Valpione S, Quaglino P, Gehl J, Kunte C, et al.

Recommendations for improving the quality of reporting clinical electrochemotherapy

studies based on qualitative systematic review. Radiol Oncol 2016; 50: 1-13. doi:

10.1515/raon-2016-0006.

41. Owen LN. (1980). TNM Classification of tumor in domestic animals, World Health

Organization.

42. Marty M, Sersa G, Garbay J, Gehl J, Collins CG, Snoj M, Billard V, Geertsen PF, Larkin

JO, Miklavcic D, Pavlovic I, Paulin-Kosir SM, Cemazar M, Morsli N, Soden DM, Rudolf

Z, Robert C, O'Sullivan GC, and Mir LM. (2006). Electrochemotherapy – an easy, highly

effective and safe treatment of cutaneous and subcutaneous metastases: results of ESOPE.

(European Standard Operating Procedures of Electrochemotherapy) study. *Eur J Cancer*

4, 3-13.

43. Eisenhauer EA1, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, Dancey

J, Arbuck S, Gwyther S, Mooney M, Rubinstein L, Shankar L, Dodd L, Kaplan

R, Lacombe D, and Verweij J. (2009). New response evaluation criteria in solid tumours:

Revised RECIST guideline (version 1.1). *Eur J Cancer* **45**, 228–247.

44. Fonseca-Alves CE, Rodrigues MMP, De Moura VMBD, Rogatto, SR, Laufer-Amorim, R. Alterations of C-MYC, NKX3.1, and E-cadherin expression in canine prostate carcinogenesis. Microscopy Research Technique. 2013; 76: 1250-1256. doi: http://dx.doi.org/10.1002/jemt.22292.

45. Fonseca-Alves CE, Kobayashi PE, Laufer-Amorim R. Evaluation of NKX3.1 and C-MYC expression in canine prostatic cancer. Res Vet Sci. 2018; 118: 365-370.

46. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal Biochem. 1976; 72:248-54.

47. Campana LG, Marconato R, Valpione S, Galuppo S, Alaibac M, Rossi CR. Basal cell carcinoma: 10-year experience with electrochemotherapy. J Transl Med. 2017; 15:122. doi. 10.1186/s12967-017-1225-5.

48. Lascelles BDX, Parry AT, Stidworthy MF, Dobson JM, White RAS. Squamous cell carcinoma of the nasal planum in 17 dogs. Vet Record. 2000; 147: 473-476.

49. Auler PA, Gamba CO, Horta RS, Lavalle GE, and Cassali GD (2014). Metastatic well differentiated squamous cell carcinoma in the prepuce of a dog: a report of clinicopathological, immunophenotypic and therapeutic approach. *Arq Bras Med Vet Zootec* **66**, 1317-1322.

50. Webster JD, Yuzbasiyan-Gurkan V, Miller RA, Kaneene JB, and Kiupel M (2007). Cellular proliferation in canine cutaneous mast cell tumors: Associations with c -KIT and its role in prognostication. *Vet Pathol* **44**, 298-308.

51. Abu Juba B, Sovrea A, Crisan D, Melincovici C, Coneac A, Badea M, and Crisan M (2013). Apoptotic markers in photoinduced cutaneous carcinoma. *Rom J Morphol Embryol* **54**, 741-747.

52. Porebska I, Kosacka M, Sobanska E, and Wyrodek (2015). Comparative expression of apoptotic markers in lug adenocarcinoma and squamous cell carcinoma. *Advs Exp Medicine Biology- Neuroscience and Respiration* **16**, 101-107.

53. Croci M, Dettwiler M, Vaughan L, and Guscetti F (2013). Immunohistochemical expression of Bax and Bak in canine non-neoplastic tissues. *Vet J* **198**, 131-140.

54. Pieper JB, Stern AW, LeClerc SM, and Campbell KL (2015). Coordinate expression of cytokeratins 7 and 14, vimentin, and Bcl-2 in canine cutaneous epithelial tumors and cysts. *J Vet Diagnost Invest* **27**, 497-503.

55. Puizina-Ivic N, Sapunar D, Marasovic D, and Miric L (2008). An overview of Bcl-2 expression in histopathological variants of basal cell carcinoma, squamous cell carcinoma, actinic keratosis and seborrheic keratosis. *Coll Antropol*, **32**, 61-65.

56. Raisova M, Hossini AM, Eberle J, Riebeling C, Wieder T, Sturm I, et al. The Bax/Bcl-2 ratio determines the susceptibility of human melanoma cells to CD95/Fas-mediated apoptosis. J Invest Dermatol. 2001; 117(2):333-40.

57. Khodapasand E, Farrokhi F, Kamalidehghan B, Houshmand M. Is Bax/Bcl-2 Ratio Considered as a Prognostic Marker with Age and Tumor Location in Colorectal Cancer? Iran Biomed J. 2015; 19(2): 69-75. doi: 10.6091/ibj.1366.201.

58. Burcombe R, Wilson GD, Dowsett M, Khan I, Richman PI, Daley F, Detre S, and Makris A (2006). Evaluation of Ki-67 proliferation and apoptotic index before, during and after neoadjuvant chemotherapy for primary breast cancer. Breast Cancer Res, 8 R31.

59. Jin G, Han Y, Liu C, Chen L, Ding B, Xuan S, Liu X, Ma G, Gao J, and Tian X (2015). Evaluation of biomarker changes after administration of various neoadjuvant in breast cancer. Int J Clin Exp Pathol 8, 914-921.