Cholesterol-dependent enrichment of understudied erythrocytic stages of human *Plasmodium* parasites

Audrey C. Brown[1], Christopher C. Moore[2], Jennifer L. Guler[1,2]

[1]Department of Biology, University of Virginia, Charlottesville, VA.

[2]Division of Infectious Diseases and International Health, University of Virginia, Charlottesville, VA

Corresponding author contact: jlg5fw@virginia.edu

Abstract

Plasmodium protozoan parasites undergo rounds of asexual replication inside human erythrocytes, progressing from ring stage, to trophozoites and schizonts, before egress and reinvasion. Given the discovery of ring-specific artemisinin tolerance and quiescence in *Plasmodium falciparum*, there is great urgency to better understand ring stage biology. However, the lack of an effective enrichment method has left rings and related parasite stages understudied compared to their late stage counterparts, which can be easily isolated due to their paramagnetic properties. Here, a method for separating *all Plasmodium* infected erythrocytes from uninfected erythrocytes is presented. This approach takes advantage of streptolysin-O (SLO) to preferentially lyse uninfected erythrocytes as previously shown by Jackson, *et al.* Following lytic treatment, Percoll gradient centrifugation removes lysed cells, leaving an intact cell population enriched in infected erythrocytes. This SLO-Percoll (SLOPE) method is effective on stages from the entire erythrocytic cycle, including previously inaccessible forms such as circulating rings from malaria-infected patients and artemisinin-induced quiescent parasites. Furthermore, the utility of SLOPE is extended to multiple media formulations used for the propagation of two human *Plasmodium* species. The alteration of external cholesterol levels modulates SLOPE effectiveness, demonstrating the role of erythrocyte membrane cholesterol in lytic discrimination. Importantly, enrichment does not impact parasite viability, which establishes the non-toxic nature of SLOPE. Targeted metabolomics of SLOPE-enriched ring stage samples confirms the impact on treated samples; parasite-derived metabolites are increased and contaminating host material is reduced compared to non-enriched samples.

Importance

Malaria is caused by infection with protozoan *Plasmodium* parasites and is responsible for over 400,000 deaths annually. The availability of effective antimalarial drugs is critical to the reduction of malaria-related mortality, yet widespread resistance highlights the need for the continued study of *Plasmodium* biology. The SLOPE method is an accessible, scalable, rapid (30-40min), and non-toxic enrichment method that is broadly effective on many erythrocytic stages. This method is ideal for use upstream of a variety of sensitive analyses, which will increase experimental quality in virtually all areas of asexual *Plasmodium* parasite research. Further, because the consumption of cholesterol is a common characteristic of other intracellular parasites (both bacteria and other protozoa), SLOPE holds potential for extension to other relevant pathogens.

Introduction

Malaria, caused by protozoan parasites of the *Plasmodium* genus, is a continuing threat to global health. A total of five *Plasmodium* species cause malaria in humans, with *Plasmodium falciparum* being responsible for the large majority of malaria morbidity and mortality (1). While the global malaria burden has decreased over the past decade, the emergence and spread of antimalarial resistant *Plasmodium* threatens to undo this progress and emphasizes the dire need to understand more about the biology of this parasite. The current World Health Organization recommendation for treatment of malaria is artemisinin combination therapy (2). However, clinical resistance has now been reported to both artemisinin and almost all of its partner drugs (3-5).

All symptoms of malaria, including cyclical fevers and hypoglycemia, occur due to the asexual replication cycle of the parasite within human erythrocytes (**Fig. 1A**). Parasites undergo rounds of replication progressing from the ring stage, to trophozoites and schizonts, before rupturing from host erythrocytes to release merozoites, which go on to invade new erythrocytes and continue the cycle of infection (6). Many studies aiming to understand the biology of asexual *Plasmodium* are performed only on late stage parasite samples (trophozoites and schizonts). This is due in part to the larger biomass of these stages but also to the existence of an effective enrichment method (7); erythrocytes infected with late stage parasites can be separated from uninfected erythrocytes using the paramagnetic properties of hemozoin, a byproduct of parasite maturation. The ability to enrich for late stages, thus limiting noise from excess uninfected erythrocytes, has fueled the recent explosion of omics-based studies of late stage *P. falciparum* biology and antimalarial drug action (8-16).

The lack of an effective enrichment method dramatically limits our ability to study the ring stage parasites, as well as ring-derived forms. For example, recent proteomics and metabolomics studies of this early erythrocytic *P. falciparum* show the heavy influence of host metabolites in non-enriched preparations, which contributes to variability between samples and obscures parasite phenotypes (16-18). This limitation is particularly acute when dealing with material directly from malaria patients as only ring forms of *P. falciparum* are collected during blood draws (19), and there is a high ratio of uninfected host cells to parasite-infected cells (typically 100:<4) (20). Ample access to clinically-relevant parasites is important for the study of antimalarial resistance especially in the context of artemisinin, which impacts rings differently than later stages through the induction of quiescence (21-23). This impactful biological discovery highlights the need to improve upstream purification steps for the study of ring stage biology.

In this study, we present a method for the enrichment of viable ring stage *Plasmodium*-infected erythrocytes that is simple to employ, rapid, and non-toxic to the parasite. This method can be scaled to the needs of individual experiments without compromising these attributes. We show

that the effective removal of uninfected erythrocytes is unaffected by standard culture media
formulations and is conserved across multiple *Plasmodium* species and parasite sources, further
highlighting utility for a range of experimental needs. To our knowledge, this is the first
enrichment method that is effective on ring stage parasites to increase parasitemia (the
percentage of erythrocytes infected with a parasite) and reduce host erythrocyte contamination.
The "SLOPE" enrichment method offers a tool to increase research quality in virtually all areas
of *Plasmodium* asexual parasite research.

Figure 1. SLOPE enrichment overview. (A) The asexual replication cycle of *Plasmodium* occurs inside erythrocytes. Both *P. falciparum* and *P. knowlesi,* which take 48 and 24 hours, respectively, to complete the replication cycle, can be propagated *in vitro* indefinitely. (B1) Hemolytic activity of Streptolysin-O (SLO) was assessed on uninfected erythrocytes to define a unit (the amount of SLO necessary for 50% lysis of 50μl of uninfected erythrocytes at 2% hematocrit in PBS for 30 min at 37∘C). (B2) Ring stage synchronized cultures were treated with a defined quantity of SLO units to preferentially lyse uninfected erythrocytes. (B3) SLO treated samples were layered over a 60% Percoll gradient and centrifuged to separate lysed ghosts from intact cells. (B4) The upper layer of Percoll containing lysed ghosts was discarded while the lower, intact, infected erythrocyte enriched fraction was collected. Uninfected erythrocytes, red circles; Infected erythrocytes, red circles with black dots; lysed membranous ghosts, white circle with dashed outline.

89
90 **Results**
91
A two-step SLOPE protocol effectively enriches ring-infected erythrocytes
We developed the Streptolysin-O (SLO)–Percoll-based protocol (termed SLOPE) for enrichment
of ring stage *P. falciparum* infected erythrocytes (**Fig. 1B,** see *Materials and Methods* and **Text
S1** for protocol details). SLO is a pore forming toxin that preferentially lyses uninfected
erythrocytes, leaving the large majority of infected erythrocytes intact (24). Using the protocol
outlined by Jackson, *et al*. with slight modifications, we were able to achieve levels of lysis
discrimination for erythrocyte populations that were comparable to this original report (93.4%
and 9.9% lysis of uninfected and infected erythrocytes, respectively, **Fig. 2A,** yellow highlight).

In addition to reproducing lysis levels, we quantified uninfected and infected erythrocyte lysis
across a gradient of SLO concentrations. More complete lysis of uninfected cells (>99%) was
obtained at the cost of greater infected erythrocyte lysis by increasing SLO quantity. For
example, we found that 47U of SLO leads to >99% lysis of uninfected erythrocytes and 40%
lysis of infected erythrocytes (**Fig. 2A,** grey highlight). Additionally, we show that SLO favors
uninfected erythrocyte lysis irrespective of parasite line or culture media (**Fig 2A**; **Fig S1**). SLO
showed comparable lysis discrimination between uninfected and infected erythrocytes in
parasites grown in two common media formulations, RPMI 1640 supplemented with AlbuMAX II
or RPMI 1640 supplemented with 20% human serum (**Fig. 2A**, maximum difference between
uninfected and infected erythrocyte lysis: AlbuMAX supplementation = 83.5%; serum
supplementation = 81.3%). Furthermore, SLO lysis showed considerable discrimination between

Figure 2. SLOPE enriches ring stage *Plasmodium* parasites irrespective of species or media formulation.
(A) SLO lysis of uninfected erythrocytes (uRBCs) and infected erythrocytes (iRBCs) from (left) ring-stage
synchronized *P. falciparum* grown in RPMI 1640 supplemented with 20% human serum (N = 6; 3 replicates each
of lines Hb3 and K1), (middle) RPMI 1640 supplemented with Albumax II (N = 9; 3 replicates each of lines Hb3,
K1, and MRA 1240), and asynchronous *P. knowlesi* grown in RPMI 1640 supplemented with Albumax II (N=3).
(B) SYBR-Green based flow cytometry measurements before and after SLOPE enrichment. Flow plots show
single cells within the erythrocyte size range. The infected erythrocyte fraction ("iRBCs") is denoted within the
dashed red gate. (C) Parasitemia fold increase upon treatment with increasing SLO units relative to untreated
controls. Represented samples were grown in RPMI 1640 supplemented with Albumax II (N = 9; 3 replicates
each of lines Hb3, K1, and MRA 1240). All error bars represent S.E.M.

uninfected and infected erythrocytes for both *P. falciparum* and the zoonotic species,
Plasmodium knowlesi (**Fig. 2A)**.

After treatment with SLO, parasite samples contained a mixture of primarily lysed erythrocyte
membranes (from uninfected erythrocyte lysis), termed ghosts, and intact erythrocytes (enriched
in infected erythrocytes) (**Fig. 1B2**). For the second part of the enrichment protocol, we exploited
the difference in density of intact erythrocytes and ghost membranes to separate the two
populations (**Fig. 1B3**). We found that intact erythrocytes travel through a 60% Percoll gradient
during centrifugation while erythrocyte ghosts remained above the Percoll layer (**Fig. 1B4**, **Fig.
3**). Collection of the intact erythrocyte population leads to a sample with up to a greater than 20-
fold increase in the parasite to erythrocyte membrane ratio. In one representative trial, we

Figure 3. Validation of ghost separation from intact erythrocytes by Percoll step. (A) Intact erythrocytes are shown as CD235a (red) positive and spectrin (green) negative. Lysed erythrocytes, termed ghosts, are shown as CD235a and spectrin double positive (yellow in merge). All images show ring-stage *P. falciparum* line MRA 1240 parasites stained with SYBR Green (cyan). 40X Magnification; bar represents 10μm. Saponin samples were treated with 0.15% saponin for 5 minutes. SLO samples were treated with 40U of SLO for 6 minutes but were not centrifuged through a Percoll gradient. SLOPE samples were also treated with 40U SLO but were subjected to Percoll gradient centrifugation (B) Proportions of lysed ghosts and intact erythrocytes quantified from fluorescence microscopy imaging across different treatments using SYBR Green dye and CD235a and spectrin antibodies (*N* = 3; 400 erythrocytes per condition per preparation, error bars represent S.E.M.).

122 demonstrate enrichment of ring stage samples to a final parasitemia over 80% (22-fold increase
123 over starting parasitemia, **Fig. 2B**). Higher levels of enrichment were attainable with higher
124 amounts of SLO (up to 48-fold with 55 units of SLO, **Fig. 2C**) but this is at the cost of parasite
125 material (up to ~40% of infected erythrocytes, **Fig. 2A**).
126
We performed SLOPE on mixed stage populations to determine whether our protocol biases
enrichment of certain asexual stages. We found that SLOPE enrichment of lightly synchronized
cultures (i.e. contains rings and some residual trophozoites) did not result in any alteration of the
ring to trophozoite proportion (**Fig. S2A**). SLOPE enrichment was also unbiased when
performed on asynchronous cultures (includes rings, trophozoites, and schizonts). However, the
protocol did result in a non-significant drop in schizont percentage (**Fig. S2B**). Centrifugation
and washes likely lysed fragile
schizont-infected erythrocytes
resulting in partial loss of the
schizont population.
137
Next, we employed
immunofluorescence detection of
erythrocyte components to
confirm that Percoll gradient
density centrifugation removed
ghost erythrocyte membranes
from SLO treated samples (**Fig.
3**). Specifically, we visualized all
erythrocyte membranes through
detection of an external
erythrocyte surface protein
(CD235a); we identified ghosts
through detection of an intra-
erythrocyte protein (spectrin).
Untreated (control) samples
contain high erythrocyte
membrane to parasite ratios
(~100:1 due to ~1% parasitemia)
as expected, and almost all
membranes (>99%) were in the
form of intact cells (spectrin
negative, **Fig. 3A** and **B**). All
erythrocyte membranes in
samples treated with a

Figure 4. SLOPE enriched parasites remain viable. (A) Mitoprobe DilC$_1$ (5) mitochondrial membrane potential (MMP) measurements obtained by flow cytometry in untreated and SLOPE enriched ring-stage *P. falciparum* line MRA 1240 parasites (*N* = 3, error bars represent S.E.M.). (B) Six day of *P. falciparum* line MRA 1240 parasite growth from untreated controls or SLOPE enriched samples diluted with uninfected erythrocytes (*N* = 4, error bars represent S.E.M.). (C) Ring-stage *P. falciparum* line Dd2 infected erythrocytes visualized by Giemsa-stain at 100X magnification.

nondiscriminatory lytic agent (saponin) were lysed ghosts (spectrin positive); however, the same
high erythrocyte membrane to parasite ratio (100:1) that was observed in untreated samples
persisted. In samples treated only with SLO (no Percoll gradient centrifugation), a subset of cells
enriched for infected erythrocytes remained intact (1 of the 4 erythrocytes in **Fig. 3A**). However,
the considerable lysed erythrocyte population remained, leaving the erythrocyte membrane to
parasite ratio at 100:1. Complete SLOPE enrichment demonstrated the effectiveness of Percoll
centrifugation for separating ghosts and intact cells by removing this lysed erythrocyte
population, leaving a >99% intact erythrocyte population that is enriched for parasites (25-fold
decrease in the erythrocyte membrane to parasite ratio; starting of 100:1, final of 4:1)**.** Image
quantification revealed that <0.2% of erythrocytes in SLOPE enriched samples were ghosts (**Fig.**
3B).
173
Parasites retain full viability after SLOPE enrichment
We sought to test the effect of SLOPE enrichment on parasite metabolism and viability. First,
parasites were stained with a mitochondrial membrane potential dependent dye to determine
the fraction of metabolically active parasites. The percentage of parasites with an active
mitochondrial membrane potential was not significantly different between SLOPE enriched
samples and untreated controls (*untreated*=97%, *SLOPE*=94%, *p*=0.46, **Fig. 4A**). To assess
parasite viability on a longer-term scale, we monitored parasite growth following SLOPE
enrichment. This was accomplished by diluting SLOPE enriched parasites with fresh
erythrocytes to reduce parasitemia to appropriate levels for culture (<5%). When compared to
that of parasitemia-matched non-enriched, untreated flasks, the growth rate of parasites from
SLOPE enriched samples showed no growth defects over the time period measured (6 days)

Figure 5. SLOPE enrichment increases detection of ring stage parasite metabolome. (A) Pipeline outlining metabolomics sample preparation and analysis. 1) Ring-stage synchronized *P. falciparum* Dd2 cultures were split into two fractions: one portion was taken for the untreated group and one portion was SLOPE enriched. 2) Equal numbers of erythrocytes from untreated and SLOPE groups were saponin lysed and washed to remove cytosolic erythrocyte metabolites. 3) Metabolism of the resulting pellet containing erythrocyte ghosts and parasites was quenched and metabolites were extracted. 4) Metabolites were identified and quantified from extracts using the AbsoluteIDQ p180 kit. These data were log transformed, centered, and scaled prior to statistical analysis. (B) Principal component analysis was performed on all metabolites detected in at least 50% of samples. Significance between untreated and SLOPE groups was determined by perMANOVA: *p*=0.037. Ellipses show 95% confidence intervals.

185 (**Fig. 4B**). Ring stage parasites retained normal morphology following SLOPE enrichment,
186 further providing evidence that SLOPE enrichment did not damage parasites (**Fig. 4C**).
187
SLOPE-enriched samples exhibit increased parasite metabolites
We hypothesized that a reduction of excess erythrocyte ghosts following SLOPE will lead to
increased parasite signal, especially in ring stage samples. In order to investigate this impact,
we compared synchronous ring stage *P. falciparum* parasites that were either untreated or
SLOPE enriched by 15-fold using a plate-based targeted metabolomics platform that detects
and quantifies up to 180 defined small molecules (*N*=4 per condition, 3×10^8 erythrocytes per
replicate) (**Fig. 5A**). One hundred and
sixteen metabolites were detected in at
least 50% of samples, representing 11
acylcarnitines, 17 amino acids, 5
polyamines, 69 glycerophospholipids,
and 14 sphingolipids. Principal
component analysis revealed distinct
separation of untreated and SLOPE
samples indicating a clear contribution
of the increased concentration of
parasite metabolites (**Fig. 5B**). In
general, lipids contribute heavily to the
separation of untreated and SLOPE
groups with phosphatidylcholines
having a large impact on the metabolic
differentiation of the two groups (**Table
S1**), which is in line with the observation
that this lipid class is dramatically
increased in infected erythrocytes (25).
213
Thirty-two metabolites were significantly
different between untreated and SLOPE
groups after multiple testing correction
(**Table 1**). While some conservation of
metabolites was expected due to the
persistence of erythrocyte membranes,
metabolites that significantly differed
between groups reflected expected
biological differences associated with
increased parasite signal. SLOPE
enriched samples exhibited a

Table 1. SLOPE enrichment leads to significant differences in detection of the ring stage metabolome.
*Denotes metabolites in the polyamine synthesis pathway.

Metabolites	Adjusted p-value	SLOPE/Control
Polyamines		x̄ = 10.3
Putrescine*	0.000848	13.9
Spermidine*	0.000848	11.5
Spermine*	0.00796	5.53
Amino Acids		x̄ = 14.8
Alanine	0.0296	13.2
Arginine*	0.0204	6.94
Asparagine	0.0125	12.3
Aspartic Acid	0.0380	20.3
Glutamic Acid	0.0204	25.0
Lysine	0.0380	3.14
Ornithine*	0.0287	2.73
Phenylalanine	0.0204	8.11
Serine	0.0485	8.92
Threonine	0.0204	17.3
Tyrosine	0.0204	20.8
Valine	0.0170	38.7
Glycerophospholipids		x̄ = 3.13
lysoPC.a.C16:0	0.0485	1.70
lysoPC.a.C17:0	0.0313	1.40
lysoPC.a.C18:1	0.0249	2.67
lysoPC.a.C18:2	0.0204	3.74
PC.aa.C32.0	0.0290	5.91
PC.aa.C36.1	0.0380	5.29
PC.aa.C38.5	0.0485	3.10
PC.aa.C40.4	0.0289	3.57
PC.ae.C38.0	0.0380	2.36
PC.ae.C38.3	0.0287	2.61
PC.ae.C40.1	0.0447	2.79
PC.ae.C40.4	0.0204	2.40
Sphingolipids		x̄ = 5.64
SM.C24.0	0.0290	7.44
SM..OH..C24.1	0.0380	4.58
SM.C24.1	0.0445	4.91
Acylcarnitines		x̄ = 2.19
C2	0.0380	3.78
C16	0.0447	0.60

substantial increase in 12 glycerophospholipids and 3 sphingolipids (3.13 and 5.64 average fold
change, respectively, **Table 1**), which was likely due to the contribution of parasite membranes
and other lipid-containing structures in parasitized erythrocytes (26). Twelve amino acids were
higher in SLOPE samples (14.8 average fold change, **Table 1**), indicative of the breakdown of
hemoglobin by the parasite into free amino acids (27). Additionally, metabolites of the polyamine
synthesis pathway, which is active in the parasite (28, 29), were dramatically increased (10.3
average fold change, **Table 1**).

S*LOPE enriches clinical samples in a cholesterol dependent manner*
In order to determine the utility of SLOPE for the enrichment of clinical *P. falciparum* parasites,
which predominantly circulate as ring stages, we tested SLO lytic discrimination between
infected and uninfected erythrocyte populations directly from human patients. Contrary to results
with *in vitro* propagated parasites (**Fig. 2A**), patient erythrocytes demonstrated reduced SLO
lytic discrimination (**Fig. 6A**, left panel). However, short-term (6h) incubation of clinical samples
in complete media (RPMI with 20% human serum) greatly increased SLO lytic discrimination,
thereby decreasing parasite loss leading to a higher possible final parasitemia (**Fig. 6A**, right

Figure 6. SLOPE is effective on clinical samples in a cholesterol dependent manner. (A) SLO lysis of erythrocytes from *P. falciparum* infected patients either directly isolated from the patient (0h) or after 6h of incubation in complete media (RPMI supplemented with 20% serum). *N*=3 patients. Error bars represent S.E.M. (B) SLO lysis of laboratory *P. falciparum* either grown for 48h in complete media or complete media supplemented with 4mM cholesterol saturated mβCD. *N*=6 (left; 3 replicates each of lines Hb3 and K1); *N*=3 (right; line Hb3). Error bars represent S.E.M. Black circles in selected graphs show iRBC lysis at 27U SLO.

panel). Restoring SLO discrimination occurred with as little as 4h of *ex vivo* incubation but the benefit peaked at ~6h (**Fig. S3**).

We hypothesized that the reduced SLO lytic discrimination observed in samples upon immediate acquisition from patients was due to the increased abundance of cholesterol *in vivo* compared to *in vitro* media formulations. As expected, the *in vivo* environment contained cholesterol at concentrations up to ~250 times higher than those found *in vitro* (**Table S2**). To test the exchange of cholesterol between serum and erythrocyte membranes, we compared the SLO lysis of laboratory *P. falciparum* grown in complete media to parasites grown in complete media supplemented with cholesterol (**Fig. 6B**). This artificial addition of cholesterol led to

265 the predicted increase in infected erythrocyte SLO lysis, closely mimicking the reduced SLO lytic
266 discrimination observed upon immediate acquisition of clinical samples from patients (47.5%
267 lysis and 47.8% lysis in clinical *ex vivo* and cholesterol supplemented *in vitro* samples,
268 respectively, when treated with ~30U SLO; **Fig. 6**, black circles).
269
SLOPE enriches in vitro-derived quiescent parasites
A low frequency of ring stage *P. falciparum* parasites has been reported to enter a quiescent
state upon artemisinin treatment. In order to explore the effectiveness of our enrichment method
on non-traditional erythrocytic forms of the parasite, we treated parasites with dihydroartemisinin
(DHA), removed actively growing parasites daily, and then performed SLOPE to enrich the
remaining metabolically quiescent parasites. When quiescent parasites were quantified using
flow cytometry (SYBR-Green positive/MitoProbe positive), we detected an enrichment in
quiescent parasites by >100-fold (pre-enrichment: undetectable but estimated based on
previous studies to be ~0.005 quiescent parasites/100 erythrocytes (30); post-enrichment: 1
quiescent parasite/100 erythrocytes). Quiescent parasites were also readily visually identifiable
in SLOPE enriched samples under microscopy (**Fig. 7A,** SYBR-Green/MitoTracker Red
positive). When compared to untreated rings, quiescent parasites showed reduced MMP
staining area, indicative of the condensed cytoplasm and reduced metabolism seen in
quiescence (31, 32). While quiescent parasites were enriched by the removal of uninfected
erythrocyte material, erythrocytes containing dead parasites were also enriched by this process

(**Fig. 7A and B**, SYBR-Green
positive, Mitotracker/Mitoprobe
negative parasites). However,
live, quiescent parasites
appeared to be enriched at
considerably higher rates
compared to their dead
counterparts. Specifically, the
dead parasite population
experienced less than 10-fold
enrichment (1.2% immediately
pre-SLOPE versus 9.1% post-
SLOPE), whereas mentioned
above, quiescent-infected
erythrocytes were enriched by
>100-fold. Overall, ~one-tenth
of total enriched parasites
(SYBR-Green positive events)
were quiescent (Mitoprobe

Figure 7. SLOPE is effective on DHA-induced quiescent parasites. (A) Erythrocytes are shown by CD235a staining and *P. falciparum* MRA 1238 parasites are shown by SYBR Green. Within the DHA-treated image, a dead parasite (left) failed to accumulate MitoTracker Deep Red, while two quiescent parasites accumulated MitoTracker Deep Red. (B) Flow cytometry plot measuring quiescent parasites as SYBR Green and Mitoprobe double positive events. Dead parasites are SYBR Green positive, but Mitoprobe negative; uninfected erythrocytes (uRBCs) are SYBR Green and Mitoprobe double negative. iRBC = infected erythrocyte.

304 positive events) (**Fig. 7B**), despite estimates that <<1% of all parasites enter quiescence upon
305 DHA treatment (30).
306
307 **Discussion**
308
For the first time, we can achieve considerable ring stage parasite enrichment that is effective
across a variety of conditions and *Plasmodium* species (**Fig. 1**; **Fig. 2**), as well as on
understudied ring-related populations (clinical isolates, **Fig. 6**; quiescent forms, **Fig. 7**). Our two-
part SLOPE enrichment relies on the cholesterol-dependent lysis of uninfected erythrocytes,
followed by the exploitation of density differences between a parasite-rich intact population and
a ghost population of primarily uninfected erythrocyte membranes (**Fig. 3; Fig. 6**). Importantly,
resulting enriched infected erythrocytes maintain active metabolism and growth (**Fig. 4; Fig. 5;**
Table 1).
317
318 The SLOPE method expands on a discovery made over a decade ago (24) and integrates our
319 metabolic knowledge about the parasite. *Plasmodium* species lack the ability to synthesize
320 cholesterol *de novo* and therefore, the parasite scavenges this lipid from host erythrocyte
321 membranes (33). This scavenging likely leads to the lower levels of cholesterol observed in
322 infected erythrocyte plasma membranes (26). Upon discovery of the preferential action of SLO

Figure 8. Proposed mechanism for decreased susceptibility of infected erythrocytes to SLO lysis. Upon invasion of an erythrocyte, the parasite salvages host membrane cholesterol leading to lower cholesterol levels on the infected erythrocyte surface (top of diagram). During *in vitro* incubation in conditions with sub-physiological cholesterol levels, cholesterol remains low via continued parasite scavenging. Upon exposure physiological levels of cholesterol, such as *in vivo,* the exchange of cholesterol between plasma and erythrocytes restores erythrocyte cholesterol to near pre-invasion levels.

on uninfected erythrocytes, reduced levels of cholesterol in the membranes was proposed as the mechanism of discrimination (24). Our study on parasites from a cholesterol rich environment and the impact of cholesterol addition on SLOPE effectiveness add validity to this hypothesis (**Fig. 6B**). We showed that when parasites were isolated from the human patient (100% serum), SLO discrimination was nominal. Since cholesterol is readily exchanged between the serum environment and erythrocyte membranes (34), we predict that this result is due to the replenishment of parasite-scavenged cholesterol

in the erythrocyte membrane (**Fig. 8**). While enrichment of parasites directly from the
bloodstream of a patient is possible, this is at the expense of parasite number (**Fig. 6A**, **Fig. S3**).
Fortunately, parasite loss can be minimized with a short-term incubation in the laboratory to
reduce erythrocyte membrane cholesterol levels; the limited duration (≤6h) and use of 20%
human serum during this step will likely allow parasites to maintain *in vivo* qualities (e.g.
transcription and metabolic program). Further studies will be required to determine if this is the
case.
351

SLOPE shows utility for the enrichment of non-traditional populations of *Plasmodium* parasites
that were previously understudied due to limited accessibility. In previous studies, clinical
parasites were often fully adapted to *in vitro* culture to generate enough material for
characterization (35, 36). This is not ideal considering parasite populations are altered by the
selective pressure and environmental changes upon transition to *in vitro* culture (37-39). Further,
a sizeable portion of clinical isolates fail entirely to adapt to *in vitro* conditions (40). With the
direct enrichment of parasites from patients, many studies can now be performed with minimal
perturbations (see above) or without culture adaptation entirely.
360

Another form of *P. falciparum* that remains understudied is the artemisinin-induced quiescent
state that is somewhat analogous to persister cells seen in bacteria (41). Populations of both
quiescent parasites and bacterial persisters demonstrate arrested growth leading to decreased
susceptibility in the face of stressful conditions, such as drug treatment. Quiescent and persister
metabolic states share similarities as well, including downregulation of DNA replication, tRNA
synthesis, and oxidative metabolism but maintenance of fatty acid synthesis (42). Following
stressor removal, both populations have the ability to resume growth leading to a recrudescent
infection. Parasite recrudescence following quiescence presents a challenge to the efficacy of
antimalarial therapies. Yet, the rarity of quiescent parasites following drug treatment (<<1% of
all parasites, (30)) limits the study of this phenomenon. Excitingly, SLOPE enrichment increases
the parasitemia of quiescent parasites to readily detectable levels (~1%) through both the
reduction of uninfected erythrocytes and the partial reduction of erythrocytes containing dead
parasites (**Fig. 7**). At detectable levels, it is feasible to further enrich quiescent parasite
populations using other methods (i.e. fluorescence activated cell sorting). The increased
enrichment preference for live, quiescent parasites over dead parasites is likely due to the loss
of cholesterol metabolism in parasites upon death. While quiescent parasites down-regulate
many aspects of metabolism (31), these data indicate that quiescent parasites continue the
utilization of cholesterol.
379

The utility of SLOPE enrichment in increasing access to both *in vitro* rings and non-traditional
parasite forms (quiescent parasites and clinical rings) will reduce many of the previous barriers
to high-quality *Plasmodium* research. Lack of an effective enrichment method for ring stage
Plasmodium has led to considerable host-contributed noise in samples, limiting the success of

sensitive downstream analyses such as proteomics and metabolomics. Specifically, our
previous work demonstrated that characteristics of the erythrocyte batch contributed to the
resultant ring parasite metabolome as heavily as drug treatment (17). Additionally, erythrocyte
noise stymied previous ring stage metabolomics and proteomics experiments as parasite-
derived signals could not reliably be determined over the host erythrocyte contribution (16-18).
Since parasitemia was low in these studies (~1-5 parasites per 100 erythrocytes), this was
undoubtedly due to the high number of erythrocyte ghost membranes that remained in the
preparations. SLOPE significantly reduces the number of ghosts in our preparations, thus
drastically increasing parasitemia (up to >80% in some cases, **Fig. 2B**). We directly observed
this increase in parasite specific metabolites over the host erythrocyte noise using targeted
metabolomics (**Fig. 5B**, **Table 1**). Most notably, we detected an increase in metabolites of the
polyamine synthesis pathway that was proportional to the parasite enrichment level (mean of
~10-fold and 15-fold, respectively). Uninfected erythrocytes lack the machinery necessary for
polyamine synthesis and contain only low levels of putrescine, spermidine, and spermine (43).
However, polyamine levels are much higher in proliferating cell, such as *Plasmodium*, which
both scavenge and synthesize polyamines (28, 29). Given the association between parasite
number and polyamine concentration, we propose to use this group of metabolites as a quality
control marker for parasite-derived metabolites in future metabolomics investigations.
402
The SLOPE method has several advantages for the isolation of material to be used for sensitive
downstream analyses. Firstly, our method can be performed rapidly (~30-40 min). This is equal
to or faster than the amount of time needed to perform the field-accepted purification method for
isolation of late stage parasites by magnetic column purification (~30min to 1hr). The rapid
purification ensures the cellular state of samples is as close as possible to the true phenotype.
Secondly, this method is highly scalable and can be used on culture amounts that range from a
fraction of a milliliter to combined pools of many flasks (hundreds of milliliters). Finally, SLOPE
enrichment is facile and only requires access to a few reagents and a standard centrifuge. While
a fluorescence-based flow cytometer and an automated brightfield cell counter were used to
perform the experiments in this paper, less costly devices, such as a microscope and
hemocytometer, respectively, can be used. The addition of SLOPE to our laboratory toolset will
be useful not only for proteomic and metabolomics studies (see above) but also genomic and
transcriptomic analyses. Reduction of contaminating host DNA and RNA that remain associated
with erythrocytes relative to parasite material will lead to increased sequencing coverage
particularly in clinical samples, which contain a considerable human contribution (44).
418
One caveat of this method may be the cost of SLO lytic agent. Researchers seeking to regularly
enrich large quantities of parasites may consider in-house isolation of SLO from *Streptococcus*
pyogenes culture or recombinant SLO expression in *Escherichia coli* in place of purchasing
commercially available SLO. An additional caveat of SLOPE enrichment is the loss of parasite
material. However, due to the scalability of this method, starting material can be increased to

compensate for this loss. When sample material is limited, researchers may wish to use lower
quantities of SLO to minimize lysis of infected erythrocytes.
426
With SLOPE enrichment as a tool, researchers can more accurately study the ring stage of the
Plasmodium erythrocytic cycle. Improvements in our understanding of parasite biology will lead
to more effective treatments for this deadly disease. Beyond this use, SLOPE has the potential
to be extended to other intracellular parasites and cell types. Many Apicomplexan parasites
scavenge cholesterol from host cells. For example, asexual replication of *Toxoplasma gondii*
within nucleated cells requires cholesterol scavenging with cholesterol being transferred from
the host cell membrane to the parasitophorous vacuole immediately upon invasion (45). This
presents the possibility that other cells types infected with cholesterol scavenging parasites will
also display increased resistance to SLO lysis. Further, SLO causes cholesterol dependent
membrane lysis in cell types other than erythrocytes (46, 47). Thus, the SLOPE method shows
potential for the enrichment of cells infected with other intracellular, cholesterol-scavenging
pathogens, including *Toxoplasma*, *Cryptosporidium*, *Theileria, Chlamydia,* and *Mycobacterium.*
439
440 **Materials and Methods**
441
Parasites and growth
Plasmodium falciparum lines (MRA-155 (Hb3), MRA-159 (K1), MRA 150 (Dd2), MRA 1240, and
MRA 1238, use of respective lines are indicated in figure legends) were obtained from the
Malaria Research and Reference Reagent Resource Center (MR4, BEI Resources); The human
erythrocyte-adapted *Plasmodium knowlesi* line (yH1, (48)) was a gift from Manoj Duraisingh
(Harvard University T.H. Chan School of Public Health). *Plasmodium* cultures were maintained
in A+ human erythrocytes at 3% hematocrit in RPMI 1640 HEPES (Sigma Aldrich, St Louis, MO)
supplemented with either 0.5% Albmumax II Lipid-Rich BSA (Sigma Aldrich, St Louis, MO) and
50 mg/L hypoxanthine (Thermo Fisher Scientific, Waltham, MA) or with 20% heat inactivated
human serum (cRPMI) (Valley Biomedical, Winchester, VA). Cultures were grown at 37$_\circ$C under
5% oxygen, 5% carbon dioxide, 90% nitrogen gas (49). Dilution of cultures with uninfected
erythrocytes to maintain parasitemia at <5% was performed every other day as was changing of
culture medium. Parasitemia was determined by flow cytometry using SYBR-Green staining
(50). Cultures were confirmed negative for mycoplasma monthly using a LookOut Mycoplasma
PCR detection kit (Sigma Aldrich, St Louis, MO).
457
Parasites from infected patients were obtained from adults admitted to the University of Virginia
Health System with clinical malaria. All patients had a recent history of travel to a malaria-
endemic African country, and *P. falciparum* infection was clinically determined by a positive rapid
diagnostic test and peripheral blood smear analysis. Samples were obtained within 24h of
phlebotomy and washed twice with RPMI 1640 HEPES to remove white blood cells. Erythrocytes
were then either immediately used for experiments or kept under short-term *in vitro* culture

conditions in cRPMI. All samples were handled in accordance with protocols approved by the
University of Virginia Institutional Review Board for Health Sciences Research.
466
SLO activation and storage
Streptolyisn-O (SLO, Sigma Aldrich, St Louis, MO) activation was performed as previously
described (24). Following activation, SLO was stored in aliquots at -20$_o$C until use. Hemolytic
units (U) of each SLO batch were quantified from a stored aliquot; one unit equals the amount
of SLO necessary for 50% lysis of 50μl uninfected erythrocytes at 2% hematocrit in PBS for 30
min at 37$_o$C. Hemolytic activity was recurrently assessed (approximately monthly) in triplicate to
control for SLO degradation over time and varying cholesterol levels contributed by different
media batches.
475
SLOPE enrichment
When required, *P. falciparum* cultures were synchronized to the ring stage prior to enrichment
using one round of 5% D-sorbitol (Sigma Aldrich, St Louis, MO) (52). SLO lysis was performed
as previously described but with modifications (24) (**Fig. 1B; Text S1**). Briefly, erythrocyte
density was measured using a Cellometer Auto T4 (Nexcelom Biosciences, Lawrence, MA). Cell
density was adjusted to 2x10$_9$ erythrocytes/mL. The desired amount of SLO (between 0U and
55U) was added in a ratio of 2 parts SLO solution to 5 parts erythrocytes. Samples were mixed
well by pipetting and incubated at room temperature for precisely 6 min. Five-10 volumes of 1X
PBS or non-cholesterol containing media (ex. RPMI 1640 HEPES) were added and cells were
centrifuged at 2,500xg for 3 min. After removal of the supernatant, cells were washed twice more
with 1X PBS or non-cholesterol containing media. Following SLO lysis, cells suspended in 1X
PBS were layered onto a 60% Percoll gradient (Sigma Aldrich, St Louis, MO) and centrifuged at
1,500xg for 5-10 min depending on the volume of the gradient. The top layer of Percoll was
discarded while the lower cell pellet was collected and washed twice with 1X PBS or media.
490
SLO lysis curves
Plasmodium were treated with a range of SLO units as described above. Flow cytometry was
used to assess equal volumes of each sample for every experiment on a BD Acurri C6 flow-
cytometer (BD Biosciences, San Jose, CA.). Total erythrocyte values were obtained by gating
for intact erythrocytes based on forward and side scatter. Infected erythrocyte values were
obtaining by gating for SYBR-Green positive intact erythrocytes. Uninfected erythrocyte values
were obtained by subtracting infected erythrocyte values from total erythrocyte values. Percent
lysis of uninfected and infected erythrocyte populations were determined by comparing the
infected and uninfected erythrocyte values for each SLO-treated sample in an experiment to the
untreated control.
501
502
503

Ghost Quantification

CD235a is constitutively present on the erythrocyte outer surface and thus, both intact erythrocytes and lysed ghosts are CD235a+. Conversely, spectrin is located in the erythrocyte cytoskeleton meaning anti-spectrin cannot access the antibody target in intact erythrocytes. However, pores in lysed erythrocyte membranes (saponin or SLO treated) allow passage of antibodies into the erythrocyte cytosolic compartment making these cells spectrin+. For fluorescent imaging, unfixed samples (average parasitemia of 1%) were blocked with 2% BSA followed by staining in suspension with a 1:100 dilution of mouse anti-alpha I spectrin antibody (Abcam, Cambridge, MA) at $2x10^7$ erythrocytes/mL. Samples were washed then incubated with the fluorescently conjugated secondary antibody, goat anti-mouse Alexa Fluor 594 (Abcam, Cambridge, MA) at 1:1000 dilution. Following additional washes, samples were incubated with the fluorescently conjugated CD235a-PE antibody (Thermo Fisher Scientific, Waltham, MA) at 1:100 and SYBR Green (Thermo Fisher Scientific, Waltham, MA) at 1:10,000. A wet mount was prepared on microscope slides, and samples were immediately imaged. Images were acquired on an EVOS FL Cell Imaging System (Thermo Fisher Scientific, Waltham, MA) using RFP, GFP, and Texas Red light cubes.

Brightfield Microscopy

Slides for brightfield imaging were prepared by fixation in methanol for 1 minute prior to the addition of Giemsa stain for 20 minutes (Sigma Aldrich, St Louis, MO). Brightfield images were obtained on an Eclipse Ci microscope (Nikon, Melville, NY) using an ImagingSource microscope camera and NIS Elements Imaging Software (Nikon, Melville, NY). For parasite stage quantification, slides were prepared for brightfield microscopy as described above both prior to and after SLOPE enrichment. Parasites were categorized by eye as either a ring, trophozoite, or schizont. In the case of a multiply infected erythrocytes, all parasites were counted separately. Gametocytes were excluded from counting.

Re-culturing of SLOPE enriched parasites

P. falciparum cultures were synchronized using 5% D-sorbitol immediately prior to use. The synchronized culture was split to <1% parasitemia and 3% hematocrit to generate an untreated control flask, while the remaining 9 mL of culture were enriched according to the SLOPE protocol described above. Following enrichment, the parasite density of was measured by SYBR-Green based flow cytometry and a "SLOPE" flask was seeded with the number of the enriched cells necessary to match the parasite density of the respective untreated control flask. Uninfected erythrocytes were added to the SLOPE flask to bring hematocrit up to 3%. The parasitemia of the flasks was measured every 48h for 3 replication cycles. Untreated and SLOPE flasks were diluted every 48h to maintain parasitemia levels under 5%. For each dilution event, both flasks were subjected to the same dilution factor. The significance between groups was calculated in Graphpad Prism using a paired t-test.

Mitochondrial membrane potential measurements

The mitochondrial membrane potential (MMP) was assessed using the MitoProbe DiIC1(5) kit (Thermo Fisher Scientific, Waltham, MA). MitoProbe DiIC1(5) accumulates in eukaryotic mitochondria in a mitochondrial membrane potential-dependent manner (53). The effect of SLOPE enrichment on MMP was tested using ring stage synchronized *P. falciparum* cultures. Following SLOPE enrichment, parasites were incubated with 50nM Mitoprobe DiIC$_1$ (5) in media at ~1×10^6 parasites/mL for 30 min at 37$_o$C. Negative controls were treated with the oxidative phosphorylation uncoupler, Carbonyl cyanide m-chlorophenyl hydrazone (CCCP) (Thermo Fisher Scientific, Waltham, MA). Untreated positive controls and were assayed alongside enriched samples. Samples were co-stained with SYBR Green and analyzed using 488nm (for SYBR-Green) and 640nm (for Mitoprobe) filters on a BD Accuri C6 flow cytometer. The percentage of MMP positive parasites was determined as the ratio of Mitoprobe positive to SYBR-Green positive events. The significance between SLOPE treated and untreated groups was calculated in Graphpad Prism using a paired t-test.

Targeted Metabolomics

P. falciparum parasites were synchronized using D-sorbitol 40h prior to sample collection and again, immediately prior to sample collection. Synchronized cultures at 3% parasitemia were either taken directly from culture for untreated samples or were SLOPE enriched to 45% parasitemia. 3.5×10^8 erythrocytes were taken per sample. Samples were lysed in 0.15% saponin as previously described (54). A series of three wash steps was then performed on all samples using PBS to remove soluble erythrocyte metabolites. Metabolites were immediately extracted from the lysed pellet according to the Biocrates p180 Cell Lysis standard operating procedures. Briefly, pellets were resuspended in ice-cold ethanol/0.01 M phosphate buffer (85:15, v/v). Samples were sonicated for 3 minutes, then snap frozen in liquid nitrogen for 30 seconds. Sonication and freezing were repeated followed by a final sonication and centrifugation to pellet insoluble material. The supernatant was taken and stored at -80$_o$C until the day of use.

Metabolite extracts were run using the AbsoluteIDQ p180 kit according to the user manual (Biocrates Life Sciences AG, Innsbruck, Austria). Samples were added to filter spots in each well along with provided internal standards and placed under a gentle stream of nitrogen gas to dry. Fresh phenyl isothiocyanate was then used for derivatization of amines, followed again by plate drying under nitrogen gas. Samples were extracted off of the filter spots using 5mM ammonium acetate in methanol. Sample extracts were either diluted 1:2 with water for LC-MS/MS analysis or diluted 1:50 with the provided flow injection analysis (FIA) mobile phase for FIA-MS/MS analysis. Liquid chromatography and mass spectrometry analysis were performed by the University of Virginia Lipidomics and Metabolomics Core. Chromatographic separation was performed using ACQUITY UPLC system (Waters Corporation, Milford, MA) with an ACQUITY 1.7 μm, 2.1 mm×75 mm BEH C18 column with an ACQUITY BEH C18 1.7 μm, 2.1mm×5mm VanGuard pre-column. Samples for both LC-MS/MS and FIA-MS/MS were

584   analyzed on a Xevo TQ-S Mass Spectrometer (Waters Corporation, Milford, MA) according to
585   the standard operating procedure provided by Biocrates for the AbsoluteIDQ p180kit. All
586   metabolites were identified and quantified against the isotopically labeled internal standards.
587   Raw data was computed in MetIDQ version Nitrogen (Biocrates Life Sciences AG, Innsbruck,
588   Austria).
589
Statistical analysis and visualization of results was performed in R version 3.5.3 with tidyverse,
vegan, and broom (55-57). Only metabolites detected in at least 50% of samples were analyzed
(116 metabolites; **Data Set S1**). Concentration values were not normalized *post-hoc* as equal
cell numbers were input for each sample. Missing values were replaced with half of the lowest
detected value for each metabolite. Values were then log transformed, centered, and scaled
(58). Statistically significant metabolites between SLOPE and untreated groups were determined
using paired t-tests and the Benjamini and Hochberg method for multiple testing correction.
597
Cholesterol manipulation and measurement
Methyl-β-cyclodextrin (mβCD) pre-saturated with cholesterol was purchased from Sigma Aldrich
and dissolved in RPMI 1640 HEPES (Sigma Aldrich, St Louis, MO) to a concentration of 5mM.
The solution was filtered then diluted with human serum (Valley Biomedical, Winchester, VA) to
make a final "cholesterol-rich media" with 20% human serum and a final concentration of 4mM
cholesterol saturated mβCD. *P. falciparum* parasites were incubated in this cholesterol-rich
media for 48h. Samples were then taken for SLO lysis curve experiments as described above.
Cholesterol levels in media formulations were quantified using the Amplex Red Cholesterol
Assay Kit (Thermo Fisher Scientific, Waltham, MA) according to the manufacturer's instructions.
The assay was read using a SpectraMax i3x microplate reader (Molecular Devices, San Jose,
CA) with Invitrogen black-walled, clear-bottom 96-well microplates (Thermo Fisher Scientific,
Waltham, MA). Each replicate measurement of media cholesterol represents a separate heat
inactivated batch of human plasma (Valley Biomedical, Winchester, VA) or a separate 0.5%
AlbuMAX II Lipid Rich BSA preparation (Sigma Aldrich, St Louis, MO).
612
Quiescent Parasite Analysis
Quiescent parasites were generated using the modified Teuscher, *et al*. protocol as described
by Breglio, *et al*. (22, 30). Briefly, synchronized *P. falciparum* cultures were treated with 700nM
dihydroartemisinin (Sigma Aldrich, St Louis, MO) in DMSO for 6h. Cells were washed three times
with RPMI to remove drug before being resuspended in media and returned to culture conditions.
Cultures were then treated with D-sorbitol every 24 hours for the following 72 hours to remove
any actively growing parasites that did not enter quiescence. Immediately following the final
sorbitol treatment, flasks were split into two aliquots. One aliquot was left unenriched to serve
as the control, while the second aliquot was SLOPE enriched with 55 SLO units as described
above. A portion of both untreated and SLOPE samples was then stained with SYBR-Green and

MitoProbe DiIC1(5) as described above for flow cytometry analysis. The remainder of each untreated and SLOPE sample was stained for fluorescence microscopy as described below.

Fluorescence Microscopy

Unenriched and SLOPE enriched samples containing quiescent parasites were stained with anti-CD235a-PE antibody, (Thermo Fisher Scientific, Waltham, MA) at 1:100 and 50nM MitoTracker Red (Thermo Fisher Scientific, Waltham, MA) at 37_oC under 5% CO_2 for 30 min. SYBR-Green (Thermo Fisher Scientific, Waltham, MA) was added for the last 15 min of incubation at 1:10,000. Erythrocytes were washed three times with PBS. A wet mount was immediately prepared on microscope slides, and samples were imaged using a Nikon Eclipse Ti spinning disk confocal microscope at 63X magnification using 488nm, 561nm, and 649nm lasers and the Nikon NIS Elements Software.

Acknowledgements

Special thanks to Michelle Warthan for culturing support and to the UVA Lipidomics and Metabolomics Core for targeted metabolomics run support. This project was supported with funding through the National Institute of Allergy and Infectious Disease R21AI119881 and the University of Virginia 3Cavaliers Grant (to JLG and CCM). ACB is supported by an institutional training grant (5T32GM008136-33).

 Supplemental Materials

644 <u>Supplemental Materials not included in this document:</u>

645 Supplemental Text 1. Detailed SLOPE Protocol (separate .docx attachment)

646 Supplemental Data Set 1. Complete p180 Metabolite Data (separate .csv attachment)

648 <u>All other Supplemental Materials are included below:</u>

649 **Supplemental Table 1. 10 most influential metabolites for principal components 1 and 2**

PC1	PC2
PC.aa.C34.1	*C18*
PC.aa.C38.5	*C18.2*
PC.aa.C38.6	*C18.1*
PC.aa.C32.2	*C16.1.OH*
PC.aa.C36.5	*SM.C16.1*
PC.aa.C30.0	*lysoPC.a.C28.1*
PC.aa.C32.1	*C16*
PC.aa.C34.2	*SM.C20.2*
PC.aa.C36.3	*PC.aa.C28.1*
PC.ae.C38.0	*lysoPC.a.C24.0*

650
651
652 **Supplemental Table 2. Quantification of cholesterol from human plasma and AlbuMAX II Lipid-Rich BSA**

Source	Average	S.E.M.	*N*
Human Plasma	3300 µM	650 µM	4
AlbuMAX II	13.5 µM	1.43 µM	4

Supplemental Figure 1. SLOPE enrichment is effective irrespective of parasite line. SLOPE enrichment performed on three different ring stage synchronized *P. falciparum* lines grown in the same blood and AlbuMAX II media batches. For each graph: *N*=3. Error bars represent S.E.M.

Supplemental Figure 2. SLOPE enrichment shows no bias across stages of the intraerythrocytic cycle. *P. falciparum* line MRA 1240 cultures either (A) synchronized by one sorbitol treatment or (B) left asynchronous were staged by microscopy before and after SLOPE enrichment. For each graph: *N*=3 (200 parasites counted per replicate). Error bars represent S.E.M.

Supplemental Figure 3. SLOPE enrichment is effective irrespective of parasite line. SLOPE enrichment performed on clinical erythroctyes incubated in cRPMI for differing lengths of time. For infected erythrocytes (iRBCs), each line represents one trial; for uninfected erythrocytes (uRBCs), lysis curves from all time points were combined as no change was observed over time: *N*=4. Error bars represent SEM.

References

654
655
1.   Organization WH. 2018. World Malaria Report 2018. WHO, Geneva, Switzerland.
2.   WHO. 2015. Guidelines for the treatment of malaria. Geneva, Switzerland.
3.   Blasco B, Leroy D, Fidock DA. 2017. Antimalarial drug resistance: linking Plasmodium
falciparum parasite biology to the clinic. Nat Med 23:917-928.
4.   Dondorp AM, Nosten F, Yi P, Das D, Phyo AP, Tarning J, Lwin KM, Ariey F, Hanpithakpong
W, Lee SJ, Ringwald P, Silamut K, Imwong M, Chotivanich K, Lim P, Herdman T, An SS,
Yeung S, Singhasivanon P, Day NP, Lindegardh N, Socheat D, White NJ. 2009. Artemisinin
resistance in Plasmodium falciparum malaria. N Engl J Med 361:455-67.
5.   Lu F, Culleton R, Zhang M, Ramaprasad A, von Seidlein L, Zhou H, Zhu G, Tang J, Liu Y,
Wang W, Cao Y, Xu S, Gu Y, Li J, Zhang C, Gao Q, Menard D, Pain A, Yang H, Zhang Q, Cao
J. 2017. Emergence of Indigenous Artemisinin-Resistant Plasmodium falciparum in Africa. N
Engl J Med 376:991-3.
6.   Prevention CfDCa. 2018. About Malaia - Biology. https://www.cdc.gov/malaria/about/biology/.
Accessed
7.   Ribaut C, Berry A, Chevalley S, Reybier K, Morlais I, Parzy D, Nepveu F, Benoit-Vical F,
Valentin A. 2008. Concentration and purification by magnetic separation of the erythrocytic
stages of all human Plasmodium species. Malar J 7:45.
8.   Allman EL, Painter HJ, Samra J, Carrasquilla M, Llinas M. 2016. Metabolomic Profiling of the
Malaria Box Reveals Antimalarial Target Pathways. Antimicrob Agents Chemother 60:6635-
6649.
9.   Cobbold SA, Chua HH, Nijagal B, Creek DJ, Ralph SA, McConville MJ. 2016. Metabolic
Dysregulation Induced in Plasmodium falciparum by Dihydroartemisinin and Other Front-Line
Antimalarial Drugs. J Infect Dis 213:276-86.
10.   Creek DJ, Chua HH, Cobbold SA, Nijagal B, MacRae JI, Dickerman BK, Gilson PR, Ralph SA,
McConville MJ. 2016. Metabolomics-Based Screening of the Malaria Box Reveals both Novel
and Established Mechanisms of Action. Antimicrob Agents Chemother 60:6650-6663.
11.   Olszewski KL, Morrisey JM, Wilinski D, Burns JM, Vaidya AB, Rabinowitz JD, Llinas M.
2009. Host-parasite interactions revealed by Plasmodium falciparum metabolomics. Cell Host
Microbe 5:191-9.
12.   Park YH, Shi YP, Liang B, Medriano CA, Jeon YH, Torres E, Uppal K, Slutsker L, Jones DP.
2015. High-resolution metabolomics to discover potential parasite-specific biomarkers in a
Plasmodium falciparum erythrocytic stage culture system. Malar J 14:122.
13.   Teng R, Lehane AM, Winterberg M, Shafik SH, Summers RL, Martin RE, van Schalkwyk DA,
Junankar PR, Kirk K. 2014. 1H-NMR metabolite profiles of different strains of Plasmodium
falciparum. Biosci Rep 34:e00150.
14.   Parvazi S, Sadeghi S, Azadi M, Mohammadi M, Arjmand M, Vahabi F, Sadeghzadeh S, Zamani
Z. 2016. The Effect of Aqueous Extract of Cinnamon on the Metabolome of Plasmodium
falciparum Using (1)HNMR Spectroscopy. Journal of Tropical Medicine 2016:3174841.
15.   Babbitt SE, Altenhofen L, Cobbold SA, Istvan ES, Fennell C, Doerig C, Llinas M, Goldberg DE.
2012. Plasmodium falciparum responds to amino acid starvation by entering into a hibernatory
state. Proc Natl Acad Sci U S A 109:E3278-87.
16.   Siddiqui G, Srivastava A, Russell AS, Creek DJ. 2017. Multi-omics Based Identification of
Specific Biochemical Changes Associated With PfKelch13-Mutant Artemisinin-Resistant
Plasmodium falciparum. J Infect Dis 215:1435-1444.

17.   Carey MA, Covelli V, Brown A, Medlock GL, Haaren M, Cooper JG, Papin JA, Guler JL. 2018.
Influential Parameters for the Analysis of Intracellular Parasite Metabolomics. mSphere 3.
18.   Dogovski C, Xie SC, Burgio G, Bridgford J, Mok S, McCaw JM, Chotivanich K, Kenny S,
Gnadig N, Straimer J, Bozdech Z, Fidock DA, Simpson JA, Dondorp AM, Foote S, Klonis N,
Tilley L. 2015. Targeting the cell stress response of Plasmodium falciparum to overcome
artemisinin resistance. PLoS Biol 13:e1002132.
19.   David PH, Hommel M, Miller LH, Udeinya IJ, Oligino LD. 1983. Parasite sequestration in
Plasmodium falciparum malaria: spleen and antibody modulation of cytoadherence of infected
erythrocytes. Proceedings of the National Academy of Sciences of the United States of America
80:5075-5079.
20.   Daily J. 2019. Treatment of uncomplicated falciparum malaria in nonpregnant adults and
children. https://www.uptodate.com/contents/treatment-of-uncomplicated-falciparum-malaria-in-
nonpregnant-adults-and-children. Accessed
21.   Codd A, Teuscher F, Kyle DE, Cheng Q, Gatton ML. 2011. Artemisinin-induced parasite
dormancy: a plausible mechanism for treatment failure. Malar J 10:56.
22.   Teuscher F, Gatton ML, Chen N, Peters J, Kyle DE, Cheng Q. 2010. Artemisinin-induced
dormancy in plasmodium falciparum: duration, recovery rates, and implications in treatment
failure. J Infect Dis 202:1362-8.
23.   Witkowski B, Lelievre J, Barragan MJ, Laurent V, Su XZ, Berry A, Benoit-Vical F. 2010.
Increased tolerance to artemisinin in Plasmodium falciparum is mediated by a quiescence
mechanism. Antimicrob Agents Chemother 54:1872-7.
24.   Jackson KE, Spielmann T, Hanssen E, Adisa A, Separovic F, Dixon MW, Trenholme KR,
Hawthorne PL, Gardiner DL, Gilberger T, Tilley L. 2007. Selective permeabilization of the host
cell membrane of Plasmodium falciparum-infected red blood cells with streptolysin O and
equinatoxin II. Biochem J 403:167-75.
25.   Gulati S, Ekland EH, Ruggles KV, Chan RB, Jayabalasingham B, Zhou B, Mantel PY, Lee MC,
Spottiswoode N, Coburn-Flynn O, Hjelmqvist D, Worgall TS, Marti M, Di Paolo G, Fidock DA.
2015. Profiling the Essential Nature of Lipid Metabolism in Asexual Blood and Gametocyte
Stages of Plasmodium falciparum. Cell Host Microbe 18:371-81.
26.   Vial HJ, Eldin P, Tielens AG, van Hellemond JJ. 2003. Phospholipids in parasitic protozoa. Mol
Biochem Parasitol 126:143-54.
27.   Krugliak M, Zhang J, Ginsburg H. 2002. Intraerythrocytic Plasmodium falciparum utilizes only a
fraction of the amino acids derived from the digestion of host cell cytosol for the biosynthesis of
its proteins. Molecular and Biochemical Parasitology 119:249-256.
28.   Phillips MA. 2018. Polyamines in protozoan pathogens. Journal of Biological Chemistry
293:18746-18756.
29.   Ramya TNC, Surolia N, Surolia A. 2006. Polyamine synthesis and salvage pathways in the
malaria parasite Plasmodium falciparum. Biochemical and Biophysical Research
Communications 348:579-584.
30.   Breglio KF, Rahman RS, Sa JM, Roberts DJ, Wellems TE. 2018. Kelch Mutations in
Plasmodium falciparum Protein K13 Do Not Modulate Dormancy after Artemisinin Exposure
and Sorbitol Selection In Vitro. Antimicrob Agents Chemother 62.
31.   Gray KA, Gresty KJ, Chen N, Zhang V, Gutteridge CE, Peatey CL, Chavchich M, Waters NC,
Cheng Q. 2016. Correlation between Cyclin Dependent Kinases and Artemisinin-Induced
Dormancy in Plasmodium falciparum In Vitro. PLoS One 11:e0157906.

32.   Peatey CL, Chavchich M, Chen N, Gresty KJ, Gray KA, Gatton ML, Waters NC, Cheng Q.
746   2015. Mitochondrial Membrane Potential in a Small Subset of Artemisinin-Induced Dormant
747   Plasmodium falciparum Parasites In Vitro. J Infect Dis 212:426-34.

748   33.   Tokumasu F, Crivat G, Ackerman H, Hwang J, Wellems TE. 2014. Inward cholesterol gradient
749   of the membrane system in P. falciparum-infected erythrocytes involves a dilution effect from
750   parasite-produced lipids. Biol Open 3:529-41.

751   34.   Quarfordt SH, Hilderman HL. 1970. Quantitation of the in vitro free cholesterol exchange of
752   human red cells and lipoproteins. J Lipid Res 11:528-35.

753   35.   Mbengue A, Bhattacharjee S, Pandharkar T, Liu H, Estiu G, Stahelin RV, Rizk SS, Njimoh DL,
754   Ryan Y, Chotivanich K, Nguon C, Ghorbal M, Lopez-Rubio JJ, Pfrender M, Emrich S,
755   Mohandas N, Dondorp AM, Wiest O, Haldar K. 2015. A molecular mechanism of artemisinin
756   resistance in Plasmodium falciparum malaria. Nature 520:683-7.

757   36.   Witkowski B, Amaratunga C, Khim N, Sreng S, Chim P, Kim S, Lim P, Mao S, Sopha C, Sam
758   B, Anderson JM, Duong S, Chuor CM, Taylor WRJ, Suon S, Mercereau-Puijalon O, Fairhurst
759   RM, Menard D. 2013. Novel phenotypic assays for the detection of artemisinin-resistant
760   Plasmodium falciparum malaria in Cambodia: in-vitro and ex-vivo drug-response studies. The
761   Lancet Infectious diseases 13:1043-1049.

762   37.   Claessens A, Affara M, Assefa SA, Kwiatkowski DP, Conway DJ. 2017. Culture adaptation of
763   malaria parasites selects for convergent loss-of-function mutants. Scientific reports 7:41303-
764   41303.

765   38.   Daily JP, Scanfeld D, Pochet N, Le Roch K, Plouffe D, Kamal M, Sarr O, Mboup S, Ndir O,
766   Wypij D, Levasseur K, Thomas E, Tamayo P, Dong C, Zhou Y, Lander ES, Ndiaye D, Wirth D,
767   Winzeler EA, Mesirov JP, Regev A. 2007. Distinct physiological states of Plasmodium
768   falciparum in malaria-infected patients. Nature 450:1091.

769   39.   Peters JM, Fowler EV, Krause DR, Cheng Q, Gatton ML. 2007. Differential Changes in
770   Plasmodium falciparum var Transcription during Adaptation to Culture. The Journal of
771   Infectious Diseases 195:748-755.

772   40.   White J, Mascarenhas A, Pereira L, Dash R, Walke JT, Gawas P, Sharma A, Manoharan SK,
773   Guler JL, Maki JN, Kumar A, Mahanta J, Valecha N, Dubhashi N, Vaz M, Gomes E, Chery L,
774   Rathod PK. 2016. In vitro adaptation of Plasmodium falciparum reveal variations in cultivability.
775   Malaria Journal 15:33.

776   41.   Barrett MP, Kyle DE, Sibley LD, Radke JB, Tarleton RL. 2019. Protozoan persister-like cells
777   and drug treatment failure. Nature Reviews Microbiology doi:10.1038/s41579-019-0238-x.

778   42.   Chen N, LaCrue AN, Teuscher F, Waters NC, Gatton ML, Kyle DE, Cheng Q. 2014. Fatty acid
779   synthesis and pyruvate metabolism pathways remain active in dihydroartemisinin-induced
780   dormant ring stages of Plasmodium falciparum. Antimicrob Agents Chemother 58:4773-81.

781   43.   Cooper KD, Shukla JB, Rennert OM. 1976. Polyamine distribution in cellular compartments of
782   blood and in aging erythrocytes. Clin Chim Acta 73:71-88.

783   44.   Shen H-M, Chen S-B, Cui Y-B, Xu B, Kassegne K, Abe EM, Wang Y, Chen J-H. 2018. Whole-
784   genome sequencing and analysis of Plasmodium falciparum isolates from China-Myanmar
785   border area. Infectious diseases of poverty 7:118-118.

786   45.   Coppens I, Joiner KA. 2003. Host but not parasite cholesterol controls Toxoplasma cell entry by
787   modulating organelle discharge. Molecular biology of the cell 14:3804-3820.

788   46.   Hirsch JG, Bernheimer AW, Weissmann G. 1963. MOTION PICTURE STUDY OF THE
789   TOXIC ACTION OF STREPTOLYSINS ON LEUCOCYTES. J Exp Med 118:223-8.

47. Launay J-M, Alouf JE. 1979. Biochemical and ultrastructural study of the disruption of blood
platelets by streptolysin O. Biochimica et Biophysica Acta (BBA) - Biomembranes 556:278-291.
48. Lim C, Hansen E, DeSimone TM, Moreno Y, Junker K, Bei A, Brugnara C, Buckee CO,
Duraisingh MT. 2013. Expansion of host cellular niche can drive adaptation of a zoonotic
malaria parasite to humans. Nature Communications 4:1638.
49. Trager W, Jensen JB. 1976. Human malaria parasites in continuous culture. Science 193:673-5.
50. Bei AK, Desimone TM, Badiane AS, Ahouidi AD, Dieye T, Ndiaye D, Sarr O, Ndir O, Mboup
S, Duraisingh MT. 2010. A flow cytometry-based assay for measuring invasion of red blood
cells by Plasmodium falciparum. Am J Hematol 85:234-7.
51. Kassaza K, Operario DJ, Nyehangane D, Coffey KC, Namugosa M, Turkheimer L, Ojuka P,
Orikiriza P, Mwanga-Amumpaire J, Byarugaba F, Bazira J, Guler JL, Moore CC, Boum Y. 2018.
Detection of <span class="named-content genus-species" id="named-
content-1">Plasmodium Species by High-Resolution Melt Analysis of
DNA from Blood Smears Acquired in Southwestern Uganda. Journal of Clinical Microbiology
56:e01060-17.
52. Lambros C, Vanderberg JP. 1979. Synchronization of Plasmodium falciparum erythrocytic
stages in culture. J Parasitol 65:418-20.
53. Grimberg BT, Jaworska MM, Hough LB, Zimmerman PA, Phillips JG. 2009. Addressing the
malaria drug resistance challenge using flow cytometry to discover new antimalarials.
Bioorganic & medicinal chemistry letters 19:5452-5457.
54. Doolan DL. 2002. Malaria methods and protocols, vol 72. Springer Science & Business Media.
55. David Robinson AH, Matthieu Gomez, Boris Demeshev, Dieter Menne, Benjamin Nutter, Luke
Johnston, Ben Bolker, Francois Briatte, Jeffrey Arnold, Jonah Gabry, Luciano Selzer, Gavin
Simpson, Jens Preussner, Jay Hesselberth, Hadley Wickham, Matthew Lincoln, Alessandro
Gasparini, Lukasz Komsta, Frederick Novometsky, Wilson Freitas, Michelle Evans, Jason Cory
Brunson, Simon Jackson, Ben Whalley, Michael Kuehn, Jorge Cimentada, Erle Holgersen, Karl
Dunkle Werner. 2019. broom: Convert Statistical Analysis Objects into Tidy Tibbles.
https://cran.r-project.org/web/packages/broom/. Accessed
56. Jari Oksanen FGB, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan McGlinn, Peter R.
Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos, M. Henry H. Stevens, Eduard Szoecs,
Helene Wagner. 2019. vegan: Community Ecology Package. https://cran.r-
project.org/web/packages/vegan/. Accessed
57. Wickman H. 2017. tidyverse: Easily Install and Load the 'Tidyverse'. https://cran.r-
project.org/web/packages/tidyverse/. Accessed
58. Sugimoto M, Kawakami M, Robert M, Soga T, Tomita M. 2012. Bioinformatics Tools for Mass
Spectroscopy-Based Metabolomic Data Processing and Analysis. Curr Bioinform 7:96-108.
826