Detection of low-frequency mutations and removal of heat-induced artifactual mutations using Duplex Sequencing

Eun Hyun Ahn[1,2,*], Seung Hyuk Lee[1,3]

[1] Department of Pathology, University of Washington, Seattle, WA 98195, USA; ahneun@uw.edu (E.H.A.); lee0504@uw.edu (S.H.L.)

[2] Institute of Stem Cell and Regenerative Medicine, University of Washington, Seattle, WA 98109, USA; ahneun@uw.edu (E.H.A.)

[3] Department of Bioengineering, University of Washington, Seattle, WA 98195, USA; lee0504@uw.edu (S.H.L.)

* Correspondence: ahneun@uw.edu

Abstract: We present a genome-wide comparative and comprehensive analysis of three different sequencing methods (conventional next generation sequencing (NGS), tag-based single strand sequencing (eg. SSCS), and Duplex Sequencing for investigating mitochondrial mutations in human breast epithelial cells. Duplex Sequencing produces a single strand consensus sequence (SSCS) and a duplex consensus sequence (DCS) analysis, respectively. Our study validates that although high-frequency mutations are detectable by all the three sequencing methods with the similar accuracy and reproducibility, rare (low-frequency) mutations are not accurately detectable by NGS and SSCS. Even with conservative bioinformatical modification to overcome the high error rate of NGS, the NGS frequency of rare mutations is 7.0×10^{-4}. The frequency is reduced to 1.3×10^{-4} with SSCS and is further reduced to 1.0×10^{-5} using DCS. Rare mutation context spectra obtained from NGS significantly vary across independent experiments, and it is not possible to identify a dominant mutation context. In contrast, rare mutation context spectra are consistently similar in all independent DCS experiments. We have systematically identified heat-induced artifactual mutations and corrected the artifacts using Duplex Sequencing. All of these artifacts are stochastically occurring rare mutations. C>A/G>T, a signature of oxidative damage, is the most increased (170-fold) heat-induced artifactual mutation type. Our results strongly support the claim that Duplex Sequencing accurately detects low-frequency mutations and identifies and corrects artifactual mutations introduced by heating during DNA preparation.

Keywords: Duplex Sequencing, Duplex Consensus Sequence (DCS), Single Strand Consensus Sequence (SSCS), Next-Generation Sequencing (NGS), sequencing error, rare mutations, oxidative DNA damage, heat-induced mutations, mitochondrial DNA, human breast cells

1. Introduction

Next-generation sequencing (NGS) has rapidly transformed entire areas of basic research and therapeutic applications by making large scale genomic studies feasible through reduced cost and faster turnaround time [1,2]. NGS has been extensively used to study clonal (high-frequency) mutations, but not subclonal (low-frequency) mutations. A major impediment in investigating subclonal (low-frequency) mutations is that conventional NGS methods have high error rates (10^{-2} to 10^{-3}), which obscure true mutations that occur less frequently than errors [3,4]. These subclonal mutations may account for the genetic heterogeneity of tumors and tumor recurrence, as well as provide a reservoir for the rapid development of resistance to chemotherapy [5].

Conventional sequencing technologies sequence only a single strand of DNA. In contrast, Duplex Sequencing examines both strands of DNA and scores mutations only if they are present on both strands of the same DNA molecule as complementary substitutions. This significantly reduces sequencing error rates to $< 5 \times 10^{-8}$ [6-9]. In the first report of Duplex Sequencing, accuracy and sensitivity of mutation detection were demonstrated mainly in M13mp2 bacteriophage by

comparing untreated/control DNA and DNA incubated with hydrogen peroxide, a radical
generator, in the presence of iron [6].

While overall frequencies and types of mutations from NGS, SSCS, and Duplex Sequencing
have been compared in previous studies [5,10], those studies focused on detection limits of
low-frequency mutations only and did not compare the sequencing methods' ability to detect
high-frequency mutations. In addition, influences of neighboring nucleotide base context on
mutations (mutation context spectra) have not been investigated.

In the current study, we systematically compared mutation frequencies, types, positions, and
sequence context spectra of the whole mitochondrial (mt) DNA in human breast epithelial cells
using three different sequencing protocols: conventional NGS, tag-based single strand consensus
sequencing (eg. SSCS), and Duplex Sequencing. We applied the three sequencing methods to
categorize and analyze high-frequency and low-frequency mutations, separately. Furthermore,
analyses were done with multiple independent DNA library preparation experiments of an identical
biological sample to evaluate the detection consistency, reproducibility, and validity of each
sequencing method. Heating samples, a common practice in preparing DNA for molecular biology
experiments, can introduce such artifactual mutations [11]. Herein, we present heat-induced
artifactual mutations identified using Duplex Sequencing and specific nucleotide contexts that
contribute to a high level of heat-induced artifactual mutations.

2. Results

Duplex Sequencing generates both SSCS and DCS analysis results. In Duplex Sequencing, both
strands of DNA are individually tagged and strands with identical tag sequence, the product of the
same DNA template, are grouped together after PCR amplification. SSCS analysis differs from DCS
analysis in that complementary tag sequences are not identified, and so complementary strands are
not grouped [6]. The SSCS method represents a tag-based single strand sequencing procedure and
is comparable to that of Safe Sequencing System (SafeSeqS) in that each single-stranded DNA
molecule is uniquely labeled before PCR amplification, allowing strands of the same derivatives to
be grouped [9,12].

The average number of nucleotides sequenced at each genome position (depth) of all NGS,
SSCS, and DCS analyses were calculated as the total number of nucleotides sequenced divided by
the mtDNA size of 16569 bases. The depths for NGS, SSCS, and DCS analyses for normal human
breast cells and immortalized cells are presented in Table S1 and Table S2. The highest depths of
NGS, SSCS and DCS that were processed under the same data processing conditions were 458441,
40421 and 6803, respectively (Table S1).

As an attempt to overcome the high error rates of NGS, more conservative bioinformatical
conditions than those applied for SSCS and DCS, referred to as NGS (Q30^r) hereinafter (See section:
Materials and Methods 4.3.2), were applied to NGS datasets. Results of NGS (the same
bioinformatical conditions as to SSCS and DCS) and NGS (Q30^r) are presented in *Supplementary
Figures S1 to S4*. Figure 1 through Figure 4 compare the results of NGS (Q30^r) with those of SSCS
and DCS.

For this study, we have defined homoplasmic (90-100%: Figures 1,2,S1,S2, Table S3) and rare
(0-1%: Figures 3-6,S3-S6, Table S4,S7) mutations based on the mutation occurrence (%) at each
genome position. Mutation frequencies were calculated by dividing the number of variants by the
total number of nucleotides sequenced.

*2.1. Homoplasmic Mutations are Detectable by All Three Methods (NGS, Tag-Based Single Strand
Sequencing, and Duplex Sequencing) with the Similar Accuracy and Reproducibility*

93   The overall frequencies (Figure 1A) of homoplasmic mutations and frequencies of each
94 mutation type (Figure 1B-D) are almost identical across all independent experiments of NGS (Q30^r),
95 SSCS, and DCS analyses.

Figure 1. Frequencies of homoplasmic point mutations in the whole mtDNA of human breast immortalized cells. (**A**) Overall frequencies determined by performing NGS, SSCS, and DCS analyses. Homoplasmic mutation frequency of specific mutation types determined using (**B**) NGS, (**C**) SSCS, and (**D**) DCS analyses. Error bars represent the Wilson score 95% confidence intervals.

Fractions (%) of each mutation type (Figure 2A) and each mutation context spectrum (Figure 2B-D) were examined for homoplasmic mutations. A mutation context spectra analysis identifies bases immediately 5' and 3' to a mutated base (i.e. the mutation appears at the second position of each trinucleotide) and enables classifying observed substitutions into 96 categories (4 bases x 6 substitutions x 4 bases) [13,14].

In our study, fractions (%) of each homoplasmic mutation type (Figure 2A) and their mutation context spectra (Figure 2B-D) are almost same across all the three sequencing methods and independent experiments (Figure 2). All of the homoplasmic mutation types are T>C/A>G or C>T/G>A transitions. T>C/A>G transitions constitute 70% of these mutations and C>T/G>A transitions make up the remaining 30% (Figure 2A). We detected 35 identical homoplasmic unique mutations in all independent experiments regardless of sequencing methods used (Table S3). Taken together, all three sequencing methods are accurate enough to study highly prevalent mutations such as germline mutations of the nuclear genome and homoplasmic mutations of the mitochondrial genome.

Figure 2. Fractions (%) of homoplasmic point mutation types and context spectra in the whole mtDNA of human breast immortalized cells. (**A**) Relative percentages of each mutation type in each experiment as determined by NGS, SSCS, and DCS analyses. Fractions of homoplasmic mutation context spectra determined by (**B**) NGS, (**C**) SSCS, and (**D**) DCS analyses.

2.2. Rarely Occurring Mutations are Neither Accurately Detectable by Conventional NGS Methods nor Tag-Based Single Strand DNA Sequencing, but are Accurately Detectable By Duplex Sequencing

Rare mutation frequencies of human breast immortalized cells were determined using NGS, SSCS, and DCS methods. The average rare mutation frequencies of the independent experiments are significantly lower in SSCS (1.30×10^{-4}) and DCS (1.04×10^{-5}) by 5-fold and 67-fold respectively than that of NGS (Q30^r) (7.00×10^{-4}) (Figure 3A, Table S1). This indicates that Duplex Sequencing removes false-positive artifactual mutations and significantly reduces the rare mutation frequencies.

The frequencies of rare mutations are highly variable in independent experiments analyzed with conventional NGS (Q30^r) (Figure 3A), whereas frequencies of rare mutations show reproducible results in independent experiments of DCS of Duplex Sequencing (Figure 3A,C). It is noted that NGS (Q30^r) datasets were processed under more conservative conditions (See section: *Materials and Methods 4.3.2*) than those of SSCS and DCS; however, these bioinformatical modifications only lowered rare mutation frequency by, on average, 35% (Figure S3,S4 and Table

S1). This indicates that the bioinformatical modification alone is not possible to overcome the high error rate of NGS.

Frequencies of each type of rare mutations reveal other significant differences between NGS (Q30^r), SSCS, and DCS analyses. In NGS (Q30^r) results, C>T/G>A transitions and C>A/G>T transversions are identified at high frequencies (Figure 3D-G). In SSCS results, C>A/G>T transversions are the most predominant mutation type followed by C>T/G>A transitions (Figure 3H,I). In contrast, DCS results indicate that C>T/G>A transitions and C>A/G>T transversions are no longer predominant and no particular type is more prominent than others (Figure 3J,K). Our data suggest that C>A/G>T transversions appear to be the most prevalent type of artifactual mutations that are scored by both NGS (Q30^r) (Figure 3D-G) and SSCS (Figure 3H,I) methods.

Figure 3. Frequencies of rare point mutations in the whole mtDNA of human breast immortalized cells. (**A-C**) Overall frequencies determined by performing NGS, SSCS, and DCS analyses. Rare mutation frequency of each mutation type as determined using (**D-G**) NGS, (**H,I**) SSCS, and (**J,K**) DCS analyses. Error bars represent the Wilson score 95% confidence intervals.

Proportions (%) of each type of rare mutations were analyzed. The prevalent rare mutation types differ under the three sequencing methods. C>A/G>T transversions are the most dominant type of rare mutation with NGS (Q30^r) data (Figure 4A); however the exact fraction (%) of C>A/G>T transversions vary widely across the four independent NGS experiments. In contrast, comparable fractions (%) of each mutation type are observed in both DCS independent experiments (Figure 4A). Significant variations are observed for rare mutation context spectra of NGS (Q30^r) data across four independent experiments (Figure 4B-E) and it is not possible to identify a dominant mutation context among them. In contrast, rare mutation context spectra are similar in all independent DCS experiments. For example, C>T transitions in contexts A<u>C</u>A and A<u>C</u>T occur at persistently high proportions in DCS data (Figure 4H,I).

Figure 4. Fractions (%) of rare mutation types and context spectra in the whole mtDNA of human breast immortalized cells. **(A)** Relative percentages (%) of mutation types in each experiment. Fractions of rare mutation context spectra determined by performing (**B-E**) NGS, (**F,G**) SSCS, and (**H,I**) DCS analyses.

2.3. Duplex Sequencing Identifies and Corrects the Heat-Induced Artifactual Mutations Introduced During DNA Sample Preparation

We investigated which specific types of artifactual mutations are introduced during DNA sample preparation such as heat treatments and to what extent these artifacts can be corrected by Duplex Sequencing. DNA was isolated from human breast primary normal cells (II) and an aliquot of DNA was incubated at 65°C for 9 hours. Unheated DNA served as the control. Libraries of heated and control DNA were prepared for Duplex Sequencing. To identify heat-induced specific mutation types, we performed both SSCS and DCS analyses. The average SSCS and DCS depths of the whole mtDNA genome were similar for control DNA and heated DNA: SSCS (control: 12,257 and heated: 11,622) and DCS (control 2,248 and heated: 2,510) (Table S2). To closely examine the DNA damage-induced artifactual mutations, which are not detectable or distinguishable by conventional sequencing methods, we investigated the rare mutations that include only variants occurring at a frequency of 1% or less using Duplex Sequencing.

The rare mutation frequencies of SSCS are significantly higher than those of DCS (Figure 5A).
This higher SSCS mutation frequency could be due to heat-induced DNA damage and/or errors
during PCR-amplification. Such artifactual mutations are present on only one of the two DNA
strands and thus they are not scored in DCS of Duplex Sequencing. While the incubation of DNA at
65°C significantly increased rare mutation frequency in SSCS analysis (Figure 5A: the first and
second bars: p-value $< 2.2 \times 10^{-16}$), both heated and control DNA displayed identical frequencies of
rare mutations in DCS analysis (Figure 5A: the third and fourth bars). Our results clearly indicate
that DCS analysis by Duplex Sequencing is not affected by heat-induced DNA damage introduced
during DNA sample preparation and correctly represents true mutations.

Figure 5. Frequencies of the heat-induced (65°C) artifactual mutations in human breast normal cells.
(A) Overall rare mutation frequency for heated versus control DNA as determined by SSCS and DCS
analyses. **(B,C)** Frequencies of specific rare mutation types for heated versus control DNA. Error bars
represent the Wilson score 95% confidence intervals. The significant differences in rare mutation
frequencies between the control DNA and the heated DNA are indicated (***$p < 5 \times 10^{-5}$ by the
Chi-square test).

2.4. Duplex Sequencing Identifies the Specific Mutation Spectra of Heat-Induced Artifacts

We further examined which specific mutation type(s) contributed to the elevated SSCS rare
mutation frequency in heated DNA. In SSCS, but not DCS, the heated DNA (Figure 5B) shows a
significant increase in rare mutation frequencies of C>A/G>T, C>T/G>A, C>G/G>C *versus* control
DNA (Figure 5B). In contrast, the 65C° incubation (heating DNA) did not affect the mutation
spectra of DCS results (Figure 5C). For example, the SSCS mutation frequency of C>A in heated
DNA is 3.26×10^{-5}. This heat-induced artifactual mutation type is significantly reduced by 170-fold to
1.88×10^{-7} in DCS analysis.
Fractions (%) of each type (Figure 6A) and each context spectrum of rare mutations (Figure
6B-E) were examined for heated *vs.* control DNA. The heat-induced DNA damage results in
increases in C>G/G>C in SSCS analysis (Figure 6A-C). Out of the 96 possible mutation sequence
contexts, 28 are significantly changed after the 65C° incubation in SSCS analysis (Figure 6B-C, Table
S4). Particularly, CCC, TCC and CCA contexts of C>G mutations showed the most significant
increase in the heated DNA compared to the control (unheated) DNA in SSCS analysis (Figure 6C,

Table S4). In contrast, these DNA damage-dependent substitutions are not observed in DCS
analysis, irrespective of neighboring nucleotides (Figure 6A,D,E).

Figure 6. Fractions (%) of the heat-induced (65°C) artifactual mutation types and context spectra in
human breast normal cells. **(A)** Relative percentages of rare mutation types for heated versus control
DNA as determined by SSCS and DCS analyses. **(B-E)** Rare mutation context spectra for heated
versus control DNA as determined by SSCS and DCS analyses. The significant differences in
percentage of each mutation context between the control untreated DNA and the heated DNA under
SSCS analyses **(C)** are indicated (* $p < 0.05$, ** $p < 5 \times 10^{-4}$, *** $p < 5 \times 10^{-5}$ by the Chi-square test).

218 *2.5 Independent Experiments of Duplex Sequencing Reproducibly Identify the Heat-Induced Artifactual*
219 *Mutaiton Profiles*

Two independent experiments (I and II) for the incubation of DNA at 65°C for 9 hours and the DNA library preparation was conducted with DNA isolated from human epithelial cells (I and II) derived from breast tissue of the same woman. The average SSCS and DCS depths were similar between the cells I and cells II: SSCS (I: 11,622 and II: 11,045) and DCS (I: 2,510 and II: 2,460) (Table S5).

Rare mutation frequencies of heated DNA of the cells I (Figure S5) were calculated for both SSCS and DCS, and the analysis reveals the same pattern observed with heated DNA of the cells II used for the main result Figure 5 experiment (Figure S5). Both the overall rare mutation frequencies and the rare mutation frequency of each mutation type are observed at similar levels between the heated DNA of the cells I and II (Figure S5). Furthermore, fractions (%) of each mutation type (Figure S6A) and mutation context spectra (Figure S6B-E) are almost identical, strengthening the finding that Duplex Sequencing is capable of identifying and correcting the heat-induced artifactual mutations and the results are reproducible in independent experiments.

2.6. All Identified Heat-Induced Artifactual Mutations are Stochastically Occurring Mutations Throughout the Whole Mitochondrial Genome

A total of 3,383 heat-induced artifactual unique mutations were identified, all these are in the mutation occurrence (%) range of 0-1% (Figure 7A). This clearly indicates that all of the heat-induced artifactual mutations introduced during DNA preparation are rarely occurring mutations, thus not accurately and reliably detectable by conventional DNA sequencing methods.

Of the all identified artifactual mutations, about 92% of them are found on coding regions of mtDNA and 69% of them are found within the 13 protein-coding regions of mtDNA (Figure 7A, Table S6). The percentages of artifactual mutations found on coding regions and 13 protein-coding regions of mtDNA closely match the percentages these two regions occupy in the whole mtDNA, which are about 92% and 68% respectively [15]. In addition, the number of artifactual mutations identified shows a strong positive correlation with the sizes of 13 protein-coding genes (Pearson's correlation coefficient R=0.99 and p=2.36x10^{-10}), which indicates that more artifactual mutations are found on larger genes (Figure 7B). We examined the 13 protein-coding genes of mtDNA to see if any particular gene is relatively more or less prone to heat-induced mutations. For each gene, we calculated the percent of variants by dividing the numbers of variants by each gene size (bases). Among the 13 genes, MT-CO2 is slightly more mutated at 23.39% and MT-ND3 is mutated the least at 13.01% (Table S7). The majority (11 out of 13) of these genes are mutated at similar mutation occurrences (average 20%), which is consistent with the finding that artifactual mutations occur stochastically.

	Human mtDNA base size	% base size of the whole mtDNA	No. of variants	% variant of total variants	A
All regions	16569	NA	3383	NA	
Coding regions	15362	92.7	3113	92.0	
Non-coding regions	1207	7.3	270	8.0	
Protein-coding regions	11341	68.4	2328	68.8	

Figure 7. Numbers of the heat-induced (65°C) artifactual mutations in the whole mtDNA of human breast normal cells identified by SSCS and DCS analyses. Variants were counted only once at each

position of the genome. (**A**) Numbers of variants identified in coding, protein-coding, and
non-coding regions of mtDNA. The percentage of variants from each region out of the total number
of identified variants is calculated (% variants). The percentage of base size that each region occupies
out of the total size of mtDNA is calculated (% base size). (**B**) The numbers of the heat-induced
artifactual rare mutations identified in each of 13 protein-coding genes of mtDNA are plotted in the
order of largest to smallest gene size. The Pearson's correlation coefficient for each gene size of the 13
genes *versus* numbers of variants of each gene was 0.99 ($p = 2.36 \times 10^{-10}$).

3. Discussion

In this study, we sequenced the entire mitochondrial DNA of human breast cells via three
different sequencing protocols: NGS, tag-based single strand sequencing (eg. SSCS), and Duplex
Sequencing. We systematically compared high-frequency and low-frequency mutations obtained
from the three methods. We demonstrate advantages of Duplex Sequencing over other sequencing
methods for studying rarely occurring mutations. In addition, we identified the heat-induced
artifactual mutations. Although Duplex Sequencing has been used in a previous study to show an
increased level of mutation frequency of small selected regions of nuclear genome in DNA incubated
at 65°C [11], the same temperature used in this study, exact identities of the heat-induced artifactual
mutations have not been presented. Moreover, while types of artifactual mutations have been
previously examined, influences of neighboring nucleotide base context on artifactual mutations
have not been investigated. To our knowledge, this is the first study to present the exact identities of
the heat-induced artifactual mutations and the specific nucleotide context spectra for these artifacts.
Our data show that rare mutation frequencies are significantly lower in DCS analysis in
comparison to NGS and SSCS analyses, suggesting that large number of rare mutations detected by
NGS and SSCS are mostly artifacts (Figure 3A). Particularly, C>A/G>T transversions, which have
been previously reported to be a predominant result of DNA oxidation [11,16], showed the greatest
decrease in frequencies with DCS analyses of Duplex Sequencing (Figure 3J,K). These findings
validate that DCS can identify and correct artifacts and be applied for accurately detecting rarely
occurring mutations. Furthermore, the comparison of rare mutation frequencies between multiple
independent experiments (Figure 3) demonstrates the ability of DCS in producing reproducible
results. However, the rare mutation data by NGS and SSCS shows high variability across
independent experiments, which indicates the lack of the capability of NGS and SSCS to produce
reliable and reproducible results. The comparison of rare mutation context spectra analyses (Figure
4) further distinguishes DCS from NGS and SSCS by showcasing its advantage to accurately and
consistently detect rarely occurring mutations.
Artifactual mutations can be generated as a result of copying damaged DNA bases. Such
mutations are present on only one of the two DNA strands and are scored by NGS and SSCS but
not by DCS. In the present study, we have identified heat-induced artifactual mutations by
performing both SSCS and DCS analyses. Our results indicate that C>A/G>T is the most
predominantly enhanced mutation artifact followed by C>T/G>A and C>G/G>C. Previous studies
reported that heating DNA can damage DNA bases by forming oxygen free radicals, (specifically
8-hydroxy-2'-deoxyguanine (8-Oxo-dG)), which deaminate cytosine to uracil, and increasing
mitochondrial superoxide anion, which also leads to oxidation of DNA [17-19]. The 8-Oxo-dG is
generated by DNA oxidation under physiopathological conditions or environmental stress. It is also
a by-product of normal cellular metabolism [20]. The formation of 8-oxoguanine, particularly
8-oxo-dG has been reported to cause a high level of C>A/G>T mutations [11,16,20-23], whereas
deamination of cytosine to uracil is known to produce high levels of C>T/G>A mutations [11,19]. In
one study, NGS analysis was done on tumors and matching normal tissues of melanoma and an
enzyme-linked immunosorbent assay (ELISA) for 8-Oxo-dG found that the CCG>CAG context have
a high potential for being a target of DNA oxidation [22]. This context is observed at high mutation
frequencies (2-fold increase) in our current study. Thus, it is likely that the high prevalence of
C>A/G>T transversions in our data is caused by 8-Oxo-dG, suggesting that the biggest contributor
of heat-induced artifactual mutations is oxidative damage to mtDNA.

To our knowledge, this study is the first to examine mutation context spectra of heat-induced
artifactual mutations in the whole mitochondrial DNA of human breast epithelial cells. Our SSCS
analysis results showed that out of 96 possible mutation sequence contexts, the fraction of 28 rare
mutation sequence contexts were significantly changed after the 65C° incubation (Figure 6B-C,
Table S4). Among the affected 28 mutation context spectra, CCC, TCC and CCA contexts of C>G
mutations showed the most significant increase. These mutation contexts could be more prone to
DNA damage and may have a high potential for being targeted by molecular reactions which result
from the heat-damaged DNA.

In summary, we present a genome-wide comprehensive and comparative analysis of
mitochondrial DNA mutations for NGS and tag-based methods (single-strand sequencing and
Duplex Sequencing) and demonstrate the identification and removal of heat-induced artifactual
mutations using Duplex Sequencing. Our results indicate that all of the heat-induced artifactual
mutations are stochastically occurring rare mutations. Thus, these artifactual mutations are not
accurately detectable by conventional sequencing methods. Even the application of more
conservative bioinformatical modification on NGS datasets is not enough to overcome the inherently
high error rate of conventional NGS methods. Our data establishes that Duplex Sequencing: 1)
accurately and reproducibly detect rare (low-frequency) mutations; 2) is not affected by damage
introduced by heating during DNA preparation; 3) identifies and removes the DNA
damage-induced artifactual mutations.

4. Materials and Methods

4.1. Cell Culture

Human breast epithelial cells were provided by Drs. Chia-Cheng Chang and James E. Trosko at
Michigan State University in East Lansing, MI, USA. Human breast normal cells used in this study
were isolated from breast tissues of a healthy (cancer-free) woman 21 years of age obtained during
reduction mammoplasty at Sparrow Hospital in Lansing, MI, USA. Human breast immortalized
cells (M13SV1) were derived from the parental normal stem cells by transforming with SV40 T
antigen [24-29]. Written consents were received from patients. The use of human breast cells was
approved by Michigan State University Institutional Review Board and a Material Transfer
Agreement was approved by both Michigan State University and University of Washington. The
cells were cultured as previously described [30] and were authenticated by short tandem repeat
(STR) DNA profiling (Genetica DNA Laboratories, Labcorp brand, Burlington, NC, USA).

4.2. DNA Extraction, Adapter Synthesis, Library Preparation

DNA extraction and purification, adapter synthesis, and DNA library preparation for Duplex
Sequencing [31] and for whole exome sequencing (WES) [32,33] were carried out as described
previously. For Duplex Sequencing, DNA was extracted using a commercially available DNA
extraction kit (Invitrogen) and the sheared DNA was subjected to end-repair with 3'-dT-tailing for
adapters with A-overhang and with 3'-dA-tailing for adapter with T-overhang. The whole
mitochondrial genome was captured using Agilent SureSelect^{XT} target enrichment (Agilent
Technologies). For WES, the DNA library was hybridized to biotinylated capture probes from the
SureSelect Human All Exon kit (Agilent Technologies). This kit covers 38 Mb of human genome,
corresponding to 23,739 genes in the National Center for Biotechnology Information Consensus
Coding Sequence database.

4.3. Data Processing

350 4.3.1 NGS Datasets and NGS, SSCS and DCS Data Processing

351   Two of four independent NGS datasets (Experiments #1 and #2 in Figure 1 through Figure 4)
352 were obtained by extracting the whole mitochondrial genome data from our WES results. WES is a
353 commonly used next-generation sequencing method that is able to read the entire exome of DNA as

well as the mitochondrial genome [34]. The fastq data files for the two WES datasets were processed as previously described [33] with some modifications. Our in-house script was modified to align the reads with the human mitochondria reference file and to include the mitochondrial genome only.

Two additional independent NGS datasets were obtained by modifying our in-house Duplex Sequencing script to simulate NGS processing on DNA libraries prepared for Duplex Sequencing (Experiments #3 and #4 in Figure 1 through Figure 4). The modified script proceeds only through alignment with a human mtDNA reference but does neither single strand consensus sequence (SSCS) nor duplex consensus sequence (DCS) data alignment steps. This negates the individuality of complementary DNA strands when processing sequence data.

SSCS and DCS datasets were processed as described previously [31]. All datasets were aligned to the Revised Cambridge Reference sequence (rCRS) reference genome, sequence number NC_012920, using BWA and genome analysis toolkit (GATK) software as described previously [31].

4.3.2. Base Quality and PCR Duplicates

An illumina® base quality score of 30 (Q30) is considered the benchmark for a correct base call in NGS. This score refers to a 1 in 1000 chance of an incorrect base call (error probability of 0.1% *vs.* error probability of 5% with the default score of 13) [35] Our NGS (Q30^r) datasets, therefore, were processed with the base quality filtering adjusted to 30 from the default value of 13 by adding "-Q30" to the pileup command (i.e. samtools mpileup –B –d 500000 –Q30 –f [reference] [input] [output]). However, introduction of artifacts in early stages of PCR amplification are not detectable as errors and are embedded in multiple PCR duplicates [6]. SAMtools software is capable of removing these PCR duplicates [36], and so they were removed for NGS (Q30^r) data by taking the combined sequence-1 and sequence-2 files in bam format and using the command "samtools rmdup –s". Both modifications bioinformatically accommodate high background error rates of conventional NGS. For DCS and SSCS data analyses, the default base quality score of 13 (error probability of 5%) was used and PCR duplicate removal was not applied since SSCS and DCS analyses have significantly lower error frequencies (1×10^{-5} and $<5\times10^{-8}$ or 1×10^{-8}, respectively) than that of conventional NGS (10^{-2} to 10^{-3}) [4,9]. Another reason the PCR duplicate removal step is not needed for SSCS and DCS analyses is that the molecular tags mark the duplicates for SSCS and DCS analyses.

The results of NGS analysis obtained under the same bioinformatical conditions as SSCS and DCS (Q13 and no PCR duplicate removal) were presented in Table S1. The effects of bioinformatical modifications (Q13 and no PCR duplicate removal *versus* Q30 and PCR duplicate removal) on NGS mutation results are presented in Supplementary Figures 1-4.

4.3.3 Comparison of Mutation Positions

To identify the heat-induced artifactual mutations, mutation positions were compared after removing the common mutations (present in both control and heated DNA) identified by SSCS and DCS analyses. Mutations found only in the heated DNA from SSCS analysis were considered as heat-induced artifactual mutations. Only genome positions that had minimum sequence read (depth) of 20 in both samples were considered.

4.3.4 Counting mutations

For calculating mutation frequencies, total number of variant reads observed were divided by total number of sequenced reads. For all other analyses, including fractions (%) of each mutation type and mutation context spectra, mutants were scored only once at each position of the genome (i.e. mutants were counted as 1 for each position regardless of number of variant reads observed in that position).

4.4 Statistical Analysis

Differences in mutation frequencies and mutation context fractions between control and heated DNA were analyzed by performing a Chi-square test using R (program version 3.4.4). Association of the number of identified unique variants in each of the thirteen protein-coding genes of mtDNA and the sizes of the corresponding thirteen genes was analyzed by Pearson correlation using Sigma Plot (version 12.0, Systat Software, Inc., San Jose, CA, USA). Differences between the groups were considered significant at $p<0.05$.

Author Contributions: EHA conceived and designed the experiment; EHA performed the experiments; SHL and EHA processed sequencing data and analyzed the data; EHA and SHL wrote the paper.

Acknowledgments: The research was supported by grants from the National Institute of Environmental Health Sciences (NIEHS) P30 ES007033 sponsored-University of Washington (UW) Center for Exposures, Diseases, Genomics and Environment (EDGE) pilot grant (to EH Ahn), UW Office of Research Royalty Research Fund (to EH Ahn), and NCI P30 CA015704-39 Fred Hutchinson Cancer Research Center-UW Cancer Consortium Support Grant (to EH Ahn), National Cancer Institute (NCI) R21 CA220111 (to EH Ahn) and National Cancer Institute of the National Institutes of Health under award number P01 AG001751 and R33 CA181771 (to LA Loeb). The content is solely the responsibility of the authors, and does not necessarily represent the official views of the National Institutes of Health. We thank Dr. Lawrence A. Loeb for critical reading of this manuscript, Dr. Tom Walsh for NGS technical assistance, Drs. Chia-Cheng Chang and James E. Trosko for providing human breast cells, Howard Nebeck for assistance in mutation context analysis and proofreading the manuscript, Clint Valentine, Dr. Michael W. Schmitt, Dr. Woo Young Kim for bioinformatics consultation, and Sujin Kwon for graphical assistance.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

DCS	Duplex Consensus Sequence
DS	Duplex Sequencing
MT	Mitochondria
NGS	Next-generation Sequencing
SSCS	Single Strand Consensus Sequence

References

1. Wetterstrand, K. A. DNA sequencing costs: data from the NHGRI Genome Sequencing Program (GSP) Available online: National Human Genome Research Institute.

2. Goodwin, S.; McPherson, J. D.; McCombie, W. R. Coming of age: ten years of next-generation sequencing technologies. *Nat. Rev. Genet.* **2016**, *17*, 333–351, doi:10.1038/nrg.2016.49.

3. Lou, D. I.; Hussmann, J. A.; McBee, R. M.; Acevedo, A.; Andino, R.; Press, W. H.; Sawyer, S. L. High-throughput DNA sequencing errors are reduced by orders of magnitude using circle sequencing. *Proc. Natl. Acad. Sci. U.S.A.* **2013**, *110*, 19872–19877, doi:10.1073/pnas.1319590110.

4. Fox, E. J.; Reid-Bayliss, K. S.; Emond, M. J.; Loeb, L. A. Accuracy of Next Generation Sequencing Platforms. *Next Gener Seq Appl* **2014**, *1*, doi:10.4172/jngsa.1000106.

5. Loeb, L.A. Human Cancers Express a Mutator Phenotype: Hypothesis, Origin, and Consequences. *Cancer Research* **2016**, *76*, 2057–2059, doi:10.1158/0008-5472.CAN-16-0794.

6. Schmitt, M. W.; Kennedy, S. R.; Salk, J. J.; Fox, E. J.; Hiatt, J. B.; Loeb, L. A. Detection of ultra-rare mutations by next-generation sequencing. *Proc. Natl. Acad. Sci. U.S.A.* **2012**, *109*, 14508–14513, doi:10.1073/pnas.1208715109.

7. Kennedy, S. R.; Schmitt, M. W.; Fox, E. J.; Kohrn, B. F.; Salk, J. J.; Ahn, E. H.; Prindle, M. J.; Kuong, K. J.; Shen, J.-C.; Risques, R.-A.; Loeb, L. A. Detecting ultralow-frequency mutations by Duplex Sequencing. *Nat Protoc* **2014**, *9*, 2586–2606, doi:10.1038/nprot.2014.170.

8. Schmitt, M. W.; Fox, E. J.; Prindle, M. J.; Reid-Bayliss, K. S.; True, L. D.; Radich, J. P.; Loeb, L. A. Sequencing small genomic targets with high efficiency and extreme accuracy. *Nature Methods* **2015**, *12*, 423–425, doi:10.1038/nmeth.3351.

9. Salk, J. J.; Schmitt, M. W.; Loeb, L. A. Enhancing the accuracy of next-generation sequencing for detecting rare and subclonal mutations. *Nature Reviews Genetics* **2018**, *19*, 269–285, doi:10.1038/nrg.2017.117.

10. Newman, A. M.; Lovejoy, A. F.; Klass, D. M.; Kurtz, D. M.; Chabon, J. J.; Scherer, F.; Stehr, H.; Liu, C. L.; Bratman, S. V.; Say, C.; Zhou, L.; Carter, J. N.; West, R. B.; Sledge Jr, G. W.; Shrager, J. B.; Loo, B. W.; Neal, J. W.; Wakelee, H. A.; Diehn, M.; Alizadeh, A. A. Integrated digital error suppression for improved detection of circulating tumor DNA. Nature Biotechnology 2016, 34, 547–555, doi:10.1038/nbt.3520.

11. Arbeithuber, B.; Makova, K. D.; Tiemann-Boege, I. Artifactual mutations resulting from DNA lesions limit detection levels in ultrasensitive sequencing applications. DNA Research 2016, 23, 547–559, doi:10.1093/dnares/dsw038.

12. Kinde, I.; Wu, J.; Papadopoulos, N.; Kinzler, K. W.; Vogelstein, B. Detection and quantification of rare mutations with massively parallel sequencing. *Proceedings of the National Academy of Sciences* **2011**, *108*, 9530–9535, doi:10.1073/pnas.1105422108.

13. Alexandrov, L. B.; Nik-Zainal, S.; Wedge, D. C.; Aparicio, S. A. J. R.; Behjati, S.; Biankin, A. V.; Bignell, G. R.; Bolli, N.; Borg, A.; Børresen-Dale, A.-L.; Boyault, S.; Burkhardt, B.; Butler, A. P.; Caldas, C.; Davies, H. R.; Desmedt, C.; Eils, R.; Eyfjörd, J. E.; Foekens, J. A.; Greaves, M.; Hosoda, F.; Hutter, B.; Ilicic, T.; Imbeaud, S.; Imielinski, M.; Imielinsk, M.; Jäger, N.; Jones, D. T. W.; Jones, D.; Knappskog, S.; Kool, M.; Lakhani, S. R.; López-Otín, C.; Martin, S.; Munshi, N. C.; Nakamura, H.; Northcott, P. A.; Pajic, M.; Papaemmanuil, E.; Paradiso, A.; Pearson, J. V.; Puente, X. S.; Raine, K.; Ramakrishna, M.; Richardson, A. L.; Richter, J.; Rosenstiel, P.; Schlesner, M.; Schumacher, T. N.; Span, P. N.; Teague, J. W.; Totoki, Y.; Tutt, A. N. J.; Valdés-Mas, R.; van Buuren, M. M.; van 't Veer, L.; Vincent-Salomon, A.; Waddell, N.; Yates, L. R.; Australian Pancreatic Cancer Genome Initiative; ICGC Breast Cancer Consortium; ICGC MMML-Seq Consortium; ICGC PedBrain; Zucman-Rossi, J.; Futreal, P. A.; McDermott, U.; Lichter, P.; Meyerson, M.; Grimmond, S. M.; Siebert, R.; Campo, E.; Shibata, T.; Pfister, S. M.; Campbell, P. J.; Stratton, M. R. Signatures of mutational processes in human cancer. *Nature* **2013**, *500*, 415–421, doi:10.1038/nature12477.

14. Pilati, C.; Shinde, J.; Alexandrov, L. B.; Assié, G.; André, T.; Hélias-Rodzewicz, Z.; Ducoudray, R.; Le Corre, D.; Zucman-Rossi, J.; Emile, J.-F.; Bertherat, J.; Letouzé, E.; Laurent-Puig, P. Mutational signature analysis identifies *MUTYH* deficiency in colorectal cancers and adrenocortical carcinomas: Mutational signature associated with *MUTYH* deficiency in cancers. *The Journal of Pathology* **2017**, *242*, 10–15, doi:10.1002/path.4880.

15. MITOMAP A human mitochondrial genome database http://www.mitomap.org/.

16. Cheng, K. C.; Cahill, D. S.; Kasai, H.; Nishimura, S.; Loeb, L. A. 8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes G----T and A----C substitutions. *J. Biol. Chem.* **1992**, *267*, 166–172.

17. Bruskov, V. I.; Malakhova, L. V.; Masalimov, Z. K.; Chernikov, A. V. Heat-induced formation of reactive oxygen species and 8-oxoguanine, a biomarker of damage to DNA. *Nucleic Acids Res.* **2002**, *30*, 1354–1363.

18. Slimen, I. B.; Najar, T.; Ghram, A.; Dabbebi, H.; Ben Mrad, M.; Abdrabbah, M. Reactive oxygen species, heat stress and oxidative-induced mitochondrial damage. A review. *Int J Hyperthermia* **2014**, *30*, 513–523, doi:10.3109/02656736.2014.971446.

19. Kang, Q.; Parkin, B.; Giraldez, M. D.; Tewari, M. Mutant DNA quantification by digital PCR can be confounded by heating during DNA fragmentation. *BioTechniques* **2016**, *60*, 175–176, 178, 180 passim, doi:10.2144/000114401.

20. Cooke, M. S.; Evans, M. D.; Dizdaroglu, M.; Lunec, J. Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J.* **2003**, *17*, 1195–1214, doi:10.1096/fj.02-0752rev.

21. Marnett, L. J. Oxyradicals and DNA damage. *Carcinogenesis* **2000**, *21*, 361–370.

22. Costello, M.; Pugh, T. J.; Fennell, T. J.; Stewart, C.; Lichtenstein, L.; Meldrim, J. C.; Fostel, J. L.; Friedrich, D. C.; Perrin, D.; Dionne, D.; Kim, S.; Gabriel, S. B.; Lander, E. S.; Fisher, S.; Getz, G. Discovery and characterization of artifactual mutations in deep coverage targeted capture sequencing data due to oxidative DNA damage during sample preparation. *Nucleic Acids Res.* **2013**, *41*, e67, doi:10.1093/nar/gks1443.

23. Chen, L.; Liu, P.; Evans, T. C.; Ettwiller, L. M. DNA damage is a pervasive cause of sequencing errors, directly confounding variant identification. *Science* **2017**, *355*, 752–756, doi:10.1126/science.aai8690.

24. Kao, C. Y.; Nomata, K.; Oakley, C. S.; Welsch, C. W.; Chang, C. C. Two types of normal human breast epithelial cells derived from reduction mammoplasty: phenotypic characterization and response to SV40 transfection. *Carcinogenesis* **1995**, *16*, 531–538.

25. Kang, K. S.; Morita, I.; Cruz, A.; Jeon, Y. J.; Trosko, J. E.; Chang, C. C. Expression of estrogen receptors in a normal human breast epithelial cell type with luminal and stem cell characteristics and its neoplastically transformed cell lines. *Carcinogenesis* **1997**, *18*, 251–257.

26. Chang, C. C.; Sun, W.; Cruz, A.; Saitoh, M.; Tai, M. H.; Trosko, J. E. A human breast epithelial cell type with stem cell characteristics as target cells for carcinogenesis. *Radiat. Res.* **2001**, *155*, 201–207.

27. Park, J.-S.; Noh, D.-Y.; Kim, S.-H.; Kim, S.-H.; Kong, G.; Chang, C.-C.; Lee, Y.-S.; Trosko, J. E.; Kang, K.-S. Gene expression analysis in SV40-immortalized human breast luminal epithelial cells with stem cell characteristics using a cDNA microarray. *Int. J. Oncol.* **2004**, *24*, 1545–1558.

28. Tai, M.-H.; Chang, C.-C.; Kiupel, M.; Webster, J. D.; Olson, L. K.; Trosko, J. E. Oct4 expression in adult human stem cells: evidence in support of the stem cell theory of carcinogenesis. *Carcinogenesis* **2005**, *26*, 495–502, doi:10.1093/carcin/bgh321.

29. Ahn, E. H.; Chang, C.-C.; Schroeder, J. J. Evaluation of sphinganine and sphingosine as human breast cancer chemotherapeutic and chemopreventive agents. *Exp. Biol. Med. (Maywood)* **2006**, *231*, 1664–1672.

30. Ahn, E. H.; Lee, S. H.; Kim, J. Y.; Chang, C.-C.; Loeb, L. A. Decreased Mitochondrial Mutagenesis during Transformation of Human Breast Stem Cells into Tumorigenic Cells. *Cancer Res.* **2016**, *76*, 4569–4578, doi:10.1158/0008-5472.CAN-15-3462.

31. Ahn, E. H.; Hirohata, K.; Kohrn, B. F.; Fox, E. J.; Chang, C.-C.; Loeb, L. A. Detection of Ultra-Rare Mitochondrial Mutations in Breast Stem Cells by Duplex Sequencing. *PLoS ONE* **2015**, *10*, e0136216, doi:10.1371/journal.pone.0136216.

32. Walsh, T.; Shahin, H.; Elkan-Miller, T.; Lee, M. K.; Thornton, A. M.; Roeb, W.; Abu Rayyan, A.; Loulus, S.; Avraham, K. B.; King, M.-C.; Kanaan, M. Whole Exome Sequencing and Homozygosity Mapping Identify Mutation in the Cell Polarity Protein GPSM2 as the Cause of Nonsyndromic Hearing Loss DFNB82. *The American Journal of Human Genetics* **2010**, *87*, 90–94, doi:10.1016/j.ajhg.2010.05.010.

33. Gulsuner, S.; Walsh, T.; Watts, A. C.; Lee, M. K.; Thornton, A. M.; Casadei, S.; Rippey, C.; Shahin, H.; Consortium on the Genetics of Schizophrenia (COGS); PAARTNERS Study Group; Nimgaonkar, V. L.; Go, R. C. P.; Savage, R. M.; Swerdlow, N. R.; Gur, R. E.; Braff, D. L.; King, M.-C.; McClellan, J. M. Spatial and temporal mapping of de novo mutations in schizophrenia to a fetal prefrontal cortical network. Cell **2013**, 154, 518–529, doi:10.1016/j.cell.2013.06.049.

34. Griffin, H. R.; Pyle, A.; Blakely, E. L.; Alston, C. L.; Duff, J.; Hudson, G.; Horvath, R.; Wilson, I. J.; Santibanez-Koref, M.; Taylor, R. W.; Chinnery, P. F. Accurate mitochondrial DNA sequencing using off-target reads provides a single test to identify pathogenic point mutations. *Genetics in Medicine* **2014**, *16*, 962–971, doi:10.1038/gim.2014.66.

35. Cliften, P. Chapter 7: Base Calling, Read Mapping, and Coverage Analysis. In *Clinical Genomics*, 1st ed.; Kulkarni, S., Pfeifer, J., Eds; Elsevier Inc.: New York, U.S., **2015**; Volume 1, pp. 91-107; 9780124041488.

36. Li, H.; Handsaker, B.; Wysoker, A.; Fennell, T.; Ruan, J.; Homer, N.; Marth, G.; Abecasis, G.; Durbin, R.; 1000 Genome Project Data Processing Subgroup The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **2009**, *25*, 2078–2079, doi:10.1093/bioinformatics/btp352.